Systematic integration of protein affecting mutations, gene fusions, and copy number alterations into a comprehensive somatic mutational profile

Shawn S Striker [1], Sierra F Wilferd [1], Erika M Lewis [1], Samantha A O'Connor [1], Christopher L Plaisier [1,2,*]

[1] School of Biological and Health Systems Engineering, Fulton Schools of Engineering, Arizona State University, Tempe, AZ 85287-9709, USA.

[2] Lead contact.

*To whom correspondence should be addressed: Christopher Plaisier, E-mail: plaisier@asu.edu, Phone: (480) 965-6832, Address: P.O. Box 879709, Tempe, AZ 87287-9709

Motivation

Genes are mutated in tumors through either PAMs, CNAs, or gene fusions, thereby splitting the signal of the somatic mutation effect across these three mutation types. We developed OncoMerge to systematically integrate the three mutation types into a single mutation profile that better captures the impact of somatic mutations on cancer phenotypes. As a tool OncoMerge fills the gap between the sophisticated variant calling pipelines and downstream analyses.

Summary

Somatic mutations occur as random genetic changes in genes through protein-affecting mutations (PAMs), gene fusions, or copy number alterations (CNAs). Mutations of different types can have a similar phenotypic effect (i.e., allelic heterogeneity) and should be integrated into a unified gene mutation profile. We developed OncoMerge to fill this niche of integrating somatic mutations to capture allelic heterogeneity, assign a function to mutations, and overcome known obstacles in cancer genetics. Application of OncoMerge to the TCGA Pan-Cancer Atlas increased the number and frequency of somatically mutated genes and improved the prediction of the somatic mutation role as either activating or loss of function. Using integrated somatic mutation matrices increased the power to infer gene regulatory networks and uncovered the enrichment of switch-like feedback motifs and delay-inducing feed-forward loops. These studies demonstrate that OncoMerge efficiently integrates PAMs, fusions, and CNAs and strengthens downstream analyses linking somatic mutations to cancer phenotypes.

Keywords: somatic mutations, allelic heterogeneity, cancer, oncogene, tumor suppressor gene

Introduction

The accumulation of somatic mutations in patient tumors drives and reinforces cancer phenotypes. The three main types of somatic mutations that modify the function of a gene or render it non-functional are: 1) protein affecting mutations (PAMs), 2) gene fusions, and 3) copy number alterations (CNAs). A PAM is a point mutation, short insertion, or short deletion inside a gene's coding region or splice sites[1]. Gene fusions occur when genomic rearrangements join two genes into a novel chimeric gene or place a promoter in front of a new gene, causing misexpression[2]. Finally, CNAs occur frequently in tumors where whole chromosomes, chromosomal arms, or localized genomic segments are duplicated or deleted[3,4]. Somatic mutation via PAM, gene fusion, or CNA can have similar effects on cancer phenotypes, i.e., allelic heterogeneity. This interchangeability and the erratic circumstances that produce somatic mutations lead to the mixture of mutation types observed in large cohorts of patient tumors[1].

Describing how somatic mutations in a gene impact cancer phenotypes requires integrating the information from all three mutation types. Most studies linking somatic mutations to cancer phenotypes focus on one mutation type. This leads to missing associations for mutations primarily found in another type and reduced power to detect associations for mutations with high allelic heterogeneity that span the mutation types. Thus, a current obstacle facing those studying the downstream effects of somatic mutations is the lack of an established method for integrating PAMs, gene fusions, and CNAs into a comprehensive gene mutation profile. The lack of integration methods is due to several complicating factors. Firstly, the allelic heterogeneity observed in and between tumors means that different mutations in the same gene can be equivalently oncogenic. Second, it is challenging to discern driver (causal) from passenger (non-causal) somatic mutations. Third, an algorithm must be able to systematically integrate the binary PAM and gene fusion (mutated or not) with the quantitative copy number from CNAs. Lastly, some tumors have drastically higher somatic mutation rates than others (e.g., microsatellite instability[5] and hypermutation[6]). These higher mutation rates confound any frequency-based integration approach and drive the discovery of spurious somatic mutations. We developed OncoMerge to fill the somatic mutation integration niche by providing an algorithm that systematically overcomes these obstacles to generate an integrated gene mutation profile. The input for the OncoMerge algorithm is the output from state-of-the-art methods for detecting PAMs (MC3[1] and MutSig2CV[7]), transcript fusions (PRADA[2,8]), and CNAs (GISTIC2.0[9]). Each method provides the likelihood that a somatic mutation happens by chance alone. Filtering on these statistics focuses integration efforts on genes most likely to harbor functional mutations. The integrated mutation profiles improve the power to detect associations with cancer phenotypes leading to a more comprehensive understanding of how genetic alterations drive cancer phenotypes.

The tremendous amount of cancer genome sequencing data generated in the last ten years has enabled efforts to discover and catalog somatic mutations across many cancers[1,10]. Many algorithms have been developed to discern which somatic mutations are drivers, how the mutations affect genes[6,7,11–17], and databases to search and view somatically mutated driver genes[16–18]. There also exist approaches for integrating somatic mutations. OncoPrint from cBioportal[18] can visually overlay somatic mutation types across patient tumors for a gene of interest. The OncodriveROLE algorithm[13] was developed to discover driver genes by systematically integrating PAMs and CNVs. However, neither OncoPrint nor OncodriveROLE provides an integrated mutational profile that can be used in downstream analyses. The impact

of somatic mutations can be classified as activating (Act) gene function (typically found in
oncogenes) or loss of function (LoF) (typically found in tumor suppressor genes)[13]. It has also
been demonstrated that the systematic integration of PAM and CNA somatic mutations for a
gene improves the ability to determine Act or LoF status[13]. These foundational studies have
created a platform to develop an algorithm that systematically integrates the three somatic
mutation types.

The systematic integration of somatic mutations requires choosing a gene-level model that
determines how the data for the three somatic mutation types will be integrated, the somatic
mutation role. We determine the somatic mutation role by employing rules similar to those in
OncodriveROLE[13] (**Figure 1**). The possible somatic mutation roles in OncoMerge are PAM,
Fusion, CNA amplification (CNAamp), CNA deletion (CNAdel), Act, or LoF. The PAM, Fusion,
CNAamp, and CNAdel somatic mutation roles use the unintegrated somatic mutation profile for
the chosen role in the final mutation matrix. The Act and LoF are integrated mutation roles that
harness allelic heterogeneity. Allelic heterogeneity is especially prevalent in tumor suppressor
genes, where mutations at many positions in a gene disrupt its function to prevent cancer
phenotypes[3]. Allelic heterogeneity is less prevalent for oncogenes where a small number of
specific gain of function alleles are needed to drive cancer phenotypes[3]. Genes underlying
CNAs can add another layer of information as tumor suppressors are often deleted, which has
an equivalent oncogenic effect as missense or truncating PAMs. The LoF role is designated
when PAMs, Fusions, and CNAdels are integrated. Oncogenes are often amplified as this
typically leads to overexpression of the underlying genes, which has a similar positive effect on
gene function as a gain of function PAM. The Act role is designated when PAMs, Fusions, and
CNAamps are integrated. Systematic determination of the somatic gene role and application of
the rules laid out above are used to integrate the three mutation types into a comprehensive
somatic mutation profile.

The lists of somatically mutated driver genes provide a set of gold standard mutations with
somatic mutation roles that can be used to assess the performance of OncoMerge. The gold
standards are classified by whether the somatic mutation of a gene was cancer-specific or not.
The TCGA consensus[6] and Cancer Gene Census (CGC) from COSMIC[11] were used to develop
gold standards with cancer-specific somatically mutated gene roles. The 20/20 rule[3],
OncodriveROLE[13], and Tokheim ensemble[12] were used to create gold standards with
somatically mutated gene roles. Comparisons of somatic mutation role between OncoMerge
and the gold standards were facilitated by converting oncogenes to Act and tumor suppressors
to LoF. Finally, a combined gene role agnostic gold standard was developed based on a union
of all somatic mutations from all five gold standards. These gold standards were used to assess
the utility of filters and the quality of the OncoMerge integrated somatic mutation matrices
through their ability to recall somatic mutations with the appropriate gene role. We chose five
gold standards that employed different algorithms for somatic mutation discovery to avoid
overfitting to any one gold standard when assessing the performance of OncoMerge.

OncoMerge is designed to construct a comprehensive somatic mutation profile that increases
the power to link mutations with cancer phenotypes. Previously, we have used the Systems
Genetics Network AnaLysis (SYGNAL) pipeline[19] to build causal and mechanistic gene
regulatory networks (GRNs) for 31 cancers from the TCGA Pan-Cancer Atlas[20]. Using
SYGNAL, we linked somatic mutations through transcription factor (TF) and miRNA regulators
to the hallmarks of cancer[21,22], thereby linking somatic mutations to cancer phenotypes. We

hypothesize that integrated somatic mutation matrices from OncoMerge will increase our power
to infer causal relationships for pan-cancer SYGNAL networks and that these will yield novel
biological insights.

Results

Establishing a baseline for the integration of somatic mutations

We developed OncoMerge as a systematic method to integrate PAM, fusion, and CNA somatic
mutations into a more comprehensive mutation matrix for subsequent analyses. OncoMerge
systematically integrates somatic mutations and defines a role for each gene (**Figure 1**): PAM,
fusion, CNA deletion (CNAdel), CNA amplification (CNAamp), Activating (Act), and Loss of
Function (LoF). The role assigned to a gene describes the rubric used to integrate the data from
the source data matrices.

A significant part of developing OncoMerge was constructing and optimizing the statistical filters
that provide an essential quality control step to identify integrated somatically mutated genes
more likely to be functional in tumor biology. The selection and optimization of OncoMerge
statistical filters were performed using the 9,584 patient tumors from 32 cancers profiled by the
TCGA Pan-Cancer Atlas[1,6,20] (cancer type abbreviations can be found in STAR⬚Methods). We
used three metrics to assess the value of potential filters: 1) impact on the number of
somatically mutated genes (**Figure 2A**); 2) impact on the distribution of the number of genes
mapping to genomic loci (**Figure 2B**); and 3) significance of the overlap between somatically
mutated genes from OncoMerge with gold standard datasets (including overlap with gene roles
and tumor-specific gene roles; **Figure 2C**; **Table S2** & **S3**). These metrics ensure that the
integrated somatic mutations are consistent with prior knowledge and that the size of CNA
mutations does not overwhelm the integration algorithm.

First, we needed to prove that the seed genes alone would not drive significant enrichment for
the gold standard comparisons. We applied OncoMerge to somatic mutation matrices with
randomized gene labels from the TCGA Pan-Cancer Atlas without any filters applied. None of
the gold standards significantly overlapped with the OncoMerge identified somatic mutations (all
p-values ≥ 0.4; **Table S4**). Thus, even though MutSig2CV and GISTIC 2.0 determine the seeds
genes, the actual somatic mutation data and filters are required to achieve the full integration
potential of OncoMerge. This result from applying OncoMerge to randomized data demonstrates
that the signal from the gold standards is not driven by the seed genes alone. Thus we can
safely use the gold standards to assess the performance of OncoMerge.

Next, we determined the integration baseline by applying OncoMerge to the TCGA Pan-Cancer
Atlas without filtering. Slightly less than one-third of the genome was considered somatically
mutated in at least 5% or greater of tumors in at least one of the 32 cancers (30% or 6,028
genes, **Figure 2A**). We observed a significant overlap between OncoMerge somatically mutated
genes and the combined gold standard (genes = 395, p-value = 1.1 x 10^{-44}, **Figure 2C**) when
gene role was not considered. Significant overlaps existed between the LoF somatic mutations
from three gold standards (TCGA consensus, CGC, and Vogelstein) with the somatic mutations
with the LoF predicted role from OncoMerge (**Figure 2C**). None of the comparisons of Act
somatic mutations were significantly overlapping (**Figure 2C**). Many of the 6,028 genes map to
the same copy number alteration genomic locus (**Figure 2B**). These unfiltered results reveal
two main integration biases. First, the overlaps were insignificant between Act somatic
mutations and previously identified Act mutations. Second, the integration with CNAs is causing

the inclusion of many passenger mutations mapping to the same genomic locus. OncoMerge
applied to the TCGA Pan-Cancer Atlas without filtering provides a baseline to benchmark
success. Addressing the integration biases we observed is our impetus for developing and
optimizing filters for OncoMerge.

Developing a filtering strategy for the integration of somatic mutations
The power of integration is that aggregating somatic mutation information can boost the
mutation frequency of a gene enough to become significant, even though the constituent
mutations do not reach significance alone. The first filter determined if the final mutation
frequency after integrating PAM, fusion, and CNA somatic mutations is larger than expected by
chance alone. A permutation-based approach empirically determined the background integrated
mutation frequency distribution. Then the observed frequencies are compared to the
randomized background distribution to calculate permuted p-values, which are corrected using
the Benjamini-Hochberg method to provide permuted q-values. A permuted q-value ≤ 0.1
denotes a significant final mutation frequency. The permuted q-value (PQ) filter reduced the
number of somatically mutated genes to 4,240 (**Figure 2A**). This filtering improved LoF somatic
mutations from three to four gold standards (TCGA consensus, CGC, Vogelstein, and
OncodriveROLE) with the somatic mutations that had the LoF predicted role from OncoMerge.
Still, the Act comparisons did not show significant enrichment (**Figure 2C**). The PQ filter had a
minimal impact on the number of genes per locus (**Figure 2B**). This lack of significant overlap
for Act somatic mutations demonstrates that further filtering is required.

A key consideration in developing OncoMerge was that integrating the somatic mutation types
should highlight the functional somatic mutations over passenger mutations. Therefore, we
created a filter to prioritize somatically mutated genes more likely to be functional. An average
CNA encompasses 3.8 ± 7.9 Mb of genomic sequence[23], and genomic segments of this size
typically include many genes. These large genomic regions make it difficult to determine which
of the affected genes are the functional gene(s) underlying the CNA locus without integrating
additional information. We assert that passenger genes underlying a CNA locus are considered
noise and can be identified by the lack of allelic heterogeneity. Thus, functional gene(s) can be
identified through allelic heterogeneity that boosts the somatic mutation frequency for a gene
above the background CNA frequency. We designed a low-pass filter that retains only the
gene(s) with the maximum final frequency (MFF). The MFF filter is only applied if a locus has
more than ten genes. Application of the MFF filter dramatically reduced the number of
somatically mutated genes from 6,028 to 1,459 (**Figure 2A**) and the number of genes per locus
(**Figure 2B**). The MFF filter also helps make the average contribution of PAMs and CNAs to the
final mutation frequency more even (**Figure S2**; **Table S5**). We additionally observed a marked
improvement in overlap with the gold standards. Significant enrichment was observed for four
Act gold standards with somatic mutations that OncoMerge predicts as Act. All five of the LoF
gold-standard versus OncoMerge predicted LoF comparisons (**Figure 2C**). Thus, the MFF filter
directly addresses the issue of too many genes in a CNA locus. Removing more than three-
quarters of the somatically mutated genes improves the overlaps with gold standards.

We then assessed the impact of applying both the PQ and MFF filters. Simultaneous application
of both filters reduced the number of somatically mutated genes beyond the MFF filter (1,145
genes; **Figure 2A**), and the number of genes per locus was further improved (**Figure 2B**).
There was also an improvement in the significant overlap with gold standards where all five LoF
gold-standard versus OncoMerge predicted LoF and significant overlap for four Act gold-

standard versus OncoMerge predicted Act (**Figure 2C**). Importantly, none of the gold standard
Act versus LoF or LoF versus Act comparisons were significant for any filter combination,
demonstrating that the OncoMerge predicted roles are consistent with prior knowledge.

Reducing biases due to microsatellite instability and hypermutation
Microsatellite instability (MSI) and hypermutation drastically increase the number of somatic
mutations in a tumor. The PQ and MFF filters and OncoMerge's core algorithm rely upon
somatic mutation frequency, which is susceptible to confounding by MSI or hypermutation.
Fortunately, all TCGA tumors used in this study are characterized for both MSI[5] and
hypermutation[6] status (**Figure 3A**). We observed a highly significant positive correlation
between MSI/hypermutation frequency and the total number of somatic mutations per cancer
after integration by OncoMerge (R = 0.68 and p-value = 2.0×10^{-5}). This strong positive
correlation demonstrates that MSI/hypermutation likely inflates the number of somatic mutations
discovered by OncoMerge. Therefore, we created the MSI and hypermutation censoring filter
(MHC) to exclude these tumors while OncoMerge determines which genes to include in the final
somatic mutation matrix. The mutation status for tumors with MSI and hypermutation are
included for genes in the final integrated mutation matrix. Applying the MHC filter alongside the
PQ and MFF filters reduced the overall number of somatically mutated genes (905 genes;
Figure 2A) and had minimal impact on the number of genes per locus (**Figure 2B**; **Data S1**).
The combined PQ, MFF, and MHC filters decreased the correlation between the
MSI/hypermutation frequency (R = 0.48 and p-value = 5.8×10^{-3}). All ten of the gold standard
Act vs. Act and LoF vs. LoF comparisons were significant. These results established that the
MHC filter is valuable for removing passenger mutations introduced by tumors with severely
increased somatic mutation rates. The PQ, MFF, and MHC filters comprise the default and final
OncoMerge filter set. The filters deal with known complications in cancer genetics and ensure
that the mutation roles in the integrated matrix are correctly assigned.

Sensitivity analyses to optimize filtering cutoffs and parameters
We then used sensitivity analyses of the filtering parameters to determine their optimal
parameterization for OncoMerge (**Figure 2D-G** and **Table S6**). First, we tested whether a
shallow (≥1) or deep (≥2) GISTIC threshold cutoff was optimal for integration purposes (**Figure
2D**). The shallow GISTIC threshold led to many more somatically mutated genes, but impaired
the discovery of meaningful activating mutations, as shown by the lack of significant overlap
with gold standard activating mutations. Therefore, the deep GISTIC threshold was chosen for
OncoMerge. Second, we varied the MFF gene number threshold of a CNA locus across the
values 1, 5, 10, 15, and 20 genes (**Figure 2E**). Significant overlap with all gold standard
activating mutations is observed up to 10 genes per loci threshold. These results demonstrate
that the optimal MFF cutoff is 10 genes per loci. Next, we tested the sensitivity of OncoMerge to
the minimum mutation frequency threshold by setting it with the values of 0.01, 0.05, 0.1, and
0.2 (**Figure 2F**). There was little impact on the significance of the gold-standard analysis across
the range of thresholds. However, the number of somatic mutations is significantly impacted by
this threshold. A 1% or smaller minimum mutation threshold would be warranted for somatic
mutation discovery. On the other hand, ensuring sufficient somatically mutated samples to
achieve statistical power for downstream analyses warrants a 5% minimum mutation threshold.
Finally, we tested the sensitivity of OncoMerge to the permuted q-value threshold from the PQ
filter across the values 0.01, 0.05, 0.1, and 0.2 (**Figure 2G**). The lowest threshold of 0.01 led to
a loss of significance for the overlap of the activating mutations for the Vogelstein et al., 2013
gold standard. The number of somatically mutated genes increased by hundreds of genes as

the permuted q-value threshold increased, and thus 0.05, 0.1, and 0.2 are all reasonable
threshold values. The permuted q-value threshold of 0.1 was chosen because it removed
integrated somatic mutations that could have happened by chance alone and retained many
somatically mutated genes. These sensitivity analyses provide a reasonable rationale for
choosing the values for the filtering cutoffs and parameters, which alternative values might be
used, and an idea of which contexts they might be useful.

Benefits of an integrated somatic mutation matrix
We evaluated the benefits of systematic somatic mutation integration by comparing OncoMerge
integrated somatic mutation matrices to those from PAMs. The PAM somatic mutation matrices
were used as a reference point because we have successfully used them as the sole source for
somatic mutations in previous studies[19,20]. We assessed the benefits of integration by tabulating
the number of somatic mutations and their roles (**Figure 3B**), the number of genes added by
integration (**Figure 3C**), and the increase in somatic mutation frequency due to integration
(**Figure 3E**). Act and LoF mutations represented the bulk of the somatic mutations in 30 cancers
(**Figure 3B**). The THCA and KICH were the only cancers that lacked Act or LoF mutations.
Consistent with Agrawal et al. 2014[24], THCA had only three mutations with a frequency ≥ 5%
BRAF, NRAS, and RET. On the other hand, KICH was under-sampled in the TCGA Pan-Cancer
atlas (n = 65), and LoF and Act mutations would likely be discovered with the inclusion of more
patient tumors.

We then investigated how many new genes the integration added for each cancer. Integration
added at least one somatically mutated gene for each cancer (**Figure 3C**) and more than sixty
somatically mutated genes for BLCA, LUAD, STAD, and UCEC (**Figure 3C**). The somatically
mutated genes added by OncoMerge make the integrated somatic mutation matrices more
comprehensive.

Next, we investigated the frequencies of the somatic mutations from the OncoMerge integrated
mutation matrices. The genes with the highest frequency map to well-known oncogenes (e.g.,
BRAF, KRAS, and EGFR) and tumor suppressors (e.g., APC, CDKN2A, and TP53; **Figure 3D**).
The APC gene was mutated in greater than eighty percent of tumors for READ. The TP53 gene
was mutated in greater than eighty percent of tumors for ESCA, LUSC, OV, READ, and UCS.
These frequently mutated genes in the OncoMerge integrated mutation matrices are consistent
with prior knowledge of somatic mutations for each cancer.

Finally, we calculated the frequency added through integration by subtracting the integrated
mutation frequency from the PAM frequency. The most substantial increases in somatic
mutation frequency were observed for TMPRSS2-ERG in PRAD and CDKN2A in ESCA, GBM,
and MESO (**Figure 3E**). Neither TMPRSS2-ERG nor CDKN2A would have been identified as
somatically mutated without incorporating fusions and CNAs, respectively. These findings
demonstrate that OncoMerge significantly improves the number and frequency of somatically
mutated genes in most cancers. Also, these results show that the systematic integration of
PAM, fusion, and CNA somatic mutations is crucial for obtaining a comprehensive mutation
matrix for each cancer.

Pan-cancer somatic mutations capture many known tumor suppressors and
oncogenes
Genes mutated in multiple cancers are of great interest as selective pressures have found a
common solution to influence cancer phenotypes in different contexts. Therefore, we searched

for genes somatically mutated in at least five cancers in the OncoMerge integrated mutation
matrices. The resulting gene list could be broken down into two groups of somatic mutations:
the LoF set (n = 23, **Figure 4A**) and the Act set (n = 13, **Figure 4B**). The genes FBXW7 and
KMT2C were somatically mutated with only PAMs. Both genes were previously classified as
tumor suppressors[25–27] and were grouped with the LoF set.

The pan-cancer somatically mutated genes harbored many well-known tumor suppressors and
oncogenes (**Figure 4C**). As expected, tumor suppressors[27] were significantly enriched in the
LoF group (overlap = 18, p-value = 5.0×10^{-20}), and oncogenes[28] were significantly enriched in
the Act group (overlap = 8, p-value = 1.6×10^{-10}). The top three most somatically mutated tumor
suppressors were TP53, PTEN, and CDKN2A. These three tumor suppressors control essential
checkpoints in the cell cycle, making them functionally interesting. The gene TP53 was
somatically mutated in 24 cancers, primarily by PAMs, but four LoFs were also observed for
GBM, LIHC, PRAD, and SARC. The top two most mutated oncogenes across cancers were
PIK3CA and KRAS, which become overactive kinases when mutated. Both PIK3CA and KRAS
have PAM and Act mutation roles across the different cancers, and only the NFE2L2 gene has
a similar mixture of PAM and Act mutation roles. The genes CASC8, CCND1, and TERT
included cancers with a CNAamp mutation role. The somatic mutation roles are all Act for the
remainder of the oncogenes. These pan-cancer analyses further validate the systematic
somatic mutation integration by OncoMerge through the unbiased recall of tumor suppressors
and oncogenes.

Improving gene regulatory network inference
Next, the integrated somatic mutation matrices for the TCGA cancer types were used to
construct gene regulatory networks (GRNs; **Table 7**) and compared to networks built using only
PAMs (legacy)[20]. The GRNs connected somatic mutations to TFs and miRNAs that regulate the
expression of a set of genes associated with cancer phenotypes or patient survival.

The average degree was the first metric we considered to compare the GRNs. The degree of a
node is the number of edges connecting it to other nodes. The average degree is a standard
network metric computed as the average of all node degrees in the network. We found that the
average degree was larger for 23 OncoMerge GRNs relative to legacy GRNs (**Figure 5A**). The
exceptions were GBM (average degree was equal) and COAD, KIRC, SKCM, and STAD
(legacy had a larger average degree). The COAD, SKCM, and STAD cancer types harbor more
MSI and hypermutation tumors (**Figure 3A**), and we observed a reduction in the number of
COAD and STAD mutations in the OncoMerge GRN relative to the legacy GRN (**Figure 5B**).
These results suggest that the MHC filter removed spurious associations. Thus, we have
increased the average degree for most networks and addressed a systematic bias in legacy
networks.

Next, we compared the number of mutations in each GRN predicted to modulate the activity of
regulators. The OncoMerge GRNs contained more somatic mutation nodes than the legacy
GRNs for all cancers but COAD and STAD, likely due to MSI and hypermutation (**Figure 5B**).
Then, we assessed the recall of somatic mutations previously associated with each cancer from
the DisGeNET database[29]. All but two OncoMerge GRNs recalled more previously associated
somatic mutations than the legacy GRNs (**Figure 5C**). The exceptions were UVM with the same
amount and COAD with fewer (**Figure 5C**). These results demonstrate that OncoMerge
integrated mutation matrices provide increased power to infer associations with somatic
mutations, especially previously associated with each cancer.

Finally, we considered the number of causal and mechanistic TFs in each GRN. The
OncoMerge GRNs contained more predicted TFs than legacy for 22 GRNs, the same number of
TF regulators for STAD, and fewer TFs for COAD, GBM, KIRC, THCA, and UCEC (**Figure 5D**).
We also assessed the recall of TFs previously associated with each cancer from the DisGeNET
database[29,30]. Twenty of the OncoMerge GRNs recalled more previously associated TFs than
legacy GRNs (**Figure 5E**). The COAD and UCEC GRNs had the same amount, and GBM,
KIRP, SKCM, and THCA had fewer, and KICH and UVM had no recall of previously associated
TFs in either GRN (**Figure 5E**). In summary, using OncoMerge integrated mutation matrices
constructs GRNs that are more extensive and biologically meaningful.

Comparing active and static TF regulatory network architectures

The interactions between TFs are important for generating the transcriptional state of a human
cell. The underlying architecture of TF regulatory networks, comprised of TFs and their
interactions, are typically explored by enumerating all three-node network motifs and computing
their enrichment or depletion into triad significance profiles (TSPs)[31]. Most studies of network
motif enrichment have relied upon unsigned interactions[31–36], which ignore whether the
interaction is activating or repressing. To facilitate comparisons, our first analysis of network
architecture uses unsigned TSPs to compare static and active TF regulatory networks. Static TF
regulatory networks were constructed using chromatin accessibility and DNA binding motifs for
41 cell types[32]. These TF regulatory networks are static because they do not incorporate gene
expression data in their construction. Active TF regulatory networks are derived from the
OncoMerge augmented SYGNAL pan-cancer GRNs, which were trained using patient tumor
transcriptional data and therefore are comprised of active TF regulatory interactions. We
calculated TSPs for twenty-five TF regulatory networks and the median TSP (**Figure 6A** & **B**;
Table S8). We excluded the cancer types DLBC, KICH, KIRP, OV, TGCT, and THYM because
they had too few inferred regulatory interactions (< 50 interactions). In addition, we recalculated
the TSPs for the static TF regulatory networks using a more recent version of the mfinder
algorithm (**Figure 6B**).

The median TSPs of the active and static TF regulatory networks were highly correlated (R =
0.75, p-value = 3.0 x 10^{-3}; **Figure 6B**). Demonstrating that the architecture of the active network
resembles the static network. However, the maximum enriched network motifs were different.
The regulated and regulating feedback motifs (motifs 108 and 46) were the most highly enriched
motifs from the static TF regulatory networks and were still enriched, although not as significant,
in the active networks. In contrast, the feed-forward loop (FFL, motif 38) is the most highly
enriched motif in the active TF regulatory networks. These two motifs are quite similar in
structure and differ only by a single edge. Feedback motifs and FFLs can be further broken
down into ten and eight signed network motifs that each have a unique functional output[37].
Thus, we can discover what functions are being selected for by evolution in general and the
microcosm of tumor biology by exploring the enrichment of signed network motifs.

Coherent feed-forward loops enriched in active TF regulatory networks

Incorporating the sign of the regulatory interactions (activating or repressing) splits the FFL motif
into eight signed network motifs classified as coherent (C1, C2, C3, C4) and incoherent (I1, I2,
I3, I4)[37]. Simulation studies have demonstrated that coherent FFLs lead to delays in target gene
expression, and incoherent FFLs accelerate target gene expression[37]. FFLs were significantly
enriched in active TF regulatory networks, which led us to question whether coherent,
incoherent, or both FFLs were enriched. In active GRNs, the sign of the correlation between the

TF regulator to TF target can be used to determine the sign of the interaction (R > 0 equates to
activation, R < 0 equates to repression). The four coherent FFLs were enriched in the active TF
regulatory networks (**Figure 6C**; **Table S9**), and incoherent FFLs were severely under-enriched
(Z << 0). In summary, coherent FFLs were enriched in our active TF regulatory networks,
suggesting that transcriptional delay mechanisms must provide a valuable function for TF
regulatory networks.

Coherent switch-like feedback motifs enriched in active TF regulatory networks
The regulated and regulating mutual feedback motifs have a two-node feedback loop at their
core. The double-positive and double-negative two-node mutual feedback loops act like
switches[38]. We tested the twenty signed regulated and regulating mutual feedback network
motif configurations for enrichment in TF regulatory networks. Three regulating and three
regulated signed mutual feedback motifs (**Figure 6C**; **Table S9**). These six enriched regulated
and regulating mutual feedback motifs had a common configuration. Firstly, all the network
motifs were coherent. Coherent regulated and regulating feedback loops have interaction signs
between the feedback loop that are either double-positive or double-negative. And the regulated
or regulating node interacts with the feedback loop nodes using the same sign for double-
positive feedback loops and the opposite sign for double-negative feedback loops. Thus, there
are three coherent configurations for both regulated and regulating mutual feedback motifs
making six total, coinciding with the six enriched configurations (**Figure 6C**; **Table S9**). The
enriched motifs containing a double-positive feedback loop had the same interactions with the
non-feedback loop node, both activating or repressing (**Figure 6C**). The enriched motif
containing a double-negative feedback loop had opposing interactions with the non-feedback
loop node, one activating and one repressing (**Figure 6C**). These enriched signed network
motifs are the configurations that function as molecular switches[39]. Again, evolution has
selected for coherent network motif configurations likely because of their function.

Discussion
We avoided overfitting while developing and optimizing parameters for OncoMerge in three
ways. First, we used five gold standards that use different methods for somatic mutation
discovery to avoid overfitting to one specific gold standard. This diversification approach was
successful because we observed variable enrichment scores across the gold standards.
Secondly, the sensitivity analyses we conducted over a plausible set of parameter values
demonstrated the robustness of OncoMerge to different parameterizations. This is important
because it shows that OncoMerge has not been parameterized into an anomalous overfit state.
Instead, the parameters were chosen based on carefully considered statistical choices and
trends in the data. Thirdly, we avoided overfitting to a specific cancer somatic mutation profile by
applying and assessing the performance of OncoMerge across thirty-two cancer types. The
ability of OncoMerge to be applied to a Pan-Cancer cohort with many different mutation profiles
strongly suggests that OncoMerge should be generalizable to new cancer cohorts. We
employed all three of these approaches to avoid overfitting and to ensure that OncoMerge could
be applied to new datasets without having to tune parameters.

Additionally, we provide sensitivity analyses that can guide users who want to change
OncoMerge parameters by observing how specific parameter values impact its performance.
For example, the minimum mutation frequency can be set to zero to conduct somatic mutation
discovery, providing a more comprehensive list of somatic mutations and their types. In this
study, we chose a five percent cutoff for the minimum mutation frequency to ensure there were

enough somatically mutated tumors to power downstream GRN inference. The sensitivity
analyses of can be used to guide the choice of OncoMerge parameters to achieve different
goals than the default parameterization.

The construction of active GRNs enabled the exploration of signed network motifs and led to the
discovery that specific signed network motif configurations are being enriched. The SYGNAL
GRN construction method identifies active gene regulatory interactions by discovering
interactions that are supported by gene expression data from patient tumors[19]. On the other
hand, prior networks were static maps of DNA binding sites constructed using digital genomic
footprinting and the similarity of the underlying sequence of the footprints for known DNA
binding motifs[32]. The active networks use a correlation-based method to determine TF
regulatory roles (activator or repressor) for the interactions, which is not possible using static
binding maps. Analyzing signed network motifs provides a leap forward in understanding how
the underlying architecture of GRNs functions in real-world biological systems. OncoMerge
integrated somatic mutations offer a more solid platform to infer active GRNs that can be used
to explore the functional architecture of TF regulatory networks.

We discovered that coherent regulated and regulating feedback and FFL network motifs were
enriched in cancer TF regulatory networks. We cannot say whether this enrichment of network
motifs will generalize to all active GRNs or if this is a cancer-specific phenomenon. In normal
organismal development, feedback motifs have been previously shown to be essential for cell
fate decision-making[40,41]. On the other hand, in tumor cells and other cells in the tumor
microenvironment, the enriched feedback motifs may be maintaining a cell fate, or the disease
could be coopting the circuit to drive tumor biology. Likewise, coherent FFL network motifs have
also been associated with enhanced drug resistance[42]. These coherent motifs are relevant for
normal and diseased cell biology, and evolution has specifically selected these motif
configurations because of their unique functional outputs.

Future improvements to the OncoMerge algorithm include a more quantitative integration
approach for the somatic mutations, a replacement for or an improved maximum final frequency
filter, aggregation across pathways, and a determination of whether other genomic features may
be integrated (ecDNA[43] or epigenomics[44]). Additionally, in future single-cell studies with both
transcriptome and genome information, it would be helpful to have an OncoMerge
implementation that integrates PAM, fusion, and CNA for every single cell. We envision
OncoMerge as a valuable tool in the somatic mutation characterization pipeline. We hope it will
facilitate multi-omic studies and lead to novel discoveries that can be translated into clinical
insights.

Limitations of the study
Currently, OncoMerge assumes that the somatic mutations will be either PAM, CNA, or gene
fusions, meaning it will miss somatic mutations such as ecDNA[43], epigenomics[44], etc. Somatic
mutations were seeded by PAMs or CNAs that were mutated more than expected by chance
alone, which may exclude mutations of lower frequency from being discovered. Future studies
could be used to come up with alternative methods of seeding somatic mutations. Using a five
percent cutoff for somatic mutation frequency means that lower-frequency mutations will be
overlooked. Setting the minimum mutation frequency cutoff to less than five percent would
provide a complete list of somatic mutations.

Acknowledgments

This work was supported by NIH-NINDS Award # 1R01NS123038-01, and 1R01NS119650-01. The authors also acknowledge Robert Schultz for assistance in preliminary studies, and the Cancer Genome Atlas Research Network for the TCGA Pan-Cancer Atlas multi-omic patient tumor profiles.

Author contributions

Conceptualization, C.L.P.; Methodology, S.S.S, S.F.W., and C.L.P.; Investigation, S.S.S and C.L.P; Visualization, S.S.S, S.F.W, E.M.L., S.A.O., and C.L.P.; Writing – Original Draft, C.L.P.; Writing – Review & Editing, S.S.S, S.F.W, E.M.L., S.A.O., and C.L.P.; Funding Acquisition, C.L.P.; Resources, C.L.P.; Supervision, C.L.P.

Declaration of interests

The authors declare no competing interests.

Figure legends

Figure 1. OncoMerge integrates PAMs, fusions, and CNAs into an integrated mutation matrix with the most suitable mutation type for each gene. The input data for OncoMerge includes the PAM, transcript fusion, and CNA matrices. OncoMerge then generates six matrices (PAM, Fusion, CNAamp, CNAdel, Act, and LoF) and uses mutational frequency and statistical filters to determine each gene's most suitable somatic mutation role.

Figure 2. Assessing the performance of OncoMerge filters and conducting sensitivity analyses. **A.** Impact of filter sets on the number of somatically mutated genes inferred by OncoMerge in at least one cancer. **B.** Impact of filter sets on the distribution of genes per CNA locus using the same set of filtering conditions (y-axis is distributed on a log scale). The dashed line indicates the ten genes per loci cutoff that invokes the MFF filter. **C.** Enrichment of the gold standard (GS) Act or LoF somatic mutations with OncoMerge (OM) Act or LoF somatic mutations for each filtering condition: no filters (None); PQ filter; MFF; combined PQ and MFF; and combined PQ, MFF, and MHC. Significant enrichments from **C** are highlighted in red and had p-value less than or equal to the Bonferroni corrected α level of 4.8×10^{-4} ($\alpha = 0.05$, number of tests = 105, Bonferroni corrected $\alpha = \alpha$ / number of tests = 0.05 / 105 = 4.8×10^{-4}). The orange arrowheads indicate OM Act vs. GS Act, and the green arrowheads indicate OM LoF vs. GS LoF. **D.** Comparing GISTIC thresholds: cutoff of one equates to shallow amplification and deletions, and cutoff of two equates to deep amplifications and deletions. **E.** Comparing the possible values for the MFF filter parameter maximum number of genes in the loci (Max. loci genes) with 1, 5, 10, 15, and 20 genes. **F.** Comparing the possible cutoff values of the minimum mutation frequency (Min. mut. freq.) with 1%, 5%, 10% and 20%. **G.** Comparing the possible cutoff values of the PQ filter permuted q-value (Perm. q-value) with 0.01, 0.05, 0.1 and 0.2. Significant enrichments from **D**-**G** are highlighted in red and had p-value less than or equal to the Bonferroni corrected α level of 2.4×10^{-3} ($\alpha = 0.05$, number of tests per parameters = 21, Bonferroni corrected $\alpha = \alpha$ / number of tests per parameter = 0.05 / 21 = 2.4×10^{-3}). The purple arrowheads indicate the final parameterization chosen for OncoMerge.

Figure 3. Summary of effect on number and frequency of somatic mutations after integrating mutation types. **A.** Frequency of hypermutation and MSI across cancers. **B.** Number and distribution of mutation types. **C.** Number of somatically mutated genes added because of integration. **D.** Integrated somatic mutation frequencies. **E.** Increases in somatic mutation frequency relative to PAM frequency after integration.

Figure 4. Pan-cancer somatic mutations with a consistent functional impact across at least five cancers. **A.** Pan-cancer somatic mutations from the loss of functions group. **B.** Pan-cancer somatic mutations from the activating group. **C.** Prior knowledge of tumor suppressor or oncogene status for each somatically mutated gene (black square indicates known tumor suppressor or oncogene activity).

Figure 5. Demonstrating improvements in downstream SYGNAL analysis by comparing GRNs constructed with an OncoMerge integrated somatic mutation matrix versus a legacy network using only PAMs. **A.** Average degree of nodes in the PanCaner SYGNAL networks. **B.** Mutations per cancer network. **C.** Mutations that overlap with genes previously associated with a specific cancer in DisGeNET. **D.** TFs per cancer network. **E.** TFs that overlap with genes previously associated with a specific cancer in DisGeNET.

Figure 6. The architecture of functional disease-specific TF regulatory networks from human
tumors. **A**. Active TF regulatory network construction pipeline: 1) TFs from all cancer regulatory
networks were identified, 2) A putative map of TF regulatory network interactions was
constructed, 3) TF $\rightarrow$ TF relationships were filtered using Pearson's correlations computed from
patient tumor data, and 4) compute the triad significance profiles using mfinder. **B**. Comparison
of active TF regulatory network based on SYGNAL GRNs (red) to the static TF regulatory
network based on ENCODE DNA binding and accessibility (blue, Neph et al., 2012). **C**.
FANMOD enrichment normalized Z-scores for the three most enriched motifs from the active TF
regulatory network after incorporating TF regulatory interaction roles (activation or repression).
The first row, titled Coherent motifs, is shaded when the motif configuration is coherent and
white when it is incoherent. Normalized Z-scores are reported for each cancer, and diagonal
dashed lines are inserted when no Z-score was returned. The network motif can be found at the
bottom of each column, colored with regulatory roles. C1, C2, C3, C4 = coherent FFLs. I1, I2,
I3, I4 = incoherent FFLs.

STAR⊡Methods

Resource availability

Lead contact
Requests for further information should be directed to the lead contact, Christopher Plaisier (plaisier@asu.edu).

Materials availability
This study did not generate new materials.

Data and code availability
This paper analyzes existing, publicly available data. All the datasets used as input for our study were deposited in Figshare (https://doi.org/10.6084/m9.figshare.21760964.v1) and they are publicly available as of the date of publication. New datasets generated from our studies were deposited in Figshare (https://doi.org/10.6084/m9.figshare.20238867.v1) and they are publicly available as of the data of publication.

The OncoMerge original code has been deposited at GitHub (https://github.com/plaisier-lab/OncoMerge) and is also accessible through Zenodo using the DOI https://doi.org/10.5281/zenodo.5519663.

Any additional information required to reanalyze the data reported in this paper is available from the lead contact upon request.

TCGA cancer abbreviations

Study Abbreviation	Study Name
ACC	Adrenocortical carcinoma
BLCA	Bladder Urothelial Carcinoma
LGG	Brain Lower Grade Glioma
BRCA	Breast invasive carcinoma
CESC	Cervical squamous cell carcinoma and endocervical adenocarcinoma
CHOL	Cholangiocarcinoma
COAD	Colon adenocarcinoma
ESCA	Esophageal carcinoma
GBM	Glioblastoma multiforme
HNSC	Head and Neck squamous cell carcinoma
KICH	Kidney Chromophobe
KIRC	Kidney renal clear cell carcinoma
KIRP	Kidney renal papillary cell carcinoma
LIHC	Liver hepatocellular carcinoma
LUAD	Lung adenocarcinoma

LUSC	Lung squamous cell carcinoma
DLBC	Lymphoid Neoplasm Diffuse Large B-cell Lymphoma
MESO	Mesothelioma
OV	Ovarian serous cystadenocarcinoma
PAAD	Pancreatic adenocarcinoma
PCPG	Pheochromocytoma and Paraganglioma
PRAD	Prostate adenocarcinoma
READ	Rectum adenocarcinoma
SARC	Sarcoma
SKCM	Skin Cutaneous Melanoma
STAD	Stomach adenocarcinoma
TGCT	Testicular Germ Cell Tumors
THYM	Thymoma
THCA	Thyroid carcinoma
UCS	Uterine Carcinosarcoma
UCEC	Uterine Corpus Endometrial Carcinoma
UVM	Uveal Melanoma

Method details

Clinical and molecular data from TCGA

These studies used standardized, normalized, batch corrected, and platform-corrected multi-omics data generated by the Pan-Cancer Atlas consortium for 11,080 participant tumors[20]. Complete multi-omic profiles were available for 9,584 patient tumors. TCGA aliquot barcodes flagged as "do not use" or excluded by pathology review from the Pan-Cancer Atlas Consortium were removed from the study. The overall survival (OS, OS.time) data used were obtained from Liu et al. 2018[45].

- Somatic protein affecting mutations (PAMs) in TCGA – Somatic PAMs were identified by the Multi-Center Mutation Calling in Multiple Cancer (MC3) project[1] and were downloaded from the ISB Cancer Gateway in the Cloud (ISB-CGC; https://isb-cgc.appspot.com/). PAMs were required to have a FILTER value of either: PASS, wga, or native_wga_mix. In addition, all PAMs needed to be protein-coding by requiring that Variant_Classification had one of the following values: Frame_Shift_Del, Frame_Shift_Ins, In_Frame_Del, In_Frame_Ins, Missense_Mutation, Nonsense_Mutation, Nonstop_Mutation, Splice_Site, or Translation_Start_Site. Additionally, mutation calls were required to be made by two or more mutation callers (NCALLERS > 1). When both normal tissue and blood were available, the blood was used as the germline reference.
- Statistical significance of PAMs in TCGA – The likelihood that a gene is somatically mutated by chance alone was determined using MutSig2CV[7] and downloaded for each

cancer from the Broad GDAC FIREHOSE (https://gdac.broadinstitute.org/). Genes with a
MutSig2CV False Discovery Rate (FDR) corrected p-value (q-value) less than or equal
to 0.1 were considered significantly mutated[7].

• Somatic transcript fusions in TCGA – The TumorFusions portal[2] provides a pan-cancer
analysis of tumor transcript fusions in the TCGA using the PRADA algorithm[8].
• Somatic copy number alterations (CNAs) in TCGA – Genomic regions that were
significantly amplified or deleted were identified using Genomic Identification of
Significant Targets in Cancer (GISTIC2.0)[9] and downloaded for each cancer from the
Broad GDAC FIREHOSE.

Somatic mutation data import and preprocessing

An essential first step in OncoMerge is loading up and binarizing the somatic mutation data
(**Figure S1**). The somatic mutation data comprised of four primary matrices: 1) PAMs, 2)
fusions, 3) CNA amplifications (CNAamps), and 4) CNA deletions (CNAdels) (**Figure 1**). In
addition, two derivative matrices Act and LoF are created by merging the PAM with the
CNAamps or CNAdels matrices, respectively (**Figure 1**). All files are formatted as comma-
separated values (CSV) files with genes as rows and patients as columns unless otherwise
noted.

• PAM matrix - The matrix values are [0 or 1]: zero indicates the gene is not mutated in a
patient tumor, and one indicates the gene is mutated in a patient tumor.
• Fusion matrix - The matrix values are [0 or 1]: zero indicates no gene fusion in a patient
tumor, and one indicates the gene fused to another genomic locus in a patient tumor.
• CNAamp and CNAdel matrices – The all_thresholded_by_genes.csv GISTIC output file
is used to populate the CNAamp and CNAdel matrices. The all_thresholeded_by_genes
matrix values range from -2 and have no positive bound, and the values indicate the
copy number relative to the background. A cutoff of greater than or equal to 2 was used
to identify deep amplifications and less than or equal to -2 for deep deletions. Only deep
amplifications or deletions were included in these studies due to heterogeneity of cell
types and tumor biopsy purity. Oncomerge allows this threshold to be modified through a
command line parameter ('-gt' or '--gistic-threshold').
○ CNAamp matrix – The matrix values are [0 or 1]: zero indicates a gene is not
amplified in a patient tumor, and one indicates the gene is amplified in a patient
tumor.
○ CNAdel matrix – The matrix values are [0 or 1]: zero indicates a gene is not
deleted in a patient tumor, and one indicates a gene is deleted in a patient tumor.
• Act matrix – The Act matrix is the bitwise OR combination of the PAM, Fusion, and
CNAamp matrices. The Act matrix has genes as rows and patients as columns. The
matrix values are [0 or 1]: zero indicates the gene is not mutated or amplified in a patient
tumor, and one indicates the gene is either mutated, fused, amplified, or some
combination in a patient tumor.
• LoF matrix – The LoF matrix is the bitwise OR combination of the PAM, Fusion, and
CNAdel matrices. The LoF matrix has genes as rows and patients as columns. The
matrix values are [0 or 1]: zero indicates the gene is not mutated or deleted in a patient
tumor, and one indicates the gene is either mutated, fused, deleted, or some
combination in a patient tumor.

Seeding OncoMerge with putative somatic mutations

OncoMerge focuses on likely causal somatic mutations by considering only somatic mutations
that were statistically shown to be mutated more often than expected by chance alone. Likely
causal somatic mutations are also required to have a mutation frequency greater than 5%, the
definition of a common mutation[46], as this ensures sufficient patient tumors will be mutated to
power downstream analyses. These statistically significant common mutations were used as
seeds for OncoMerge integration. PAMs used as seeds were identified with MutSig2CV q-
values less than or equal to 0.1[14] and a mutation frequency greater than 5%. Gene fusions used
as seeds were identified as significant in PRADA[2,8] and had a mutation frequency greater than
5%. CNAamps or CNAdels used as seeds were identified as significantly amplified or deleted
from the amplified genes (amp_genes) or deleted genes (del_genes) GISTIC output files with
residual q-values less than or equal to 0.05[47]. CNAs from sex chromosomes (X and Y) were
excluded. Genes from sex chromosomes can enter OncoMerge as seeds from PAMs or fusions.
These seed genes become the starting point of the OncoMerge integration. Subsequent steps
determine if Act or LoF merged mutation profiles or their component PAM, Fusion, CNAamp, or
CNAdel mutation roles are the most appropriate integration model for a gene.

Merging somatic mutations in OncoMerge
The mutation role for each seed gene is assigned based on the frequencies of the mutation
types for a gene from the original (PAM, Fusion, CNAamp, CNADel) and merged (Act and LoF)
somatic mutation matrices and statistical thresholds for PAM (MutSig2CV) and CNAs (GISTIC).
The function g is applied to each seed gene to choose the mutation role using the following
parameters: f_{MMF} the minimum mutation frequency (defaults to 5%), f_{Act} the frequency of the
merged Act mutations, f_{LoF} the frequency of the merged LoF mutations, f_{PAM} the frequency of
the PAM mutations, f_{Fusion} the frequency of the gene fusion mutations, f_{CNAamp} the frequency
of CNA amplification mutation, f_{CNAdel} the frequency of the CNA deletions mutations,
$qv_{MutSig2CV}$ significance of PAM mutations as MutSig2CV q-value, and qv_{GISTIC} significance of
CNA mutations as GISTIC residual q-value.

$$g\left(f_{MMF}, f_{Act}, f_{LoF}, f_{PAM}, f_{Fusion}, f_{CNAamp}, f_{CNAdel}, qv_{MutSig2CV}, qv_{GISTIC}\right)$$
$$= \begin{cases} Act, & if \left(f_{Act} > \max\left(f_{LoF}, f_{PAM}, f_{Fusion}, f_{CNAamp}, f_{CNAdel}\right)\right) and \left(f_{Act} \geq f_{MMF}\right) \\ LoF, & elif \left(f_{LoF} > \max\left(f_{Act}, f_{PAM}, f_{Fusion}, f_{CNAamp}, f_{CNAdel}\right)\right) and \left(f_{Lof} \geq f_{MMF}\right) \\ PAM, & elif \left(f_{PAM} \geq \max\left(f_{Act}, f_{LoF}, f_{Fusion}, f_{CNAamp}, f_{CNAdel}\right)\right) and \left(f_{PAM} \geq f_{MMF}\right) and \left(qv_{MutSig2CV} \leq 0.1\right) \\ Fusion, & elif \left(f_{Fusion} \geq \max\left(f_{Act}, f_{LoF}, f_{PAM}, f_{CNAamp}, f_{CNAdel}\right)\right) and \left(f_{Fusion} \geq f_{MMF}\right) \\ CNAamp, & elif \left(f_{CNAamp} \geq \max\left(f_{Act}, f_{LoF}, f_{PAM}, f_{Fusion}, f_{CNAdel}\right)\right) and \left(f_{CNAamp} \geq f_{MMF}\right) and \left(qv_{GISTIC} \leq 0.05\right) \\ CNAdel, & elif \left(f_{CNAdel} \geq \max\left(f_{Act}, f_{LoF}, f_{PAM}, f_{Fusion}, f_{CNAamp}\right)\right) and \left(f_{CNAdel} \geq f_{MMF}\right) and \left(qv_{GISTIC} \leq 0.05\right) \end{cases}$$

The mutation role for each seed gene is chosen using this decision tree based on mutational
frequencies and statistical significance. The Act and LoF are first in the decision tree because
merging mutation types should lead to a larger mutation frequency than any individual source
mutation frequency ($f_{LoF}, f_{PAM}, f_{Fusion}, f_{CNAamp}, f_{CNAdel}$). A strict inequality (greater than) is used so
that the Act or Lof is disregarded if it has the same mutation frequency as a source mutation
frequency. If an Act or LoF integrated mutation role is not chosen, then the source mutation with
the highest frequency is chosen. And in the case of ties the non-strict inequalities (greater than
or equal to) determine the order of preference for the tied mutational roles: PAM > Fusion >
CNAamp > CNAdel. This ordering ensures that integrated mutation roles are chosen when
possible and that the most frequent source mutation role is otherwise chosen. The PQ, MFF,

and MHC filters further modify the assigned gene mutation roles to determine the final gene
mutation role.

Permuted q-value (PQ) filter

672
For putative Act and LoF mutations, a permuted q-value is computed by randomizing the order
of rows in the PAM, Fusion, and CNA mutation matrices' and then calculating the randomized
frequency distribution for Acts and LoFs. The observed frequency for an Act or Lof mutation is
then compared to the randomized frequency distribution to compute the permuted p-value.
Permuted p-values are corrected into q-values using the multiple-test Benjamini-Hochberg FDR-
based correction method. Only Acts or LoFs that had a permuted q-value ≤ 0.1 were retained.
Any Act or LoF with a permuted q-value > 0.1 was set to the mutation role of either PAM,
Fusion, CNAamp, or CNAdel based on which mutation role had the highest frequency. This
modifies the function g into g_{PQ} that includes the permuted q-value as a new input variable
$qv_{permuted}$, and is included as a constraint for the calls of Act and LoF.

$$g_{PQ}\left(f_{MMF}, f_{Act}, f_{LoF}, f_{PAM}, f_{Fusion}, f_{CNAamp}, f_{CNAdel}, qv_{MutSig2CV}, qv_{GISTIC}, qv_{permuted}\right)$$

$$= \begin{cases} Act, & if\ \left(f_{Act} > \max\left(f_{LoF}, f_{PAM}, f_{Fusion}, f_{CNAamp}, f_{CNAdel}\right)\right)\ and\ \left(f_{Act} \geq f_{MMF}\right)\ and\ \left(qv_{permuted} \leq 0.1\right) \\ LoF, & elif\ \left(f_{LoF} > \max\left(f_{Act}, f_{PAM}, f_{Fusion}, f_{CNAamp}, f_{CNAdel}\right)\right)\ and\ \left(f_{Lof} \geq f_{MMF}\right)\ and\ \left(qv_{permuted} \leq 0.1\right) \\ PAM, & elif\ \left(f_{PAM} \geq \max\left(f_{Act}, f_{LoF}, f_{Fusion}, f_{CNAamp}, f_{CNAdel}\right)\right)\ and\ \left(f_{PAM} \geq f_{MMF}\right)\ and\ \left(qv_{MutSig2CV} \leq 0.1\right) \\ Fusion, & elif\ \left(f_{Fusion} \geq \max\left(f_{Act}, f_{LoF}, f_{PAM}, f_{CNAamp}, f_{CNAdel}\right)\right)\ and\ \left(f_{Fusion} \geq f_{MMF}\right) \\ CNAamp, & elif\ \left(f_{CNAamp} \geq \max\left(f_{Act}, f_{LoF}, f_{PAM}, f_{Fusion}, f_{CNAdel}\right)\right)\ and\ \left(f_{CNAamp} \geq f_{MMF}\right)\ and\ \left(qv_{GISTIC} \leq 0.05\right) \\ CNAdel, & elif\ \left(f_{CNAdel} \geq \max\left(f_{Act}, f_{LoF}, f_{PAM}, f_{Fusion}, f_{CNAamp}\right)\right)\ and\ \left(f_{CNAdel} \geq f_{MMF}\right)\ and\ \left(qv_{GISTIC} \leq 0.05\right) \end{cases}$$

683   The permuted q-value cutoff defaults to 0.1 and can be set to another value through a command
684   line parameter ('-pq', --perm_qv').

Maximum final frequency (MFF) filter

685
686   The maximum final frequency (MFF) filter is a low-pass genomic filter designed to remove
687   passenger genes from frequently mutated CNA loci that contain many underlying genes. By
688   default, the filter is applied when there are 10 or more genes in a locus. Let L_{MFF} be the set of
689   loci with greater than 10 genes, $locus$ be defined as the set of genes in a CNA locus, and len a
690   function that returns the number of genes in a set.

$$L_{MFF} = \{\ locus \mid locus\ in\ L, and\ len(locus) \geq 10\ \}$$

691   Let G_{locus} be the set of all n genes underlying a CNA locus.

$$G_{locus} = \{gene_1, gene_2, gene_3, \dots, gene_n\}$$

692   The first step of the filter defines the maximum mutation frequency (f_{MFF}) for the genes of $locus$.
693   This requires using two functions: $freq$ which returns the mutation frequency for a gene, and
694   max which returns the maximum value from a set.

$$f_{MFF} = \max\left(\{freq(gene_1), freq(gene_2), freq(gene_3), \dots, freq(gene_n)\}\right)$$

695   Only the gene(s) that have a mutation frequency equal to the f_{MFF} are retained for $locus_{MFF}$.

$$locus_{MFF} = \{\ gene \mid gene\ in\ G_{locus}, and\ freq(gene) == f_{MFF}\ \}$$

The genes from each $locus_{MFF}$ are included in the final mutation matrix. The number of genes
underlying a CNA locus can be set through a command line parameter ('-mlg', --
max_loci_genes').

Microsatellite hypermutation censoring (MHC) filter
The TCGA tumors used in this study have been characterized for both MSI[5] and hypermutation[6]
(**Table S1**). The tumors with MSI or hypermutation are loaded as a blocklist of patient IDs
through a command line parameter ('-bl' or '--blocklist'). All tumors in the blocklist are excluded
from consideration by the PQ and MFF filters while determining the genes to include in the final
somatic mutation matrix. The mutation status for blocklist tumors are included in the final
integrated mutation matrix.

Optimizing OncoMerge filtering parameters
Sensitivity analyses of filtering parameters (GISTIC threshold, maximum loci genes, minimum
mutation frequency, and permuted q-value cutoff) for the OncoMerge algorithm was conducted
by varying one input parameter while fixing all others. The number of somatically mutated genes
and enrichment of gold standards from each parameterization of OncoMerge was evaluated to
determine the optimal values for each input parameter.

OncoMerge outputs
OncoMerge provides four output files that provide valuable information about the integration
process and the final integrated mutation matrix that can be used in downstream studies. Here
is a brief description of each file and its contents:
• oncoMerge_mergedMuts.csv – The integrated mutation matrix is comprised of genes
717   (rows) by patient tumors (columns) of mutation status after integration by OncoMerge.
718   The matrix values are [0 or 1]: zero indicates that the gene is not mutated in a patient
719   tumor, and one indicates that the gene was mutated in a patient tumor.
720   • oncoMerge_CNA_loci.csv – A list of the genes mapping to each CNAamp or CNAdel
721   locus included in the OncoMerge integrated mutation matrix.
722   • oncoMerge_ActLofPermPV.csv – List of all significant Act and LoF genes, their
723   OncoMerge mutation role, frequency, empirical p-value, and empirical q-value. This
724   output is before the application of the low-pass frequency filter.
725   • oncoMerge_summaryMatrix.csv – Matrix of genes (rows) by all information gathered by
726   OncoMerge.

To aid in comparisons between runs, we provide the save permutation option ('-sp' or '--
save_permutation') to output permutation results so that the same permuted distribution can be
used with different parameters in separate runs. We also provide the load permutation option ('-
lp' or '--load_permutation') to load up the permuted distribution from a previous run. The
permuted distributions are saved in the following files if requested:
• oncomerge_ampPerm.npy, oncomerge_delPerm.npy – Snapshot of the non-
733   deterministic permutation results from combining PAM, Fusion, and CNAamp or PAM,
734   Fusion, and CNAdel frequencies, respectively.

735 **Gold standard cancer-specific gene role validation datasets**
736   Gold standard datasets are vital to validating the usefulness of each feature in OncoMerge. Two
737   different sources of gold standard cancer-specific gene role (Act or LoF) datasets were used to
738   validate the OncoMerge predicted tumor-specific gene roles:

739  •  TCGA consensus: The TCGA consensus is a list of driver genes identified from the
740    TCGA Pan-Cancer Atlas labeled with somatic mutation role (oncogene or tumor
741    suppressor) and cancer type. The TCGA consensus was constructed by Bailey et al.,
742    2018 wherein they catalog a list of 299 unique oncogenesis associated genes[6]. In the
743    TCGA consensus 280 cancer-specific oncogene roles were identified, and 417 cancer-
744    specific tumor suppressor roles were identified (**Table S2**).
745  •  Cancer Gene Census (CGC): The CGC from COSMIC is an expert-curated database of
746    human cancer driver genes labeled with somatic mutation role (oncogene and tumor
747    suppressor) and cancer type. The CGC was developed by Catalogue of Somatic
748    Mutations in Cancer (COSMIC) as an expert-curated database of human cancer-driving
749    genes[11]. CGC cancers were mapped to the TCGA cancers by manual curation (**Table
750    S2**). In the CGC 205 cancer-specific oncogene roles were identified, and 304 cancer-
751    specific tumor suppressor roles were identified (**Table S2**).

752 **Gold standard gene role validation datasets**
753 Three different sources of gold standard gene role (Act or LoF) datasets were used to validate
754 the OncoMerge predicted gene roles:
755  •  20/20 rule: The 20/20 rule defines oncogenes (Act) by requiring >20% of mutations in
756    recurrent positions, and tumor suppressors (LoF) as >20% of recorded mutations are
757    inactivating (missense or truncating)[3]. With the 20/20 rule, 54 oncogene roles were
758    identified, and 71 tumor suppressor roles were identified (**Table S2**).
759  •  OncodriveROLE: OncodriveROLE is a machine learning algorithm that classifies genes
760    according to their role (Act or LoF) based on well-curated genomic features[13]. With
761    OncodriveROLE, 76 oncogene (Act) roles were identified, and 109 tumor suppressor
762    (LoF) roles were identified (**Table S2**).
763  •  Tokheim Ensemble: Ensemble-based method from Tokheim et al., 2016[12], which
764    integrates MutSigCV, 20/20+, and TUSON methods for predicting gene roles (oncogene
765    and tumor suppressor). With the Tokheim Ensemble, 78 oncogene (Act) roles were
766    identified, and 212 tumor suppressor (LoF) roles were identified (**Table S2**).

Computing overlap between OncoMerge and gold standards
A hypergeometric enrichment statistic was used to compute the significance of overlap
observed between each gene role in OncoMerge versus the gold standards. When possible, the
tumor specificity of the gene role was taken into consideration (TCGA consensus and CGC). A
total of 105 hypergeometric enrichment tests were conducted for the comparison to gold
standards to test out different filters (5 combined gold standard tests + 5 gold standard datasets
* 5 filter conditions [None, PQ, MFF, PQ MFF, and PQ MFF MHC] * 4 GS & OM functional tests
[Act vs. Act, Act vs. LoF, LoF vs. Act, and LoF vs. LoF] = 105 tests). An α level of 0.05 was
chosen, and significant overlaps were determined as p-values less than or equal to the
Bonferroni multiple hypothesis corrected alpha level of 4.8 x 10^{-4} (α / number of tests = 0.05 /
105 = 4.8 x 10^{-4}). This cutoff ensures that the comparisons to the gold standards are not likely to
have occurred by chance alone, even though we conducted 105 independent tests.

For each sensitivity analysis we conducted 21 tests against the gold standards (1 combined
gold standard test + 5 gold standard datasets * 4 GS & OM functional tests [Act vs Act, Act vs
LoF, LoF vs Act, and LoF vs LoF] = 21 tests). An α level of 0.05 was chosen, and significant
overlaps for sensitivity analyses across the potential parameter values for OncoMerge were
determined as p-values less than or equal to the Bonferroni multiple hypothesis corrected alpha

level of 2.4 x 10^{-3} (α / number of tests = 0.05 / 21 = 2.4 x 10^{-3}). This cutoff addresses the impact
of the 21 independent tests for each parameter value in the sensitivity analysis.

Availability of OncoMerge
We provide the Oncomerge software and data in several standard distribution formats to
facilitate future studies that aim to integrate somatic mutations. The source code for OncoMerge
is available on GitHub (https://github.com/plaisier-lab/OncoMerge). Finally, an OncoMerge
Docker image was created that can be run as a virtual machine with all dependencies pre-
installed (https://hub.docker.com/r/cplaisier/oncomerge). Detailed documentation is provided,
along with a tutorial that describes the use of OncoMerge. The goal of disseminating
OncoMerge in these ways is to give end-users flexibility to choose what distribution method best
fits their computational platform.

OncoMerge TCGA Pan-Cancer Atlas input and output files
We also provide the Pan-Cancer Atlas TCGA somatic mutation data used as input for
OncoMerge (https://doi.org/10.6084/m9.figshare.21760964.v1). And the resulting OncoMerge
integrated somatic mutation matrices for those planning studies that use somatic mutations from
the TCGA Pan-Cancer Atlas (https://doi.org/10.6084/m9.figshare.20238867). These integrated
somatic mutation matrices can be used for any downstream analyses incorporating somatic
mutations and will provide the same power boost observed in our studies. In addition, we also
offer the pan-cancer SYGNAL GRNs and TF regulatory networks as supplementary tables
(**Tables S7**–**S9**) to expedite systems genetics studies of TCGA cancers.

TCGA Pan-Cancer SYstems Genetics Network AnaLysis (SYGNAL)
The mRNA and miRNA expression data required to run SYGNAL were obtained from Thorsson
et al., 2018[20]. The SYGNAL pipeline is composed of 4 steps and command-line parameters for
all programs are described in detail in Plaisier et al., 2016[19]. Each cancer was run separately
through the pipeline to reduce the confounding from tissue of origin differences. Highly
expressed genes were discovered for each cancer by requiring that genes have greater than or
equal to the median expression of all genes across all conditions in ≥ 50% of patients[19]. These
gene sets were then used as input to SYGNAL to construct the gene regulatory networks
(GRNs) for each cancer.

The underlying cMonkey2 biclustering results are identical to those from Thorsson et al., 2018[20]
as they do not rely upon genetic information. All immune-specific filters were removed for these
analyses, and all bicluster filtering was done as described in Plaisier et al., 2016[19]. Using
Network Edge Orienting (NEO)[48] somatic mutations are integrated with bicluster and regulator
expression in the next step. Two networks were constructed by applying systems genetics
analysis with NEO to the biclusters: 1) GRNs were inferred using PAM-only somatic mutation
matrices as a baseline; 2) GRNs were inferred using OncoMerge integrated somatic mutation
matrices. Importantly, the PAM-only somatic mutation matrices used were the same ones used
as input for OncoMerge.

TF regulatory network construction for PanCan-SYGNAL networks
A TF regulatory network was built for each cancer in three steps (**Figure 6A**). First, the TFs
regulating survival-associated biclusters were extracted from each cancer's SYGNAL GRN.
Second, a preliminary $TF_{regulator} \rightarrow TF_{target}$ regulatory network was constructed based on the
presence of a binding site for a putative $TF_{regulator}$ in the promoter of a TF_{target} from the
Transcription Factor Target Gene Database[19] (http://tfbsdb.systemsbiology.net). TF family

expansion[19] was used to supplement TFs that did not have an experimentally determined DNA
recognition motif in the database. The assumption was that the motifs within a TF family would
not vary significantly. Therefore, TF family members from the TFClass database[49] with a known
DNA recognition motif can be used as a proxy for a TF with no known DNA recognition motif.
Finally, the putative $TF_{regulator} \rightarrow TF_{target}$ regulatory network was filtered by requiring a significant
Pearson correlation between the mRNA expression of the $TF_{regulator}$ and TF_{target} (Pearson's $|R| \geq$
0.3 and p-value $\leq$ 0.05; **Figure 6A**; **Table S9**). The sign of the correlation coefficient can be
used to determine the role of a regulatory interaction: a positive correlation coefficient equates
to the $TF_{regulator}$ being an activator, and a negative correlation coefficient equates to the $TF_{regulator}$
being a repressor. Networks with fewer than 50 interactions were not included in the analyses
as they were not sufficiently powered to run the network motif analysis. The cancer regulatory
networks for DLBC, KICH, KIRP, OV, TGCT, and THYM were excluded from further studies.
840
TF regulatory network motif analysis
Three-node network motifs were enumerated from the TF regulatory networks using mfinder[50] in
the same manner as Neph et al., 2012[32] and used to compute triad significance profiles
(TSPs)[31]. The parameters used with mfinder v1.20 were[32]: motif size set at 3 (-s 3), requested
250 random networks to be generated (-r 250), and the Z-score threshold was set at -2000 to
ensure all motifs are reported (-z -2000). All Z-scores were extracted for each cancer and
converted to triad significance profiles using the methods of Milo et al., 2004[31].
848
849 For consistency, the TF regulatory networks for the 41 different cell types from Neph et al., 2012
850 were downloaded from http://www.regulatorynetworks.org/ and analyzed using the same
851 approach described above.
852
Signed network motif analysis incorporating TF regulator interaction roles
The enrichment of signed feed-forward loops (FFLs), regulated feedback, and regulating
feedback network motifs was computed using FANMOD[51], which takes into consideration TF
regulatory roles (activation and repression). The command line version of FANMOD from
IndeCut[52] was used with default parameters, except for the inclusion of regulatory role (colored
edges)[51] (fanmod 3 100000 1 <input_file> 1 0 1 2 0 1 0 1000 3 3 <output_file> 1 1). Z-scores for
signed FFLs, regulated feedback, and regulating feedback network motifs were extracted for
each cancer and converted to triad significance profiles using the methods of Milo et al., 2004[31].
The signed FFL network motifs are broken down into C1, C2, C3, C4, I1, I2, I3, and I4, as
described previously[37].
863
Quantification and statistical analysis

A nominal alpha value (p-value or q-vaule cutoff) of 0.05 was used unless otherwise stated.
Statistical analyses are described in detail in the methods sections where they were used, and
we provide a brief synopsis of the statistical methods below. Hypergeometric enrichment
analysis was used to identify significant overlaps of OncoMerge-derived gene sets with gold-
standards gene sets. When appropriate for the gold-standard analyses, Benjamini-Hochberg
FDR multiple hypothesis correction was applied to the hypergeometric p-values. Cutoff were as
described in the methods or results. Permuted p-values were computed for each integrated
somatic mutation and Benjamini-Hochberg FDR multiple hypothesis correction was applied to
generate permuted q-values. Pearson correlations were used to compare two sets of

quantitative values (e.g., number of somatic mutations and MSI/hypermutation frequency) and
the correlation coefficient (R) and p-value are reported. Triad significance profiles (TSPs) were
used to quantify the enrichment of three node network motifs.
877

Supplementary tables

Table S1. Blocklist of MSI and hypermutator phentoype tumors. Related to **Figure 2**.

Table S2. Gold standard datasets. Related to **Figure 2**.

Table S3. Gold standard enrichment analysis with OncoMerge predicted somatic mutations.
Related to **Figure 2**.

Table S4. Gold standard enrichment analysis with OncoMerge predicted somatic mutations
using randomized somatic mutation matrices to show that overlap is not due only to seed
genes. Related to **Figure 2**.

Table S5. Relative mean contribution of each mutation type to final mutation frequency. Related
to **Figure 2**.

Table S6. Gold standard enrichment analysis from OncoMerge sensitivity analyses for cutoffs
and parameters. Related to **Figure 3**.

Table S7. Mechanistic and causal regulatory interactions from the TCGA Pan-Cancer SYGNAL
network. Related to **Figure 5**.

Table S8. Active TF regulatory network from TCGA PanCancer SYGNAL network. Related to
Figure 6.

Table S9. Active TF regulatory network motifs. Related to **Figure 6**.

Supplementary data

Data S1. OncoMerge summaries for all cancers. Related to **Figure 3**.

References

1. Ellrott, K., Bailey, M.H., Saksena, G., Covington, K.R., Kandoth, C., Stewart, C., Hess, J., Ma, S., Chiotti, K.E., McLellan, M., et al. (2018). Scalable Open Science Approach for Mutation Calling of Tumor Exomes Using Multiple Genomic Pipelines. Cell Syst *6*, 271-281.e7. 10.1016/j.cels.2018.03.002.

2. Hu, X., Wang, Q., Tang, M., Barthel, F., Amin, S., Yoshihara, K., Lang, F.M., Martinez-Ledesma, E., Lee, S.H., Zheng, S., et al. (2018). TumorFusions: an integrative resource for cancer-associated transcript fusions. Nucleic Acids Res *46*, D1144–D1149. 10.1093/nar/gkx1018.

3. Vogelstein, B., Papadopoulos, N., Velculescu, V.E., Zhou, S., Diaz, L.A., and Kinzler, K.W. (2013). Cancer genome landscapes. Science *339*, 1546–1558. 10.1126/science.1235122.

4. Beroukhim, R., Mermel, C.H., Porter, D., Wei, G., Raychaudhuri, S., Donovan, J., Barretina, J., Boehm, J.S., Dobson, J., Urashima, M., et al. (2010). The landscape of somatic copy-number alteration across human cancers. Nature *463*, 899–905. 10.1038/nature08822.

5. Bonneville, R., Krook, M.A., Kautto, E.A., Miya, J., Wing, M.R., Chen, H.-Z., Reeser, J.W., Yu, L., and Roychowdhury, S. (2017). Landscape of Microsatellite Instability Across 39 Cancer Types. JCO Precis Oncol *2017*. 10.1200/PO.17.00073.

6. Bailey, M.H., Tokheim, C., Porta-Pardo, E., Sengupta, S., Bertrand, D., Weerasinghe, A., Colaprico, A., Wendl, M.C., Kim, J., Reardon, B., et al. (2018). Comprehensive Characterization of Cancer Driver Genes and Mutations. Cell *173*, 371-385.e18. 10.1016/j.cell.2018.02.060.

7. Lawrence, M.S., Stojanov, P., Polak, P., Kryukov, G.V., Cibulskis, K., Sivachenko, A., Carter, S.L., Stewart, C., Mermel, C.H., Roberts, S.A., et al. (2013). Mutational heterogeneity in cancer and the search for new cancer-associated genes. Nature *499*, 214–218. 10.1038/nature12213.

8. Torres-García, W., Zheng, S., Sivachenko, A., Vegesna, R., Wang, Q., Yao, R., Berger, M.F., Weinstein, J.N., Getz, G., and Verhaak, R.G.W. (2014). PRADA: pipeline for RNA sequencing data analysis. Bioinformatics *30*, 2224–2226. 10.1093/bioinformatics/btu169.

9. Mermel, C.H., Schumacher, S.E., Hill, B., Meyerson, M.L., Beroukhim, R., and Getz, G. (2011). GISTIC2.0 facilitates sensitive and confident localization of the targets of focal somatic copy-number alteration in human cancers. Genome Biol *12*, R41. 10.1186/gb-2011-12-4-r41.

10. ICGC/TCGA Pan-Cancer Analysis of Whole Genomes Consortium (2020). Pan-cancer analysis of whole genomes. Nature *578*, 82–93. 10.1038/s41586-020-1969-6.

11. Sondka, Z., Bamford, S., Cole, C.G., Ward, S.A., Dunham, I., and Forbes, S.A. (2018). The COSMIC Cancer Gene Census: describing genetic dysfunction across all human cancers. Nat Rev Cancer *18*, 696–705. 10.1038/s41568-018-0060-1.

12. Tokheim, C.J., Papadopoulos, N., Kinzler, K.W., Vogelstein, B., and Karchin, R. (2016). Evaluating the evaluation of cancer driver genes. Proc Natl Acad Sci U S A *113*, 14330–14335. 10.1073/pnas.1616440113.

13. Schroeder, M.P., Rubio-Perez, C., Tamborero, D., Gonzalez-Perez, A., and Lopez-
Bigas, N. (2014). OncodriveROLE classifies cancer driver genes in loss of function and
activating mode of action. Bioinformatics *30*, i549-555. 10.1093/bioinformatics/btu467.

14. Lawrence, M.S., Stojanov, P., Mermel, C.H., Robinson, J.T., Garraway, L.A., Golub,
T.R., Meyerson, M., Gabriel, S.B., Lander, E.S., and Getz, G. (2014). Discovery and
saturation analysis of cancer genes across 21 tumour types. Nature *505*, 495–501.
10.1038/nature12912.

15. Davoli, T., Xu, A.W., Mengwasser, K.E., Sack, L.M., Yoon, J.C., Park, P.J., and Elledge,
S.J. (2013). Cumulative haploinsufficiency and triplosensitivity drive aneuploidy patterns and
shape the cancer genome. Cell *155*, 948–962. 10.1016/j.cell.2013.10.011.

16. Tamborero, D., Gonzalez-Perez, A., and Lopez-Bigas, N. (2013). OncodriveCLUST:
exploiting the positional clustering of somatic mutations to identify cancer genes.
Bioinformatics *29*, 2238–2244. 10.1093/bioinformatics/btt395.

17. Gonzalez-Perez, A., Perez-Llamas, C., Deu-Pons, J., Tamborero, D., Schroeder, M.P.,
Jene-Sanz, A., Santos, A., and Lopez-Bigas, N. (2013). IntOGen-mutations identifies cancer
drivers across tumor types. Nat. Methods *10*, 1081–1082. 10.1038/nmeth.2642.

18. Cerami, E., Gao, J., Dogrusoz, U., Gross, B.E., Sumer, S.O., Aksoy, B.A., Jacobsen, A.,
Byrne, C.J., Heuer, M.L., Larsson, E., et al. (2012). The cBio cancer genomics portal: an
open platform for exploring multidimensional cancer genomics data. Cancer Discov *2*, 401–
404. 10.1158/2159-8290.CD-12-0095.

19. Plaisier, C.L., O'Brien, S., Bernard, B., Reynolds, S., Simon, Z., Toledo, C.M., Ding, Y.,
Reiss, D.J., Paddison, P.J., and Baliga, N.S. (2016). Causal Mechanistic Regulatory Network
for Glioblastoma Deciphered Using Systems Genetics Network Analysis. Cell Syst *3*, 172–
186. 10.1016/j.cels.2016.06.006.

20. Thorsson, V., Gibbs, D.L., Brown, S.D., Wolf, D., Bortone, D.S., Ou Yang, T.-H., Porta-
Pardo, E., Gao, G.F., Plaisier, C.L., Eddy, J.A., et al. (2018). The Immune Landscape of
Cancer. Immunity *48*, 812-830.e14. 10.1016/j.immuni.2018.03.023.

21. Hanahan, D. (2022). Hallmarks of Cancer: New Dimensions. Cancer Discov *12*, 31–46.
10.1158/2159-8290.CD-21-1059.

22. Hanahan, D., and Weinberg, R.A. (2011). Hallmarks of cancer: the next generation. Cell
144, 646–674. 10.1016/j.cell.2011.02.013.

23. Harbers, L., Agostini, F., Nicos, M., Poddighe, D., Bienko, M., and Crosetto, N. (2021).
Somatic Copy Number Alterations in Human Cancers: An Analysis of Publicly Available Data
From The Cancer Genome Atlas. Front Oncol *11*, 700568. 10.3389/fonc.2021.700568.

24. Cancer Genome Atlas Research Network (2014). Integrated genomic characterization of
papillary thyroid carcinoma. Cell *159*, 676–690. 10.1016/j.cell.2014.09.050.

25. Gala, K., Li, Q., Sinha, A., Razavi, P., Dorso, M., Sanchez-Vega, F., Chung, Y.R.,
Hendrickson, R., Hsieh, J.J., Berger, M., et al. (2018). KMT2C mediates the estrogen

dependence of breast cancer through regulation of ERα enhancer function. Oncogene *37*,
4692–4710. 10.1038/s41388-018-0273-5.

26.   Hillman, R.T., Celestino, J., Terranova, C., Beird, H.C., Gumbs, C., Little, L., Nguyen, T.,
Thornton, R., Tippen, S., Zhang, J., et al. (2018). KMT2D/MLL2 inactivation is associated
with recurrence in adult-type granulosa cell tumors of the ovary. Nat Commun *9*, 2496.
10.1038/s41467-018-04950-x.

27.   Zhao, M., Kim, P., Mitra, R., Zhao, J., and Zhao, Z. (2016). TSGene 2.0: an updated
literature-based knowledgebase for tumor suppressor genes. Nucleic Acids Res *44*, D1023-
1031. 10.1093/nar/gkv1268.

28.   Liu, Y., Sun, J., and Zhao, M. (2017). ONGene: A literature-based database for human
oncogenes. J Genet Genomics *44*, 119–121. 10.1016/j.jgg.2016.12.004.

29.   Piñero, J., Ramírez-Anguita, J.M., Saüch-Pitarch, J., Ronzano, F., Centeno, E., Sanz,
F., and Furlong, L.I. (2020). The DisGeNET knowledge platform for disease genomics: 2019
update. Nucleic Acids Res. *48*, D845–D855. 10.1093/nar/gkz1021.

30.   Lambert, S.A., Jolma, A., Campitelli, L.F., Das, P.K., Yin, Y., Albu, M., Chen, X., Taipale,
J., Hughes, T.R., and Weirauch, M.T. (2018). The Human Transcription Factors. Cell *172*,
650–665. 10.1016/j.cell.2018.01.029.

31.   Milo, R., Itzkovitz, S., Kashtan, N., Levitt, R., Shen-Orr, S., Ayzenshtat, I., Sheffer, M.,
and Alon, U. (2004). Superfamilies of evolved and designed networks. Science *303*, 1538–
1542. 10.1126/science.1089167.

32.   Neph, S., Stergachis, A.B., Reynolds, A., Sandstrom, R., Borenstein, E., and
Stamatoyannopoulos, J.A. (2012). Circuitry and dynamics of human transcription factor
regulatory networks. Cell *150*, 1274–1286. 10.1016/j.cell.2012.04.040.

33.   Li, Y., Shao, T., Jiang, C., Bai, J., Wang, Z., Zhang, J., Zhang, L., Zhao, Z., Xu, J., and
Li, X. (2015). Construction and analysis of dynamic transcription factor regulatory networks in
the progression of glioma. Sci Rep *5*, 15953. 10.1038/srep15953.

34.   Stergachis, A.B., Neph, S., Sandstrom, R., Haugen, E., Reynolds, A.P., Zhang, M.,
Byron, R., Canfield, T., Stelhing-Sun, S., Lee, K., et al. (2014). Conservation of trans-acting
circuitry during mammalian regulatory evolution. Nature *515*, 365–370. 10.1038/nature13972.

35.   Gerstein, M.B., Kundaje, A., Hariharan, M., Landt, S.G., Yan, K.-K., Cheng, C., Mu, X.J.,
Khurana, E., Rozowsky, J., Alexander, R., et al. (2012). Architecture of the human regulatory
network derived from ENCODE data. Nature *489*, 91–100. 10.1038/nature11245.

36.   Boyle, A.P., Araya, C.L., Brdlik, C., Cayting, P., Cheng, C., Cheng, Y., Gardner, K.,
Hillier, L.W., Janette, J., Jiang, L., et al. (2014). Comparative analysis of regulatory
information and circuits across distant species. Nature *512*, 453–456. 10.1038/nature13668.

37.   Mangan, S., and Alon, U. (2003). Structure and function of the feed-forward loop
network motif. Proc Natl Acad Sci U S A *100*, 11980–11985. 10.1073/pnas.2133841100.

38.   Alon, U. (2007). Network motifs: theory and experimental approaches. Nat Rev Genet *8*,
1014   450–461. 10.1038/nrg2102.

1015   39.   Gardner, T.S., Cantor, C.R., and Collins, J.J. (2000). Construction of a genetic toggle
1016   switch in Escherichia coli. Nature *403*, 339–342. 10.1038/35002131.

1017   40.   McCauley, B.S., Weideman, E.P., and Hinman, V.F. (2010). A conserved gene
1018   regulatory network subcircuit drives different developmental fates in the vegetal pole of highly
1019   divergent echinoderm embryos. Dev Biol *340*, 200–208. 10.1016/j.ydbio.2009.11.020.

1020   41.   Davidson, E.H., Rast, J.P., Oliveri, P., Ransick, A., Calestani, C., Yuh, C.-H., Minokawa,
1021   T., Amore, G., Hinman, V., Arenas-Mena, C., et al. (2002). A genomic regulatory network for
1022   development. Science *295*, 1669–1678. 10.1126/science.1069883.

1023   42.   Charlebois, D.A., Balázsi, G., and Kærn, M. (2014). Coherent feedforward transcriptional
1024   regulatory motifs enhance drug resistance. Phys Rev E Stat Nonlin Soft Matter Phys *89*,
1025   052708. 10.1103/PhysRevE.89.052708.

1026   43.   Kim, H., Nguyen, N.-P., Turner, K., Wu, S., Gujar, A.D., Luebeck, J., Liu, J., Deshpande,
1027   V., Rajkumar, U., Namburi, S., et al. (2020). Extrachromosomal DNA is associated with
1028   oncogene amplification and poor outcome across multiple cancers. Nat Genet *52*, 891–897.
1029   10.1038/s41588-020-0678-2.

1030   44.   Saghafinia, S., Mina, M., Riggi, N., Hanahan, D., and Ciriello, G. (2018). Pan-Cancer
1031   Landscape of Aberrant DNA Methylation across Human Tumors. Cell Rep *25*, 1066-1080.e8.
1032   10.1016/j.celrep.2018.09.082.

1033   45.   Liu, J., Lichtenberg, T., Hoadley, K.A., Poisson, L.M., Lazar, A.J., Cherniack, A.D.,
1034   Kovatich, A.J., Benz, C.C., Levine, D.A., Lee, A.V., et al. (2018). An Integrated TCGA Pan-
1035   Cancer Clinical Data Resource to Drive High-Quality Survival Outcome Analytics. Cell *173*,
1036   400-416.e11. 10.1016/j.cell.2018.02.052.

1037   46.   Gibson, G. (2012). Rare and common variants: twenty arguments. Nat Rev Genet *13*,
1038   135–145. 10.1038/nrg3118.

1039   47.   Camacho, N., Van Loo, P., Edwards, S., Kay, J.D., Matthews, L., Haase, K., Clark, J.,
1040   Dennis, N., Thomas, S., Kremeyer, B., et al. (2017). Appraising the relevance of DNA copy
1041   number loss and gain in prostate cancer using whole genome DNA sequence data. PLoS
1042   Genet *13*, e1007001. 10.1371/journal.pgen.1007001.

1043   48.   Aten, J.E., Fuller, T.F., Lusis, A.J., and Horvath, S. (2008). Using genetic markers to
1044   orient the edges in quantitative trait networks: the NEO software. BMC Syst Biol *2*, 34.
1045   10.1186/1752-0509-2-34.

1046   49.   Wingender, E., Schoeps, T., and Dönitz, J. (2013). TFClass: an expandable hierarchical
1047   classification of human transcription factors. Nucleic Acids Res. *41*, D165-170.
1048   10.1093/nar/gks1123.

1049   50.   Milo, R., Shen-Orr, S., Itzkovitz, S., Kashtan, N., Chklovskii, D., and Alon, U. (2002).
1050   Network motifs: simple building blocks of complex networks. Science *298*, 824–827.
1051   10.1126/science.298.5594.824.

1052   51.   Wernicke, S., and Rasche, F. (2006). FANMOD: a tool for fast network motif detection.
1053   Bioinformatics *22*, 1152–1153. 10.1093/bioinformatics/btl038.

1054   52.   Ansariola, M., Megraw, M., and Koslicki, D. (2018). IndeCut evaluates performance of
1055   network motif discovery algorithms. Bioinformatics *34*, 1514–1521.
1056   10.1093/bioinformatics/btx798.