Systematic integration of protein affecting mutations, gene fusions, and copy number alterations into a comprehensive somatic mutational profile

Shawn S Striker [1], Sierra F Wilferd [1], Erika M Lewis [1], Samantha A O'Connor [1], Christopher L Plaisier [1]*

[1] School of Biological and Health Systems Engineering, Arizona State University, Tempe, AZ

*To whom correspondence should be addressed: Christopher Plaisier, E-mail: plaisier@asu.edu, Phone: (480) 965-6832, Address: P.O. Box 879709, Tempe, AZ 87287-9709

Abstract

A gene can be mutated across a tumor cohort by protein affecting mutations (PAMs), gene fusions, or copy number alterations (CNAs). These mutations can have a similar phenotypic effect (i.e., allelic heterogeneity) and should be integrated into a unified gene mutation profile. We provide OncoMerge as a somatic mutation integration platform that tames allelic heterogeneity, discovers causal mutations, integrates binary PAM and fusion with quantitative CNA data types, and overcomes known obstacles in cancer genetics. OncoMerge was applied to the 9,584 patient tumors from 32 cancers profiled by the TCGA Pan-Cancer Atlas to validate the novel integration methods. Integration increased the number and frequency of somatically mutated genes and improved the prediction of the somatic mutation role as either activating or loss of function. Using OncoMerge integrated somatic mutations boosts the power to infer active gene regulatory networks that increase the connectedness of the networks and incorporate more somatic mutations and regulators associated with cancer biology. We extracted transcription factor (TF) regulatory networks and found that they were enriched with feedback and feed-forward loop network motifs. Subsequent, signed network motif analysis demonstrated that switch-like feedback motifs and delay-inducing feed-forward loops were the only enriched configurations. This enrichment pattern suggests that evolution in general or in the tumor microenvironment is selecting for these particular functions. The OncoMerge integrated somatic mutations provide a more comprehensive platform for studies linking somatic mutations to downstream cancer phenotypes and will lead to novel biological insights in clinical samples.

Keywords: somatic mutations, allelic heterogeneity, cancer, oncogene, tumor suppressor gene

Introduction

The selective pressures driving the accrual of somatic mutations that affect cancer phenotypes are shared across cancers. This phenomenon leads to genes being somatically mutated across multiple cancers, e.g., oncogenes and tumor suppressors. The three main types of somatic mutations that modify the function of a gene or render it non-functional are: 1) protein affecting mutations (PAMs), 2) gene fusions, and 3) copy number alterations (CNAs). A PAM is a point mutation, short insertion, or short deletion inside a gene's coding region or splice sites[1]. Gene fusions occur when genomic rearrangements join two genes into a novel chimeric gene or place a promoter in front of a new gene, causing misexpression[2]. Finally, CNAs occur frequently in tumors where whole chromosomes, chromosomal arms, or localized genomic segments are duplicated or deleted[3,4]. Somatic mutation via PAM, gene fusion, or CNA can have similar effects on cancer phenotypes, i.e., allelic heterogeneity. This interchangeability and the erratic circumstances that produce somatic mutations lead to the mixture of mutation types observed in large cohorts of patient tumors[1].

Describing how somatic mutations in a gene impact cancer phenotypes requires integrating the information from all three mutation types. Currently, most studies linking somatic mutations to cancer phenotypes focus on one mutation type. This leads to missing associations for mutations primarily found in another type and reduced power to detect associations for mutations with high allelic heterogeneity that span the mutation types. Thus, a current obstacle facing those studying the downstream effects of somatic mutations is the lack of an established method for integrating PAMs, gene fusions, and CNAs into a comprehensive gene mutation profile. The lack of integration methods is due to several complicating factors. Firstly, the allelic heterogeneity observed in and between tumors means that different mutations in the same gene can be equivalently oncogenic. Second, it is challenging to discern driver (causal) from passenger (non-causal) somatic mutations. Third, an algorithm must be able to systematically integrate the binary PAM and gene fusion (mutated or not) with the quantitative copy number from CNAs. Lastly, some tumors have drastically higher somatic mutation rates than other tumors (e.g., microsatellite instability[5] and hypermutation[6]). These higher mutation rates confound any frequency-based integration approach and drive the discovery of spurious somatic mutations. We developed OncoMerge to fill the somatic mutation integration niche by providing an algorithm that systematically overcomes these obstacles to generate an integrated gene mutation profile. The input for the OncoMerge algorithm is the output from state-of-the-art methods for detecting PAMs (MC3[1] and MutSig2CV[7]), transcript fusions (PRADA[2,8]), and CNAs (GISTIC2.0[9]). Each of these methods provides the likelihood that a somatic mutation would happen by chance alone. Filtering on this statistic focuses integration efforts on genes most likely to harbor functional mutations. The integrated mutation profiles will give more power to detect associations with cancer phenotypes by capturing all tumors with a somatically mutated gene leading to a more comprehensive understanding of how genetic alterations drive cancer phenotypes.

The tremendous amount of cancer genome sequencing data generated in the last ten years has enabled efforts to discover and catalog somatic mutations across many cancers[1,10]. Many algorithms have been developed to discern which somatic mutations are drivers and how the mutations affect genes[6,7,11–17], and databases to search and view somatically mutated driver genes[16–18]. There also exist approaches for integrating somatic mutations. OncoPrint from cBioportal[18] can visually overlay somatic mutation types across patient tumors for a gene of

interest. The OncodriveROLE algorithm[13] was developed to discover driver genes by
systematically integrating PAMs and CNVs. However, neither OncoPrint nor OncodriveROLE
provides an integrated mutational profile that can be used in downstream analyses. The impact
of somatic mutations can be classified as activating (Act) gene function (typically found in
oncogenes), or loss of function (LoF) (typically found in tumor suppressor genes)[13]. It has also
been demonstrated that the systematic integration of PAM and CNA somatic mutations for a
gene improves the ability to determine Act or LoF status[13]. These foundational studies have
created a platform to develop an algorithm that systematically integrates the three somatic
mutation types.

The systematic integration of somatic mutations requires choosing a gene-level model that
determines how the data for the three somatic mutation types will be integrated, which in
OncoMerge is called the somatic mutation role. We determine the somatic mutation role by
employing rules similar to those used in OncodriveROLE[13] (**Figure 1**). The possible somatic
mutation roles in OncoMerge are PAM, Fusion, CNA amplification (CNAamp), CNA deletion
(CNAdel), Act, or LoF. The PAM, Fusion, CNAamp, and CNAdel somatic mutation roles use the
somatic mutation profile of the role in the integrated mutation matrix. The Act and LoF are
integrated mutation roles that harness allelic heterogeneity. Allelic heterogeneity is especially
prevalent in tumor suppressor genes, where mutations at many positions in a gene can impede
its function to prevent cancer phenotypes[3]. Allelic heterogeneity is less prevalent for oncogenes
where a small number of specific gain of function alleles are needed to drive cancer
phenotypes[3]. Genes underlying CNAs can add another layer of information as tumor
suppressors are often deleted, which has an equivalent oncogenic effect as missense or
truncating PAMs. The LoF role is designated when PAMs, Fusions, and CNAdels are
integrated. Oncogenes are often amplified as this typically leads to overexpression of the
underlying genes, which has a similar positive effect on gene function as a gain of function
PAM. The Act role is designated when PAMs, Fusions, and CNAamps are integrated.
Systematic determination of the somatic gene role and application of the rules laid out above
will be used to integrate the three mutation types into a comprehensive somatic mutation profile.

The algorithms developed to discern somatic mutation drivers for cancers provide a set of gold
standard mutations with gene roles that can be used to assess the performance of the new
OncoMerege algorithm. The gold standards are classified by whether the somatic mutation of a
gene was cancer-specific or not. The TCGA consensus[6] and Cancer Gene Census (CGC) from
COSMIC[11] were used to develop gold standards with cancer-specific somatically mutated gene
roles. The TCGA consensus is a list of driver genes identified from the TCGA Pan-Cancer Atlas
labeled with somatic mutation role (oncogene or tumor suppressor) and cancer type. The CGC
from COSMIC is an expert-curated database of human cancer driver genes labeled with
somatic mutation role (oncogene and tumor suppressor) and cancer type. The 20/20 rule[3],
OncodriveROLE[13], and Tokheim ensemble[12] were used to develop gold standards with
somatically mutated gene roles. The 20/20 rule defines oncogenes by requiring >20% of
mutations in recurrent positions and tumor suppressors as >20% of recorded mutations are
inactivating (missense or truncating)[3]. OncodriveROLE is a machine learning algorithm that
classifies genes according to their role (Act or LoF) based on well-curated genomic features[13].
The Tokheim ensemble is an ensemble-based method that integrates MutSigCV, 20/20+, and
TUSON methods for predicting gene roles (oncogene and tumor suppressor)[12]. Comparisons of
somatic mutation role between OncoMerge and the gold standards were facilitated by
converting oncogenes to Act and tumor suppressors to LoF. Finally, a combined gene role

agnostic gold standard was developed based on a union of all somatic mutations from all five
gold standards. These gold standards were used to assess the utility of filters and the quality of
the OncoMerege integrated somatic mutation matrices through their ability to recall somatic
mutations with the appropriate gene role.

A primary goal of OncoMerge is to construct a comprehensive somatic mutation profile that will
increase the power to identify how mutations modulate cancer phenotypes. Previously, we have
used the Systems Genetics Network AnaLysis (SYGNAL) pipeline[19] to build causal and
mechanistic gene regulatory networks (GRNs) for 31 cancers from the TCGA Pan-Cancer
Atlas[20]. Using SYGNAL, we link somatic mutations through the GRN to the hallmarks of
cancer[21,22], thereby linking somatic mutations to cancer phenotypes. These SYGNAL GRNs
describe how somatic mutations influence transcription factor (TF) or miRNA expression, which
modulates the expression of downstream genes. In SYGNAL, somatic mutations are used as
input for the Network Edge Orienting (NEO) portion of the pipeline that infers causal flows of
information (somatic mutation $\rightarrow$ TF or miRNA regulator $\rightarrow$ bicluster of co-regulated genes).
Thus, we use the OncoMerge integrated somatic mutation matrices in SYGNAL GRN inference
to demonstrate the increased power to identify how mutations modulate cancer phenotypes.

TFs are a significant factor in regulating gene expression in a cell, and interactions between TFs
could be used to explain much of the overall transcriptional state of a cell. Neph et al., 2012
constructed a human TF gene regulatory network by integrating genome-wide digital genomic
footprinting with DNA recognition motifs across 41 cell types[23]. The network architecture of
three-node network motifs was investigated[24–26] and shown to have a pattern similar to other
biologically derived networks. Because these TF regulatory networks were generated based on
DNA binding alone, they are not an active representation of the effect on transcript levels but
static DNA binding maps. On the other hand, SYGNAL GRNs are trained using coexpression as
an integral element of network construction. Therefore, SYGNAL GRNs can be considered
active because transcriptional effects support regulatory interactions. We compare and contrast
the underlying architecture of active TF regulatory networks from SYGNAL relative to static TF
regulatory networks from DNA binding maps.

As proof of principle, we apply OncoMerge to the multi-omic characterization of 32 cancers by
the TCGA PanCancer Atlas to develop filters and demonstrate a meaningful benefit for
downstream analyses. We demonstrate the power of using an integrated mutation matrix in
downstream analysis by re-analyzing the causal relationships for pan-cancer SYGNAL
networks[20]. We constructed transcription factor (TF) regulatory networks[23] and generated triad
significance profiles[24] to investigate the underlying network architecture[24–26]. We provide the
complete OncoMerge code, comprehensive mutation matrices for 32 TCGA cancers, regulatory
networks for 31 cancers, and TF regulatory network architecture for 25 cancers. These studies
demonstrate that OncoMege efficiently integrates PAMs, fusions, and CNAs into a
comprehensive mutational profile that strengthens downstream analyses linking somatic
mutations to cancer phenotypes.

Results

Establishing a baseline for the integration of somatic mutations

Somatic mutations play a significant role in cancer pathogenesis, and the main mutation types
are PAMs, fusions, and CNAs (amplifications and deletions). Somatic mutation of the same
gene with different mutation types can have similar downstream effects on cancer phenotypes.

We have developed OncoMerge as a systematic method to integrate PAM, fusion, and CNA
somatic mutations into a more comprehensive mutation matrix for subsequent analyses.
OncoMerge systematically integrates somatic mutations and defines a role for each gene
(**Figure 1**): PAM, fusion, CNA deletion (CNAdel), CNA amplification (CNAamp), Activating
(Act), and Loss of Function (LoF). The role assigned to a gene describes the rubric used to
integrate the data from the source data matrices.

A significant part of developing OncoMerge was constructing and optimizing the statistical filters
that provide an essential quality control step to identify somatically mutated genes that are more
likely to be functional in tumor biology. The selection and optimization of OncoMerge statistical
filters were performed using the 9,584 patient tumors from 32 cancers profiled by the TCGA
Pan-Cancer Atlas[1,6]. We used three metrics to assess the value of potential filters: 1) impact on
the number of somatically mutated genes (**Figure 2A**); 2) impact on the distribution of the
number of genes mapping to genomic loci (**Figure 2B**); and 3) significance of the overlap
between somatically mutated genes from OncoMerge with gold standard datasets (including
overlap with gene roles and tumor-specific gene roles; **Figure 2C**; **Supplementary Table 3**).
These metrics ensure that the integrated somatic mutations are consistent with prior knowledge
and that the size of CNA mutations does not overwhelm the integration algorithm.

Next, we determined the integration baseline by applying OncoMerge to the TCGA Pan-Cancer
Atlas without filtering. Slightly less than one-third of the genome was considered somatically
mutated in at least 5% or greater of tumors in at least one of the 32 cancers (30% or 6,028
genes, **Figure 2A**). We observed a highly significant overlap between OncoMerge somatically
mutated genes and the combined gold standard (genes = 395, p-value = 1.1×10^{-44}, **Figure 2C**)
when gene role was not considered. Significant overlaps existed between the LoF somatic
mutations from three gold standards (TCGA consensus, CGC, and Vogelstein) with the somatic
mutations with the LoF predicted role from OncoMerge (**Figure 2C**). None of the comparisons of
Act somatic mutations were significantly overlapping (**Figure 2C**). Many of the 6,028 genes map
to the same copy number alteration genomic locus (**Figure 2B**). These unfiltered results reveal
two main integration biases. First, there is no overlap of Act somatic mutations with previously
identified Act mutations. Second, the integration with CNAs is causing the inclusion of many
passenger mutations mapping to the same genomic locus. OncoMerge applied to the TCGA
Pan-Cancer Atlas without filtering provides a baseline to benchmark success. Addressing the
integration biases we observed is the impetus we had for developing and optimizing filters for
OncoMerge.

Developing a filtering strategy for the integration of somatic mutations
The power of integration is that aggregating somatic mutation information can boost the
mutation frequency of a gene enough to become significant, even though the constituent
mutations do not reach significance alone. The first filter determined if the final mutation
frequency after integrating PAM, fusion, and CNA somatic mutations is larger than expected by
chance alone. A permutation-based approach empirically determined the background integrated
mutation frequency distribution. Then the observed frequencies are compared to the
randomized background distribution to calculate permuted p-values, which are corrected using
the Benjamini-Hochberg method to provide permuted q-values. A permuted q-value ≤ 0.1
denotes a significant final mutation frequency. The permuted q-value (PQ) filter reduced the
number of somatically mutated genes to 4,240 (**Figure 2A**). This filtering improved LoF somatic
mutations from three to four gold standards (TCGA consensus, CGC, Vogelstein, and

OncodriveROLE) with the somatic mutations that had the LoF predicted role from OncoMerge.
Still, the Act comparisons did not show significant enrichment (**Figure 2C**). The PQ filter had a
minimal impact on the number of genes per locus (**Figure 2B**). This lack of significant overlap
for Act somatic mutations demonstrates that further filtering is required.

A key consideration in developing OncoMerge was that integrating the somatic mutation types
should highlight the functional somatic mutations over passenger mutations. Therefore, we
created a filter designed to prioritize somatically mutated genes that are more likely to be
functional. An average CNA encompasses 3.8 ± 7.9 Mb of genomic sequence[27], and genomic
segments of this size typically include many genes. These large genomic regions make it
difficult to determine which of the affected genes are the functional gene(s) underlying the CNA
locus without integrating additional information. We assert that passenger genes underlying a
CNA locus can be considered noise and can be identified by the lack of allelic heterogeneity.
Thus, functional gene(s) can be identified through allelic heterogeneity that boosts the somatic
mutation frequency for a gene above the background CNA frequency. We designed a low-pass
filter that retains only the gene(s) with the maximum final frequency (MFF). The MFF filter is
only applied if a locus has more than ten genes. Application of the MFF filter dramatically
reduced the number of somatically mutated genes from 6,028 to 1,459 (**Figure 2A**) and the
number of genes per locus (**Figure 2B**). The MFF filter also helps to make the average
contribution of PAMs and CNAs to the final mutation frequency more even (**Supplementary**
Figure 2). We additionally observed a marked improvement in overlap with the gold standards.
Significant enrichment was observed for four Act gold standards with somatic mutations that
OncoMerge predicts to be Act, and all five of the LoF gold-standard versus OncoMerge
predicted LoF comparisons (**Figure 2C**). The MFF filter directly addresses the issue of too many
genes in a CNA locus. Removing more than three-quarters of the somatically mutated genes
improves the overlaps with gold standards.

We then assessed the impact of applying both the PQ and MFF filters. Simultaneous application
of both filters led to a reduction in the number of somatically mutated genes beyond the MFF
filter (1,145 genes; **Figure 2A**), and the number of genes per locus was further improved
(**Figure 2B**). There was also an improvement in the significant overlap with gold standards
where all five LoF gold-standard versus OncoMerge predicted LoF and four Act gold-standard
versus OncoMerge predicted Act were significant (**Figure 2C**). Importantly, none of the gold
standard Act versus LoF or LoF versus Act comparisons were significant for any filter
combination, demonstrating that the OncoMerge predicted roles are consistent with prior
knowledge.

Reducing biases due to microsatellite instability and hypermutation
Microsatellite instability (MSI) and hypermutation phenotypes drastically increase the number of
somatic mutations in a tumor. The PQ and MFF filters and OncoMerge's core algorithm rely
upon somatic mutation frequency which is susceptible to confounding by MSI or hypermutation.
Fortunately, all TCGA tumors used in this study are characterized for both MSI[5] and
hypermutation[6] status (**Figure 3A**). We observed a highly significant positive correlation
between MSI/hypermutation frequency and the total number of somatic mutations per cancer
after integration by OncoMerge (R = 0.68 and p-value = 2.0 x 10[-5]). This strong positive
correlation demonstrates that MSI/hypermutation is likely inflating the number of somatic
mutations discovered by OncoMerge. Therefore, we created the MSI and hypermutation
censoring filter (MHC) to exclude these tumors while OncoMerge determines which genes to

include in the final somatic mutation matrix. The mutation status for tumors with MSI and
hypermutation are included for genes in the final integrated mutation matrix. Applying the MHC
filter alongside the PQ and MFF filters reduced the overall number of somatically mutated genes
(905 genes; **Figure 2A**) and had minimal impact on the number of genes per locus (**Figure 2B**;
Supplementary Table 4). The combined PQ, MFF, and MHC filters decreased the correlation
between the MSI/hypermutation frequency (R = 0.48 and p-value = 5.8 x 10^{-3}). All ten of the
gold standard Act vs. Act and LoF vs. LoF comparisons were significant. These results
established that the MHC filter is valuable for removing passenger mutations introduced by
tumors with severely increased somatic mutation rates. The PQ, MFF, and MHC filters comprise
the default and final OncoMerge filter set. The filters deal with known complications in cancer
genetics and ensure that the mutation roles in the integrated matrix are correctly assigned.

Sensitivity anlaysis to optimize filtering cutoffs and parameters
We then used sensitivity analysis of the filtering parameters to determine their optimal
parameterization for OncoMerge (**Figure 2D-G** and **Supplementary Table 4**). First, we tested
whether the GISTIC threshold cutoff for shallow or deep amplification or deletion (shallow >=1
and deep >=2) was optimal for integration purposes (**Figure 2D**). The shallow GISTIC threshold
led to many more somatically mutated genes, but this had a negative impact on the discovery of
meaningful activating mutations, as shown by the lack of significant overlap with gold standard
activating mutations across the board. Based on these results, we demonstrated that the deep
GISTIC threshold was optimal for OncoMerge. Second, we varied the gene number threshold of
a CNA locus for the MFF with the values 1, 5, 10, 15, and 20 genes (**Figure 2E**). After the 10
genes per loci threshold, the significance for the gold standard activating mutations for Tokheim
et al., 2016 was lost, and after 15 genes per loci for the TCGA consensus. These results
demonstrate that a 10 genes per loci cutoff for the MFF is optimal. Next, we tested the
sensitivity of OncoMerge to the minimum mutation frequency threshold by setting it with the
values of 0.01, 0.05, 0.1, and 0.2 (**Figure 2F**). Impressively, there was little impact on the
significance of the gold-standard analysis across the range of thresholds. However, the number
of somatic mutations is significantly impacted by this threshold. Somatic mutation discovery
could warrant a minimum mutation threshold of 1% or less. On the other hand, downstream
analyses warrant a 5% minimum mutation threshold to ensure sufficient somatically mutated
samples to achieve statistical power. Finally, we tested the sensitivity of OncoMerge to the
permuted q-value threshold from the PQ filter across the values 0.01, 0.05, 0.1, and 0.2 (**Figure
2G**). The lowest threshold of 0.01 led to a loss of significance for the overlap of the activating
mutations for the Vogelstein et al., 2013 gold standard. The number of somatically mutated
genes increased as the permuted q-value threshold increased, and thus 0.05, 0.1, and 0.2 are
all reasonable threshold values. The threshold of 0.1 was chosen based on the statistical
meaning of the PQ filter to remove most of the integrated somatic mutations that could have
happened by chance alone while not being too restrictive. These sensitivity analyses provide a
reasonable rationale for choosing the values for the filtering cutoffs and parameters, for which
alternative values might be used, and in which contexts.

Benefits of an integrated somatic mutation matrix
We evaluated the benefits of systematic somatic mutation integration by comparing OncoMerge
integrated somatic mutation matrices to those from PAMs. The PAM somatic mutation matrices
were used as a reference point because we have successfully used them as the sole source for
somatic mutations in previous studies[19,20]. We assessed the benefits of integration by tabulating
the number of somatic mutations and their roles (**Figure 3B**), the number of genes added by

integration (**Figure 3C**), and the increase in somatic mutation frequency due to integration (**Figure 3E**). Impressively, Act and LoF mutations represented the bulk of the somatic mutations in 30 cancers (**Figure 3B**). The papillary thyroid carcinoma (THCA) and kidney chromophobe (KICH) were the only cancers that lacked Act or LoF mutations. Consistent with Agrawal et al. 2014[28], THCA had only three mutations with a frequency ≥ 5% BRAF, NRAS, and RET. On the other hand, KICH was under-sampled in the TCGA Pan-Cancer atlas (n = 65), and LoF and Act mutations would likely be discovered with the inclusion of more patient tumors.

We then investigated how many new genes the integration added for each cancer. Integration added at least one somatically mutated gene for each cancer (**Figure 3C**), and more than eighty somatically mutated genes for BLCA, LUAD, and UCEC (**Figure 3C**). The somatically mutated genes added by OncoMerge make the integrated somatic mutation matrices more comprehensive.

Next, we investigated the frequencies of the somatic mutations from the OncoMerge integrated mutation matrices. The genes with the highest frequency map to well-known oncogenes (e.g., BRAF) and tumor suppressors (e.g., APC and TP53; **Figure 3D**). The two tumor suppressor genes APC and TP53 were mutated in greater than eighty percent of the tumors for multiple cancers (**Figure 3D**). The APC gene was mutated in greater than eighty percent of tumors for colon adenocarcinoma (COAD) and rectal adenocarcinoma (READ). The TP53 gene was mutated in greater than eighty percent of tumors for esophageal carcinoma (ESCA), lung squamous carcinoma (LUSC), ovarian serous cystadenocarcinoma (OV), rectal carcinoma (READ), and uterine carcinosarcoma (UCS). These frequently mutated genes in the OncoMerge integrated mutation matrices are consistent with prior knowledge of somatic mutations for each cancer.

Finally, we calculated the frequency added through integration by subtracting the integrated mutation frequency from the PAM frequency. The most substantial increases in somatic mutation frequency were observed for TMPRSS2-ERG in prostate adenocarcinoma (PRAD) and CDKN2A in mesothelioma (MESO), glioblastoma (GBM), diffuse large B-cell lymphoma (DLBC), and esophageal carcinoma (ESCA; **Figure 3E**). Neither TMPRSS2-ERG nor CDKN2A would have been identified as somatically mutated without incorporating fusions and CNAs, respectively. These findings demonstrate that OncoMerge significantly improves the number and frequency of somatically mutated genes in most cancers. Also, these results show that the systematic integration of PAM, fusion, and CNA somatic mutations is crucial for obtaining a comprehensive mutation matrix for each cancer.

Pan-cancer somatic mutations capture many known tumor suppressors and oncogenes

Genes mutated in multiple cancers are of great interest as selective pressures have found a common solution in different contexts to influence cancer phenotypes. Therefore, we searched for genes somatically mutated in at least five cancers in the OncoMerge integrated mutation matrices. The resulting gene list could be broken down into two groups of somatic mutations: the LoF set (n = 23, **Figure 4A**) and the Act set (n = 14, **Figure 4B**). The FBXW7, KMT2C, and KMT2D somatic mutations were challenging to classify as LoF or Act. The genes FBXW7 and KMT2D were somatically mutated with PAMs in six and seven cancers, respectively (**Figure 4A**). The gene KMT2C (also known as MLL3) was primarily LoF and PAM but had the mutation role of Act for ovarian cancer (OV) (**Figure 4B**). Based on a literature search, all three genes

have been classified as tumor suppressors[29–31]. Therefore, we grouped FBXW7, KMT2C, and
KMT2D mutations with the LoF set.

The pan-cancer somatically mutated genes harbored many well-known tumor suppressors and
oncogenes (**Figure 4C**). As expected, tumor suppressors[31] were significantly enriched in the
LoF group (overlap = 18, p-value = 5.0 x 10^{-20}), and oncogenes[32] were significantly enriched in
the Act group (overlap = 8, p-value = 4.4 x 10^{-10}). The top three most somatically mutated tumor
suppressors were TP53, PTEN, and CDKN2A. These three tumor suppressors control important
checkpoints in the cell cycle making them functionally interesting. The gene TP53 was
somatically mutated in 24 cancers, primarily by PAMs, but four LoF were also observed for
glioblastoma (GBM), liver hepatocellular carcinoma (LIHC), prostate adenocarcinoma (PRAD),
and sarcoma (SARC). The top three most mutated oncogenes across cancers were PIK3CA,
KRAS, and CCNE1. Two of these genes (PIK3CA and KRAS) become overactive kinases when
mutated, and CCNE1 is a fundamental part of the cell cycle regulatory machinery. Both PIK3CA
and KRAS have PAM and Act mutation roles across the different cancers, and only the NFE2L2
gene has a similar mixture of PAM and Act mutation roles. The remainder of the oncogenes are
like CCNE1 in that the gene somatic mutation roles are all Act. These pan-cancer analyses
further validate the systematic somatic mutation integration by OncoMerge through the
unbiased recall of tumor suppressors and oncogenes.

Improving gene regulatory network inference
A major goal of developing OncoMerge was to construct an integrated somatic mutation profile
that would increase the power to identify how mutations modulate cancer phenotypes.
Previously we used PAMs from the cancers in the TCGA Pan-Cancer Atlas as input for
SYGNAL to construct gene regulatory networks (GRNs)[20]. SYGNAL GRNs are composed of
causal and mechanistic interactions linking somatic mutations to a TF or miRNA regulator to a
co-regulated set of genes (bicluster). Somatic mutations in SYGNAL are used as input for the
Network Edge Orienting (NEO) portion of the pipeline that infers causal flows of information
(somatic mutation → TF or miRNA regulator → bicluster of co-regulated genes). Therefore, we
recomputed NEO analyses using the OncoMerge integrated somatic mutation matrices for each
cancer in the TCGA Pan-Cancer Atlas to demonstrate the increased power to detect causal
flows of information. The resulting networks were filtered to include only biclusters with good
quality coexpression that were significantly associated with patient survival. Regulatory
interactions were required to be both causal (significant evidence of information flow between a
mutation → regulator → bicluster) and mechanistic (enrichment of regulator binding sites in the
promoter or 3' UTR of the bicluster genes). We compare SYGNAL GRNs inferred using
OncoMerge integrated mutation matrices (**Supplementary Table 5**) with SYGNAL GRNs
inferred using PAM-based mutation matrices (legacy).

The GRNs are comprised of nodes and edges. The degree of a node is the number of edges
connecting it to other nodes. The average degree is a standard network metric computed as the
average of all node degrees in the network. We found that the average degree was larger for 23
OncoMerge GRNs relative to legacy GRNs (**Figure 5A**). The exceptions were GBM (average
degree was equal) and COAD, KIRC, SKCM, and STAD (legacy had a larger average degree).
COAD, SKCM, and STAD have many MSI and hypermutation tumors (**Figure 3A**), suggesting
that the MHC filter removed spurious associations. Furthermore, the hypothesis that MSI and
hypermutation inflated the average degree of GRNs is supported by the reduction in the number
of COAD and STAD mutations in the OncoMerge GRN relative to the legacy GRN (**Figure 5B**).

Thus, we have increased the average degree in the most networks and addressed a systematic
bias found in legacy networks.

Next, we considered the number of mutations in each GRN predicted to modulate the activity of
regulators. The OncoMerge GRNs contained more somatic mutation nodes than the legacy
GRNs for all cancers but COAD and STAD, likely due to MSI and hypermutation as described
above (**Figure 5B**). Then, we assessed the recall of somatic mutations previously associated
with each cancer from the DisGeNET database[33]. All but two OncoMerge GRNs recalled more
previously associated somatic mutations than the legacy GRNs (**Figure 5C**). The exceptions
were UVM with the same amount and COAD with fewer (**Figure 5C**). This demonstrates that
OncoMerge integrated mutation matrices provide increased power for linking somatic mutation
matrices into GRNs, and improve the capture of somatic mutations previously associated with
each cancer.

Finally, we considered the number of predicted causal and mechanistic transcription factor (TF)
regulators in each GRN. The OncoMerge GRNs contained more predicted TF regulators than
legacy for 22 GRNs, the same number of TF regulators for STAD, and less TF regulators for
COAD, GBM, KIRC, THCA, and UCEC (**Figure 5D**). We also assessed the recall of TFs
previously associated with each cancer from the DisGeNET database[33,34]. Twenty of the
OncoMerge GRNs recalled more previously associated TFs than legacy GRNs (**Figure 5E**).
The COAD and UCEC GRNs had the same amount, and GBM, KIRP, SKCM, and THCA had
fewer, and KICH and UVM had no recall of previously associated TFs in either GRN (**Figure
5E**). In summary, using OncoMerge integrated mutation matrices in GRN construction builds
more extensive and biologically meaningful networks.

Comparing active and static TF regulatory network architectures
The interactions between TFs are important for generating the transcriptional state of a human
cell. The underlying architecture of TF regulatory networks, comprised of TFs and their
interactions, are typically explored by enumerating all three-node network motifs and computing
their enrichment or depletion into triad significance profiles (TSPs)[24]. Most studies of network
motif enrichment have relied upon unsigned interactions[23,24,35–38] which ignore whether the
interaction is activating or repressing. To facilitate comparisons, our first analysis of network
architecture uses unsigned TSPs to compare static and active TF regulatory networks. Static TF
regulatory networks were constructed using chromatin accessibility and DNA binding motifs for
41 cell types[23]. These TF regulatory networks are static because they do not incorporate gene
expression data in their construction. Active TF regulatory networks are derived from the
OncoMerge augmented SYGNAL pan-cancer GRNs, that were trained using patient tumor
transcriptional data and therefore are comprised of active TF regulatory interactions. Using the
following steps, we constructed TF regulatory networks for each cancer from the pan-cancer
SYGNAL GRNs (**Figure 6A**). First, we extracted all the TF regulators from the pan-cancer
GRNs. Interactions between TFs were inferred based on the presence of DNA binding motifs
from the TF target gene database[19], and a significant correlation between the TF regulator and
TF target in patient tumor expression (Pearson's $|R| \geq 0.3$ and p-value ≤ 0.05; **Figure 6A**;
Supplementary Table 6). The enrichment (or depletion) of motifs in the network was computed
using triad significance profiles (TSPs)[24]. Triad significance profiles were calculated for twenty-
five TF regulatory networks and summarized as the median TSP (**Figure 6A** & **B**). We excluded
the cancer types DLBC, KICH, KIRP, OV, TGCT, and THYM because they had too few inferred

regulatory interactions (< 50 interactions). Finally, we recomputed the TSPs for the static TF
regulatory networks using a more recent version of the mfinder algorithm (**Figure 6B**).

The median TSPs of the active and static TF regulatory networks were highly correlated (R =
0.75, p-value = 3.0 x 10^{-3}; **Figure 6B**). Demonstrating that the architecture of the active network
resembles the static network. However, the maximum enriched network motifs were different.
The regulated and regulating feedback motifs (motifs 108 and 46) were the most highly enriched
motifs from the static TF regulatory networks and were still enriched, although not as significant
as in the active networks. In contrast, the feed-forward loop (FFL, motif 38) is the most highly
enriched motif in the active TF regulatory networks. These two motifs are quite similar in
structure and differ only by a single edge. Feedback motifs and FFLs can be further broken
down into ten and eight signed network motifs that each have a unique functional output[39]. Thus
exploring the enrichment of signed network motifs allows the discovery of what functions are
being selected for by evolution in general and the microcosm of tumor biology.

Coherent feed-forward loops enriched in active TF regulatory networks
Incorporating the sign of the regulatory interactions (activating or repressing) splits the FFL motif
into eight signed network motifs classified as coherent (C1, C2, C3, C4) and incoherent (I1, I2,
I3, I4)[39]. Simulation studies have demonstrated that coherent FFLs lead to delays in target gene
expression, and incoherent FFLs accelerate target gene expression[39]. FFLs were significantly
enriched in active TF regulatory networks, which led us to question whether coherent,
incoherent, or both FFLs were enriched. In active GRNs, the sign of the correlation between the
TF regulator to TF target can be used to determine the sign of the interaction (R > 0 equates to
activation, R < 0 equates to repression). The four coherent FFLs were enriched in the active TF
regulatory networks (**Figure 6C**; **Supplementary Table 7**), and incoherent FFLs were severely
under-enriched (Z << 0). In summary, coherent FFLs were enriched in our active TF regulatory
networks, suggesting that transcriptional delay mechanisms must provide a valuable function for
TF regulatory networks.

Coherent switch-like feedback motifs enriched in active TF regulatory networks
The regulated and regulating mutual feedback motifs have a two-node feedback loop at their
core. The double-positive and double-negative two-node mutual feedback loops act like
switches[40]. We tested the twenty signed regulated and regulating mutual feedback network
motif configurations for enrichment in TF regulatory networks. Three regulating and three
regulated signed mutual feedback motifs (**Figure 6C**; **Supplementary Table 7**). These six
enriched regulated and regulating mutual feedback motifs had a commonality in their
configuration. Firstly, all the network motifs were coherent. Coherent regulated and regulating
feedback loops have interaction signs between the feedback loop that are either double-positive
or double-negative. And the regulated or regulating node interacts with the feedback loop nodes
using the same sign for double-positive feedback loops and the opposite sign for double-
negative feedback loops. Thus, there are three coherent configurations for both regulated and
regulating mutual feedback motifs making six total, coinciding with the six enriched
configurations (**Figure 6C**; **Supplementary Table 7**). The enriched motifs containing a double-
positive feedback loop had the same interactions with the non-feedback loop node, both
activating or repressing (**Figure 6C**). The enriched motif containing a double-negative feedback
loop had opposing interactions with the non-feedback loop node, one activating and one
repressing (**Figure 6C**). These enriched signed network motifs are the configurations that

function as molecular switches[41]. Again, evolution has selected for coherent network motif
configurations likely because of their function.

Discussion

We developed OncoMerge to integrate PAMs, fusions, and CNAs into a more accurate
representation of the somatic mutation landscape of patient tumors. The OncoMerge integration
algorithm and three filters (PQ, MFF, and MHC) effectively address the issues of allelic
heterogeneity and the unification of binary and quantitative mutation data. These issues have
forced most studies of somatic mutations to focus on one somatic mutation type and were the
impetus for us to develop OncoMerge for the integration of the three most common somatic
mutation types. We tested OncoMerge by integrating the somatic mutation data from 32 cancers
from the TCGA Pan-Cancer Atlas. Comparison to gold standards confirmed that the genes and
roles selected by OncoMerge were accurate. The integration of somatic mutation types had
several quantifiable benefits for somatically mutated genes. First, most somatically mutated
genes had an integrated role of Act or LoF, demonstrating that consolidation of allelic
heterogeneity is vital to achieving a complete picture of somatic mutations for a patient cohort.
Second, genes somatically mutated primarily by fusions and CNAs were added by the
integration. Lastly, the frequency of many somatically mutated genes increased due to the
integration of the three somatic mutation types. We used the integrated somatic mutations as
input to SYGNAL to demonstrate improvements in power for systems genetics-based inference
of GRNs. Using integrated somatic mutations increased the average connectedness of the
GRNs by incorporating more somatic mutations and regulators previously linked to cancer
biology. Next, we found that while the underlying architecture of active SYGNAL TF regulatory
networks and static DNA binding TF regulatory networks were similar overall, the top most
enriched network motifs were different. We discovered that switch-like feedback and delay-
inducing feed-forward loop motifs were enriched in TF regulatory networks. We developed and
tested a novel systematic integration tool and demonstrated that integrated somatic mutations
improve our ability to link somatic mutations with cancer phenotypes.

The construction of active GRNs enabled the exploration of signed network motifs and led to the
discovery that specific signed network motif configurations are being enriched. The SYGNAL
GRNs construction method identifies active gene regulatory interactions by discovering
interactions that are supported by gene expression data from patient tumors[19]. On the other
hand, prior networks were static maps of DNA binding sites constructed using digital genomic
footprinting and the similarity of the underlying sequence of the footprints for known DNA
binding motifs[23]. The active networks use a correlation-based method to determine TF
regulatory roles (activator or repressor) for the interactions, which is not possible using static
binding maps. Analyzing signed network motifs provides a leap forward in understanding how
the underlying architecture of GRNs functions in real-world biological systems. OncoMerge
integrated somatic mutations offer a more solid platform to infer active GRNs that can be used
to explore the functional architecture of TF regulatory networks.

We discovered that coherent regulated and regulating feedback and FFL network motifs were
enriched in cancer TF regulatory networks. We cannot say whether this enrichment of network
motifs will generalize to all active GRNs or if this is a cancer-specific phenomenon. In normal
organismal development, feedback motifs have been previously shown to be important for cell
fate decision-making[42,43]. On the other hand, in tumor cells and other cells in the tumor
microenvironment, the enriched feedback motifs may be maintaining a cell fate, or the disease

could be coopting the circuit to drive tumor biology. Likewise, coherent FFL network motifs have
also been associated with enhanced drug resistance[44]. These coherent motifs are relevant for
normal and diseased cell biology, and evolution has specifically selected these motif
configurations because of their unique functional outputs.

We provide the Oncomerge software and data in several standard distribution formats to
facilitate future studies that aim to integrate somatic mutations. The source code for OncoMerge
is available on GitHub (https://github.com/plaisier-lab/OncoMerge). Finally, an OncoMerge
Docker image was created that can be run as a virtual machine with all dependencies pre-
installed (https://hub.docker.com/r/cplaisier/oncomerge). Detailed documentation is provided,
along with a tutorial that describes the use of OncoMerge. The goal of disseminating
OncoMerge in these ways is to give end-users flexibility to choose what distribution method best
fits their computational platform.

We also provide the Pan-Cancer Atlas TCGA somatic mutation data used as input for
OncoMerge (https://doi.org/10.6084/m9.figshare.21760964.v1). And the resulting OncoMerge
integrated somatic mutation matrices for those planning studies that use somatic mutations from
the TCGA Pan-Cancer Atlas (https://doi.org/10.6084/m9.figshare.20238867). These integrated
somatic mutation matrices can be used for any downstream analyses incorporating somatic
mutations and will provide the same power boost observed in our studies. In addition, we also
offer the pan-cancer SYGNAL GRNs and TF regulatory networks as supplementary tables to
expedite systems genetics studies of TCGA cancers. We hope these accessible results will
facilitate studies linking somatic mutations to downstream cancer phenotypes and lead to novel
biological insights in clinical samples.

Future improvements to the OncoMerge algorithm include a more quantitative integration
approach for the somatic mutations, a replacement for or an improved maximum final frequency
filter, aggregation across pathways, and a determination of whether other genomic features may
be integrated (ecDNA[45] or epigenomics[46]). Additionally, in future single-cell studies with both
transcriptome and genome information, it would be helpful to have an OncoMerge
implementation that integrates PAM, fusion, and CNA for every single cell. We envision
OncoMerge as a valuable tool in the somatic mutation characterization pipeline. We hope that it
will facilitate multi-omic studies and lead to novel discoveries that can be translated into clinical
insights.

Acknowledgments

555   This work was supported by NIH-NINDS Award # 1R01NS123038-01, and 1R01NS119650-01.
556   The authors also acknowledge Robert Schultz for assistance in preliminary studies, and the
557   Cancer Genome Atlas Research Network for the TCGA Pan-Cancer Atlas multi-omic patient
558   tumor profiles.

References

1. Ellrott, K., Bailey, M.H., Saksena, G., Covington, K.R., Kandoth, C., Stewart, C., Hess, J., Ma,
S., Chiotti, K.E., McLellan, M., et al. (2018). Scalable Open Science Approach for Mutation
Calling of Tumor Exomes Using Multiple Genomic Pipelines. Cell Syst *6*, 271-281.e7.
10.1016/j.cels.2018.03.002.

2. Hu, X., Wang, Q., Tang, M., Barthel, F., Amin, S., Yoshihara, K., Lang, F.M., Martinez-
Ledesma, E., Lee, S.H., Zheng, S., et al. (2018). TumorFusions: an integrative resource for
cancer-associated transcript fusions. Nucleic Acids Res *46*, D1144–D1149.
10.1093/nar/gkx1018.

3. Vogelstein, B., Papadopoulos, N., Velculescu, V.E., Zhou, S., Diaz, L.A., and Kinzler, K.W.
(2013). Cancer genome landscapes. Science *339*, 1546–1558. 10.1126/science.1235122.

4. Beroukhim, R., Mermel, C.H., Porter, D., Wei, G., Raychaudhuri, S., Donovan, J., Barretina,
J., Boehm, J.S., Dobson, J., Urashima, M., et al. (2010). The landscape of somatic copy-
number alteration across human cancers. Nature *463*, 899–905. 10.1038/nature08822.

5. Bonneville, R., Krook, M.A., Kautto, E.A., Miya, J., Wing, M.R., Chen, H.-Z., Reeser, J.W.,
Yu, L., and Roychowdhury, S. (2017). Landscape of Microsatellite Instability Across 39
Cancer Types. JCO Precis Oncol *2017*. 10.1200/PO.17.00073.

6. Bailey, M.H., Tokheim, C., Porta-Pardo, E., Sengupta, S., Bertrand, D., Weerasinghe, A.,
Colaprico, A., Wendl, M.C., Kim, J., Reardon, B., et al. (2018). Comprehensive
Characterization of Cancer Driver Genes and Mutations. Cell *173*, 371-385.e18.
10.1016/j.cell.2018.02.060.

7. Lawrence, M.S., Stojanov, P., Polak, P., Kryukov, G.V., Cibulskis, K., Sivachenko, A., Carter,
S.L., Stewart, C., Mermel, C.H., Roberts, S.A., et al. (2013). Mutational heterogeneity in
cancer and the search for new cancer-associated genes. Nature *499*, 214–218.
10.1038/nature12213.

8. Torres-García, W., Zheng, S., Sivachenko, A., Vegesna, R., Wang, Q., Yao, R., Berger, M.F.,
Weinstein, J.N., Getz, G., and Verhaak, R.G.W. (2014). PRADA: pipeline for RNA
sequencing data analysis. Bioinformatics *30*, 2224–2226. 10.1093/bioinformatics/btu169.

9. Mermel, C.H., Schumacher, S.E., Hill, B., Meyerson, M.L., Beroukhim, R., and Getz, G.
(2011). GISTIC2.0 facilitates sensitive and confident localization of the targets of focal
somatic copy-number alteration in human cancers. Genome Biol *12*, R41. 10.1186/gb-2011-
12-4-r41.

10.   ICGC/TCGA Pan-Cancer Analysis of Whole Genomes Consortium (2020). Pan-cancer
analysis of whole genomes. Nature *578*, 82–93. 10.1038/s41586-020-1969-6.

11.   Sondka, Z., Bamford, S., Cole, C.G., Ward, S.A., Dunham, I., and Forbes, S.A. (2018).
The COSMIC Cancer Gene Census: describing genetic dysfunction across all human
cancers. Nat Rev Cancer *18*, 696–705. 10.1038/s41568-018-0060-1.

12.   Tokheim, C.J., Papadopoulos, N., Kinzler, K.W., Vogelstein, B., and Karchin, R. (2016).
Evaluating the evaluation of cancer driver genes. Proc Natl Acad Sci U S A *113*, 14330–
14335. 10.1073/pnas.1616440113.

13.   Schroeder, M.P., Rubio-Perez, C., Tamborero, D., Gonzalez-Perez, A., and Lopez-
Bigas, N. (2014). OncodriveROLE classifies cancer driver genes in loss of function and
activating mode of action. Bioinformatics *30*, i549-555. 10.1093/bioinformatics/btu467.

14.   Lawrence, M.S., Stojanov, P., Mermel, C.H., Robinson, J.T., Garraway, L.A., Golub,
T.R., Meyerson, M., Gabriel, S.B., Lander, E.S., and Getz, G. (2014). Discovery and
saturation analysis of cancer genes across 21 tumour types. Nature *505*, 495–501.
10.1038/nature12912.

15.   Davoli, T., Xu, A.W., Mengwasser, K.E., Sack, L.M., Yoon, J.C., Park, P.J., and Elledge,
S.J. (2013). Cumulative haploinsufficiency and triplosensitivity drive aneuploidy patterns and
shape the cancer genome. Cell *155*, 948–962. 10.1016/j.cell.2013.10.011.

16.   Tamborero, D., Gonzalez-Perez, A., and Lopez-Bigas, N. (2013). OncodriveCLUST:
exploiting the positional clustering of somatic mutations to identify cancer genes.
Bioinformatics *29*, 2238–2244. 10.1093/bioinformatics/btt395.

17.   Gonzalez-Perez, A., Perez-Llamas, C., Deu-Pons, J., Tamborero, D., Schroeder, M.P.,
Jene-Sanz, A., Santos, A., and Lopez-Bigas, N. (2013). IntOGen-mutations identifies cancer
drivers across tumor types. Nat. Methods *10*, 1081–1082. 10.1038/nmeth.2642.

18.   Cerami, E., Gao, J., Dogrusoz, U., Gross, B.E., Sumer, S.O., Aksoy, B.A., Jacobsen, A.,
Byrne, C.J., Heuer, M.L., Larsson, E., et al. (2012). The cBio cancer genomics portal: an
open platform for exploring multidimensional cancer genomics data. Cancer Discov *2*, 401–
404. 10.1158/2159-8290.CD-12-0095.

19.   Plaisier, C.L., O'Brien, S., Bernard, B., Reynolds, S., Simon, Z., Toledo, C.M., Ding, Y.,
Reiss, D.J., Paddison, P.J., and Baliga, N.S. (2016). Causal Mechanistic Regulatory Network
for Glioblastoma Deciphered Using Systems Genetics Network Analysis. Cell Syst *3*, 172–
186. 10.1016/j.cels.2016.06.006.

20.   Thorsson, V., Gibbs, D.L., Brown, S.D., Wolf, D., Bortone, D.S., Ou Yang, T.-H., Porta-
Pardo, E., Gao, G.F., Plaisier, C.L., Eddy, J.A., et al. (2018). The Immune Landscape of
Cancer. Immunity *48*, 812-830.e14. 10.1016/j.immuni.2018.03.023.

21.   Hanahan, D. (2022). Hallmarks of Cancer: New Dimensions. Cancer Discov *12*, 31–46.
10.1158/2159-8290.CD-21-1059.

22.   Hanahan, D., and Weinberg, R.A. (2011). Hallmarks of cancer: the next generation. Cell
144, 646–674. 10.1016/j.cell.2011.02.013.

23.   Neph, S., Stergachis, A.B., Reynolds, A., Sandstrom, R., Borenstein, E., and
Stamatoyannopoulos, J.A. (2012). Circuitry and dynamics of human transcription factor
regulatory networks. Cell *150*, 1274–1286. 10.1016/j.cell.2012.04.040.

24.   Milo, R., Itzkovitz, S., Kashtan, N., Levitt, R., Shen-Orr, S., Ayzenshtat, I., Sheffer, M.,
and Alon, U. (2004). Superfamilies of evolved and designed networks. Science *303*, 1538–
1542. 10.1126/science.1089167.

25.   Milo, R., Shen-Orr, S., Itzkovitz, S., Kashtan, N., Chklovskii, D., and Alon, U. (2002).
Network motifs: simple building blocks of complex networks. Science *298*, 824–827.
10.1126/science.298.5594.824.

26.   Wernicke, S., and Rasche, F. (2006). FANMOD: a tool for fast network motif detection.
Bioinformatics *22*, 1152–1153. 10.1093/bioinformatics/btl038.

27.   Harbers, L., Agostini, F., Nicos, M., Poddighe, D., Bienko, M., and Crosetto, N. (2021).
Somatic Copy Number Alterations in Human Cancers: An Analysis of Publicly Available Data
From The Cancer Genome Atlas. Front Oncol *11*, 700568. 10.3389/fonc.2021.700568.

28.   Cancer Genome Atlas Research Network (2014). Integrated genomic characterization of
papillary thyroid carcinoma. Cell *159*, 676–690. 10.1016/j.cell.2014.09.050.

29.   Gala, K., Li, Q., Sinha, A., Razavi, P., Dorso, M., Sanchez-Vega, F., Chung, Y.R.,
Hendrickson, R., Hsieh, J.J., Berger, M., et al. (2018). KMT2C mediates the estrogen
dependence of breast cancer through regulation of ERα enhancer function. Oncogene *37*,
4692–4710. 10.1038/s41388-018-0273-5.

30.   Hillman, R.T., Celestino, J., Terranova, C., Beird, H.C., Gumbs, C., Little, L., Nguyen, T.,
Thornton, R., Tippen, S., Zhang, J., et al. (2018). KMT2D/MLL2 inactivation is associated
with recurrence in adult-type granulosa cell tumors of the ovary. Nat Commun *9*, 2496.
10.1038/s41467-018-04950-x.

31.   Zhao, M., Kim, P., Mitra, R., Zhao, J., and Zhao, Z. (2016). TSGene 2.0: an updated
literature-based knowledgebase for tumor suppressor genes. Nucleic Acids Res *44*, D1023-
1031. 10.1093/nar/gkv1268.

32.   Liu, Y., Sun, J., and Zhao, M. (2017). ONGene: A literature-based database for human
oncogenes. J Genet Genomics *44*, 119–121. 10.1016/j.jgg.2016.12.004.

33.   Piñero, J., Ramírez-Anguita, J.M., Saüch-Pitarch, J., Ronzano, F., Centeno, E., Sanz,
F., and Furlong, L.I. (2020). The DisGeNET knowledge platform for disease genomics: 2019
update. Nucleic Acids Res. *48*, D845–D855. 10.1093/nar/gkz1021.

34.   Lambert, S.A., Jolma, A., Campitelli, L.F., Das, P.K., Yin, Y., Albu, M., Chen, X., Taipale,
J., Hughes, T.R., and Weirauch, M.T. (2018). The Human Transcription Factors. Cell *172*,
650–665. 10.1016/j.cell.2018.01.029.

35.   Li, Y., Shao, T., Jiang, C., Bai, J., Wang, Z., Zhang, J., Zhang, L., Zhao, Z., Xu, J., and
Li, X. (2015). Construction and analysis of dynamic transcription factor regulatory networks in
the progression of glioma. Sci Rep *5*, 15953. 10.1038/srep15953.

36.   Stergachis, A.B., Neph, S., Sandstrom, R., Haugen, E., Reynolds, A.P., Zhang, M.,
Byron, R., Canfield, T., Stelhing-Sun, S., Lee, K., et al. (2014). Conservation of trans-acting
circuitry during mammalian regulatory evolution. Nature *515*, 365–370. 10.1038/nature13972.

37.   Gerstein, M.B., Kundaje, A., Hariharan, M., Landt, S.G., Yan, K.-K., Cheng, C., Mu, X.J.,
Khurana, E., Rozowsky, J., Alexander, R., et al. (2012). Architecture of the human regulatory
network derived from ENCODE data. Nature *489*, 91–100. 10.1038/nature11245.

38. Boyle, A.P., Araya, C.L., Brdlik, C., Cayting, P., Cheng, C., Cheng, Y., Gardner, K.,
Hillier, L.W., Janette, J., Jiang, L., et al. (2014). Comparative analysis of regulatory
information and circuits across distant species. Nature *512*, 453–456. 10.1038/nature13668.

39. Mangan, S., and Alon, U. (2003). Structure and function of the feed-forward loop
network motif. Proc Natl Acad Sci U S A *100*, 11980–11985. 10.1073/pnas.2133841100.

40. Alon, U. (2007). Network motifs: theory and experimental approaches. Nat Rev Genet *8*,
450–461. 10.1038/nrg2102.

41. Gardner, T.S., Cantor, C.R., and Collins, J.J. (2000). Construction of a genetic toggle
switch in Escherichia coli. Nature *403*, 339–342. 10.1038/35002131.

42. McCauley, B.S., Weideman, E.P., and Hinman, V.F. (2010). A conserved gene
regulatory network subcircuit drives different developmental fates in the vegetal pole of highly
divergent echinoderm embryos. Dev Biol *340*, 200–208. 10.1016/j.ydbio.2009.11.020.

43. Davidson, E.H., Rast, J.P., Oliveri, P., Ransick, A., Calestani, C., Yuh, C.-H., Minokawa,
T., Amore, G., Hinman, V., Arenas-Mena, C., et al. (2002). A genomic regulatory network for
development. Science *295*, 1669–1678. 10.1126/science.1069883.

44. Charlebois, D.A., Balázsi, G., and Kærn, M. (2014). Coherent feedforward transcriptional
regulatory motifs enhance drug resistance. Phys Rev E Stat Nonlin Soft Matter Phys *89*,
052708. 10.1103/PhysRevE.89.052708.

45. Kim, H., Nguyen, N.-P., Turner, K., Wu, S., Gujar, A.D., Luebeck, J., Liu, J., Deshpande,
V., Rajkumar, U., Namburi, S., et al. (2020). Extrachromosomal DNA is associated with
oncogene amplification and poor outcome across multiple cancers. Nat Genet *52*, 891–897.
10.1038/s41588-020-0678-2.

46. Saghafinia, S., Mina, M., Riggi, N., Hanahan, D., and Ciriello, G. (2018). Pan-Cancer
Landscape of Aberrant DNA Methylation across Human Tumors. Cell Rep *25*, 1066-1080.e8.
10.1016/j.celrep.2018.09.082.

47. Liu, J., Lichtenberg, T., Hoadley, K.A., Poisson, L.M., Lazar, A.J., Cherniack, A.D.,
Kovatich, A.J., Benz, C.C., Levine, D.A., Lee, A.V., et al. (2018). An Integrated TCGA Pan-
Cancer Clinical Data Resource to Drive High-Quality Survival Outcome Analytics. Cell *173*,
400-416.e11. 10.1016/j.cell.2018.02.052.

48. Gibson, G. (2012). Rare and common variants: twenty arguments. Nat Rev Genet *13*,
135–145. 10.1038/nrg3118.

49. Camacho, N., Van Loo, P., Edwards, S., Kay, J.D., Matthews, L., Haase, K., Clark, J.,
Dennis, N., Thomas, S., Kremeyer, B., et al. (2017). Appraising the relevance of DNA copy
number loss and gain in prostate cancer using whole genome DNA sequence data. PLoS
Genet *13*, e1007001. 10.1371/journal.pgen.1007001.

50. Aten, J.E., Fuller, T.F., Lusis, A.J., and Horvath, S. (2008). Using genetic markers to
orient the edges in quantitative trait networks: the NEO software. BMC Syst Biol *2*, 34.
10.1186/1752-0509-2-34.

51.   Wingender, E., Schoeps, T., and Dönitz, J. (2013). TFClass: an expandable hierarchical
classification of human transcription factors. Nucleic Acids Res. *41*, D165-170.
10.1093/nar/gks1123.

52.   Ansariola, M., Megraw, M., and Koslicki, D. (2018). IndeCut evaluates performance of
network motif discovery algorithms. Bioinformatics *34*, 1514–1521.
10.1093/bioinformatics/btx798.

Figure legends

Figure 1. OncoMerge integrates PAMs, fusions, and CNAs into an integrated mutation matrix with the most suitable mutation type for each gene. The input data for OncoMerge includes the PAM, transcript fusion, and CNA matrices. OncoMerge then generates six matrices (PAM, Fusion, CNAamp, CNAdel, Act, and LoF) and uses the mutational frequency and statistical filters to determine each gene's most suitable mutation type.

Figure 2. OncoMerge inferred activating and loss of function mutations overlap significantly with prior knowledge from five independent gold standard datasets. **A.** Impact of filter sets on the number of somatically mutated genes inferred by OncoMerge in at least one cancer. **B.** Impact of filter sets on the distribution of genes per CNA locus using the same set of filtering conditions (y-axis is distributed on a log scale). The dashed line indicates the ten genes per loci cutoff that invokes the MFF filter. **C.** Enrichment of the gold standard (GS) activating (Act) or loss of function (LoF) somatic mutations with OncoMerge (OM) Act or LoF somatic mutations for each filtering condition: no filters (None); permuted q-value filter (PQ); maximum final frequency (MFF); combined PQ and MFF; and combined PQ, MFF, and microsatellite and hypermutation censoring filter (MHC). After Bonferroni multiple hypothesis correction, significant enrichments are highlighted in red (p-value ≤ 4.8×10^{-4}). The orange arrowheads indicate OM Act vs. GS Act, and the green arrowheads indicate OM LoF vs. GS LoF.

Figure 3. Summary of effect on number and frequency of somatic mutations after integrating mutation types. **A.** Frequency of hypermutation and microsatellite instability across cancers. **B.** Number and distribution of mutation types. **C.** Number of somatically mutated genes with a frequency ≥5% added after integration. **D.** Integrated somatic mutation frequencies. **E.** Increases in somatic mutation frequency relative to PAM frequency after integration.

Figure 4. Pan-cancer somatic mutations with a consistent functional impact across at least five cancers. **A.** Pan-cancer somatic mutations from the loss of functions group. **B.** Pan-cancer somatic mutations from the activating group. **C.** Prior knowledge of tumor suppressor or oncogene status for each somatically mutated gene (black square indicates known tumor suppressor or oncogene activity).

Figure 5. Demonstrating improvements in downstream SYGNAL analysis by comparing GRNs constructed with an OncoMerge integrated somatic mutation matrix versus a legacy network using only PAMs. **A.** Average degree of nodes in the PanCaner SYGNAL networks. OncoMerge = orange, legacy = yellow. **B.** Mutations per cancer network. OncoMerge = red, legacy = blue. **C.** Mutations that overlap with genes previously associated with a specific cancer in DisGeNET. OncoMerge = red, legacy = blue. **D.** TFs per cancer network. OncoMerge = green, legacy = purple. **E.** TFs that overlap with genes previously associated with a specific cancer in DisGeNET. OncoMerge = green, legacy = purple.

Figure 6. The architecture of functional disease-specific TF regulatory networks from human tumors. **A.** Active TF regulatory network construction pipeline: 1) TFs from all cancer regulatory networks were identified, 2) A putative map of TF regulatory network interactions was constructed, 3) TF → TF relationships were filtered using Pearson's correlations computed from patient tumor data, and 4) compute the triad significance profiles using mfinder. **B.** Comparison of active TF regulatory network based on SYGNAL GRNs (red) to the static TF regulatory network based on ENCODE DNA binding and accessibility (blue, Neph et al., 2012). **C.**

FANMOD enrichment normalized Z-scores for the three most enriched motifs from the active TF
regulatory network after incorporating TF regulatory interaction roles (activation or repression).
The first row, titled Coherent motifs, is shaded when the motif configuration is coherent and
white when it is incoherent. Normalized Z-scores are reported for each cancer, and diagonal
dashed lines are inserted when no Z-score was returned. The network motif can be found at the
bottom of each column, colored with regulatory roles (activation = green arrow, repression = red
perpendicular line). C1, C2, C3, C4 = coherent FFLs. I1, I2, I3, I4 = incoherent FFLs.

STAR✳Methods

Key resources table

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Existing Data Sources		
Somatic protein affecting mutations (PAMs) in TCGA	ISB Cancer Gateway in the Cloud (ISB-CGC; https://isb-cgc.appspot.com/)	https://doi.org/10.1016/j.cels.2018.03.002
Mutsig2CV for PAMs in TCGA	Broad GDAC FIREHOSE (https://gdac.broadinstitute.org)	TCGA
Somatic transcript fusions in TCGA	TumorFusions portal (https://www.tumorfusions.org/)	https://doi.org/10.1093/nar/gkx1018
Somatic copy number alterations (CNAs) in TCGA	Broad GDAC FIREHOSE (https://gdac.broadinstitute.org)	TCGA
TCGA consensus gold-standard		https://doi.org/10.1016/j.cell.2018.02.060
COSMIC Cancer Gene Census (CGC) gold-standard		https://doi.org/10.1038/s41568-018-0060-1
20/20 rule gold-standard		https://doi.org/10.1126/science.1235122
OncodriveROLE gold-standard		https://doi.org/10.1093/bioinformatics/btu467
Tokheim Ensemble gold-standard		https://doi.org/10.1073/pnas.1616440113
TF regulatory networks for 41 different cell types	http://www.regulatorynetworks.org/	https://doi.org/10.1016/j.cell.2012.04.040
Transcription Factor Target Gene	http://tfbsdb.systemsbiology.net	https://doi.org/10.1016/j.cels.2016.06.006

Database		
MSI in TCGA		https://doi.org/10.1200/po.17.00073
Hypermutation in TCGA		https://doi.org/10.1016/j.cell.2018.02.060
Deposited Data		
OncoMerge TCGA inputs	This manuscript	https://doi.org/10.6084/m9.figshare.21760964.v1
OncoMerge TCGA integrated somatic mutation matrices	This manuscript	https://doi.org/10.6084/m9.figshare.20238867.v1
Software and Algorithms		
MutSig2CV	https://doi.org/10.1038/nature12213	RRID:SCR_010779
GISTIC2.0	https://doi.org/10.1186/gb-2011-12-4-r41	RRID:SCR_000151
OncoMerge	https://github.com/plaisier-lab/OncoMerge	https://doi.org/10.5281/zenodo.5519663
cMonkey2	https://github.com/baliga-lab/cmonkey2	https://doi.org/10.1093/nar/gkv300
SYGNAL	https://github.com/plaisier-lab/sygnal	https://doi.org/10.1016/j.cels.2016.06.006
Network Edge Orienting (NEO)	https://horvath.genetics.ucla.edu/html/aten/NEO/	https://doi.org/10.1186/1752-0509-2-34
mfinder	https://www.weizmann.ac.il/mcb/UriAlon/download/network-motif-software	https://doi.org/10.1126/science.298.5594.824
FANMOD	https://www.softpedia.com/get/Network-Tools/Misc-Networking-Tools/FANMOD.shtml	https://doi.org/10.1093/bioinformatics/btl038
IndeCut	https://github.com/megrawlab/IndeCut	https://doi.org/10.1093/bioinformatics/btx798

Resource availability

Lead contact

Requests for further information should be directed to the lead contact, Christopher Plaisier (plaisier@asu.edu).

Materials availability

This study did not generate new materials

Data and code availability

This paper analyzes existing, publicly available data. All the datasets used as input for our study were deposited in Figshare (https://doi.org/10.6084/m9.figshare.21760964.v1) and they are publicly available as of the date of publication. New datasets generated from our studies were deposited in Figshare (https://doi.org/10.6084/m9.figshare.20238867.v1) and they are publicly available as of the data of publication.

All original code has been deposited at GitHub (https://github.com/plaisier-lab/OncoMerge) and is publicly available as of the date of publication.

787 Any additional information required to reanalyze the data reported in this paper is available from
788 the lead contact upon request.

Method details

Clinical and molecular data from TCGA
These studies used standardized, normalized, batch corrected, and platform-corrected multi-
omics data generated by the Pan-Cancer Atlas consortium for 11,080 participant tumors[20].
Complete multi-omic profiles were available for 9,584 patient tumors. TCGA aliquot barcodes
flagged as "do not use" or excluded by pathology review from the Pan-Cancer Atlas Consortium
were removed from the study. The overall survival (OS, OS.time) data used were obtained from
Liu et al. 2018[47].

• Somatic protein affecting mutations (PAMs) in TCGA – Somatic PAMs were identified by
the Multi-Center Mutation Calling in Multiple Cancer (MC3) project[1] and were
downloaded from the ISB Cancer Gateway in the Cloud (ISB-CGC; https://isb-
cgc.appspot.com/). PAMs were required to have a FILTER value of either: PASS, wga,
or native_wga_mix. In addition, all PAMs needed to be protein-coding by requiring that
Variant_Classification had one of the following values: Frame_Shift_Del,
Frame_Shift_Ins, In_Frame_Del, In_Frame_Ins, Missense_Mutation,
Nonsense_Mutation, Nonstop_Mutation, Splice_Site, or Translation_Start_Site.
Additionally, mutation calls were required to be made by two or more mutation callers
(NCALLERS > 1). When both normal tissue and blood were available, the blood was
used as the germline reference.
• Statistical significance of PAMs in TCGA – The likelihood that a gene is somatically
mutated by chance alone was determined using MutSig2CV[7] and downloaded for each
cancer from the Broad GDAC FIREHOSE (https://gdac.broadinstitute.org/). Genes with a
MutSig2CV False Discovery Rate (FDR) corrected p-value (q-value) less than or equal
to 0.1 were considered significantly mutated[7].
• Somatic transcript fusions in TCGA – The TumorFusions portal[2] provides a pan-cancer
analysis of tumor transcript fusions in the TCGA using the PRADA algorithm[8].
• Somatic copy number alterations (CNAs) in TCGA – Genomic regions that were
significantly amplified or deleted were identified using Genomic Identification of
Significant Targets in Cancer (GISTIC2.0)[9] and downloaded for each cancer from the
Broad GDAC FIREHOSE.

Somatic mutation data import and preprocessing
An essential first step in OncoMerge is loading up and binarizing the somatic mutation data. The
somatic mutation data comprised of four primary matrices: 1) PAMs, 2) fusions, 3) CNA
amplifications (CNAamps), and 4) CNA deletions (CNAdels) (**Figure 1**). In addition, two
derivative matrices Act and LoF are created by merging the PAM with the CNAamps or
CNAdels matrices, respectively (**Figure 1**). All files are formatted as comma-separated values
(CSV) files with genes as rows and patients as columns unless otherwise noted.

• PAM matrix - The matrix values are [0 or 1]: zero indicates the gene is not mutated in a
patient tumor, and one indicates the gene is mutated in a patient tumor.
• Fusion matrix - The matrix values are [0 or 1]: zero indicates no gene fusion in a patient
tumor, and one indicates the gene fused to another genomic locus in a patient tumor.

• CNAamp and CNAdel matrices – The all_thresholded_by_genes.csv GISTIC output file
is used to populate the CNAamp and CNAdel matrices. The all_thresholeded_by_genes
matrix values range from -2 and have no positive bound, and the values indicate the
copy number relative to the background. A cutoff of greater than or equal to 2 was used
to identify deep amplifications and less than or equal to -2 for deep deletions. Only deep
amplifications or deletions were included in these studies due to heterogeneity of cell
types and tumor biopsy purity. Oncomerge allows this threshold to be modified through a
command line parameter ('-gt' or '--gistic-threshold').
○ CNAamp matrix – The matrix values are [0 or 1]: zero indicates a gene is not
amplified in a patient tumor, and one indicates the gene is amplified in a patient
tumor.
○ CNAdel matrix – The matrix values are [0 or 1]: zero indicates a gene is not
deleted in a patient tumor, and one indicates a gene is deleted in a patient tumor.
• Act matrix – The Act matrix is the bitwise OR combination of the PAM, Fusion, and
CNAamp matrices. The Act matrix has genes as rows and patients as columns. The
matrix values are [0 or 1]: zero indicates the gene is not mutated or amplified in a patient
tumor, and one indicates the gene is either mutated, fused, amplified, or some
combination in a patient tumor.
• LoF matrix – The LoF matrix is the bitwise OR combination of the PAM, Fusion, and
CNAdel matrices. The LoF matrix has genes as rows and patients as columns. The
matrix values are [0 or 1]: zero indicates the gene is not mutated or deleted in a patient
tumor, and one indicates the gene is either mutated, fused, deleted, or some
combination in a patient tumor.

Seeding OncoMerge with putative somatic mutations
853 OncoMerge focuses on likely causal somatic mutations by considering only somatic mutations
854
that were statistically shown to be mutated more often than expected by chance alone. Likely
causal somatic mutations are also required to have a mutation frequency greater than 5%, the
definition of a common mutation[48], as this ensures sufficient patient tumors will be mutated to
power downstream analyses. These statistically significant common mutations were used as
seeds for OncoMerge integration. PAMs used as seeds were identified with MutSig2CV q-
values less than or equal to 0.1[14] and a mutation frequency greater than 5%. Gene fusions used
as seeds were identified as significant in PRADA[2,8] and had a mutation frequency greater than
5%. CNAamps or CNAdels used as seeds were identified as significantly amplified or deleted
from the amplified genes (amp_genes) or deleted genes (del_genes) GISTIC output files with
residual q-values less than or equal to 0.05[49]. CNAs from sex chromosomes (X and Y) were
excluded. Genes from sex chromosomes can enter OncoMerge as seeds from PAMs or fusions.
These seed genes become the starting point of the OncoMerge integration. Subsequent steps
determine if Act or LoF merged mutation profiles or their component PAM, Fusion, CNAamp, or
CNAdel mutation roles are the most appropriate integration model for a gene.

Merging somatic mutations in OncoMerge
869
The mutation role for each seed gene is assigned based on the frequencies of the mutation
types for a gene from the original (PAM, Fusion, CNAamp, CNADel) and merged (Act and LoF)
somatic mutation matrices and statistical thresholds for PAM (MutSig2CV) and CNAs (GISTIC).
The function g is applied to choose the mutation role for a gene using the following parameters:
f_{MMF} the minimum mutation frequency (defaults to 5%), f_{Act} the frequency of the merged Act
mutations, f_{LoF} the frequency of the merged LoF mutations, f_{PAM} the frequency of the PAM

mutations, f_{Fusion} the frequency of the gene fusion mutations, f_{CNAamp} the frequency of CNA
amplification mutation, f_{CNAdel} the frequency of the CNA deletions mutations, $qv_{MutSig2CV}$
significance of PAM mutations as MutSig2CV q-value, and qv_{GISTIC} significance of CNA
mutations as GISTIC residual q-value.

$$g\big(f_{MMF}, f_{Act}, f_{LoF}, f_{PAM}, f_{Fusion}, f_{CNAamp}, f_{CNAdel}, qv_{MutSig2CV}, qv_{GISTIC}\big)$$

$$= \begin{cases} Act, & if\ \big(f_{Act} > \max(f_{LoF}, f_{PAM}, f_{Fusion}, f_{CNAamp}, f_{CNAdel})\big) \wedge (f_{Act} > f_{MMF}) \\ LoF, & elif\ \big(f_{LoF} > \max(f_{Act}, f_{PAM}, f_{Fusion}, f_{CNAamp}, f_{CNAdel})\big) \wedge (f_{Lof} > f_{MMF}) \\ PAM, & elif\ \big(f_{PAM} \geq \max(f_{Act}, f_{LoF}, f_{Fusion}, f_{CNAamp}, f_{CNAdel})\big) \wedge (f_{PAM} > f_{MMF}) \wedge \big(qv_{MutSig2CV} \leq 0.1\big) \\ Fusion, & elif\ \big(f_{Fusion} \geq \max(f_{Act}, f_{LoF}, f_{PAM}, f_{CNAamp}, f_{CNAdel})\big) \wedge (f_{Fusion} > f_{MMF}) \\ CNAamp, & elif\ \big(f_{CNAamp} \geq \max(f_{Act}, f_{LoF}, f_{PAM}, f_{Fusion}, f_{CNAdel})\big) \wedge \big(f_{CNAamp} > f_{MMF}\big) \wedge \big(qv_{GISTIC} > 0.05\big) \\ CNAdel, & elif\ \big(f_{CNAdel} \geq \max(f_{Act}, f_{LoF}, f_{PAM}, f_{Fusion}, f_{CNAamp})\big) \wedge \big(f_{CNAdel} > f_{MMF}\big) \wedge \big(qv_{GISTIC} > 0.05\big) \end{cases}$$

The mutation role is chosen using this decision tree based on mutational frequencies and
statistical significance. The Act and LoF are first in the decision tree because the useful
integration of PAM, Fusion, CNAamp, and CNAdel and should lead to a larger mutation
frequency than any individual source mutation frequency ($f_{LoF}, f_{PAM}, f_{Fusion}, f_{CNAamp}, f_{CNAdel}$). If the
Act or LoF integrated mutation role is not chosen then the source mutation with the highest
frequency is chosen. This ordering ensures that integrated mutation roles are chosen when
possible, and that the most frequent source mutation role is otherwise chosen. The assigned
gene mutation roles are further modified by the PQ, MFF, and MHC filters to determine the final
gene mutation role.

Permuted q-value (PQ) filter
For putative Act and LoF mutations, a permuted q-value is computed by randomizing the order
of rows in the PAM, Fusion, and CNA mutation matrices' and then calculating the randomized
frequency distribution for Acts and LoFs. The observed frequency for an Act or Lof mutation is
then compared to the randomized frequency distribution to compute the permuted p-value.
Permuted p-values are corrected into q-values using the multiple-test Benjamini-Hochberg FDR-
based correction method. Only Acts or LoFs that had a permuted q-value ≤ 0.1 were retained.
Any Act or LoF with a permuted q-value > 0.1 was set to the mutation role of either PAM,
Fusion, CNAamp, or CNAdel based on which mutation role had the highest frequency. This
modifies the function g into g_{PQ} that includes the permuted q-value as a new input variable
$qv_{permuted}$, and is included as a constraint for the calls of Act and LoF.

$$g_{PQ}\big(f_{MMF}, f_{Act}, f_{LoF}, f_{PAM}, f_{Fusion}, f_{CNAamp}, f_{CNAdel}, qv_{MutSig2CV}, qv_{GISTIC}, qv_{permuted}\big)$$

$$= \begin{cases} Act, & if\ \big(f_{Act} > \max(f_{LoF}, f_{PAM}, f_{Fusion}, f_{CNAamp}, f_{CNAdel})\big) \wedge (f_{Act} > f_{MMF}) \wedge \big(qv_{permuted} \leq 0.1\big) \\ LoF, & elif\ \big(f_{LoF} > \max(f_{Act}, f_{PAM}, f_{Fusion}, f_{CNAamp}, f_{CNAdel})\big) \wedge (f_{Lof} > f_{MMF}) \wedge \big(qv_{permuted} \leq 0.1\big) \\ PAM, & elif\ \big(f_{PAM} \geq \max(f_{Act}, f_{LoF}, f_{Fusion}, f_{CNAamp}, f_{CNAdel})\big) \wedge (f_{PAM} > f_{MMF}) \wedge \big(qv_{MutSig2CV} \leq 0.1\big) \\ Fusion, & elif\ \big(f_{Fusion} \geq \max(f_{Act}, f_{LoF}, f_{PAM}, f_{CNAamp}, f_{CNAdel})\big) \wedge (f_{Fusion} > f_{MMF}) \\ CNAamp, & elif\ \big(f_{CNAamp} \geq \max(f_{Act}, f_{LoF}, f_{PAM}, f_{Fusion}, f_{CNAdel})\big) \wedge \big(f_{CNAamp} > f_{MMF}\big) \wedge \big(qv_{GISTIC} > 0.05\big) \\ CNAdel, & elif\ \big(f_{CNAdel} \geq \max(f_{Act}, f_{LoF}, f_{PAM}, f_{Fusion}, f_{CNAamp})\big) \wedge \big(f_{CNAdel} > f_{MMF}\big) \wedge \big(qv_{GISTIC} > 0.05\big) \end{cases}$$

The permuted q-value cutoff defaults to 0.1 and can be set to another value through a command
line parameter ('-pq', '--perm_qv').

Maximum final frequency (MFF) filter

The maximum final frequency (MFF) filter is a low-pass genomic filter designed to remove
passenger genes from frequently mutated CNA loci that contain many underlying genes. By
default, the filter is applied when there are 10 or more genes in a locus. Let L_{MFF} be the set of
loci with greater than 10 genes, $locus$ be defined as the set of genes in a locus, and len a
function that returns the length of a set.

$$L_{MFF} = \{\, locus \mid locus \text{ in } L, \text{ and } len(locus) \geq 10 \,\}$$

908　Let G_{locus} be the set of all n genes underlying a CNA locus.

$$locus = \{gene_1, gene_2, gene_3, \dots, gene_n\}$$

909　The first step is to define f_{MFF} the *maximum mutation frequency* from the genes of $locus$. This
910　requires using two functions: $freq$ which returns the mutation frequency for a gene, and max
911　which returns the maximum value from a set.

$$f_{MFF} = \max\left(\{freq(gene_1), freq(gene_2), freq(gene_3), \dots, freq(gene_n)\}\right)$$

912　Only the gene(s) that have a mutation frequency equal to the f_{MFF} are retained in the set
913　$locus_{MFF}$.

$$locus_{MFF} = \{\, gene \mid gene \text{ in } G_{locus}, \text{ and } freq(gene) == f_{MFF} \,\}$$

914　The genes from each $locus_{MFF}$ are used in generating the final mutation matrix. The number of
915　genes underlying a CNA locus can be set through a command line parameter ('-mlg', --
916　max_loci_genes').

Microsatellite hypermutation censoring (MHC) filter
The TCGA tumors used in this study have been characterized for both MSI[5] and hypermutation[6]
(**Supplementary Table 1**). The tumors with MSI or hypermutation are loaded as a blocklist of
patient IDs through a command line parameter ('-bl' or '--blocklist'). All tumors in the blocklist are
excluded from consideration by the PQ and MFF filters while determining the genes to include in
the final somatic mutation matrix. The mutation status for blocklist tumors are included in the
final integrated mutation matrix.

Optimizing OncoMerge filtering parameters
925　Sensitivity analysis of filtering parameters (GISTIC threshold, maximum loci genes, minimum
926　mutation frequency, and permuted q-value cutoff) for the OncoMerge algorithm was conducted
927　by varying one input parameter while fixing all others. The number of somatically mutated genes
928　and enrichment of gold standards from each parameterization of OncoMerge was evaluated to
929　determine the optimal values for each input parameter.

OncoMerge outputs
OncoMerge provides four output files that provide valuable information about the integration
process and the final integrated mutation matrix that can be used in downstream studies. Here
is a brief description of each file and its contents:
● oncoMerge_mergedMuts.csv – The integrated mutation matrix is comprised of genes
(rows) by patient tumors (columns) of mutation status after integration by OncoMerge.
The matrix values are [0 or 1]: zero indicates that the gene is not mutated in a patient
tumor, and one indicates that the gene was mutated in a patient tumor.

•   oncoMerge_CNA_loci.csv – A list of the genes mapping to each CNAamp or CNAdel
locus included in the OncoMerge integrated mutation matrix.
•   oncoMerge_ActLofPermPV.csv – List of all significant Act and LoF genes, their
OncoMerge mutation role, frequency, empirical p-value, and empirical q-value. This
output is before the application of the low-pass frequency filter.
•   oncoMerge_summaryMatrix.csv – Matrix of genes (rows) by all information gathered by
OncoMerge.

To aid in comparisons between runs, we provide the save permutation option ('-sp' or '--
save_permutation') to output permutation results so that the same permuted distribution can be
used with different parameters in separate runs. We also provide the load permutation option ('-
lp' or '--load_permutation') to load up the permuted distribution from a previous run. The
permuted distributions are saved in the following files if requested:
•   oncomerge_ampPerm.npy, oncomerge_delPerm.npy – Snapshot of the non-
deterministic permutation results from combining PAM, Fusion, and CNAamp or PAM,
Fusion, and CNAdel frequencies, respectively.

Gold standard cancer-specific gene role validation datasets
Gold standard datasets are vital to validating the usefulness of each feature in OncoMerge. Two
different sources of gold standard cancer-specific gene role (Act or LoF) datasets were used to
validate the OncoMerge predicted tumor-specific gene roles:
•   TCGA consensus: The TCGA consensus was constructed by Bailey et al., 2018
wherein they catalog a list of 299 unique oncogenesis associated genes[6]. In the TCGA
consensus 280 cancer-specific oncogene roles were identified, and 417 cancer-specific
tumor suppressor roles were identified (**Supplementary Table 2**).
•   Cancer Gene Census (CGC): The CGC was developed by Catalogue of Somatic
Mutations in Cancer (COSMIC) as an expert-curated database of human cancer-driving
genes[11]. CGC cancers were mapped to the TCGA cancers by manual curation
(Supplementary Table 2). In the CGC 205 cancer-specific oncogene roles were
identified, and 304 cancer-specific tumor suppressor roles were identified
(**Supplementary Table 2**).

Gold standard gene role validation datasets
Three different sources of gold standard gene role (Act or LoF) datasets were used to validate
the OncoMerge predicted gene roles:
•   20/20 rule: The 20/20 rule defines oncogenes (Act) by requiring >20% of mutations in
recurrent positions, and tumor suppressors (LoF) as >20% of recorded mutations are
inactivating (missense or truncating)[3]. With the 20/20 rule, 54 oncogene roles were
identified, and 71 tumor suppressor roles were identified (**Supplementary Table 2**).
•   OncodriveROLE: The OncodriveROLE is a machine learning algorithm that classifies
genes according to their role based on well-curated genomic features[13]. With
OncodriveROLE, 76 oncogene (Act) roles were identified, and 109 tumor suppressor
(LoF) roles were identified (**Supplementary Table 2**).
•   Tokheim Ensemble: Ensemble-based method from Tokheim et al., 2016[12], which
integrates MutSigCV, 20/20+, and TUSON methods for predicting gene roles. With the
Tokheim Ensemble, 78 oncogene (Act) roles were identified, and 212 tumor suppressor
(LoF) roles were identified (**Supplementary Table 2**).

Computing overlap between OncoMerge and gold standards

982 A hypergeometric enrichment statistic was used to compute the significance of overlap
983 observed between each gene role in OncoMerge versus the gold standards. When possible, the
984 tumor specificity of the gene role was taken into consideration (TCGA consensus and CGC).
985 Enrichment p-values less than the Bonferroni corrected alpha value of 0.002 were considered
986 significant.

TCGA Pan-Cancer SYstems Genetics Network AnaLysis (SYGNAL)

The mRNA and miRNA expression data required to run SYGNAL were obtained from Thorsson
et al., 2018[20]. The SYGNAL pipeline is composed of 4 steps and command-line parameters for
all programs are described in detail in Plaisier et al., 2016[19]. Each cancer was run separately
through the pipeline to reduce the confounding from tissue of origin differences. Highly
expressed genes were discovered for each cancer by requiring that genes have greater than or
equal to the median expression of all genes across all conditions in ≥ 50% of patients[19]. These
gene sets were then used as input to SYGNAL to construct the gene regulatory networks
(GRNs) for each cancer.

The underlying cMonkey2 biclustering results are identical to those from Thorsson et al., 2018[20]
as they do not rely upon genetic information. All immune-specific filters were removed for these
analyses, and all bicluster filtering was done as described in Plaisier et al., 2016[19]. Using
Network Edge Orienting (NEO)[50] somatic mutations are integrated with bicluster and regulator
expression in the next step. Two networks were constructed by applying systems genetics
analysis with NEO to the biclusters: 1) GRNs were inferred using PAM-only somatic mutation
matrices as a baseline; 2) GRNs were inferred using OncoMerge integrated somatic mutation
matrices. Importantly, the PAM-only somatic mutation matrices used were the same ones used
as input for OncoMerge.

TF regulatory network construction for PanCan-SYGNAL networks

A TF regulatory network was built for each cancer in three steps (**Figure 6A**). First, the TFs
regulating survival-associated biclusters were extracted from each cancer's SYGNAL GRN.
Second, a preliminary $TF_{regulator} \rightarrow TF_{target}$ regulatory network was constructed based on the
presence of a binding site for a putative $TF_{regulator}$ in the promoter of a TF_{target} from the
Transcription Factor Target Gene Database[19] (http://tfbsdb.systemsbiology.net). TF family
expansion[19] was used to supplement TFs that did not have an experimentally determined DNA
recognition motif in the database. The assumption was that the motifs within a TF family would
not vary significantly. Therefore TF family members from the TFClass database[51] with a known
DNA recognition motif can be used as a proxy for a TF with no known DNA recognition motif.
Finally, the putative $TF_{regulator} \rightarrow TF_{target}$ regulatory network was filtered by requiring a significant
Pearson correlation between the mRNA expression of the $TF_{regulator}$ and TF_{target} (Pearson's $|R| \geq$
0.3 and p-value ≤ 0.05). The sign of the correlation coefficient can be used to determine the role
of a regulatory interaction: a positive correlation coefficient equates to the $TF_{regulator}$ being an
activator, and a negative correlation coefficient equates to the $TF_{regulator}$ being a repressor.
Networks with fewer than 50 interactions were not included in the analyses as they were not
sufficiently powered to run the network motif analysis. The cancer regulatory networks for
DLBC, KICH, KIRP, OV, TGCT, and THYM were excluded from further studies.

TF regulatory network motif analysis

Three-node network motifs were enumerated from the TF regulatory networks using mfinder[25] in
the same manner as Neph et al., 2012[23] and used to compute triad significance profiles
(TSPs)[24]. The parameters used with mfinder v1.20 were[23]: motif size set at 3 (-s 3), requested
250 random networks to be generated (-r 250), and the Z-score threshold was set at -2000 to
ensure all motifs are reported (-z -2000). All Z-scores were extracted for each cancer and
converted to triad significance profiles using the methods of Milo et al., 2004[24].
1033
1034   For consistency, the TF regulatory networks for the 41 different cell types from Neph et al., 2012
1035   were downloaded from http://www.regulatorynetworks.org/ and analyzed using the same
1036   approach described above.
1037
Signed network motif analysis incorporating TF regulator interaction roles
The enrichment of signed feed-forward loops (FFLs), regulated feedback, and regulating
feedback network motifs was computed using FANMOD[26], which takes into consideration TF
regulatory roles (activation and repression). The command line version of FANMOD from
IndeCut[52] was used with default parameters, except for the inclusion of regulatory role (colored
edges)[26] (fanmod 3 100000 1 <input_file> 1 0 1 2 0 1 0 1000 3 3 <output_file> 1 1). Z-scores for
signed FFLs, regulated feedback, and regulating feedback network motifs were extracted for
each cancer and converted to triad significance profiles using the methods of Milo et al., 2004[24].
The signed FFL network motifs are broken down into C1, C2, C3, C4, I1, I2, I3, and I4, as
described previously[39].
1048

Quantification and statistical analysis

A nominal alpha value (p-value or q-vaule cutoff) of 0.05 was used unless otherwise stated.
Statistical analyses are described in detail in the methods sections where they were used, and
we provide a brief synopsis of the statistical methods below. Hypergeometric enrichment
analysis was used to identify significant overlaps of OncoMerge-derived gene sets with gold-
standards gene sets. When appropriate for the gold-standard analyses, Benjamini-Hochberg
FDR multiple hypothesis correction was applied to the hypergeometric p-values. Cutoff were as
describe in the methods or results. Permuted p-values were computed for each integrated
somatic mutation and Benjamini-Hochberg FDR multiple hypothesis correction was applied to
generate permuted q-values. Pearson correlations were used to compare two sets of
quantitative values (e.g., number of somatic mutations and MSI/hypermutation frequency) and
the correlation coefficient (R) and p-value are reported. Triad significance profiles (TSPs) were
used to quantify the enrichment of three node network motifs.

Figures

 Figure 1.

Figure 2.

Figure 3.

Figure 4.

1072   **Figure 5.**

Figure 6.