Nanoscale phosphoinositide distribution on cell membranes of mouse

cerebellar neurons

Authors

Kohgaku Eguchi[1*] and Ryuichi Shigemoto[1*]

[1]Institute of Science and Technology Austria (ISTA), Am Campus 1, 3400 Klosterneuburg,

Austria

[*]For correspondence: kohgaku.eguchi@ist.ac.at (KE); ryuichi.shigemoto@ist.ac.at (RS)

Abstract

Phosphatidylinositol-4,5-bisphosphate (PI(4,5)P$_2$) plays an essential role in neuronal

activities through interaction with various proteins involved in signaling at membranes.

However, the distribution pattern of PI(4,5)P$_2$ and the co-clustering with these proteins on the

neuronal cell membranes remain elusive at the electron microscopic level. In this study, we

quantitatively investigated the nanoscale distribution of PI(4,5)P$_2$ on the neuronal cell

membranes by SDS-digested freeze-fracture replica labeling with cryo-fixed mouse

cerebellum. We demonstrate that PI(4,5)P$_2$ makes clusters with a mean size of ~ 1,000 nm^2

and these clusters show preferential accumulation in specific membrane compartments

depending on cell types. Purkinje cell (PC) spines and granule cell (GC) presynaptic active

zones are particularly rich in the PI(4,5)P$_2$ clusters. Furthermore, these clusters are

extensively associated with clusters of Ca$_V$2.1 and GIRK3 throughout different membrane

compartments of PCs, GCs, and molecular layer interneurons. In contrast, these clusters

showed compartment-specific association with mGluR1α in PC spines. These results suggest

that visualization of the nanoscale PI(4,5)P$_2$ distribution may provide insight into the

physiological functions of PI(4,5)P$_2$ in neurons.

Introduction

Phosphoinositides (PIs) are minor components on the cytoplasmic side of eukaryotic cell membranes, but they play essential roles in a wide variety of cellular functions. In neuronal cells, each stereoisomer of PIs is distributed in different subcellular compartments (Haucke, 2005; Idevall-Hagren and De Camilli, 2015; Wenk and De Camilli, 2004): PI(4)P is enriched in the membrane of the Golgi apparatus and synaptic vesicles (SVs), PI(4,5)P$_2$ and PI(3,4,5)P$_3$ mainly exist in the plasma membrane, PI(3)P and PI(3,5)P$_2$ are selectively concentrated on early and late endosomes, respectively. PIs contribute to various aspects of neuronal activity, such as synaptic transmission and maintenance of membrane excitability by regulating ion channels and intracellular signaling pathways. At chemical synapses, PIs regulate exo- and endocytosis of synaptic vesicles at the presynaptic sites (Lei et al., 2017; Martin, 2001; Posor et al., 2015; Ueda and Hayashi, 2013)Many proteins related to SV exo- and endocytosis have binding domains to one or several stereoisomers of PIs (Di Paolo and De Camilli, 2006; Falkenburger et al., 2010), indicating a potential role of PIs as an anchor for these presynaptic proteins to regulate their localization and dynamics during synaptic transmission. At the postsynaptic sites, PIs regulate the morphological plasticity of dendritic spines through actin remodeling (Lei et al., 2017; Martin, 2001; Posor et al., 2015; Ueda and Hayashi, 2013)during long-term potentiation (Lei et al., 2017; Martin, 2001; Posor et al., 2015; Ueda and Hayashi, 2013).

Although it is essential to know the distribution pattern of PIs and their association with signaling proteins at neuronal cell membranes to understand the roles of PIs in neuronal activities, it has been poorly investigated due to several technical issues in the conventional methods (Tsuji et al., 2019). Immunoelectron microscopy using anti-PI(4,5)P$_2$ antibody allows for observing the nanoscale PI(4,5)P$_2$ distribution on the cell membrane, but the use of aldehyde-fixed cell preparations has a potential issue because aldehydes cannot fix the lateral diffusion of most membrane lipids across the membrane (Tanaka et al., 2010). Fluorescent probes for PIs based on a PI-binding domain (PBD) of proteins have been developed to visualize PIs in living cells (Idevall-Hagren and De Camilli, 2015; Maekawa and Fairn, 2014). However, this method does not have sufficient spatial resolution to observe the nanoscale distribution of PIs in small compartments in neuronal cell membranes, such as presynaptic active zones (AZs) and postsynaptic densities (PSDs). In addition, the overexpressed PBD-based probes mask PIs and competitively interfere with their interactions

with proteins in the living cells (Suh and Hille, 2008).

To solve these issues and visualize the nanoscale distribution of PIs on cell membranes,

sodium dodecyl sulfate-digested freeze-fracture replica labeling (SDS-FRL) combined with

cryofixation has been utilized (Aktar et al., 2017; Cheng et al., 2014; Fujita et al., 2009;

Ozato-Sakurai et al., 2011; Tsuji et al., 2019). This immunoelectron microscopic method

enables nanoscale membrane phospholipid labeling by physically fixing membrane

molecules without chemical fixation. In this study, we optimized this SDS-FRL method to

acute mouse cerebellar slices and investigated the nanoscale distribution of $PI(4,5)P_2$ on

neuronal cell membranes using recombinant GST-tagged pleckstrin homology (PH) domain

of phospholipase Cδ1 (PLCδ1) as a specific probe for $PI(4,5)P_2$. This approach allowed us to

quantitatively examine the numbers, densities, and distribution patterns of $PI(4,5)P_2$ on

somatodendritic and axonal membranes, including post- and presynaptic sites in mouse

cerebellar neurons. We show that $PI(4,5)P_2$ makes small clusters on neuronal membranes and

specifically co-clusters with P/Q-type voltage-gated calcium channels, G-protein-coupled

inwardly rectifying potassium channels, and metabotropic glutamate receptors in distinct

membrane compartments, giving insights into the physiological functions of $PI(4,5)P_2$ in the

regulation of neuronal excitability and neurotransmitter release.

78   Results

79   Visualization of nanoscale $PI(4,5)P_2$ distribution on SDS-digested freeze-fracture replica

80   preparations of mouse brain tissues

To observe the nanoscale two-dimensional distribution of $PI(4,5)P_2$, SDS-FRL has been

utilized on cultured human fibroblasts, mouse smooth muscle cells, and rat pancreatic

exocrine acinar cells (Fujita et al., 2009; Ozato-Sakurai et al., 2011). In this study, we

optimized this method for mouse cerebellar tissues to quantitatively analyze the nanoscale

$PI(4,5)P_2$ distribution on neuronal cell membranes. We used a recombinant GST-tagged PH

domain of PLCδ1 (GST-PH) as a specific probe of $PI(4,5)P_2$. To verify the specificity of

GST-PH among the PI stereoisomers, we labeled freeze-fracture replicas of liposomes

containing either PhdIns or a PI stereoisomer (5 mol%) with GST-PH (50 ng/ml), anti-GST

primary antibody, and gold particle-conjugated secondary antibody. The density of

immunogold particles was much higher on the replica prepared from PI(4,5)P$_2$ liposome

compared to those from other liposomes (Figure 1A), indicating a high specificity of the

PI(4,5)P$_2$ probe (Figure 1B).

To visualize the two-dimensional (2D) distribution of PI(4,5)P$_2$ on neuronal cell membranes,

we labeled PI(4,5)P$_2$ in the mouse cerebellum using SDS-FRL with GST-PH. Acute slices

prepared at physiological temperature (Eguchi et al., 2020) were frozen under high pressure

with 15% PVP as a cryoprotectant (Figure 1C) to minimize damage caused by ice crystals

during freezing (Borges-Merjane et al., 2020). After the SDS-digestion to remove cytosolic

proteins and extracellular matrix, cerebellar replicas were sequentially incubated with GST-

PH, primary antibodies, and gold particle-conjugated secondary antibodies. Ca$_V$2.1 was co-

labeled using an anti-Ca$_V$2.1 antibody directed against its intracellular domain (Althof et al.,

2015), giving specific labeling on the cytoplasmic leaflets (P-face) of neuronal cell

membranes. Immunogold particles that label PI(4,5)P$_2$ by GST-PH (PI(4,5)P$_2$ particles)

showed higher density on the P-face of the somatic membrane of Purkinje cells (PCs) than on

the E-face (Chi-square likelihood-ratio test (Chi-LRT): $p < 0.001$; Figures 1D and 1E),

indicating the dominance of PI(4,5)P$_2$ distribution on the cytoplasmic leaflet of the cell

membranes. The particle density without GST-PH was significantly lower on both E- and P-

face of the PC somatic membranes than that with GST-PH (Figures S1A and S1B), indicating

that even the low-density PI(4,5)P$_2$ labeling on the E-face is ascribable to GST-PH binding.

To quantitatively investigate whether PI(4,5)P$_2$ is clustered on somatic membranes of PCs,

we performed Monte-Caro random simulations and compared the nearest neighbor distance

(NND) between the observed and simulated PI(4,5)P$_2$ particles from all images (Szoboszlay

et al., 2017). The mean NNDs of the observed PI(4,5)P$_2$ particles were around half of the

simulated one (Chi-LRT: $p < 0.001$, Figure 1F), indicating clustering of PI(4,5)P$_2$ on the

membrane. The distribution of NNDs between PI(4,5)P$_2$ particles obtained from a PC somatic

membrane (Figure 1G) is highly right-skewed (skewness = 3.55). Fitting this distribution

with Gaussian mixture modeling shows that the NNDs between PI(4,5)P$_2$ particles are

produced from a mixture of short and long NND populations. These results indicate that

PI(4,5)P$_2$ on the PC soma has two distinct distribution patterns, clustered and scattered, and

that most PI(4,5)P$_2$ constitute clusters (Figure 1G). The PI(4,5)P$_2$ particle clusters were also

observed on the E-face of the somatic membrane (Figure S1C). The number of particles in a

cluster was significantly lower without GST-PH than that with GST-PH on both E- and P-

face (Figure S1C). These results indicate that PI(4,5)P$_2$ forms clusters on both the outer and

inner leaflets of the somatic membranes.

To examine whether the density of PI(4,5)P$_2$ differs between neuronal cell types in the

cerebellum, we observed somatic membranes of cerebellar granule cells (GCs) and molecular

layer interneurons (MLIs). PI(4,5)P$_2$ particles were also distributed on the P-face making

clusters in somatic membranes of both cell types (Figure 1H). Although the density of

PI(4,5)P$_2$ particles in the somatic membrane of GCs was significantly higher than that of

MLIs (multiple pairwise comparisons with Tukey adjustment (Tukey): $p < 0.001$; Figure 1I),

the NNDs of the PI(4,5)P$_2$ particles on somatic membranes of the cerebellar neurons did not

differ between the cell types (Chi-LRT: $p = 0.12$; Figure 1J), indicating a similar local

concentration of PI(4,5)P$_2$ in these clusters across the examined cerebellar neuronal cell

types.

PI(4,5)P$_2$ distribution on somatodendritic compartments of PCs

We next investigate whether the distribution of PI(4,5)P$_2$ differs among somatodendritic

compartments of PCs: somata, main shafts (MS), smooth branchlets (SmB), spiny branchlets

(SpB), and spines. EM pictures of each dendritic component were taken in the molecular

layer (ML) divided into proximal (MS), intermediate (SmB) and distal one-third (SpB,

spines) of the PC layer. Gold particles for PI(4,5)P$_2$ were observed throughout all the

dendritic compartments (Figure 2A). The density of PI(4,5)P$_2$ particles was found to increase

gradually from the proximal to the distal dendrites (Chi-LRT: $p < 0.001$; Figure 2B): spine

membranes showed a ~1.5 times higher density of PI(4,5)P$_2$ particles than somatic, MS, and

SmB membranes (Tukey: $p < 0.001$), whereas no significant difference compared to SpB

membranes was detected (Tukey: $p = 0.43$). Mean NNDs of PI(4,5)P$_2$ particles were not

significantly different between the compartments (Figure 2B, Chi-LRT: $p = 0.55$), indicating

that the local concentration of PI(4,5)P$_2$ does not differ between different somatodendritic

compartments of PCs.

PI(4,5)P$_2$ particles on the dendritic membranes formed clusters as seen on the somatic

membranes. To quantitatively compare the PI(4,5)P$_2$ cluster profiles (cluster area, particle

number in cluster, and intra-cluster density) between all compartments of somatodendritic

membranes, we detected the clusters using the Density-Based Spatial Clustering of

Applications with Noise (DBSCAN) algorithm (Methods, (Szoboszlay et al., 2017). The

DBSCAN detected variable sizes of $PI(4,5)P_2$ particle clusters on the somatodendritic

membranes (Figure 2C). The density of $PI(4,5)P_2$ clusters on the PC cell membranes

gradually increased from the proximal to the distal compartments (Chi-LRT: $p < 0.001$;

Figure 2D); spine membranes showed a higher density of $PI(4,5)P_2$ clusters than somatic,

MS, and SmB membranes (Tukey: $p < 0.001$), whereas no significant difference with SpB

membranes was detected (Tukey: $p = 0.21$). The cluster area was not significantly different

between all compartments of somatodendritic membranes (Chi-LRT: $p = 0.10$; Figure 2D),

whereas intra-cluster $PI(4,5)P_2$ particle density in spines was significantly higher than those

in other compartments (Chi-LRT: $p < 0.001$; Figure 2D). These results suggest that $PI(4,5)P_2$

are distributed throughout all compartments of somatodendritic membranes of PCs, and the

density is higher at the distal dendrites and spines than at the soma and proximal dendrites.

This proximo-distal gradient of $PI(4,5)P_2$ along the somatodendritic compartments of PC

membranes is ascribable to an increase in the density of $PI(4,5)P_2$ clusters rather than the

cluster size, and a higher $PI(4,5)P_2$ density within clusters on spine membranes.

PI(4,5)P₂ distribution on membrane compartments of GCs and MLIs

$PI(4,5)P_2$ distribution on membrane compartments of GCs and MLIs

We next examined the $PI(4,5)P_2$ distributions in different compartments of GCs (somata,

dendrites, parallel fiber (PF) axons, presynaptic PF boutons with PC spines (PF-PC) or MLI

dendrites (PF-MLI)) and MLIs (somata, dendrites, presynaptic basket cell boutons on PC

somata (BC-PC)). Gold particles for $PI(4,5)P_2$ were observed throughout all compartments of

GCs and MLIs (Figures 3A and 3B). In the GC membranes, the density of $PI(4,5)P_2$ particles

was not significantly different between the compartments (Chi-LRT: $p = 0.095$; Figure 3C),

though the mean NNDs of $PI(4,5)P_2$ particles were slightly but significantly different

between somatic and axonal membranes (Tukey: $p = 0.027$, Figure 3C). The cluster

parameters of $PI(4,5)P_2$ were also not significantly different between the compartments

(Figure 3D). These results suggest that $PI(4,5)P_2$ in GC membranes shows a similar

distribution pattern throughout the subcellular compartments.

In MLIs, the $PI(4,5)P_2$ density was not significantly different between the membrane

compartments, but the mean NNDs were larger in dendritic and presynaptic bouton

membranes than in somatic membranes (Figure 3E). This difference is caused by larger

population of scattered $PI(4,5)P_2$ in these compartments as shown in Figure 3 – Figure

supplement 1. The intra-cluster density of PI(4,5)P$_2$ in the bouton membrane was

significantly higher than in the somatic membrane, whereas the cluster density and area were

similar between the compartments (Figure 3F).

189   PI(4,5)P$_2$ distribution on synaptic membranes

Next, we focused on PI(4,5)P$_2$ distribution in the presynaptic membrane of PF (Figure 4A).

Since many presynaptic proteins related to neurotransmitter release contain PI(4,5)P$_2$-binding

domains (Martin, 2001), PI(4,5)P$_2$ is expected to be localized at the presynaptic active zones

(AZs). The PI(4,5)P$_2$ particle density was significantly higher in the AZs than in the whole

bouton and extra-AZ membranes at both PF-PC and PF-MLI synapses (Tukey: $p < 0.001$;

Figure 4B). The observed density of PI(4,5)P$_2$ particles in AZs was significantly higher than

that of randomly distributed PI(4,5)P$_2$ particles on the bouton membrane by Monte-Carlo

simulation (Chi-LRT: $p < 0.001$ (PF-PC) and $p = 0.002$ (PF-MLI); Figure 4C). Next, we

examined how the PI(4,5)P$_2$ particles are distributed within the AZs using the center-

periphery index (CPI), where a CPI of 0 indicates that the particle is at the center of gravity,

and a CPI of 1 indicates the particle is at the edge of the AZ (Kleindienst et al., 2020). The

histogram of CPI in the bouton membranes suggests that the particles are distributed at a high

density within the AZ and that the distribution probability decreases with distance from the

AZ (Figure 4D, top). The CPI of the particles within AZs was higher at around CPI = 1.0

compared to the values of the random simulation (Figure 4D, bottom), indicating that

PI(4,5)P$_2$ is preferentially located in the periphery of the AZs. The mean CPI of the PI(4,5)P$_2$

particles in AZs was significantly higher than that of randomly distributed particles (Chi-

LRT: $p < 0.001$, Figure 4E) in both PF-PC and PF-MLI synapses. These results suggest that

AZ, in particular, its periphery is PI(4,5)P$_2$-enriched regardless of postsynaptic cell types.

Next, we investigated the PI(4,5)P$_2$ distribution in PSD of dendritic spines. The PI(4,5)P$_2$

particles were distributed throughout the spine membranes (Figure 4F), and the density in the

PSDs was not significantly different compared to the whole spine and extra-PSD membranes

(Chi-LRT: $p = 0.77$; Figure 4G). The observed PI(4,5)P$_2$ particle density in PSDs was not

significantly higher than that of the particles randomly distributed in the spine (Chi-LRT: $p =$

0.25; Figure 4H). Furthermore, the CPI of the PI(4,5)P$_2$ particles in the spines was uniformly

distributed, and the CPI in the PSDs was similar to the simulated particles randomly

distributed (Figure 4I). These results suggest that $PI(4,5)P_2$ is not specifically accumulated in

the postsynaptic site of PC dendritic spines.

The association of ion channels and receptors with $PI(4,5)P_2$ on cell membranes of

cerebellar neurons

The association of $Ca_V2.1$ with $PI(4,5)P_2$

$Ca_V2.1$ is an α-subunit of P/Q-type voltage-gated calcium channels and is regulated by

$PI(4,5)P_2$ directly or indirectly through Cav β-subunit (Suh et al., 2012; Suh and Hille, 2008).

Here, we examined whether $Ca_V2.1$ is associated with $PI(4,5)P_2$ in the neuronal membranes

by co-immunolabeling of $Ca_V2.1$ with the $PI(4,5)P_2$ labeling. To assess the association, we

compared NNDs from $Ca_V2.1$ to $PI(4,5)P_2$ particles (NND_{C-P}) between observed and

simulated $Ca_V2.1$ particles. To reproduce the $Ca_V2.1$ clustering in the simulation, we

performed a fitted simulation of $Ca_V2.1$ particle distribution (Methods, (Kleindienst et al.,

2020; Luján et al., 2018b)). In PCs, $Ca_V2.1$ is broadly expressed in somatodendritic

membranes (Figure 5A) as previously reported (Indriati et al, 2013). The real values of

NND_{C-P} in somatic, SpB, and spine membranes were significantly shorter than the simulated

ones (Figure 5B). The mean NND_{C-P} was not significantly different between the

compartments of PC membranes (Figure 5C). In GCs, $Ca_V2.1$ was highly expressed in

somatic and presynaptic AZ membranes (Figure 5D). The comparison of NND_{C-P} between

observed and simulated $Ca_V2.1$ particles showed significant differences in the somatic and

PF-PC/MLI AZ membranes (Figure 5E), indicating the association of $Ca_V2.1$ with $PI(4,5)P_2$.

The mean NND_{C-P} values in PF-PC and PF-MLI AZs were not significantly different (Figure

5F). In MLI, $Ca_V2.1$ was expressed in somatic and presynaptic basket cell (BC)-PC bouton

membranes (Figure 6A). The NND_{C-P} in somatic and BC-PC bouton membranes was

significantly shorter than the simulated one (Figure 6B). These results suggest the ubiquitous

association of $Ca_V2.1$ with $PI(4,5)P_2$ in neuronal cell membranes across various cell types

and their compartments.

 The association of GIRK3 channels with PI(4,5)P$_2$

PI(4,5)P$_2$ is associated with some ion channels and regulates their activities, e.g., open

channel probability (Suh and Hille, 2008). G protein-coupled inwardly rectifying K$^+$ (GIRK)

channels are a family of lipid-gated potassium channels that are activated by PI(4,5)P$_2$ and G-

protein βγ-subunits (G$_{βγ}$) released from G-protein coupled receptors (GPCRs) (Whorton and

MacKinnon, 2011). Thus, GIRK channels are expected to be associated with PI(4,5)P$_2$

clusters in order to be efficiently activated. To examine whether PI(4,5)P$_2$ and GIRK

channels are co-clustered on dendritic membranes of mouse cerebellar PCs, we co-labeled

PI(4,5)P$_2$ and GIRK3, the most predominant subunit of GIRK channels in PCs (Aguado et al.,

2008; Fernández-Alacid et al., 2009), on the P-face of the dendritic membranes. The GIRK3

particles were observed throughout the dendritic SpBs and spines of the PCs (Figure 7A). We

compared the NNDs from GIRK3 to PI(4,5)P$_2$ particles (NND$_{G-P}$) between observed and

simulated GIRK3 particles in spines and SpBs spines of PCs. The real NND$_{G-P}$ in both spines

and SpBs was significantly shorter than the simulated one (Chi-LRT: $p < 0.001$, Figure 7B),

suggesting that GIRK3 channels co-clustered with PI(4,5)P$_2$ in the distal dendritic

membranes of PCs. GIRK channels are also expressed on presynaptic PF-PC boutons,

including AZs (Figure 7A, (Fernández-Alacid et al., 2009; Luján et al., 2018b), and may

regulate presynaptic excitability. In the presynaptic AZs, the real NND$_{G-P}$ was significantly

shorter than the simulated one (Chi-LRT: $p < 0.001$, Figure 7B), indicating the coexistence of

GIRK3 with PI(4,5)P$_2$. The mean NND$_{G-P}$ in spine and AZ membranes was significantly

shorter than that in SpB membranes (Figure 7C), suggesting the tighter association of GIRK3

and PI(4,5)P$_2$ in synaptic membranes.

The association of mGluR1α with PI(4,5)P$_2$

Group I metabotropic glutamate receptors (mGluRs) are GPCRs coupled with Gα$_q$ subunit

and hydrolyze PI(4,5)P$_2$ into inositol 1,4,5-trisphosphate (IP$_3$) and diacylglycerol (DAG)

through the activation of PLCβ. A subtype of Group I mGluRs mGluR1α is highly expressed

on the dendritic shafts and spines but avoids PSDs in PCs, and forms clusters (Luján et al.,

2018a). To effectively produce the second messengers from PI(4,5)P$_2$, a mGluR1α cluster is

expected to be located close to the PI(4,5)P$_2$ clusters on the dendritic membrane of PCs. To

address this possibility, we co-labeled mGluR1α with PI(4,5)P$_2$ on the dendritic membranes

of PCs and compared the NNDs between mGluR1α and PI(4,5)P$_2$ particles (NND$_{M-P}$) with the fitted simulation of the mGluR1α particle distribution. The mGluR1α particles were distributed as clusters on the membranes of spines and SpBs (Figure 7D-E) as previously reported (Luján et al., 2018a). The real NND$_{M-P}$ was significantly shorter than the simulated one on the spine membrane (Chi-LRT: p < 0.001) but not on the SpB membrane (Chi-LRT: p = 0.36, Figure 7F). These results suggest compartment-specific co-clustering of mGluR1α with PI(4,5)P$_2$ on PC spines.

Discussion

Visualization of PI(4,5)P$_2$ distribution on neuronal cell membranes using SDS-FRL

To understand how and where PI(4,5)P$_2$ exerts the physiological functions in neurons, it is essential to know its nanoscale distribution in the neuronal cell membranes. To visualize the distribution of PI(4,5)P$_2$ on the cell membrane, cryofixation of phospholipids is necessary because they diffuse laterally even after chemical fixation (Tanaka et al., 2010). Detergents and organic solvents used in conventional immunoEM methods disrupt the cell membrane and should be avoided. In regions with densely packed proteins such as AZs and PSDs, limited accessibility of the probes could hamper quantitative analysis of the PI(4,5)P$_2$ distribution. The fluorescent protein-tagged PH domain of PLCδ1 has been developed as a specific probe for PI(4,5)P$_2$ (Idevall-Hagren and De Camilli, 2015). This probe allows us to perform single-molecule imaging of PI(4,5)P$_2$ on the plasma membrane of the cultured cells using super-resolution microscopies such as Stimulated Emission Depletion (STED) microscopy and Direct Stochastic Optical Reconstruction Microscopy (dSTORM) (Milovanovic et al., 2016; Wang and Richards, 2012). However, it is still difficult to observe the PI(4,5)P$_2$ distribution on neuronal cell membranes in brain tissues because of the limited accessibility of the probes in AZs and PSDs. In addition, overexpressed fluorescent probes interfere with endogenous protein binding to PI(4,5)P$_2$ by masking PI(4,5)P$_2$ on the cell membrane (Suh and Hille, 2008), making it difficult to quantify the changes in PI(4,5)P$_2$. Thus, the SDS-FRL with cryo-fixed brain tissue used in this study has the advantage of visualizing accurately the native PI(4,5)P$_2$ distribution. In this method, phospholipids are physically immobilized by high-pressure freezing and replication by deposition of evaporated carbon/platinum, so the phospholipids do not diffuse after freezing. Since cytosolic proteins

are removed by SDS treatment, $PI(4,5)P_2$ and transmembrane proteins are exposed on the

replica surface, making high accessibility of the probes and antibodies. Due to a low temporal

resolution, it is still challenging to examine the dynamics of $PI(4,5)P_2$ during neuronal

activity, but it is a beneficial method to study the nanoscale distribution of $PI(4,5)P_2$ and

other phospholipids on neuronal cell membranes.

This study demonstrates that $PI(4,5)P_2$ is broadly distributed on neuronal cell membranes in

the mouse cerebellum. Comparisons of the $PI(4,5)P_2$ distribution between cell types revealed

slightly different densities of $PI(4,5)P_2$ particles on the somatic membrane, but the differences

in distribution patterns between subcellular compartments were more pronounced; in PCs, the

$PI(4,5)P_2$ density was higher at the distal dendrites than at the somatic and proximal

dendrites; in GCs, the density was similar at the somata, dendrites, axons, and presynaptic

boutons; in MLIs, the density of $PI(4,5)P_2$ in clusters was lower in presynaptic bouton

membranes than somatic and dendritic membranes. These differences in the $PI(4,5)P_2$

distribution patterns may reflect different roles of $PI(4,5)P_2$ in cell functions between

different cell types.

$PI(4,5)P_2$ makes clusters on neuronal cell membranes in mouse cerebellum

Previous studies using super-resolution microscopy have demonstrated that $PI(4,5)P_2$ forms

clusters of 50-100 nm in diameter on the cell membrane, as shown in PC12 cells (Aoyagi et

al., 2005; van den Bogaart et al., 2011; Wang and Richards, 2012) and mouse myoblast cells

(Petersen et al., 2016). Our study reveals that $PI(4,5)P_2$ forms clusters of around 1,000 nm^2

(corresponding to 35 nm in diameter) on the cell membrane of neurons in the acute cerebellar

tissues. The area of the clusters showed almost the same mean values in the somatodendritic

and axonal membranes of different neuronal cell types in the cerebellum. The cluster density

and the intra-cluster particle density of $PI(4,5)P_2$ in the PC spine membrane are higher than in

other somatodendritic compartments. These observations suggest the accumulation of

densely-packed $PI(4,5)P_2$ clusters in the spine membranes, supporting the contribution of

$PI(4,5)P_2$ and other PIs to spine formation and morphological long-term plasticity (Lei et al.,

2017; Ueda and Hayashi, 2013). Although it is still unknown why $PI(4,5)P_2$ forms clusters,

the clustered $PI(4,5)P_2$ localization indicates the importance of a close spatial relationship

between the clusters and effector proteins for their effective interaction, as discussed in the

following section.

PI(4,5)P$_2$ distribution in the synaptic membrane

In this study, we found the accumulation of PI(4,5)P$_2$ in AZ membranes of the presynaptic PF

boutons regardless of the postsynaptic cell types. This result suggests the roles of PI(4,5)P$_2$

on presynaptic activities, including vesicle exocytosis and endocytosis. At the presynaptic

membrane, Rab6-interacting proteins (RIMs) have PI(4,5)P$_2$-binding sites at C2A and C2B

domains, and their interactions are essential for efficient exocytosis (de Jong et al., 2018).

PI(4,5)P$_2$ also interacts with a SNARE protein syntaxin-1A and an SV Ca^{2+} sensor protein

synaptotagmin-1 (Aoyagi et al., 2005; Honigmann et al., 2013; Milovanovic et al., 2016).

Since Ca$_V$2.1 directly binds to RIMs and dominantly induces Ca^{2+} influx into PF boutons, the

association of Ca$_V$2.1 with PI(4,5)P$_2$ supports the idea that PI(4,5)P$_2$ anchors SVs and Ca^{2+}

channels through interaction with these proteins. PI(4,5)P$_2$ also regulates vesicle endocytosis

via interactions with endocytic proteins e.g., AP-2, AP180, clathrin, and dynamin (Haucke,

2005; Martin, 2001; Posor et al., 2015). At calyx of Held synapses of rat brainstem,

upregulation of PI(4,5)P$_2$ by the retrograde signals through the nitric oxide/cyclic GMP-

activated protein kinase/RhoA/Rho-associated protein kinase pathway accelerates vesicle

endocytosis, which strengthens the homeostatic plasticity for the maintenance of high-

frequency synaptic transmission (Eguchi et al., 2012; Taoufiq et al., 2013). PI(4,5)P$_2$ clusters

may work as anchors of presynaptic proteins involved in this pathway providing a site for

their interactions.

Co-clustering of PI(4,5)P$_2$ and ion channels and receptors

PI(4,5)P$_2$ regulates ion channels such as inwardly rectifying K$^+$ channels, KCNQ channels,

transient receptor potential (TRP) channels, and voltage-gated Ca^{2+} channels (Suh and Hille,

2008). Therefore, clarifying the distance relationship between these ion channels and

PI(4,5)P$_2$ may provide insight into the regulation mechanism of neuronal excitability by

PI(4,5)P$_2$. In this study, we found that GIRK3 was associated with PI(4,5)P$_2$ in SpB and

spines of PCs and also AZs of PF boutons. Because PI(4,5)P$_2$ is essential to activate GIRK

channels by βγ subunit of G protein (Whorton and MacKinnon, 2011), the association

between GIRK3 and PI(4,5)P_2 suggests that GIRK3 on SpB and spines is constantly ready to

be activated by G-protein coupled receptors. Since GIRK channels are also associated with

GABA$_B$ receptors at different compartments of the cerebellar neurons, such as dendritic

shafts and spines of PCs and AZs of PF boutons (Fernández-Alacid et al., 2009; Luján et al.,

2018b), the co-assembly of PI(4,5)P_2, GIRK channels and GABA$_B$ receptors may contribute

to effective inhibitory postsynaptic transmission on Purkinje cells (Luján et al., 2018b).

Group I mGluRs are GPCRs coupled with Gα_q subunit and hydrolyze PI(4,5)P_2 into inositol

1,4,5-trisphosphate (IP$_3$) and diacylglycerol (DAG) through the activation of PLCβ. PCs

express mGluR1α, a subtype of Group I mGluRs, on the dendritic shafts and spines

concentrated in the peri-PSD areas (Luján et al., 2018a). We found a significant association

of mGluR1α with PI(4,5)P_2 clusters specific to the spine membrane in PCs, which may

contribute to the effective production of the second messengers from PI(4,5)P_2 in response to

the glutamate release from presynaptic PF boutons. On the other hand, no particular

concentration of PI(4,5)P_2 has been found in the peri-PSD areas. The diffusion dynamics of

PI(4,5)P_2 on the spine membrane may be enough to support the interaction with mGluR1α.

Thus, the tight spatial association of PI(4,5)P_2 as observed for Ca$_V$2.1 and GIRK3 may not be

necessary for the coupling of mGluR1α activation and intracellular signal cascade, though the

distance between mGluR1α (and the coupled PLCβ) and PI(4,5)P_2 may affect the kinetics of

the second messenger (IP$_3$ and DAG) production from PI(4,5)P_2.

The present study demonstrated that the SDS-FRL method could be successfully applied to

neuronal cells in acute mouse brain tissues to visualize the distribution of PI(4,5)P_2. We

quantitatively analyzed the nanoscale two-dimensional distribution of PI(4,5)P_2 on the cell

membrane of mouse cerebellar neurons, and its spatial relationship with GIRK3, Ca$_V$2.1, and

mGluR1α in PC dendrites and PF boutons. Notably, we have demonstrated a higher density

of PI(4,5)P_2 in distal PC dendrites, showing a specific association of mGluR1α with PI(4,5)P_2

in spines. PI(4,5)P_2 also showed concentration and association with Ca$_V$2.1 and GIRK3 in

presynaptic PF AZs. These results will help to elucidate the physiological functions of

PI(4,5)P_2 in neuronal activities, including SV fusion and retrieval in presynaptic membranes

and long-term plasticity in postsynaptic spine membranes. This method could also be a

powerful tool for the analysis of other PIs and phospholipid distribution in neuronal cell

membranes using specific probes.

Materials and Methods

Animals

Animal experiments were conducted in accordance with the guideline of the Institute of
Science and Technology Austria (Animal license number: BMWFW-66.018/0012-
WF/V/3b/2016). Mice were initially purchased from Jackson Laboratory (Bar Harbor, ME,
USA) and were bred at the Preclinical Facility of IST Austria on 12:12 light-dark cycle with
access to food and water *ad libitum*. All experiments were performed in the light phase of the
cycle. C57BL/6J (stock number 000664) mice of either sex at postnatal (P) 5-7 weeks were
used in this study.

Liposome preparation and high-pressure freezing

Phosphatidylcholine (18:1 (Δ9-Cis) PC), phosphatidylethanolamine (18:1 (Δ9-Cis) PE),
phosphatidylserine (18:1 (Δ9-Cis) PS), phosphatidylinositol (18:1 PI) and phosphoinositides
(18:1) were from Avanti Polar Lipids. All liposomes contained 45 mol % PC, 30 mol % PE,
20 mol % PS, and either 5 mol % PI or a phosphoinositide. Solutions of PC, PE, and PS in
chloroform and PI or phosphoinositides in chloroform:methanol:H_2O:HCl (1N) (20:9:1:0.1)
were mixed in the required proportion in amber-color glass vials. To improve
phosphoinositide homogenization with other lipids, the chloroform:methanol 2:1 ratio was
maintained in the mixture. A lipid film was produced by evaporation of solvents in the vial
under the stream of nitrogen gas and then drying using a vacuum desiccator for 2 h. The dried
lipid film was stored at -20 °C with argon gas and used within 2 days.

The lipid film was resuspended in a buffer containing 220 mM sucrose and 20 mM HEPES
(pH 7.4, adjusted with NaOH). The suspension was vortexed well and then freeze-thawed 5
times in liquid nitrogen and warm water (~60 °C). Unilamellar liposomes were produced by
extrusion through a 0.4 μm pore size polycarbonate filter using an extrusion apparatus
(Avanti Polar Lipids). After extrusion, the liposomes were diluted 5 times in 120 mM NaCl,
20 mM HEPES buffer (pH 7.4, adjusted with NaOH), and centrifuged for 20 min at 15,000 g.
The same amount of 100% glycerol was applied to the pellets of the liposomes as a

cryoprotectant. The liposome/glycerol mixture was placed on a copper carrier with a ring of

double-sided tape (140-μm thickness), covered with another carrier, and then frozen by a

high-pressure freezing machine (HPM010, BAL-TEC). The frozen samples were stored in

liquid nitrogen until use.

High-pressure freezing of acute cerebellar slices

Acute slices of mouse cerebellum were prepared at physiological temperature (PT) as

described previously (Eguchi et al., 2020). Briefly, mice were decapitated under isoflurane

anesthesia and their brains were quickly removed from the skull and immersed into a cutting

solution containing (mM): 300 sucrose, 2.5 KCl, 10 glucose, 1.25 NaH_2PO_4, 2 Na Pyruvate, 3

myo-inositol, 0.5 Na ascorbate, 26 $NaHCO_3$, 0.1 $CaCl_2$, 6 $MgCl_2$ (pH 7.4 when gassed with

95% O_2/5% CO_2) at PT (35-37 °C). The cerebellum was dissected from the whole brain and

immediately glued on a cutting stage of a tissue slicer (Linear Slicer Pro7, Dosaka EM,

Kyoto, Japan) and sliced (sagittal, 140-160 μm thickness) in the cutting solution kept at PT.

Slices were then maintained in the artificial cerebrospinal fluid (ACSF) containing (in mM):

125 NaCl, 2.5 KCl, 10 glucose, 1.25 NaH_2PO_4, 2 sodium pyruvate, 3 *myo*-inositol, 0.5

sodium ascorbate, 26 $NaHCO_3$, 2 $CaCl_2$, 1 $MgCl_2$ (pH 7.4 when gassed with 95% O_2/5%

CO_2) at 37 °C until use. Small blocks containing lobule IV-VII were trimmed from the slices

in the cutting solution using a micro scalpel (#10316-14, FST). The trimmed block was

transferred into cryoprotectant buffer (15% polyvinylpyrrolidone in ACSF with 10 mM

HEPES, pH 7.3 adjusted with NaOH), sandwiched between two copper carriers with a ring of

double-sided tape (140-μm thickness), and then frozen by a high pressure freezing machine.

The frozen samples were stored in liquid nitrogen until use.

SDS-digested freeze-fracture replica labeling (SDS-FRL)

The frozen samples were fractured into two parts at -130 °C and replicated by carbon (4-5 nm

thick), carbon-platinum (uni-direction from 60°, 2 nm), and carbon (20-25 nm) deposition in

a freeze-fracture machine (JFD-V, JOEL, Tokyo, Japan). The samples were digested with

2.5% SDS solution containing 0.1 M Tris-HCl (pH 8.3) at 80°C for 18-22 h. The replicas

were washed in the SDS solution and then a washing buffer (50 mM Tris-buffered saline

(TBS, pH 7.4) containing 0.1% BSA) at RT. To avoid non-specific binding of the probes and

antibodies, the replicas were blocked with 3% BSA, 2% cold fish skin gelatin, and 0.05%

Tween-20 in TBS for 1 h at RT. The replicas were incubated with 50 ng/ml GST-tagged PH

domain of phospholipase C δ1 (PI(4,5)P$_2$-Grip, Echelon Inc.) in a dilution buffer (1% BSA,

1% cold fish skin gelatin, and 0.05% Tween-20 in TBS) at 4°C overnight. Then the replica

was incubated with anti-GST antibody (rabbit IgG, 5 μg/ml, Bethyl Laboratories) and anti-

Ca$_V$2.1 antibody (guinea pig, 2.5 μg/ml, Synaptic Systems) as a marker of neurons at 15°C

overnight, and then gold-nanoparticle conjugated secondary antibodies (goat anti-rabbit IgG,

5 nm, 1:50, BBI; donkey anti-guinea pig IgG, 12 nm, 1:30, Jackson Immnoresearch)

dissolved in the dilution buffer at 15°C overnight. For double labeling of PI(4,5)P$_2$ with

proteins, the replicas were incubated with a mixture of primary antibodies (GluD2: rabbit

anti-GST IgG + guinea pig anti-GluD2 IgG (2.0 μg/ml, Frontier Institute); GIRK3: chicken

anti-GST IgY (5 μg/ml, Bethyl Laboratories) + rabbit anti-GIRK3 IgG (XX μg/ml, Frontier

Institute); mGluR1α: rabbit anti-GST IgG + guinea pig anti-mGluR1α IgG (XX μg/ml,

Frontier Institute)) at 15°C 1-2 overnight and then with gold-nanoparticle conjugated

secondary antibodies at 15°C overnight. After washing the replicas with the washing buffer,

they were picked up onto a grid coated with formvar in distilled water. Images were obtained

under TEM (Tecnai 10) with RADIUS software at magnifications of 65,000 and 39,000.

Image analysis

Images were analyzed with Darea software (Kleindienst et al., 2020), Fiji, and R. The gold

particle detection and the demarcation of the region of interest were performed on Darea

software. AZs on the P-face of PF boutons were indicated with the aggregation of

intramembrane particles on the replica at the electron microscopic level as described

previously (Eguchi et al., 2022, 2020; Harris and Landis, 1986; Landis and Reese, 1974;

Masugi-Tokita et al., 2007). Gold particles inside or < 30 nm away from the demarcation

border of AZs (outer rim) were counted as the particles in the AZs (Kleindienst et al., 2020).

Because the PSD area on the P-face of dendritic spines of PCs cannot be identified based on

morphological features, the largest cluster of GluD2-labeling gold particles on the spines was

identified as the PSD area (Eguchi et al., 2020; Luján et al., 2018a). The demarcated region

of the images was imported to R via FIJI/ImageJ for the following point pattern analysis.

Point pattern analysis of the gold particles described below was performed using spatstat

package (version 2.3-0) of R (version 4.1.0). Nearest neighbor distances (NNDs) to both

particles of the same size (e.g. from a 5 nm particle to the nearest 5 nm particle) and the other

size (e.g. from a 5 nm particle to the nearest 10 nm particle) were computed to evaluate the

distribution pattern of the particles. Center Periphery Index (CPI), indicating the location of

the particles in the AZs or PSDs, was calculated as the square of the normalized distances

from the center of the region of interest. When particles are randomly distributed in a circle,

the mean CPI is near 0.5 (Kleindienst et al., 2020). To assess the randomness of the particle

distribution, we performed two types of Monte-Carlo simulations, termed random and fitted

simulations, following the methods described in the previous publications (Kleindienst et al.,

2020; Luján et al., 2018b) using R. For the random simulation, particles were randomly

placed on the demarcated area. The simulated particles were placed to keep the minimum

distance of 10 nm from any other particles and then randomly shifted within a disc with a 30-

nm diameter to reproduce the immunolabeling with a probe and antibodies (Tabata et al.,

2019). In the fitted simulation, a constraint was added that the distance distribution between

the simulated particles should not differ significantly from the distance distribution between

the original particles. The particle distribution pattern was modeled and simulated as a

Matern Cluster point process, and the goodness-of-fit between the real and simulated

distribution was assessed by comparing both the all pairwise distances (APD) and NNDs of

the particles using the two-sample *Kolmogorov-Smirnov* test (KS test), respectively, and

considered them similar if the p-value was equal or above 0.1 for both. To avoid excessive

statistical power due to the large sample size caused by a large number of particles,

parametric bootstrapping was performed when the number of values to be compared

exceeded 100. Specifically, we first randomly selected 100 distance values (NND or APD)

from the simulated distribution and compared them to the empirical cumulative distribution

function of the distance values of the real distribution using the KS test. This process was

repeated 1,000 times, and the average of the p-values was used to assess the goodness-of-fit.

Gold particle clusters were detected with a hierarchical clustering algorithm called Ward

linkage (Figure 1 - Figure Supplement 1C) and a density-based clustering algorithm called

DBSCAN (Figures 2 and 3). For Ward linkage, the threshold distance was set as 50 nm. For

DBSCAN, we set the minimum number of particles consisting of a cluster as 3 and the

maximum distance between particles as the sum of the median and 1.5 times the interquartile

range (IQR) of the NNDs. The area of the convex polygon connecting the outermost particles

forming the cluster was defined as the cluster area.

Statistical analysis

To consider the hierarchical structure, correlation, and probability distribution, data were
analyzed with either a linear mixed-effects model (LMM) or its generalized form (GLMM)
using the lme4 package (version 1.1-27.1) of R (Aarts et al., 2014; Yu et al., 2021).
Probability distributions for models were chosen by the goodness of fit to Poisson (for
discrete variables e.g. the number of particles), normal (e.g. CPI), or gamma distribution (for
continuous variables e.g. NNDs). Appropriate to the particular experiment and statistical
model, fixation methods (perfusion- and cryo-fixation), the faces of cell membranes (P- and
E-face), components of neurons (e.g. somata, spines, AZs on PF boutons), genotypes
(wildtype and Ts65Dn), and potentially their interactions were used as fixed effects, while
experiments, animals, replicas, and cells were used as random effects to consider the nested
and/or crossed data structure in the statistical analysis. For all experiments, at least four
animals per condition were used. All data are presented as estimated marginal means with
95% confidence intervals (CI, in figures) or standard error of means (s.e.m., in text) estimated
using the emmeans package. The goodness-of-fit of the models was assessed by second-order
Akaike Information Criterion (AICc) for the elimination of the random factors (MuMIn
package and then by likelihood ratio chi-square tests (Chi-LRT) with models in which the
fixed effects of interest had been dropped. For multiple pairwise comparisons between 3 or
more groups, posthoc comparisons to assume the significance of differences between pairs of
group means were performed using emmeans packages with Tukey (for all pairwise) or
Benjamini-Hochberg (BH, for selected pairwise) method when Chi-LRT detected a
significant difference ($p < 0.05$). Statistical significance was assumed if $p < 0.05$ (indicated
with blue or single asterisk), $p < 0.01$ (green or double asterisk), and $p < 0.001$ (red or triple
asterisk).

Author Contributions

550 KE and RS designed and conducted the experiments and wrote the manuscript. KE analyzed
551 the data.

Acknowledgment

We would like to thank Nicoleta Condruz (IST Austria, Klosterneuburg, Austria) for technical assistance with sample preparation; the Electron Microscopy Facility of IST Austria (Klosterneuburg, Austria) for technical support with EM works; Natalia Baranova (University of Vienna, Vienna, Austria) and Martin Loose (IST Austria, Klosterneuburg, Austria) for advice on liposome preparation; Yugo Fukazawa (University of Fukui, Fukui, Japan) for comments. This work was financially supported by funding from the Institute of Science and Technology Austria, the European Union's Horizon 2020 research and innovation program under the Marie Skłodowska-Curie grant agreement No. 793482 (to KE), and the European Research Council (ERC) grant agreement No. 694539 (to RS).

References

Aarts E, Verhage M, Veenvliet JV, Dolan CV, Van Der Sluis S. 2014. A solution to dependency: Using multilevel analysis to accommodate nested data. *Nature Neuroscience*. doi:10.1038/nn.3648

Aguado C, Colón J, Ciruela F, Schlaudraff F, Cabañero MJ, Perry C, Watanabe M, Liss B, Wickman K, Luján R. 2008. Cell type-specific subunit composition of G protein-gated potassium channels in the cerebellum. *J Neurochem* **105**:497–511.

Aktar S, Takatori S, Tsuji T, Orii M, Ohsaki Y, Cheng J, Fujimoto T. 2017. A New Electron Microscopic Method to Observe the Distribution of Phosphatidylinositol 3,4-bisphosphate. *Acta Histochem Cytochem* **50**:141–147.

Althof D, Baehrens D, Watanabe M, Suzuki N, Fakler B, Kulik Á. 2015. Inhibitory and excitatory axon terminals share a common nano-architecture of their Cav2.1 (P/Q-type) Ca(2+) channels. *Front Cell Neurosci* **9**:315.

Aoyagi K, Sugaya T, Umeda M, Yamamoto S, Terakawa S, Takahashi M. 2005. The activation of exocytotic sites by the formation of phosphatidylinositol 4,5-bisphosphate microdomains at syntaxin clusters. *J Biol Chem* **280**:17346–17352.

Borges-Merjane C, Kim O, Jonas P. 2020. Functional Electron Microscopy, "Flash and

Freeze," of Identified Cortical Synapses in Acute Brain Slices. *Neuron* **105**:992–1006.e6.

Cheng J, Fujita A, Yamamoto H, Tatematsu T, Kakuta S, Obara K, Ohsumi Y, Fujimoto T.

2014. Yeast and mammalian autophagosomes exhibit distinct phosphatidylinositol 3-

phosphate asymmetries. *Nat Commun* **5**. doi:10.1038/ncomms4207

de Jong APH, Roggero CM, Ho MR, Wong MY, Brautigam CA, Rizo J, Kaeser PS. 2018.

RIM C2B Domains Target Presynaptic Active Zone Functions to PIP2-Containing

Membranes. *Neuron* **98**:335–349.e7.

Di Paolo G, De Camilli P. 2006. Phosphoinositides in cell regulation and membrane

dynamics. *Nature* **443**:651–657.

Eguchi K, Montanaro J, Le Monnier E, Shigemoto R. 2022. The Number and Distinct

Clustering Patterns of Voltage-Gated Calcium Channels in Nerve Terminals. *Front*

Neuroanat **16**. doi:10.3389/fnana.2022.846615

Eguchi K, Nakanishi S, Takagi H, Taoufiq Z, Takahashi T. 2012. Maturation of a PKG-

Dependent Retrograde Mechanism for Exoendocytic Coupling of Synaptic Vesicles. *Neuron*

74:517–529.

Eguchi K, Velicky P, Hollergschwandtner E, Itakura M, Fukazawa Y, Danzl JG, Shigemoto

R. 2020. Advantages of Acute Brain Slices Prepared at Physiological Temperature in the

Characterization of Synaptic Functions. *Front Cell Neurosci* **14**:63.

Falkenburger BH, Jensen JB, Dickson EJ, Suh BC, Hille B. 2010. Phosphoinositides: Lipid

regulators of membrane proteins. *J Physiol* **588**:3179–3185.

Fernández-Alacid L, Aguado C, Ciruela F, Martín R, Colón J, Cabañero MJ, Gassmann M,

Watanabe M, Shigemoto R, Wickman K, Bettler B, Sánchez-Prieto J, Luján R. 2009.

Subcellular compartment-specific molecular diversity of pre- and post-synaptic GABA-

activated GIRK channels in Purkinje cells: Pre- and post-synaptic GABAB-activated GIRK

channels in PCs. *J Neurochem* **110**:1363–1376.

Fujita A, Cheng J, Tauchi-Sato K, Takenawa T, Fujimoto T. 2009. A distinct pool of

phosphatidylinositol 4,5-bisphosphate in caveolae revealed by a nanoscale labeling

technique. *Proc Natl Acad Sci U S A* **106**:9256–9261.

Harris KM, Landis DM. 1986. Membrane structure at synaptic junctions in area CA1 of the
rat hippocampus. *Neuroscience* **19**:857–872.

Haucke V. 2005. Phosphoinositide regulation of clathrin-mediated endocytosis. *Biochem Soc*
Trans **33**:1285–1289.

Honigmann A, van den Bogaart G, Iraheta E, Risselada HJ, Milovanovic D, Mueller V,
Müllar S, Diederichsen U, Fasshauer D, Grubmüller H, Hell SW, Eggeling C, Kühnel K,
Jahn R. 2013. Phosphatidylinositol 4,5-bisphosphate clusters act as molecular beacons for
vesicle recruitment. *Nat Struct Mol Biol* **20**:679–686.

Idevall-Hagren O, De Camilli P. 2015. Detection and manipulation of phosphoinositides.
Biochim Biophys Acta **1851**:736–745.

Kleindienst D, Montanaro J, Bhandari P, Case MJ, Fukazawa Y, Shigemoto R. 2020. Deep
Learning-Assisted High-Throughput Analysis of Freeze-Fracture Replica Images Applied to
Glutamate Receptors and Calcium Channels at Hippocampal Synapses. *Int J Mol Sci* **21**.
doi:10.3390/ijms21186737

Landis DM, Reese TS. 1974. Differences in membrane structure between excitatory and
inhibitory synapses in the cerebellar cortex. *J Comp Neurol* **155**:93–125.

Lei W, Myers KR, Rui Y, Hladyshau S, Tsygankov D, Zheng JQ. 2017. Phosphoinositide-
dependent enrichment of actin monomers in dendritic spines regulates synapse development
and plasticity. *J Cell Biol* **216**:2551–2564.

Luján R, Aguado C, Ciruela F, Arus XM, Martín-Belmonte A, Alfaro-Ruiz R, Martínez-
Gómez J, de la Ossa L, Watanabe M, Adelman JP, Shigemoto R, Fukazawa Y. 2018a. SK2
Channels Associate With mGlu1α Receptors and CaV2.1 Channels in Purkinje Cells. *Front*
Cell Neurosci **12**:311.

Luján R, Aguado C, Ciruela F, Cózar J, Kleindienst D, de la Ossa L, Bettler B, Wickman K,
Watanabe M, Shigemoto R, Fukazawa Y. 2018b. Differential association of GABAB
receptors with their effector ion channels in Purkinje cells. *Brain Struct Funct* **223**:1565–
1587.

Maekawa M, Fairn GD. 2014. Molecular probes to visualize the location, organization and
dynamics of lipids. *J Cell Sci* **127**:4801–4812.

Martin TFJ. 2001. PI(4,5) P2 regulation of surface membrane traffic. *Curr Opin Cell Biol*
13:493–499.

Masugi-Tokita M, Tarusawa E, Watanabe M, Molnar E, Fujimoto K, Shigemoto R. 2007.
Number and Density of AMPA Receptors in Individual Synapses in the Rat Cerebellum as
Revealed by SDS-Digested Freeze-Fracture Replica Labeling. *Journal of Neuroscience*
27:2135–2144.

Milovanovic D, Platen M, Junius M, Diederichsen U, Schaap IAT, Honigmann A, Jahn R,
Van Den Bogaart G. 2016. Calcium promotes the formation of syntaxin 1 mesoscale domains
through phosphatidylinositol 4,5-bisphosphate. *J Biol Chem* **291**:7868–7876.

Ozato-Sakurai N, Fujita A, Fujimoto T. 2011. The distribution of phosphatidylinositol 4,5-
bisphosphate in acinar cells of rat pancreas revealed with the Freeze-Fracture replica labeling
method. *PLoS One* **6**. doi:10.1371/journal.pone.0023567

Petersen EN, Chung H-W, Nayebosadri A, Hansen SB. 2016. Kinetic disruption of lipid rafts
is a mechanosensor for phospholipase D. *Nat Commun* **7**:13873.

Posor Y, Eichhorn-Grünig M, Haucke V. 2015. Phosphoinositides in endocytosis. *Biochimica*
et Biophysica Acta - Molecular and Cell Biology of Lipids **1851**:794–804.

Suh B-C, Hille B. 2008. PIP2 is a necessary cofactor for ion channel function: how and why?
Annu Rev Biophys **37**:175–195.

Suh B-C, Kim D-I, Falkenburger BH, Hille B. 2012. Membrane-localized -subunits alter the
PIP2 regulation of high-voltage activated Ca2+ channels. *Proceedings of the National*
Academy of Sciences **109**:3161–3166.

Szoboszlay M, Kirizs T, Nusser Z. 2017. Objective quantification of nanoscale protein
distributions. *Sci Rep* **7**:15240.

Tabata S, Jevtic M, Kurashige N, Fuchida H, Kido M, Tani K, Zenmyo N, Uchinomiya S,
Harada H, Itakura M, Hamachi I, Shigemoto R, Ojida A. 2019. Electron Microscopic
Detection of Single Membrane Proteins by a Specific Chemical Labeling. *iScience* **22**:256–
268.

Tanaka KAK, Suzuki KGN, Shirai YM, Shibutani ST, Miyahara MSH, Tsuboi H, Yahara M,
Yoshimura A, Mayor S, Fujiwara TK, Kusumi A. 2010. Membrane molecules mobile even

after chemical fixation. *Nat Methods* **7**:865–866.

Taoufiq Z, Eguchi K, Takahashi T. 2013. Rho-Kinase Accelerates Synaptic Vesicle
Endocytosis by Linking Cyclic GMP-Dependent Protein Kinase Activity to
Phosphatidylinositol-4,5-Bisphosphate Synthesis. *Journal of Neuroscience* **33**:12099–12104.

Tsuji T, Takatori S, Fujimoto T. 2019. Definition of phosphoinositide distribution in the
nanoscale. *Current Opinion in Cell Biology*. doi:10.1016/j.ceb.2018.10.008

Ueda Y, Hayashi Y. 2013. PIP3 Regulates Spinule Formation in Dendritic Spines during
Structural Long-Term Potentiation. *Journal of Neuroscience* **33**:11040–11047.

van den Bogaart G, Meyenberg K, Risselada HJ, Amin H, Willig KI, Hubrich BE, Dier M,
Hell SW, Grubmüller H, Diederichsen U, Jahn R. 2011. Membrane protein sequestering by
ionic protein-lipid interactions. *Nature* **479**:552–555.

Wang J, Richards DA. 2012. Segregation of PIP2 and PIP3 into distinct nanoscale regions
within the plasma membrane. *Biol Open* **1**:857–862.

Wenk MR, De Camilli P. 2004. Protein-lipid interactions and phosphoinositide metabolism in
membrane traffic: Insights from vesicle recycling in nerve terminals. *Proceedings of the*
National Academy of Sciences **101**:8262–8269.

Whorton MR, MacKinnon R. 2011. Crystal structure of the mammalian GIRK2 K+ channel
and gating regulation by G proteins, PIP2, and sodium. *Cell* **147**:199–208.

Yu Z, Guindani M, Grieco SF, Chen L, Holmes TC, Xu X. 2021. Beyond t test and ANOVA:
applications of mixed-effects models for more rigorous statistical analysis in neuroscience
research. *Neuron*. doi:10.1016/j.neuron.2021.10.030

Figure 1. SDS-FRL of PI(4,5)P₂ on somatic membranes of mouse cerebellar neurons.

(A-B) Specificity of PI(4,5)P₂ labeling. Replicas of liposomes containing 5% PtdIns or

different stereoisomers of PIs were labeled using GST-PH, anti-GST antibody, and 5-nm gold

particle-conjugated secondary antibody (**A**, Scale bar = 200 nm). The density of gold

particles was highest in the liposome containing PI(4,5)P$_2$ (**B**).

(**C**) Acute cerebellar slice preparation for high-pressure freezing (HPF) and replica

preparation. Left-top, an acute sagittal slice of the mouse cerebellum. The dashed line

indicates the trimmed region for HPF. Left-bottom, a trimmed cerebellar slice on a copper

carrier with double-sided tape for HPF. Right, low-magnification transmission electron

microscopic (TEM) image of the mouse cerebellar replica containing granule cell layer

(GCL), Purkinje cell layer (PCL), and molecular layer (ML). Scale bar = 20 μm.

(**D**) Example TEM image of 5-nm gold particle labeling for PI(4,5)P$_2$ with immunogold

labeling for Ca$_V$2.1 (12 nm) on P- (left) and E-face (right) of the PC somatic membranes of

cerebellar PC. Scale bar = 200 nm.

(**E**) Statistical comparison of the PI(4,5)P$_2$ particle density on the E-face and P-face of the PC

somatic membranes. Open and closed circles indicate the means of the PI(4,5)P$_2$ particle

density in each PC and each mouse, respectively, with different colors. Black horizontal bars

indicate estimated marginal means (emmeans, thick bars) and 95% confidence intervals (CIs,

error bars) of the density estimated by GLMM (Methods). The PI(4,5)P$_2$ density was

significantly higher on the P-face than on the E-face of the PC somatic membranes (P-face:

51.2 ± 8.5 particles/μm^2, E-face: 8.0 ± 1.2 particles/μm^2, n = 213 images/12 cells/4 mice, p <

0.001, Chi-square likelihood ratio test (Chi-LRT)).

(**F**) Comparison of nearest neighbor distances (NND) between real (Data NND, x-axis) and

simulated (Sim NND, y-axis) PI(4,5)P$_2$ particles on PC somatic membranes. Data-NNDs are

significantly smaller than Sim NNDs (Data: 35.1 ± 3.6 nm, Sim: 73.3 ± 7.0 nm, n = 206

images/11 cells/4 mice, p < 0.001, Chi-LRT).

(**G**) Distribution of NNDs of the PI(4,5)P$_2$ particles obtained from a single PC somatic

membrane (n = 2,929 particles). Red and blue lines indicate the distinct components of the

NND distribution estimated from the Gaussian mixture modeling.

(**H**) PI(4,5)P$_2$ labeling (5 nm) with Ca$_V$2.1 (12 nm) on the P-face of GC (left) and molecular

layer interneuron (MLI, right) somatic membranes. Scale bar = 200 nm.

(**I**) The PI(4,5)P$_2$ particle density on the somatic membranes (P-face) of different neuronal

cell types in the mouse cerebellum (PC: 51.8 ± 9.6 particles/μm^2, GC: 69.8 ± 11.3

particles/μm^2, MLI: 42.9 ± 6.7 particles/μm^2, n = 273 images/57 cells/4 mice, p = 0.003, Chi-
LRT).

(**J**) NND of the PI(4,5)P$_2$ particles on the somatic membranes of different neuronal cell types
in the mouse cerebellum (PC: 34.4 ± 3.0 nm, GC: 31.0 ± 2.4 nm, MLI: 34.8 ± 2.6 nm, n =
65,607 values/57 cells/4 mice, p = 0.12, Chi-LRT).

Figure Supplement 1. Non-specific labeling on the somatic membranes of PCs

Figure 2. PI(4,5)P₂ particle distribution on somatodendritic membranes of PCs.

(**A**) Example images obtained from the different compartments of PC dendritic membranes: main shaft (MS, top-left), smooth branchlet (SmB, top-right), spiny branchlet (SpB, bottom-left), and spine (bottom-right). Scale bar = 200 nm. Inset: low-magnification images of MS, SmB, and SpB, respectively, indicated with blue.

(**B**) Comparison of the PI(4,5)P₂ particle density (top) and NNDs (bottom) in the somatodendritic membrane compartments of PCs. The PI(4,5)P₂ density gradually increased

from soma and proximal dendrites to distal dendritic components (top, n = 611 images/134

components/4 mice, p < 0.001, Chi-LRT), whereas no significant difference in NNDs was

detected between these compartments (bottom, n = 67,380 values/134 components/4 mice, p

= 0.49, Chi-LRT).

(**C**) Example images of the PI(4,5)P$_2$ clusters on the somatic (left) and spine (right)

membrane of PCs. Colored polygons indicate individual PI(4,5)P$_2$ clusters detected by the

DBSCAN algorithm. Scale bars = 200 nm.

(**D**) Quantitative analysis of the PI(4,5)P$_2$ clusters on the somatodendritic compartments of

PCs. Top, the number of clusters per area (cluster density, n = 611 values/134 components/4

mice). Middle, the cluster area (10,479 clusters/124 components/4 mice). Bottom, the intra-

cluster particle density (10,343 clusters/134 components/4 mice). The cluster density and the

intra-cluster particle density are significantly higher in spine membranes compared to the

somatic and proximal dendritic membranes (p < 0.001, Chi-LRT), whereas there is no

significant difference in the cluster area (p = 0.17, Chi-LRT).

Table Supplement 1. PI(4,5)P$_2$ distribution in somatodendritic membranes of PCs

Figure 3. PI(4,5)P₂ particle distribution in GCs and MLIs.

(**A**) Example images obtained from the different compartments of GC membranes: dendrites (top-left), parallel fiber (PF) axons (top-right), presynaptic PF boutons on the PC spine (PC-PF, bottom-left) and on the MLI dendrite (PF-MLI, bottom-right). Yellow and orange in the

PF boutons indicate the cross-fracture and synaptic vesicle membranes, respectively. Scale bars = 200 nm.

(**B**) Example images obtained from different compartments of MLI membranes: dendrites (top) and basket cell (BC) presynaptic bouton on PC soma (bottom). Scale bar = 200 nm.

(**C**) Comparison of the PI(4,5)P$_2$ particle density (left) and NNDs (right) in the different GC membrane compartments. NNDs show significant differences between the compartments (n = 67,380 values/134 components/4 mice, p < 0.001, Chi-LRT), whereas there is no significance in the density (n = 376 images/320 components/4 mice, p = 0.09, Chi-LRT).

(**D**) Quantitative analysis of the PI(4,5)P$_2$ particle clusters in different GC membrane compartments. There are no significant differences in the cluster density (n = 376 images/320 components/4 mice, p = 0.37, Chi-LRT), the cluster area (n = 4070 clusters/254 components/4 mice, p = 0.18, Chi-LRT), and the intra-cluster particle density (n = 4004 clusters/253 compartments/4 mice, p = 0.50, Chi-LRT).

(**E**) Comparison of the PI(4,5)P$_2$ particle density (left) and NNDs (right) in different membrane compartments of MLIs. NNDs in dendritic and BC bouton membranes are significantly larger than that in somatic membranes (n = 24,012 values/152 components/4 mice, p < 0.001, Chi-LRT), whereas there is no significant difference in the density between the compartments (n = 230 images/153 components/4 mice, p = 0.46, Chi-LRT).

(**F**) Quantitative analysis of the PI(4,5)P$_2$ particle clusters in different MLI membrane compartments. The intra-cluster particle density in BC bouton membranes is significantly smaller than the other components (n = 3,766 clusters/149 components/4 mice, p = 0.03, Chi-LRT), whereas no significant difference was detected in the cluster density (n = 230 images/153 components/4 mice, p = 0.43, Chi-LRT) and the cluster area (n = 3,822 clusters/146 components/4 mice, p = 0.12, Chi-LRT) between these compartments.

Figure Supplement 1. Clustered and scattered PI(4,5)P$_2$ distribution in different membrane compartments of MLIs

Table Supplement 1. PI(4,5)P$_2$ distribution in different membrane compartments of GCs

Table Supplement 2. PI(4,5)P$_2$ distribution in different membrane compartments of MLIs

Figure 4. PI(4,5)P$_2$ distribution in pre- and post-synaptic membranes of PF-PC synapses.

(**A**) Example images for PI(4,5)P$_2$ particle distribution on the membranes of PF-PC (top) and PF-MLI bouton (bottom). Red and black circles indicate gold particles for PI(4,5)P$_2$ and Ca$_V$2.1, respectively. The Blue area and dotted line indicate AZ and outer-rim (30 nm from the edge of AZ, Methods), respectively. Scale bar = 200 nm.

(**B**) Beeswarm plot of the PI(4,5)P$_2$ density in the whole bouton (Bt), AZs, and extra-AZ region (exAZ) of the PF-PC (left) and PF-MLI (right) bouton membranes. The PI(4,5)P$_2$ density was significantly higher in AZs than in the whole bouton and the exAZ in both PF-PC and PF-MLI boutons (n = 111 boutons/4 mice, p < 0.001, Tukey method).

(**C**) Comparison of the PI(4,5)P$_2$ density in the AZs between real and simulated random distribution on the PF bouton membranes. The density of the real particle distribution was significantly higher than that of the simulated one in both PF-PC (real: 97.2 ± 19.7 particles/μm^2, sim: 66.7 ± 13.5 particles/μm^2, n = 55 boutons/4 mice, p < 0.001, Chi-LRT) and PF-MLI (real 86.7 ± 12.4 particles/μm^2, sim: 73.2 ± 10.4 particles/μm^2, n = 56 boutons/4 mice, p = 0.002, Chi-LRT) AZ membranes.

(**D**) Distribution of center-periphery index (CPI) of PI(4,5)P$_2$ particles in boutons (top) and AZs (bottom). Blue in the bottom graph indicates the CPI distribution of the simulated particles that are randomly distributed in AZs.

(**E**) Comparison of CPIs of the PI(4,5)P$_2$ particles in AZs between PF-PC and PF-MLI AZs. There is no significant difference in the CPIs between PF-PC and PF-MLI AZs (PF-PC: 0.67 ± 0.02, PF-MLI: 0.67 ± 0.02, n = 107 boutons/4 mice, p = 0.999, Chi-LRT). Dashed lines with red show the mean CPIs of the simulated randomly-distributed particles. Asterisks on the bars indicate statistical differences in CPIs between real and simulated particles (PF-PC: 0.59 ± 0.02, PF-MLI: 0.58 ± 0.02, *p < 0.05, **p < 0.01, Tukey method).

(**F**) Example images of the PI(4,5)P$_2$ particle distribution on the PC spine membrane. Red and black circles indicate PI(4,5)P$_2$ and GluD2, respectively. The green area indicates postsynaptic density (PSD) based on the cluster of GluD2. Scale bar = 200 nm.

(**G**) Beeswarm plot of the PI(4,5)P$_2$ density in the whole spine (spine), PSD, and extra-PSD region (exPSD) of the PC spine membranes. No significant difference in the density was detected between these compartments (n = 108 spines/6 mice, p = 0.77, Chi-LRT).

(**H**) Comparison of the PI(4,5)P$_2$ density in the PSDs between real and simulated random distribution on the PC spine membranes. The density of the real and simulated particle distribution was not significantly different (real: 0.59 ± 0.01, sim: 0.55 ± 0.01, n = 108 spines/6 mice, p = 0.25, Chi-LRT).

(**I**) Distribution of CPIs of PI(4,5)P$_2$ particles in spines (top) and PSDs (bottom). Blue in the bottom graph indicates the CPI distribution of the simulated particles that are randomly distributed in PSDs. The CPI of the PI(4,5)P$_2$ particles is uniformly distributed in the spines and PSDs, suggesting the random distribution of PI(4,5)P$_2$.

Table Supplement 1. PI(4,5)P$_2$ particle density in synaptic membranes of cerebellar neurons

Figure 5. Ubiquitous association of Ca$_V$2.1 with PI(4,5)P$_2$ on cell membranes of PCs and GCs

(**A**) Example images for co-immunolabeling of PI(4,5)P$_2$ and Ca$_V$2.1 on somatic (left), SpB (middle), and spine (right) membranes of the PC. Red and purple (closed, open) circles indicate PI(4,5)P$_2$ and Ca$_V$2.1 (real, fitted-simulated) particles, respectively. Scale bars = 200 nm.

(**B**) Comparison of the NNDs from $Ca_V2.1$ to $PI(4,5)P_2$ particles (NND_{C-P}) between real and fitted-simulated $Ca_V2.1$ distribution on somatic (left), SpB (middle), and spine (right) membranes of PCs. The NND_{C-P} of the real distribution was significantly smaller than that of the simulated one in somatic (real: 89.7 ± 9.6 nm, sim: 112.9 ± 12.1 nm, n = 129 images/14 cells/5 mice, p < 0.001, Chi-LRT), SpB (real: 82.3 ± 8.1 nm, sim: 100.5 ± 9.9 nm, n = 158 images/20 dendrites/4 mice, p < 0.001, Chi-LRT), and spine membrane (real: 80.9 ± 8.7 nm, sim: 92.8 ± 9.9 nm, 69 spines/4 mice, p = 0.006, Chi-LRT).

(**C**) Comparison of NND_{C-P} between somatodendritic compartments. There is no significant difference between the PC compartments (n = 12,956 values/92 components/5 mice, p = 0.70, Chi-LRT).

(**D**) Example images for co-immunolabeling of $PI(4,5)P_2$ and $Ca_V2.1$ on somatic (left) and presynaptic PF-PC (middle) and PF-MLI (right) AZ membranes of the GC. Red and purple (closed, open) circles indicate $PI(4,5)P_2$ and $Ca_V2.1$ (real, fitted-simulated) particles, respectively. The Blue area and dotted lines indicate AZs and the outer-rim, respectively. Scale bars = 200 nm.

(**E**) Comparison of the NND_{C-P} between real and fitted-simulated $Ca_V2.1$ distribution on somatic (left), PF-PC AZ (middle), and PF-MLI AZ (right) membranes of GCs. The NND_{C-P} of the real distribution was significantly smaller than that of the simulated one in somatic (real: 76.0 ± 9.0 nm, sim: 93.0 ± 13.1 nm, n = 81 images/25 cells/5 mice, p = 0.017, Chi-LRT), PF-PC AZ (real: 54.0 ± 7.9 nm, sim: 64.0 ± 9.3 nm, n = 54 AZs/4 mice, p < 0.001, Chi-LRT), and PF-MLI AZ membrane (real: 60.8 ± 6.7 nm, sim: 67.0 ± 7.3 nm, 52 AZs/4 mice, p = 0.006, Chi-LRT).

(**F**) Comparison of NND_{C-P} between presynaptic AZs of PF-PC and PF-MLI synapses. There is no significant difference in NND_{C-P} between the AZs (n = 1,110 values/68 AZs/4 mice, p = 0.60, Chi-LRT).

Figure 6. Association of Ca$_V$2.1 with PI(4,5)P$_2$ on cell membranes of MLIs

(**A**) Example images for co-immunolabeling of PI(4,5)P$_2$ and Ca$_V$2.1 on somatic (left) and

basket cell (BC)-PC bouton (right) membranes of the MLI. Red and purple (closed, open)

circles indicate PI(4,5)P$_2$ and Ca$_V$2.1 (real, fitted-simulated) particles, respectively. P-face of

BC-PC boutons were identified based on Ca$_V$2.1 clusters and the surrounding E-face of the

PC somatic membranes with Ca$_V$2.1 clusters. Scale bars = 200 nm.

(**B**) Comparison of the NND$_{C-P}$ between real and fitted-simulated Ca$_V$2.1 distribution on

somatic (left) and BC-PC bouton (right) membranes of MLIs. The NND$_{C-P}$ of the real

distribution was significantly smaller than that of the simulated one in somatic (real: $97.5 \pm$

13.1 nm, sim: 115.4 ± 15.5 nm, n = 102 images/26 cells/4 mice, p < 0.001, Chi-LRT) and

BC-PC bouton membrane (real: 106.0 ± 7.7 nm, sim: 143.0 ± 10.4 nm, 37 boutons/3 mice, p

= 0.006, Chi-LRT).

(**C**) Comparison of NND$_{C-P}$ between somatic and bouton membranes. No significant

difference was shown between these compartments (n = 2,854 values/63 components/4 mice,

p = 0.29, Chi-LRT).

Figure 7. Association of GIRK3 channels and mGluR1α receptors with PI(4,5)P₂ on cell membranes of cerebellar neurons

(**A**) Example images for co-labeling of PI(4,5)P$_2$ and GIRK3 channels on PC SpB (left), PC spine (middle), and PF-PC AZ (right) membranes. Red and blue (closed, open) circles indicate PI(4,5)P$_2$ and GIRK3 (real, fitted-simulated) particles, respectively. The Blue area and a dotted line on the left indicate the AZ and the outer-rim, respectively. Scale bars = 200 nm.

(**B**) Comparison of the NNDs from GIRK3 to PI(4,5)P$_2$ particles (NND$_{G-P}$) between real and fitted-simulated GIRK3 distribution on PC SpB (left), PC spine (middle), and PF-PC AZ (right) membranes. The NND$_{G-P}$ of the real distribution was significantly smaller than that of the simulated one in SpB (real: 89.3 ± 11.4 nm, sim: 120.7 ± 15.4 nm, n = 88 images/19 dendrites/5 mice, $p < 0.001$, Chi-LRT), spine (real: 63.6 ± 8.4 nm, sim: 83.0 ± 11.0 nm, n = 57 spines/4 mice, $p < 0.001$, Chi-LRT), and PF-PC AZ membrane (real: 58.6 ± 3.9 nm, sim: 81.1 ± 5.3 nm, 65 AZs/4 mice, $p < 0.001$, Chi-LRT).

(**C**) Comparison of NND$_{G-P}$ between the post- and presynaptic compartments of cerebellar neurons. The NND$_{G-P}$ was significantly shorter in the spine and AZ membranes than in the SpB membrane (4,730 values/90 compartments/4 mice, $p < 0.001$, Chi-LRT).

(**D**) Example images for co-labeling of PI(4,5)P$_2$ and mGluR1α receptors on PC SpB (left) and spine (right) membranes. Red and green (closed, open) circles indicate PI(4,5)P$_2$ and mGluR1α (real, fitted-simulated) particles, respectively. Scale bars = 200 nm.

(**E**) Comparison of the NNDs from mGluR1α to PI(4,5)P$_2$ particles (NND$_{M-P}$) between real and fitted-simulated mGluR1α distribution on PC SpB (left) and spine (right) membranes. The NND$_{M-P}$ of the real distribution was significantly smaller than that of the simulated one in the spine membranes (real: 81.8 ± 18.4 nm, sim: 91.2 ± 20.6 nm, n = 140 spines/7 mice, $p < 0.001$, Chi-LRT), but not in the SpB membrane (n = 147 images/35 dendrites/7 mice, $p = 0.36$, Chi-LRT).

(**F**) Comparison of NND$_{M-P}$ between the SpB and spine membranes of PCs. The NND$_{M-P}$ was shorter in the spines than in the SpB membrane (real: 110.0 ± 5.6 nm, sim: 112.0 ± 5.6 nm, n = 22,391 values/140 compartments/7 mice, $p < 0.001$, Chi-LRT).

Figure 1 - Figure supplement 1. Non-specific labeling on the somatic membranes of PCs

(**A**) Example images of the $PI(4,5)P_2$ labeling without incubation with GST-PH on the P-
(left) and E-face of PC somatic membranes. The replicas were subsequently incubated with
anti-GST/anti-$Ca_V2.1$ primary antibodies and gold-conjugated secondary antibodies (5 nm for
anti-GST, 12 nm for anti-$Ca_V2.1$). Scale bar = 200 nm.

(**B**) Comparison of the 5-nm gold particle densities with and without GST-PH on the E- and
P-face of PC somatic membranes. The particle density without GST-PH was significantly
lower than that with GST-PH in both faces. Without the GST-PH incubation, no significant
difference in the density was detected between the E- and P-face (p = 0.86, multiple
comparisons with the Benjamini-Hochberg (BH) method).

(**C**) Comparison of the number of 5-nm gold particles per cluster on the E- and P-face of the
PC somatic membranes incubated with or without GST-PH. The clusters were detected using
Ward Linkage hierarchical clustering method. The threshold distance to separate each cluster
was set as 50 nm, which is nearby the maximum distance for cluster detection on the PC
somatic membrane using DBSCAN (Figure 2, Method). The particle number per cluster
without GST-PH incubation was significantly lower than that with GST-PH in both faces
(multiple comparisons with BH method), suggesting that the $PI(4,5)P_2$ particle clusters

observed on the neuronal membranes are also ascribable to specific clustering of GST-PH

binding sites but not to non-specific aggregation of the primary and secondary antibodies.

Figure 3 - Figure supplement 1. Clustered and scattered PI(4,5)P$_2$ distribution on MLI compartments

(**A**) Distribution of NNDs distribution of the PI(4,5)P$_2$ particles obtained from the somatic (left, n = 20,164 particles), dendritic (middle, n = 3,175 particles), and BC bouton membranes (right, n = 673 particles) of MLIs. Red and blue lines indicate the shorter and longer components of the NND distribution, respectively, estimated from the Gaussian mixture modeling.

(**B**) Proportion (left) and mean values of shorter (red, middle) and longer (blue, left) components of the NND of PI(4,5)P$_2$ particles on MLI compartments estimated by the Gaussian mixture model. Since there are no significant differences in the mean values of both shorter and longer components of the NND (p = 0.21 and 0.24, respectively, Chi-LRT), the higher mean NND values in the dendritic and BC bouton membranes compared to the somatic membranes (Figure 3E) are ascribable to larger proportions of the scattered PI(4,5)P$_2$ particles in these compartments.

Figure 2 - Table Supplement 1. PI(4,5)P₂ distribution in somatodendritic membrane

compartments of PCs

Particle density (μm⁻²)

Compartment[*]	mean[**]	s.e.m	95% CI		P-value[†] / Z-score				
			Lower	Upper	Spine	SpB	SmB	MS	Soma
Soma	52.3	7.0	40.2	68.1	< .001	0.341	0.949	1.000	-
MS	52.3	7.1	40.2	68.2	< .001	0.346	0.949	-	-0.002
SmB	48.7	6.0	38.2	62.1	< .001	0.019	-	0.732	0.732
SpB	62.7	7.8	49.2	80.0	0.149	-	-3.059	-1.848	-1.858
Spine	73.7	8.6	58.6	92.5	-	-2.287	-5.882	-3.917	-3.917

NND (nm)

Compartment[*]	mean[**]	s.e.m	95% CI		P-value[†] / Z-score				
			Lower	Upper	Spine	SpB	SmB	MS	Soma
Soma	34.3	2.3	30.2	39.0	0.877	1.000	0.871	0.992	-
MS	33.5	2.3	29.3	38.4	0.983	0.994	0.662	-	0.444
SmB	35.9	2.3	31.6	40.8	0.419	0.755	-	-1.347	-0.964
SpB	34.2	2.1	30.3	38.6	0.873	-	1.195	-0.416	0.089
Spine	32.4	2.4	28.0	37.6	-	0.960	1.724	0.544	0.950

Cluster density (μm⁻²)

Compartment[*]	mean[**]	s.e.m	95% CI		P-value[†] / Z-score				
			Lower	Upper	Spine	SpB	SmB	MS	Soma
Soma	8.28	1.00	6.55	10.5	< .001	0.483	0.859	0.990	-
MS	7.95	0.96	6.27	10.1	< .001	0.221	0.991	-	0.467
SmB	7.67	0.88	6.13	9.60	< .001	0.027	-	0.455	0.993
SpB	9.41	1.08	7.51	11.8	0.054	-	-2.943	-2.098	-1.622
Spine	11.7	1.42	9.22	14.8	-	-2.701	-5.321	-4.349	-3.945

Cluster area (nm²)

Compartment[*]	mean[**]	s.e.m	95% CI		P-value[†] / Z-score				
			Lower	Upper	Spine	SpB	SmB	MS	Soma
Soma	936	76	798	1098	0.265	0.960	0.999	1.000	-
MS	914	88	757	1105	0.385	0.927	0.991	-	0.185
SmB	965	64	847	1099	0.155	0.992	-	-0.457	-0.288
SpB	1006	68	882	1148	0.085	-	-0.443	-0.812	-0.684
Spine	668	99	500	892	-	2.524	2.270	1.780	2.003

Intra-cluster particle density (nm⁻²)

Compartment[*]	mean[**]	s.e.m	95% CI		P-value[†] / Z-score				
			Lower	Upper	Spine	SpB	SmB	MS	Soma
Soma	0.0093	0.0007	0.0080	0.0108	< .001	0.244	0.996	0.910	-
MS	0.0100	0.0008	0.0086	0.0117	0.006	0.826	0.977	-	-0.865
SmB	0.0096	0.0007	0.0084	0.0109	< .001	0.319	-	0.588	-0.376
SpB	0.0109	0.0008	0.0095	0.0125	0.055	-	-1.896	-1.063	-2.047
Spine	0.0137	0.0011	0.0117	0.0160	-	-2.693	-4.370	-3.394	-4.308

[*]MS = main shaft, SmB = smooth branchlet, SpB = spiny branchlet

[**]Marginal means estimated from generalized mixed-effects models

[†]P-values are calculated using mutiple comparison with Tukey method

952 **Figure 3 - Table Supplement 1. PI(4,5)P₂ distribution in different membrane**

953 **compartments of GCs**

Particle density (μm^{-2})

Compartment[*]	mean[**]	s.e.m	95% CI Lower	95% CI Upper	P-value[†] / Z-score PF-MLI	PF-PC	PF axon	Dendrite	Soma
Soma	68.8	10.8	51.6	94.4	0.961	1.000	0.339	0.658	-
Dendrite	57.7	7.2	45.3	73.6	0.916	0.309	0.916	-	1.353
PF axon	53.2	6.9	41.2	68.7	0.540	0.088	-	0.845	1.860
PF-PC bouton	70.9	8.9	55.3	90.7	0.864	-	-2.510	-1.916	-0.108
PF-MLI bouton	63.0	8.5	48.4	82.1	-	0.980	-1.534	-0.847	0.679

NND (nm)

Compartment[*]	mean[**]	s.e.m	95% CI Lower	95% CI Upper	P-value[†] / Z-score PF-MLI	PF-PC	PF axon	Dendrite	Soma
Soma	30.8	1.8	27.6	34.5	0.153	0.054	0.027	0.521	-
Dendrite	34.5	2.4	30.2	39.4	0.889	0.848	0.298	-	-1.563
PF axon	43.7	5.1	34.8	54.9	0.775	0.726	-	-1.937	-2.944
PF-PC bouton	37.4	2.6	32.7	42.8	1.000	-	1.243	-1.017	-2.701
PF-MLI bouton	37.5	3.1	31.9	44.2	-	-0.023	1.158	-0.922	-2.274

Cluster density (μm^{-2})

Compartment[*]	mean[**]	s.e.m	95% CI Lower	95% CI Upper	P-value[†] / Z-score PF-MLI	PF-PC	PF axon	Dendrite	Soma
Soma	10.94	1.51	8.35	14.3	0.887	0.994	0.357	0.541	-
Dendrite	9.11	1.20	7.04	11.8	0.997	0.832	0.937	-	1.534
PF axon	8.14	1.39	5.82	11.4	0.885	0.582	-	0.780	1.828
PF-PC bouton	10.37	1.47	7.86	13.7	0.982	-	-1.470	-1.050	0.414
PF-MLI bouton	9.5	1.51	6.99	13.0	-	0.552	-0.931	-0.347	0.926

Cluster area (nm^{2})

Compartment[*]	mean[**]	s.e.m	95% CI Lower	95% CI Upper	P-value[†] / Z-score PF-MLI	PF-PC	PF axon	Dendrite	Soma
Soma	1016	144	770	1340	0.990	0.971	0.772	0.053	-
Dendrite	696	127	488	995	0.974	0.340	1.000	-	2.705
PF axon	673	215	360	1260	0.992	0.749	-	0.091	1.164
PF-PC bouton	1208	442	590	2475	0.869	-	-1.205	-1.859	-0.924
PF-MLI bouton	851	327	400	1808	-	0.969	-0.439	-0.610	0.558

Intra-cluster particle density (nm^{-2})

Compartment[*]	mean[**]	s.e.m	95% CI Lower	95% CI Upper	P-value[†] / Z-score PF-MLI	PF-PC	PF axon	Dendrite	Soma
Soma	0.0092	0.0009	0.0077	0.0110	0.989	0.991	0.993	0.119	-
Dendrite	0.0123	0.0017	0.0094	0.0160	0.845	0.370	0.703	-	-2.385
PF axon	0.0082	0.0022	0.0048	0.0139	0.976	1.000	-	1.281	0.432
PF-PC bouton	0.0086	0.0014	0.0063	0.0117	0.968	-	-0.163	1.807	0.455
PF-MLI bouton	0.0101	0.0023	0.0065	0.0157	-	-0.642	-0.593	1.023	-0.480

[*]PF= parallel fiber, PC = Purkinje cell, MLI = molecular layer interneuron

[**]Marginal means estimated from generalized mixed-effects models

954 [†]P-values are calculated using mutiple comparison with Tukey method

Figure 3 - Table Supplement 2. PI(4,5)P$_2$ distribution in different membrane

compartments of MLIs

Particle density (μm^{-2})

Compartment*	mean**	s.e.m	95% CI Lower	95% CI Upper	P-value† / Z-score BC-PC	Dendrite	Soma
Soma	42.4	6.5	31.4	57.1	0.449	0.789	-
Dendrite	39.1	5.6	29.5	51.9	0.649	-	0.656
BC-PC bouton	35.7	5.8	26.1	49.0	-	0.887	1.208

NND (nm)

Compartment*	mean**	s.e.m	95% CI Lower	95% CI Upper	P-value† / Z-score BC-PC	Dendrite	Soma
Soma	34.9	2.6	30.3	40.3	< .001	< .001	-
Dendrite	44.9	3.5	38.5	52.2	0.358	-	-3.612
BC-PC bouton	52.2	6.1	41.5	65.7	-	-1.368	-3.650

Cluster density (μm^{-2})

Compartment*	mean**	s.e.m	95% CI Lower	95% CI Upper	P-value† / Z-score BC-PC	Dendrite	Soma
Soma	6.73	1.08	4.91	9.21	0.417	0.635	-
Dendrite	6.02	0.96	4.40	8.24	0.791	-	0.908
BC-PC bouton	5.49	1.05	3.77	8.00	-	0.652	1.261

Cluster area (nm^2)

Compartment*	mean**	s.e.m	95% CI Lower	95% CI Upper	P-value† / Z-score BC-PC	Dendrite	Soma
Soma	979	303	534	1794	0.994	0.307	-
Dendrite	690	115	498	955	0.667	-	1.468
BC-PC bouton	931	263	535	1619	-	-0.857	0.109

Intra-cluster particle density (nm^{-2})

Compartment*	mean**	s.e.m	95% CI Lower	95% CI Upper	P-value† / Z-score BC-PC	Dendrite	Soma
Soma	0.0103	0.0011	0.0083	0.0127	0.008	0.747	
Dendrite	0.0111	0.0009	0.0095	0.0129	0.016		-0.727
BC-PC bouton	0.0038	0.0015	0.0017	0.0084		2.763	2.975

*BC = basket cell

**Marginal means estimated from generalized mixed-effects models

†P-values are calculated using mutiple comparison with Tukey method

Figure 4 - Table Supplement 1. PI(4,5)P$_2$ particle density in synaptic membranes of

cerebellar neurons (particles/µm^2)

PF-PC boutons

Compartment[*]	mean[**]	s.e.m	95% CI		P-value[†] / Z-score		
			Lower	**Upper**	**exAZ**	**AZ**	**Bouton**
Bouton	70.2	10.2	52.9	93.3	< .001	< .001	-
AZ	102.6	15.2	76.7	137.2	< .001	-	-5.908
exAZ	55.1	8.3	40.9	74.1	-	6.952	4.018

PF-MLI boutons

Compartment[*]	mean[**]	s.e.m	95% CI		P-value[†] / Z-score		
			Lower	**Upper**	**exAZ**	**AZ**	**Bouton**
Bouton	63.8	9.9	47.2	86.4	< .001	< .001	-
AZ	118.0	18.9	86.2	161.3	< .001	-	-8.089
exAZ	40.3	6.6	29.2	55.6	-	6.952	5.950

PC spines

Compartment[*]	mean[**]	s.e.m	95% CI		P-value[†] / Z-score		
			Lower	**Upper**	**exPSD**	**PSD**	**Spine**
Spine	114.0	12.9	91.0	142	0.749	0.840	-
PSD	124.0	16.5	95.6	161	0.805	-	-0.564
exPSD	111.0	14.8	85.0	144	-	0.628	0.724

[*]AZ = active zone, exAZ = extra AZ, PSD = postsynaptic density, exPSD = extra PSD

[**]Marginal means estimated from generalized mixed-effects models

[†]P-values are calculated using mutiple comparison with Tukey method