1   Direct Submission Research Report

3   Resurrection genomics provides molecular and phenotypic evidence of rapid adaptation to
4   salinization in a keystone aquatic species

5   Matthew J. Wersebe[1,2] and Lawrence J. Weider[1]

7   [1] Program in Ecology and Evolutionary Biology, Department of Biology, University of
8   Oklahoma

9   [2]Corresponding Author: 730 Van Vleet Oval, Richards Hall 304, Norman, OK 73019

10   matthew.wersebe@ou.edu

14   Keywords:

15   *Daphnia,* Salinization, Whole Genome Sequencing, LC_{50}, Rapid Adaptation

Abstract:

Ecologists and evolutionary biologists are increasingly cognizant of rapid adaptation in wild populations. Rapid adaptation to anthropogenic environmental change is critical for maintaining biodiversity and ecosystems services into the future. Anthropogenic salinization of freshwater ecosystems is quickly emerging as a primary threat, which is well documented in the northern temperate ecoregion. Specifically, many northern temperate lakes have undergone extensive salinization because of urbanization, including increased impervious surfaces for runoff, and the extensive use of road deicing salts (e.g., NaCl). It remains unclear if increasing salinization will lead to extirpation of species from these systems. Using a "resurrection genomics" approach, we investigated whether the keystone aquatic herbivore, *Daphnia pulicaria,* has evolved increased salinity tolerance in a severely salinized lake located in Minnesota, USA. Whole genome resequencing of 54 *Daphnia* clones hatched from resting eggs that represent a 25-year temporal contrast demonstrates that many regions of the genome containing genes related to osmoregulation are under selection in the study population. Tolerance assays of clones revealed that the most recent clones are more tolerant to salinity than older clones; this pattern is concomitant with the temporal pattern of increasing salinity in this lake. Together, our results demonstrate that keystone species such as *Daphnia* can rapidly adapt to increasing freshwater salinization. Further, our results indicate that rapid adaptation to salinity may allow lake *Daphnia* populations to persist in the face of anthropogenic salinization maintaining the food webs and ecosystem services they support despite global environmental change.

Significance Statement:

Rapid adaptation to human-induced environmental change is critical for preserving biodiversity and ecosystem services into the future. A key question is whether populations of keystone species can rapidly adapt to maintain the ecosystems they support. We investigated rapid adaptation to anthropogenic salinization in *Daphnia pulicaria,* a keystone aquatic herbivore in lake ecosystems. By resuscitating decades-old resting eggs, we investigate genomic changes across an approximately 25-year temporal contrast from a severely salinized lake. We report that the genes showing signatures of natural selection throughout the genome are related to osmoregulation. Phenotyping clones for salinity tolerance revealed evidence that genetic changes underlie rapid evolution. We provide molecular genetic and phenotypic evidence for rapid adaptation to salinity in *D. pulicaria.*

Introduction:

Ecologists and evolutionary biologists now recognize many examples of wild populations rapidly evolving in the face of environmental change (1, 2). A population's ability to rapidly evolve is critical for survival in the face of ever-increasing anthropogenic environmental change. This capacity is especially important for organisms that provision key ecosystem services or are keystone species because their extirpation would fundamentally alter ecosystem dynamics. Despite this, studies demonstrating a mechanistic basis for rapid adaptation that integrates information from the genome to the phenome of a population are rare (3). A key reason for the paucity of such studies is that many loci of small effect are thought to contribute most to rapid evolutionary change and do not align with classical "hard-sweep" models making their identification difficult (4, 5). Additional analytical challenges are exacerbated by different population-specific parameters (e.g., effective size or N_e) that influence the supply of new potentially beneficial mutations ($\theta = 4N_e\mu$) and the resulting distribution of fitness effects ($N_e s$) (6, 7).

One way that rapid adaptation can be studied is using temporal genomic contrasts. Most commonly, temporal contrasts take the form of so-called "evolve and re-sequence studies" which follow a population across time during experimental evolution trials that typically employ contrasting selection regimes (8–10). By finding the alleles that increase in frequency rapidly in different treatments, the molecular basis of phenotypic shifts can be explored (8). Such studies have been largely restricted to organisms such as bacteria (11) or *Drosophila* (10), which have rapid generation times and are easily manipulated in the lab. Other types of temporal contrasts encompass more natural experiments, such as the isolation and sequencing of ancient DNA, which can give insight into past selection (12). However, with few exceptions, the genomes sampled are divorced from the phenomes they produced; thus, inference is based solely on the change in allele frequencies. A third way that temporal contrasts can be studied is through "resurrection" studies that seek to hatch or germinate seeds, cysts or other resting stages of organisms and compare genotypes and phenotypes from different points in time (13–15). For instance, resurrection ecology (16, 17), commonly used in animals from the freshwater crustacean genus *Daphnia* has provided insight into the genetic basis of various traits (15, 18, 19). Thus far, however, this method has not allowed the identification of loci that can be plausibly related to the phenotype under study because genetic markers are either too sparse (20) or the traits under study are too highly integrated across the genome (18).

Daphnia are keystone species in freshwater food webs, connecting the flow of energy from algal production to higher trophic levels such as fish (21, 22). Specific to many North American lakes, *Daphnia pulicaria,* maintains water clarity, a key ecosystem service and supports recreational fisheries with values in the millions of USD per lake per annum (23). Freshwater ecosystems are among the most threatened ecosystems worldwide, impacted by various anthropogenic stressors such as pollution, climate change and invasive species (24). One issue threatening many freshwater ecosystems is salinization due to human activities (25), within northern temperate lakes specifically salinization is particularly acute (26, 27). The causes and scope of salinization have been well known (28, 29), while more recent studies have focused on

deciphering the ecological impacts of salinization (30, 31). In addition, we lack a more general
understanding of this widespread environmental issue from an evolutionary perspective, and of
the specific genetic architecture of adaptative responses. Such a perspective is critical because
recent research has shown that current water quality guidelines do not sufficiently protect aquatic
life from salinization (32).

To address this short coming, we sought to use resurrection ecology to study the
evolutionary response of *D. pulicaria* from a severely salinized lake located in Minnesota, USA.
Previous work on this lake has demonstrated the ecological dynamics of this population over the
last 150 years (30). Towards this goal, we resurrected genotypes from across approximately 25
years from the sediment egg bank isolated from a dated sediment core. Using whole genome
sequencing (WGS) of resurrected and extant individuals, we conducted numerous population
genomic analyses to depict population structure over time, reconstruct the demographic history,
and identify outlier genomic regions in the data. Additionally, we assayed a subset of genotypes
for tolerance to salinity. Specifically, we were interested in testing two main hypotheses; first,
we believed that the outlier regions would contain genes related to osmoregulation as selection
would favor higher salinity tolerance. Second, this would be reflected in a higher mean tolerance
of the more recent subpopulations. In addition, we identify a list of candidate genes, which likely
influence phenotypic variation throughout the genome, and that can be targeted for further study.
111

Results:

We studied the *D. pulicaria* population of Tanners Lake (TL; 44°57'02.2"N 92°58'54.2"W), a
small suburban hardwater lake located in Oakdale, Minnesota. The watershed includes
approximately 32% impervious surfaces including parking lots, interstate highways and
residential development (30). TL has received significant inputs of chloride from the watershed
primarily in the form of the road deicer NaCl (524 kg $Cl^- ha^{-1} yr^{-1}$) (33). The chloride
concentration of TL has increased significantly in the last 75 years, from approximately 1-2 mg
$Cl^- L^{-1}$ on average to over 150 mg $Cl^- L^{-1}$ (34, 35).

Population Structure and Genetic Divergence:

The 54 clones selected for sequencing were separated into two clusters based on the first two
principal component axes explaining 7.6% and 4.6% of the variance in the SNP data,
respectively (Figure 1A). These two clusters largely separated the clones by depth with the older
clones (layers 16-18 cm, 18-20 cm, and 22-24cm) and more recent clones (Lake Clones, 2-4 cm,
6-8 cm, and 10-12 cm) forming the two groupings. Subsequently, we decided to assign the
clones into three groups for the remainder of the analyses. We designated these as DEEP,
encompassing all clones from 16-24 cm in depth (n = 18) and date from the mid to late 1990s.
The second designation, called MID, encompassed all clones from 6-12 cm in depth (n = 18) and
date from the mid to late 2000s. The third and final designation, referred to as TOP hereafter,
included all clones collected from the water column and 2-4 cm in depth in the core (n = 18)
spanning from 2016 to 2019. Although somewhat arbitrary in our assignment, we based the
groupings on two observations: firstly, the TOP and DEEP subpopulations are delineated by the

PCA clusters, and thus warranted separate assignments. Secondly, the MID subpopulation, while
closely related to TOP, was intermediate in the time-scale –approximately 10 years prior to TOP
and approximately 10 years after DEEP– and thus formed an appropriate intermediate grouping.
Conveniently, this scheme allowed each grouping to achieve equal sample size (i.e., 18 samples).
Site-wise F_{st} was low across time, with an overall estimate of approximately 0.016 (Figure 1B).
However, F_{st} ranged from around 0 to as high as 0.398, with most sites having an F_{st} of
essentially 0. The pairwise genetic distance between the temporal subpopulations was related to
the temporal distance with the TOP vs DEEP comparison being highest (0.019) and the MID
population being intermediate to both (Figure 1C).

Estimation of Effective Population Size and Simulations:

We were able to estimate the effective population size of the TL *D. pulicaria* population using
both linkage disequilibrium (LD) and coalescent simulations. The LD-based results using just
samples from the lake clone (LC) isolates showed that over the last few hundred generations,
there was a period of population expansion and contraction (Figure S1). The population reached
a peak N_e of 4000-6500 approximately 150 generations ago and the population has contracted
recently to an N_e of around 2000. We interpret "generations" here to be sexual generations (i.e.,
LD is related to recombination and asexual generations are ameiotic). Since sex may occur once
or at most a few times per year in stable lake habitats (36, 37), we interpret a single generation to
be equivalent to one year. The FSC2.7 (38) run with the highest likelihood estimated N_e to be
2931 individuals and had a signal of population contraction with a population growth rate of
2.629×10^{-5}. The estimate of effective population size was within the 95% confidence for 100
parametric bootstraps, however the estimate for population growth rate was not and the
confidence intervals included zero (table S1). We ran 100 simulations of FSC2.7 using the
maximum likelihood estimates established above and simulations were pooled to develop a
distribution of expected F_{st} values from ~110000 SNPs. Testing each LD-pruned SNP against
this distribution and correcting for multiple testing using false discovery rate (FDR) resulted in
178 outlier SNPs with corrected one-tailed p-values above a p = 0.05 significance threshold
(Figure 2). There were outliers on every chromosome, ranging from a low of 2 SNPs (CHR 08)
to a high of 42 (CHR 04).

Genes surrounding F_{st} outliers and GO term enrichment and Variant Annotation:

We extracted 286 genes near the outlier SNPs with known function in the *D. pulicaria* genome.
GO term enrichment analysis with PantherDB webtool (39) yielded 59 enriched terms for this
list of genes after correction for FDR (Figure 3). The enriched terms and p-values are available
in table S2. Notable among the enriched terms for molecular function are chloride channel
activity (GO:0005254; p = 0.00925); however, many different ion and channel terms were
enriched. After running variant effect prediction (40) on all the SNPs called in the population, we
identified 17181 variants with "high" predicted effects. Intersecting this list with outlier genes,
we found 78 of the 286 outlier genes had high effect variants, only one of these, Chloride
Channel 2 isoform X2 (*clcn2-x2*), located on chromosome 5 had any obvious relation to
osmoregulation or salinity tolerance. *Clcn2-x2* has five SNPs of high effect, including four
premature stop codon changes and a splice donor change that would likely severely interrupt

protein function. It is tightly linked to the F_{st} outlier site at CHR05:12238, which is an intronic
SNP within *clcn2-x2*. In addition to the high effect mutations, this gene has a total of 28
missense mutations and 37 synonymous mutations classified as moderate and low impact
respectively. Of the clones surveyed for chloride tolerance (see next section), we identified that
the two most tolerant clones are homozygous for the wildtype (i.e., functional) allele at *clcn2-x2*.
Overall, 252 of the outlier genes had effects that included moderate impacts to function such as a
missense SNP.

However, *clcn2-x2*'s behavior as an outlier is not consistent with a single large effect
locus. The SNP at CHR05:12238 is the 88[th] most differentiated SNPs among the 178 outliers
with an overall F_{st} of just 0.1985 (FDR corrected p = 0.0269). The top 11 outlier SNPs (F_{st}
0.3983 – 0.3175) all had F_{st} unobserved under neutral demography. One of the SNPs is an
intergenic variant, while the 10 remaining SNPs are intronic SNPs associated with 13 different
genes, 10 of which have annotated functions (table S3). The SNP located at CHR04:13394600 is
the most differentiated SNP observed and is associated with the genes *pank4* and *tda6*. The
remaining genes span from functional categories including the structural protein collagen alpha-1
chain and the calcium-binding protein *fstl5*, to several genes related to development comprising
still life (41)*, daam1* (42)*, and pnt* (43), which contains two outlier SNPs*, rap1gtp* (44)*, and
rotund (45)*.

Chloride Tolerance LC$_{50}$:

Clone-specific estimates of 96-hour LC$_{50}$ ranged from a low of 584.91 mg/L Cl$^-$ (clones 10-12-
12A & 18-20-05A) to a high of 1047.16 mg/L Cl$^-$ for (Clone LC-06). We observed a main effect
of subpopulation in the Kruskal-Wallis test (p = 0.006; Figure 4). Post-hoc testing using a
pairwise Wilcox test found that the Lake subpopulation (i.e., TOP) was more tolerant on average
than either the DEEP (22-24 cm & 18-20 cm) or MID (10-12 cm) subpopulations (p = 0.0076).
The DEEP and MID subpopulations did not differ from one another (p = 0.9073).

Discussion:

Our resurrection ecology (RE) study across ~25-year temporal contrast has provided
phenotypic and molecular evidence of rapid evolution of salinity tolerance in the TL *Daphnia*
pulicaria population. Using whole genome resequencing to analyze population genetic structure
and demographic history, we have identified genomic regions putatively under selection, with
support from LC$_{50}$ chloride tolerance assays. We found that Chloride Channel-2 isoform X2
(*clcn2-x2*) has a unique mutational history and may be affecting salinity tolerance in this
population, as one of many genes under selection in this population. We find support for both of
our main hypotheses — genes related to osmoregulation are enriched in outlier regions and that
the population shows increasing tolerance to salinity over time.

Demographic History and Subpopulation Structure:

A goal of our study was to establish a demographic model for the TL population. This
would enable the modeling of population genetic summary statistics (i.e., F_{st}) for establishment
of a distribution of expected or null estimates of F_{st} based on the demographic history (discussed

below). We used two complementary methods – LD-based methods and coalescent simulation to understand the recent demographic history of the TL population. Each method provides incomplete information, and each has unique biases that need to be considered. Furthermore, despite N_e being a critical parameter in population genetics, it is notoriously difficult to estimate (46). For instance, it appears that SFS (coalescent) based methods are underpowered for recent demographic history; however larger sample sizes may ameliorate some of these effects (47). This may also explain why it was difficult to accurately estimate the population growth rate parameter. The LD-based method we employed here appears to be strongly affected by the number of homozygotes in the sample (48), and thus, is sensitive to analyzing genomes from different generations. This prevented us from modeling each subpopulation separately. Additionally, it also appears to be inaccurate for the first few generations, giving unrealistically small numbers (see Figure S1). As such, we used these methods in tandem to increase the confidence in our estimation of the demographic parameters. Both methods were concordant that recent N_e was approximately 2000-3000 individuals, and both showed signatures of recent population contraction. This effective population size is not unexpected for *D. pulicaria* because populations are thought to delay sex and recombination for long periods of time (a year or more) (36, 37). During this time, clonal selection is thought to winnow down the population to a collection of ecologically-equivalent clones (49). These facts, taken together with the current understanding of zooplankton metapopulation dynamics (50, 51) suggests that *Daphnia* populations should be small and insular.

Despite the small and insular nature of the TL population, it appears that drift does not always predominate. Our PCA results demonstrate at least on a decadal scale (i.e., between MID and TOP) that population structure is consistent. This likely reflects the dominance of the few ecologically equivalent clones predicted by Lynch and Spitze (37). It is likely that only after a rapid or pronounced change in environment one can expect to see a population structure across time, as reflected in other studies of *Daphnia* utilizing RE (15, 52). Salinization in TL is ongoing and likely started in the mid-20th century with the onset of widespread use of road deicing salts (53). The sediment intervals from which we recovered the deepest samples reflect a period of change in the sediment egg bank (e.g., low ephippial fluxes) that have been previously reported (30), and suggest a period of rapid environmental change in this lake. However, because we were unable to hatch eggs from before the mid-1990s in TL, it is impossible to determine if the structuring we observed was ongoing or more sudden. Regardless, since the clones continue closely related across time at both the nuclear and mitochondrial genomic levels, it remains unlikely that the population became extinct and was recolonized by migrant genotypes (Fig. 1C, S2).

Outliers and the Identification of Candidate Genes:

The specification of a demographic model for the TL lake population allowed us to establish the presence of statistical significance for each site in the genome. This analysis revealed 178 regions with F_{st} beyond what we could reasonably expect based on demography alone. Genes surrounding these outliers had many functions, but as expected, genes involved in osmoregulation were among the most enriched in the dataset according to GO terms for

molecular function. The presence of hundreds of outlier genes suggests that salinity tolerance has
a complex genetic architecture and that tolerance to increased salinity requires changes at many
loci of small effect. Transcriptomic studies of clonal isolates from within the *D. pulex* complex
(to which *D. pulicaria* belongs) supports this assertion (54). Indeed, cross-referencing these
outlier genes with different mutational types (e.g., missense SNPs) showed that many of these
genes have sites that may be under selection. However, the only gene with known functional role
in osmoregulation with high effect mutations was *clcn2-x2*. *Daphnia* are known as hypo-
osmoregulators, meaning they attempt to maintain their hemolymph solute concentration above
the ambient media concentration (55). Osmoregulation occurs within the gill-epithelium of
Daphnia, and chloride channels like *clcn2* play a major role in shuttling Cl^- ions into the
hemolymph across the apical membrane of the gill epithelium (56). The constellation of
premature stop codon mutations in this gene means that the protein is very unlikely to function
properly because the channel is too short to pass through the cell membrane. Individuals are at
least heterozygous for each of the four premature stop codon SNPs and the wild-type gene
sequence annotated in the reference. Thus, they can produce a functioning protein. The two most
tolerant clones in this study were both homozygous for the functional allele. This suggests that
clcn2 is a critical gene requiring at least one functional copy and it explains some portion of the
variance in acute salinity tolerance. However, a larger sample of genotyped and phenotyped
individuals would be required to further validate this statistically.

The most differentiated genes in the genome were involved in a few key functions. There
was a concentration of outlier SNPs in genes involved in regulating Rho-like GTPases which are
related in the regulation of actin filaments and neuronal development. The reason why these
genes are the most differentiated is not initially clear and may involve a few mechanisms that
require further explanation. First, these genes may be related to ongoing evolution of *D.
pulicaria* during its recent invasion into lake ecosystems. Studies of the comparative genomics
of *D. pulicaria* and its ecological-sister species *D. pulex* have noted that the differing growth and
development phenotypes in these two species is underlain by distinct changes in developmental
genes (57, 58). Thus, we may be picking up a global signal of adaptation in our study. Second,
increased salinity tolerance may require the accommodation of new developmental trajectories
through phenotypic plasticity. Evolution via plasticity (59) requires that central developmental
pathways serve as fuel for phenotypic differentiation (60). Furthermore, plasticity is thought to
play an important role in rapid adaptation in *Daphnia*, a pattern we observe in the TL population
(61–63). Regardless, testing either of these assertions would be difficult because unlike other
model arthropods, *Daphnia* development is not well explored.

Conclusions:

Resurrection ecology (RE) when paired with whole genome sequencing provides a
unique and powerful way to study rapid evolution of populations *in situ.* Integration of candidate
loci identified with RE into study designs including breeding experiments (64), forward mutation
screens (65) or CRISPR technology (66) will provide unrivaled insight for the genetic
architecture of complex, ecologically-relevant traits in the wild. Such an approach would be
beneficial here, for example in validating the effect of the null *clcn2-x2* allele observed in the TL

population on salinity tolerance. Overall, here RE provides both molecular and phenotypic evidence of rapid adaptation in the TL population. Thus, the persistence of *D. pulicaria* in this severely salinized lake is likely the result of this rapid adaptation. Our findings indicate that keystone aquatic species such as lake *Daphnia* may continue to thrive in lakes that exceed current water quality limits for salinization and thus maintaining the stability of the food webs and ecosystem services they support.

Methods:

Clone bank and Sequencing:

On 2 July 2019, we collected duplicate sediment cores in TL from a 14 m deep station following Wright (67). During the same period, we also collected *D. pulicaria* from the active plankton community using several vertical tows of a Wisconsin net at the core sampling station. Animals were isolated as single individuals in 125 mL plastic (screw-capped) cups in COMBO media (68). A total of 10 of these clones were established in laboratory culture. Resting eggs (ephippia) collected from throughout the cores were collected according to the methods outlined in Wersebe et al. (30). Ephippia identified as *D. pulicaria* were subjected to hatching protocol described (15) (see supplemental). These hatchling individuals were expanded in culture to establish upwards of 10 clones per sediment layer to establish a clone bank. 54 individuals from the clone bank were selected for DNA extraction and whole genome sequencing (average 10X) on an Illumina NovaSeq by the Oklahoma Medical Research Foundation.

Bioinformatics:

Raw sequencing reads were quality trimmed and adaptor contamination removed using Trimmomatic (69). Quality trimmed reads were aligned to the chromosome-level *D. pulicaria* genome assembly (57) using the BWA mem algorithm (70). The resulting files were piped through samtools (71) to mark duplicates, fix mates and sort the bam files. We called variants using the bcftools mpileup and call pipeline using all individuals together (72). Using bcftools, the resulting BCF files were concatenated together into a single genome-wide file and quality filtered to a set of high confidence biallelic single nucleotide polymorphisms (SNPs).

Population Structure and Genetic Divergence:

The variants in the final quality-filtered VCF file were pruned for linkage disequilibrium (LD) using Plink (73) independent pairwise function. The resulting variants were further filtered to SNPs with 0% missingness to a set 27854 genome wide SNPs. We conducted Principal Components Analysis (PCA) in R (Version 4.2.0; R core team 2022) using the packages *adegenet* and *vcfr* (74, 75). From the PCA results, we determined that the samples could be assigned to three "sub-populations" according to the depth of their recovery (see results). Using these three subpopulations as designations, we estimated site-wise F_{st} using the basic.stats function and mean pairwise genetic distance genet.dist function in the R package *hierfstat* (76).

Estimation of Effective Population Size and Simulations:

To parameterize our tests for selection, we sought to identify the recent demographic history of
the TL *D. pulicaria* population. To accomplish this, we used two different methods to estimate
effective population size (N_e) and growth trajectory of the population. The first method,
implemented in GONE (48) uses LD to estimate the recent population history. This method is
robust to non-equilibrium histories such as selection (77), however is suitable only for sample
pools collected from the same generation. Thus, for this method we used only the samples
collected from the water column in 2019. We ran six independent runs of this method using
random subsets of 600,000 SNPs called in the population and a constant recombination rate of
7.2 cm/mb estimated from the *D. pulicaria* genetic map (57). In addition to LD-based methods,
we estimated demographic parameters of the TL population - estimating N_e and the population
growth rate - by fitting a demographic model to the folded site frequency spectrum (SFS)
implemented in FastSimcoal 2.7 (FSC2.7) (38). With the LD pruned SNPs used above, we
created folded 2-D SFS from the three temporal subpopulations. Using 100 independent runs of
FSC2.7, we chose the run that maximized the likelihood of the observed data. Each run
conducted 1million coalescent simulations and used 40 brent maximization cycles.

Further, we estimated empirical p-values for the site-wise F_{st} estimates using simulation (78). We
choose the best fitting demographic model estimated in FSC2.7 and conducted 100 separate runs
of this model to simulate approximately 1100 SNPs for each run. For each run, we estimated F_{st}
using basic.stats function in HIERSTAT (76). We pooled each simulation into an empirical
distribution of probable F_{st} values under the best fitting demographic model. We tested each
observed site-wise F_{st} estimate against this empirical distribution to estimate a p-value. We
corrected these p-values for multiple testing using a false discovery rate in R using the p.adjust
function.

Genes surrounding Fst outliers, GO term enrichment, and Variant Effects:

We extracted the genes surrounding F_{st} outliers ($p \leq 0.05$ after correction) in 10 Kb windows
using the *D. pulicaria* RefSeq annotation (release 100, SC_F0-13Bv2) using bedtools (79). Next,
we created Panther Generic Mappings for the genes with known annotations using each gene's
protein sequences following the method outlined in (39). Using the generic mappings, we tested
for molecular function Gene Ontology (GO) term enrichment by testing against the *Daphnia*
pulex gene list using the PantherDB webtool using a false discovery rate correction (39). In
addition to testing for GO term enrichment, we also surveyed all genes for potential SNPs and
small indels mutations potentially driving selection using Ensembl's Variant Effect Predictor
(VEP) (40). We built a custom database using the RefSeq annotations in GFF format following
the developer's protocol. We extracted all "High" (e.g., premature stops) and "Moderate" (e.g.,
missense SNPs) impact mutations predicted and cross-referenced these with the genes near F_{st}
outliers.

Chloride Tolerance:

We selected a subset of 30 (out of the 54 sequenced) clones from the Lake (n = 10), 10-12 cm
sediment layer (n =10), 22-24 cm (n = 7), and 18-20 cm (n = 3) subpopulations for estimating
clone-specific tolerance to chloride. See the Supplemental Materials for details on the

experimental set-up. We estimated LC_{50} for each clone separately by fitting a reduced-bias
generalized linear model (GLM) to the survival curve using the R package *brglm* (80). Data were
not normally distributed, nor did they meet the assumption of equal variances, so we performed a
non-parametric ANOVA (Kruskal-Wallis) test on the LC_{50} estimates to test for differences in the
mean LC_{50} value for each subpopulation.

Acknowledgments:

We would like to thank M. Edlund who facilitated the collection of the Tanners Lake sediment
cores. T. Curb, Z. Arnold, A. Twumasi-Mensah, and A. Hemani, helped with the LC_{50}
experiments. G. Wiley aided in the preparation of sequencing libraries. The University of
Oklahoma Biological Station and the St. Croix Watershed Research Station facilitated field
work. Funding for this study was provided by the University of Oklahoma Department of
Biology Adams Summer Scholarship, Robberson Graduate College Grant, Hill Fund for
Research in Biology, Graduate Student Senate Research Grant, and the Biogeography of
Behavior seed grant (NSF DBI-2021880) awarded to MJW in support of graduate research. Any
opinions, findings and conclusions or recommendations expressed in this material are those of
the authors and do not necessarily reflect the views of the National Science Foundation. This
manuscript represents a portion of MJW's doctoral dissertation at The University of Oklahoma.

Data Availability:

390 Sequencing reads will be made available on the NCBI SRA upon acceptance. Data and
391 appropriate metadata from this study will be archived in the University of Oklahoma's ShareOK
392 data repository (https://shareok.org). Code used in the analysis is available on GitHub at
393 https://github.com/mwersebe/Tanner_lake_genomics.

Competing Interests:

395 The authors declare no competing interests.

396
397

Works Cited:

1. S. M. Rudman, *et al.*, What genomic data can reveal about eco-evolutionary dynamics. *Nat. Ecol. Evol.* **2**, 9–15 (2018).

2. N. G. Hairston, *et al.*, Lake ecosystems: Rapid evolution revealed by dormant eggs. *Nature* **401**, 446–446 (1999).

3. F. Mallard, V. Nolte, R. Tobler, M. Kapun, C. Schlötterer, A simple genetic basis of adaptation to a novel thermal environment results in complex metabolic rewiring in Drosophila. *Genome Biol.* **19**, 119 (2018).

4. P. W. Messer, S. P. Ellner, N. G. Hairston, Can Population Genetics Adapt to Rapid Evolution? *Trends Genet.* **32**, 408–418 (2016).

5. P. W. Messer, D. A. Petrov, Population genomics of rapid adaptation by soft selective sweeps. *Trends Ecol. Evol.* **28**, 659–669 (2013).

6. A. Eyre-Walker, P. D. Keightley, The distribution of fitness effects of new mutations. *Nat. Rev. Genet. 2007 88* **8**, 610–618 (2007).

7. P. D. Keightley, A. Eyre-Walker, What can we learn about the distribution of fitness effects of new mutations from DNA sequence data? *Philos. Trans. R. Soc. B Biol. Sci.* **365**, 1187–1193 (2010).

8. A. Long, G. Liti, A. Luptak, O. Tenaillon, Elucidating the molecular architecture of adaptation via evolve and resequence experiments. *Nat. Rev. Genet.* **16**, 567–582 (2015).

9. C. Schlötterer, R. Kofler, E. Versace, R. Tobler, S. U. Franssen, Combining experimental evolution with next-generation sequencing: A powerful tool to study adaptation from standing genetic variation. *Heredity (Edinb).* **114**, 431–440 (2015).

10. T. L. Turner, A. D. Stewart, A. T. Fields, W. R. Rice, A. M. Tarone, Population-Based Resequencing of Experimentally Evolved Populations Reveals the Genetic Basis of Body Size Variation in Drosophila melanogaster. *PLoS Genet.* **7**, e1001336 (2011).

11. R. E. Lenski, Convergence and divergence in a long-term experiment with bacteria. *Am. Nat.* **190**, S57–S68 (2017).

12. M. Dehasque, *et al.*, Inference of natural selection from ancient DNA. *Evol. Lett.* **4**, 94–108 (2020).

13. L. J. Weider, P. D. Jeyasingh, D. Frisch, Evolutionary aspects of resurrection ecology: Progress, scope, and applications-An overview. *Evol. Appl.* **11**, 3–10 (2018).

14. M. Gros-Balthazard, *et al.*, The genomes of ancient date palms germinated from 2,000 y old seeds. *Proc. Natl. Acad. Sci. U. S. A.* **118** (2021).

15. D. Frisch, *et al.*, A millennial-scale chronicle of evolutionary responses to cultural eutrophication in *Daphnia*. *Ecol. Lett.* **17**, 360–368 (2014).

16. W. C. Kerfoot, J. A. Robbins, L. J. Weider, A new approach to historical reconstruction: Combining descriptive and experimental paleolimnology. *Limnol. Oceanogr.* **44**, 1232–

435   1247 (1999).

17. W. C. Kerfoot, L. J. Weider, "Experimental paleoecology (resurrection ecology): Chasing Van Valen's Red Queen hypothesis" (2004).

18. A. Chaturvedi, *et al.*, Extensive standing genetic variation from a small number of founders enables rapid adaptation in Daphnia. *Nat. Commun.* **12**, 4306 (2021).

19. C. Cousyn, *et al.*, Rapid, local adaptation of zooplankton behavior to changes in predation pressure in the absence of neutral genetic changes. *Proc. Natl. Acad. Sci. U. S. A.* **98**, 6256–60 (2001).

20. D. Frisch, *et al.*, Paleogenetic records of *Daphnia pulicaria* in two North American lakes reveal the impact of cultural eutrophication. *Glob. Chang. Biol.* **23**, 708–718 (2017).

21. W. Lampert, U. Sommer, *Limnoecology: The Ecology of Lakes and Streams*, 2nd Ed. (Oxford University Press, 2007).

22. Ž. Ogorelec, C. Wunsch, A. J. Kunzmann, P. Octorina, J. I. Navarro, Large daphniids are keystone species that link fish predation and phytoplankton in trophic cascades. *Fundam. Appl. Limnol.* **194**, 297–309 (2021).

23. J. R. Walsh, S. R. Carpenter, M. J. Vander Zanden, Invasive species triggers a massive loss of ecosystem services through a trophic cascade. *Proc. Natl. Acad. Sci. U. S. A.* **113**, 4081–5 (2016).

24. A. J. Reid, *et al.*, Emerging threats and persistent conservation challenges for freshwater biodiversity. *Biol. Rev.* **94**, 849–873 (2019).

25. S. S. Kaushal, *et al.*, Freshwater salinization syndrome on a continental scale. *Proc. Natl. Acad. Sci.* (2018) https:/doi.org/10.1073/pnas.1711234115.

26. H. A. Dugan, *et al.*, Lakes at Risk of Chloride Contamination. *Environ. Sci. Technol.* **54**, 6639–6650 (2020).

27. H. A. Dugan, *et al.*, Salting our freshwater lakes. *Proc. Natl. Acad. Sci. U. S. A.* **114**, 4453–4458 (2017).

28. W. D. Hintz, R. A. Relyea, A review of the species, community, and ecosystem impacts of road salt salinisation in fresh waters. *Freshw. Biol.* **64**, 1081–1097 (2019).

29. D. Cunillera-Montcusí, *et al.*, Freshwater salinisation: a research agenda for a saltier world. *Trends Ecol. Evol.* **0** (2022).

30. M. J. Wersebe, M. B. Edlund, L. J. Weider, Does salinization impact long-term Daphnia assemblage dynamics? Evidence from the sediment egg bank in a small hard-water lake. *Limnol. Oceanogr. Lett.* **n/a** (2021).

31. S. Arnott, *et al.*, Road salt impacts freshwater zooplankton at concentrations below current water quality guidelines. *Environ. Sci. Technol.* (2020) https:/doi.org/10.1021/acs.est.0c02396 (August 11, 2020).

32. H. W. D., *et al.*, Current water quality guidelines across North America and Europe do not

472   protect lakes from salinization. *Proc. Natl. Acad. Sci.* **119**, e2115033119 (2022).

33.   E. V. Novotny, D. Murphy, H. G. Stefan, Increase of urban lake salinity by road deicing salt. *Sci. Total Environ.* **406**, 131–144 (2008).

34.   J. M. Ramstack, S. C. Fritz, D. R. Engstrom, S. A. Heiskary, The Application of a Diatom-based Transfer Function to Evaluate Regional Water-Quality Trends in Minnessota Since 1970. *J. Paleolimnol.* **29**, 79–94 (2003).

35.   J. M. Ramstack, S. C. Fritz, D. R. Engstrom, Twentieth century water quality trends in Minnesota lakes compared with presettlement variability. *Can. J. Fish. Aquat. Sci.* **61**, 561–576 (2004).

36.   D. E. Allen, M. Lynch, THE EFFECT OF VARIABLE FREQUENCY OF SEXUAL REPRODUCTION ON THE GENETIC STRUCTURE OF NATURAL POPULATIONS OF A CYCLICAL PARTHENOGEN. *Evolution (N. Y).* **66**, 919–926 (2012).

37.   C. E. Cáceres, A. J. Tessier, Incidence of diapause varies among populations of Daphnia pulicaria. *Oecologia* **141**, 425–431 (2004).

38.   L. Excoffier, *et al.*, fastsimcoal2: demographic inference under complex evolutionary scenarios. *Bioinformatics* **37**, 4882–4885 (2021).

39.   H. Mi, *et al.*, PANTHER version 16: A revised family classification, tree-based classification tool, enhancer regions and extensive API. *Nucleic Acids Res.* **49**, D394–D403 (2021).

40.   W. McLaren, *et al.*, The Ensembl Variant Effect Predictor. *Genome Biol.* **17**, 122 (2016).

41.   S. Masaki, *et al.*, Still life, a Protein in Synaptic Terminals of Drosophila Homologous to GDP-GTP Exchangers. *Science (80-.).* **275**, 543–547 (1997).

42.   P. Aspenström, N. Richnau, A.-S. Johansson, The diaphanous-related formin DAAM1 collaborates with the Rho GTPases RhoA and Cdc42, CIP4 and Src in regulating cell morphogenesis and actin dynamics. *Exp. Cell Res.* **312**, 2180–2194 (2006).

43.   C. Klambt, The Drosophila gene pointed encodes two ETS-like proteins which are involved in the development of the midline glial cells. *Development* **117**, 163–176 (1993).

44.   C. Fangli, B. Margaret, R. K. T., Q. Adrian, H. I. K., Biological characterization of Drosophila Rapgap1, a GTPase activating protein for Rap1. *Proc. Natl. Acad. Sci.* **94**, 12485–12490 (1997).

45.   S. E. St Pierre, M. I. Galindo, J. P. Couso, S. Thor, Control of Drosophila imaginal disc development by rotund and roughened eye: differentially expressed transcripts of the same gene encoding functionally distinct zinc finger proteins. *Development* **129**, 1273–1281 (2002).

46.   J. Wang, E. Santiago, A. Caballero, Prediction and estimation of effective population size. *Heredity (Edinb).* **117**, 193–206 (2016).

47.   J. D. Robinson, A. J. Coffman, M. J. Hickerson, R. N. Gutenkunst, Sampling strategies for frequency spectrum-based population genomic inference. *BMC Evol. Biol.* **14**, 254 (2014).

48. E. Santiago, *et al.*, Recent Demographic History Inferred by High-Resolution Analysis of
Linkage Disequilibrium. *Mol. Biol. Evol.* **37**, 3642–3653 (2020).

49. M. Lynch, K. Spitze, "Evolutionary Genetic of Daphnia" in *Ecological Genetics*, L. Real,
Ed. (Princeton University Press, 1994).

50. L. De Meester, A. Gómez, B. Okamura, K. Schwenk, The Monopolization Hypothesis and
the dispersal-gene flow paradox in aquatic organisms. *Acta Oecologica* **23**, 121–135
(2002).

51. L. De Meester, J. Vanoverbeke, K. De Gelas, R. Ortells, P. Spaak, Genetic structure of
cyclic parthenogenetic zooplankton populations a conceptual framework. *Arch. für*
Hydrobiol., 217–244 (2006).

52. D. Frisch, *et al.*, Paleogenetic records of *Daphnia pulicaria* in two North American lakes
reveal the impact of cultural eutrophication. *Glob. Chang. Biol.* **23**, 708–718 (2017).

53. S. E. G. Findlay, V. R. Kelly, Emerging indirect and long-term road salt effects on
ecosystems. *Ann. N. Y. Acad. Sci.* **1223**, 58–68 (2011).

54. L. C. Latta, L. J. Weider, J. K. Colbourne, M. E. Pfrender, The evolution of salinity
tolerance in *Daphnia*⍰: a functional genomics approach. *Ecol. Lett.* **15**, 794–802 (2012).

55. J. Hervé Lignot, G. Charmantier, "Chapter 8: Osmoregulation and Excretion" in *Natural*
History of Crustacea Volume IV: Physiology, E. S. Chang, M. Thiel, Eds. (Oxford
University Press, 2015).

56. M. B. Griffith, Toxicological perspective on the osmoregulation and ionoregulation
physiology of major ions by freshwater animals: Teleost fish, crustacea, aquatic insects,
and Mollusca. *Environ. Toxicol. Chem.* **36**, 576–600 (2017).

57. M. J. Wersebe, R. E. Sherman, P. D. Jeyasingh, L. J. Weider, The roles of recombination
and selection in shaping genomic divergence in an incipient ecological species complex.
Mol. Ecol. **n/a** (2022).

58. C. E. Jackson, *et al.*, Chromosomal rearrangements preserve adaptive divergence in
ecological speciation. *bioRxiv*, 2021.08.20.457158 (2021).

59. N. A. Levis, D. W. Pfennig, Evaluating "Plasticity-First" Evolution in Nature: Key
Criteria and Empirical Approaches. *Trends Ecol. Evol.* **31**, 563–574 (2016).

60. M. J. West-Eberhard, Developmental plasticity and the origin of species differences. *Proc.*
Natl. Acad. Sci. **102**, 6543–6549 (2005).

61. K. I. Brans, L. De Meester, City life on fast lanes: Urbanization induces an evolutionary
shift towards a faster lifestyle in the water flea *Daphnia*. *Funct. Ecol.* **32**, 2225–2240
(2018).

62. N. G. Hairston, *et al.*, Natural selection for grazer resistance to toxic cyanobacteria:
Evolution of phenotypic plasticity? *Evolution (N. Y).* **55**, 2203–2214 (2001).

63. A. Scoville, M. Pfrender, Phenotypic plasticity facilitates recurrent rapid adaptation to
introduced predators. *Proc. Natl. Acad. Sci.* **107**, 4260–4263 (2010).

64. R. E. Sherman, R. Hartnett, E. L. Kiehnau, L. J. Weider, P. D. Jeyasingh, Quantitative
genetics of phosphorus content in the freshwater herbivore, *Daphnia pulicaria. J. Anim.*
Ecol. **00**, 1365-2656.13419 (2021).

65. M. Snyman, T. V Huynh, M. T. Smith, S. Xu, The genome-wide rate and spectrum of
EMS-induced heritable mutations in the microcrustacean Daphnia: on the prospect of
forward genetics. *Heredity (Edinb).* **127**, 535–545 (2021).

66. C. Hiruta, K. Kakui, K. E. Tollefsen, T. Iguchi, Targeted gene disruption by use of
CRISPR/Cas9 ribonucleoprotein complexes in the water flea Daphnia pulex. *Genes to*
Cells **23**, 494–502 (2018).

67. H. E. Wright, Coring tips. *J. Paleolimnol.* **6**, 37–49 (1991).

68. S. S. Kilham, D. A. Kreeger, S. G. Lynn, C. E. Goulden, L. Herrera, COMBO: a defined
freshwater culture medium for algae and zooplankton. *Hydrobiologia* **377**, 147–159
(1998).

69. A. M. Bolger, M. Lohse, B. Usadel, Trimmomatic: a flexible trimmer for Illumina
sequence data. *Bioinformatics* **30**, 2114–2120 (2014).

70. H. Li, R. Durbin, Fast and accurate short read alignment with Burrows-Wheeler
transform. *Bioinformatics* **25**, 1754–1760 (2009).

71. H. Li, *et al.*, The Sequence Alignment/Map format and SAMtools. *Bioinformatics* **25**,
2078–2079 (2009).

72. P. Danecek, *et al.*, Twelve years of SAMtools and BCFtools. *Gigascience* **10**, 1–4 (2021).

73. C. C. Chang, *et al.*, Second-generation PLINK: Rising to the challenge of larger and richer
datasets. *Gigascience* **4**, 7 (2015).

74. T. Jombart, adegenet: a R package for the multivariate analysis of genetic markers.
Bioinformatics **24**, 1403–1405 (2008).

75. B. J. Knaus, N. J. Grünwald, vcfr: a package to manipulate and visualize variant call
format data in R. *Mol. Ecol. Resour.* **17**, 44–53 (2017).

76. J. Goudet, hierfstat, a package for r to compute and test hierarchical F-statistics. *Mol.*
Ecol. Notes **5**, 184–186 (2005).

77. I. Novo, E. Santiago, A. Caballero, The estimates of effective population size based on
linkage disequilibrium are virtually unaffected by natural selection. *PLOS Genet.* **18**,
e1009764 (2022).

78. K. E. Lotterhos, M. C. Whitlock, The relative power of genome scans to detect local
adaptation depends on sampling design and statistical method. *Mol. Ecol.* **24**, 1031–1046
(2015).

79. A. R. Quinlan, I. M. Hall, BEDTools: a flexible suite of utilities for comparing genomic
features. *Bioinformatics* **26**, 841–842 (2010).

80. I. Kosmidis, E. C. Kenne Pagui, N. Sartori, Mean and median bias reduction in

generalized linear models. *Stat. Comput.* **30**, 43–59 (2020).

Figures:

Figure 1: Genetic structure, divergence, and distance across time in Tanners Lake. A) PCA biplot of all clones color-coded according to the depth of recovery in the core (cm). PC1 and PC2 explain 7.6 % and 4.6 % of variance observed in the SNP data, respectively. Older clones (22-24 cm, 18-20 cm, & 16-18 cm) form a cluster distinct from more recent clones (LC, 2-4 cm, 6-8 cm and 10-12 cm). B) Observed Weir and Cockerham site-wise F_{st} estimates. Overall F_{st} was low (red vertical line) at only 0.0169. C) Pair-wise genetic chord distances between the three temporal subpopulations (TOP, MID and DEEP).

Figure 2: Genome-wide F_{st} Manhattan plots. Each panel represents a chromosome (1-12), each point designates a Single Nucleotide Polymorphism (SNP), the horizontal line indicates genome-wide significance at p = 0.05. Points above the red line are statistical outliers.

Figure 3: Enriched Gene Ontology (GO) terms for molecular function. Enriched terms are set
along the X-axis. The height of each bar indicates the term's fold enrichment in the analyzed
gene list, and the bar color denotes its FDR corrected p-value. In total, 59 terms were enriched
among the 286 genes found near outlier SNPs.

606
607
608
609

613 **Figure 4:** 96-hour lethal concentration-50% assay (LC_{50}) estimates for clones evaluated in the
614 salinity tolerance experiments. Box plots indicate population medians and variances, each point
615 is a clone. Overall, non-parametric ANOVA (Kruskal-Wallis) indicated a significant main effect
616 of population. Post-hoc testing (pairwise Wilcox) confirmed that the TOP population was more
617 tolerant than either the MID or DEEP populations, which were not different from one another.