Angiogenesis precedes myogenesis during regeneration

following biopsy injury in skeletal muscle

Nicole L. Jacobsen[1†], Aaron B. Morton[1†], Steven S. Segal[1-4*]

[1]Department of Medical Pharmacology and Physiology, University of Missouri; Columbia,

MO

[2]Dalton Cardiovascular Research Center; Columbia, MO

[3]Department of Biomedical Sciences, University of Missouri; Columbia, MO

[4]Department of Biomedical, Biological, and Chemical Engineering, University of Missouri;

Columbia, MO

*Corresponding author. Email: segalss@health.missouri.edu

[†]These authors contributed equally to this work.

Keywords

Skeletal muscle, biopsy injury, angiogenesis, myogenesis

Abstract

Background: Acute injury to skeletal muscle damages myofibers and fragments

capillaries, impairing contractile function and local perfusion. Myofibers and microvessels

regenerate from satellite cells and from surviving microvessel fragments, respectively, to

restore intact muscle. However, it is unknown whether myofiber regeneration and

microvascular regeneration reflect interdependent processes or may proceed

sequentially.

Methods: To investigate the temporal relationship between myogenesis and

angiogenesis during regeneration, a punch biopsy (diameter, 2 mm) was performed

through the center of the gluteus maximus (GM) muscle. Complete removal of all tissue

components created a void into which regeneration was evaluated through 21 days post

injury (dpi). Confocal imaging and histological analyses of whole-mount GM preparations

and GM cross sections assessed the growth of microvessels and myofibers into the

wound. Regeneration of perfused microvessels was evaluated *in vivo* by injecting

fluorescent dextran into the circulation during intravital imaging.

Results: A provisional matrix filled with PDGFRα$^+$ and CD45$^+$ cells spanned the wound

within 1 dpi. Regenerating microvessels advanced into the matrix by 7 dpi. At 10 dpi,

sprouting and intussusceptive angiogenesis produced disorganized microvascular

networks and spanned the wound with perfusion by 14 dpi. In striking contrast, the wound

remained devoid of myofibers at 7 and 10 dpi. Myogenesis into the wound began by 14

dpi with nascent myofibers traversing the wound by 21 dpi. Regenerating myofibers and

microvessels were less well organized than in the surrounding (uninjured) muscle.

Conclusions: Angiogenesis precedes myogenesis following punch biopsy injury of adult

skeletal muscle. Regenerating microvessels encompass the wound and become

perfused with blood prior to colocalization with regenerating myofibers. These findings

infer that a microvascular supply supports the metabolic demands of regenerating skeletal

muscle. Finding that regenerated microvascular networks and myofibers are disorganized

within the biopsy site suggests that loss of guidance cues upon complete tissue removal

impairs re-establishment of canonical skeletal muscle structure.

Background

Acute injury to skeletal muscle damages myofibers, triggers an inflammatory

response, and fragments capillary networks (1-3). Although the events and time course

of the inflammatory cascade (3) and the regeneration of myofibers from resident stem

cells [satellite cells (SCs)] following injury are well defined (1, 4-6), data regarding the

recovery of the microcirculation following injury are limited. Following exposure to

myotoxins, freezing, or physical trauma, capillary blood flow is abolished within 1 day post

injury (dpi) but begins to recover by 5 dpi as capillary networks reform (1, 7). Although

initially disorganized, newly-formed networks remodel coincident with the maturation of

regenerated myofibers by 21 dpi (2, 7, 8).

In accord with their spatial proximity, crosstalk between SCs and capillary

endothelial cells (ECs) is integral to the regeneration of intact skeletal muscle and SC

self-renewal (9, 10). Moreover, as the immune response transitions from inflammation

and degeneration to a milieu that promotes healing (3, 11), SC proliferation is stimulated

(12) along with gene co-regulation between angiogenesis and myogenesis (13).

However, it is unknown whether microvascular regeneration and myofiber regeneration

reflect interdependent processes or may proceed sequentially following injury.

To test whether a temporal relationship exists between myogenesis and

angiogenesis during regeneration in adult skeletal muscle, we developed a model of

complete tissue removal using punch biopsy of the mouse gluteus maximus (GM) muscle.

We hypothesized that following removal of all tissue components, angiogenesis precedes

the formation of new myofibers to support the metabolic demands of regenerating skeletal

muscle. We report here that creating a void into which regeneration proceeds reveals that

angiogenesis occurs prior to myogenesis after biopsy injury. These findings establish a

foundation for detailed studies of distinct versus shared regulation between regeneration

of myofibers and their vascular supply.

Methods

Animal care and use

Male and female mice (C57BL/6, Jackson Laboratory; Bar Harbor, ME) were bred

and housed in animal care facilities of the University of Missouri. Mice (weight, ~30 g)

were studied the age of ~4 months. In reporter mice bred on a C57BL/6 background

[Cdh5-Cre^{ERT2} (14) x ROSA26^{mTmG} (#007676, Jackson Laboratory)], Cre recombination

for expression of membrane bound green fluorescent protein (GFP) in ECs was induced

through intraperitoneal injection of 100 uL tamoxifen (10 mg/mL + 5% ethanol in corn oil;

#T5648, Sigma-Aldrich; St. Louis, MO) on 3 consecutive days; at least 1 week elapsed

after the first injection prior to study. All mice were maintained under a 12:12 h light/dark

cycle at 22-24 ⁰C with fresh food and water *ad libitum*. To avoid any order effect, collection

of data at criterion timepoints was randomized. Prior to muscle injury, intravital

microscopy, or tissue collection, a mouse was anesthetized [ketamine (100 mg/kg) +

xylazine (10 mg/kg) in sterile saline; intraperitoneal injection]. Mice were euthanized at

the end of an experiment by anesthetic overdose and cervical dislocation.

Punch biopsy

A mouse was anesthetized and positioned on an aluminum warming plate to

maintain body temperature at 37 °C. As needed, supplemental injections of anesthesia

(~20% of initial) were given to maintain a stable plane of anesthesia confirmed by lack of

withdrawal to a toe pinch (monitored every 15 min). Skin covering the left GM was shaved

and sterilized by swabbing 3x with Betadine and 70% alcohol. While viewing through a

stereomicroscope, the mouse was positioned on its abdomen and a ~1 cm incision was

made through the skin overlying the GM near the spine. The skin was retracted to expose

the GM and irrigated with sterile saline. A hole was made through the center of the GM

with a sterile 2 mm diameter biopsy punch (MediChoice #DP0200, Owens & Minor;

Mechanicsville, VA); this size is below the limits of muscle loss for effective regeneration

(15) while large enough to create a void into which regeneration could be reliably studied.

Anatomical landmarks provided a reference for consistency in the site of biopsy. The skin

was closed with sterile 6-0 nylon sutures (UNIFY #S-N618R13, AD Surgical; Sunnyvale,

CA). This entire procedure required ~20 min. For recovery, the mouse was placed on a

heated platform, monitored until consciousness and ambulation were restored (~2-3 h),

then returned to its original cage. Normal activity and behavior were routinely observed

within 24 h. Regeneration of tissue components into the wound was evaluated at key

timepoints through 21 days post injury (dpi) with uninjured mice (0 dpi) as controls.

To evaluate cellular damage surrounding the biopsy site (Fig S1), Evans Blue dye

(EBD) (1% solution in sterile saline; #E2129, Sigma) was injected intraperitoneally (10

µL/g body mass) following the surgical procedure. At 1 dpi, the GM was dissected (as

below) to image EBD. Images were acquired with a 4x objective on an E800 microscope

coupled to DS-Fi3 camera using Elements software (all from Nikon; Tokyo, Japan).

Dissection of the gluteus maximus muscle

A mouse was anesthetized, the surgical area was shaved, the mouse was placed

on a warming plate, and the skin overlying the GM was removed with scissors. Exposed

tissue was continuously superfused with a bicarbonate-buffered physiological salt

solution (bbPSS; pH 7.4, 34-35°C) containing (in mM) 131.9 $NaCl_2$ (Fisher Scientific;

Pittsburg, PA), 4.7 KCl (Fisher), 2 $CaCl_2$ (Sigma), 1.17 $MgSO_4$ (Sigma), and 18 $NaHCO_3$

(Sigma) equilibrated with 5% CO_2/95% N_2. While viewing through a stereomicroscope,

excess connective tissue was removed using microdissection while avoiding the site of

injury. The GM was cut from its origin along the lumbar fascia, sacrum, and iliac crest

then reflected away from the body to view its vascular supply from the ventral surface (7,

16).

Intravital microscopy

The exteriorized GM was spread onto a transparent rubber pedestal (Sylgard 184;

Dow Corning; Midland, MI) and pinned at its edges to approximate *in situ* dimensions.

Exposed tissue on the body of the mouse was covered with plastic film (Glad Press n'

Seal) to prevent dehydration, the preparation was transferred to the stage of a Nikon

E600FN microscope, and the GM equilibrated for 30 min while continuously superfused

with bbPSS at 3 mL/min. Supplemental doses of anesthetic were given throughout the

experimental protocol (duration, 2-3 h) to maintain a stable plane of anesthesia (as

above). To assess vascular perfusion, fluorescein isothiocyanate (FITC) conjugated

dextran (70 kDa; 10 mg/mL sterile saline) was injected (200 µL) into the systemic

circulation via the retroorbital sinus and allowed to circulate for ~10 min. The GM was

illuminated with a mercury lamp for fluorescence imaging using a filter cube appropriate

for circulating FITC or transgenic eGFP. Images were acquired through Nikon Plan Fluor

4x/0.13 or Plan Fluor 10x/0.3 objectives coupled to a low light CMOS FP-Lucy camera

[Stanford Photonics, Inc.; Palo Alto, CA (SPI)] and displayed on a digital monitor. Time

lapse images were recorded at 40 frames/s using Piper Control software (SPI).

Confocal imaging of fresh whole mount muscle preparations

The GM was removed from a Cdh5-mTmG mouse and placed in a custom imaging

chamber with the ventral surface facing the objective. To optimize resolution of the

microvasculature, a drop of PBS (~10 µL) was added to the chamber and the GM was

flattened by placing a glass block (2 cm x 2.5 cm x 1 cm; mass, 7.8 g) on the dorsal

surface. Images were acquired with a HC PL APO 10/0.40 CS2 objective on an inverted

laser scanning confocal microscope (TCS SP8) using LASX software (all from , Leica

Microsystems; Buffalo Grove, IL). A Dragonfly High Speed Confocal Microscope System

[Leica DMi8 microscope coupled to an Andor Dragonfly 200 and integrated Fusion

software (Oxford Instruments; Abingdon, United Kingdom) was also used to acquire

images. To image the entire wound, tile scans (3 x 3 grid encompassing ~3.5 mm x 3.5

mm) were acquired with a Leica HC PL APO 10x/0.45 objective and stitched using

integrated software. Using maximum projection z-stacks acquired with a Leica HC PL

IRAPO 20x/0.75 objective, 10 random microvessels were chosen within a region of

interest (ROI; 493 x 493 μm). Diameters of regenerating microvessels were measured at

the midpoint between two branch points and averaged for a given GM; 5-7 GM were

analyzed at respective dpi.

Immunostaining

An excised GM was immobilized by pinning the edges in the well of a 12-well plate

coated with Sylgard 184. After washing with PBS, a region of muscle (~5 mm x 5 mm)

containing the injury surrounded by undamaged tissue was trimmed and prepared for

either whole mount immunostaining or frozen in Tissue Tek OCT compound (VWR

International LLC; Radnor, PA) to obtain tissue cross sections for histology.

Whole mount preparations were fixed in 2% paraformaldehyde for 30 min, washed

in PBS, and placed in blocking buffer (2% bovine serum albumin, 4% normal donkey

serum, 0.5% triton X-100 in PBS) for 30 min. Preparations were then incubated overnight

at 4 °C with validated primary antibodies: anti-rat CD31 [platelet endothelial cell adhesion

molecule, [PECAM-1; 1:400, #550274, BD Pharmingen; San Diego, CA; (17)] to identify

endothelial cells; anti-rabbit CD45 [leukocyte common antigen; 1:200, #ab10558, Abcam;

Cambridge, UK; (18)] to identify inflammatory cells; and anti-goat PDGFRα [1:200,

#AF1062, R&D Systems; Minneapolis, MN; (19)] to identify fibroadipogenic precursor

cells (FAPs) (20-22). Preparations were washed in blocking buffer, incubated with

secondary antibodies for 30 min, washed again, then rinsed in PBS before transfer to an

imaging chamber for confocal image acquisition (as above).

For tissue cross sections, trimmed and washed GM samples were transferred to a

cryomold containing OCT compound with black silk suture (2-0 thickness; length, 2 mm)

positioned adjacent to the injury for spatial reference. The cryomold was frozen in

isopentane cooled in liquid nitrogen and stored at -80 °C until sectioning. Cross sections

of frozen GM were cut (thickness, 10 µm) with a cryostat (HM 550 Cryostat, Thermo

Scientific; Waltham, MA) at the center of the injury referencing to the silk suture.

For fluorescence imaging, sections were fixed in 4% paraformaldehyde for 10 min

and then stained with CD31 (1:500, BD Pharmingen), myosin heavy chain [1:5, MF-20,

Developmental Studies Hybridoma Bank, The University of Iowa Department of Biology;

Iowa City, IA; (23)], laminin [(1:200, #PA1-16730, Invitrogen; Waltham, MA; (24)], and

appropriate secondary antibodies (1:400, AlexaFluor, Fisher). Prolong Gold containing

DAPI (Fisher) was added before slides were coverslipped. Sections were imaged using

appropriate filters on a Nikon E800 microscope with a DS-Qi2 camera and Nikon

Elements Software.

The distance devoid of myofibers (e.g., diameter of remaining visible wound) was

measured along GM cross sections accounting for curvatures within the specimen (Fig.

4). Within this region, the total area of CD31[+] staining was measured to evaluate

microvascular density within the wound preceding myofiber ingrowth.

Histochemistry

To visualize collagen deposited within the provisional matrix in whole mount

preparations, the GM was removed, permeabilized in 0.5% Triton X-100 in PBS, and

incubated for 1 h with PicroSirius Red (1% Direct Red in saturated picric acid). Thereafter,

GM were treated for 30 min with 0.5% acetic acid in ddH$_2$O and rinsed in 100% EtOH.

Images were acquired as described for EBD.

Statistics

Summary data are displayed for individual GM preparations along with means ±

s.e.m. Statistical analyses were performed using Prism 9 software (GraphPad Software,

Inc.; La Jolla, CA). One-way ANOVA with Tukey's post hoc tests were used to determine

statistical significance between time points with $P < 0.05$ considered significant. Values

for 'n' refer to the number of GM analyzed; one GM was studied per mouse.

Results

Punch biopsy creates a void into which regeneration advances

Muscle injury was created using a biopsy punch (diameter, 2 mm) to remove a

volume of muscle below the critical threshold for regeneration [Fig. 1A; (15)]. A circular

hole through the center of the GM resulted in minimal collateral damage to the

surrounding tissue as evidenced by nominal uptake of EBD into the ends of myofibers

bordering the wound at 1 dpi (Supp. Fig. 1). Observed in anesthetized mice, intravascular

injection of FITC dextran confirmed the perfusion void within the wound while blood flow

was preserved to tissue surrounding the injury (Fig. 1B). Sirius red staining of whole

mount GM preparations revealed the deposition of collagen to create a provisional matrix

(25) that nearly spanned the wound at 1 dpi (Fig. 1C); immunostaining demonstrated

invasion of this matrix by PDGFRα⁺ cells (fibroadipogenic progenitor cells, FAPs)

accompanied by CD45⁺ (inflammatory) cells (Fig. 1D).

Microvascular growth into the wound

Using an EC-specific Cre driver [Cdh5-CreERT2; (14)] and tamoxifen-induced

recombination of the Rosa26 mTmG locus to genetically label the endothelium with

membrane-bound eGFP (Cdh5-mTmG mice), we evaluated microvascular regeneration

following punch biopsy. Angiogenic sprouts reflecting endothelial tip cells (26) were

apparent at 5 dpi (not shown) and extended into the wound at 7 dpi (Fig. 2A&B). Nascent

microvessels also underwent internal division (i.e., intussusceptive angiogenesis) as

networks expanded to extend across the wound (Fig. 2C-E). Through 14 dpi,

microvessels split into daughter vessels generating numerous branches with short

interconnections. The angiogenic response from 7-14 dpi progressed as a centripetal

gradient of microvessel growth from the edges towards the center of the wound.

By 14 dpi, microvascular networks spanning the wound were perfused with blood

but lacked hierarchical structure or definitive patterns of blood flow (Fig. 3, Supp. Fig. 2).

Instead, nascent microvessels were clustered into zones of robust angiogenic activity with

random orientation (Fig. 2D). In addition, microvessel diameter significantly increased

from 4.1 ± 0.3 µm in uninjured muscle to 7.4 ± 0.4 µm at 7 dpi and 9.0 ± 0.6 µm at 14 dpi

(Fig. 3D). Intravital microscopy revealed that only the microvessels having the largest

diameter were continuously perfused; most segments contained stationary red blood cells

or plasma alone at 14 dpi (Supp. Fig. 2). In perfused microvessels, the direction of blood

flow oscillated and was interspersed with periods of flow cessation.

By 21 dpi, extensive microvascular networks encompassed the site of injury (Fig.

3C) with diameters (6.6 ± 0.8 µm) returning towards control. Nevertheless, the parallel

arrangement of microvessels observed in the surrounding uninjured muscle was not

restored. Instead, numerous anastomoses created circuitous pathways of local perfusion.

Regeneration of myofibers into the wound

In striking contrast to the time course of revascularization, the wound remained

devoid of myofibers at 7 dpi, a time at which myofiber regeneration would be well

advanced in other models of skeletal muscle injury (1, 7). As myofibers began to penetrate

the provisional matrix and regenerate into the wound at 10-14 dpi (Fig. 3), they were

densely packed, interwoven, and coursed along the wound edge before integrating with

undamaged myofibers. Additionally, regenerating myofibers were smaller in width (i.e.,

cross sectional area) than in surrounding (uninjured) tissue, consistent with other muscle

injury models (1, 5, 7, 27). Newly-formed myofibers intertwined as they spanned the

wound at 21 dpi. In some preparations, clusters of adipocytes occupied space where

myofibers had not regrown (Fig. 3C, Supp. Fig. 3). In turn, regenerating capillary networks

remodeled in accord with adipocyte morphology (Fig. 3C).

To confirm that revascularization occurred prior to myofiber regeneration, GM

cross sections (thickness, 10 µm) were immunolabeled for CD31 to identify ECs, myosin

heavy chain (MyHC) for contractile myofibers, laminin to detect basal laminae, and DAPI

for nuclei. Congruent with data from intravital and confocal imaging, healing progressed

into the wound from 7 dpi through 21 dpi with regeneration of microvessels preceding that

of myofibers (Fig. 4A-C). The length of the injured region devoid of myofibers

(corresponding to the diameter of existing wound) was used as an index of healing over

time. This length was 1512 ± 153 µm at 7 dpi which progressively decreased to 475 ±

134 µm at 14 dpi and 71 ± 71 at 21 dpi (Fig. 4D). We quantified the area occupied by ECs

within the region lacking myofibers as an indicator of microvascular density that preceded

myogenesis. This EC area was 2794 ± 1126 µm^2 at 7 dpi, 3366 ± 930 µm^2 at 14 dpi, and

decreased to 700 ± 700 µm^2 at 21 dpi (Fig. 4E), reflecting the growth of myofibers into

the wound, colocalizing with ECs (microvessels) as regeneration advanced. In contrast

to centrally-located nuclei identifying regenerated myofibers (1, 4, 5), adjacent fibers

within the injury often had nuclei localized to their periphery (Fig. 4F), suggesting that the

removal of stroma affects localization of myonuclei during myogenesis post injury.

Discussion

Punch biopsy is an acute injury that is shown here to dissociate the time course of

angiogenesis from that of myogenesis in regenerating skeletal muscle for the first time.

In contrast to myotoxins, freezing, or physical trauma, where efferocytosis removes

cellular debris while leaving a residual extracellular matrix [ECM (28-31)], punch biopsy

removes all tissue components to create an empty void. In the GM of adult male and

female mice, creating such a void through the center of the muscle resulted in the

formation of a provisional matrix (25) filled with immune cells and FAPs at 1 dpi.

Endothelial cells proliferate and migrate from the edges of the wound into the provisional

matrix by 7 dpi; a combination of sprouting and intussusceptive angiogenesis resulted in

the development of microvascular networks that spanned the wound and became

perfused with blood by 14 dpi. Myofibers grew into the wound by 14 dpi and filled the

wound at 21 dpi, approximately 1 week after comparable events in the regenerating

microvasculature. Microvessels and myofibers within the wound are disorganized

compared to uninjured muscle suggesting that guidance cues for regeneration may be

lost upon complete tissue removal that would otherwise promote restoration of intact

muscle.

Microvessel and myofiber regeneration following injury

The GM is a thin muscle comprised of mixed fiber type (32) rendering it well-suited

for studying microvascular networks in the mouse (16). Previous studies evaluating the

GM microcirculation following injury with $BaCl_2$ show that the microcirculation regenerates

and recovers function concomitant with myofibers through 21-35 dpi (2, 7, 8). These

findings in the GM are consistent with injuries to other rodent hindlimb muscles (1, 4, 5,

13, 27). Myofibers and microvessels regenerate *in situ* from SCs and from surviving

capillary fragments (i.e., ECs), respectively, to restore intact muscle (1, 2, 4, 7). Myofiber

regeneration ensues as the milieu transitions from proinflammatory to one that promotes

healing (3, 33). Earlier studies reported that capillarization is governed by the size and

growth of regenerating myofibers (34), with an upregulation of angiogenic gene

signatures during the early stage of muscle regeneration (35). Supporting this view were

findings that ECs enhanced SC growth through secretion of an array of growth factors

[VEGF, IGF-1, HGF, bFGF, PDGF-BB (9)]. Complementary studies found that SCs

modulate angiogenesis through hypoxia-inducible factor-1 (HIF-1), a transcriptional

regulator of VEGF gene expression (36). With SCs remaining in juxtavascular locations,

differentiating myogenic cells were both proangiogenic and spatiotemporally associated

with angiogenesis (9).

Sensing attractive and repulsive signals in the microenvironment, ECs invade the

provisional matrix and began to revascularize the wound at 7 dpi (Fig. 2). With sprouting

angiogenesis, endothelial tip cells guide the formation of new capillaries while stalk cells

proliferate behind the tip cells to extend the newly forming microvessels (26).

Intussusceptive angiogenesis further contributes to microvascular proliferation within the

wound (Fig. 2) by forming pillars within the lumen of a vessel that extend to split a

preexisting vessel into two daughter segments and modify branch angles at bifurcations

(37). During regeneration following hindlimb ischemia in mice, intussusception was the

dominant mechanism of angiogenesis (38). The findings presented herein illustrate that

both sprouting and intussusceptive angiogenesis contribute to microvascular

regeneration following punch biopsy injury. The formation of new microvessels through

vasculogenesis (39) may also contribute to revascularization but was not resolved in the

present experiments.

Modulation of SC-EC crosstalk by immune cells and FAPs

Crosstalk between surviving SCs and ECs occurs under the influence of immune

cells and FAPs (3, 6, 9-13, 20, 40, 41). FAPs are identified by the expression of

mesenchymal stem cell surface proteins (e.g., Sca-1, CD34, PDGFRα) and reside in the

interstitial space adjacent to the microvasculature (21, 22, 42). Upon injury, FAPs enter

the cell cycle to proliferate, localize to the site of tissue damage, differentiate into

fibroblasts and adipocytes [but not into SCs or myofibers (20, 22, 40)], and work in concert

with macrophages and ECs to promote myofiber regeneration (6, 22, 41, 43). Our finding

that CD45[+] and PDGFRα[+] cells invade the provisional matrix within 1 dpi following punch

biopsy (Fig. 1) is consistent with earlier studies (13, 20, 22, 43). Activated FAPs deposit

ECM proteins (e.g., collagen), secrete myogenic differentiation factors, and recruit CD45[+]

immune cells via paracrine signaling (20, 43). The ensuing cascade of granulocytes,

monocytes, and macrophages produce cytokines and chemokines aimed at clearing

cellular debris, promoting self-renewal of SCs, and preventing premature myogenic cell

differentiation (3, 44).

The ECM aids in cell adhesion, cell-to-cell communication, and differentiation (28,

45). In addition to providing matrix structure, collagen fibers guide the growth of capillaries

during regeneration (46). In such manner, the microenvironment of the provisional matrix

established by FAPs and immune cells serves as the foundation for the regeneration of

intact skeletal muscle (25). However, if the injury exceeds a critical threshold of muscle

loss [e.g., a 3-mm diameter biopsy in the mouse quadriceps (15)], inflammation and

fibrosis persist. Furthermore, if myoblast migration and fusion are delayed during

regeneration, FAPs may also contribute to myosteatosis (42).

Novel features of punch biopsy injury

Previous studies of muscle injury have not resolved whether regeneration of

myofibers and microvessels following injury are interdependent processes or whether

angiogenesis and myogenesis proceed as sequential events. Indeed, earlier findings

have led to the conclusion that angiogenesis and matrix remodeling occur during the last

phase of myogenesis and growth of new myofibers (5, 33). However, data presented

herein show that matrix generation and angiogenesis occur first. Such ambiguity may be

attributed to the persistence of basal laminae following efferocytosis, an outcome that

prevails in established models (1, 4, 5, 11, 27). Preexisting basal laminae can provide

guidance during regeneration of microvessels and myofibers (28-31, 46, 47), but a biopsy

punch removes the entire tissue. A key finding here is that, in contrast to the concomitant

regeneration of microvessels and myofibers discussed above, the onset of microvascular

growth into the wound and perfusion across the wound precedes that of myofibers by

approximately 1 week. We attribute this difference from other injury models to the removal

of the entire tissue, with loss of primed progenitor cells and guidance cues that would

otherwise be retained in the residual stroma.

In addition to distinguishing between the time course of angiogenesis versus

myogenesis, key differences emerged during regeneration following punch biopsy when

compared to other models of muscle injury. In contrast to the onset of sprouting

angiogenesis 2-3 dpi following exposure to myotoxins (1, 2, 7), EC sprouts were not

apparent until ~1 week following punch biopsy (Fig. 2A). This longer delay may be

attributable to prioritized healing around the edges of the open wound prior to EC invasion

of the provisional matrix. A delay in angiogenesis (14 dpi) was also observed following

freezing (1)], which kills all cells within the injured region.

Nascent microvessels were nonuniformly enlarged through 21 dpi compared to

capillaries supplying healthy myofibers in the surrounding tissue. This finding is

inconsistent with data from the GM following $BaCl_2$ injury, where vessel diameter returned

to baseline values and the structure of capillary networks had nearly recovered at this

time (8). The irregular blood flow patterns observed within the wound (Supp. Fig. 2) can

be attributed to the absence of hierarchical organization of newly-formed networks or

established arterio-venous pressure gradients (48), reflecting the absence of structural

and functional cues that may otherwise be provided by the residual ECM.

Whereas centrally located myonuclei are a hallmark of regenerated myofibers (1,

4, 5), including the GM following $BaCl_2$ injury (7), regenerating myofibers filling the void

contained myonuclei positioned centrally in some myofibers while positioned at the

periphery of other myofibers, often located adjacent to each other (Fig. 4F). This

difference in nuclear localization implies a modified mechanism of myogenesis when

compared to other models of skeletal muscle injury and myofiber regeneration (49), which

may reflect a distinct microenvironment for regeneration after biopsy injury. That

regenerating myofibers intertwine instead of exhibiting their characteristic parallel

alignment further indicates that the provisional matrix lacks the structural basis to dictate

the canonical arrangement of myofibers (29, 30).

Summary and Conclusions

The present study demonstrates that creating a hole through the GM muscle

enabled definitive temporal resolution of angiogenesis versus myogenesis for the first

time following injury in adult skeletal muscle. Following punch biopsy injury to remove all

tissue components, a provisional matrix invested with inflammatory cells and fibroblasts

is deposited within 1 dpi. Sprouting and intussusceptive angiogenesis revascularize the

wound at 7-14 dpi, a time frame that precedes the onset and completion of myofiber

regeneration by ~ 1 week. Consistent with other models of injury, myofibers invested with

a microvascular supply are restored by 21 dpi. However, regenerated capillary networks

and myofibers within the site of injury remain disorganized relative to uninjured muscle,

suggesting that the total loss of skeletal muscle ultrastructure disrupts the morphology of

regenerating tissue components.

These experiments are the first to illustrate that revascularization precedes the

replacement of myofibers in adult skeletal muscle following injury. Our findings are

consistent with the interpretation that establishing a vascular supply in the provisional

matrix generates conditions that support the regeneration of myofibers (6). The outcome

of this study contrasts with injuries in which the basal laminae (and progenitor cells)

persist and angiogenesis occurs concomitant with myogenesis (9, 13, 47). The present

findings thereby provide a foundation for selectively manipulating variables known to

affect respective components of regeneration to restore the functional integrity of skeletal

muscle.

List of abbreviations

Cdh5-mTmG: Cdh5-Cre[ERT2] x Rosa[mTmG] transgenic mice; dpi: days post injury; EBD:

Evans Blue dye; ECs: endothelial cells; ECM: extracellular matrix; GFP: green fluorescent

protein; FAPs: fibroadipogenic precursor cells; FITC: fluorescein isothiocyanate; GM:

gluteus maximus; PDGFRα: platelet derived growth factor receptor alpha; MyHC: myosin

heavy chain; SCs: satellite cells.

Declarations

Ethical approval

All procedures were approved by the Animal Care and Use Committee at the University of Missouri (protocol #17720). Experiments were performed in accordance with the National Research Council's Guide for the Care and Use of Laboratory Animals.

Consent for publication

Not applicable.

Availability of data and materials

The data generated and analyzed during the current study are available from the corresponding author on reasonable request.

Competing interests

The authors declare that they have no competing interests.

Funding

The funding sources that supported the work completed in the current study had no role in 1) the design of the study, 2) collection, analysis, and interpretation of data, or 3) in writing the manuscript. This study was supported by National Institutes of Health grant F32 HL-152558 (NLJ), American Physiological Society Postdoctoral Fellowship (ABM), National Institutes of Health Loan Repayment award (ABM), Margaret Proctor Mulligan Fellowship, University of Missouri School of Medicine (NLJ), Margaret Proctor

Mulligan Professorship, University of Missouri School of Medicine (SSS), School of

Medicine Development Award, University of Missouri School of Medicine (SSS), National

Institutes of Health MERIT Award R37 HL-041026 (SSS).

Acknowledgments

Yuki Yang contributed to preliminary experiments. Drs. Erika Boerman and Scott

Zawieja provided use of their confocal microscopes and imaging processing software.

Drs. DDW Cornelison and Charles Norton contributed thoughtful discussion throughout

data collection and analysis.

Author contributions

Conceptualization: NLJ, ABM, SSS; Methodology: NLJ, ABM, SSS; Investigation:

NLJ, ABM, SSS; Visualization: NLJ, ABM; Funding acquisition: NLJ, ABM, SSS; Project

administration: SSS; Supervision: SSS; Writing – original draft: NLJ; Writing – review &

editing: NLJ, ABM, SSS

References

1. Hardy D, Besnard A, Latil M, Jouvion G, Briand D, Thepenier C, et al. Comparative Study of Injury Models for Studying Muscle Regeneration in Mice. PLoS One. 2016;11(1):e0147198.

2. Morton AB, Norton CE, Jacobsen NL, Fernando CA, Cornelison DDW, Segal SS. Barium chloride injures myofibers through calcium-induced proteolysis with fragmentation of motor nerves and microvessels. Skelet Muscle. 2019;9(1):27.

3. Tidball JG. Regulation of muscle growth and regeneration by the immune system. Nat Rev Immunol. 2017;17(3):165-78.

4. Yin H, Price F, Rudnicki MA. Satellite cells and the muscle stem cell niche. Physiol Rev. 2013;93(1):23-67.

5. Baghdadi MB, Tajbakhsh S. Regulation and phylogeny of skeletal muscle regeneration. Developmental Biology. 2018;433(2):200-9.

6. Anderson JE. Key concepts in muscle regeneration: muscle "cellular ecology" integrates a gestalt of cellular cross-talk, motility, and activity to remodel structure and restore function. Eur J Appl Physiol. 2022;122(2):273-300.

7. Fernando CA, Pangan AM, Cornelison D, Segal SS. Recovery of blood flow regulation in microvascular resistance networks during regeneration of mouse gluteus maximus muscle. J Physiol. 2019;597(5):1401-17.

8. Jacobsen NL, Norton CE, Shaw RL, Cornelison DDW, Segal SS. Myofibre injury induces capillary disruption and regeneration of disorganized microvascular networks. J Physiol. 2022;600(1):41-60.

9.   Christov C, Chretien F, Abou-Khalil R, Bassez G, Vallet G, Authier FJ, et al. Muscle

satellite cells and endothelial cells: close neighbors and privileged partners. Mol Biol

Cell. 2007;18(4):1397-409.

10. Verma M, Asakura Y, Murakonda BSR, Pengo T, Latroche C, Chazaud B, et al.

Muscle Satellite Cell Cross-Talk with a Vascular Niche Maintains Quiescence via

VEGF and Notch Signaling. Cell Stem Cell. 2018;23(4):530-43 e9.

11. Juban G, Chazaud B. Efferocytosis during Skeletal Muscle Regeneration. Cells.

2021;10(12).

12. Ratnayake D, Nguyen PD, Rossello FJ, Wimmer VC, Tan JL, Galvis LA, et al.

Macrophages provide a transient muscle stem cell niche via NAMPT secretion.

Nature. 2021;591(7849):281-7.

13. Latroche C, Weiss-Gayet M, Muller L, Gitiaux C, Leblanc P, Liot S, et al. Coupling

between Myogenesis and Angiogenesis during Skeletal Muscle Regeneration Is

Stimulated by Restorative Macrophages. Stem Cell Reports. 2017;9(6):2018-33.

14. Wang Y, Nakayama M, Pitulescu ME, Schmidt TS, Bochenek ML, Sakakibara A, et

al. Ephrin-B2 controls VEGF-induced angiogenesis and lymphangiogenesis. Nature.

2010;465(7297):483-6.

15. Anderson SE, Han WM, Srinivasa V, Mohiuddin M, Ruehle MA, Moon JY, et al.

Determination of a Critical Size Threshold for Volumetric Muscle Loss in the Mouse

Quadriceps. Tissue Eng Part C Methods. 2019;25(2):59-70.

16. Bearden SE, Payne GW, Chisty A, Segal SS. Arteriolar network architecture and

vasomotor function with ageing in mouse gluteus maximus muscle. J Physiol.

2004;561(Pt 2):535-45.

17. Baldwin HS, Shen HM, Yan HC, DeLisser HM, Chung A, Mickanin C, et al. Platelet
endothelial cell adhesion molecule-1 (PECAM-1/CD31): alternatively spliced,
functionally distinct isoforms expressed during mammalian cardiovascular
development. Development. 1994;120(9):2539-53.

18. Biscetti F, Straface G, De Cristofaro R, Lancellotti S, Rizzo P, Arena V, et al. High-
mobility group box-1 protein promotes angiogenesis after peripheral ischemia in
diabetic mice through a VEGF-dependent mechanism. Diabetes. 2010;59(6):1496-
505.

19. Gilbertson DG, Duff ME, West JW, Kelly JD, Sheppard PO, Hofstrand PD, et al.
Platelet-derived growth factor C (PDGF-C), a novel growth factor that binds to PDGF
alpha and beta receptor. J Biol Chem. 2001;276(29):27406-14.

20. Biferali B, Proietti D, Mozzetta C, Madaro L. Fibro–Adipogenic Progenitors Cross-
Talk in Skeletal Muscle: The Social Network. Frontiers in Physiology. 2019;10(1074).

21. Sono T, Hsu CY, Wang Y, Xu J, Cherief M, Marini S, et al. Perivascular Fibro-
Adipogenic Progenitor Tracing during Post-Traumatic Osteoarthritis. Am J Pathol.
2020;190(9):1909-20.

22. Joe AW, Yi L, Natarajan A, Le Grand F, So L, Wang J, et al. Muscle injury activates
resident fibro/adipogenic progenitors that facilitate myogenesis. Nat Cell Biol.
2010;12(2):153-63.

23. Bader D, Masaki T, Fischman DA. Immunochemical analysis of myosin heavy chain
during avian myogenesis in vivo and in vitro. J Cell Biol. 1982;95(3):763-70.

24. Bal NC, Singh S, Reis FCG, Maurya SK, Pani S, Rowland LA, et al. Both brown

adipose tissue and skeletal muscle thermogenesis processes are activated during

mild to severe cold adaptation in mice. J Biol Chem. 2017;292(40):16616-25.

25. Barker TH, Engler AJ. The provisional matrix: setting the stage for tissue repair

outcomes. Matrix Biol. 2017;60-61:1-4.

26. Blanco R, Gerhardt H. VEGF and Notch in tip and stalk cell selection. Cold Spring

Harb Perspect Med. 2013;3(1):a006569.

27. Carlson BM, Faulkner JA. The regeneration of skeletal muscle fibers following injury:

a review. Med Sci Sports Exerc. 1983;15(3):187-98.

28. Grounds MD. Complexity of Extracellular Matrix and Skeletal Muscle Regeneration.

In: Schiaffino S, Partridge T, editors. Skeletal Muscle Repair and Regeneration.

Dordrect, The Netherlands: Springer; 2008. p. 269-301.

29. Sanes JR. The basement membrane/basal lamina of skeletal muscle. J Biol Chem.

2003;278(15):12601-4.

30. Vracko R, Benditt EP. Basal lamina: the scaffold for orderly cell replacement.

Observations on regeneration of injured skeletal muscle fibers and capillaries. J Cell

Biol. 1972;55(2):406-19.

31. Webster MT, Manor U, Lippincott-Schwartz J, Fan CM. Intravital Imaging Reveals

Ghost Fibers as Architectural Units Guiding Myogenic Progenitors during

Regeneration. Cell Stem Cell. 2016;18(2):243-52.

32. Lampa SJ, Potluri S, Norton AS, Laskowski MB. A morphological technique for

exploring neuromuscular topography expressed in the mouse gluteus maximus

muscle. J Neurosci Methods. 2004;138(1-2):51-6.

33. Al-Zaeed N, Budai Z, Szondy Z, Sarang Z. TAM kinase signaling is indispensable for proper skeletal muscle regeneration in mice. Cell Death Dis. 2021;12(6):611.

34. Luque E, Pena J, Alonso PJ, Jimena I. Microvascular pattern during the growth of regenerating muscle fibers in the rat. Ann Anat. 1992;174(3):245-9.

35. Wagatsuma A. Endogenous expression of angiogenesis-related factors in response to muscle injury. Molecular and Cellular Biochemistry. 2007;298(1):151-9.

36. Rhoads RP, Johnson RM, Rathbone CR, Liu X, Temm-Grove C, Sheehan SM, et al. Satellite cell-mediated angiogenesis in vitro coincides with a functional hypoxia-inducible factor pathway. Am J Physiol Cell Physiol. 2009;296(6):C1321-8.

37. Mentzer SJ, Konerding MA. Intussusceptive angiogenesis: expansion and remodeling of microvascular networks. Angiogenesis. 2014;17(3):499-509.

38. Arpino JM, Yin H, Prescott EK, Staples SCR, Nong Z, Li F, et al. Low-flow intussusception and metastable VEGFR2 signaling launch angiogenesis in ischemic muscle. Sci Adv. 2021;7(48):eabg9509.

39. Risau W, Flamme I. Vasculogenesis. Annu Rev Cell Dev Biol. 1995;11:73-91.

40. Uezumi A, Ito T, Morikawa D, Shimizu N, Yoneda T, Segawa M, et al. Fibrosis and adipogenesis originate from a common mesenchymal progenitor in skeletal muscle. J Cell Sci. 2011;124(Pt 21):3654-64.

41. Panci G, Chazaud B. Inflammation during post-injury skeletal muscle regeneration. Seminars in Cell & Developmental Biology. 2021;119:32-8.

42. Hamrick MW, McGee-Lawrence ME, Frechette DM. Fatty Infiltration of Skeletal Muscle: Mechanisms and Comparisons with Bone Marrow Adiposity. Front Endocrinol (Lausanne). 2016;7:69.

43. Collins BC, Kardon G. It takes all kinds: heterogeneity among satellite cells and fibro-

adipogenic progenitors during skeletal muscle regeneration. Development.

2021;148(21).

44. Chazaud B. Macrophages: supportive cells for tissue repair and regeneration.

Immunobiology. 2014;219(3):172-8.

45. Frantz C, Stewart KM, Weaver VM. The extracellular matrix at a glance. J Cell Sci.

2010;123(Pt 24):4195-200.

46. Senk A, Djonov V. Collagen fibers provide guidance cues for capillary regrowth during

regenerative angiogenesis in zebrafish. Sci Rep. 2021;11(1):19520.

47. Hansen-Smith FM, Carlson BM, Irwin KL. Revascularization of the freely grafted

extensor digitorum longus muscle in the rat. Am J Anat. 1980;158(1):65-82.

48. Pries AR, Secomb TW. Origins of heterogeneity in tissue perfusion and metabolism.

Cardiovasc Res. 2009;81(2):328-35.

49. Folker ES, Baylies MK. Nuclear positioning in muscle development and disease.

Front Physiol. 2013;4:363.

Figure captions

Figure 1. Punch biopsy removes all skeletal muscle tissue components. **A.** A biopsy punch was used to create a local injury (diameter, 2 mm) through the GM (muscle perimeter outlined by yellow broken line). **B.** Microvascular perfusion was disrupted at the site of the injury at 1 dpi. Intravascular injection of fluorescent dextran showed the empty biopsy site in addition to a small unperfused region downstream of the injury. **C.** Representative Sirius red staining of collagen at 1 dpi highlighting deposition of a provisional matrix in the wound; * indicates residual gap. **D.** Stitched tile scan images of immunostained whole mount GM at 1 dpi showing masses of inflammatory cells (CD45[+], red) and fibroblasts (PDGFRα[+], magenta) invading the provisional matrix; capillaries (CD31[+], green) retain parallel orientation along surrounding uninjured myofibers.

Figure 2. The microcirculation regenerates through sprouting and intussusceptive angiogenesis. **A.** At 7 dpi, nascent microvessels emigrate from existing vasculature at wound edge. **B.** At 7 dpi, microvessels are nonuniformly enlarged (arrowheads) and extend into the wound with random orientation. **C.** Intussusceptive angiogenesis (white arrowhead) at 10 dpi is integral to revascularization while sprouting continues (yellow arrowhead). **D.** Sites of robust angiogenic activity at 10 dpi create irregular microvascular plexuses. **E.** At 14 dpi, sprouting (yellow arrowhead) and intussusceptive angiogenesis (white arrowhead) expand regenerating microvascular networks. Images are from Cdh5-mTmG reporter mice in which ECs are green (eGFP) and other cells are red.

Figure 3. Tile scanned images showing time course of microvascular and myofiber regeneration after punch biopsy. **A.** At 7 dpi, angiogenesis begins around edges of wound. **B.** At 14 dpi, regenerating microvessels have spanned the wound which remains devoid of myofibers **C.** At 21 dpi, myofibers traverse the wound while capillary networks are disorganized. Dark areas within the regenerated GM reflect interweaving among adjacent myofibers in contrast to parallel (flat) organization of surrounding (uninjured) myofibers. Within the wound, microvessels remodel (arrowhead) to supply a cluster of adipocytes (not visible) where myofibers have not formed. Images are from Cdh5-mTmG mice in which ECs are green (eGFP expression) and all other cells are red (tdTomato expression). **D.** Summary data of microvessel diameters at criterion time points; n = 5-7 mice per time point, *P < 0.05 versus 0 dpi.

Figure 4. Angiogenesis precedes myogenesis during regeneration. Representative images of GM cross sections from the center of the wound at **A.** 7 dpi, **B.** 14 dpi, and **C.** 21 dpi. MyHC (myofibers): red; laminin (basal laminae): white; CD31 (ECs): green; nuclei: blue. **D.** Summary data for length of region devoid of myofibers (shown by broken yellow lines in A and B) and **E.** Vascular area within the wound region devoid of myofibers (outlined by broken yellow boxes in A and B). n = 4 per time point, *P < 0.05 vs 0 dpi. **F.** Cross section of from regenerated region within the wound at 21 dpi stained for MyHC (red) and myonuclei (blue, DAPI). Some myofibers exhibit centrally located nuclei (white arrowheads) while others exhibit peripheral nuclei (yellow arrowheads). Yellow asterisk: region within punch biopsy where myofibers did not regenerate.

Supp. Figure 1. At 1 dpi, Evans Blue dye uptake is restricted to the edges of biopsied

myofibers.

Supp. Figure 2. Blood flow within the regenerating wound at 10 dpi was visualized by

red blood cell transit during intravital microscopy. **A.** Monochrome image of

regenerating microvessels in Cdh5- mTmG mice. **B.** Color coding denotes the presence

and direction of red blood cell flow at 10 dpi.

Supp. Figure 3. Adipocytes accumulate in the biopsy void if myofibers fail to regenerate

at 21 dpi.

Fig 1

A Injury

B 1 dpi
FITC-dextran
1 mm

C 1 dpi
Sirius Red
*
1 mm

D 1 dpi
CD45
PDGFRα
CD31
Merge
1 mm

Fig 2

A 7 dpi

ECs
Other cells

500 µm

B 7 dpi

100 µm

C 10 dpi

100 µm

D 10 dpi

100 µm

E 14 dpi

100 µm

Fig 3

A 7 dpi

B 14 dpi

C 21 dpi

D

Fig 4

Supp Fig 1

1 dpi

Supp Fig 2

A   **10 dpi**

250 μm

B   **10 dpi**

Inflow
Outflow
Both
No flow

250 μm

Supp Fig 3

21 dpi