Plastid phylogenomics clarifies broad-level relationships in *Bulbophyllum*

(Orchidaceae) and provides insights into range evolution of Australasian

section *Adelopetalum*

Lalita Simpson[a,b*], Mark A. Clements[c], Harvey K. Orel[d, e], Darren M. Crayn[a],

Katharina Nargar[a,e]

[a] Australian Tropical Herbarium, James Cook University, PO Box 6811, Cairns, QLD 4870,

Australia

[b] College of Science and Engineering, James Cook University, PO Box 6811, Cairns, QLD

4870, Australia

[c] Centre for Australian National Biodiversity Research, GPO Box 1700, Canberra, ACT

2601, Australia

[d] School of BioSciences, The University of Melbourne, Parkville, Victoria 3052, Australia

[e] National Research Collections Australia, Commonwealth Industrial and Scientific Research

Organisation (CSIRO), GPO Box 1700, Canberra, ACT 2601, Australia

* Correspondence: Lalita Simpson: lalita.simpson1@jcu.edu.au

Keywords

Ancestral area reconstruction, Australasia, Bulbophyllinae, Divergence time estimation,

High-throughput sequencing, Orchidaceae

Abstract

The hyperdiverse orchid genus *Bulbophyllum* is the second largest genus of flowering plants

and exhibits a pantropical distribution with a center of diversity in tropical Asia. The only

Bulbophyllum section with a centre of diversity in Australasia is sect. *Adelopetalum*.

However, phylogenetic placement, interspecific relationships, and spatio-temporal evolution

of the section have remained largely unclear. To infer broad-level relationships within

Bulbophyllum and interspecific relationships within sect. *Adelopetalum*, a genome skimming

dataset was generated for 89 samples, yielding 70 plastid coding regions and the nuclear

ribosomal DNA cistron. For 18 additional samples, Sanger data from two plastid loci (*mat*K,

*ycf*1) and nuclear ITS were added using a supermatrix approach. The study provided new

insights into broad-level relationships in *Bulbophyllum,* including phylogenetic evidence for

the non-monophyly of sections *Beccariana, Brachyantha, Brachypus, Cirrhopetaloides,*
Cirrhopetalum, Desmosanthes, Minutissima, Oxysepala, Polymeres and *Sestochilos*. Section
Adelopetalum and sect. *Minutissima s.s.* formed a highly supported clade that was resolved in
sister group position to the remainder of the genus. Divergence time estimations based on a
relaxed molecular clock model placed the origin of *Bulbophyllum* in the early Oligocene (ca.
33.2 Ma) and of sect. *Adelopetalum* in the late Oligocene (ca. 23.6 Ma). Ancestral range
estimations based on a BAYAREALIKE model identified the Australian continent as
ancestral area of sect. *Adelopetalum.* The section underwent crown diversification during the
mid-Miocene to the late Pleistocene, predominantly in continental Australia. At least two
independent long-distance dispersal events were inferred eastwards from the Australian
continent to New Zealand, and New Caledonia from the early Pliocene onwards, likely
mediated by the predominantly westerly winds of the southern hemisphere. Retraction and
fragmentation of eastern Australian rainforests from the early Miocene onwards are discussed
as likely drivers of lineage divergence within sect. *Adelopetalum*, facilitating allopatric
speciation.

1. Introduction

The hyperdiverse orchid genus *Bulbophyllum* Thouars (Epidendroideae) is the second largest
genus of flowering plants with more than 2,100 species that exhibit exceptional
morphological and ecological diversity (Frodin, 2004; Pridgeon et al., 2014, WCSP 2022).
Species of this predominantly epiphytic genus occur in a wide range of tropical and
subtropical habitats, from montane rainforests to dry deciduous forests, savannah woodlands,
and rocky fields with shrubby vegetation (Pridgeon et al., 2014). *Bulbophyllum* is distributed
pantropically, occupying all botanical continents defined by Brummit (2001) except for
Antarctic and Eurasia. The genus is most diverse on the botanical continent of tropical Asia
(1562 species), also occurring on the botanical continents of Africa (305), temperate Asia
(152), Southern America (88), the Pacific (49), Australasia (Australia and New Zealand; 35),
and Northern America (7) (WCSP, 2022). Centres of diversity are found in tropical Asia in
the floristic regions of Malesia (667) and Papuasia (656) and in the Afrotropics in the western
Indian Ocean region, on the islands of Madagascar, and the Mascarenes (218) (WCSP, 2022).

The high number of species and complex patterns of morphological variation has presented
significant challenges for resolving relationships in *Bulbophyllum* and this is reflected in
substantial taxonomic revisions that have been proposed. Traditionally, the subtribe

Bulbophyllinae Schltr. (tribe Dendrobieae Endl.) included the large genus *Bulbophyllum*

along with smaller genera, such as *Cirrhopetalum* Lindl., *Drymoda* Lindl., *Pedilochilus*

Schltr., *Sunipia* Buch. -Ham. ex Sm., and *Trias* Lindl. (Dressler, 1993; Garay et al., 1994;

Szlachetko and Margonska, 2001). Recent revisions treat all genera within the subtribe

Bulbophyllinae in a more broadly defined *Bulbophyllum* and recognise 97 sections within the

genus (Pridgeon et al., 2014; Vermeulen et al., 2014). Molecular phylogenetic studies have

largely focused on species from specific geographic regions such as Madagascar and the

Mascarenes (Fischer et al., 2007; Gamisch et al., 2015), the Neotropics (Smidt et al., 2013,

2011), and Peninsular Malaysia (Hosseini et al., 2012) or on taxonomic groups such as the

Cirrhopetalum alliance (Hu et al., 2020). Few have taken a global perspective (e.g., Gamisch

and Comes, 2019). These studies revealed a strong biogeographic pattern and four main

clades were identified that include species largely confined or endemic within one broader

geographical area: 1) continental Africa, 2) Madagascar and the Mascarene Islands, 3)

Southern America, or 4) Asia (Fischer et al., 2007; Gamisch et al., 2015; Gamisch and

Comes, 2019; Smidt et al., 2011). The Southern American clade, the Madagascan clade, and

the continental African clade together form a highly supported lineage (Fischer et al., 2007;

Gamisch et al., 2015; Gamisch and Comes, 2019; Smidt et al., 2013; 2011), in sister group

position to the Asian clade (Fischer et al., 2007; Gamisch et al., 2015; Gamisch and Comes,

2019). Previous molecular phylogenetic studies have mainly elucidated relationships within

Madagascan, continental African and Neotropical sections and within the *Cirrhopetalum*

alliance (Fischer et al., 2007; Gamisch et al., 2015; Smidt et al., 2011; Hu et al., 2020).

However, evolutionary relationships within the Asian clade, which also includes taxa from

the Australasian, and Pacific regions, are still poorly understood, and the monophyly of

sections within this clade has remained largely untested.

The study of hyperdiverse groups such as *Bulbophyllum* relies on a robust phylogenetic

framework to ascertain monophyly of its infrageneric taxa. High throughput sequencing

approaches facilitate the establishment of a robust phylogenetic framework that can be used

to investigate broader evolutionary relationships and assess the monophyly of infrageneric

taxa and their trait evolution (Hassemer et al., 2019; van Kleinwee et al., 2022, Nargar et al.

2022). However, phylogenomic studies which provide insights into broad-level evolutionary

relationships within the Asian clade of *Bulbophyllum* are still lacking. This hampers progress

in understanding diversification of its evolutionary lineages in time and space and trait

evolution within this highly diverse genus.

Within *Bulbophyllum*, section *Adelopetalum* has a unique distribution, being the only section with a centre of diversity in Australia (Brummit, 2001; Pridgeon et al., 2014, Vermeulen, 1993), and thus presents an interesting study case for range evolution within *Bulbophyllum*. The section comprises twelve tropical to temperate epi-lithophytic species. Nine species occur along Australia's east coast in the montane forest communities of the Great Dividing Range, with one species (*B. argyropus*) also found on Australian islands (Lord Howe Island, and Norfolk Island). Two species are endemic to the montane forests of New Caledonia (*B. corythium*, *B. lingulatum*) and one to the lowland coastal forests of New Zealand (*B. tuberculatum*). The section was circumscribed based on morphological affinities recognised among ten species from Australia and New Caledonia previously assigned to *Bulbophyllum* sections *Desmosanthes, Racemosae* and *Sestochilus* (Dockrill, 1969; 1992; Vermeulen, 1993). Subsequent treatments recognised two additional species within the section, *B. weinthalii* and *B. exiguum* (Jones and Clements, 2002; Clements and Jones, 2006). Section *Adelopetalum* is characterised by plants having thin creeping rhizomes adpressed to the host, anchored by filamentous roots with small pseudobulbs that are crowded to widely spaced, and a small single flat leaf arising from the apex of the pseudobulb. The inflorescence is single to few-flowered, with small white, cream or yellow flowers, sometimes with red or purple patterns. The petals are smaller than the sepals but similar in shape, with the bases of the lateral sepals fused to the column foot. The fleshly three-lobed labellum is firmly hinged to the apex of the column foot. Previous cladistic analysis of sect. *Adelopetalum* based on morphological characters resolved two main clades within the section, differentiated by the size and shape of the lower margin of the stelidia: the filiform column appendages typical for most *Bulbophyllum* (Vermeulen, 1993). Previous molecular phylogenetic studies based on the nuclear ribosomal ITS region (ITS1 + 5.8S + ITS2) included two to three representatives of the section (Gamisch et al., 2015; Gamisch and Comes, 2019), placing these in an early diverging position within the Asian clade. However, phylogenetic placement of the section within *Bulbophyllum* was not strongly supported (PP<90, BS 97) and thus requires further study. Further, phylogenetic relationships within sect. *Adelopetalum* and its ancestral range evolution are poorly understood and have not yet been investigated using a molecular phylogenetic approach.

The aims of this study were to 1) build a phylogenomic framework for *Bulbophyllum* with focus on the Asian clade 2) assess the monophyly and phylogenetic placement of sect.

Adelopetalum within *Bulbophyllum*; 3) to infer interspecific relationships within sect.

Adelopetalum 4) and to reconstruct the range evolution of sect. *Adelopetalum*.

2. Methods

2.1 Sampling

In total, 136 orchid samples representing 114 species were included in the study.

Representatives of the Asian and Afrotropical/Neotropical clades of *Bulbophylllum* were

included based on previous phylogenetic studies (Fischer et al., 2007; Smidt et al., 2011;

Gamisch and Comes, 2019). Species names were standardised to accepted taxonomy based

on WCSP (2022) and sectional taxonomy followed IOSPE (2022). Exceptions were made for

B. exiguum which was placed in section *Adelopetalum* and *B. wolfei* which was placed in

section *Polymeres* based on Jones and Clements (2002) and Clements and Jones (2006).

From the Asian, Australasian and Pacific regions a broad sampling was included representing

41 sections, i.e. 60% of sections recognised from these regions in the most recent treatment

of the group (Pridgeon et al., 2014). From the Australasian region, all *Bulbophyllum* species

were sampled (Australia: 30, New Zealand: 2). For *Bulbophyllum* sect. *Adelopetalum,* 28

samples were included, representing all 12 species recognised for the section. The

morphologically closely related sect. *Minutissima* was included with nine samples

representing five species comprising all four Australasian and Pacific species and two (of ca.

19) tropical Asian species. The outgroup comprised representatives of subtribe Dendrobineae

which is sister to Bulbophyllinae, and tribes Malaxideae, Arethuseae, Nervilleae, and

Neottieae based on previous molecular phylogenetic studies (Givnish et al., 2015; Górniak et

al., 2010, Serna-Sánchez et al. 2021). Details of the plant material studied, voucher

information and number of sequences generated for each sample are provided in

Supplementary Material S1 and a complete list of loci analysed is provided in Supplementary

Material S2.

2.2 DNA extraction, amplification, and sequencing

Total genomic DNA was extracted from ca. 10 to 20 mg silica-dried leaf material.

Extractions were carried out with commercial extraction kits (Qiagen DNeasy plant kit,

Venlo, Netherlands; ChargeSwitch gDNA plant kit, Invitrogen, Carlsbad, USA) following

the manufacturer's protocols or using the CTAB method (Doyle and Doyle, 1990), with

modifications as described in Weising et al. (2005). Sequence data was generated using both

Sanger sequencing (46 samples) for the nuclear ribosomal ITS region (ITS1, 5.8s, ITS2) and

two plastid genes (*mat*K, *ycf*1) and shotgun high-throughput sequencing (89 samples) to

recover 70 plastid coding sequences (CDS) and the nuclear ribosomal DNA cistron

(Supplementary Material S1). Libraries for high-throughput sequencing were constructed

from 50 to 100 ng total DNA using the TruSeq Nano DNA LT library preparation kit

(Illumina, San Diego, USA) for an insert size of 350 base pairs (bp) and paired-end reads

following the manufacturer's protocol. Libraries were multiplexed 96 times and DNA

sequencing with 125 bp paired-end reads was carried out on an Illumina HiSeq 2500 platform

at the Australian Genomic Research Facility, Melbourne (Australia).

For Sanger sequencing, amplifications for ITS were carried out with primers 17F and 26SER

(Sun et al., 1994), for *mat*K with the primers 19F and, 1326R (Cuénoud et al., 2002), and for

ycf1 with primers 3720F, intR, intF, 5500R (Neubig et al., 2008). PCR reaction protocols are

provided in Supplementary Material S3. Sequencing reactions were carried out using the

amplification primers and sequencing was conducted on an AB3730xl 96-capillary sequencer

(Australian Genome Research Facility, Brisbane, Australia).

2.3 Assembly and alignment

Sequences were assembled and edited in Geneious R10 (Kearse et al., 2012). Illumina

sequences were assembled to a reference set of plastid CDS extracted from *Dendrobium*

catenatum (GenBank accession numbers KJ862886) and for *ycf*68 from *Anoectochilus*

roxburghii (KP776980). To build a reference for the nuclear ribosomal ITS-ETS region,

Illumina reads of *B. boonjee* (CNS_G07175) were first mapped to the ITS-ETS region of

Corallorhiza trifida (JVF2676a). To extend the region assembled, the *B. boonjee* Illumina

reads were then mapped to the *B. boonjee* consensus sequence generated in the initial step,

yielding a *B. boonjee* reference of the nuclear ribosomal DNA cistron (5'ETS, 18s, ITS1,

5.8s, ITS2, 28s, 3'ETS). Illumina sequences for all other samples were assembled against the

B. boonjee nuclear ribosomal DNA cistron reference. Assemblies were carried out with the

highest quality threshold and a minimum coverage of ten reads. The quality of the assemblies

was checked and edited manually where required. Sequences were deposited in GenBank and

ENA. For Sanger sequences, bidirectional reads were assembled in Geneious and edited

manually. Additional sequences were sourced from DRYAD

(https://doi.org/10.5061/dryad.n9r58) for *Coelogyne flaccida* (Givnish et al., 2015). DNA

sequences were combined into supermatrices for the nuclear ribosomal DNA cistron and each

plastid region, aligned using MAFFT v.7.222 (Katoh et al., 2005, 2002) with the default

settings and manually inspected. Plastid regions were concatenated, excluding regions with

low species coverage or problematic alignments such as the NDH genes. The nuclear

supermatrix included 136 accessions, partitioned into coding and non-coding regions

(alignment length: 6,341bp, number of parsimony informative sites: 995 (16%)); and the

plastid supermatrix included 130 accessions, and 70 plastid coding regions, partitioned by

gene and codon position (alignment length: 61,553bp, number of parsimony informative

sites: 5,789 (9%)).

For divergence time estimations, the plastid supermatrix was reduced to one representative

per species (indicated by an asterisk in Supplementary Material S1), comprising 111

accessions (alignment length: 60,984 bp, number of parsimony informative sites: 5,755

(9%)).

2.4 Phylogenetic analysis

Phylogenetic relationships were inferred using maximum likelihood (ML) in IQ-TREE v.

1.6.12 (Nguyen et al., 2015). The best-fit partition scheme and nucleotide substitution model

for each partition was determined with IQ-TREE's ModelFinder (Kalyaanamoorthy et al.

2017) based on the Akaike information criterion (AIC) (Akaike, 1974). Nodal support was

assessed based on 1000 replicates of ultrafast bootstrap approximation with clades receiving

>95 ultrafast bootstrap support (UFBS) considered as well supported (Minh et al., 2013;

Hoang et al., 2018).

2.4 Divergence time estimation

Divergence times were estimated based on the plastid dataset in BEAST2 v. 2.4.8 (Bouckaert

et al., 2014) applying the best fit partition scheme and substitution model as determined by

IQ-TREE's ModelFinder. We tested two molecular clock models: 1) strict clock

(Zuckerkandl and Pauling, 1965) and 2) relaxed lognormal clock (Drummond et al., 2006)

and two models of speciation and extinction: 1) Yule and 2) birth-death (Yule, 1925;

Gernhard et al., 2008). Three secondary calibration points were used applying priors with a

normal distribution and mean ages and 95% higher posterior density (HDP) intervals based

on the results of a family-wide molecular clock analysis by Chomicki et al. (2015). The root

age was set to 55.02 Ma (HDP: 42.0–68.0). The next secondary calibration point was applied

to the last common ancestor of Dendrobineae, Malaxideae, and Arethuseae and was set to

47.77 Ma (HDP: 36.4–59.1). Monophyly was constrained for this node consistent with

relationships reconstructed in previous phylogenetic analyses (Chomicki et al. 2015; Givnish

et al., 2015). The last secondary calibration was set at the stem node of Dendrobieae and

Malaxideae with 38.68 Ma (HDP: 30.8–46.6). An additional calibration based on the fossil

Dendrobium winikaphyllum (Conran et al., 2009) was applied to the stem node of the

Australasian *Dendrobium* clade (*D. macropus, D. cunninghamii,* and *D. muricatum*), using a

uniform distribution with an infinite maximum age and the minimum age constrained to 20.4

Ma, based on the minimum age of the strata containing the fossil (Mildenhall et al. 2014).

Ten independent BEAST analyses were run for 30 million MCMC generations, with trees

sampled every 3×10^4 generations. To assess convergence of independent runs and determine

burn-in fractions, log files were assessed in Tracer v.1.7.1 (Rambaut and Drummond, 2007).

Log and trees files from independent runs were combined in LogCombiner (from the BEAST

package) with a cumulative burn-in fraction of 10%-31% and the sampling frequency set to

generate at least 10,000 tree and log files (Drummond and Bouckaert, 2015). The combined

log file was assessed in Tracer to ensure the effective sample size of all parameters was

above 200. An additional five independent BEAST runs were conducted for the final analysis

using a relaxed log normal clock with birth death speciation to achieve an effective sample

size above 200 for the ucldmean parameter. A maximum clade credibility tree was generated

in TreeAnnotator (BEAST package) with median node heights. To compare clock and

speciation models, the Akaiki information criteria by MCMC app from the BEAST 2

package v 2.6.2 was used to measure the AICM for the combined MCMC runs generated in

the BEAST analysis for each model (Supplementary Material S4).

2.5 Ancestral range analysis

Species distributions were determined from the World Checklist of Selected Plant Families

(WCSP, 2022). Biogeographic areas were largely delineated based on botanical continents

defined by Brummit (2001). The subcontinental regions of Papuasia, Australia and New

Zealand were also recognised to allow a more fine-scaled resolution of range evolution in

these regions (Brummit 2001). The following seven biogeographic areas were coded: a,

Africa; b, temperate Asia; c, tropical Asia; d, Papuasia; e, Australia; f, New Zealand, and g,

Pacific. Ancestral ranges were estimated in RASP v. 4.0 (Yu et al., 2015) with the

BIOGEOBEARS package (Matzke, 2013) based on the maximum clade credibility tree

obtained from the BEAST analysis of the plastid supermatrix, pruned of the outgroups to

Dendrobieae. Three models of range evolution were tested: the dispersal-extinction

cladogenesis model (DEC) (Ree and Smith, 2008), a ML version of Ronquist's parsimony

dispersal-vicariance (DIVA; Ronquist, 1997), termed DIVALIKE (Matzke, 2013), and a

simplified likelihood interpretation of the Bayesian "BayArea" program (Landis et al., 2013)

known as BAYAREALIKE (Matzke, 2013). No constraints were applied to dispersal

direction and the maximum number of ranges was set to five based on the maximum number

of observed areas in extant species. Likelihood values were compared and the model of best

fit determined by AIC score (Akaike, 1974) was used to infer the relative probabilities of

ancestral ranges at each node in the phylogeny (Supplementary Material S5).

3. Results

3.1.1 Phylogenetic relationships – Plastid data

The ML phylogeny inferred from the 70 loci plastid supermatrix provided strong support for

the monophyly of *Bulbophyllum* and its sister group relationship to *Dendrobium* (Fig. 1).

Section *Adelopetalum* and *Minutissima s.s.* formed a highly supported clade, here termed the

Adelopetalum/Minutissima clade, which was resolved in sister group position to the

remainder of the genus (ultrafast bootstrap support/UFBS 98) (Fig. 1). Within the

Adelopetalum/Minutissima clade, all *Adelopetalum* species plus *B. pygmaeum* (sect.

Minutissima) formed a highly supported lineage (UFBS 100), here termed the Adelopetalum

clade. Within the Adelopetalum clade several highly supported groups were resolved: 1) the

argyropus clade consisting of *B. argyropus*, *B. corythium* and *B. tuberculatum* (UFBS 100),

reconstructed in a highly supported sister group relationship to *B. weinthalii* (UFBS 100)*;* 2)

the bracteatum clade, including *B. boonjee*, *B. bracteatum*, and *B. elisae* (UFBS 99); and 3)

the newportii clade comprised of *B. exiguum*, *B. lageniforme*, *B. lilianae*, *B. lingulatum*, and

B. newportii (UFBS 100). Relationships among *B. pygmaeum*, the argyropus clade + *B.*

weinthalii, bracteatum and newportii clades received weak support. Sister to the

Adelopetalum clade was the highly supported Minutissimum clade comprised of three

species of sect. *Minutissima* (*B. globuliforme, B. keekee, B. minutissimum*), including the type

species of the section (UFBS 100) (Fig. 1). Section *Minutissima* was identified as

polyphyletic, with sect. *Minutissima* species also placed within the Adelopetalum clade (*B.*

pygmaeum) and the Asian clade (*B. mucronatum, B. moniliforme*). Within the Asian clade,

sections *Beccariana, Brachyantha, Brachypus, Cirrhopetaloides, Cirrhopetalum,*

Desmosanthes, Oxysepala, Polymeres and *Sestochilos* were also identified as polyphyletic or

paraphyletic.

Our analyses showed that sect. *Adelopetalum* does not share a close relationship with other

Australasian *Bulbophyllum* species, such as those in sect. *Brachypus (B. nematopodum),* sect.

Brachystachyae (B. evasum), sect. *Cirrhopetalum (B. longiflorum),* sect. *Ephippium (B.*

gracillimum), sect. *Monanthes (B. macphersonii),* sect. *Oxysepala (B. gadgarrense, B.*

grandimesense, B. lamingtonense, B. lewisense, B. schillerianum, B. shepherdii, B.

wadsworthii, B. windsorense), sect. *Polymeres (B. bowkettiae, B. johnsonii, B. radicans, B. wolfei),* and sect. *Sestochilus (B. baileyi).* Australian species from each of these sections were placed in nine different positions within the Asian clade. Australian species from section *Polymeres* formed a highly supported clade (UFBS 100), while Australian species from sect. *Oxysepala* formed a moderately supported clade (UFBS 91) and together with the type species of section *Oxysepala* from Papuasia (*B. cladistinum*) formed a close relationship with the Australian representative of section *Monanthes* (*B. macphersonii*) (UFBS 100).

Fig. 1. Maximum likelihood phylogenetic reconstruction of *Bulbophyllum* based on the supermatrix of 70 plastid coding regions. Ultrafast bootstrap values are given adjacent to nodes. Australian species are shown with an asterisk.

Fig. 2. Maximum likelihood phylogenetic reconstruction of *Bulbophyllum* based on the supermatrix of 70 plastid coding regions. Ultrafast bootstrap values are given adjacent to nodes. Australian species are shown with an asterisk.

3.1.2 Phylogenetic relationships – Nuclear data

The ML phylogeny based on the nuclear ribosomal DNA cistron was resolved with overall lower support when compared to analyses based on 70 plastid loci supermatrix (Fig. 2). Relationships among outgroup taxa were concordant with the plastid phylogeny and *Bulbophyllum* was resolved with maximum support. Within *Bulbophyllum,* the Afrotropical (UFBS 97), Asian (UFBS 100) and Adelopetalum/Minutissima (UFBS 100) clades were resolved with high to maximum support, however the relationships among them were poorly supported. Within the Adelopetalum/Minutissima clade the highly supported clades revealed in the plastid phylogeny were also reconstructed based on the nuclear dataset albeit with poor support overall: argyropus clade (UFBS 100), bracteatum clade (UFBS 93), minutissimum clade (UFBS 86), and newportii clade (UFBS 83). Similar to reconstructions based on the plastid phylogeny: relationships among the argyropus, bracteatum and newportii clades, *B. pygmaeum* and *B. weinthalii* were poorly supported; sections *Beccariana, Brachyantha, Brachypus, Cirrhopetaloides, Cirrhopetalum, Desmosanthes, Minutissima*, *Oxysepala, Polymeres,* and *Sestochilos* were identified as polyphyletic or paraphyletic; and Australian species from sect. *Brachystachyae,* sect. *Cirrhopetalum,* sect. *Monanthes,* sect. *Oxysepala,* sect. *Brachypus,* sect. *Polymeres,* sect. *Ephippium,* sect. *Stenochilus* were placed in nine clades across the Asian clade.

329 **Fig. 3.** Maximum likelihood phylogenetic reconstruction of *Bulbophyllum* based on the

330 nuclear ribosomal DNA cistron (5'ETS, 18s, ITS1, 5.8s, ITS2, 28s, 3'ETS). Ultrafast

331 bootstrap values are given adjacent to nodes. Australian species are shown with an asterisk.

Fig. 4. Maximum likelihood phylogenetic reconstruction of *Bulbophyllum* based on nuclear
ribosomal DNA cistron (5'ETS, 18s, ITS1, 5.8s, ITS2, 28s, 3'ETS). Ultrafast bootstrap
values are given adjacent to nodes. Australian species are shown with an asterisk.

3.2 Divergence time estimation

The divergence time analysis based on a relaxed log normal clock and birth death prior with speciation and extinction, which was identified as the model of best fit based on the Akaike information criterion (Supplementary Material S4), is presented here with the Asian and Afrotropical clades collapsed and the complete chronogram provided in Supplementary Material S7. The divergence time analysis based on the plastid dataset was well resolved and highly supported (Fig. 3, Supplementary Material S6). The divergence between *Bulbophyllum* and *Dendrobium* was estimated to have occurred during the early Oligocene, ca. 33.2 Ma (95% highest posterior probability density, HPD: 27.7–39.0). The crown of *Bulbophyllum*, constituting the divergence of the Adelopetalum/Minutissima clade from the remainder of the genus, was dated to the late Oligocene, ca. 24.9 Ma (HPD: 20.1–30.7). Divergence between the Asian clade and the Afrotropical clade was estimated to have taken place during the late Oligocene, ca. 24.3 Ma (HPD: 19.4–29.8) and diversification within the Asian clade was estimated from the mid Miocene 21.4 Ma (HPD: 17.2–26.4 Ma) and the Afrotropical clade from 16.0 Ma (HPD: 10.4–21.7). The crown of the Adelopetalum/Minutissima clade was dated to the late Oligocene, ca. 23.6 Ma (HPD: 18.6–29.1), with the split of the Minutissimum clade from the Adelopetalum clade. The crown of the Adelopetalum clade was dated to the mid Miocene, ca. 15.3 Ma (HPD: 10.6–21.2). The stem branches of major lineages within the Adelopetalum clade were estimated to have diversified during the mid-Miocene: the bracteatum clade was dated to ca. 14.5 Ma (HPD: 10.0–20.4); the lineage giving rise to *B. weinthalii* to ca. 12.1Ma (6.9–17.6); the argyropus clade to ca. 12.1 Ma (6.9–17.6); and the newportii clade to ca. 14.9 Ma (HPD 10.3–20.7). Diversification among species within these lineages took place from the mid-Miocene onwards with the most recent divergence identified during the late Pleistocene among *B. argyropus*, *B. corythium,* and *B. tuberculatum.*

Fig 5. Maximum clade credibility chronogram for *Bulbophyllum* section *Adelopetalum* based on 70 plastid coding sequences. Divergence dates and 95% highest posterior density values are indicated adjacent to nodes. Grey bars indicate 95% highest posterior density. The asterisk denotes the node constrained with a fossil calibration point; the diamond shape denotes nodes which were constrained by secondary calibration points.

3.3 Ancestral range analysis

Model testing of the three biogeographic models (DEC, DIVALIKE, BAYAREALIKE) using the Akaike information criterion identified the BAYAREALIKE model as the model of best fit for the ancestral range estimation (Supplementary Material S5). Ancestral ranges estimated with the BAYAREALIKE model are presented here with the Asian and Afrotropical clades collapsed. The complete chronogram is provided in Supplementary Material S7 and range probabilities for all nodes in Supplementary Material S8.

Australia was reconstructed as the most likely ancestral range for the MRCA of the Adelopetalum clade (range probability (RP) 82) and all nodes within this lineage (RP 72-99) except for the argyropus clade. Range shifts from Australia were inferred from the early Pliocene to the Pacific region (New Caledonia) in the newportii clade, in the lineage giving rise to *B. lingulatum*. Range shifts were also inferred from Australia to the Pacific region

(New Caledonia) and New Zealand either in the lineage giving rise to the MRCA of the

argyropus clade or subsequently within this lineage. Three alternative ancestral ranges were

reconstructed for the MRCA of the argyropus clade: Australia (RP 36), or widespread

distributions including Australia and New Zealand (RP 33) or Australia and New Caledonia

(RP 26). Two alternative ranges were also reconstructed for the MRCA of *B. corythium* and

B. tuberculatum: New Zealand (RP 41) and New Caledonia (RP 34). Taking into account

these alternative scenarios, range shifts within the argyropus clade were estimated to have

occurred sometime between the mid Miocene and late Pliocene (12.1–0.5 Ma). The ancestral

range of MRCA of the Adelopetalum/Minutissima clade and *Bulbophyllum* remained

unresolved in the ancestral range reconstruction. The most likely ancestral range for the

MRCA of the Adeloptalum/Minutissima clade was a widespread distribution across Australia

and tropical Asia (RP 26), while alternative ranges reconstructed included a widespread range

including Australia, tropical Asia and Papuasia (RP 13) and Australia (RP 11). Two

alternative ancestral ranges were reconstructed for the MRCA of *Bulbophyllum*, both

widespread distributions including Australia and tropical Asia (RP 25.9) or Australia, tropical

Asia and Papuasia (RP 21).

Fig 6. Range evolution of *Bulbophyllum* sect. *Adelopetalum*. A) ancestral area reconstruction based on a the BAYAREALIKE model with species extant distributions shown within the grid*;* b) geographic regions delineated in the biogeographic analysis; c) legend of colour-coded geographic regions and shared ancestral area, black with an asterisk represents other ancestral ranges.

4. Discussion

4.1 Phylogenetic relationships

This study provided a broad plastid phylogenetic framework for Asian and Australasian sections of *Bulbophyllum* and revealed a close relationship between sections *Adelopetalum* and *Minutissima s.s.*, that together form a highly supported early diverging lineage within the

genus (Fig. 1, Fig. 2). Relationships based on 70 plastid genes support a sister group

relationship between the *Adelopetalum*/*Minutissima* clade and the remainder of the genus

(Asian + Afrotropical clades). Within the Adelopetalum/Minutissima clade, analyses based

on our 70 plastid loci supermatrix showed a dichotomous split between the highly supported

Minutissima s.s. and Adelopetalum clades. Species were reconstructed in each of these clades

according to their sectional placement with the exception of New Zealand endemic *B.*

pygmaeum (sect. *Minutissima*)*,* which was nested within the Adelopetalum clade, rendering

section *Adelopetalum* paraphyletic. Section *Minutissima* was identified as polyphyletic, with

the Australian (*B. minutissimum, B. globuliforme*) and Pacific species (*B. keekee*) placed in

the Minutissima clade and New Zealand species (*B. pygmaeum*) in the Adelopetalum clade

while the Asian species, *B. mucronatum* and *B. moniliforme* were resolved within the Asian

clade. Section *Minutissima* has undergone numerous taxonomic changes with treatments

ranging from a narrower circumscription recognising species from Australia (Jones and

Clements, 2001), to broader classifications including 23 species from Thailand, Indonesia,

Australia, New Zealand, New Caledonia and New Guinea (Pridgeon et al. 2014). Our

phylogenetic analysis based on plastid and nuclear markers did not reconstruct a close

relationship between sect. *Minutissima* species from the Australasian/Pacific region and

Asian species *B. mucronatum* and *B. moniliforme*. Rather, in our analyses the Australasian

species fell within the Adelopetalum/Minutissima clade while the Asian species were nested

within the Asian clade. The results support morphological studies differentiating sect.

Minutissima species from Australasia and Asia (Jones and Clements, 2001) and show minute

pseudobulbs are a trait that has evolved more than once independently.

While plastid phylogenomics has clarified broad level relationships within *Bulbophyllum*,

further studies are required. Non monophyletic relationships identified in the present study

(e.g., in sections *Beccariana, Brachyantha, Brachypus, Cirrhopetaloides, Cirrhopetalum,*

Desmosanthes, and *Polymeres*) and in previous molecular phylogenetic studies (Fischer et

al., 2007; Smidt et al., 2011; Pridgeon et al. 2014; Hu 2020) highlight the need for further

taxonomic revision within *Bulbophyllum.* Further studies are required with an expanded

sampling of the diverse Asian and Pacific taxa to improve understanding of evolutionary

relationships and test sectional classifications within a phylogenetic framework. Phylogenetic

relationships reconstructed from the nuclear ribosomal DNA cistron were not strongly

supported overall here and in previous studies (Gamisch et al., 2015; Gamisch and Comes,

2019, Hu 2020). Further studies utilising approaches yielding higher coverage from the

nuclear genome, such as target sequence capture methods, provide an opportunity to improve

the understanding of evolutionary relationships in future studies. While assembling datasets

with comprehensive species coverage within mega diverse groups such as *Bulbophyllum*

remains a challenge, the present study provides an example of the use of a broad

phylogenetic framework with targeted sampling within a section, to test the monophyly and

phylogenetic placement of groups of interest.

4.2 Spatio-temporal evolution of Bulbophyllum sect. Adelopetalum

Our divergence time analysis and ancestral range estimations showed that *Bulbophyllum*

section *Adelopetalum* represents an Australasian lineage that originated on the Australian

continent during the late Oligocene to early Miocene. The Australian ancestral range is

largely conserved within the lineage, indicating diversification among species has

predominantly taken place on the Australian continent. The conservation of ancestral range

observed within sect. *Adelopetalum* is consistent with previous phylogenetic analyses of

Bulbophyllum that have shown a strong biogeographic signal among clades, being largely

confined to biogeographic regions such as Madagascar, continental Africa and South

America (Fischer et al., 2007; Gamisch et al., 2015; Gamisch and Comes, 2019; Smidt et al.,

2011). The evolution of *Bulbophyllum* during the early Oligocene occurred subsequently to

the breakup of Gondwana (Matthews, et al. 2016; Zahirovic et al. 2016), implicating long-

distance dispersal (LDD) in the evolution of biogeographical lineages within the genus (Van

den Berg, 2003, Smidt et al., 2011; Gamisch et al., 2015; Gamisch and Comes, 2019).

Nevertheless, the conservation of ancestral ranges observed within the *Adelopetalum* lineage

in this study and strong biogeographic signal among clades identified in previous studies

indicate LDD with successful establishment and persistence has been relatively infrequent

within *Bulbophyllum*. Although the minute wind-dispersed seeds of orchids have a high

dispersal potential, successful establishment in a new area are limited by several factors, such

as the presence of mycorrhizal partners necessary for germination and development, a

suitable host or substrate and microclimatic conditions, and the availability of pollinators

(Arditti and Ghani, 2000; Jersáková and Malinová, 2007; McCormick et al., 2012). Our

results are consistent with previous studies that have identified in situ diversification as the

dominant biogeographic process, despite evidence for LDD, and provide further support for

the hypothesis that the complex requirements for successful establishment, rather than

dispersal limitations, play an important role in constraining the geographic distribution of

orchids (Perez-Escobar and Chomicki et al. 2017; Givnish et al., 2016).

Our phylogenetic analysis further resolved interspecific relationships among sect.

Adelopetalum species (Fig. 1). Divergence time estimation showed that divergence among

species occurred mainly during the Miocene and Pliocene (Fig. 3), during a period of

extensive changes to the distribution of forest vegetation on the Australian continent in

response to drastic climatic changes. During the early Miocene, Australian vegetation

diversified in response to aridification of the Australian continent and the abrupt shift to a

cool dry climate during the mid-Miocene resulted in considerable fragmentation of rainforest

habitats (Martin, 2006, Byrne et al. 2011). *Bulbophyllum* sect. *Adelopetalum* comprises

epiphytic species that occur in mesic forest habitats and thus diversification and

fragmentation of these habitats were likely drivers of allopatric lineage divergence within this

group. Sister group relationships were identified between two species pairs with disjunct

distributions in Australia's northern wet tropical rainforests and south-eastern rainforests (*B.*

boonjee/B. bracteatum and *B. newportii/B. exiguum*). These relationships support the

hypothesis that the diversification and fragmentation of forest habitats in Australia has been

an important driver of lineage divergence in Australia's mesic biome (Byrne et al., 2011,

Simpson et al., 2018).

Whilst the ancestral range was predominantly conserved within the *Adelopetalum* lineage,

range expansion events were inferred from continental Australia across the Coral and Tasman

Seas, to New Caledonia in the lineage giving rise to *B. lingulatum* and to New Zealand and

New Caledonia in the argyropus clade (Fig. 4). New Caledonia and New Zealand each have a

long history of isolation from Australia that predates the evolution of *Bulbophyllum*,

indicating colonisation of these islands by *Bulbophyllum* species has been via LDD

(Matthews et al., 2016). It remains unclear if LDD to New Zealand and New Caledonia in the

argyropus clade occurred from the early Miocene in the lineage giving rise to the MRCA of

the group or subsequently within this clade during the late Pleistocene, thus the spatio

temporal evolution of this lineage remains unresolved. The group shares morphological

affinities and taxonomic treatments recognise one to three species within the group: *B.*

argyropus (Australia's east coast and off shore islands: Lord Howe Island, and Norfolk

Island), *B. corythium* (New Caledonia), and *B. tuberculatum* (New Zealand) (Clements and

Jones, 2002; Halle, 1981; Vermeulen, 1993). Further studies are required to clarify species

delimitation and dispersal patterns utilising population level sampling and molecular

sequencing techniques suited to resolving relationships among closely related species, such

as reduced library representation high-throughput sequencing approaches or target sequence

capture methods recovering highly variable non-coding regions (Peterson et al., 2012;

Weitemier et al., 2014; Folk et al., 2015; Bagley et al. 2020).

The pattern of eastward dispersal observed in range shifts from Australia, across the Coral

and Tasman Seas, is consistent with dispersal patterns inferred in other angiosperms,

including *Abrotanella, Dendrobium, Dracophyllum, Hebe, Korthalsella, Leucopogon,*

Northofagus, Oreobolus, Pterostylis, Rytidosperma (Chacón et al., 2006; Lavarack et al.,

2000; Linder, 1999; Molvray et al., 1999; Puente-Lelièvre et al., 2013; Swenson et al., 2001;

Wagstaff et al., 2010, 2006, 2002, Nargar et al. 2022). The bias towards eastward dispersal

observed within section *Adelopetalum* among other plant groups may be facilitated by the

predominant westerly winds occurring in the southern hemisphere that initiated after the

rifting of Australia and South America from Antarctica during the Eocene (Sanmartín et al.,

2007).

Acknowledgements

This work was supported by the Australian Biological Resources Study (BBR 210-34 to

K.N., D.M.C.., and M.AC..), the Hermon Slade Foundation (HSF 16-04 to KN. and M.AC.),

Bioplatforms Australia (CITES1-33 to DMC and KS), and the Australian Orchid Foundation

(AOF 320/17 to L.S.). The authors are grateful to the KuKu Nyungkal traditional owners and

the Queensland National Parks and Wildlife for permission to sample from wild populations.

We thank B. Gray, D.L. Jones, G. McCraith, M.T. Mathieson, B.P. Molloy, L. Roberts, and

P.D. Ziesing Michael Harrison for contributing plant material to the study. We thank

Guillaume Chomicki for providing files utilised for secondary calibration of our divergence

dating analyses. We gratefully acknowledge the valuable comments and suggestions of three

anonymous reviewers on previous versions of this manuscript.

References

Akaike, H., 1974. A new look at the statistical model identification. IEEE Trans. Automat.
Contr. 19, 716–723. doi:10.1109/TAC.1974.1100705

Arditti, J., Ghani, A.K.A., 2000. Numerical and physical properties of orchid seeds and their
biological implications. New Phytol. 145, 367–421. doi:10.1046/j.1469-8137.2000.00587.x

Bagley, J. C., Uribe-Convers, S., Carlsen, M. M., & Muchhala, N. (2020). Utility of targeted
sequence capture for phylogenomics in rapid, recent angiosperm radiations: Neotropical
Burmeistera bellflowers as a case study. *Molecular Phylogenetics and Evolution*, *152*,
106769.

Bouckaert, R., Heled, J., Kühnert, D., Vaughan, T., Wu, C.-H., Xie, D., Suchard, M.A.,
Rambaut, A., Drummond, A.J., 2014. BEAST 2: A software platform for Bayesian
evolutionary analysis. PLoS Comput. Biol. 10, e1003537. doi:10.1371/journal.pcbi.1003537

Brummitt, R.K., 2001. World geographical scheme for recording plant distributions, edition
2. *Hunt Institute for Botanical Documentation,* Carnegie Mellon University*,* Pittsburgh.

Byrne, M., Steane, D.A., Joseph, L., Yeates, D.K., Jordan, G.J., Crayn, D., Aplin, K., Cantrill,
D.J., Cook, L.G., Crisp, M.D., Keogh, J.S., Melville, J., Moritz, C., Porch, N., Sniderman,
J.M.K., Sunnucks, P., 2011. Decline of a biome : evolution, contraction, fragmentation,
extinction and invasion of the Australian mesic zone biota. J. Biogeogr. 38, 1–22.
doi:10.1111/j.1365-2699.2011.02535.x

Chacón, J., Madriñán, S., Chase, M.W., Bruhl, J.J., 2006. Molecular phylogenetics of *Oreobolus*
(Cyperaceae) and the origin and diversification of the American species. Taxon 55, 359–366.
doi:10.2307/25065583

Chomicki, G., Bidel, L.P.R., Ming, F., Coiro, M., Zhang, X., Wang, Y., Baissac, Y., Jay-
Allemand, C., Renner, S.S., 2015. The velamen protects photosynthetic orchid roots against
UV-B damage, and a large dated phylogeny implies multiple gains and losses of this function
during the Cenozoic. New Phytol. 205, 1330–1341. doi:10.1111/nph.13106

Conran, J.G., Bannister, J.M., Lee, D.E., 2009. Earliest orchid macrofossils: early Miocene
Dendrobium and *Earina* (Orchidaceae: Epidendroideae) from New Zealand. Am. J. Bot. 96,
466–74. doi:10.3732/ajb.0800269

Cuénoud, P., Savolainen, V., Chatrou, L.W., Powell, M., Grayer, R.J., Chase, M.W., 2002.
Molecular phylogenetics of Caryophyllales based on nuclear 18S rDNA and plastid *rbcL*,
atpB, and *matK* DNA sequences. Am. J. Bot. 89, 132–44. doi:10.3732/ajb.89.1.132

Dockrill, A. W. (1969, 1992). *Australian Indigenous Orchids: The epiphytes, the tropical*

terrestrial species. Surrey Beatty & Sons.

Doyle, J.J., Doyle, J.L., 1990. Isolation of plant DNA from fresh tissue. Focus 12, 13–15.

Dressler, R.L., 1993. Phylogeny and classification of the orchid family. Cambridge University

Press.

Drummond, A. J., & Bouckaert, R. R. (2015). *Bayesian evolutionary analysis with BEAST*.

Cambridge University Press.

Drummond, A.J., Ho, S.Y.W., Phillips, M.J., Rambaut, A., Rambaut, A., 2006. Relaxed

Phylogenetics and dating with confidence. PLoS Biol. 4, e88.

doi:10.1371/journal.pbio.0040088

Fischer, G. a, Gravendeel, B., Sieder, A., Andriantiana, J., Heiselmayer, P., Cribb, P.J., Smidt,

E.D.C., Samuel, R., Kiehn, M., 2007. Evolution of resupination in Malagasy species of

Bulbophyllum (Orchidaceae). Mol. Phylogenet. Evol. 45, 358–76.

doi:10.1016/j.ympev.2007.06.023

Folk, R. A., Mandel, J. R., & Freudenstein, J. V. (2015). A protocol for targeted enrichment of

intron-containing sequence markers for recent radiations: A phylogenomic example from

Heuchera (Saxifragaceae). *Applications in Plant Sciences*, *3*(8), 1500039.

Frodin, D.G., 2004. History and concepts of big plant genera. Taxon 53, 753–756.

doi:10.2307/4135449

Gamisch, A., Fischer, G.A., Comes, H.P., 2015. Multiple independent origins of auto-

pollination in tropical orchids (*Bulbophyllum*) in light of the hypothesis of selfing as an

evolutionary dead end. BMC Evol. Biol. 15, 192–209. doi:10.1186/s12862-015-0471-5

Gamisch, A., Comes, H.P., 2019. Clade-age-dependent diversification under high species

turnover shapes species richness disparities among tropical rainforest lineages of

Bulbophyllum (Orchidaceae). BMC Evol. Biol. 19, 93. doi:10.1186/s12862-019-1416-1

Garay, L.A., Hamer, F., Siegerist, E.S., 1994. The genus *Cirrhopetalum* and the genera of the

Bulbophyllum alliance. Nord. J. Bot. 14, 609–646. doi:10.1111/j.1756-1051.1994.tb01080.x

Gernhard, T., 2008. The conditioned reconstructed process. J. Theor. Biol. 253, 769–778.

doi:10.1016/j.jtbi.2008.04.005

Givnish, T.J., Spalink, D., Ames, M., Lyon, S.P., Hunter, S.J., Zuluaga, A., Doucette, A., Caro,
G.G., McDaniel, J., Clements, M.A., Arroyo, M.T.K., Endara, L., Kriebel, R., Williams,
N.H., Cameron, K.M., 2016. Orchid historical biogeography, diversification, Antarctica and
the paradox of orchid dispersal. J. Biogeogr. 43, 1905–1916. doi:10.1111/jbi.12854

Givnish, T.J., Spalink, D., Ames, M., Lyon, S.P., Hunter, S.J., Zuluaga, A., Iles, W.J.D.,
Clements, M.A., Arroyo, M.T.K., Leebens-Mack, J., Endara, L., Kriebel, R., Neubig, K.M.,
Whitten, W.M., Williams, N.H., Cameron, K.M., 2015. Orchid phylogenomics and multiple
drivers of their extraordinary diversification. Proc. R. Soc. B Biol. Sci. 282, 20151553–
20151562. doi:10.1098/rspb.2015.1553

Górniak, M., Paun, O., Chase, M.W., 2010. Phylogenetic relationships within Orchidaceae
based on a low-copy nuclear coding gene, *Xdh*: Congruence with organellar and nuclear
ribosomal DNA results. Mol. Phylogenet. Evol. 56, 784–95.
doi:10.1016/j.ympev.2010.03.003

Halle, N., 1981. New data and carpograms of orchids from New Caledonia. Adansonia 20, 355–
368.

Hassemer, G., Bruun-Lund, S., Shipunov, A. B., Briggs, B. G., Meudt, H. M., & Rønsted, N.
(2019). The application of high-throughput sequencing for taxonomy: the case of Plantago
subg. Plantago (Plantaginaceae). *Molecular phylogenetics and evolution*, *138*, 156-173.
doi.org/10.1016/j.ympev.2019.05.013

Hoang, D.T., Chernomor, O., von Haeseler A., Minh B.Q., and Vinh L.S., 2018. UFBoot2:
Improving the ultrafast bootstrap approximation. Mol. Biol. Evol. 35, 518–522.
https://doi.org/10.1093/molbev/msx281

Hosseini, S., Go, R., Dadkhah, K., & Nuruddin, A. A. (2012). Studies on maturase K sequences
and systematic classification of Bulbophyllum in Peninsular Malaysia. *Pak. J. Bot*, *44*(6),
2047-2054.

Hu, A. Q., Gale, S. W., Liu, Z. J., Suddee, S., Hsu, T. C., Fischer, G. A., & Saunders, R. M.
(2020). Molecular phylogenetics and floral evolution of the Cirrhopetalum alliance
(Bulbophyllum, Orchidaceae): Evolutionary transitions and phylogenetic signal variation.
Molecular phylogenetics and evolution, *143*, 106689.

IOSPE (2022, 6th July) Internet Orchid Species Photo Encyclopedia.
http://www.orchidspecies.com/

Jersáková, J. and Malinová, T., 2007, Spatial aspects of seed dispersal and seedling recruitment
in orchids. New Phytol., 176, 237–241. doi:10.1111/j.1469-8137.2007.02223.x

Jones, D.L., Clements, M.A., 2001. *Oncophyllum*, a new genus of Orchidaceae from Australia.
The Orchadian 13, 420–425.

Jones, D.L., Clements, M.A., 2002. Nomenclatural changes in the Australian and New Zealand
Bulbophyllinae and Eriinae (Orchidaceae). The Orchadian 13, 498–501

Kalyaanamoorthy S., Minh B.Q., Wong T.K.F., von Haeseler A., and Jermiin L.S., 2017.
ModelFinder: fast model selection for accurate phylogenetic estimates. Nat. Methods. 14,
587–589. DOI: 10.1038/nmeth.4285

Katoh, K., Kuma, K., Toh, H., Miyata, T., 2005. MAFFT version 5: improvement in accuracy of
multiple sequence alignment. Nucleic Acids Res. 33, 511–518. doi:10.1093/nar/gki198

Katoh, K., Misawa, K., Kuma, K., Miyata, T., 2002. MAFFT: a novel method for rapid multiple
sequence alignment based on fast Fourier transform. Nucleic Acids Res. 30, 3059–66.

Kearse, M., Moir, R., Wilson, A., Stones-Havas, S., Cheung, M., Sturrock, S., Buxton, S.,
Cooper, A., Markowitz, S., Duran, C., Thierer, T., Ashton, B., Meintjes, P., Drummond, A.,
2012. Geneious Basic: An integrated and extendable desktop software platform for the
organization and analysis of sequence data. Bioinformatics 28, 1647–1649.
doi:10.1093/bioinformatics/bts199

Landis, M. J., Matzke, N. J., Moore, B. R., & Huelsenbeck, J. P. (2013). Bayesian analysis of
biogeography when the number of areas is large. *Systematic biology*, *62*(6), 789-804.

Lavarack, B., Harris, W., Stocker, G., 2000. *Dendrobium* and its relatives. Timber Press,
Portland.

Linder, H.P., 1999. *Rytidosperma vickeryae* — a new danthonioid grass from Kosciuszko (New
South Wales, Australia): Morphology, phylogeny and biogeography. Aust. Syst. Bot. 12,
743–755. doi:10.1071/SB97046

Martin, H.A., 2006. Cenozoic climatic change and the development of the arid vegetation in
Australia. J. Arid Environ. 66, 533–563. doi:10.1016/J.JARIDENV.2006.01.009

Matthews, K.J., Maloney, K.T., Zahirovic, S., Williams, S.E., Seton, M., Müller, R.D., 2016.
Global plate boundary evolution and kinematics since the late Paleozoic. Glob. Planet.
Change 146, 226–250. doi:10.1016/J.GLOPLACHA.2016.10.002

Matzke, N.J., 2013. BioGeoBEARS: Biogeography with Bayesian (and Likelihood)

evolutionary analysis in R Scripts.

McCormick, M.K., Taylor, D., Juhaszova, K., Burnett, R.K., Whigham, D.F., O'neill, J.P.,

2012. Limitations on orchid recruitment: not a simple picture. Mol. Ecol. 21, 1511–1523.

doi:10.1111/j.1365-294X.2012.05468.x

Mildenhall, D.C., Kennedy, E.M., Lee, D.E., Kaulfuss, U., Bannister, J.M., Fox, B. and Conran,

J.G., 2014. Palynology of the early Miocene Foulden Maar, Otago, New Zealand: Diversity

following destruction. *Review of Palaeobotany and Palynology*, *204*, pp.27-42.

Minh, B. Q., Nguyen, M. A. T., & von Haeseler, A. (2013). Ultrafast approximation for

phylogenetic bootstrap. *Molecular biology and evolution*, *30*(5), 1188-1195.

Molvray, M., Kores, P.J., Chase, M.W., 1999. Phylogenetic relationships within *Korthalsella*

(Viscaceae) based on nuclear ITS and plastid *trnL-F* sequence data. Am. J. Bot. 86, 249–260.

doi:10.2307/2656940

Nargar, K., O'Hara, K., Mertin, A., Bent, S., Nauheimer, L., Simpson, L., ... & Clements, M. A.

(2022). Evolutionary relationships and range evolution of greenhood orchids (subtribe

Pterostylidinae): insights from plastid phylogenomics. Front. Plant Sci. In press

doi.org/10.3389/fpls.2022.912089

Neubig, K.M., Whitten, A.W.M., Carlsward, B.S., Blanco, M.A., Endara, L., Williams, N.H.,

Moore, M., 2008. Phylogenetic utility of *ycf1* in orchids: a plastid gene more variable than

matK. Plant Syst. Evol. 277, 75–84. doi:10.1007/s00606-008-0105-0

Nguyen, L. T., Schmidt, H. A., Von Haeseler, A., & Minh, B. Q. (2015). IQ-TREE: a fast and

effective stochastic algorithm for estimating maximum-likelihood phylogenies. *Molecular*

biology and evolution, *32*(1), 268-274. doi.org/10.1093/molbev/msu300

Peterson, B. K., Weber, J. N., Kay, E. H., Fisher, H. S., & Hoekstra, H. E. (2012). Double digest

RADseq: an inexpensive method for de novo SNP discovery and genotyping in model and

non-model species. *PloS one*, *7*(5), e37135.

Pridgeon, A.M., Cribb, P.J., Chase, M.W., Rasmussen, F.N., 2014. Genera Orchidacearum

Volume 6 Epidendroideae. Oxford University Press, New York, USA.

Puente-Lelièvre, C., Harrington, M.G., Brown, E.A., Kuzmina, M., Crayn, D.M., 2013.

Cenozoic extinction and recolonization in the New Zealand flora: The case of the fleshy-

fruited epacrids (Styphelieae, Styphelioideae, Ericaceae). Mol. Phylogenet. Evol. 66, 203–

214. doi:10.1016/J.YMPEV.2012.09.027

Drummond, A. J., & Rambaut, A. (2007). BEAST: Bayesian evolutionary analysis by sampling

trees. *BMC evolutionary biology*, *7*(1), 1-8.

Rambaut, A., Drummond, A.J., 2007. Tracer v.1.5. Available from:

http://beast.bio.ed.ac.uk/Tracer

Ree, R.H., Smith, S.A., 2008. Maximum likelihood inference of geographic range evolution by

dispersal, local extinction, and cladogenesis. Syst. Biol. 57, 4–14.

doi:10.1080/10635150701883881

Ronquist, F., 1997. Dispersal-vicariance analysis: a new approach to the quantification of

historical biogeography. Syst. Biol 46, 195–203.

Ronquist, F., 2001. DIVA version 1.2. Computer program for MacOS and Win32.

Sanmartín, I., Wanntorp, L., Winkworth, R.C., 2007. West Wind Drift revisited: testing for

directional dispersal in the Southern Hemisphere using event-based tree fitting. J. Biogeogr.

34, 398–416. doi:10.1111/j.1365-2699.2006.01655.x

Serna-Sánchez, M. A., Pérez-Escobar, O. A., Bogarín, D., Torres-Jimenez, M. F., Alvarez-Yela,

A. C., Arcila-Galvis, J. E., ... & Arias, T. (2021). Plastid phylogenomics resolves ambiguous

relationships within the orchid family and provides a solid timeframe for biogeography and

macroevolution. *Scientific reports*, *11*(1), 1-11.

Simpson, L., Clements, M.A., Crayn, D.M., Nargar, K., 2018. Evolution in Australia's mesic

biome under past and future climates: insights from a phylogenetic study of the Australian

Rock Orchids (*Dendrobium speciosum* complex, Orchidaceae). Mol. Phylogenet. Evol. 118,

32–46. doi:10.1016/J.YMPEV.2017.09.004

Smidt, E.C., Gallo, L.W., Scatena, V.L., 2013. Leaf anatomical and molecular studies in

Bulbophyllum section *Micranthae* (Orchidaceae) and their implications for systematics.

Brazilian J. Bot. 36, 75–82. doi:10.1007/s40415-013-0008-3

Smidt, E.C., Borba, E.L., Gravendeel, B., Fischer, G.A., Berg, C. Van Den, 2011. Molecular

phylogeny of the Neotropical sections of *Bulbophyllum* (Orchidaceae) using nuclear and

plastid spacers. Taxon 60, 1050–1064.

Sun, Y., Skinner, D.Z., Liang, G.H., Hulbert, S.H., 1994. Phylogenetic analysis of *Sorghum* and

related taxa using internal transcribed spacers of nuclear ribosomal DNA. Theor. Appl.

Genet. 89, 26–32. doi:10.1007/BF00226978

Swenson, U., Backlund, A., McLoughlin, S., Hill, R.S., 2001. *Nothofagus* biogeography

revisited with special emphasis on the enigmatic distribution of subgenus *Brassospora* in

New Caledonia. Cladistics 17, 28–47. doi:10.1111/j.1096-0031.2001.tb00109.x

Szlachetko, D.L., Margonska, H.B., 2001. Genera et species Orchidalium. 3. Polish Bot. J. 461,

113–121.

Pérez-Escobar, O.A., Chomicki, G., Condamine, F.L., Karremans, A.P., Bogarín, D., Matzke,

N.J., Silvestro, D., Antonelli, A., 2017. Recent origin and rapid speciation of Neotropical

orchids in the world's richest plant biodiversity hotspot. New Phytol. 215, 891–905.

doi:10.1111/nph.14629

Van den Berg, C. 2003. Estudos em sistemática molecular na família Orchidaceae. Thesis

presented for obtaining the title of Full Pro-fessor, Universidade Estadual de Feira de

Santana, Bahia, Brazil.

van Kleinwee, I., Larridon, I., Shah, T., Bauters, K., Asselman, P., Goetghebeur, P., ... &

Veltjen, E. (2022). Plastid phylogenomics of the Sansevieria Clade of Dracaena

(Asparagaceae) resolves a recent radiation. *Molecular Phylogenetics and Evolution*, 107404.

doi.org/10.1016/j.ympev.2022.107404

Vermeulen, J.J., 1993. A taxonomic revision of *Bulbophyllum* sections *Adelopetalum*,

Lepanthanthe, *Macrouris*, *Pelma*, *Peltopus*, and *Unicifera* (Orchidaceae). Orchid Monogr. 7.

Vermeulen, J.J., Schuiteman, A., De Vogel, E.F., 2014. Nomenclatural changes in *Bulbophyllum*

(Orchidaceae; Epidendroideae). Phytotaxa 166, 101–113. doi:10.11646/phytotaxa.166.2.1

Wagstaff, S.J., Bayly, M.J., Garnock-Jones, P.J., ip Garnock-Jones, P.J., Albach, D.C., 2002.

Classification, origin, and diversification of the New Zealand *Hebes* (Scrophulariaceae). Ann.

Missouri Bot. Gard. 89, 38–63.

Wagstaff, S.J., Breitwieser, I., Swenson, U., 2006. Origin and relationships of the Austral genus

Abrotanella (Asteraceae) inferred from DNA sequences. Taxon 55, 95–106.

doi:10.2307/25065531

Wagstaff, S.J., Dawson, M.I., Venter, S., Munzinger, J., Crayn, D.M., Steane, D.A., Lemson,

K.L., 2010. Origin, diversification, and classification of the Australasian genus

Dracophyllum (Richeeae, Ericaceae). Ann. Missouri Bot. Gard. 97, 235–258.

doi:10.3417/2008130

Weitemier, K., Straub, S. C., Cronn, R. C., Fishbein, M., Schmickl, R., McDonnell, A., &

Liston, A. (2014). Hyb-Seq: Combining target enrichment and genome skimming for plant

phylogenomics. *Applications in plant sciences*, *2*(9), 1400042.

WCSP (6th July, 2022). World Checklist of Selected Plant Families. *Facilitated by the Royal*

Botanic Gardens, Kew. http://wcsp.science.kew.org/ Retrieved 2022, 6th July.

Weising, K., Nybom, H., Pfenninger, M., Wolff, K., Kahl, G., 2005. DNA fingerprinting in

plants: principles, methods, and applications 2nd. ed. CRC Press, BocaRaton, Florida.

Yu, Y., Harris, A.J., Blair, C., He, X., 2015. RASP (Reconstruct Ancestral State in

Phylogenies): A tool for historical biogeography. Mol. Phylogenet. Evol. 87, 46–49.

doi:10.1016/J.YMPEV.2015.03.008

Yule, G.U., 1925. A Mathematical Theory of Evolution, Based on the Conclusions of Dr. J. C.

Willis, F.R.S. Philos. Trans. R. Soc. B Biol. Sci. 213, 21–87. doi:10.1098/rstb.1925.0002

Zahirovic, S., Matthews, K. J., Flament, N., Müller, R. D., Hill, K. C., Seton, M., & Gurnis, M.

(2016). Tectonic evolution and deep mantle structure of the eastern Tethys since the latest

Jurassic. *Earth-Science Reviews*, *162*, 293-337.

Zuckerkandl, E., Pauling, L., 1965. Molecules as documents of evolutionary history. J. Theor.

Biol. 8, 357–366. doi:10.1016/0022-5193(65)90083-4

Supplementary Material

Supplementary Material S3. PCR reaction protocols.

Supplementary Material S4. Comparison of likelihood scores for divergence dating analyses

Supplementary Material S5. Comparison of likelihood scores for three models of range

evolution.

Supplementary Material S6. Maximum clade credibility chronogram for *Bulbophyllum* based

on 70 plastid coding sequences.

Supplementary Material S7. Ancestral area reconstruction based on a the BAYAREALIKE

model.