Trophoblast glycoprotein is required for efficient synaptic vesicle exocytosis from retinal

rod bipolar cells.

Colin M Wakeham[1], Qing Shi[1], Gaoying Ren[1], Tammie L Haley[1], Robert M Duvoisin[1], Henrique

von Gersdorff[2], Catherine W Morgans[1]

[1]Department of Chemical Physiology & Biochemistry, Oregon Health & Science University,

Portland, OR 97239, USA

[2]Vollum Institute, Oregon Health & Science University, Portland, OR 97239, USA

Corresponding Author

Catherine W Morgans

Chemical Physiology & Biochemistry, L-334

Oregon Health & Science University

3181 SW Sam Jackson Park Rd

Portland, OR 97239

Email: morgansc@ohsu.edu

Acknowledgements

This work was supported by NIH grants to CWM (R01 EY029985 and P30 EY010572) and HvG

(R01 EY014043 and DC012938). The authors would like to thank R Lane Brown, Marc A

Meadows, and André A Dagostin for assistance with patch clamp electrophysiology and data

analysis. We would also like to thank the OHSU Neuroscience Imaging Center's Ultrastructure

and Single Particle Microscopy Core (P30 NS061800) for FIB-SEM data collection.

Author contributions

Study design: CMW, RD, HvG, CWM; Experimentation: CMW, HS, GR, TLH; Data analysis:

CMW, HS, RD, HvG, CWM; Figure and manuscript preparation: CMW, HS, RD, HvG, CWM;

Editing and review: CMW, RD, HvG, CWM

Conflict of interest statement

The authors declare no competing financial conflicts of interest.

33
34
35

Abstract

Rod bipolar cells (RBCs) faithfully transmit light-driven signals from rod photoreceptors in the outer retina to third order neurons in the inner retina. Recently, significant work has focused on the role of leucine-rich repeat (LRR) proteins in synaptic development and signal transduction at RBC synapses. We previously identified trophoblast glycoprotein (TPBG) as a novel transmembrane LRR protein localized to the dendrites and axon terminals of RBCs. We now examine the effects on RBC physiology and retinal processing of TPBG genetic knockout in mice using immunofluorescence and electrophysiological techniques. The scotopic electroretinogram showed a modest increase in the b-wave and a marked attenuation in oscillatory potentials in the TPBG knockout. No effect of TPBG knockout was observed on the RBC dendritic morphology, TRPM1 currents, or RBC excitability. Because scotopic oscillatory potentials primarily reflect RBC-driven rhythmic activity of the inner retina, we investigated the contribution of TPBG to downstream transmission from RBCs to third-order neurons. Using electron microscopy, we found shorter synaptic ribbons in TPBG knockout axon terminals in RBCs. Time-resolved capacitance measurements indicated that TPBG knockout reduces synaptic vesicle exocytosis and subsequent GABAergic reciprocal feedback without altering voltage-gated Ca^{2+} currents. Thus, TPBG is required for normal synaptic ribbon development and efficient neurotransmitter release from RBCs to downstream cells. Our results highlight a novel synaptic role for TPBG at RBC ribbon synapses and support further examination into the mechanisms by which TPBG regulates RBC physiology.

Graphical abstract

Introduction

Rod bipolar cells (RBCs) receive light-dependent synaptic input from rod photoreceptors in the outer plexiform layer (OPL) and drive retinal output by synapsing with AII amacrine cells (AII-ACs) in the inner plexiform layer (IPL). As the only dedicated bipolar cell in the primary rod visual pathway, RBCs must be able to reliably transmit visual signals over the whole range of rod

sensitivity. Accurate transmission of light signals requires precise synaptic targeting and tight
control of signal transduction between RBCs and its synaptic partners. Identifying and
characterizing the proteins required for the optimization and regulation of RBC synaptic
development and function is essential for expanding our understanding of how RBCs contribute
to vision.

Leucine-rich repeat (LRR) proteins form a large class of transmembrane proteins, each
containing multiple leucine-rich motifs arranged to form extracellular protein binding domains
(Kobe & Deisenhofer, 1994; Kobe & Kajava, 2001). Several LRR proteins identified in RBCs have
been implicated in the development and maintenance of synaptic morphology and function. At
synapses between photoreceptors and ON-bipolar cells, LRIT3 and nyctalopin are required for
the localization of signal transduction components to the post-synaptic compartment (Cao,
Posokhova, & Martemyanov, 2011; Hasan et al., 2019, 2020; Neuillé et al., 2017, 2015; Pearring
et al., 2011). In RBC dendrites, LRRTM4 forms a trans-synaptic complex with rod spherules
(Agosto & Wensel, 2020), whereas in RBC axon terminals, LRRMT4 is required for functional
GABA receptor clustering, reciprocal feedback inhibition, and dyad synapse formation (Sinha et
al., 2020). The ELFN family of LRR proteins form trans-synaptic complexes between
photoreceptors and bipolar cells without which functioning connections fail to develop and retinal
circuitry is mis-wired (Cao et al., 2015, 2020).

We recently identified trophoblast glycoprotein (TPBG, also called 5T4 or WAIF1 (Wnt-
activated inhibitory factor 1)) in a phosphoproteomics screen as a novel LRR transmembrane
glycoprotein in mouse RBCs that undergoes PKCα-dependent phosphorylation (Wakeham et al.,
2019). TPBG contains eight N-terminal extracellular LRR motifs and a short intracellular C-
terminal tail with a PDZ-interacting domain (Zhao, Malinauskas, Harlos, & Jones, 2014). TPBG
localizes to the dendrites and synaptic terminals of RBCs and its expression is closely linked to
eye opening and TRPM1 expression, suggesting that TPBG may play a role in RBC development
or physiology (Wakeham, Ren, & Morgans, 2020).

Because the role of TPBG in the retina has not been established, we sought to
characterize the effects of genetic knockout of TPBG in mouse RBCs using immunofluorescence
confocal microscopy, electron microscopy, and electrophysiological techniques. RBC synapses
form specialized active zones, each containing a synaptic ribbon, that permit a greater bandwidth
of information transfer than conventional synapses. Ribbon synapses are noted for tight vesicle-
Ca^{2+} coupling and a large supply of readily-available synaptic vesicles tethered close to the active
zone (Dieck & Brandstätter, 2006; Morgans, 2000; Moser, Grabner, & Schmitz, 2020). In this
study, we report that TPBG is required for normal synaptic ribbon morphology and efficient

synaptic vesicle exocytosis at RBC synapses, implicating TPBG in ribbon synapse development

and function.

Methods

Animals and statistical analysis

Constitutive TPBG knockout (TPBG-KO; KO) mice were B6;129S5-Tpbg^{tm1Lex}/Mmucd

(MMRRC; UC Davis; Davis, CA, USA; Cat# 031630-UCD; RRID: MMRRC_031630-UCD). KO

mice were purchased as cryopreserved sperm and reconstituted into a C57BL/6J (The Jackson

Laboratory; Bar Harbor, M; Cat# 000664; RRID: IMSR_JAX:000664) background. Age-matched

homozygous wild type (WT) littermates were used as controls. We used mice of both sexes and

all mice were maintained on a 12-hour light/dark cycle and provided food and water ad libitum.

Animals for immunofluorescence and electroretinogram experiments were 3-6 months old while

animals used for patch clamp electrophysiology experiments were three months old. At least three

mice were used from each condition for each experiment except for electron microscopy, which

used one block per genotype. All animal procedures were in accordance with the National

Institutes of Health guidelines and approved by the Oregon Health & Science University

Institutional Animal Care and Use Committee.

Statistical analyses were performed in Prism 9 (GraphPad; San Diego, CA, USA) and

reported as mean $\pm$ SEM. We detected significance between conditions using unpaired Welch's

t-tests and Brown-Forsythe ANOVA tests. Statistical significance is noted by asterisks (not

significant (ns): $p > 0.05$; *: $p < 0.05$; **: $p < 0.01$; ***: $p < 0.001$; ****: $p < 0.0001$).

Immunofluorescence

Fresh eyecups were prepared from dissected eyes by cutting behind the ora serrata and

removing the cornea and lens. For sectioning, eyecups were fixed for 30 min by immersion in

fresh 4% paraformaldehyde (PFA) in PBS. The fixed eyecups were washed in PBS and then

cryoprotected via sequential immersion in 10, 20, and 30% sucrose in PBS. Sections were cut at

20 μm thickness. Retina sections were post-fixed for 10 minutes in 4% PFA, blocked and

permeabilized by incubation at room temperature for one hour in antibody incubation solution

(AIS; 3% normal horse serum, 0.5% Triton X-100, 0.025% NaN$_3$ in PBS), and then incubated in

primary antibodies diluted in AIS for one hour at room temperature. After washing with PBS, the

sections were incubated in secondary antibodies diluted in AIS for one hour at room temperature.

The slides were washed again in PBS and then mounted with VECTASHIELD PLUS antifade

mounting media with DAPI (Vector Laboratories; Cat# H-2000-10).

For whole mount immunofluorescence, eyecups were fixed for 45 min by immersion in

fresh 4% PFA in PBS. The fixed eyecups were washed in PBS then retinas were removed prior

to blocking and permeabilization for one hour in AIS. The retinas were incubated in primary

antibodies diluted in AIS for four days at 4° C then washed overnight in PBST (PBS + 0.05%

Triton X-100) at 4° C. Retinas were incubated in secondary antibodies diluted in AIS overnight at

4° C then washed again in PBST. Finally, the retinas were mounted with VECTASHIELD PLUS

antifade mounting media.

Immunofluorescence images were taken with a Leica TCS SP8 X confocal microscope

(Leica Biosystems; Wetzlar, Germany) using a Leica HC PL APO CS2 63x/1.40 oil immersion

objective (Leica; Cat# 15506350). Brightness and contrast were adjusted equally across

genotypes using Fiji (Schindelin et al., 2012). A Gaussian blur using a radius of 0.5 pixels was

used to remove graininess from images. See figure legends for specific information on Z-step

distance and number of steps. To compare the spatial relationship between presynaptic and

postsynaptic proteins in the OPL, ROI lines were drawn across $Ca_v1.4$-GPR179 pairs and the

intensity of both fluorescent channels was plotted against the line length in Fiji. The distance

between the two fluorescence peaks approximates the distance across the synapse. To examine

fluorescent puncta in RBC axon terminals in the IPL, PKCα binaries were created in Fiji using

default auto-thresholds to isolate RBC axon terminals. These binaries were then merged with the

other channels to remove PKCα-negative puncta. Fluorescent puncta were counted in PKCα-

positive axon terminals and normalized to axon terminal area. Mander's colocalization coefficients

were calculated using the JACoP plugin in Fiji to compare colocalization patterns between $Ca_v1.4$

and RIBEYE puncta after thresholding with Otsu's method. The M1 coefficient was used for the

fraction of $Ca_v1.4$ containing RIBEYE and the M2 coefficient was for the fraction of RIBEYE

containing $Ca_v1.4$. Colocalization of puncta in PKCα-negative areas was also calculated as a

control.

Immunoblotting

Retinas were extracted from freshly dissected eyes, suspended in chilled lysis buffer (50

mM Tris pH 7.4, 150 mM NaCl, 1 mM EDTA, 1% Triton X-100, 1% deoxycholate, 0.1% SDS) with

1X protease/phosphatase inhibitor cocktail (Cell Signaling Technology; Danvers, MA, USA; Cat#

5872) and homogenized with a Teflon-glass homogenizer. The lysate was centrifuged for 15

minutes at 16,400 rpm and 4° C, and the pellet was discarded. Retinal lysates were diluted to a

protein concentration of 1 μg/μl in lysis buffer and brought to 1X NuPAGE LDS Sample Buffer

(Thermo Fisher Scientific; Cat# NP0007) and 1X NuPAGE Sample Reducing Agent (Thermo

Fisher Scientific; Cat# NP0009). Pre-cast NuPAGE 1 mm 4-12% Bis-Tris gels (Thermo Fisher

Scientific; Cat# NP0322BOX) were loaded and run at 200 V and 140 mA for 55 minutes in 1X

NuPAGE BOLT SDS Running Buffer (Thermo Fisher Scientific; Cat# B0001).

Proteins were transferred onto PVDF membranes using a wet transfer system and 1X

NuPAGE Transfer Buffer (Thermo Fisher Scientific; Cat# NP00061) with 5% MeOH at 300 mA

for two hours. The membranes were then rinsed with methanol and blocked for one hour in

Odyssey Blocking Buffer TBS (LI-COR Biosciences; Cat# 927-50003) on a shaker at room

temperature, before being incubated in primary antibody diluted in Odyssey buffer at 4° C

overnight. The membranes were washed in TBST (Tris-buffered saline with 0.1% Tween-20),

then incubated in secondary antibody diluted in Odyssey buffer for one hour at room temperature

before being washed in TBST. Dried blots were imaged using a LI-COR Odyssey CLx Imaging

System at 680 and 800 nm.

Electron microscopy

Mice were euthanized and enucleated and the retinas were quicky dissected and fixed for

30 min by immersion in 3% PFA and 1% glutaraldehyde. The tissue was then processed for

electron microscopy using the Dresden protocol (Paridaen, Wilsch-Bräuninger, & Huttner, 2013).

Three-dimensional data was acquired using the Helios G3 NanoLab DualBeam FIB-SEM as

previous described. Briefly, the resin blocks were trimmed to contain the ganglion cell layer and

the IPL near the top edge of the block, mounted to a 45° pre-tilted stub, and coated with 8 nm of

carbon. A focused beam of gallium atoms ablated 4 nm off the surface of the sample (FIB

conditions: 30 keV accelerating voltage, 790 pA beam current) over the SEM imaging area of 25

μm x 10 μm. The freshly ablated surface was then imaged by backscattered electrons using the

In-Column Detector (SEM conditions: 3 keV accelerating voltage, 400 pA beam current, 4 nm per

pixel, 3 μs dwell time; image resolution: 6144 x 4086, 4 nm isotropic). Ablation and imaging cycles

were run over a three-day period resulting in a sample depth of roughly 5 μm (1373 images for

the WT dataset and 1284 images for the TPBG-KO dataset). Image stacks were registered using

Linear Stack Alignment with the SIFT algorithm in Fiji and cropped to remove alignment artifacts.

RBC synaptic terminals were identified by their morphology (size, shape, presence of synaptic

ribbons, and arrangement of postsynaptic elements) and their proximity to the ganglion cell layer.

Electroretinogram recording

Mice were dark-adapted overnight and prepared for recording under dim red light.
Anesthesia was initiated via IP injection of ketamine:xylazine (100:10 mg/kg) and maintained with
booster injections (30:3 mg/kg) at approximately 30 minute intervals until completion of the
experiment. Body temperature was maintained at 36.5-38° C. Before recording, the pupils were
dilated with 2.5% phenylephrine and 1% tropicamide, and the cornea was anesthetized with 1%
topical proparacaine. A wire loop placed under the upper teeth was used to draw the mouse into
a custom-made holder that stabilized the head and permitted delivery of 95% O_2/5% CO_2
(carbogen; ~0.25 l/min). The recording was made using a custom contact lens electrode featuring
a central platinum wire placed against the cornea with a small drop of hypromellose to prevent
drying. A loop electrode placed behind the eye served as a reference electrode and a needle
electrode placed in the tail served as a ground. The mice were placed in a Ganzfield sphere and
light stimuli were provided by custom 502 nm LED photostimulators. Light stimulus intensity was
controlled with neutral density filters and by altering flash duration and was measured post-hoc
with an ILT5000 radiometer (International Light; Newburyport, MA; USA) using a scotopic filter.
Traces were recorded with customized software (ERGLab; Richard Weleber; Casey Eye Institute;
Portland, OR, USA).

Full-field scotopic electroretinograms (ERGs) from both eyes were recorded
simultaneously after a series light flashes ranging from 8.71 x 10^{-1} to 8.51 x 10^1 cd•s/m^2 (-0.06 to
1.93 log(cd•s/m^2) (Xiong et al., 2015). For dimmer intensities (-1 to 0 log(cd•s/m^2)), three trials
were averaged with inter-flash intervals of 20 seconds. For brighter intensities (0 to 2
log(cd•s/m^2)), two trials were averaged with inter-flash intervals of 30 to 120 seconds.

ERGLab data was exported for offline processing using a custom Python script and the
SciPy package (https://docs.scipy.org/doc). Statistical analyses were performed in Prism 9
(GraphPad; San Diego, CA, USA). The start of the flash stimulus was set to time zero and traces
were baselined. A-waves and b-waves were isolated using a low-pass filter (60 Hz) and OPs were
isolated with a Butterworth band-pass filter (30-300 Hz) and the "signal.filtfilt" function in SciPy
(**Figure 2: Supplementary Figure 1**). The a-wave amplitudes were measured from baseline-to-
trough, b-wave amplitudes were measured from a-wave trough-to-b-wave peak, and the b/a-wave
ratio was calculated to normalize responses. OP amplitude was measured from trough-to-peak
of the largest wave and the OP/b-wave ratio was calculated using b-wave amplitudes from the
same recording.

Patch clamp electrophysiology

Retinas were isolated and embedded in 3% low melting point agarose (Sigma; Cat#
A0701) in carbogenated bicarbonate-buffered Ames medium (Ames & Nesbett, 1981; Ames
Medium w/L-Glutamate; US Biological; Swampscott, MA, USA; Cat# A1372-25) and 200-250 μm
slices were prepared with a Leica VT1200 vibrating microtome. For chemically simulated light
response experiments, retinas were sliced in carbogenated Ames. For all other recordings, slicing
was performed in chilled and carbogenated low-Na$^+$, high-sucrose cutting solution containing (in
mM): 210 sucrose, 35 NaHCO$_3$, 20 HEPES, 10 glucose, 6 MgCl$_2$, 3 sodium pyruvate, 2.5 KCl and
0.5 CaCl$_2$ and buffered to pH 7.4 with NaOH. Retina slices were stored at room temperature in
carbogenated Ames medium before being transferred to the recording chamber. Heated (31-33°
C) Ames medium was continuously perfused over the retinal slices during all patch-clamp
recordings.

Thick-walled borosilicate glass pipettes (World Precision Instruments; Sarasota, Fl, USA;
Cat# 1B150F-4) were pulled using a Narishige PP-830 electrode puller (Tokyo, Japan). Ultrapure
salts were purchased from Sigma-Aldrich (St. Louis, MO, USA). For RBC visualization, the
fluorescent tracer Alexa Fluor 594 Hydrazide (100 μM; Thermo Fisher Scientific; Cat# A10438)
was added to the pipettes. Whole-cell voltage-clamp and current-clamp recordings were
performed in photopic conditions using a double EPC-10 patch clamp amplifier controlled by
Patchmaster software (HEKA Elektronik, Lambrecht/Pfalz, Germany) or an Axon Axopatch 200B
controlled by pCLAMP software (Molecular Devices; San Jose, CA, USA). Data was acquired at
a 20 kHz sampling rate and filtered with a 2.9 kHz low-pass Bessel filter. RBCs were targeted
visually using Dodt gradient contrast microscopy and confirmed post-hoc with cell-filling and
epifluorescence microscopy. Both pCLAMP and Patchmaster data sets were exported to Igor Pro
(WaveMetrics, Lake Oswego, OR, USA) for offline analysis with custom Igor scripts and statistical
analysis was performed in Prism. We used a P/10 leak subtraction protocol (-80 mV to -73 mV)
in voltage-clamp to isolate voltage-gated Ca^{2+} currents and GABAergic feedback currents; leak
traces for each trial were baselined, summated, and subtracted from the raw pulse trace.

To record chemically simulated light responses, pipettes were pulled to 10-12 MΩ and
backfilled with a potassium gluconate solution containing (in mM): 125 K-gluconate, 8 KCl, 5
HEPES, 1 MgCl$_2$, 1 CaCl$_2$, 3 adenosine 5'-triphosphate magnesium salt (Mg-ATP), 0.5 guanosine
5'-triphosphate disodium salt (Na$_2$-GTP), and 5 EGTA, and pH was adjusted to 7.3 with KOH. The
Ames medium was supplemented with 4 μM L-(+)-2-amino-4-phosphonobutyric acid (L-AP4), an
mGluR6 agonist, to maintain a simulated dark-adapted state. A puffer pipette (12-18 MΩ) was
maneuvered into the dendritic field of the target cell, and the cell was patched in voltage-clamp

and held at -60 mV. The mGluR6 antagonist [RS]-α-cyclopropyl-4-phophophenylglycine (CPPG)

was applied directly to the dendrites using a Picospritzer II (Parker Hannifin; Hollis, NH, USA; 600

μM at 20 psi) for five seconds and the RBC response was recorded. For each cell, four sweeps

were recorded at 15-second intervals.

For current-clamp experiments, pipettes were pulled to 9-12 MΩ and backfilled with a

potassium gluconate solution containing (in mM): 104 K-gluconate, 10 KCl, 10 HEPES, 4 Mg-

ATP, 0.4 Na_2-GTP, 10 Na_2-Phosphocreatine, and 2 EGTA, and pH was adjusted to 7.3 with KOH.

The Ames medium was supplemented with 1 mM $CaCl_2$ and the synaptic blockers (in μM): 0.5

strychnine (glycine receptor blocker), 3 gabazine (SR 95531; $GABA_A$ receptor blocker), 50

[1,2,5,6-tetrahydropyridin-4-yl]methylphosphinic acid (TPMPA; $GABA_{A\rho}$ receptor blocker), and 4

L-AP4 to block excitatory and inhibitory inputs. The total external $CaCl_2$ concentration was

estimated to be 2.15 mM. Cells were first patched in voltage-clamp and then the amplifier was

switched to fast current-clamp mode using the EPC-10 Gentle CC-Switch feature. Resting

membrane voltage was recorded at 0 pA and then current was injected for 100 ms from -80 pA

to 80 pA in 20 pA steps. Peak voltage was measured as the peak (min or max) voltage

immediately after the onset of the current pulse, and sustained voltage was measured as the

average over the final 10 ms of the pulse. For each cell, five sets of sweeps were recorded at

five-second intervals.

For the remaining experiments, the pipettes were pulled to 5-8 MΩ and backfilled with a

cesium gluconate solution containing (in mM): 94 Cs-gluconate, 10 HEPES, 20 tetraethyl

ammonium chloride (TEA-Cl), 4 Mg-ATP, 0.4 Na_2-GTP, 10 Na_2-Phosphocreatine, and 2 EGTA,

and pH was adjusted to 7.3 with CsOH.

For voltage-clamp time-resolved capacitance recordings, the Ames medium was

supplemented with 1 mM $CaCl_2$ and (in μM): 0.5 strychnine, 3 gabazine, 50 TPMPA, and 4 L-AP4

to isolate voltage-gated Ca^{2+} currents and maximize membrane resistance. Pipettes were

wrapped with parafilm to reduce pipette capacitance and electrical noise. Membrane capacitance

(C_m), membrane resistance (R_m), and series resistance (R_s) were measured using the "sine + DC"

method implemented in Patchmaster software (Balakrishnan, Puthussery, Kim, Taylor, &

von Gersdorff, 2015; Gillis, 2000; Oltedal & Hartveit, 2010). Exocytosis was evoked in voltage-

clamp mode with a 100 ms depolarizing pulse from -80 mV to -10 mV to activate voltage-gated

Ca^{2+} currents. Resting C_m was measured using a sinusoidal voltage command (200 Hz, 30 mV

trough-to-peak) superimposed over the membrane holding potential (-80 mV) and one

measurement was generated per cycle (**Figure 5A**). An increase in exocytosis (ΔC_m) was

measured from the change in C_m values before and after the depolarizing pulse. A baseline was
established by averaging the resting C_m over the one second immediately preceding the voltage
step. C_m measurements immediately after repolarization were excluded from analysis to allow for
the full decay of evoked conductances and C_m was averaged over the following 500 ms. Cells
were discarded if changes in C_m correlated significantly with changes in R_m or R_s. For each cell,
five sets of sweeps were recorded with 15 second intervals. To remove noise, C_m data was filtered
through a 5 Hz low-pass FIR filter. When different frequency sinusoidal voltage commands were
used, recordings were resampled to 200 Hz for ease of comparisons.

To record reciprocal inhibitory currents, the Ames medium was supplemented with 1 mM
$CaCl_2$, 4 μM L-AP4, and 0.5 μM strychnine. Feedback current was evoked in voltage-clamp mode
with a 100 ms depolarizing pulse from -80 mV to -10 mV to activate voltage-gated Ca^{2+} currents
and subsequent outward GABAergic inhibitory currents. We isolated inhibitory currents (I_{Inhib}) by
subtracting recordings with GABA blockers (I_{Ca}) were from recordings without GABA blockers
(I_{Total}), and calculated the charge transfer across the duration of the depolarizing pulse. Five sets
of sweeps were recorded at five-second intervals.

Results

TPBG is not required for normal gross RBC morphology

We used the constitutive Tpbg knockout mouse line B6;129S5-Tpbg[tm1Lex]/Mmucd (referred
to as TPBG-KO or KO) to analyze the consequences of genetic deletion of TPBG in the retina.
Wild type (WT) littermates were used as controls. Expression and localization of TPBG were
examined by western blotting and immunofluorescence confocal microscopy (**Figure 1**). As
previously reported (Wakeham et al., 2020), the TPBG antibody labels RBC cell bodies, dendrites,
and axon terminals in the WT retina, as well as a narrow band in the middle of the IPL
corresponding to the processes of TPBG-expressing amacrine cells. TPBG immunoreactivity was
absent throughout the KO retina (**Figure 1A**); this was confirmed by immunoblotting (**Figure 1B**)
of WT and KO retinal lysates with the TPBG antibody using β-actin as a loading control. Using
an antibody against PKCα in WT and TPBG-KO retina, we found no discernable disruption in
morphology of RBC dendrites or axon terminals in KO retinas (**Figure 1C-D**).

Figure 1: TPBG-KO does not alter gross RBC morphology. **(A)** Immunofluorescence confocal microscopy of WT (left) and TPBG-KO (KO; right) retina sections labeled with an antibody against TPBG (green). The nuclear stain DAPI (bottom; magenta) was used to highlight the neuronal layers of the retina. **(B)** Immunoblot of WT and TPBG-KO (KO) retinal lysates probed for TPBG and β-actin. The two channels were imaged separately and merged. **(C)** Confocal microscopy of WT (left) and TPBG-KO (KO; right) retina sections labeled with an antibody against PKCα. **(D)** Close-up images depicting the OPL (top) and IPL (bottom) for better visualization. For all images, Z-projections of 11 sections were used with a Z-step distance of 0.415 μm for a total thickness of 4.57 μm. Intensity and contrast was adjusted equally across genotypes. OPL: outer plexiform layer; IPL: inner plexiform layer. Scale bars: **(A and C)** 20 μm; **(D)** 10 μm.

Genetic knockout of TPBG alters the scotopic ERG light response

The light response of RBCs is reflected in the b-wave of the dark-adapted (scotopic) full-field electroretinogram (ERG), a commonly-used technique for recording the electrical responses of the whole retina *in vivo*. Different components of the ERG waveform correspond to the activity of specific cell types in the retina. The a-wave is an initial negative deflection corresponding to hyperpolarization of the photoreceptors (Robson & Frishman, 2014), whereas the b-wave, a larger and slower positive deflection, primarily corresponds to the subsequent depolarization of the ON-bipolar cells (Stockton & Slaughter, 1989). Rod driven activity can be isolated by dark-adapting the mouse before testing; therefore, the dark-adapted ERG b-wave represents the activity of RBCs in response to light stimuli.

Figure 2: TPBG-KO alters the scotopic ERG after bright stimuli. Scotopic electroretinograms (ERGs) were recorded from dark-adapted WT (blue; n = 12) and TPBG-KO (KO; orange; n = 14) mice after light stimuli (-0.06 to 1.93 log(cd•s/m^2)). **(A)** After baselining the raw, unfiltered ERG traces, a 60 Hz low-pass filter was used to remove OPs to better visualize the a- and b-waves. A 30-300 Hz Butteworth band-pass filter was used to isolate OPs from the ERG for quantification. **(B)** Average ERG recordings after light stimuli of 0.63 log(cd•s/m^2). A-wave amplitudes were measured from baseline to the a-wave trough and b-wave amplitudes were measured from the a-wave trough to the b-wave peak. **(C)** Quantification of b-wave amplitudes normalized to a-wave amplitudes. **(D)** Average oscillatory potentials (OPs) after light stimuli of 0.63 log(cd•s/m^2). **(E)** Quantification of OP amplitudes normalized to b-wave amplitudes. OP amplitudes were measured from the trough to peak of the largest wave. Quantified data is presented as the mean ± SEM. Statistical significance was determined using Welch's t-tests; ns: $p > 0.05$; *: $p < 0.05$; ** $p < 0.01$.

We recorded the scotopic ERG light responses in dark-adapted anesthetized WT (blue; n = 12) and TPBG-KO (KO; orange; n = 14) mice across several stimulus intensities, from dim (-0.06 log(cd•s/m^2)) to bright (1.93 log(cd•s/m^2)). We applied a 60 Hz low-pass filter to remove oscillatory potential (OP) contamination from the ERG a- and b-waves, and a 30-300 Hz Butterworth band-pass filter to isolate OPs for further analysis (**Figure 2A**). There was no difference in a-wave amplitudes across intensities (**Figure 2: Supplementary Figure**), indicating that the KO did not affect the rod photoreceptor light responses, but there was an enlarged b-wave in the TPBG-KO mouse (**Figure 2B**). Because the ERG b-wave is dependent on upstream photoreceptor activation, which is represented by the a-wave, it is common practice for ERG b-wave amplitudes to be normalized to a-wave amplitudes for comparison across animals. The TPBG-KO increased the normalized b-waves (b/a-wave ratio) across the range of stimuli intensities (**Figure 2C**), indicating a larger and more prolonged RBC light response in TPBG-KO mice.

Oscillatory potentials (OPs) are low amplitude, high frequency waves superimposed over the rising phase of the ERG b-wave and observed after bright stimuli. Previous studies indicate

that OPs originate from rhythmic activity in the IPL downstream of the bipolar cells (Ogden, 1973).

Because the light response of a dark-adapted retina is primarily rod-driven, the OPs are indirectly

correlated to RBC output. Therefore, we normalized the OP amplitude by the b-wave amplitude

(OP/b-wave ratio) of the corresponding ERG recording. Normalized TPBG-KO OPs were reduced

compared to WT (**Figure 2D and E**). TPBG is primarily expressed in RBCs, therefore an effect of

TPBG-KO on the OPs is most likely to be due to reduced RBC output onto downstream neurons,

though an effect mediated by the TPBG-expressing amacrine cells cannot be ruled out.

Figure 3: TPBG is not required for mGluR6-driven responses in RBC dendrites. (**A**) Immunofluorescence confocal
microscopy from WT and TPBG-KO whole mount retinas using antibodies against the rod presynaptic calcium channel
$Ca_v1.4$ (green) and the RBC post-synaptic protein GPR179 (magenta). Square boxes in the leftmost images indicate
regions of interest expanded in the other images. Z-projections of two sections were used with a Z-step distance of
0.297 μm for a total thickness of 0.594 μm. Scale bar is 5 μm. (**B**) Plot of fluorescence intensity over distance from
$Ca_v1.4$ (left) and GPR179 (right) channels from WT (blue; n = 66 synapses) and TPBG-KO (KO; orange; n = 54
synapses). (**C**) Quantification of synaptic distance. Box plot extends from the first to the third quartile and the whiskers
extend across all points. The line in the box represents the median. (**D-F**) Chemically simulated light responses were
recorded in WT (blue; n = 10) and TPBG-KO (KO; orange; n = 6) RBCs following a pulse of 600 μM CPPG for 5

seconds. **(D)** Example CPPG-induced TRPM1 currents. The gray bar represents the region across which charge
transfer was calculated. **(E)** Quantification of peak CPPG-induced TRPM1 current (I_{TRPM1}) and **(F)** charge transfer
(Q_{TRPM1}) over the five-second duration of the CPPG pulse. For **E** and **F**, open circles represent single cells and colored
bars represent the mean $\pm$ SEM. Statistical significance was determined using Welch's *t*-tests; ns: *p* > 0.05.
398

TPBG is not required for mGluR6-driven responses in RBC dendrites

We examined the expression and localization patterns of key pre- and post-synaptic

proteins that are required for synaptic transmission between rod photoreceptors and RBCs. RBCs

receive synaptic input from rod photoreceptors via a sign-inverting signal transduction cascade

localized to the tips of RBC dendrites. Glutamate released from rod photoreceptors is detected

by the group III metabotropic glutamate receptor mGluR6 (Masu et al., 1995; Nakajima et al.,

1993; Vardi, Duvoisin, Wu, & Sterling, 2000; Xu et al., 2012), that is negatively coupled to the

TRPM1 cation channel (Koike, Numata, Ueda, Mori, & Furukawa, 2010; Koike, Obara, et al.,

2010; Morgans et al., 2009; Shen et al., 2009). Thus, a light stimulus relieves TRPM1 tonic

inhibition, permitting cation influx and subsequent RBC depolarization.

First, we looked at the spatial relationship between presynaptic elements in the rod

spherule and the tips of the RBC dendrites (**Figure 3A**) using an antibody against the rod L-type

Ca^{2+} channel subunit $Ca_v1.4$ to label the horseshoe-shaped active zone and an antibody that

labels GPR179 in RBC dendritic tips (Hasan, Ray, & Gregg, 2016). In normal rod-RBC synapses,

GPR179 labeling appears as puncta within the concavity of the horseshoe-shaped $Ca_v1.4$

immunofluorescence. We quantified the spatial relationship between rod active zones and RBC

dendritic tips by measuring the distance between the peak intensity for each fluorescent channel

(**Figure 3B**). We found no difference between WT (blue; *n* = 66 synapses from eight animals) and

TPBG-KO (KO; orange; *n* = 54 synapses from seven animals) synaptic distance (**Figure 3C**; WT:

0.43 ± 0.006 μm; KO: 0.43 ± 0.004 μm; Welch's *t*-test: *p* = 0.96, *t*(117.60) = 0.052), suggesting

that RBC dendrites form morphologically normal invaginations into rod spherules in the TPBG-

KO mouse.

Scotopic ERG recordings revealed increased b-waves in TPBG-KO mice (**Figure 2**)

indicating an enhanced light response in KO RBCs. Because the RBC light response is initiated

by TRPM1, we compared TRPM1 currents between WT (blue; *n* = 10) and TPBG-KO (KO;

orange; *n* = 6) RBCs by recording chemically simulated light responses in retinal slices in whole-

cell voltage-clamp (**Figure 3D-F**). In this technique, bath application of the mGluR6 agonist L-

AP4 maintains the retina in a simulated dark-adapted state, and brief puffs of the mGluR6

antagonist, CPPG, onto the dendrites simulates a light flash and evokes TRPM1 currents (**Figure**

3D). TPBG-KO did not alter the CPPG-induced TRPM1 peak current (**Figure 3E**; WT: -88.24 $\pm$

8.92 pA; KO: -98.04 $\pm$ 11.28 pA; Welch's *t*-test: *p* = 0.51, *t*(10.85) = 0.68) or charge transfer

(**Figure 3F**; WT: -157.10 ± 47.01 pC; KO: -173.10 ± 38.62 pC; Welch's *t*-test: p = 0.47, $t(12.40)$ =

0.74). These results suggest that TPBG is not required for the formation of functional rod-RBC

synapses and that the larger scotopic ERG b-waves do not reflect enhanced TRPM1 currents in

the TPBG-KO.

Synaptic ribbons are smaller in TPBG-KO RBCs compared to WT

Oscillatory potentials are thought to be generated by the inner retina, and RBCs drive the

majority of downstream activity under scotopic conditions; thus, the attenuation of scotopic OPs

in the TPBG-KO suggests a deficit in RBC output to third-order neurons. Like rods, RBCs signal

to downstream cells using ribbon synapses. A depolarized potential entering the axon terminal

prompts the opening of L-type voltage-gated Ca^{2+} channels clustered beneath the ribbons and

the resulting influx of Ca^{2+} triggers the fusion (exocytosis) of ribbon-tethered synaptic vesicles

(Gersdorff, Vardi, Matthews, & Sterling, 1996). This process requires a close spatial relationship

between the Ca^{2+} channels the synaptic ribbons (Jarsky, Tian, & Singer, 2010). We used triple

immunofluorescence microscopy to examine the relative localization and expression patterns of

the L-type Ca^{2+} channel subunit, $Ca_v1.4$, and the primary structural synaptic ribbon protein

RIBEYE (**Figure 4A-C**). Because neither $Ca_v1.4$ nor RIBEYE are specific to RBC axon terminals,

we isolated RBC-specific immunofluorescence using a binary mask derived from PKCα

immunofluorescence to exclude labeling from other bipolar cells in the IPL. $Ca_v1.4$ and RIBEYE

show punctate immunofluorescence in RBC axon terminals and are co-localized or closely

apposed (**Figure 4A**). Mander's coefficients (M1 and M2) were calculated to examine

colocalization between $Ca_v1.4$ and RIBEYE puncta in both control (PKCα-negative) terminals and

RBC (PKCα-positive) terminals (**Figure 4B**). We found no differences in co-localization

coefficients between WT and TPBG-KO control or RBC axon terminals: WT (blue; n = 134 control

puncta and 100 RBC puncta) and TPBG-KO (orange; n = 327 control puncta and 119 RBC

puncta), control M1 (WT: 0.53 ± 0.022 arbitrary units (AU); KO: 0.52 ± 0.013 AU; Welch's *t*-test:

p = 0.60, $t(237.2)$ = 0.53), control M2 (WT: 0.40 ± 0.015 AU; KO: 0.42 ± 0.011 AU; Welch's *t*-test:

p = 0.28, $t(288.1)$ = 1.09), RBC M1 (WT: 0.66 ± 0.016 AU; KO: 0.64 ± 0.015 AU; Welch's *t*-test: p

= 0.47, $t(208.3)$ = 0.73), RBC M2 (WT: 0.32 ± 0.01 AU; KO: 0.31 ± 0.01 AU; Welch's *t*-test: p =

0.76, $t(216.9)$ = 0.31). Additionally, we found that the number of ribeye puncta per unit area was

unchanged in WT (blue; n = 90 axon terminals) and TPBG-KO (KO; orange; n = 109 axon

terminals) RBC terminals (**Figure 4C**; WT: 2.07 ± 0.66 puncta/μm^2; KO: 2.03 ± 0.58 puncta/μm^2;

Welch's *t*-test: p = 0.70, $t(177.8)$ = 3.92). We also compared synaptotagmin I/II

immunofluorescence between RBC axon terminals in WT and TPBG-KO retinas and saw no
difference (not shown; WT: 1.352 ± 0.0486 AU; KO: 1.446 ± 0.0560; Welch's t-test: p-value =
0.209, t(52.63) = 1.271).

467
Figure 4: RBC synaptic ribbons are smaller in the TPBG-KO retina. (**A**) Triple immunofluorescence confocal
microscopy of WT and TPBG-KO (KO) whole mount retinas using antibodies against PKCα (blue), Ca$_v$1.4 (Green) and
RIBEYE (magenta). To isolate labeling in RBC axon terminals, binary images were created using the PKCα channel
and overlayed over the other two channels to generate binary masks. Z-projections of two sections were used with a
Z-step distance of 0.297 µm for a total thickness of 0.547 µm. The scale bar is 5 µm. (**B**) Quantification of Ca$_v$1.4 and
RIBEYE colocalization in WT (blue) and TPBG-KO (KO; orange) using M1 and M2 Mander's coefficients in control
(WT: *n* = 134 puncta; KO: *n* = 327 puncta) and RBC axon terminals (WT: *n* = 100 puncta; KO: *n* = 119 puncta). (**C**)

To examine synaptic ribbons in RBC axon terminals, serial electron micrographs were

collected from a volume each of WT and TPBG-KO retina using a focused ion beam scanning

electron microscope (FIB-SEM). Bipolar cell terminals closest to the ganglion cell layer were

identified as putative RBC axon terminals. The general ultrastructure of the RBC axon terminals,

including the synaptic vesicles, synaptic ribbons, and postsynaptic elements was similar in WT

and TPBG-KO retina. To estimate synaptic ribbon size, we measured the maximum distance that

each ribbon extended into the cytoplasm from its attachment site on the plasma membrane in WT

(blue; *n* = 33 ribbons) and TPBG-KO (KO; orange; *n* = 38 ribbons) and found that synaptic ribbons

were 15% smaller in TPBG-KO RBCs compared to WT RBCs (**Figure 4E**; WT: 416.8 ± 13.78 nm;

KO: 355.7 ± 8.85 nm; Welch's *t*-test: *p* = 0.0004, $t(55.65)$ = 3.74).

TPBG is required for efficient exocytosis in RBC synaptic terminals.

To examine the effect of TPBG-KO on RBC synaptic output, we recorded time-resolved

membrane capacitance (C_m) before and after a depolarizing pulse using the "sine + DC" method

(Gillis, 2000; Oltedal & Hartveit, 2010). High-resolution membrane capacitance changes (ΔC_m)

after a depolarizing pulse strongly correlate with vesicle exocytosis in neuroendocrine cells

(Klyachko & Jackson, 2002; Neher & Marty, 1982) and glutamate release from bipolar cell axon

terminals (Kim & von Gersdorff, 2016). A sinusoidal voltage command (200 Hz at 30 mV trough-

to-peak) was superimposed over the holding voltage of -80 mV and exocytosis was evoked by

depolarizing the cell to -10 mV to open the voltage-gated Ca^{2+} channels (**Figure 5:**

Supplementary Figure A). Membrane capacitance (C_m), membrane resistance (R_m), and series

resistance (R_s) were recorded before and after the pulse and ΔC_m was measured as the change

in C_m after exocytosis. Changes in R_m and R_s were not correlated with changes in C_m (**Figure 5:**

Supplementary Figure B).

C_m measurements in RBCs are complicated by the attenuation and filtering of the

sinusoidal voltage command amplitude down their narrow axons (Oltedal & Hartveit, 2010;

Oltedal, Veruki, & Hartveit, 2009). We sought to confirm that a low-frequency sinusoidal command

reduces attenuation and maximizes RBC C_m changes compared to higher frequencies (**Figure 5:**

Supplementary Figure C-F) using 200 (n = 6), 400 (n = 4), 800 (n = 4), and 1600 (n = 4) Hz

commands. We found no difference in the voltage-gated Ca^{2+} current amplitude (I_{Ca}) or charge

transfer (Q_{Ca}) across frequencies (not shown), but the ΔC_m (**Figure 5: Supplementary Figure C-**

D; 200 Hz: 41.94 ± 4.95 fF; 400 Hz: 19.13 ± 4.06 fF; 800 Hz: 4.05 ± 2.91 fF; 1600 Hz: -4.23 ±

1.40; Brown-Forsythe ANOVA: p < 0.0001, 34.12(2.00, 10.87)) and the resting C_m (**Figure 5:**

Supplementary Figure E-F; 200 Hz: 5.97 ± 0.23 pF, 400 Hz: 5.73 ± 0.12 pF, 800 Hz: 4.81 ± 0.15

pF, 1600 Hz: 4.04 ± 0.12 pF; Brown-Forsythe ANOVA: p < 0.0001, 25.06(3.00, 11.34)) were

greater at 200 Hz compared to higher frequencies. These supplementary results support our

decision to examine exocytosis in RBCs using a 200 Hz "sine + DC" protocol.

To examine RBC synaptic vesicle exocytosis, we used whole-cell voltage-clamp to

compare WT (blue; n = 6) and TPBG-KO (KO; orange; n = 7) voltage-gated Ca^{2+} currents and

subsequent changes in membrane capacitance (**Figure 5**). Cells where C_m, R_m, and R_s were

highly correlated were discarded. The synaptic blockers L-AP4, strychnine, TPMPA, and

gabazine were used to block spontaneous activity and help increase and stabilize R_m during

recordings. The passive properties of R_m (WT: 1.603 ± 0.24 GΩ; KO: 1.630 ± 0.16 GΩ; Welch's

t-test: p = 0.93, *t*(8.84) = 0.097) and R_s (WT: 50.54 ± 2.33 MΩ; KO: 49.61 ± 1.91 MΩ; Welch's *t*-

test: p = 0.76, *t*(10.16) = 0.31) were not different between WT and TPBG-KO RBCs. As long as

R_m is significantly greater than R_s, ΔC_m should be a faithful approximation of overall exocytosis

(Gillis, 2000).

Figure 5: TPBG is required for efficient synaptic vesicle release in RBC axon terminals. Whole-cell voltage-clamp
recordings were performed in WT (blue; *n* = 6) and TPBG-KO (KO; orange; *n* = 7) RBCs. **(A)** Epifluorescence
micrograph of an RBC filled with Alexa Fluor 594 Hydrazide after patch-clamp recording. RBCs were patched and cell
type was verified *post-hoc* using epifluorescence microscopy. The scale bar is 10 μm. **(B)** Average voltage-gated Ca^{2+}
current traces recorded after a depolarizing pulse from -80 mV to -10 mV for 100 ms. **(C)** Quantification of peak voltage-
gated Ca^{2+} current amplitudes (I_{Ca}) and **(D)** charge transfer (Q_{Ca}) over the duration of the pulse. **(E)** Average membrane
capacitance (C_m) recordings using a 200 Hz sine + DC protocol. ΔC_m was measured as the difference between
baselined resting C_m before and after the depolarizing pulse. Data immediately after the pulse was excluded until C_m

stabilized. The gray bar represents the region across which C_m values were averaged. **(F)** Quantification of ΔC_m. **(G)**
Quantification of exocytosis efficiency calculated as the absolute value of the change in C_m per unit Q_{Ca}. Ames medium
was supplemented with 1 mM $CaCl_2$ and (in µM): 0.5 strychnine, 3 gabazine, 50 TPMPA, and 4 L-AP4. Open circles in
bar graphs represent single cells and bars represent the mean ± SEM. Statistical significance was determined using
Welch's t-tests; ns: $p > 0.05$; *: $p < 0.05$; **: $p < 0.01$.

TPBG-KO had no effect on the voltage-gated Ca^{2+} currents (**Figure 5B-C**; WT: -74.99 ±

4.49 pA, KO: -74.24 ± 3.41 pA; Welch's t-test: $p = 0.90$, $t(9.73) = 0.13$) or charge transfer (**Figure**

5D; WT: -7.0 ± 0.38 pC, KO: -6.7 ± 0.33 pC; Welch's t-test: $p = 0.57$, $t(10.43) = 0.58$). However,

TPBG-KO resulted in a reduction in ΔC_m compared to WT (**Figure 5E-F**; WT: 41.94 ± 4.95 fF,

KO: 25.90 ± 3.91 fF; Welch's t-test: $p = 0.029$, $t(9.95) = 2.55$), suggesting a decrease in synaptic

vesicle exocytosis in the KO. Most interestingly, the efficiency of exocytosis (the ratio of ΔC_m per

unit Q_{Ca}) is also reduced in the KO (**Figure 5G**; WT: 5.93 ± 0.50 fF/-pC, KO: 3.76 ± 0.44 fF/-pC;

Welch's t-test: $p = 0.0077$, $t(10.52) = 3.28$), indicating that TPBG-KO alters the efficiency of RBC

vesicle exocytosis without changing Ca^{2+} influx.

551
Figure 6: TPBG does not alter RBC input resistance or excitability*. Whole-cell current-clamp recordings in WT
(blue; $n = 9$) and TPBG-KO (KO; orange; $n = 11$) RBCs were used to construct a current-voltage relationship and
calculate input resistance (R_{in}). A current injection protocol from -80 pA to 80 pA in 20 pA steps was used with a K-
gluconate-based internal solution. **(A)** Average voltage traces following -80 pA, 0 pA, and 80 pA current injections for
100 ms. **(B)** Resting membrane voltages recorded immediately after entering current clamp mode. **(C)** A plot of the
voltage-current (V-I) relationship measured at the peak voltage during current injections. **(D)** Quantification of peak R_{in}.

(E) A plot of the sustained V-I relationship measured as the average voltage calculated over the last 10 ms of the
current pulse. **(F)** Quantification of the sustained R_{in}. Peak and sustained R_{in} values were calculated by measuring the
slope of the linear portion of the V-I curves (from -40 to 40 pA). Ames medium was supplemented with 1 mM $CaCl_2$
and (in µM): 0.5 strychnine, 3 gabazine, 50 TPMPA, and 4 L-AP4. For V-I curves, open circles represent the mean and
error bars represent SEM. For quantification, open circles represent single cells and bars represent the mean ± SEM.
Statistical significance was determined using Welch's *t*-tests; ns: *p* > 0.05.

TPBG does not alter RBC input resistance or excitability

TPBG-KO amplifies the RBC light response and reduces synaptic vesicle exocytosis,

suggesting that TPBG-KO may reduce the gain of RBC synaptic release by shifting RBC

excitability. To compare the excitability between WT (blue; *n* = 9) and TPBG-KO (KO; orange; *n*

= 11) RBCs, we used current injections (from -80 to 80 pA in 20 pA intervals) in whole-cell current-

clamp mode to construct voltage-current relationships and calculate the peak and sustained input

resistances (R_{in}). For these experiments, we used the synaptic blockers L-AP4, strychnine,

TPMPA, and gabazine to isolate RBC-specific voltage responses. **Figure 6A** shows voltage

responses after representative current injections of -80 pA, 0 pA, and +80 pA. The resting

membrane potential (V_m) was unchanged in the KO (**Figure 6B**; WT: -58.02 ± 4.91 mV, KO: -

56.35 ± 4.66 mV; Welch's *t*-test: *p* = 0.81, *t*(17.53) = 0.25). We found no difference in peak (**Figure**

6C-D; WT: 1.40 ± 0.13 GΩ, KO: 1.39 ± 0.09 GΩ; Welch's *t*-test: *p* = 0.97, *t*(14.94) = 0.04) or

sustained R_{in} (**Figure 6E-F**; WT: 1.13 ± 0.11 GΩ, KO: 1.05 ± 0.07 GΩ; Welch's *t*-test: *p* = 0.58,

t(14.28) = 0.56) between the WT and KO RBCs. These results indicate that a shift in R_{in} and RBC

excitability in the TPBG-KO do not explain the reduced exocytosis.

TPBG-KO reduces reciprocal inhibitory feedback

RBCs drive retinal output via glutamate release into the IPL, which stimulates Ca^{2+}-

permeable AMPA receptors in downstream AII amacrine cells (AII-ACs). RBC glutamate release

is also detected by A17 amacrine cells (A17-ACs), which when depolarized, release GABA back

onto RBC synaptic terminals where it is sensed by $GABA_A$ and $GABA_{A}\rho$ (also called $GABA_C$) Cl^-

channels (**Figure 7A**). An influx of Cl^- into the axon terminal through open GABA channels

provides inhibitory feedback by hyperpolarizing the RBC and shaping further RBC glutamate

release (Chávez, Grimes, & Diamond, 2010; Chávez, Singer, & Diamond, 2006; Euler & Masland,

2000; E. Hartveit, 1996; Espen Hartveit, 1999; Singer & Diamond, 2003). Each RBC active zone

forms a dyad synapse with one AII-AC and one A17-AC (Famiglietti & Kolb, 1975; Grimes, Zhang,

Graydon, Kachar, & Diamond, 2010; McGuire, Stevens, & Sterling, 1984; Raviola & Dacheux,

1987). As A17-mediated reciprocal feedback is driven by RBC glutamate release, the inhibitory

feedback currents in the RBCs are an indirect measure of glutamate release from RBCs and
would be expected to be reduced concomitantly with reduced RBC exocytosis in the TPBG-KO.

595
Figure 7: TPBG-KO reduces reciprocal inhibitory feedback. **(A)** The dyad synapse arrangement between one RBC,
one AII-AC, and one A17-AC in the IPL. Glutamate release by RBCs is detected by Ca^{2+}-permeable AMPA receptors
(AMPARs) in AII-AC and A17-AC dendrites. A17-ACs then release GABA back onto the synaptic cleft which is sensed
by $GABA_A$ receptors ($GABA_A$Rs) and $GABA_{A\rho}$ receptors ($GABA_{A\rho}$Rs) in the RBC axon terminals. **(B)** Inhibitory currents
are superimposed over voltage-gated Ca^{2+} currents. Reciprocal feedback in RBCs was evoked with a 100 ms
depolarizing pulse from -80 mV to -10 mV. In the absence of GABA receptor blockers, this depolarizing pulse elicits an
inward voltage-gated Ca^{2+} current that is quickly subsumed by outward inhibitory currents (I_{Total}; green). These outward
currents are abolished by the application of the GABA receptor blockers gabazine and TPMPA (I_{Ca}; magenta). Inhibitory
currents (I_{Inhib}; magenta) can be isolated by subtracting I_{Ca} (maroon) from I_{Total} (green). **(C)** Average whole-cell patch-
clamp recordings from WT (blue; n = 6) and TPBG-KO (KO; orange; n = 5) RBCs to compare reciprocal inhibitory
feedback ($I_{Inhib} = I_{Total} - I_{Ca}$) after a depolarizing pulse from -80 mV to -10 mV for 100 ms. The gray bar represents the
region across which the charge transfer was calculated. **(D)** Quantification of the charge transfer (Q_{Inhib}) measured from
baseline over the duration of the 100 ms stimulus. Ames medium was supplemented with 1 mM $CaCl_2$, 0.5 µM
strychnine and 4 µM L-AP4. Open circles in bar graphs represent single cells and bars represent the mean ± SEM.
Statistical significance was determined using Welch's t-tests; *: p < 0.05.
611

To test the hypothesis that reduced RBC glutamate release results in smaller inhibitory
feedback currents in the TPBG-KO, we used whole-cell voltage-clamp to examine the reciprocal
inhibitory response to a depolarizing pulse from -80 mV to -10 mV for 100 ms in WT (blue; n = 6)
and TPBG-KO (KO; orange; n = 5) RBCs. L-AP4 was used to block spontaneous excitatory inputs
and strychnine was used to block glycinergic inhibition. In the absence of GABA receptor blockers,
outward currents can be seen superimposed on the evoked voltage-gated Ca^{2+} current (**Figure
7B; Green;** I_{Total}). Inhibitory currents (magenta; I_{Inhib}) can be isolated by subtracting voltage-gated
Ca^{2+} currents (maroon; I_{Ca})from the total current. The charge transfer across the duration of the
stimulus was reduced in the TPBG-KO compared to WT (**Figure 7D**; WT: 2.35 ± 0.37 pC; KO:
1.13 ± 0.22 pC; Welch's t-test: p = 0.022, t(7.83) = 2.83). This supports our previous result that
the TPBG-KO suppresses RBC vesicle exocytosis and glutamate release (see **Figure 5F**).

Figure 8: TPBG is not required for localization of GABA receptor immunofluorescence to RBC axon terminals.
Immunofluorescence confocal microscopy of WT and TPBG-KO (KO) whole mount retinas using an antibody against
PKCα (magenta) and antibodies against either the GABA$_A$α1 subunit (**A**; green; WT: n = 18 terminals; KO n = 28
terminals) or the GABA$_A$ρ1 subunit (**C**; green; WT: n = 20 terminals; KO n = 27 terminals). To isolate labeling in RBC
axon terminals, binaries were created using the PKCα channel and overlayed over the other channels to generate
binary masks. For all images, Z-projections of two sections were used with a Z-step distance of 0.297 μm for a total
thickness of 0.547 μm. Scale bar is 5 μm. **(B)** Quantification of GABA$_A$α1 subunit puncta/μm^2. **(D)** Quantification of
GABA$_A$α1 subunit puncta/μm^2. Box plots extend from the first to the third quartile and the whiskers extend across all
points. The line in the box represents the median. Statistical significance was determined using Welch's t-tests; ns: p
> 0.05.

It is possible that reciprocal feedback is altered in TPBG-KO due to disrupted expression

or clustering of GABA receptors in RBC axon terminals. RBCs express both GABA$_A$α1 and

GABA$_A$ρ1 receptor subunits in their axon terminals (Fletcher, Koulen, & Wässle, 1998; Palmer,

2006). To see whether TPBG-KO alters GABA receptor expression patterns, we used double

immunofluorescence labeling of WT and TPBG-KO whole mount retinas using an antibody

against PKCα and antibodies against GABA$_A$α1 (WT: n = 18 terminals from three animals; KO n

= 28 terminals from four animals) and GABA$_A$ρ1 (WT: n = 20 terminals from three animals; KO n

= 27 terminals from three animals) receptor subunits (**Figure 8**). As previously, we isolated RBC

axon terminal-specific labeling using a binary mask derived from PKCα immunofluorescence.

Next, we counted the GABA puncta per unit area in PKCα-positive axon terminals. We did not

observe any major differences in puncta/μm^2 between WT and TPBG-KO (KO) retinas labeled

against either GABA$_A\alpha$1 (**Figure 8A-B**; WT: 1.21 $\pm$ 0.05 puncta/μm^2; KO: 1.40 $\pm$ 0.09 puncta/μm^2;

Welch's *t*-test: *p* = 0.064, *t*(39.73) = 1.91) or GABA$_A\rho$1 (**Figure 8C-D**; WT: 1.03 $\pm$ 0.03 puncta/μm^2;

KO: 0.98 $\pm$ 0.03 puncta/μm^2; Welch's *t*-test: *p* = 0.20, *t*(41.11) = 1.29), indicating that impaired

GABA receptor expression is likely not reducing inhibitory feedback. Taken together, our data

supports the conclusion that TPBG is required for normal synaptic ribbon development and

efficient glutamate release from RBC axon terminals.

Discussion

In this study, we characterized the functional consequences of genetic knockout of TPBG

in the mouse retina using immunofluorescence and electrophysiological approaches. TPBG-KO

did not alter gross RBC morphology in the OPL or IPL as PKCα-labeled RBC dendrites and axon

terminals appear normal in both conditions. Because TPBG is localized to both the dendrites and

axon terminals of RBCs (Wakeham et al., 2020), we sought to examine the effect of TPBG-KO

on the RBC synaptic structure and physiology in both compartments. Though ERG b-waves were

larger in the TPBG-KO, dendritic TRPM1 currents activated by CPPG did not change. In contrast

to the larger b-waves, the scotopic OPs were smaller in the KO, suggesting changes in RBC

ribbon synapse function. Accordingly, time-resolved C$_m$ measurements and whole-cell recording

indicated that synaptic vesicle exocytosis and GABAergic feedback are reduced at TPBG-KO

RBC synapses, and synaptic ribbons are shorter. Importantly, we did not detect differences in

presynaptic voltage-gated Ca^{2+} current amplitude or changes in Ca^{2+} channel, GABA receptor, or

RIBEYE expression and localization. We propose that TPBG plays an important role in fine tuning

the synaptic architecture of RBC ribbon synapses to increase exocytosis efficiency.

TPBG and RBC dendrites

We found that TPBG is not required for the normal insertion of RBC dendritic tips inside

rod spherules or for normal depolarizing currents through TRPM1 cation channels. Our

immunofluorescence and electrophysiological examination of RBC dendrites suggest that TPBG

is not essential for basic synaptic function in that compartment. That being said, TPBG

immunofluorescence reveals that large amounts of TPBG are present in RBC dendrites,

suggesting some role in dendritic development or physiology.

TPBG regulates dendritic arborization in mouse olfactory bulb granule cells (Yoshihara et

al., 2012). In these neurons, TPBG-KO reduced dendritic branching and dendritic migration to

synaptic targets. Lentiviral overexpression of TPBG increased dendritic branching complexity and
restored dendritic arborization deficits induced by nasal occlusion. In the retina, TPBG may also
regulate RBC dendritic branching complexity in the OPL. TPBG is a known regulator of Wnt
signaling (He et al., 2015; Kagermeier-Schenk et al., 2011), which is required for functional
synaptic targeting and OPL lamination in RBCs, therefore TPBG may promote RBC dendritic
development through control of the Wnt pathway (Tsuboi, 2020). We didn't detect a structural or
physiological phenotype in RBC dendrites, but thorough evaluation of RBC dendritic maturation
and branching using sparse labeling techniques might reveal a more subtle effect of TPBG-KO
on dendritic architecture.
687
TPBG and RBC synaptic vesicle exocytosis
We found a reduction in the OPs isolated from scotopic ERG recordings in TPBG-KO
mice, which we interpret as evidence for decreased RBC output to third-order neurons. While we
cannot rule out a contribution of the TPBG-expressing amacrine cells to the reduced OP
amplitudes in the TPBG-KO, the similarity of the OP phenotype to that seen in the PKCα-KO
(Xiong et al., 2015) suggests that it is due to an RBC defect. Furthermore, pharmacological
dissection of ERG components points to the RBC to AII/A17 reciprocal synapse as the origin of
the OPs (Liao, Liu, Milla-Navarro, Villa, & Germain, 2023; Wachtmeister, 1998). Early studies
implicated interactions between bipolar cell axon terminals, amacrine cell processes, and
ganglion cell dendrites as potential generators of the OPs (Ogden, 1973). Intravitreal injections of
glycine into rabbit retinas resulted in morphological changes in many ACs and disappearance of
the OPs from the ERG (Korol, Leuenberger, Englert, & Babel, 1975) while application of
neurotoxic kainate to ganglion cell dendrites diminishes OP amplitude (Vaegan & Millar, 1994). A
more recent study unequivocally implicates multiple distinct generators in the production of early,
intermediate, and late OP peaks (Dong, Agey, & Hare, 2004). The use of APB and CNQX to block
synaptic inputs to ON and OFF-bipolar cells significantly diminished intermediate and later OPs
but left early OPs intact, suggesting that early OPs are generated by photoreceptors. Intermediate
and late OPs have differential sensitivity to tetrodotoxin, which abolished only late OPs, indicating
that intermediate OPs are generated by action-potential-independent interactions and late OPs
are generated by action-potential-dependent ones (Dong et al., 2004). Our OP data shows a
much larger attenuation of TPBG-KO OP amplitude in intermediate and late OPs compared to
early ones. Thus, the results of these studies support our conclusion that reduced scotopic OPs
are likely due to suppressed rhythmic activity of third-order neurons due to reduced RBC
glutamate release.

712    By simultaneously recording the presynaptic voltage-gated Ca^{2+} currents and subsequent

713 membrane capacitance changes evoked by RBC depolarization, we showed that reduced

714 exocytosis in TPBG-KO RBCs compared to WT, despite no change in presynaptic Ca^{2+} currents,

715 indicating reduced exocytosis efficiency in the TPBG-KO. The Ca^{2+} current amplitudes we

716 recorded (-75 pA) following a depolarizing step are significantly larger than those previously

717 reported (-30 pA by (Protti & Llano, 1998) and -20 pA by (Singer & Diamond, 2003)). We also

718 recorded larger C_m jumps (42 fF in WT) than those reported from dissociated RBCs (30 fF in

719 (Zhou, Wan, Thakur, & Heidelberger, 2006)). We believe these differences can be attributed to

720 higher recording temperatures (33° C vs 21° C) and our use of older animals. With a mean resting

721 C_m of 5.97 pF and a mean R_s of 50.04 MΩ, we can calculate an average voltage-clamp speed of

722 298.74 μs – sufficiently fast to clamp the membrane to a 200 Hz sine wave with a 5 ms period.

723 The reduced exocytosis in the TPBG-KO, despite the same Ca^{2+} currents as in WT, suggests that

724 calcium influx is not as tightly coupled to the synaptic vesicle exocytosis in the TPBG-KO, resulting

725 in reduced exocytosis efficiency.

726    Exocytosis efficiency is calculated as the exocytosis per unit Ca^{2+} charge transfer. We

727 calculated an exocytosis efficiency of approximately 6 fF/pC in adult WT mouse RBCs after a 100

728 ms depolarizing pulse at 33° C. In adult vertebrate auditory hair cells, which also form ribbon

729 synapses, similar exocytosis efficiencies of 5 to 6 fF/pC have been reported after 100 ms pulses

730 (Chen & von Gersdorff, 2019; Moser et al., 2020). Rod bipolar cell ribbon synapses operate via

731 nanodomain coupling between Ca^{2+} channels and docked vesicles (Jarsky et al., 2010), whereas

732 immature calyx of Held synapses contain conventional active zones that operate via Ca^{2+}

733 microdomains where Ca^{2+} influx triggers vesicle exocytosis from greater distances (Kushmerick,

734 Renden, & von Gersdorff, 2006). Nevertheless, exocytosis efficiency can still be as high as 16 to

735 20 fF/pC at immature calyx of Held axon terminals after a short 1 ms pulse. Further studies will

736 be required to verify the relationship between TPBG and nanodomain coupling at RBC ribbon

737 synapses.

739 *TPBG and synaptic ribbon proteins*

740    Neurotransmitter release from synaptic ribbons can be simplified into three major steps:

741 synaptic docking, Ca^{2+} influx and its detection by Ca^{2+} sensors, and Ca^{2+}-dependent fusion of

742 synaptic vesicles with the presynaptic membrane (Dieck & Brandstätter, 2006; Morgans, 2000).

743 These are all potential points at which TPBG may be supporting efficient vesicle release from

744 RBCs. Vesicle accumulation and transport in RBC synaptic terminals is dependent on F-actin

745 polymerization and is regulated by PKCα (Berglund, Midorikawa, & Tachibana, 2002; Doussau &

Augustine, 2000; Job & Lagnado, 1998; Minami, Berglund, Sakaba, Kohmoto, & Tachibana, 1998;

Tachibana, 1999). We originally identified TPBG as a PKCα-dependent phosphoprotein

(Wakeham et al., 2019); thus, it is possible that PKCα controls vesicle docking via actin dynamics

in RBC axon terminals through TPBG.

The tight temporal control of vesicle release that is a trademark of synaptic ribbons

requires nanodomain-coupling between L-type voltage-gated Ca^{2+} channels and Ca^{2+} binding

proteins that trigger synaptic vesicle fusion (Balakrishnan et al., 2015; Eggermann, Bucurenciu,

Goswami, & Jonas, 2012; Jarsky et al., 2010). Synaptic ribbons are primarily composed of the

scaffold protein RIBEYE and are anchored to the presynaptic active zone by a number of

cytoskeletal anchoring proteins, including Bassoon and Piccolo (Zanazzi & Matthews, 2009). We

did not observe a difference in RIBEYE immunofluorescence in WT and TPBG-KO axon

terminals, but immunofluorescence does not have the resolution to confirm the correct localization

or orientation of the ribbon at the active zone, If TPBG is required for correct localization of the

ribbon, exocytosis efficiency may be significantly disrupted in KO RBCs, but electron microscopy

showed normal positioning of ribbons at RBC active zones in the KO. $Ca_v1.4$ and RIBEYE appear

co-localized by immunofluorescence in both WT and TPBG-KO RBC axon terminals, but a small

disruption of this tight coupling might not be detected with immunofluorescence and could have a

large effect on glutamate release by slowing the accumulation of Ca^{2+} at the sensor. Nanodomain

coupling likely depends on a close spatial relationship between the L-type channels and the

vesicle release machinery, yet it is not fully known how this clustering is achieved in RBCs. It is

possible that TPBG facilitates Ca^{2+} channel clustering or association between the channels and

the ribbon such that, in its absence, normal nanodomain coupling is disrupted and synaptic vesicle

fusion is reduced.

Ultrastructural analysis of RBC axon terminals revealed that the synaptic ribbons are

approximately 15% smaller in TPBG-KO RBCs compared to WT. Shorter ribbons could result in

a smaller number of ribbon-attached vesicles that may constitute a smaller ready releasable pool

(RRP), resulting in the observed C_m jump attenuation. The density of ribbons was not reduced in

the TPBG-KO as assayed by immunofluorescence imaging, nor were Ca^{2+} currents changed;

thus, the smaller C_m jumps not could be due to a smaller number of ribbons or reduced Ca^{2+} influx.

We used a large 100 ms depolarizing pulse to evoke vesicle release which likely exhausts the

fast RRP (docked vesicles) and potentially depletes the secondary RRP (vesicles attached to the

ribbon in upper rows). The fast RRP probably consists of the first row of vesicles lining the bottom

of the ribbon closest to the active zone (von Gersdorff & Matthews, 1997; von Gersdorff et al.,

1996) and this pool may not be affected by a shorter ribbon. In contrast, a shorter ribbon may

significantly reduce the size of the secondary RRP, presumably corresponding to vesicles
tethered higher on the ribbon, leading to reduced C_m jumps. Indeed, in RIBEYE-KO mice, where
the synaptic ribbon is absent, exocytosis is greatly reduced in RBC and rod photoreceptor
synapses (Grabner & Moser, 2021; Maxeiner, Luo, Tan, Schmitz, & Südhof, 2016). More
experimentation is required to determine exactly how the shorter ribbons in the TPBG-KO RBC
axon terminals affects the different vesicle pools.

TPBG and reciprocal feedback inhibition

To confirm that the TPBG-KO reduces RBC vesicle fusion and glutamate release, we
compared GABAergic feedback inhibition between WT and TPBG-KO RBCs. We found
attenuation of inhibitory charge transfer in the TPBG-KO. With glycinergic inhibition blocked by
strychnine, the primary remaining source of feedback inhibition in RBCs is A17-mediated
GABAergic reciprocal feedback. GABA released from A17-ACs is sensed by two kinetically
distinct ionotropic GABA receptors in RBC axon terminals: $GABA_A$ and $GABA_{A\rho}$ (Fletcher et al.,
1998). $GABA_A$ receptor-mediated current is rapidly-activating, while $GABA_{A\rho}$-mediated current is
slowly-activating (Grimes et al., 2015). We detected two different inhibitory components: a large,
quickly-inactivating current and a smaller sustained one. Though we did not pharmacologically
isolate each current, the kinetics of the inhibitory currents suggest that peak I_{Inhib} is likely mediated
by $GABA_A$ receptors and sustained I_{Inhib} is likely mediated by $GABA_{A\rho}$ receptors. We note that our
peak WT $GABA_A$ receptor currents of ~125 pA and sustained $GABA_{A\rho}$ receptor currents of ~50
pA are considerably larger than reported previously in mouse RBCs (Chávez et al., 2010). Again,
we attribute this to our use of older animals and higher recording temperatures and, perhaps, the
use of Ames recording solutions: Ames contains ascorbic acid, which is an enhancer of GABA
currents (Calero et al., 2011).

GABAergic reciprocal feedback from A17-ACs might be altered if GABA receptor
expression or localization was disrupted by TPBG-KO. $GABA_A$ receptors are large pentameric ion
channels usually containing α, β, and γ subunits. Early work indicates that RBC axon terminals
express $GABA_A\alpha1$, $\alpha3$, and $\gamma2$ receptor subunits (Fletcher et al., 1998), but a recent study found
that the $\alpha1$ subunit dominates in adult mice (Sinha et al., 2021). $GABA_{A\rho}$ receptors contain only
ρ subunits (Fletcher et al., 1998). Therefore, we used antibodies against $GABA_A\alpha1$ and $GABA_{A\rho}1$
to visualize $GABA_A$ and $GABA_{A\rho}$ receptor localization and an antibody against PKCα to isolate
RBC-specific axon terminals. We did not detect any difference in GABA receptor

immunofluorescence in RBC axon terminals between WT and TPBG-KO, indicating that reduced
GABAergic feedback is likely not due to disrupted GABA receptor localization.

We found that TPBG-KO increased scotopic ERG b-waves, suggesting an increase in
RBC depolarization. The scotopic ERG b-wave is initiated by current through TRPM1 non-specific
cation channels in the RBC dendrites. Yet, we did not find any difference in CPPG-induced
TRPM1 currents between WT and TPBG-KO retinas and the input resistance and excitability of
the RBC was unchanged. However, the scotopic b-wave is also shaped by reciprocal inhibition.
Pharmacological studies show that blocking inhibitory currents from third-order neurons changes
the scotopic b-wave. In particular, injections of a $GABA_A$ antagonist (bicuculline) or a $GABA_{A\rho}$
antagonist (TPMPA) into the IPL significantly increases the amplitude and duration of the scotopic
b-wave (Dong & Hare, 2000, 2003), suggesting that GABAergic feedback from A17-ACs
contributes to the shape of the RBC light response. We suggest that the enlarged ERG b-wave
in the TPBG-KO is due to reduced inhibitory feedback onto the RBC terminals, secondary to
reduced RBC glutamate release by the RBC.

Summary

Our findings show that genetic deletion of TPBG attenuates scotopic ERG oscillatory
potentials, synaptic vesicle exocytosis from RBCs, and reciprocal feedback onto RBCs. We
propose that TPBG is required for efficient coupling of Ca^{2+} influx and glutamate release at RBC
active zones, possibly through its effect on synaptic ribbons. Further evaluation of TPBG is critical
for producing a more comprehensive understanding of its role in fine-tuning RBC synaptic ribbon
physiology.

Figure Supplements
835

836
837   ***Figure 2 Supplementary Figure: TPBG-KO does not alter the scotopic ERG a-wave***. Quantification of a-wave
838   amplitudes recorded after light stimuli (-0.06 to 1.93 log(cd•s/m^2)). A-waves are measured from baseline to the a-wave
839   trough. Data is presented as the mean $\pm$ SEM.
840

841
842
Figure 4 Supplementary Figure: FIB-SEM imaging of RBC axon terminals. Focused ion beam scanning electron
microscopy (FIB-SEM) was used to image RBC axon terminals in the distal IPL and examine synaptic ribbon location
and morphology. Zoomed-out examples of WT and TPBG-KO (KO) RBC axon terminals in the IPL. The scale bar is 5
μm.

Figure 5 Supplementary Figure: Measuring membrane capacitance in RBCs. **(A)** Whole-cell voltage-clamp
capacitance recordings were made using the "sine + DC" method. A sinusoidal voltage command (200 Hz at 30 mV
trough-to-peak) was superimposed on the holding potential of -80 mV. Membrane capacitance (C_m), membrane
resistance (R_m), and series resistance (R_s) were recorded before and after a depolarizing pulse from -80 mV to -10 mV
for 100 ms. **(B)** A sample recording from a WT RBC using the sine + DC method as described. The change in membrane
capacitance (ΔC_m) was measured as the difference between the C_m before and after the depolarizing pulse. Changes
to R_m (middle) and R_s (bottom) were small and do not correlate with changes in C_m. **(C)** Average baseline-subtracted
C_m recordings using 200 Hz (blue; *n* = 6), 400 Hz (orange; *n* = 4), 800 Hz (green; *n* = 4), or 1600 Hz (magenta; *n* = 4)
sinusoidal voltage commands in WT RBCs. Data for the 200 Hz condition was reused from WT in Figure 4. Traces with
higher frequencies were resampled to 200 Hz for ease of comparisons. **(D)** Quantification of ΔC_m. **(E)** Average C_m
recordings and **(F)** quantification of resting C_m across frequencies. Resting C_m was measured before the depolarizing
pulse. Open circles in bar graphs represent single cells and bars represent the mean ± SEM. Statistical significance
was determined using Brown-Forsythe ANOVA tests; ****: *p* < 0.0001.

References

Agosto, M. A., & Wensel, T. G. (2020). LRRTM4 is a member of the transsynaptic complex between rod photoreceptors and bipolar cells. *Journal of Comparative Neurology*, *529*(1), 221–233. doi: 10.1002/cne.24944

Balakrishnan, V., Puthussery, T., Kim, M.-H., Taylor, W. R., & von Gersdorff, H. (2015). Synaptic Vesicle Exocytosis at the Dendritic Lobules of an Inhibitory Interneuron in the Mammalian Retina. *Neuron*, *87*(3), 563–575. doi: 10.1016/j.neuron.2015.07.016

Berglund, K., Midorikawa, M., & Tachibana, M. (2002). Increase in the Pool Size of Releasable Synaptic Vesicles by the Activation of Protein Kinase C in Goldfish Retinal Bipolar Cells. *The Journal of Neuroscience*, *22*(12), 4776–4785. doi: 10.1523/jneurosci.22-12-04776.2002

Calero, C. I., Vickers, E., Cid, G. M., Aguayo, L. G., von Gersdorff, H., & Calvo, D. J. (2011). Allosteric modulation of retinal GABA receptors by ascorbic acid. *The Journal of Neuroscience : The Official Journal of the Society for Neuroscience*, *31*(26), 9672–9682. doi: 10.1523/jneurosci.5157-10.2011

Cao, Y., Posokhova, E., & Martemyanov, K. A. (2011). TRPM1 Forms Complexes with Nyctalopin In Vivo and Accumulates in Postsynaptic Compartment of ON-Bipolar Neurons in mGluR6-Dependent Manner. *The Journal of Neuroscience*, *31*(32), 11521–11526. doi: 10.1523/jneurosci.1682-11.2011

Cao, Y., Sarria, I., Fehlhaber, K. E., Kamasawa, N., Orlandi, C., James, K. N., … Martemyanov, K. A. (2015). Mechanism for Selective Synaptic Wiring of Rod Photoreceptors into the Retinal Circuitry and Its Role in Vision. *Neuron*, *87*(6), 1248–1260. doi: 10.1016/j.neuron.2015.09.002

Cao, Y., Wang, Y., Dunn, H. A., Orlandi, C., Shultz, N., Kamasawa, N., … Martemyanov, K. A. (2020). Interplay between cell-adhesion molecules governs synaptic wiring of cone photoreceptors. *Proceedings of the National Academy of Sciences*, *117*(38), 23914–23924. doi: 10.1073/pnas.2009940117

Chávez, A. E., Grimes, W. N., & Diamond, J. S. (2010). Mechanisms Underlying Lateral GABAergic Feedback onto Rod Bipolar Cells in Rat Retina. *The Journal of Neuroscience*, *30*(6), 2330–2339. doi: 10.1523/jneurosci.5574-09.2010

Chávez, A. E., Singer, J. H., & Diamond, J. S. (2006). Fast neurotransmitter release triggered by Ca influx through AMPA-type glutamate receptors. *Nature*, *443*(7112), 705–708. doi: 10.1038/nature05123

Chen, M., & von Gersdorff, H. (2019). How to Build a Fast and Highly Sensitive Sound Detector That Remains Robust to Temperature Shifts. *The Journal of Neuroscience*, *39*(37), 7260–7276. doi: 10.1523/jneurosci.2510-18.2019

Dieck, S. T., & Brandstätter, J. H. (2006). Ribbon synapses of the retina. *Cell and Tissue Research*, *326*(2), 339–346. doi: 10.1007/s00441-006-0234-0

Dong, C.-J., Agey, P., & Hare, W. A. (2004). Origins of the electroretinogram oscillatory
potentials in the rabbit retina. *Visual Neuroscience*, *21*(4), 533–543. doi:
10.1017/s0952523804214043

Dong, C.-J., & Hare, W. A. (2000). Contribution to the kinetics and amplitude of the
electroretinogram b-wave by third-order retinal neurons in the rabbit retina. *Vision Research*,
40(6), 579–590. doi: 10.1016/s0042-6989(99)00203-5

Dong, C.-J., & Hare, W. A. (2003). Temporal Modulation of Scotopic Visual Signals by A17
Amacrine Cells in Mammalian Retina In Vivo. *Journal of Neurophysiology*, *89*(4), 2159–2166.
doi: 10.1152/jn.01008.2002

Doussau, F., & Augustine, G. J. (2000). The actin cytoskeleton and neurotransmitter release: An
overview. *Biochimie*, *82*(4), 353–363. doi: 10.1016/s0300-9084(00)00217-0

Eggermann, E., Bucurenciu, I., Goswami, S. P., & Jonas, P. (2012). Nanodomain coupling
between Ca2+ channels and sensors of exocytosis at fast mammalian synapses. *Nature*
Reviews Neuroscience, *13*(1), 7–21. doi: 10.1038/nrn3125

Euler, T., & Masland, R. H. (2000). Light-Evoked Responses of Bipolar Cells in a Mammalian
Retina. *Journal of Neurophysiology*, *83*(4), 1817–1829. doi: 10.1152/jn.2000.83.4.1817

Famiglietti, E. V., & Kolb, H. (1975). A bistratified amacrine cell and synaptic cirucitry in the
inner plexiform layer of the retina. *Brain Research*, *84*(2), 293–300.

Fletcher, E. L., Koulen, P., & Wässle, H. (1998). GABAA and GABAC receptors on mammalian
rod bipolar cells. *Journal of Comparative Neurology*, *396*(3), 351–365. doi:
10.1002/(sici)1096-9861(19980706)396:3<351::aid-cne6>3.0.co;2-1

Gillis, K. D. (2000). Admittance-based measurement of membrane capacitance using the EPC-9
patch-clamp amplifier. *Pflügers Archiv*, *439*(5), 655–664. doi: 10.1007/s004249900173

Grabner, C. P., & Moser, T. (2021). The mammalian rod synaptic ribbon is essential for Cav
channel facilitation and ultrafast synaptic vesicle fusion. *ELife*, *10*, e63844. doi:
10.7554/elife.63844

Grimes, W. N., Zhang, J., Graydon, C. W., Kachar, B., & Diamond, J. S. (2010). Retinal Parallel
Processors: More than 100 Independent Microcircuits Operate within a Single Interneuron.
Neuron, *65*(6), 873–885. doi: 10.1016/j.neuron.2010.02.028

Grimes, W. N., Zhang, J., Tian, H., Graydon, C. W., Hoon, M., Rieke, F., & Diamond, J. S.
(2015). Complex inhibitory microcircuitry regulates retinal signaling near visual threshold.
Journal of Neurophysiology, *114*(1), 341–353. doi: 10.1152/jn.00017.2015

Hartveit, E. (1996). Membrane currents evoked by ionotropic glutamate receptor agonists in rod
bipolar cells in the rat retinal slice preparation. *Journal of Neurophysiology*, *76*(1), 401–422.
doi: 10.1152/jn.1996.76.1.401

Hartveit, E. (1999). Reciprocal Synaptic Interactions Between Rod Bipolar Cells and Amacrine
Cells in the Rat Retina. *Journal of Neurophysiology*, *81*(6), 2923–2936. doi:
10.1152/jn.1999.81.6.2923

Hasan, N., Pangeni, G., Cobb, C. A., Ray, T. A., Nettesheim, E. R., Ertel, K. J., … Gregg, R. G.
(2019). Presynaptic Expression of LRIT3 Transsynaptically Organizes the Postsynaptic
Glutamate Signaling Complex Containing TRPM1. *Cell Reports*, *27*(11), 3107-3116.e3. doi:
10.1016/j.celrep.2019.05.056

Hasan, N., Pangeni, G., Ray, T. A., Fransen, K. M., Noel, J., Borghuis, B. G., … Gregg, R. G.
(2020). LRIT3 is required for Nyctalopin expression and normal ON and OFF pathway
signaling in the retina. *ENeuro, 7*(1), ENEURO.0002-20.2020. doi: 10.1523/eneuro.0002-
20.2020

Hasan, N., Ray, T. A., & Gregg, R. G. (2016). CACNA1S expression in mouse retina: Novel
isoforms and antibody cross-reactivity with GPR179. *Visual Neuroscience*, *33*, E009. doi:
10.1017/s0952523816000055

He, P., Jiang, S., Ma, M., Wang, Y., Li, R., Fang, F., … Zhang, Z. (2015). Trophoblast
glycoprotein promotes pancreatic ductal adenocarcinoma cell metastasis through Wnt/planar
cell polarity signaling. *Molecular Medicine Reports*, *12*(1), 503–509. doi:
10.3892/mmr.2015.3412

Jarsky, T., Tian, M., & Singer, J. H. (2010). Nanodomain Control of Exocytosis Is Responsible
for the Signaling Capability of a Retinal Ribbon Synapse. *Journal of Neuroscience*, *30*(36),
11885–11895. doi: 10.1523/jneurosci.1415-10.2010

Job, C., & Lagnado, L. (1998). Calcium and Protein Kinase C Regulate the Actin Cytoskeleton
in the Synaptic Terminal of Retinal Bipolar Cells. *The Journal of Cell Biology*, *143*(6), 1661–
1672. doi: 10.1083/jcb.143.6.1661

Kagermeier-Schenk, B., Wehner, D., Özhan-Kizil, G., Yamamoto, H., Li, J., Kirchner, K., …
Weidinger, G. (2011). Waif1/5T4 Inhibits Wnt/β-Catenin Signaling and Activates
Noncanonical Wnt Pathways by Modifying LRP6 Subcellular Localization. *Developmental
Cell*, *21*(6), 1129–1143. doi: 10.1016/j.devcel.2011.10.015

Kim, M.-H., & von Gersdorff, H. (2016). Postsynaptic Plasticity Triggered by Ca^{2+}-Permeable
AMPA Receptor Activation in Retinal Amacrine Cells. *Neuron*, *89*(3), 507–520. doi:
10.1016/j.neuron.2015.12.028

Klyachko, V. A., & Jackson, M. B. (2002). Capacitance steps and fusion pores of small and
large-dense-core vesicles in nerve terminals. *Nature*, *418*(6893), 89–92. doi:
10.1038/nature00852

Kobe, B., & Deisenhofer, J. (1994). The leucine-rich repeat: a versatile binding motif. *Trends in
Biochemical Sciences*, *19*(10), 415–421. doi: 10.1016/0968-0004(94)90090-6

Kobe, B., & Kajava, A. V. (2001). The leucine-rich repeat as a protein recognition motif. *Current
Opinion in Structural Biology*, *11*(6), 725–732. doi: 10.1016/s0959-440x(01)00266-4

Koike, C., Numata, T., Ueda, H., Mori, Y., & Furukawa, T. (2010). TRPM1: A vertebrate TRP
channel responsible for retinal ON bipolar function. *Cell Calcium*, *48*(2–3), 95–101. doi:
10.1016/j.ceca.2010.08.004

Koike, C., Obara, T., Uriu, Y., Numata, T., Sanuki, R., Miyata, K., … Furukawa, T. (2010).
TRPM1 is a component of the retinal ON bipolar cell transduction channel in the mGluR6
cascade. *Proceedings of the National Academy of Sciences*, *107*(1), 332–337. doi:
10.1073/pnas.0912730107

Korol, S., Leuenberger, P. M., Englert, U., & Babel, J. (1975). In vivo effects of glycine on retinal
ultrastructure and averaged electroretinogram. *Brain Research*, *97*(2), 235–251. doi:
10.1016/0006-8993(75)90447-3

Kushmerick, C., Renden, R., & von Gersdorff, H. (2006). Physiological Temperatures Reduce
the Rate of Vesicle Pool Depletion and Short-Term Depression via an Acceleration of Vesicle
Recruitment. *The Journal of Neuroscience*, *26*(5), 1366–1377. doi: 10.1523/jneurosci.3889-
05.2006

Liao, F., Liu, H., Milla-Navarro, S., Villa, P. de la, & Germain, F. (2023). Origin of Retinal
Oscillatory Potentials in the Mouse, a Tool to Specifically Locate Retinal Damage.
International Journal of Molecular Sciences, *24*(4), 3126. doi: 10.3390/ijms24043126

Masu, M., Iwakabe, H., Tagawa, Y., Miyoshi, T., Yamashita, M., Fukuda, Y., … Nakanishi, S.
(1995). Specific deficit of the ON response in visual transmission by targeted disruption of
the mGluR6 gene. *Cell*, *80*(5), 757–765. doi: 10.1016/0092-8674(95)90354-2

Maxeiner, S., Luo, F., Tan, A., Schmitz, F., & Südhof, T. C. (2016). How to make a synaptic
ribbon: RIBEYE deletion abolishes ribbons in retinal synapses and disrupts neurotransmitter
release. *The EMBO Journal*, *35*(10), 1098–1114. doi: 10.15252/embj.201592701

McGuire, B., Stevens, J., & Sterling, P. (1984). Microcircuitry of bipolar cells in cat retina. *The
Journal of Neuroscience*, *4*(12), 2920–2938. doi: 10.1523/jneurosci.04-12-02920.1984

Minami, N., Berglund, K., Sakaba, T., Kohmoto, H., & Tachibana, M. (1998). Potentiation of
transmitter release by protein kinase C in goldfish retinal bipolar cells. *The Journal of
Physiology*, *512*(1), 219–225. doi: 10.1111/j.1469-7793.1998.219bf.x

Morgans, C. W. (2000). Neurotransmitter release at ribbon synapses in the retina. *Immunology
and Cell Biology*, *78*(4), 442–446. doi: 10.1046/j.1440-1711.2000.00923.x

Morgans, C. W., Zhang, J., Jeffrey, B. G., Nelson, S. M., Burke, N. S., Duvoisin, R. M., & Brown,
R. L. (2009). TRPM1 is required for the depolarizing light response in retinal ON-bipolar
cells. *Proceedings of the National Academy of Sciences*, *106*(45), 19174–19178. doi:
10.1073/pnas.0908711106

Moser, T., Grabner, C. P., & Schmitz, F. (2020). Sensory Processing at Ribbon Synapses in the
Retina and the Cochlea. *Physiological Reviews*, *100*(1), 103–144. doi:
10.1152/physrev.00026.2018

Nakajima, Y., Iwakabe, H., Akazawa, C., Nawa, H., Shigemoto, R., Mizuno, N., & Nakanishi, S.
(1993). Molecular characterization of a novel retinal metabotropic glutamate receptor
mGluR6 with a high agonist selectivity for L-2-amino-4-phosphonobutyrate. *The Journal of*
Biological Chemistry, *268*(16), 11868–11873.

Neher, E., & Marty, A. (1982). Discrete changes of cell membrane capacitance observed under
conditions of enhanced secretion in bovine adrenal chromaffin cells. *Proceedings of the*
National Academy of Sciences, *79*(21), 6712–6716. doi: 10.1073/pnas.79.21.6712

Neuillé, M., Cao, Y., Caplette, R., Guerrero-Given, D., Thomas, C., Kamasawa, N., … Zeitz, C.
(2017). LRIT3 Differentially Affects Connectivity and Synaptic Transmission of Cones to ON-
and OFF-Bipolar CellsLRIT3's Role in Cone to Bipolar Cell Transmission. *Investigative*
Ophthalmology & Visual Science, *58*(3), 1768–1778. doi: 10.1167/iovs.16-20745

Neuillé, M., Morgans, C. W., Cao, Y., Orhan, E., Michiels, C., Sahel, J., … Zeitz, C. (2015).
LRIT3 is essential to localize TRPM1 to the dendritic tips of depolarizing bipolar cells and
may play a role in cone synapse formation. *European Journal of Neuroscience*, *42*(3), 1966–
1975. doi: 10.1111/ejn.12959

Ogden, T. E. (1973). The oscillatory waves of the primate electroretinogram. *Vision Research*,
13(6), 1059-IN4. doi: 10.1016/0042-6989(73)90144-2

Oltedal, L., & Hartveit, E. (2010). Transient release kinetics of rod bipolar cells revealed by
capacitance measurement of exocytosis from axon terminals in rat retinal slices. *The Journal*
of Physiology, *588*(9), 1469–1487. doi: 10.1113/jphysiol.2010.186916

Oltedal, L., Veruki, M. L., & Hartveit, E. (2009). Passive membrane properties and electrotonic
signal processing in retinal rod bipolar cells. *The Journal of Physiology*, *587*(4), 829–849.
doi: 10.1113/jphysiol.2008.165415

Palmer, M. J. (2006). Functional segregation of synaptic GABAA and GABAC receptors in
goldfish bipolar cell terminals. *The Journal of Physiology*, *577*(1), 45–53. doi:
10.1113/jphysiol.2006.119560

Paridaen, J. T. M. L., Wilsch-Bräuninger, M., & Huttner, W. B. (2013). Asymmetric Inheritance of
Centrosome-Associated Primary Cilium Membrane Directs Ciliogenesis after Cell Division.
Cell, *155*(2), 333–344. doi: 10.1016/j.cell.2013.08.060

Pearring, J. N., Bojang, P., Shen, Y., Koike, C., Furukawa, T., Nawy, S., & Gregg, R. G. (2011).
A Role for Nyctalopin, a Small Leucine-Rich Repeat Protein, in Localizing the TRP
Melastatin 1 Channel to Retinal Depolarizing Bipolar Cell Dendrites. *The Journal of*
Neuroscience, *31*(27), 10060–10066. doi: 10.1523/jneurosci.1014-11.2011

Protti, D. A., & Llano, I. (1998). Calcium Currents and Calcium Signaling in Rod Bipolar Cells of
Rat Retinal Slices. *Journal of Neuroscience*, *18*(10), 3715–3724. doi: 10.1523/jneurosci.18-
10-03715.1998

Raviola, E., & Dacheux, R. F. (1987). Excitatory dyad synapse in rabbit retina. *Proceedings of*
the National Academy of Sciences, *84*(20), 7324–7328. doi: 10.1073/pnas.84.20.7324

Robson, J. G., & Frishman, L. J. (2014). The rod-driven a-wave of the dark-adapted mammalian
electroretinogram. *Progress in Retinal and Eye Research*, *39*, 1–22. doi:
10.1016/j.preteyeres.2013.12.003

Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T., … Cardona,
A. (2012). Fiji: an open-source platform for biological-image analysis. *Nature Methods*, *9*(7),
676–682. doi: 10.1038/nmeth.2019

Shen, Y., Heimel, J. A., Kamermans, M., Peachey, N. S., Gregg, R. G., & Nawy, S. (2009). A
Transient Receptor Potential-Like Channel Mediates Synaptic Transmission in Rod Bipolar
Cells. *The Journal of Neuroscience*, *29*(19), 6088–6093. doi: 10.1523/jneurosci.0132-
09.2009

Singer, J. H., & Diamond, J. S. (2003). Sustained Ca2+ Entry Elicits Transient Postsynaptic
Currents at a Retinal Ribbon Synapse. *Journal of Neuroscience*, *23*(34), 10923–10933. doi:
10.1523/jneurosci.23-34-10923.2003

Sinha, R., Grimes, W. N., Wallin, J., Ebbinghaus, B. N., Luu, K., Cherry, T., … Hoon, M. (2021).
Transient expression of a GABA receptor subunit during early development is critical for
inhibitory synapse maturation and function. *Current Biology*, *31*(19), 4314-4326.e5. doi:
10.1016/j.cub.2021.07.059

Sinha, R., Siddiqui, T. J., Padmanabhan, N., Wallin, J., Zhang, C., Karimi, B., … Hoon, M.
(2020). LRRTM4: A Novel Regulator of Presynaptic Inhibition and Ribbon Synapse
Arrangements of Retinal Bipolar Cells. *Neuron*, *105*(6), 1007-1017.e5. doi:
10.1016/j.neuron.2019.12.028

Stockton, R. A., & Slaughter, M. M. (1989). B-wave of the electroretinogram. A reflection of ON
bipolar cell activity. *The Journal of General Physiology*, *93*(1), 101–122. doi:
10.1085/jgp.93.1.101

Tachibana, M. (1999). Regulation of transmitter release from retinal bipolar cells. *Progress in
Biophysics and Molecular Biology*, *72*(2), 109–133. doi: 10.1016/s0079-6107(99)00003-6

Tsuboi, A. (2020). LRR-Containing Oncofetal Trophoblast Glycoprotein 5T4 Shapes Neural
Circuits in Olfactory and Visual Systems. *Frontiers in Molecular Neuroscience*, *13*, 581018.
doi: 10.3389/fnmol.2020.581018

Vaegan, & Millar, T. J. (1994). Effect of kainic acid and NMDA on the pattern electroretinogram,
the scotopic threshold response, the oscillatory potentials and the electroretinogram in the
urethane anaesthetized cat. *Vision Research*, *34*(9), 1111–1125. doi: 10.1016/0042-
6989(94)90294-1

Vardi, N., Duvoisin, R., Wu, G., & Sterling, P. (2000). Localization of mGluR6 to dendrites of ON
bipolar cells in primate retina. *Journal of Comparative Neurology*, *423*(3), 402–412. doi:
10.1002/1096-9861(20000731)423:3<402::aid-cne4>3.0.co;2-e

von Gersdorff, H., & Matthews, G. (1997). Depletion and Replenishment of Vesicle Pools at a
Ribbon-Type Synaptic Terminal. *Journal of Neuroscience*, *17*(6), 1919–1927. doi:
10.1523/jneurosci.17-06-01919.1997

von Gersdorff, H., Vardi, E., Matthews, G., & Sterling, P. (1996). Evidence That Vesicles on the
Synaptic Ribbon of Retinal Bipolar Neurons Can Be Rapidly Released. *Neuron*, *16*(6), 1221–
1227. doi: 10.1016/s0896-6273(00)80148-8

Wachtmeister, L. (1998). Oscillatory potentials in the retina: what do they reveal. *Progress in
Retinal and Eye Research*, *17*(4), 485–521. doi: 10.1016/s1350-9462(98)00006-8

Wakeham, C. M., Ren, G., & Morgans, C. W. (2020). Expression and distribution of trophoblast
glycoprotein in the mouse retina. *Journal of Comparative Neurology*, *528*(10), 1660–1671.
doi: 10.1002/cne.24850

Wakeham, C. M., Wilmarth, P. A., Cunliffe, J. M., Klimek, J. E., Ren, G., David, L. L., &
Morgans, C. W. (2019). Identification of PKCα-dependent phosphoproteins in mouse retina.
Journal of Proteomics, *206*, 103423. doi: 10.1016/j.jprot.2019.103423

Xiong, W.-H., Pang, J.-J., Pennesi, M. E., Duvoisin, R. M., Wu, S. M., & Morgans, C. W. (2015).
The Effect of PKCα on the Light Response of Rod Bipolar Cells in the Mouse Retina.
Investigative Ophthalmology & Visual Science, *56*(8), 4961–4974. doi: 10.1167/iovs.15-
16622

Xu, Y., Dhingra, A., Fina, M. E., Koike, C., Furukawa, T., & Vardi, N. (2012). mGluR6 deletion
renders the TRPM1 channel in retina inactive. *Journal of Neurophysiology*, *107*(3), 948–957.
doi: 10.1152/jn.00933.2011

Yoshihara, S.-I., Takahashi, H., Nishimura, N., Naritsuka, H., Shirao, T., Hirai, H., … Tsuboi, A.
(2012). 5T4 Glycoprotein Regulates the Sensory Input-Dependent Development of a Specific
Subtype of Newborn Interneurons in the Mouse Olfactory Bulb. *The Journal of Neuroscience*,
32(6), 2217–2226. doi: 10.1523/jneurosci.5907-11.2012

Zanazzi, G., & Matthews, G. (2009). The Molecular Architecture of Ribbon Presynaptic
Terminals. *Molecular Neurobiology*, *39*(2), 130–148. doi: 10.1007/s12035-009-8058-z

Zhao, Y., Malinauskas, T., Harlos, K., & Jones, E. Y. (2014). Structural Insights into the
Inhibition of Wnt Signaling by Cancer Antigen 5T4/Wnt-Activated Inhibitory Factor 1.
Structure, *22*(4), 612–620. doi: 10.1016/j.str.2014.01.009

Zhou, Z.-Y., Wan, Q.-F., Thakur, P., & Heidelberger, R. (2006). Capacitance Measurements in
the Mouse Rod Bipolar Cell Identify a Pool of Releasable Synaptic Vesicles. *Journal of
Neurophysiology*, *96*(5), 2539–2548. doi: 10.1152/jn.00688.2006

1116