Parallel duplication and loss of aquaporin-coding genes during the 'out of the sea' transition paved the way for animal terrestrialization

Gemma I. Martínez-Redondo[1*], Carolina Simón Guerrero[1,2], Leandro Aristide[1], Pau Balart-García[1], Vanina Tonzo[1,3], Rosa Fernández[1*]

[1] Metazoa Phylogenomics Lab, Biodiversity Program, Institute of Evolutionary Biology (CSIC-Universitat Pompeu Fabra), Passeig Marítim de la Barceloneta 37-49, 08003 Barcelona, Spain

[2] Current institution: Biocomputing Unit, National Centre of Biotechnology (CNB-CSIC), Campus de Cantoblanco, 28049 Madrid, Spain.

[3] Current institution: Centre d'Ecologie Fonctionnelle et Evolutive (CEFE-CNRS), 1919 Route de Mende, 34293 Montpellier 5, France.

[*] Corresponding authors: Gemma I. Martínez Redondo, gemma.martinez@ibe.upf-csic.es, and Rosa Fernández, rosa.fernandez@ibe.upf.csic.es

Abstract

One of the most important physiological challenges animals had to overcome during terrestrialization (i.e., the transition from sea to land) is water loss, which alters their osmotic and hydric homeostasis. Aquaporins are a superfamily of membrane water transporters heavily involved in osmoregulatory processes. Their diversity and evolutionary dynamics in most animal lineages remain unknown, hampering our understanding of their role in marine-terrestrial transitions. Here, we interrogated aquaporin gene repertoire evolution across the main terrestrial animal lineages. We annotated aquaporin-coding genes in genomic data from 458 species from 7 animal phyla where terrestrialization episodes occurred. We then explored aquaporin gene evolutionary dynamics to assess differences between terrestrial and aquatic species through phylogenomics and phylogenetic comparative methods. Our results revealed parallel aquaporin-coding gene duplications in aquaporins transporting water, glycerol and ammonia in the transition from marine to non-marine environments (e.g., brackish, freshwater and terrestrial), rather than from aquatic to terrestrial ones, with some notable duplications in ancient lineages. Contrarily, we also recovered a significantly lower number of oxygen peroxide-transporting aquaporin-coding genes in terrestrial arthropods, suggesting that more efficient oxygen homeostasis in land arthropods might be linked to a reduction in this type of aquaporins. Our results thus indicate

that aquaporin-coding gene duplication and loss might have been one of the key steps towards the evolution of osmoregulation across animals, facilitating the 'out of the sea' transition and ultimately the colonisation of land.

Introduction

The ecological transition from water to land (i.e. terrestrialization) is one of the most remarkable episodes that shaped the evolution of biodiversity as we know it today. Throughout the eukaryote tree of life, living beings have shifted their habitat from water to land multiple times (Jamy et al.). Among them, animals have successfully terrestrialized independently in 10 different phyla: chordates, nematodes, nematomorphs, tardigrades, arthropods, onychophorans, platyhelminthes, molluscs, annelids, and nemerteans. Within most phyla, more than one independent habitat shift has occurred, ranging from only one terrestrialization event in the onychophorans to more than 30 in nematodes (Holterman et al. 2019), at different times along Earth's history. The most studied terrestrialization event is, without a shadow of a doubt, the one that gave rise to terrestrial tetrapods during the Devonian period from euryhaline aquatic ancestors (Goedert et al. 2018; Qvarnström et al. 2018). Nevertheless, the most ancient terrestrialization events occurred in arthropods (arachnids, myriapods and hexapods) from marine ancestors during the late Cambrian-early Ordovician (Rota-Stabelli et al. 2013). More than 4 additional independent terrestrialization events have occurred in the following millions of years within Pancrustacea (isopods, amphipods and anomura and brachyura decapods) (Sharma 2017; Watson-Zink 2021). Other terrestrialization events that have recently gained interest are the ones in molluscs and annelids. In the case of molluscs, gastropods have independently colonised land more than 9 times (Kameda and Kato 2011), with the latest estimates suggesting up to 30 times (Vermeij and Watson-Zink 2022), either following a freshwater route or through marginal environments such as brackish water (van Straalen 2021; Krug et al. 2022; Vermeij and Watson-Zink 2022). Among them, perhaps the most successful terrestrial gastropod lineage is the Stylommatophora, an order of air-breathing pulmonate land gastropods that contains most living snails and slugs. Regarding annelids, earthworms (Crassiclitellata) derive from a single terrestrialization event, while other events have occurred within Hirudinida and Enchytraeidae (Erséus et al. 2020; van Straalen 2021). Another example is that of onychophorans, or velvet worms, which are the only entirely terrestrial animal phylum. This lineage closely related to arthropods has been hypothesised to have a marine origin based on the similar osmotic pressure of their haemolymph to terrestrial arthropods (Little 1990). All these independent colonisation events (Fig 1A) make animal terrestrialization such a

complex and intriguing area of study, especially since the genomic underpinnings facilitating these transitions remain highly unexplored.

Terrestrialization required many complex physiological changes in different biological processes, such as respiration, locomotion or reproduction, that allowed them to overcome the environmental barriers and adapt to life on land. One of the most important challenges species faced during this transition was avoiding water loss, which alters their osmotic and hydric homeostasis. Most marine animals are osmoconformers, which means that they cannot regulate their internal osmotic pressure under environmental shifts. When marine animals transitioned to euryhaline or freshwater habitats, they faced osmotic stress due to the differences in the ionic composition between their extracellular fluids and the environment. The development of the ability to osmoregulate, i.e. to maintain the osmotic pressure of their extracellular fluids (Rankin and Davenport 1981), allowed them to overcome this osmotic shock. Accordingly, terrestrial animals have acquired several physiological adaptive mechanisms to regulate their internal osmotic balance and water loss. Such is the case of closing spiracles in arthropods with trachea or the pneumostome in terrestrial pulmonate gastropods to regulate gas exchange, the evolution of a kidney structure and non-ammonia excretion in tetrapods, or the development of Malpighian tubules in terrestrial arthropods for regulating osmotic pressure.

At the molecular level, animals can modify the osmotic composition of their fluids by regulating the transport of ions or water through biological membranes (Uchiyama and Konno 2006; Takvam et al. 2021). Previous experiments have shown that cells of these osmoregulatory organs express protein channels called aquaporins in their membranes (Spring et al. 2009; Yi et al. 2011; Finn et al. 2014; Wittekindt and Dietl 2019; Cabrero et al. 2020). Aquaporins are a superfamily of small membrane intrinsic proteins (24-36kDa) that are heavily involved in osmoregulatory processes by forming pores in the cell membrane that facilitate the transport of water and other small molecules (Kruse et al. 2006; Abascal et al. 2014). Structurally, the basic units of aquaporins are monomers characterised by six transmembrane domains that surround a narrow pore, and two selectivity filters: two NPA (Asn-Pro-Ala) motifs, responsible for proton exclusion, and an aromatic/arginine (ar/R) region that determines the pore size (Fig 1B). Variations in these two selectivity filters slightly change the pores' size, structure and hydrophobicity, and ultimately determine different subtypes of aquaporins and their ability to transport molecules different from water (Fig 1C) (de Maré et al. 2020). In the membrane, four aquaporin monomers form a tetramer, creating a fifth pore (Kreida and Törnroth-Horsefield 2015). This general structure is conserved between bacteria and eukaryotic aquaporins, as well as between different subtypes of

aquaporins, despite the low sequence similarity of these proteins (less than 30%) (Kruse et al. 2006).

Phylogenetic analysis of aquaporins across all organisms, together with the permeability differences of aquaporins, have allowed their classification into two main subtypes: aquaporins that transport water (Aqps), and aquaglyceroporins that are also permeable to glycerol and other molecules (Glps). Further studies have shown that aquaporins are also permeable to other molecules (CO_2, H_2O_2 or ammonia, among others) (Finn and Cerdà 2015). Bacteria and archaea generally have one gene copy of each subtype, but some groups retain only Glps or Aqps. Conversely, eukaryote aquaporins have recurrently expanded, giving rise to a large aquaporin diversity. In animals, Aqps can be further divided into three subtypes: classical aquaporins, aquaammoniaporins, and unorthodox aquaporins or superaquaporins, whose origin dates back to the base of the eukaryotic tree (Fig 1D) (Abascal et al. 2014).

Previous studies have analysed the repertoire of aquaporin-coding genes (Aqp-coding genes hereafter) in selected animal species (e.g., *Danio rerio* (Tingaud-Sequeira et al. 2010), *Caenorhabditis elegans (Huang et al. 2007)*, *Milnesium tardigradum (Grohme et al. 2013),* blood-feeding arthropods (Benoit et al. 2014), dust mites (Peng et al. 2018) or the salmon louse and its host (Stavang et al. 2015), to name a few). Meanwhile, others have focused on the diversity and evolution of animal aquaporins in groups of human interest, such as vertebrate (Finn et al. 2014) and hemipteran aquaporins (Van Ekert et al. 2016), or insect entomoglyceroporins (Finn et al. 2015). Recent studies have leveraged the newly available transcriptomic and genomic resources to expand taxon sampling and dig deeper into aquaporin (or aquaglyceroporin) diversity in groups of animals that are not widely studied, such as copepods and crustaceans (Catalán-García et al. 2021), gastropods (Colgan and Santos 2018), annelids (Mucciolo et al. 2021), and non-vertebrate deuterostomes (Yilmaz et al. 2020). However, the whole aquaporin gene repertoire in other animal groups, such as molluscs and less studied orders of arthropods (i.e. chelicerates other than spiders, scorpions and mites), has not yet been explored, let alone in a comparative framework within the context of animal terrestrialization. Only a couple of previous studies have explored the potential key role of aquaporin gene duplications in the adaptation of vertebrates to life on land, with the identification of 3 tetrapod-exclusive classical aquaporins functionally related to water absorption mechanisms (Finn et al. 2014), and some signals of positive selection in semiterrestrial mudskippers' aquaporins (Lorente-Martínez et al. 2018). Nevertheless, even though the importance of gene duplication in the origin of new functions and driving adaptation has long been reported

(Kondrashov 2012), the potential role of aquaporin gene duplication and loss in the terrestrialization of lineages other than tetrapods has not yet been explored.

Here, we test the hypothesis that gene duplication and loss in Aqp-coding genes played a key role in animal terrestrialization and facilitated marine-to-land transition in other phyla beyond chordates. To investigate this, we mined and annotated more than 6,300 Aqp-coding gene sequences from 458 genomic and transcriptomic data sets, maximising taxon sampling to ensure a good representation of most animal lineages with terrestrial representatives. To assess whether aquaporins may have had a role during animal terrestrialization, we explored aquaporin gene evolutionary dynamics to assess differences between terrestrial and aquatic species through phylogenomics and phylogenetic comparative methods. We interpreted and discussed our results in the context of animal terrestrialization.

Results and Discussion

An expanded catalogue of aquaporins in arthropods, molluscs and onychophorans unveils the complex dynamics of aquaporin evolution

Our knowledge of animal aquaporins' evolution is patchy. To close some gaps that affect our understanding of their potential involvement in terrestrialization, we first explored the aquaporin gene repertoire in arthropods, molluscs and onychophorans, since we had an extensive taxon sampling for each phylum. We identified a total of 2,303 arthropodan, 1,225 molluscan, and 52 onychophoran Aqps (Supp. Data 1) after annotating 310 arthropodan, 101 molluscan and 4 onychophoran genomic data sets. Only some Aqps from a few arthropod species and even fewer mollusc species had been previously annotated as aquaporins (Finn et al. 2015; Stavang et al. 2015; Colgan and Santos 2018; Catalán-García et al. 2021). We also report here the first identification of Aqps in the phylum Onychophora.

As we wanted to include additional phyla to have a better understanding of the evolution of Aqp-coding genes in animals that have transitioned from the sea to the land, we searched the literature for additional aquaporin sequences in representatives from other phyla that underwent terrestrialization episodes (Nematoda, Tardigrada, Craniata and Annelida), obtaining a total of 4,230 aquaporins. Aqps of *C. elegans (Huang et al. 2007)* and *M. tardigradum (Grohme et al. 2013)* were classified into the four main subtypes by building a phylogenetic tree with the backbone sequences. For annelids and vertebrates, aquaporins

had already been classified into the main groups, which were later confirmed in our phylogeny.

Our inferred phylogeny of all animal Aqps (Fig S1) supports the four aquaporin clades previously defined in the literature (Abascal et al. 2014): aquaglyceroporins (which we have termed Aqp3-like), classical aquaporins (Aqp1-like), aquaammoniaporins (Aqp8-like) and superaquaporins (Aqp11-like). Moreover, the loss of Aqp8-like in arthropods and Aqp1-like in *C. elegans* is recovered, as previously reported (Finn and Cerdà 2015; Stavang et al. 2015), together with the identification of Aqp8-like loss in onychophorans, suggesting that the loss of Aqp8 may have occurred before the split between Onychophora and Arthropoda. Hexapod-specific glyceroporins (i.e. entomoglyceroporins or Eglps) clustered within arthropod Aqp1-like, showing their Prip-like origin as previously reported (Finn et al. 2015).

Parallel aquaporin-coding gene duplications in aquaporins transporting water, glycerol and ammonia may have facilitated the transition from marine to non-marine environments in animals

The presence of Aqp clades specific to terrestrial animals, such as the Eglps in insects (Finn et al. 2015) or AQP2, AQP5 and AQP6 in tetrapods (Finn et al. 2014; Lorente-Martínez et al. 2018), suggests that the number of aquaporins found in terrestrial animals might be different than the number found in aquatic ones. To test whether the number of aquaporins was related to the habitat in which extant species live, we obtained Aqp-coding genes identified in each species and performed a phylogenetic Poisson regression using the habitat as an independent variable. We tested two different hypotheses using this model. The first one is that the number of Aqp-coding genes in terrestrial species is different to that of aquatic ones (including freshwater, brackish and marine), supporting that a copy number variation of Aqp-coding genes through gene gain, duplication or loss facilitated terrestrialization during aquatic-terrestrial transitions. The second hypothesis tested is that the number of Aqp-coding genes in marine species is different to that in non-marine species (including in this case brackish, freshwater and terrestrial species). The rationale behind this second test is the fact that non-marine species are osmoregulators compared to marine species, which are osmoconformers (except for teleosts due to their hypothetical freshwater origin (Rankin and Davenport 1981; Carrete Vega and Wiens 2012). This may suggest that the number of aquaporins in species that live in marine environments may be different than in non-marine species, hence with this hypothesis we are testing that

gene gain, duplication or loss in Aqp-coding genes may underlie the emergence of osmoregulation.

We first explored two independent tests with genomic data of the five phyla (Onychophora, Craniata (Vertebrata), Mollusca, Annelida and Arthropoda; five-phyla data set hereafter; Fig 2A). Tardigrades and nematodes were not included as we only analysed one species per phylum and lacked the representation we had in the others, which could bias the results. Moreover, since we had a dense taxon representation of most main lineages in arthropods and molluscs, we performed the tests independently for each of these two phyla as well. These tests were explored both taking into account all Aqp-coding genes and in each type of Aqp-coding genes independently, following the main four types described above (i.e., Aqp1-like, Aqp3-like, Aqp8-like and Aqp11-like).

Overall, our results showed statistical differences in the number of several aquaporin-coding gene subfamilies when we compared marine and non-marine species, pointing to a significantly higher number in non-marine species than in marine ones in most cases (Fig 2B and C). First, in the analyses of the five-phyla data set we observed that non-marine animals have a significantly higher number of aquaglycerolporins (i.e. Aqp3-like) compared to marine species (Fig 2B, left). The role of polyols (*e.g.* glycerol) and sugars (*e.g.* trehalose) as osmolytes in animal osmoregulation has been known for years (Lamitina et al. 2004; Christoph et al. 2007; Thorat et al. 2012), as well as the ability of Aqp3-like to transport these molecules under osmotic stress (Laforenza et al. 2013; Gil et al. 2017). Thus, during animal paths out of the sea towards land or freshwater habitats, a higher number of aquaglyceroporins may have provided a more efficient osmolyte flux in fluids to counteract water loss. A new nomenclature for Aqp3-like subclades in the different animal phyla and the implications of its phylogenetic history in the context of animal terrestrialization is discussed in detail below (see also Table 1).

Regarding the phylum-specific analyses in molluscs, we recovered a significantly higher number of Aqp-coding genes in non-marine compared to marine species (Fig 2C). Independent analyses for each type of Aqp-coding gene revealed a significantly higher number in non-marine species of mainly water-transporting aquaporins (Aqp1-like, also referred to as classical aquaporins) and ammonia-transporting aquaporins (Aqp8-like, also referred to as aquammioporins) (Fig 2B, right). As the number of aquaporins in terrestrial species compared to non-terrestrial ones is not significant, this suggests that, as in the case of aquaglyceroporins in the five-phyla analysis, aquaporin-coding gene duplication potentially enabled to cope with osmotic stress during the transition of molluscs to freshwater

and land. In fact, the role of Aqp1-like expansions in the transition from the sea to freshwater was previously suggested for freshwater bivalves (Calcino et al. 2019), as well as their critical role during the early cleavage of the freshwater bivalve *Dreissena rostriformis* (Zieger et al. 2022). A description of the phylogenetic relationships of Aqp1-like and Aqp8-like as well as their potential role in mollusc terrestrialization is discussed below.

Gene loss in oxygen peroxide-transporting aquaporin-coding genes may underlie more efficient oxygen homeostasis in terrestrial arthropods

Our results indicated that terrestrial arthropods have a significantly lower mean number of Aqp11-like aquaporins than aquatic ones (Fig 2B, centre). Aqp11-like (also called superaquaporins) are less studied than other aquaporins, with virtually no functional studies to understand their role at the cellular and physiological levels. This type of Aqps is expressed intracellularly and is involved in keeping redox/pH homeostasis in the endoplasmic reticulum (ER) (Nozaki et al. 2008; Bestetti et al. 2020; Ishibashi et al. 2021). During hypoxia, Reactive Oxygen Species (ROS), such as H_2O_2, are generated in the ER. As the levels of oxygen are higher on land, the transition from the sea may have removed the need for H_2O_2 transport through Aqp11-like due to a lower production of ROS, which may have triggered gene loss in Aqp11-like coding genes. Hence, we may hypothesise a potential link of gene loss in Aqp11-like coding genes with more efficient oxygen homeostasis in land arthropods. This hypothesis is further reinforced by the identification of several independent duplications of Aqp11-like coding genes in aquatic arthropod species -i.e. xiphosurans and mayflies (Fig S2), all of them occurring in species that had habitat transitions back to aquatic habitats (from land to sea, and land to freshwater, respectively). In addition, we detect Aqp11-like duplications in freshwater copepods and crayfish, which may be related to their transition from sea to freshwater. Further functional studies are most needed to understand the functional role of this enigmatic type of aquaporins.

Ancient independent duplications in Aqp-coding genes may have paved the way for the 'out of the sea' transition across animal phyla

   a) *Classical aquaporins (Aqp1-like) and aquaamnioporins (Aqp8-like)*

The phylogeny of protostome classical aquaporins (Fig 3, Fig S3) reveals a complex pattern of gains and losses in the different aquaporin-coding gene clusters, with many originating early during animal evolution. We highlight two clades of Aqps1-like, which we

coined as Aqp1-bila1 and Aqp1-bila2, and that include the majority of Aqp1-like animal sequences (except for Aqp1-ann1, Prip and Drip). Since deuterostomes (mostly vertebrates, Fig 3, in turquoise) cluster inside Aqp1-bila2, while being absent in Aqp1-bila1, we can infer that the duplication that gave rise to both Aqp1-bila clades occurred at the base of Bilateria, before the diversification into Deuterostomia and Protostomia. Both Aqp1-bila copies were further duplicated or lost independently across animal evolution. Aqp1-bila1 contains mainly genes from species within Spiralia, together with Onychophora and Tardigrada, while Aqp1-bila2 includes sequences from all animal phyla included in this study, except for nematodes, for which a lack of Aqp1-like sequences had previously been reported (Finn and Cerdà 2015). It is worth mentioning that lineages that have the most profound adaptations to life on land (i.e. tetrapods and arthropods) underwent the loss of Aqp1-bila1 and only retain Aqp1-bila2.

Classical aquaporins were also independently duplicated in terrestrial lineages (Fig 2B and C). Our data show a not fully resolved clade containing the insect-exclusive Drip and arthropod Prip-like clusters (Fig 3B). It appears that there were a series of expansions in the arthropod ancestor followed by a differential loss and independent duplications in the different arthropod lineages. The Prip-like pauropodan, symphylan and chilopodan myriapod classes retain two ancestral copies, while the origin of the second copy in diplopodan myriapods that retain more than one is more recent. Although previous studies have suggested that there has been an ancient whole-genome duplication (WGD) in Arachnopulmonata (Schwager et al. 2017), the Prip-like duplications that we recover do not suggest that as the origin of the Arachnopulmonata clades. On the contrary, the high clustering of all xiphosuran Prip-like duplicate sequences is indeed consistent with them being onhologs that originated after 3 rounds of WGD (Nong et al. 2021). Nevertheless, an independent phylogenetic tree we inferred using only arthropod sequences recovers a different topology, with hexapod Drip (including the previously-mentioned Drip insect-exclusive cluster and some additional early-splitting hexapod sequences) and Prip-like having emerged after the divergence of Pancrustacea from Myriapoda. However, this topology that suggests independent duplications in each of the three terrestrial arthropod groups is not supported by bootstrapping, and therefore these interrelationships should be taken with caution (Fig S4). In addition to Drip and Prip-like clades, it is noteworthy that the Aqp1-arth cluster (Fig 3) contains chelicerate and myriapod sequences, while it lacks pancrustacean sequences (except for some sequences from Malacostraca, Ostracoda, Cirripedia and a couple of earliest insect-splitting species -Fig S3). This loss of Aqp1-arth in most pancrustacea (especially in hexapods -i.e. the remaining arthropod group with an ancient terrestrialization event-) could be a consequence of the origin of the Drip cluster.

Functional studies in *D. melanogaster* have demonstrated the high efficiency of water transport in Malpighian tubule cells expressing Prip and Drip, as well as their importance in desiccation resistance (Cabrero et al. 2020), which may or may not be the case for arachnid and myriapod Prip-like duplicates.

Onychophorans, the only fully-terrestrial animal phyla, have three clusters of Aqp1-like sequences (Fig 3, in purple). One of them, Aqp1-onych2, harbours a gene duplication at the level of phylum, followed by several duplications and gene losses. However, their role in onychophoran physiology can only be hypothesised based on comparisons with other animals in the absence of functional studies.

Regarding annelids, we observe 6 different clusters named Aqp1-ann1 to Aqp1-ann6 (Fig 3, in yellow). Among them are the Aqp1-ann6, Aqp1-ann4 and Aqp1-ann2 clusters (Fig 3E, 3G and 3D, respectively). Aqp1-ann6 include a wide expansion in terrestrial Crassiclitellata (that include most terrestrial annelids, i. e. earthworms), and loss of ancient copies in non-Crassiclitellata annelids, putatively related to their terrestrialization (Fig 3E). In addition, we observe a duplication of Aqp1-ann4 in Clitellata with a subsequent loss of one sub-copy in secondarily marine clitellates (*Olavius algarvensis*), as shown in Fig 3G. Regarding Aqp1-ann2 (Fig 3D), there is an expansion at the level of Clitellata with the subsequent loss of all copies but one in non-Crassiclitelata annelids. The transition from a marine to a freshwater environment in clitelates (Erséus et al. 2020) may have been accompanied by the reshaping of their osmoregulatory capabilities putatively facilitated by this duplication. In the lack of functional studies of these aquaporins or the organs in which they are expressed, we can only suggest putative roles in annelid osmoregulation based on results from other organisms.

In molluscs, we find 6 supported clades (Fig 3, in indigo), which we have named Aqp1-mol1-5, Maqp and Mglp (Malacoaquaporins and Malacoglyceroporins, respectively, in accordance with previous research on the stylommatophoran snail *Helix pomatia* (Kosicka et al. 2016). As mentioned before, gastropods are the only molluscs that have colonised land. When looking at the mollusc-specific classical aquaporin clusters, we find three groups of special relevance. The first one is a Stylommatophora-specific duplication within Aqp1-mol1 (Fig 3F). The origin of this duplication dates back to right before the origin of Panpulmonata (Tenctipleura), with one copy being lost in all non-Stylommatophoran lineages. (Kosicka et al. 2016) identified the two duplicates of this clade in *H. pomatia*: HpAqp1 and HpAqp2. They reported the high expression of *HpAqp2* in the intestine, an organ whose main function is water reabsorption (Forester 1977). However, the expression of *HpAqp1* was not high in any

of the organs they examined. If we assume that *HpAqp2* and *HpAqp1* orthologues preserve the same function in the rest of stylommatophoran species studied here, the sub- or neofunctionalization of the stylommatophoran Aqp1-mol1 aquaporins may have contributed to the success of this order in its transition to land potentially by maximising water reabsorption in the intestine. Furthermore, we also report the expansion of Aqp1-mol2 in terrestrial Stylommatophora, freshwater panpulmonates, and the freshwater bivalves *Margaritifera margaritifera* and *Cristaria plicata* (Fig 3, Fig S3). This last expansion, together with the previously reported expansion of aquaporins in *D. rostriformis (Calcino et al. 2019)*, some of whose aquaporins we recover here as sister group to this expansion, and the reported functional role of a *D. rostriformis* Aqp1-like during embryonic development (Zieger et al. 2022), hints at the role aquaporins may have had during the sea-to-freshwater transition of bivalves. Moreover, an additional duplication within Maqp (Fig 3A) occurred before the split of Gastropoda and Scaphopoda, with only one copy being retained in Scaphopoda, and two within panpulmonate gastropods. The rest of gastropods Maqp are not clustered within this clade. One of these clades formed after the duplication contains the aestivation-related *HpAqp6*, which was mentioned above to be expressed in the foot during aestivation (Kosicka et al. 2020). This functional relationship to the drought-related behaviour evokes the putative role this pulmonate-specific duplication might have had on the panpulmonate terrestrialization events.

In contrast to Aqp1-like, not much is known about aquaamnioporins (Aqp8-like), except for their ability to transport water and ammonia and their expression in the inner membrane of mammalian mitochondria (Calamita et al. 2005; Lee and Thévenod 2006; Soria et al. 2010). Its relevant contribution to mitochondrial respiration (Ikaga et al. 2015)implies its importance for normal cell functioning and stress tolerance (Sokolova 2018), especially during the 'out of the sea' transition. We identified several lineage-specific duplications of Aqp8-like in *C. elegans*, annelids and molluscs (Fig S5). In the case of annelids both terrestrial and aquatic species show duplications within this aquaporin subtype. However, not all molluscan lineages showed duplications. For instance, gastropods contain only one copy of Aqp8-like, except for the terrestrial gastropod *Pomatias elegans*. In addition, while marine bivalves contain only 1 or 2 copies, freshwater bivalves contain 3 or even 4. These two examples point to the putative role of these expansions in their transition from the marine environment.

b) Aquaglycerolporins (Aqp3-like)

Compared to the phylogeny of Aqp1-like (Fig 3), where deuterostome aquaporins were nested within protostome aquaporins showing a shared origin, deuterostome and protostome Aqp3-like form two separate clades with independent evolutionary histories (Fig 4, Fig S6). Despite using a different methodology, our results of vertebrate Aqp3-like are mostly consistent with previous studies (Finn et al. 2014), except for the recovery of AQP10 and AQP7 as sister clades. Within protostomes, there is a further split into two clades, which we named Aqp3-prot1 and Aqp3-prot2. Aqp3-prot1 includes representatives of all invertebrate phyla included in this study. On the contrary, Aqp3-prot2 is mostly composed of arthropod sequences but includes as well some sequences from some mollusc and nematode species.

Zooming in the phylogenetic tree inferred for Aqp3-like, we recovered several ancient duplications in the nodes where a habitat transition occurred. Previous studies have reported independent duplications of Aqp3-like within chelicerates (Finn et al. 2015), myriapods and copepods (Catalán-García et al. 2021). However, their results could have been biassed due to the omission of other major arthropod lineages in their inferences. Here, we overcome this limitation by having representatives of virtually all arthropod orders analysed together. Our results pointed to an ancestral duplication that was independently retained in some groups and lost in others (Fig 4 and Fig 4C), instead of inferring independent within-lineage duplications. For instance, (Finn et al. 2015) report that two clades of Aqp3-like evolved in chelicerates based on their phylogeny with only spider, scorpion and acari sequences. When including the rest of the arachnid orders, such as xiphosurans and pycnogonids, results show that the duplication they reported is arachnid-specific (with xiphosurans nested within arachnids, Fig 4D) and derives from one of the two ancient copies of Aqp3-like that are present in arthropods, while having lost the other one. On the contrary, Pycnogonida, the earliest-splitting chelicerate lineage, has retained both copies. This is consistent with a single terrestrialization event in arachnids (considering a secondary later colonisation of water by xiphosurans as discussed in (Ballesteros et al. 2022)), where the duplication of Aqp3-like may have helped chelicerates to transition to a different niche. We could hypothesise that this new copy might have replaced the ancient Aqp3-like if it were neo- or subfunctionalized into a more efficient transporter, as reported with other aquaporins (Finn et al. 2015). These Aqp3-like copies subsequently underwent independent duplications in both clades, with two subclades in Xiphosura, Ricinulei, Opiliones and Arachnopulmonata (which we named Aqp3-araq1, not likely the result of their WGD; Fig. 4D), and two subclades in Acariformes, Scorpiones, Pseudoscorpiones and Araneae (which we named Aqp3-araq2, Fig 4D). Regarding myriapods, (Catalán-García et al. 2021) reported that Aqp3-like genes within Pauropoda, Symphyla and Diplopoda clustered in two separate clades, the ones that we

inferred in this study as ancient duplications. Symphyla and Diplopoda retain both ancient copies (named Aqp3-arth1 and Aqp3-arth2 in Fig 4C), while Chilopoda and Pauropoda only retain Aqp3-arth1. Last, but not least, we reproduce the pancrustacean results obtained by (Catalán-García et al. 2021) regarding the presence of two copepod Aqp3-like clusters, one of them as a sister group to Thecostraca and Arguloida Aqp3-like (in which we also report a close relationship to Collembola), and the other clustered within the rest of Pancrustacea (Fig 4C). The subfunctionalization of *Lepeophtheirus salmonis*' Aqp3-like in the enterocytes reported by (Catalán-García et al. 2021) may suggest a similar role during intestine water absorption in other arthropods. We also obtain an early-branching clade of Aqp3-arth1 composed of freshwater copepods and terrestrial diplurans, albeit recovered with low bootstrap support. Noteworthy, we also confirm that holometabolous insects have lost Aqp3-like sequences in favour of a potentially more efficient insect-specific Aqp1-like-derived glycerol transporter, as previously reported by (Finn et al. 2015). Moreover, we observe a Branchiopoda-specific duplication in Aqp3-arth1 (Fig 4C) that was also reported in previous studies (Finn et al. 2015; Stavang et al. 2015; Catalán-García et al. 2021). Branchiopoda are crustaceans of marine origin that have colonised brackish and freshwater environments (Oakley et al. 2013; Lozano-Fernandez et al. 2016). Thus, the presence of this duplication (with Thecostraca having lost one of the copies) reinforces our hypothesis of the role of aquaporins in osmoregulation.

Aqp3-like sequences from the other two members of Panarthropoda (Onychophora and Tardigrada) form a separate cluster from Arthropoda, with an intertwined evolutionary history. Fig 4B shows a shared duplication between Onychophora and Tardigrada, which suggests independent gene loss of both copies in arthropods. Each of these subclades underwent subsequent duplications, with an expansion within the onychophoran family Peripatopsidae. However, if the tardigrade copies have a species-specific or a much older origin cannot be pinpointed with the current taxonomic representation. Despite this, the role of *M. tardigradum* Aqp4 in anhydrobiosis has been previously put forward (Schokraie et al. 2012), hinting at the putative role onychophoran Aqp3-like may have in desiccation resistance and, therefore, in non-arthropod Panarthropoda terrestrialization.

Terrestrial annelids belong to the Clitellata class, characterised by the presence of a clitellum that secretes a cocoon. Cocoons are egg sacs that nourish the developing embryos, whose structure and composition differ between clitellates. Previous studies have shown that earthworm but not enchytraeid cocoons accumulate polyols when exposed to cold or drought (Anon 1995; Bauer et al. 2001). Thus, successive highly supported gene duplications of putative polyol transporters Aqp3-ann2 in Crassiclitellata, but not

enchytraeids (Fig 4E), suggest their putative role in this cocoon's physiological response to drought and cold. Resistance to drought and cold might have facilitated their transition to the heterogeneous environmental conditions of land. It is noteworthy that the exclusively terrestrial Aqp3-ann2 is clustered together with Aqp3-mol3, which contains aquaporins from panpulmonate gastropods (most of which are terrestrial), Caudofoveata and Cephalopoda. This phylogenetic relationship, in addition to the putative physiological role of Aqp3-ann2 mentioned before, suggests that Aqp3-mol3 may have also participated in the terrestrialization of Panpulmonata.

On the topic of molluscs, we report a duplication of Aqp3-mol2 in the freshwater-origin molluscan subclass Caenogastropoda, that was subsequently lost in the only marine species we included in our data set (*Hinea brasiliana*). The presence or absence of this duplication in other Caenogastropoda marine species remains to be studied, but the presence of both copies in the terrestrial *P. elegans* suggests that it was at least retained in the common ancestors of this species with other marine members of the group. Interestingly, contrary to Stylommatophora, *P. elegans* has not lost Aqp3-mol1.

Convergent evolution of Aqp1-like glycerol transporters in Panpulmonata and Insecta via gene duplication predates major evolutionary transitions

Major evolutionary transitions, such as the colonization of land habitats or the emergence of metamorphosis, resulted in the organisms needing new solutions to face new physiological and metabolic challenges. In both transitions, osmoregulation is, without a shadow of a doubt, one of these main challenges.

In addition to Aqp3-like, previous studies have proven the ability of some terrestrial-specific Aqp1-like proteins to transport glycerol. Such is the case of entomoglyceroporins, insect-specific Aqp1-like derived glycerol transporters (Finn et al. 2015), and AQP6, a tetrapod-specific classical aquaporin that can also transport glycerol (Holm et al. 2004). Moreover, the rise of entomoglyceroporins was accompanied by the loss of Aqp3-like in holometabolous insects. Here, we describe a similar pattern in molluscs: the rise of Aqp1-like malacoglyceroporins (Mglp, Fig 3C), and the lack of panpulmonate representatives in Aqp3-mol1 and Aqp3-mol2 (Fig 4). Mglps are a group of glycerol transporter aquaporins which have been previously described as mollusc-specific by (Kosicka et al. 2016). Notwithstanding, our larger taxonomic representation of molluscs restricts its specificity to gastropods, more specifically to Euthyneura. In addition, the clade contains a duplication at its root, right before the origin of Panpulmonata (Euthyneura),

gastropods that have mantle-derived lungs. Non-marine Panpulmonata are in fact the only ones that retain both copies after the duplication, suggesting their putative importance in their transition out of the sea. Previous studies located Aqp1-like and Aqp3-like protein expression in respiratory organs of vertebrates (Wittekindt and Dietl 2019) and insects (Yi et al. 2011), where they have a role in water excretion during respiration. Although no expression studies of gastropod aquaporins in pallial lungs have been carried out yet, malacoglyceroporins seem to be promising candidates for the functionality of this gastropod respiratory organ. Nevertheless, (Kosicka et al. 2020) reported the increased expression of *HpAqp3* (which clustered within the Mglp clade) and *HpAqp6* (Aqp8-like gene) in the foot during snail aestivation, one of the most relevant mechanisms of adaptation to humidity stress (Schweizer et al. 2019), indicating that Mglps may have a key role during aestivation, and therefore could have facilitated mollusc terrestrialization.

Regarding entomoglyceroporins, we observe that the most diverse holometabolous insects (e.g., Diptera, Coleoptera, Lepidoptera and Hymenoptera) contain two or more groups of entomoglyceroporins (Fig S7), compared to non-holometabolous insects. This expansion in holometabolous insects was suggested to be associated with the emergence of metamorphosis via the accumulation of polyols for cold and drought resistance (Finn et al. 2015). Further investigations including a denser taxonomic representation of insects will help shed light on this observation and pinpoint the exact origin of each duplication. Moreover, some species or groups of insects that colonised freshwater or the sea contain additional copies of Eglps (i.e. *Cloeon dipterum, Gerris buenoi* and *Anopheles gambiae* species and the Ephrydidae dipteran family). The secondarily marine *Anopheles merus* lacks Eglps, which can be the result of a secondary loss. Due to their expression in the Malpighian tubules (Cabrero et al. 2020), these duplications could be associated with osmoregulation, as suggested before for the inverse transition with Aqp1-like.

In the light of this, gene duplication of Aqp1-like glycerol transporters in Panpulmonata and Insecta may represent an example of convergent evolution that might have facilitated major evolutionary transitions (i.e., the 'out of sea' pulmonate gastropod transition and the origin of insect complete metamorphosis) through their involvement in osmoregulatory processes. Further genomic and functional studies will shed light on this hypothesis.

Conclusions

The transition of life from the sea to the land involved the acquisition of physiological osmoregulatory adaptations to avoid water loss and keep the osmotic homeostasis of internal fluids. The molecular function of aquaporins as a transporter of water and other molecules makes them an ideal system to start interrogating the evolution of osmoregulation during terrestrialization. Here, we provide phylogenetically-informed evolutionary evidence of the role that gene duplication and loss in Aqp-coding genes may have had during the evolution of terrestrial animal life. By comparing our phylogenetic results with previously described functional evidence, we were able to hypothesise in which physiological processes aquaporins might have contributed to the transition of the different animal phyla from the sea to freshwater or land, summarised in Table 1. Overall, our results suggest that ancient differential expansions of aquaporins in non-marine animals may have played a key role in the adaptation to life on land through the evolution of more sophisticated osmoregulatory mechanisms, such as the role of classical aquaporins (Aqp1-like) in molluscan aestivation or the putative role of aquaglycerolporins (Aqp3-like) in earthworms cocoons resistance to drought. Our results also show the complexity of aquaporin evolution in animals, highlighting the importance of taxon sampling when studying ancient gene families. All in all, our study sets the road towards a deeper understanding of one of the many pieces in the evolutionary puzzle of animal terrestrialization: the emergence and reshaping of osmoregulation.

Materials and methods

Obtaining genomic and transcriptomic data and habitat assignment

We analysed the aquaporin gene repertoire of 485 species from 7 animal phyla: Arthropoda, Onychophora, Tardigrada, Nematoda, Mollusca, Annelida and Vertebrata. The full list of species included and the source they were obtained from is available at Supp. Data S2.

Data from the remaining 415 invertebrate species were obtained from MATEdb (Fernández et al. 2022), which provides standard comparable proteomes for phylogenomic studies. We downloaded the gene annotation of those species (longest isoform for each gene), and their taxonomic and habitat information. Vertebrate, *M. tardigradum*, and *C. elegans* aquaporins were directly obtained from the literature (Huang et al. 2007; Grohme et al. 2013; Finn et al. 2014) respectively). The only exceptions were the aquaporins from the amphibian *Leptobrachium leishanense*, for which we downloaded the genome annotation (Cunningham et al. 2021). In the case of annelids, (Mucciolo et al. 2021) provided a fairly good representation of aquaporins across Pleistoannelida, the most abundant clade within annelids. However, as noted by the authors, the methodology followed could have underestimated the identification of aquaporins due to the incompleteness or low quality of the source databases' data. For instance, An example of this is *Eisenia andrei,* for which only one Aqp was identified. To verify this information, we downloaded the genome annotation of *E. andrei* (Shao et al. 2020) to annotate its Aqp-coding genes ourselves as described below. We identified a total of 27 Aqp-coding genes, in contrast with the one reported by (Mucciolo et al. 2021).

Habitat information for species was obtained from the World Register of Marine Species (WoRMS) database (WoRMS Editorial Board 2022). A species was considered to be aquatic when it spent at least one stage of its life cycle in an aquatic environment.

Annotating genes encoding for aquaporins in downloaded proteomes

From the whole set of proteomes, genes encoding for aquaporins were annotated using BITACORA version 1.3 (Vizueta et al. 2020) in "protein mode". Briefly, BITACORA combines BLAST and HMMER (Eddy 2011) searches against a custom curated database of genes belonging to the gene family of interest. Due to the conservation of aquaporins across

all animals, our aquaporin curated database contained a list of 123 metazoan aquaporin sequences that have been experimentally confirmed in previous studies (Supp. Data 3). To benchmark this methodology, we used *Drosophila melanogaster* to compare the number of aquaporins identified using BITACORA with the number of aquaporins annotated in its genome, obtaining the same result.

Phylogenetic tree inference of aquaporin-coding genes

As we needed to know which aquaporins belonged to which subtype (i.e., classical aquaporins - Aqp1-like, aquaammoniaporins - Aqp3-like, unorthodox aquaporins - Aqp8-like - or superaquaporins - Aqp11-like) for posterior analyses, we first independently classified aquaporins of all animal species included in this study, except for the ones that had already been classified in previous publications (Finn et al. 2014; Mucciolo et al. 2021). We based this classification on the description of aquaporin phylogenetic clusters and their relative position in the phylogenetic tree to the already classified sequences, which were considered as a backbone.

For arthropods, onychophorans and molluscs (i.e. lineages whose aquaporins we identified de novo in this study) we manually curated the classification by checking unclustered aquaporin sequences and long branches that could represent outliers using BLASTP against the non-redundant NCBI database (NCBI nr). This was done to reclassify identified aquaporins that had wrongly fallen into the wrong subtype and remove contaminants from species outside of the studied phylum that had escaped filtering in the original data sets. In total, 41 arthropod and mollusc sequences were removed from the initial set.

To build all phylogenetic trees, amino acid sequences were aligned using PASTA v1.8.6 (Mirarab et al. 2015). The alignments included 202 sequences previously described as aquaporins (Finn et al. 2014; Finn et al. 2015) to use as a backbone for identifying aquaporin subfamilies in our data set (Supp. Data 4). In addition, 3 previously identified malacoglyceroporins and malacoaquaporin (Kosicka et al. 2016) were also included when building aquaporin phylogenetic trees. All resultant alignments were trimmed using trimAL v1.4.1 (Capella-Gutiérrez et al. 2009). We tested different trimming methods and gap thresholds to select the best one in terms of keeping fewer alignment positions to reduce computational time while keeping biologically informative sites (Fig S8). Final alignments were trimmed with a gap threshold of 0.01.

We created guiding trees for each of the alignments using the LG model in FastTree v2.1.11 (Price et al. 2010), which were later used as starting trees for the ML inference of aquaporin trees with IQ-TREE v2.1.2 (Minh et al. 2020). We applied an LG protein matrix with a profile mixture model of 10 classes (C10), plus an additional class of empirical amino acid profile (F) and rate heterogeneity across sites using a discrete Gamma model with 4 categories (G). We also included ultrafast bootstrap (UFBoot) (Minh et al. 2013) and SH-aLRT test (Guindon et al. 2010) to obtain support values of resultant trees. The AqpM sequence of *Methanococcus maripaludis* (NCBI accession NP_988083), included in the backbone, was always used as an outgroup for tree rooting, as suggested in previous studies (Finn et al. 2014).

To avoid long branch attraction (LBA) due to their higher divergence and because our preliminary results using all aquaporins showed that they formed a monophyletic clade, entomoglyceroporins were analysed separately and excluded from the phylogeny that combined all animal Aqp1-like sequences.

We visualised the trees using iTOL (Letunic and Bork 2021) and created the final figures with the R packages ggtree (Yu et al. 2017) and ggtreeExtra (Yu et al. 2021). A full list of packages used is included in the code available on GitHub (see Data availability section).

Inference of a combined dated species tree

Time-scaled ultrametric trees were inferred independently for arthropods, molluscs and annelids by using a node-dating approach on species trees that followed the most recent topologies (Richter et al. 2001; Colgan et al. 2007; Lindgren 2010; Kocot et al. 2011; Smith et al. 2011; Wiegmann et al. 2011; Kocot et al. 2013; Misof et al. 2014; González et al. 2015; Razkin et al. 2015; Salvi and Mariottini 2016; Uribe et al. 2016; Kocot et al. 2017; Peters et al. 2017; Uribe and Zardoya 2017; Wang et al. 2017; Fernández et al. 2018; Johnson et al. 2018; Pabst and Kocot 2018; Schwentner et al. 2018; Yang et al. 2018; Cunha and Giribet 2019; Kawahara et al. 2019; Kocot et al. 2019; Lee et al. 2019; Lozano-Fernandez et al. 2019; McKenna et al. 2019; Doğan et al. 2020; Erséus et al. 2020; Kocot et al. 2020; Sun et al. 2020; Anderson and Lindgren 2021; Kallal et al. 2021; Moles and Giribet 2021; Ontano et al. 2021; Smith 2021; Ballesteros et al. 2022). Full details for the dating analysis of arthropods and molluscs are provided elsewhere (Aristide et al. in prep; Tonzo et al. in prep). (*Note: for reviewing purposes, see details at the end of the manuscript*).

In the case of annelids, we dated the species tree using the Aqp1-like gene sequences in MCMCtree (Yang 2007). For that, we first inferred the annelid Aqp1-like gene tree using aquaporins annotated in (Mucciolo et al. 2021) and our inferred aquaporins for *E. andrei*, as our preliminary analyses showed that the aquaporin phylogeny is broadly concordant with the species tree topology at the level of aquaporin subclusters. Second, from this gene tree, we derived 16 sets of 1:1 orthologues with at least four species represented using PhyloPyPruner (Thálen 2019), considering the phylogenetic interrelationships of annelid species as described in (Erséus et al. 2020). These orthologous sequences were then concatenated into a single matrix of 2,694 sites for their analysis and split into two partitions based on their evolutionary rate (measured as the sum of branch lengths divided by the number of tips in the respective gene tree). Prior distributions for node ages (see Supplementary Information in Github) were based on (Chen et al. 2020) for the annelid MRCA and in the divergence time intervals and fossil calibrations in (Erséus et al. 2020). Species *Romanchella perrieri* was removed by PhyloPyPruner and was therefore not included in the tree.

Vertebrate and onychophoran time trees were obtained from the literature (Irisarri et al. 2017; Baker et al. 2021) and pruned to keep only a selection of 13 vertebrate species whose aquaporins have previously been studied (Finn et al. 2014) and the 4 onychophorans (or closely-related species) that were included in MATEdb (Fernández et al. 2022), respectively.

Finally, the ultrametric trees of all 5 phyla were grafted into a manually-built backbone topology depicting the relationships among them using the RRphylo R package (Castiglione et al. 2018). Divergence times for the ancestral nodes in the backbone tree were obtained from (Dohrmann and Wörheide 2017).

Phylogenetic comparative methods to explore aquaporin evolution

In order to test whether there were differences in the mean total number of aquaporins between terrestrial and aquatic species, we implemented a phylogenetic generalised linear model using the phyloglm R package (Ho and Ané 2014). Briefly, we fitted a phylogenetic Poisson regression model to our data describing the relationship between the number of aquaporins and habitat, accounting for the phylogenetic history of the species. For habitat, we classified species in two different ways: marine vs non-marine (i.e., freshwater, brackish and terrestrial), and terrestrial vs non-terrestrial (i.e., freshwater, brackish and marine). This classification allowed us to identify shifts in aquaporin number

during marine to freshwater and terrestrial transitions, and marine and freshwater to terrestrial transitions. We tested for differences in each of the 4 main subtypes of aquaporins and in all aquaporins together.

Data availability

All supplementary information and code are included in the Github repository: https://github.com/MetazoaPhylogenomicsLab/Martinez_Redondo_et_al_2022_aquaporins

Author contribution

GIMR and RF conceived the study. GIMR performed the analyses and designed the figures. GIMR and CSG annotated aquaporins. PBG and VT assisted aquaporin annotation. LA designed the Poisson regression analysis. LA and VT inferred the ultrametric trees. LA and GIMR performed the Poisson regression analysis. GIMR and RF interpreted the results. GIMR and RF wrote the first version of the manuscript. RF provided resources and supervised the study. All authors revised and approved the final version of the manuscript.

Acknowledgements

GIMR acknowledges the support of Secretaria d'Universitats i Recerca del Departament d'Empresa i Coneixement de la Generalitat de Catalunya and European Social Fund (ESF) 'Investing in your future' (grant 2021 FI_B 00476) and the CSIC 2020 JAE Intro programme. LA acknowledges funding from a Juan de la Cierva-Formación (grant agreement FJC2019-042184-I funded by MCIN/AEI/10.13039/501100011033). PBG was supported by an FPI grant (grant agreement no. BES-2017-081050) financed by MCIN/AEI /10.13039/501100011033 and by ESF 'Investing in your Future'. RF acknowledges support from the following sources of funding: Ramón y Cajal fellowship (grant agreement no. RYC2017-22492 funded by MCIN/AEI /10.13039/501100011033 and ESF 'Investing in your future'), Agencia Estatal de Investigación (project PID2019-108824GA-I00 funded by MCIN/AEI/10.13039/501100011033) and the European Research Council (this project has received funding from the European Research Council (ERC) under the European's Union's Horizon 2020 research and innovation programme (grant agreement no. 948281)). We also thank Centro de Supercomputación de Galicia (CESGA) for access to computer resources, and particularly Pablo Rey for his assistance and guidance. We thank Julien Clavel for helping us with the poisson regression analyses.

References

Abascal F, Irisarri I, Zardoya R. 2014. Diversity and evolution of membrane intrinsic proteins. *Biochim. Biophys. Acta* 1840:1468–1481.

Anderson FE, Lindgren AR. 2021. Phylogenomic analyses recover a clade of large-bodied decapodiform cephalopods. *Mol. Phylogenet. Evol.* 156:107038.

Anon. 1995. Polyol accumulation in earthworm cocoons induced by dehydration. *Comp. Biochem. Physiol. A Physiol.* 111:251–255.

Baker CM, Buckman-Young RS, Costa CS, Giribet G. 2021. Phylogenomic analysis of velvet worms (Onychophora) uncovers an evolutionary radiation in the Neotropics. *Mol. Biol. Evol.* 38:5391–5404.

Ballesteros JA, Santibáñez-López CE, Baker CM, Benavides LR, Cunha TJ, Gainett G, Ontano AZ, Setton EVW, Arango CP, Gavish-Regev E, et al. 2022. Comprehensive species sampling and sophisticated algorithmic approaches refute the monophyly of Arachnida. *Mol. Biol. Evol.* 39:msac021.

Bauer R, Worland MR, Block W. 2001. Experimental studies on cold survival of enchytraeid cocoons. *Pedobiologia*. 46:561-571.

Benoit JB, Hansen IA, Szuter EM, Drake LL, Burnett DL, Attardo GM. 2014. Emerging roles of aquaporins in relation to the physiology of blood-feeding arthropods. *J. Comp. Physiol. B*. 184:811–825.

Bestetti S, Galli M, Sorrentino I, Pinton P, Rimessi A, Sitia R, Medraño-Fernandez I. 2020. Human aquaporin-11 guarantees efficient transport of H_2O_2 across the endoplasmic reticulum membrane. *Redox Biol*. 28:101326.

Cabrero P, Terhzaz S, Dornan AJ, Ghimire S, Holmes HL, Turin DR, Romero MF, Davies SA, Dow JAT. 2020. Specialized stellate cells offer a privileged route for rapid water flux in *Drosophila* renal tubule. *Proc. Natl. Acad. Sci. U.S.A.* 117:1779–1787.

Calamita G, Ferri D, Gena P, Liquori GE, Cavalier A, Thomas D, Svelto M. 2005. The inner mitochondrial membrane has aquaporin-8 water channels and is highly permeable to water. *J. Biol. Chem.* 280:17149–17153.

Calcino AD, de Oliveira AL, Simakov O, Schwaha T, Zieger E, Wollesen T, Wanninger A. 2019. The quagga mussel genome and the evolution of freshwater tolerance. *DNA Res.* 26:411–422.

Capella-Gutiérrez S, Silla-Martínez JM, Gabaldón T. 2009. trimAl: a tool for automated alignment trimming in large-scale phylogenetic analyses. *Bioinformatics* 25:1972–1973.

Carrete Vega G, Wiens JJ. 2012. Why are there so few fish in the sea? *Proc. Biol. Sci.* 279:2323–2329.

Castiglione S, Tesone G, Piccolo M, Melchionna M, Mondanaro A, Serio C, Di Febbraro M, Raia P. 2018. A new method for testing evolutionary rate variation and shifts in phenotypic evolution. *Methods Ecol. Evol.* 9:974–983.

Catalán-García M, Chauvigné F, Stavang JA, Nilsen F, Cerdà J, Finn RN. 2021. Lineage-level divergence of copepod glycerol transporters and the emergence of isoform-specific trafficking regulation. *Commun Biol* 4:643.

Chen H, Parry LA, Vinther J, Zhai D, Hou X, Ma X. 2020. A Cambrian crown annelid reconciles phylogenomics and the fossil record. *Nature* 583:249–252.

Christoph K, Beck F-X, Neuhofer W. 2007. Osmoadaptation of Mammalian cells - an orchestrated network of protective genes. *Curr. Genomics* 8:209–218.

Colgan DJ, Ponder WF, Beacham E, Macaranas J. 2007. Molecular phylogenetics of Caenogastropoda (Gastropoda: Mollusca). *Mol. Phylogenet. Evol.* 42:717–737.

Colgan DJ, Santos RP. 2018. A phylogenetic classification of gastropod aquaporins. *Mar. Genomics* 38:59–65.

Cunha TJ, Giribet G. 2019. A congruent topology for deep gastropod relationships. *Proc. Royal Soc. B.* 286:20182776.

Cunningham F, Allen JE, Allen J, Alvarez-Jarreta J, Amode MR, Armean IM, Austine-Orimoloye O, Azov AG, Barnes I, Bennett R, et al. 2021. Ensembl 2022. *Nucleic Acids Res.* 50:D988–D995.

Doğan Ö, Schrödl M, Chen Z. 2020. The complete mitogenome of Moquin-Tandon, 1855 (Gastropoda: Stylommatophora): mitochondrial genome architecture, evolution and phylogenetic considerations within Stylommatophora. *PeerJ* 8:e8603.

Dohrmann M, Wörheide G. 2017. Dating early animal evolution using phylogenomic data. *Sci. Rep.* 7:3599.

Eddy SR. 2011. Accelerated Profile HMM Searches. *PLoS Comput. Biol.* 7:e1002195.

Erséus C, Williams BW, Horn KM, Halanych KM, Santos SR, James SW, des Châtelliers MC, Anderson FE. 2020. Phylogenomic analyses reveal a Palaeozoic radiation and support a freshwater origin for clitellate annelids. *Zool. Scr.* 49:614–640.

Fernández R, Edgecombe GD, Giribet G. 2018. Phylogenomics illuminates the backbone of the Myriapoda Tree of Life and reconciles morphological and molecular phylogenies. *Sci. Rep.* 8:83.

Fernández R, Tonzo V, Guerrero CS, Lozano-Fernandez J, Martínez-Redondo GI, Balart-García P, Aristide L, Eleftheriadi K, Vargas-Chávez C. 2022. MATEdb, a data repository of high-quality metazoan transcriptome assemblies to accelerate phylogenomic studies. Available from: http://dx.doi.org/10.1101/2022.07.18.500182

Finn RN, Cerdà J. 2011. Aquaporin evolution in fishes. *Front. Physiol.* 2:44.

Finn RN, Cerdà J. 2015. Evolution and functional diversity of aquaporins. *Biol. Bull.* 229:6–23.

Finn RN, Chauvigné F, Hlidberg JB, Cutler CP, Cerdà J. 2014. The lineage-specific evolution of aquaporin gene clusters facilitated tetrapod terrestrial adaptation. *PLoS One* 9:e113686.

Finn RN, Chauvigné F, Stavang JA, Belles X, Cerdà J. 2015. Insect glycerol transporters evolved by functional co-option and gene replacement. *Nat. Commun.* 6:7814.

Forester AJ. 1977. The function of the intestine in the pulmonate mollusc Helix pomatia L. *Experientia* 33:465–467.

Gil CI, Jarius M, von Kries JP, Rohlfing A-K. 2017. Neuronal Chemosensation and Osmotic

Stress Response Converge in the Regulation of aqp-8 in C. elegans. *Front. Physiol.* 8:380.

Goedert J, Lécuyer C, Amiot R, Arnaud-Godet F, Wang X, Cui L, Cuny G, Douay G, Fourel F, Panczer G, et al. 2018. Euryhaline ecology of early tetrapods revealed by stable isotopes. *Nature* 558:68–72.

González VL, Andrade SCS, Bieler R, Collins TM, Dunn CW, Mikkelsen PM, Taylor JD, Giribet G. 2015. A phylogenetic backbone for Bivalvia: an RNA-seq approach. *Proc. Biol. Sci.* 282:20142332.

Grohme MA, Mali B, Wełnicz W, Michel S, Schill RO, Frohme M. 2013. The aquaporin channel repertoire of the tardigrade *Milnesium tardigradum*. *Bioinform. Biol. Insights* 7:BBI.S11497

Guindon S, Dufayard J-F, Lefort V, Anisimova M, Hordijk W, Gascuel O. 2010. New algorithms and methods to estimate Maximum-Likelihood Phylogenies: assessing the performance of PhyML 3.0. *Syst. Biol.* 59:307–321.

Holm LM, Klaerke DA, Zeuthen T. 2004. Aquaporin 6 is permeable to glycerol and urea. *Pflugers Arch.* 448:181–186.

Ho L si T, Ané C. 2014. A linear-time algorithm for Gaussian and non-Gaussian trait evolution models. *Syst. Biol.* 63:397–408.

Holterman M, Schratzberger M, Helder J. 2019. Nematodes as evolutionary commuters between marine, freshwater and terrestrial habitats. *Biol. J. Linn. Soc.* 128:756–767.

Huang CG, Lamitina T, Agre P, Strange K. 2007. Functional analysis of the aquaporin gene family in*Caenorhabditis elegans*. *Am. J. Physiol. Cell Physiol.* 292:C1867–C1873.

Ikaga R, Namekata I, Kotiadis VN, Ogawa H, Duchen MR, Tanaka H, Iida-Tanaka N. 2015. Knockdown of aquaporin-8 induces mitochondrial dysfunction in 3T3-L1 cells. *Biochem Biophys Rep* 4:187–195.

Irisarri I, Baurain D, Brinkmann H, Delsuc F, Sire J-Y, Kupfer A, Petersen J, Jarek M, Meyer A, Vences M, et al. 2017. Phylotranscriptomic consolidation of the jawed vertebrate timetree. *Nat. Ecol. Evol.* 1:1370–1378.

Ishibashi K, Tanaka Y, Morishita Y. 2021. The role of mammalian superaquaporins inside the cell: An update. *Biochim. Biophys. Acta - Biomembr.* 1863:183617.

Jamy M, Biwer C, Vaulot D, Obiol A, Jing H, Peura S, Massana R, Burki F. Global patterns and rates of habitat transitions across the eukaryotic tree of life. Available from: http://dx.doi.org/10.1101/2021.11.01.466765

Johnson KP, Dietrich CH, Friedrich F, Beutel RG, Wipfler B, Peters RS, Allen JM, Petersen M, Donath A, Walden KKO, et al. 2018. Phylogenomics and the evolution of hemipteroid insects. *Proc. Natl. Acad. Sci. U.S.A.* 115:12775–12780.

Kallal RJ, Kulkarni SS, Dimitrov D, Benavides LR, Arnedo MA, Giribet G, Hormiga G. 2021. Converging on the orb: denser taxon sampling elucidates spider phylogeny and new analytical methods support repeated evolution of the orb web. *Cladistics* 37:298–316.

Kameda Y, Kato M. 2011. Terrestrial invasion of pomatiopsid gastropods in the heavy-snow region of the Japanese Archipelago. *BMC Evol. Biol.* 11:118.

Kawahara AY, Plotkin D, Espeland M, Meusemann K, Toussaint EFA, Donath A, Gimnich F, Frandsen PB, Zwick A, Dos Reis M, et al. 2019. Phylogenomics reveals the evolutionary timing and pattern of butterflies and moths. *Proc. Natl. Acad. Sci. U. S. A.* 116:22657–22663.

Kocot KM, Cannon JT, Todt C, Citarella MR, Kohn AB, Meyer A, Santos SR, Schander C, Moroz LL, Lieb B, et al. 2011. Phylogenomics reveals deep molluscan relationships. *Nature* 477:452–456.

Kocot KM, Halanych KM, Krug PJ. 2013. Phylogenomics supports Panpulmonata: Opisthobranch paraphyly and key evolutionary steps in a major radiation of gastropod molluscs. *Mol. Phylogenet. Evol.* 69:764–771.

Kocot KM, Poustka AJ, Stöger I, Halanych KM, Schrödl M. 2020. New data from Monoplacophora and a carefully-curated dataset resolve molluscan relationships. *Sci. Rep.* 10.

Kocot KM, Struck TH, Merkel J, Waits DS, Todt C, Brannock PM, Weese DA, Cannon JT, Moroz LL, Lieb B, et al. 2017. Phylogenomics of Lophotrochozoa with Consideration of Systematic Error. *Syst. Biol.* 66:256–282.

Kocot KM, Todt C, Mikkelsen NT, Halanych KM. 2019. Phylogenomics of Aplacophora (Mollusca, Aculifera) and a solenogaster without a foot. *Proc. Royal Soc. B.* 286:20190115.

Kondrashov FA. 2012. Gene duplication as a mechanism of genomic adaptation to a changing environment. *Proc. Biol. Sci.* 279:5048–5057.

Kosicka E, Grobys D, Kmita H, Lesicki A, Pieńkowska JR. 2016. Putative new groups of invertebrate water channels based on the snail Helix pomatia L. (Helicidae) MIP protein identification and phylogenetic analysis. *Eur. J. Cell Biol.* 95:543–551.

Kosicka E, Lesicki A, Pieńkowska JR. 2020. Molluscan aquaporins: an overview, with some notes on their role in the entry into aestivation in gastropods. *Molluscan Res.* 40:101–111.

Kreida S, Törnroth-Horsefield S. 2015. Structural insights into aquaporin selectivity and regulation. *Curr. Opin. Struct. Biol.* 33:126–134.

Krug PJ, Caplins SA, Algoso K, Thomas K, Valdés ÁA, Wade R, Wong NLWS, Eernisse DJ, Kocot KM. 2022. Phylogenomic resolution of the root of Panpulmonata, a hyperdiverse radiation of gastropods: new insight into the evolution of air breathing. *Proc. Biol. Sci.* 289:20211855.

Kruse E, Uehlein N, Kaldenhoff R. 2006. The aquaporins. *Genome Biol.* 7:206.

Laforenza U, Scaffino MF, Gastaldi G. 2013. Aquaporin-10 represents an alternative pathway for glycerol efflux from human adipocytes. *PLoS One* 8:e54474.

Lamitina ST, Morrison R, Moeckel GW, Strange K. 2004. Adaptation of the nematode Caenorhabditis elegans to extreme osmotic stress. *Am. J. Physiol. Cell Physiol.* 286:C785–C791.

Lee H, Chen W-J, Puillandre N, Aznar-Cormano L, Tsai M-H, Samadi S. 2019. Incorporation of deep-sea and small-sized species provides new insights into gastropods phylogeny. *Mol. Phylogenet. Evol.* 135:136–147.

Lee W-K, Thévenod F. 2006. A role for mitochondrial aquaporins in cellular life-and-death decisions? *Am. J. Physiol. Cell Physiol.* 291:C195–C202.

Letunic I, Bork P. 2021. Interactive Tree Of Life (iTOL) v5: an online tool for phylogenetic tree display and annotation. *Nucleic Acids Res.* 49:W293–W296.

Lindgren AR. 2010. Molecular inference of phylogenetic relationships among Decapodiformes (Mollusca: Cephalopoda) with special focus on the squid order Oegopsida. *Mol. Phylogenet. Evol.* 56:77–90.

Little C. 1990. The terrestrial invasion: an ecophysiological approach to the origins of land animals. Cambridge University Press

Lorente-Martínez H, Agorreta A, Torres-Sánchez M, Mauro DS. 2018. Evidence of positive selection suggests possible role of aquaporins in the water-to-land transition of mudskippers. *Org. Divers. Evol.* 18:499–514.

Lozano-Fernandez J, Carton R, Tanner AR, Puttick MN, Blaxter M, Vinther J, Olesen J, Giribet G, Edgecombe GD, Pisani D. 2016. A molecular palaeobiological exploration of arthropod terrestrialization. *Philos. Trans. R. Soc. Lond. B Biol. Sci.* 371.

Lozano-Fernandez J, Giacomelli M, Fleming JF, Chen A, Vinther J, Thomsen PF, Glenner H, Palero F, Legg DA, Iliffe TM, et al. 2019. Pancrustacean evolution illuminated by taxon-rich genomic-scale data sets with an expanded remipede sampling. *Genome Biol. Evol.* 11:2055–2070.

de Maré SW, Venskutonytė R, Eltschkner S, de Groot BL, Lindkvist-Petersson K. 2020. Structural Basis for Glycerol Efflux and Selectivity of Human Aquaporin 7. *Structure* 28:215–222.e3.

McKenna DD, Shin S, Ahrens D, Balke M, Beza-Beza C, Clarke DJ, Donath A, Escalona HE, Friedrich F, Letsch H, et al. 2019. The evolution and genomic basis of beetle diversity. *Proc. Natl. Acad. Sci. U. S. A.* 116:24729–24737.

Minh BQ, Nguyen MAT, von Haeseler A. 2013. Ultrafast approximation for phylogenetic bootstrap. *Mol. Biol. Evol.* 30:1188–1195.

Minh BQ, Schmidt HA, Chernomor O, Schrempf D, Woodhams MD, von Haeseler A, Lanfear R. 2020. IQ-TREE 2: New models and efficient methods for phylogenetic inference in the genomic era. *Mol. Biol. Evol.* 37:1530–1534.

Mirarab S, Nguyen N, Guo S, Wang L-S, Kim J, Warnow T. 2015. PASTA: ultra-large multiple sequence alignment for nucleotide and amino-acid sequences. *J. Comput. Biol.* 22:377–386.

Misof B, Liu S, Meusemann K, Peters RS, Donath A, Mayer C, Frandsen PB, Ware J, Flouri T, Beutel RG, et al. 2014. Phylogenomics resolves the timing and pattern of insect evolution. *Science* 346:763–767.

Moles J, Giribet G. 2021. A polyvalent and universal tool for genomic studies in gastropod molluscs (Heterobranchia). *Mol. Phylogenet. Evol.* 155:106996.

Mucciolo S, Desiderato A, Salonna M, Mamos T, Prodocimo V, Di Domenico M, Mastrototaro F, Lana P, Gissi C, Calamita G. 2021. Finding aquaporins in annelids: an evolutionary analysis and a case study. *Cells* 10:3562.

Nesverova V, Törnroth-Horsefield S. 2019. Phosphorylation-dependent regulation of

mammalian aquaporins. *Cells* 8.

Nong W, Qu Z, Li Y, Barton-Owen T, Wong AYP, Yip HY, Lee HT, Narayana S, Baril T, Swale T, et al. 2021. Horseshoe crab genomes reveal the evolution of genes and microRNAs after three rounds of whole genome duplication. *Commun Biol* 4:83.

Nozaki K, Ishii D, Ishibashi K. 2008. Intracellular aquaporins: clues for intracellular water transport? *Pflugers Arch.* 456:701–707.

Oakley TH, Wolfe JM, Lindgren AR, Zaharoff AK. 2013. Phylotranscriptomics to bring the understudied into the fold: monophyletic Ostracoda, fossil placement, and pancrustacean phylogeny. *Mol. Biol. Evol.* 30:215–233.

Ontano AZ, Gainett G, Aharon S, Ballesteros JA, Benavides LR, Corbett KF, Gavish-Regev E, Harvey MS, Monsma S, Santibáñez-López CE, et al. 2021. Taxonomic sampling and rare genomic changes overcome long-branch attraction in the phylogenetic placement of pseudoscorpions. *Mol. Biol. Evol.* 38:2446–2467.

Pabst EA, Kocot KM. 2018. Phylogenomics confirms monophyly of Nudipleura (Gastropoda: Heterobranchia). *J. Molluscan Stud.* 84:259–265.

Peng J, Zhou Y, Jia H, Li L, Qian J, Han F, Yin H, Cui Y. 2018. Transcriptomics-based identification of aquaporin diversity in the house dust mite Dermatophagoides farinae (Acariformes: Pyroglyphidae). *J. Insect Sci.* 18.

Peters RS, Krogmann L, Mayer C, Donath A, Gunkel S, Meusemann K, Kozlov A, Podsiadlowski L, Petersen M, Lanfear R, et al. 2017. Evolutionary history of the Hymenoptera. *Curr. Biol.* 27:1013–1018.

Price MN, Dehal PS, Arkin AP. 2010. FastTree 2 – Approximately Maximum-Likelihood trees for large alignments. *PLoS one* 5:e9490.

Qvarnström M, Szrek P, Ahlberg PE, Niedźwiedzki G. 2018. Non-marine palaeoenvironment associated to the earliest tetrapod tracks. *Sci. Rep.* 8.

Rankin JC, Davenport J. 1981. Animal osmoregulation. New York: Wiley.

Razkin O, Gómez-Moliner BJ, Prieto CE, Martínez-Ortí A, Arrébola JR, Muñoz B, Chueca LJ, Madeira MJ. 2015. Molecular phylogeny of the western Palaearctic Helicoidea (Gastropoda, Stylommatophora). *Mol. Phylogenet. Evol.* 83:99–117.

Richter, Richter, Scholtz. 2001. Phylogenetic analysis of the Malacostraca (Crustacea). *J. Zool. Syst. Evol. Res.* 39:113–136.

Rota-Stabelli O, Daley AC, Pisani D. 2013. Molecular timetrees reveal a Cambrian colonization of land and a new scenario for ecdysozoan evolution. *Curr. Biol.* 23:392–398.

Salvi D, Mariottini P. 2016. Molecular taxonomy in 2D: a novel ITS2 rRNA sequence-structure approach guides the description of the oysters' subfamily Saccostreinae and the genus *Magallana* (Bivalvia: Ostreidae). *Zool. J. Linn. Soc.* 179

Schokraie E, Warnken U, Hotz-Wagenblatt A, Grohme MA, Hengherr S, Förster F, Schill RO, Frohme M, Dandekar T, Schnölzer M. 2012. Comparative proteome analysis of Milnesium tardigradum in early embryonic state versus adults in active and anhydrobiotic state. *PLoS One* 7:e45682.

Schwager EE, Sharma PP, Clarke T, Leite DJ, Wierschin T, Pechmann M, Akiyama-Oda Y, Esposito L, Bechsgaard J, Bilde T, et al. 2017. The house spider genome reveals an ancient whole-genome duplication during arachnid evolution. *BMC Biol.* 15:62.

Schweizer M, Triebskorn R, Köhler H-R. 2019. Snails in the sun: Strategies of terrestrial gastropods to cope with hot and dry conditions. *Ecol. Evol.* 9:12940–12960.

Schwentner M, Richter S, Rogers DC, Giribet G. 2018. Tetraconatan phylogeny with special focus on Malacostraca and Branchiopoda: highlighting the strength of taxon-specific matrices in phylogenomics. *Proc. Biol. Sci.* 285.

Shao Y, Wang X-B, Zhang J-J, Li M-L, Wu S-S, Ma X-Y, Wang X, Zhao H-F, Li Y, Zhu HH, et al. 2020. Genome and single-cell RNA-sequencing of the earthworm Eisenia andrei identifies cellular mechanisms underlying regeneration. *Nat. Commun.* 11:2656.

Sharma PP. 2017. Chelicerates and the Conquest of Land: A View of Arachnid Origins Through an Evo-Devo Spyglass. *Integr. Comp. Biol.* 57:510–522.

Smith CH. 2021. A high-quality reference genome for a parasitic bivalve with doubly uniparental inheritance (Bivalvia: Unionida). *Genome Biol. Evol.* 13.

Smith SA, Wilson NG, Goetz FE, Feehery C, Andrade SCS, Rouse GW, Giribet G, Dunn CW. 2011. Resolving the evolutionary relationships of molluscs with phylogenomic tools. *Nature* 480:364–367.

Sokolova I. 2018. Mitochondrial Adaptations to Variable Environments and Their Role in Animals' Stress Tolerance. *Integr. Comp. Biol.* 58:519–531.

Soria LR, Fanelli E, Altamura N, Svelto M, Marinelli RA, Calamita G. 2010. Aquaporin-8-facilitated mitochondrial ammonia transport. *Biochem. Biophys. Res. Commun.* 393:217–221.

Spring JH, Robichaux SR, Hamlin JA. 2009. The role of aquaporins in excretion in insects. *J. Exp. Biol.* 212:358–362.

Stavang JA, Chauvigné F, Kongshaug H, Cerdà J, Nilsen F, Finn RN. 2015. Phylogenomic and functional analyses of salmon lice aquaporins uncover the molecular diversity of the superfamily in Arthropoda. *BMC Genomics* 16:618.

van Straalen NM. 2021. Evolutionary terrestrialization scenarios for soil invertebrates. *Pedobiologia* 87-88:150753.

Sun X, Yu D, Xie Z, Dong J, Ding Y, Yao H, Greenslade P. 2020. Phylomitogenomic analyses on collembolan higher taxa with enhanced taxon sampling and discussion on method selection. *PLoS One* 15:e0230827.

Takvam M, Wood CM, Kryvi H, Nilsen TO. 2021. Ion transporters and osmoregulation in the kidney of teleost fishes as a function of salinity. *Front. Physiol.* 12:664588.

Thálen F. 2019. PhyloPyPruner. Available at https://github.com/fethalen/phylopypruner.

Thorat LJ, Gaikwad SM, Nath BB. 2012. Trehalose as an indicator of desiccation stress in Drosophila melanogaster larvae: a potential marker of anhydrobiosis. *Biochem. Biophys. Res. Commun.* 419:638–642.

Tingaud-Sequeira A, Calusinska M, Finn RN, Chauvigné F, Lozano J, Cerdà J. 2010. The zebrafish genome encodes the largest vertebrate repertoire of functional aquaporins

with dual paralogy and substrate specificities similar to mammals. *BMC Evol. Biol.* 10:38.

Uchiyama M, Konno N. 2006. Hormonal regulation of ion and water transport in anuran amphibians. *Gen. Comp. Endocrinol.* 147:54–61.

Uribe JE, Colgan D, Castro LR, Kano Y, Zardoya R. 2016. Phylogenetic relationships among superfamilies of Neritimorpha (Mollusca: Gastropoda). *Mol. Phylogenet. Evol.* 104:21–31.

Uribe JE, Zardoya R. 2017. Revisiting the phylogeny of Cephalopoda using complete mitochondrial genomes. *J. Molluscan Stud.* 83:133–144.

Van Ekert E, Chauvigné F, Finn RN, Mathew LG, Hull JJ, Cerdà J, Fabrick JA. 2016. Molecular and functional characterization of Bemisia tabaci aquaporins reveals the water channel diversity of hemipteran insects. *Insect Biochem. Mol. Biol.* 77:39–51.

Vermeij GJ, Watson-Zink VM. 2022. Terrestrialization in gastropods: lineages, ecological constraints and comparisons with other animals. *Biol. J. Linn. Soc.* 136:393-404.

Vizueta J, Sánchez-Gracia A, Rozas J. 2020. bitacora: A comprehensive tool for the identification and annotation of gene families in genome assemblies. *Mol. Ecol. Resour.* 20:1445–1452.

Wang Z, Shi Y, Qiu Z, Che Y, Lo N. 2017. Reconstructing the phylogeny of Blattodea: robust support for interfamilial relationships and major clades. *Sci. Rep.* 7:3903.

Watson-Zink VM. 2021. Making the grade: Physiological adaptations to terrestrial environments in decapod crabs. *Arthropod Struct. Dev.* 64:101089.

Wiegmann BM, Trautwein MD, Winkler IS, Barr NB, Kim J-W, Lambkin C, Bertone MA, Cassel BK, Bayless KM, Heimberg AM, et al. 2011. Episodic radiations in the fly tree of life. *Proc. Natl. Acad. Sci. U. S. A.* 108:5690–5695.

Wittekindt OH, Dietl P. 2019. Aquaporins in the lung. *Pflug. Arch. Eur. J. Physiol.* 471:519–532.

WoRMS Editorial Board. 2022. World Register of Marine Species. Available from https://www.marinespecies.org at VLIZ. Last accessed 2022-05-03.

Yang H, Zhang J-E, Xia J, Yang J, Guo J, Deng Z, Luo M. 2018. Comparative characterization of the complete mitochondrial genomes of the three apple snails (Gastropoda: Ampullariidae) and the phylogenetic analyses. *Int. J. Mol. Sci.* 19.

Yang Z. 2007. PAML 4: phylogenetic analysis by maximum likelihood. *Mol. Biol. Evol.* 24:1586–1591.

Yilmaz O, Chauvigné F, Ferré A, Nilsen F, Fjelldal PG, Cerdà J, Finn RN. 2020. Unravelling the complex duplication history of deuterostome glycerol transporters. *Cells* 9:1663.

Yi S-X, Benoit JB, Elnitsky MA, Kaufmann N, Brodsky JL, Zeidel ML, Denlinger DL, Lee RE Jr. 2011. Function and immuno-localization of aquaporins in the Antarctic midge Belgica antarctica. *J. Insect Physiol.* 57:1096–1105.

Yu G, Smith DK, Zhu H, Guan Y, Lam TT. 2017. ggtree : an r package for visualization and annotation of phylogenetic trees with their covariates and other associated data. *Methods Ecol. Evol.* 8:28–36.

Yu G, Xu S, Dai Z, Guo P, Fu X, Liu S, Zhou L, Tang W, Feng T, Chen M, et al. 2021. ggtreeExtra: Compact visualization of richly annotated phylogenetic data. *Mol. Biol. Evol.* 38:4039-4042.

Zieger E, Schwaha T, Burger K, Bergheim I, Wanninger A, Calcino AD. 2022. Midbody-localized aquaporin mediates intercellular lumen expansion during early cleavage of an invasive freshwater bivalve. *Front. Cell Dev. Biol.* 10:894434.

Figures and Tables

Figure 1. **Terrestrialization and aquaporins. A.** Schematic representation of the main terrestrialization events in the phyla included in this study and their habitat of origin. Arthropods and onychophorans have mostly colonised land from a marine ancestor; some terrestrial gastropods and Crassiclitellata annelids (earthworms) from a freshwater ancestor; while tetrapods and other terrestrial molluscs (such as Stylommatophora) have a euryhaline and mangrove/intertidal origin, respectively. In the case of hexapods, their terrestrialization has been suggested to have been mediated through an intermediate subterranean state, represented by a cave in the figure, due to their relatedness with the subterranean aquatic Remipedia. **B.** Aquaporin tertiary and quaternary structure and representation of transmembrane domains adapted from (Nesverova and Törnroth-Horsefield 2019). Water molecules are represented in red, and the two half-helices formed by loops B and E are in yellow. **C.** Schematic representation of pore size and structure changes allow different molecules to pass through. Molecules are ordered from the smallest to the largest (water, ammonia, and glycerol, respectively) from left to right. **D.** General aquaporin phylogeny based on previous studies.

Figure 2. **Phylogenetic comparative methods to understand aquaporin evolution. A.** Ultrametric calibrated tree of the 483 species from 5 phyla included in the analyses. **B.** Phylogenetic Poisson regression results for marine (dark blue) vs non-marine (yellow) comparison in all animals, arthropods and molluscs in the four aquaporin subtypes (i.e. Aqp1-like, Aqp3-like, Aqp8-like and Aqp11-like). Significant results are highlighted in bold with an asterisk and inside a square. **C.** Phylogenetic Poisson regression results for terrestrial (dark orange) vs aquatic (blue) comparison in all animals, arthropods and molluscs in the four aquaporin subtypes (i.e. Aqp1-like, Aqp3-like, Aqp8-like and Aqp11-like). Significant results are highlighted in bold with an asterisk and inside a square. **D.** Phylogenetic Poisson regression results for marine (dark blue) vs non-marine (yellow) in all animals, arthropods and molluscs in all aquaporins. Significant results are highlighted in bold with an asterisk and inside a square. **E.** Phylogenetic Poisson regression results for terrestrial (dark orange) vs aquatic (blue) in all animals, arthropods and molluscs in all aquaporins.

Figure 3. **Consensus phylogenetic tree of 2,202 Aqp1-like plus 63 backbone sequences in 6 studied phyla. A.** Both copies of malacoaquaporins (Maqp) in Panpulmonata. **B.** Prip-like and Drip clades in arthropods. **C.** Malacoglyceroporins' duplication in Panpulmonata. **D**. Aqp1-ann2 expansion within Crassiclitellata. **E.** Aqp1-ann6 clade expansion in Crassiclitellata. **F.** Aqp1-mol1 duplication in Stylommatophora. **G.** Aqp1-ann4 subclade in Clitellata. Duplications are marked with a green dot. Bootstrap values are included only in nodes with support values higher than 70.

Figure 4. Consensus phylogenetic tree of 1,108 Aqp3-like plus 68 backbone sequences in the 7 studied animal phyla. A. Aqp3-mol2 clade in Caenogastropoda shows a duplication in this freshwater-origin subclass. **B.** Aqp3-like paralogs in Onychophora and Tardigrada. **C.** Aqp3-arth1 and Aqp3-arth2. **D.** Aqp3-araq1 and Aqp3-araq2 arachnid paralogs. **E.** Aqp3-ann2 expansions within Crassiclitellata. Duplications are marked with a green dot. Bootstrap values are included only in nodes with support values higher than 70.

Table 1. The role of aquaporin evolution in animal terrestrialization. Summary table of the results obtained in this paper. All chordate results (marked with a) are proposed by previous studies (Finn et al. 2014; Lorente-Martínez et al. 2018). The putative function of positively selected Aqp10a (Aqp3-like) in mudskippers' gills (marked with b) is proposed here based on the differential gene expression of its ortholog Aqp10a and paralog Aqp10b in zebrafish in gills (Finn and Cerdà 2011), and the enlarged gill chambers mudskippers have whose function is related to their semiterrestrial lifestyle (i.e. they close the gill chambers when exposed to air to keep gills moist). Nevertheless, no functional studies have looked at Aqps in mudskippers' gills. Information from the phylum Nematoda is not included as only one terrestrial species was included in our analyses.

	Aqp1-like		Aqp3-like		Aqp8-like		Aqp11-like	
Phylum[a]	Evidence	Putative role in terrestrialization	Evidence	Putative role in terrestrialization	Evidence	Putative role in terrestrialization	Evidence	Putative role in terrestrialization
Arthropoda	Independent duplications	Increased efficiency in excretion	Ancestral duplication and differential retention	Water absorption in intestine		-	Reduction in terrestrial species	Redox homeostasis in ER
Onychophora	Expansion	Unknown	Ancient duplication and independent lineage-specific expansions	Resistance to desiccation		-		-
Tardigrada	-			Anhydrobiosis		-		-
Annelida	Duplication in Clitellata and loss in secondarily marine species Duplication in Clitellata and loss in non-Crassiclitellata species Expansions in terrestrial annelids	Sea-to-freshwater transition that predated terrestrialization of Enchytreida and Crassiclitellata	Duplication in Crassiclitellata	Accumulation of polyols in cocoons when exposed to cold or drought		-		-
Mollusca	Evolution of glycerol transporters and duplication in Stylommatophora Duplication before Panpulmonata and retention in Stylommatophora	Aestivation Water absorption in the intestine	Duplication in Caenogastropoda	Sea-to-freshwater transition that predated terrestrialization of Pomatiidae	Expansion in Panpulmonata	Aestivation, redox stress		-
Chordata[b]	Tetrapod-specific aquaporins	Evolution of water conservation mechanisms	Positive selection in mudskippers	Gill function?[c]		-	Positive selection in mudskippers	Redox homeostasis in ER

[a] Information from the phylum Nematoda is not discussed as only one terrestrial species was included in our analyses.

[b] Chordate results are proposed by previous studies (Finn et al. 2014; Lorente-Martínez et al. 2018).

[c] The putative function of positively selected Aqp10a (Aqp3-like) in mudskippers' gills is proposed here based on the differential gene expression of its ortholog Aqp10a and paralog Aqp10b in zebrafish in gills (Finn and Cerdà 2011), and the enlarged gill chambers mudskippers have whose function is related to their semiterrestrial lifestyle (i.e. they close the gill chambers when exposed to air to keep gills moist). Nevertheless, no functional studies have looked at Aqps in mudskippers' gills.