(Formatted for VACCINES) -Research Article

Title: Survivin (BIRC5) peptide vaccine in the 4T1 murine mammary tumor model: a potential neoadjuvant T-cell immunotherapy for triple negative breast cancer.

Authors: Scott R Burkholz, MS1†, Charles V Herst, PhD1†, Richard T Carback, PhD1, Paul E Harris PhD1,2, Reid M Rubsamen, MD1,3,4

Affiliations

1. Flow Pharma Inc., Warrensville Heights, Ohio

2.Vagelos College of Physicians and Surgeons, Columbia University, New York, New York.

3. University Hospitals, Cleveland Medical Center, Cleveland, Ohio

4. Case Western Reserve School of Medicine, Cleveland, Ohio

†These authors contributed equally to this work.

Address correspondence to:

Reid M Rubsamen, MD, MS, MHCM

University Hospitals, Cleveland Medical Center

Department of Anesthesia and Perioperative Medicine

11100 Euclid Avenue, Cleveland, Ohio 44016

Email: reid.rubsamen@uhhospitals.org

Telephone: 925 708 8000

Abstract

A triple negative breast cancer (TNBC) model using the murine 4T1 tumor cell line was used to explore the efficacy of an adjuvanted survivin (SVN)-peptide microparticle vaccine using tumor growth as the outcome metric. We first performed tumor cell dose titration studies to determine a tumor cell dose that resulted in sufficient tumor takes but minimal morbidity/mortality within the required study period. Later, in a second cohort of mice, the SVN-peptide microparticle vaccine was administered via intraperitoneal injection at study start with a second dose given 14 days later. An orthotopic injection of 4T1 cells into the mammary tissue was performed on the same day as the administration of the second vaccine dose. The mice were followed for up to 41 days with subcutaneous measurements of tumor volume made every 3-4 days. Vaccination with SVN peptides was associated with a peptide antigen-specific γIFN Elispot response in the murine splenocyte population but was absent from the control microparticle group. At end of study, we found that vaccination with adjuvanted SVN-peptide microparticles resulted in statistically significant slower primary tumor growth rates in BALB/c mice challenged with 4T1 cells relative to the control peptide-less vaccination group. These studies suggest that T-cell immunotherapy specifically targeting survivin might be an applicable neoadjuvant immunotherapy therapy for TNBC. More pre-clinical studies and clinical trials will be needed to explore this concept further.

Keywords: Breast Cancer, Immunotherapy, Bioinformatics, Next-Generation Sequencing, Neoadjuvant Therapy, tumor-associated antigen, survivin, BIRC5,

Introduction

The survivin (SVN), also known as BIRC5, protein is an inhibitor of apoptosis and is overexpressed in many malignancies, including breast cancer stem cells and breast tumor tissue [1], relative to adjacent normal adult cells and tissues (reviewed in [2] and [3]). These observations suggest that SVN might be an ideal tumor cell target. In fact the NCI declared SVN as a research priority more than twenty years ago[4]. Since then, SVN as a target of humoral and cellular immunity has been studied in animal models and in cancer patients. The structure of many immunogenic B cell and T cell epitopes of SVN have been characterized [2,5-8] and applied to the development of synthetic survivin-peptide cancer vaccines. These studies have provided proof of principle data supporting several currently active human immunotherapy/vaccine clinical trials targeting SVN in glioblastoma, neuroendocrine tumors, ovarian cancer, and hormone receptor positive breast cancer (e.g., NCT05163080, NCT02334865, NCT04895761).   In glioblastoma, immunotherapy targeting survivin has provided significant prolongation of overall survival compared with standard radiation therapy and chemotherapy [9].

Estrogen, progesterone receptor negative, HER2 "negative" breast cancer (TNBC) accounts for up to 15%-20% of all breast malignancies, has a generally worse prognosis and fewer treatment options relative to more common forms of breast cancer. Current standard of care treatment options for recurrent TNBC have been limited to conventional cytotoxic chemotherapeutic agents due the lack of expression of molecular targets found on the more common types of breast malignancies. Adjuvant immunotherapy for TNBC is currently being explored and the approaches fall into several diverse categories (reviewed in [10]). One approach, recently approved by the FDA, combines chemotherapy with immune checkpoint inhibitors (ICI) with the goal of boosting existing adaptive immune response to breast tumor antigens. However, immunohistochemistry studies show that PD-L1 expression, one target of ICI therapy, is limited to about 20% of TNBC [11]. Clinical and histochemical data has confirmed that efficacy of anti PD-L1 mAb ICI as neoadjuvant and adjuvant treatment for TNBC patients whose tumors and/or stromal cells (including tumor infiltrating lymphocytes) express PD-L1. While the PD-L1 positive subgroup show better overall survival, the PD-L1 negative subgroup

faired equally poorly, regardless of anti-PD-L1 treatment[12], suggesting that additional approaches to adjuvant immunotherapy need to exploration.

Adjuvant vaccine-based immunotherapy of TBNC has been recently reviewed [13]. Out of 42 clinical studies reviewed, only one limited exploratory human study[14] has targeted SVN in TBNC. With the goal of developing novel immunotherapies for TNBC, we turned to the 4T1 murine model of TNBC [15] to evaluate the therapeutic potential of a synthetic SVN peptide based microparticle vaccine for adjuvant immunotherapy of TNBC. We previously developed an adjuvanted, synthetic peptide - poly (lactic-co-glycolic acid) (PLGA) copolymer microparticle vaccine platform [16] capable of inducing robust therapeutic CD8+ cytotoxic T lymphocyte responses to viral antigens presented by MHC Class I molecules in non-human primate [17] and murine models [18]. In this preclinical study, we report on the effects of vaccination with adjuvanted survivin-peptide loaded microspheres on orthotopically implanted tumor growth in a 4T1 murine model of TBNC.

Materials and Methods

Characterization of survivin expression in 4T1 cells and normal mouse mammary tissue.

To fully characterize SVN in the TNBC model mouse 4T1 cell line (ATCC, Manassas, VA), frozen 4T1 cell samples were sent to Complete Genomics (Beijing, China) for sequencing. DNA was extracted from the snap-frozen cryopreserved 4T1 cell line, and BALB/c mouse tails. Sample DNA was prepared using the Agilent SureSelect XT Mouse All Exon Kit (Agilent Technologies, Santa Clara, CA). RNA was extracted from the frozen 4T1 cell line and from also from snap-frozen, normal mammary tissue samples harvested from BALB/c mice. RNA samples were prepared for mRNA sequencing via poly-A tail capture with MGI Tech Company reagents (MGI, Shenzhen, China). Samples were sequenced by Complete Genomics on the BGISEQ-500 (BGI, Beijing, China) at 100 base pair paired end reads. DNA read coverage for normal tissue and tumor was 100x and 300x respectively, with RNA read count at 80 million paired end reads. To process the data, FASTP was used to perform quality control and adapter trimming and BWA-MEM aligned reads to the GRCm38 reference mouse genome [19-22]. The germline sequences were confirmed to match the peptides identified for vaccine administration. RNA expression of survivin was examined by adapter trimming and quality control with FASTP

[19], followed by pseudoalignment with Kallisto on the GENCODE v25 mouse transcriptome for quantification in Sleuth expressed as transcripts per million [21].

Vaccine Design/Peptide Selection

The overall strategy and rationale for the selection of synthetic peptides used to stimulate potential CTL immune responses to SVN targets presented by murine 4T1 cells has been previously described[16,18,23]. Briefly, we identified survivin peptide antigens potentially capable of stimulating MHC Class I and Class II restricted, tumor-specific T-cell response in BALB/c mice. These peptide sequences were determined by reviewing previous publications describing SVN specific MHC Class I restricted T cells response using various peptide vaccine formulations [24,25]. In addition, NetMHCIIpan and NetMHCII [26] were utilized to identify QP19, a region predicted to bind to class II MHC to stimulate CD4 helper T-cells. The sequences of these murine class I H-2 K, D, and L binding peptides and murine class II H-2 I-E and I-A binding peptides are given in Table 1.

Table 1

Epitope Name	Peptide Sequence	MHC Class	Position	MHC Match Screening Method
AL9	ATFKNWPFL	I	20-28	[26,27]
AM9	AFLTVKKQM	I	85-93	[25]
GI9	GWEPDDNPI	I	66-74	[24,25]
TI9	TAKTTRQSI	I	127-135	[25]
QP19	QIWQLYLKNYRIATFKNWP	I/II	8-26	[26,28]
PADRE	AKFVAAWTLKAAA	2	N/A	[29,30]

Table 1 legend. Literature references and prediction tools supporting a BALB/c MHC match to each of the six peptides microencapsulated into the adjuvanted microsphere vaccine platform described here

Vaccine Manufacture

A blend of PLGA microspheres, prepared as described in a previous publication [17], was manufactured containing individual synthetic GMP-grade peptides (Peptides International, Louisville, KY, USA) , selected from the primary amino acid sequence of the murine survivin protein (Table 1). The adjuvanted microsphere vaccine formulation contained 3 µg/mg of survivin-specific class 1 and class 2 peptides and 0.5 µg/mg of the TLR-9 oligonucleotide agonist CpG (ODN-1018) (Trilink Biosciences, San Diego, CA, USA). The microspheres, approximately 6 microns in diameter, were delivered in a 200 µl volume of PBS / Tween 20 (0.01% v/v) injectate solution containing the TLR-4 agonist MPLA (Avanti Polar Lipids, Alabaster, AL, USA) at a concentration of 100 µg/ml.

Post vaccination evaluation of peripheral T cell responses

Enzyme-linked immunosorbent spot for interferon gamma (ELISpot) assays were performed with BALB/c splenocytes obtained from control and vaccinated surviving tumor bearing mice on day 41 after 4T1 tumor inoculation. Splenocytes were prepared as previously described [31]. The peptide antigens used in the ELISpot assays were the same as those used in the peptide vaccine and individually added to ELISpot wells at 10 µg/ml final concentration. ELISpot assay plates were prepared and processed as per the manufacturer's instructions (3321-4HPT-10, Mabtech Inc., Cincinnati, OH). ELISpots were enumerated by machine (CTL S6 Entry M2, Shaker Heights, Cleveland, OH) to calculate the frequency of gamma interferon producing T-cells in the splenocyte populations.

Animal Immunizations

Previous studies with the adjuvanted microsphere platform have shown that T-cell expansion capable of providing protection against viral challenge is present 14 days in mouse models [16,18]. A 4T1 inoculation dose ranging study was undertaken to find the maximum 4T1 cell count that would show limited tumor growth during the study and allow evaluation of the vaccine's possible efficacy, as shown in the study design schematic in supplementary materials Figure S1. Based on the 4T1 dose ranging data shown in supplementary materials

Figure S2, the challenge study was designed with one cohort receiving a 250 4T1 cell inoculation dose and the second cohort receiving 500 cells of orthotopically injected 4T1 breast cancer cells. Ten of the animals in each cohort received two doses of intra-peritoneally delivered adjuvanted peptide microspheres (2.5 mg/dose), and the control mice were given two doses of blank microspheres (2.5 mg/dose, control microspheres; without peptide antigen and adjuvants) given both fourteen days before 4T1 cell inoculation, and again at the same time as orthotopic injection of 4T1cells. Subcutaneous tumor volumes were measured every three to four days with a 41-day endpoint after implantation as shown in the study design schematic in Figure 1.

Figure 1

Figure 1 legend. Study design schematic for a BALB/c mouse model 4T1 TNBC tumor challenge microsphere peptide vaccine efficacy study.

Mouse tumor volume was measured non-invasively every three to four days with a micrometer applied to the tumor growing in the subcutaneous space. The tumor volume (TV) was expressed in mm^3 using the modified ellipsoid formula: TV = ½(length x width^2)[32,33]. This data was used to calculate the tumor growth rate expressed as Δ mm^3/day. Tumor-take frequencies were defined as the number of mice with measurable tumors on the indicated day divided by the total number of mice inoculated with 4T1 cells.

Results

Wild-type Survivin mRNA was highly expressed in all 4T1 cell line samples studied and was seen only at very low background transcription levels in the BALB/c normal mammary tissue samples (Figure 2).

Figure 2

Figure 2 legend. Expression level from mRNA-Seq of mouse wild-type Survivin transcript (ENST0000081387)[34] in normal breast tissue and the 4T1 cell line used in this study. The mean transcripts per millions (TPM) are shown by black bars. The significance of the difference between the average TPM found normal murine breast tissue and 4T1 cells was calculated by the method of Student.

Of the six peptides loaded into the adjuvanted microsphere formulation shown in Table 1, only QP19 (QIWQLYLKNYRIATFKNWP), produced a positive ELISpot response as shown in Figure 3. The mice who were not vaccinated did not produce a detectable response to the survivin QP19 peptide antigen. Although published studies suggested that the administered MHC class I peptide epitopes were immunogenic in BALB/c mice, we observed that only QP19 produced a T-cell response as measured by ELISpot.

Figure 3

Figure 3 legend. ELISpot response to QP19 in splenocytes harvested from vaccinated and unvaccinated mice that received 250 cells of 4T1. Statistical significance of the difference in the average number of ELISpots in vaccinated and control groups was determined using the unpaired, non-parametric, Mann-Whitney T-test.

The inoculation dose of 4T1 at both the 250 and 500 cell levels did not result in a tumor-take frequency of 100%, but were similar to previous reports [35] and the tumor-take frequencies measured in the dose ranging study (Figure S2). However, vaccination with survivin peptide antigens was associated with statistically significant slower primary 4T1 mammary tumor growth rates compared to tumors in control mice (Figure 4).

This effect was particularly evident in the 500 4T1 cells dose group, but only at later time points / tumor volumes in the 250 4T1 cell inoculum dose group. We observed γ IFN ELISpot responses to peptide QP19 in 9/10 vaccinated mice, and noted that of these 10 mice only two mice developed growing tumors.

Figure 4

Figure 4 Legend. Comparison of average tumor growth rates (vaccinated versus control) for the 500 cell and 250 cell 4T1 tumor inoculation challenge groups. The sample size was n=10 for each group (Open diamonds- SVN microsphere vaccinated mice, Filled circles – control mice). The mean tumor growth rates (Red Bars – SVN microsphere vaccinated mice, Black Bars-Control mice) were calculated from growing tumors only. The statistical significance of the differences in growth rates (only non-zero data was considered) for each tumor

inoculum size and treatment group (i.e., control vs. SVN peptide microsphere) was assessed using ANCOVA with Tukeys HSD and excel software according to the general linear model with time treated as a covariate. For either experiment, 500 cells or 250 cells, treatment (vaccination or control) was the significant variable (p<0.0001and p<0.009, respectively) at the 95% confidence interval.

Discussion

The targeting of survivin peptides that are HLA-matched to the host for evoking a cell-mediated immune response to kill tumor cells is of particular interest. MHC-restricted responses to peptides located on the survivin protein have been shown to elicit an immune response, including immunotherapies targeting survivin in a number of clinical trials [4]. A comprehensive list of various approaches used clinically to elicit a host immune response against survivin-expressing tumors is shown ins1s1 Table S1. A collection of HLA-restricted survivin peptide antigens identified across these various studies also raises the possibility of a broadly applicable immunotherapy for tumors that express survivin. As these studies also illustrate, eliciting a reliable, clinically significant immune response to peptide antigens is challenging.

Ensuring that the correct peptide sequence is selected and delivered effectively for T-cell expansion to occur have been obstacles to the development of safe and effective targeted immunotherapies. For example, small peptides injected on their own, even when combined with adjuvants known to enhance a T-cell response, have not been shown to trigger particularly robust T cells response [16]. As we describe here, microspheres can be manufactured that encapsulate potentially immunogenic peptides and the TLR-9 agonist CpG in a biodegradable PLGA polymer, delivered after reconstitution in a saline solution with the TLR-4 agonist MPLA to produce a cellular immune response to the administered peptide antigens as demonstrated by ELISpot [16].

In this study, we found that multiple putative peptide antigens, derived from the primary sequence of survivin and predicted to bind to MHC Class I molecules of BALB/c mice, did not elicit a detectable *ex-vivo* immune response. The QP19 peptide antigen, predicted to bind to I-A^d/I-E^d, the MHC Class II molecules of

BALB/c, however, was able to elicit an *ex-vivo* gamma-IFN T-cell ELISpot response and this response was associated with slower tumor growth rates. This observation suggests that protective T-cell responses were vaccine-induced and operant during tumor growth.

One possible explanation for this observation would be the presence of one or more CD8+ epitopes co-localized within the QP19 nineteen-mer, producing the observed T-cell response and anti-tumor growth activity associated with vaccination. Analysis of all possible overlapping peptides of 8-9 amino acids in length within QP19 using NetMHC and NetMHCpan found 4 potential BALB/c MHC matched peptide antigens as listed in Table S2 [26,27]. The identification of possible survivin-protective CD8+ T cell epitopes demonstrated by *ex-vivo* ELISpot response, and the formal demonstration of CD4+ T cell helper activity evoked by vaccination with QP19 await further experimentation. If such CD8+ T cell epitopes within QP19 do indeed exist, they maybe immunodominant relative to the other MHC Class I peptide SVN antigens used in these experiments. Immunodominant CD8+ T-clones may prevail over subdominant clones, masking their response. The use of anti PD-1 mAbs have been shown to promote epitope spreading in the context of anti-tumor CD8+ T cell responses [36] and certainly suggest a potential role for ICIs in SVN peptide microsphere vaccination for TNBC immunotherapy.

The use of immunotherapy for breast cancer has gained attention recently [37]. Tumor associated antigens, in contrast to neoantigens, provide the opportunity to develop immunotherapy targeting a fixed set of peptides with collective HLA restrictions predicted to provide broad population coverage that could be administered to breast cancer patients without the need for patient-specific tumor gene sequencing and manufacturing of personalized immunotherapy. Targeted T-cell immunotherapy triggering a T-cell immune response against survivin as neoadjuvant therapy has the potential to reduce recurrence after surgical excision of breast tumor tissue if the number of cells remaining after tumor debulking is low enough to result in CTL attack sufficient to blunt tumor-take and tumor growth rate. Previous studies have seen mixed efficacy with unprotected peptides used as immunotherapy[4,38]. A delivery system such as the adjuvanted microsphere

encapsulation described herein, may be able to effectively deliver peptides to produce T-cell expansion against

TAA   targets   such   as   survivin   expressed   by   TNBC   patients.

Supplementary Materials

Table S1

Cancer Type	Clinical Stage	Peptide	HLA	Delivery	Results	Citations
Glioblastoma	Phase II	DLAQCFFMFKELEGW	A*02, A*03, A*24	A peptide mutated from the wild type sequence conjugated to the adjuvant keyhole limpet haemocyanin, injected subcutaneous 4 times biweekly along with Montanide ISA 51 with sargramostim.	Increased survivin-specific IgG antibodies and CD8+ T-cells generated in majority of patients with some efficacy for PFS and OS along with temozolomide.	[9,39]
Multiple	Phase II	AYACNTSTL	A*24:02	Subcutaneous injection of peptide at multiple intervals, 14 days apart; addition of IFN-α or IFN-β in later studies; addition of IFA in later studies	Increased levels of peptide specific CTL in a portion of patients, but adjuvants are added due to lack of clinical efficacy, however, progression free survival was not increased.	[7,40-42]
Melanoma	Phase II	LMLGEFLKL, and a peptide from IDO	A*02	Multiple subcutaneous injection of peptide mutated from survivin and indoleamine 2,3-dioxygenase with Montanide ISA 51. Topical 5% imiquimod cream and GM-CSF to assist in delivery and response.	The majority of patients had an immune response, with an increase of memory CD4+ and CD8+. No significant results seen, but tumor regression in one individual.	[43]
Solid Tumors	Phase I	FTELTLGEF, LMLGEFLKL, RISTFKNWPK, STFKNWPFL, LPPAWQPFL	A*01, A*02, A*03, A*24, B*7	Subcutaneous injection of peptide, some mutated from wild-type, multiple times with Montanide ISA 51.	Survivin-specific T-cells in the majority of patients, with most of those not detected before vaccination. Clinical results not evaluated.	[44]
Melanoma	Phase II	FTELTLGEF, LMLGEFLKL, EPDLAQCFY	A*01, A*02, B*35	Deep Subcutaneous injection of 3 peptides differing from wild type sequence along with Montanide ISA 51. Different regimen used for vaccination frequency and addition of cyclophosphamide.	Increase survivin-specific CD8+ T-cells in the majority of individuals. Significant extension seen in overall survival.	[45]
Prostate	Phase I/II	TLGEFLKLDRERAKN, TLPPAWQPFL,	DRB1*0X, A*02	Peptides subcutaneously injected or loaded onto	Several HLA-DRB1 present this peptide for a Th1 CD4+ response,	[46,47]

		ELTLGEFLKL, and multiple non-survivin class 1 and class 2 peptides		autologous DC at intervals with one of the following: Imiquimod, GM☐CSF, local hyperthermia or the TLR☐7/8 agonist mRNA/protamine complex.	however, one patient had an anaphylactic reaction to this peptide after multiple doses.	

Table S1 legend. Clinical trial list detailing survivin peptide immunotherapy studies conducted over the last two decades for a variety of tumor types.

Table S2

Epitope Name	Peptide Sequence	Class	Position
LI8	LYLKNYRI	1	12-19
LA9	LYLKNYRIA	1	12-20
KF8	KNYRIATF	1	15-22
LF9	LKNYRIATF	1	14-21

Table S2 legend. NetMHC and NetMHCpan were used to scan all possible overlapping peptides of 8-9 amino acids in length within QP19 (32,42). The four potential BALB/c MHC matched class I peptide antigens found within QP19 are listed here.

Figure S1

4T1 Cell Orthotopic Inoculation Dose Escalation Study Design

Tumor Volume Measurements on Days 7, 11, 14, 18, 21, 25, and 28 After Innoculation

10,000 cells / 20 µl

7500 cells / 20 µl

1500 cells / 20 µl

750 cells / 20 µl

500 cells / 20 µl

Inoculation of 4T1 Cells into Mammary Fat Pad

Analysis of Tumor Growth Kinetics

Tumor Volume (mm3)

Time (days)

Supplementary Figure 1 legend. Study design schematic for 4T1 TNBC cell line inoculation dose escalation study. Varying numbers of 4T1 tumor cells were injected into mouse mammary tissue.

Figure S2

Supplementary Figure 2 legend. Average growth rate curves shown after injection of escalating numbers of

4T1 TNBC cancer cells into BALB/c mouse mammary tissue at T0. Sample size was n=10 mice per group.

Funding

This research did not receive any specific external grant from funding agencies.

Acknowledgements

A pre-print of this manuscript was posted on BioRxiv.

Author Contributions

All authors made substantial contributions to: (1) the conception and design of the study (R.M.R., S.B., C.V.H., and P.E.H.), the acquisition of data (R.M.R., R.C. and S.B.), or the analysis and interpretation of data (P.E.H., R.M.R., S.B. and C.V.H.); and (2) drafting the article or revising it critically for important intellectual content (P.E.H., S.B., C.V.H., R.C. and R.M.R.). All authors have read and agreed to the published version of the manuscript.

Institutional Review Board Statement

The study was conducted according to the guidelines of the Declaration of Helsinki and approved by the Institutional Animal Care and Use Committee (IACUC) of GemPharmatech Co,Ltd. (Nanjing, Jiangsu) (protocol code: SYXK (SU) 2018-0027), approval date 08/05/2019). The care and use of animals was conducted in accordance with the regulations of the Association for Assessment and Accreditation of Laboratory Animal Care (AAALAC) and the National Institutes of Health guidelines.

Data Availability Statement

DNA and RNA-seq data can be accessed at NIH SRA under accession # PRJNA868747.

Conflicts of Interest:

R.M.R., S.B., R.C., P.E.H. and C.V.H. are employees of Flow Pharma, Inc. compensated in cash and stock, and are named inventors on various issued and pending patents assigned to Flow Pharma. Some of these patents pending are directly related to the study presented here.

References

1. Schwarz-Cruz, Y.C.A.; Ceballos-Cancino, G.; Vazquez-Santillan, K.; Espinosa, M.; Zampedri, C.; Bahena, I.; Ruiz, V.; Maldonado, V.; Melendez-Zajgla, J. Basal-Type Breast Cancer Stem Cells Over-Express Chromosomal Passenger Complex Proteins. *Cells* **2020**, *9*, doi:10.3390/cells9030709.

2. Andersen, M.H.; Svane, I.M.; Becker, J.C.; Straten, P.T. The universal character of the tumor-associated antigen survivin. *Clin Cancer Res* **2007**, *13*, 5991-5994, doi:10.1158/1078-0432.CCR-07-0686.

3. Wheatley, S.P.; Altieri, D.C. Survivin at a glance. *J Cell Sci* **2019**, *132*, doi:10.1242/jcs.223826.

4. Li, F.; Aljahdali, I.; Ling, X. Cancer therapeutics using survivin BIRC5 as a target: what can we do after over two decades of study? *J Exp Clin Cancer Res* **2019**, *38*, 368, doi:10.1186/s13046-019-1362-1.

5. Andersen, M.H.; Pedersen, L.O.; Capeller, B.; Brocker, E.B.; Becker, J.C.; thor Straten, P. Spontaneous cytotoxic T-cell responses against survivin-derived MHC class I-restricted T-cell epitopes in situ as well as ex vivo in cancer patients. *Cancer Res* **2001**, *61*, 5964-5968.

6. Bachinsky, M.M.; Guillen, D.E.; Patel, S.R.; Singleton, J.; Chen, C.; Soltis, D.A.; Tussey, L.G. Mapping and binding analysis of peptides derived from the tumor-associated antigen survivin for eight HLA alleles. *Cancer Immun* **2005**, *5*, 6.

7. Hirohashi, Y.; Torigoe, T.; Maeda, A.; Nabeta, Y.; Kamiguchi, K.; Sato, T.; Yoda, J.; Ikeda, H.; Hirata, K.; Yamanaka, N.; et al. An HLA-A24-restricted cytotoxic T lymphocyte epitope of a tumor-associated protein, survivin. *Clin Cancer Res* **2002**, *8*, 1731-1739.

8. Schmidt, S.M.; Schag, K.; Muller, M.R.; Weck, M.M.; Appel, S.; Kanz, L.; Grunebach, F.; Brossart, P. Survivin is a shared tumor-associated antigen expressed in a broad variety of malignancies and recognized by specific cytotoxic T cells. *Blood* **2003**, *102*, 571-576, doi:10.1182/blood-2002-08-2554.

9. Ahluwalia, M.S.; Reardon, D.A.; Abad, A.P.; Curry, W.T.; Wong, E.T.; Figel, S.A.; Mechtler, L.L.; Peereboom, D.M.; Hutson, A.D.; Withers, H.G.; et al. Phase IIa Study of SurVaxM Plus Adjuvant

Temozolomide for Newly Diagnosed Glioblastoma. *J Clin Oncol* **2022**, JCO2200996, doi:10.1200/JCO.22.00996.

10. Luo, C.; Wang, P.; He, S.; Zhu, J.; Shi, Y.; Wang, J. Progress and Prospect of Immunotherapy for Triple-Negative Breast Cancer. *Front Oncol* **2022**, *12*, 919072, doi:10.3389/fonc.2022.919072.

11. Mittendorf, E.A.; Philips, A.V.; Meric-Bernstam, F.; Qiao, N.; Wu, Y.; Harrington, S.; Su, X.; Wang, Y.; Gonzalez-Angulo, A.M.; Akcakanat, A.; et al. PD-L1 expression in triple-negative breast cancer. *Cancer Immunol Res* **2014**, *2*, 361-370, doi:10.1158/2326-6066.CIR-13-0127.

12. Schmid, P.; Rugo, H.S.; Adams, S.; Schneeweiss, A.; Barrios, C.H.; Iwata, H.; Dieras, V.; Henschel, V.; Molinero, L.; Chui, S.Y.; et al. Atezolizumab plus nab-paclitaxel as first-line treatment for unresectable, locally advanced or metastatic triple-negative breast cancer (IMpassion130): updated efficacy results from a randomised, double-blind, placebo-controlled, phase 3 trial. *Lancet Oncol* **2020**, *21*, 44-59, doi:10.1016/S1470-2045(19)30689-8.

13. Hosseini, M.; Seyedpour, S.; Khodaei, B.; Loghman, A.H.; Seyedpour, N.; Yazdi, M.H.; Rezaei, N. Cancer Vaccines for Triple-Negative Breast Cancer: A Systematic Review. *Vaccines (Basel)* **2023**, *11*, doi:10.3390/vaccines11010146.

14. Ohtake, J.; Ohkuri, T.; Togashi, Y.; Kitamura, H.; Okuno, K.; Nishimura, T. Identification of novel helper epitope peptides of Survivin cancer-associated antigen applicable to developing helper/killer-hybrid epitope long peptide cancer vaccine. *Immunol Lett* **2014**, *161*, 20-30, doi:10.1016/j.imlet.2014.04.010.

15. Pulaski, B.A.; Ostrand-Rosenberg, S. Mouse 4T1 breast tumor model. *Curr Protoc Immunol* **2001**, *Chapter 20*, Unit 20 22, doi:10.1002/0471142735.im2002s39.

16. Rubsamen, R.M.; Herst, C.V.; Lloyd, P.M.; Heckerman, D.E. Eliciting cytotoxic T-lymphocyte responses from synthetic vectors containing one or two epitopes in a C57BL/6 mouse model using

peptide-containing biodegradable microspheres and adjuvants. *Vaccine* **2014**, *32*, 4111-4116, doi:10.1016/j.vaccine.2014.05.071.

17. Harris, P.E.; Brasel, T.; Massey, C.; Herst, C.V.; Burkholz, S.; Lloyd, P.; Blankenberg, T.; Bey, T.M.; Carback, R.; Hodge, T.; et al. A Synthetic Peptide CTL Vaccine Targeting Nucleocapsid Confers Protection from SARS-CoV-2 Challenge in Rhesus Macaques. *Vaccines (Basel)* **2021**, *9*, doi:10.3390/vaccines9050520.

18. Herst, C.V.; Burkholz, S.; Sidney, J.; Sette, A.; Harris, P.E.; Massey, S.; Brasel, T.; Cunha-Neto, E.; Rosa, D.S.; Chao, W.C.H.; et al. An effective CTL peptide vaccine for Ebola Zaire Based on Survivors' CD8+ targeting of a particular nucleocapsid protein epitope with potential implications for COVID-19 vaccine design. *Vaccine* **2020**, *38*, 4464-4475, doi:10.1016/j.vaccine.2020.04.034.

19. Chen, S.; Zhou, Y.; Chen, Y.; Gu, J. fastp: an ultra-fast all-in-one FASTQ preprocessor. *Bioinformatics* **2018**, *34*, i884-i890, doi:10.1093/bioinformatics/bty560.

20. Pimentel, H.; Bray, N.L.; Puente, S.; Melsted, P.; Pachter, L. Differential analysis of RNA-seq incorporating quantification uncertainty. *Nat Methods* **2017**, *14*, 687-690, doi:10.1038/nmeth.4324.

21. Bray, N.L.; Pimentel, H.; Melsted, P.; Pachter, L. Erratum: Near-optimal probabilistic RNA-seq quantification. *Nat Biotechnol* **2016**, *34*, 888, doi:10.1038/nbt0816-888d.

22. Li, H.; Durbin, R. Fast and accurate short read alignment with Burrows-Wheeler transform. *Bioinformatics* **2009**, *25*, 1754-1760, doi:10.1093/bioinformatics/btp324.

23. Cunha-Neto, E.; Rosa, D.S.; Harris, P.E.; Olson, T.; Morrow, A.; Ciotlos, S.; Herst, C.V.; Rubsamen, R.M. An Approach for a Synthetic CTL Vaccine Design against Zika Flavivirus Using Class I and Class II Epitopes Identified by Computer Modeling. *Front Immunol* **2017**, *8*, 640, doi:10.3389/fimmu.2017.00640.

24. NoeDominguez-Romero, A.; Zamora-Alvarado, R.; Servin-Blanco, R.; Perez-Hernandez, E.G.; Castrillon-Rivera, L.E.; Munguia, M.E.; Acero, G.; Govezensky, T.; Gevorkian, G.; Manoutcharian, K.

Variable epitope library carrying heavily mutated survivin-derived CTL epitope variants as a new class of efficient vaccine immunogen tested in a mouse model of breast cancer. *Hum Vaccin Immunother* **2014**, *10*, 3201-3213, doi:10.4161/hv.29679.

25.   Siegel, S.; Wagner, A.; Schmitz, N.; Zeis, M. Induction of antitumour immunity using survivin peptide-pulsed dendritic cells in a murine lymphoma model. *Br J Haematol* **2003**, *122*, 911-914, doi:10.1046/j.1365-2141.2003.04535.x.

26.   Reynisson, B.; Alvarez, B.; Paul, S.; Peters, B.; Nielsen, M. NetMHCpan-4.1 and NetMHCIIpan-4.0: improved predictions of MHC antigen presentation by concurrent motif deconvolution and integration of MS MHC eluted ligand data. *Nucleic Acids Res* **2020**, *48*, W449-W454, doi:10.1093/nar/gkaa379.

27.   Andreatta, M.; Nielsen, M. Gapped sequence alignment using artificial neural networks: application to the MHC class I system. *Bioinformatics* **2016**, *32*, 511-517, doi:10.1093/bioinformatics/btv639.

28.   Jensen, K.K.; Andreatta, M.; Marcatili, P.; Buus, S.; Greenbaum, J.A.; Yan, Z.; Sette, A.; Peters, B.; Nielsen, M. Improved methods for predicting peptide binding affinity to MHC class II molecules. *Immunology* **2018**, *154*, 394-406, doi:10.1111/imm.12889.

29.   Ghaffari-Nazari, H.; Tavakkol-Afshari, J.; Jaafari, M.R.; Tahaghoghi-Hajghorbani, S.; Masoumi, E.; Jalali, S.A. Improving Multi-Epitope Long Peptide Vaccine Potency by Using a Strategy that Enhances CD4+ T Help in BALB/c Mice. *PLoS One* **2015**, *10*, e0142563, doi:10.1371/journal.pone.0142563.

30.   Zamani, P.; Teymouri, M.; Nikpoor, A.R.; Navashenaq, J.G.; Gholizadeh, Z.; Darban, S.A.; Jaafari, M.R. Nanoliposomal vaccine containing long multi-epitope peptide E75-AE36 pulsed PADRE-induced effective immune response in mice TUBO model of breast cancer. *Eur J Cancer* **2020**, *129*, 80-96, doi:10.1016/j.ejca.2020.01.010.

31.   Lim, J.F.; Berger, H.; Su, I.H. Isolation and Activation of Murine Lymphocytes. *J Vis Exp* **2016**, doi:10.3791/54596.

32.	Euhus, D.M.; Hudd, C.; LaRegina, M.C.; Johnson, F.E. Tumor measurement in the nude mouse. *J Surg Oncol* **1986**, *31*, 229-234, doi:10.1002/jso.2930310402.

33.	Tomayko, M.M.; Reynolds, C.P. Determination of subcutaneous tumor size in athymic (nude) mice. *Cancer Chemother Pharmacol* **1989**, *24*, 148-154, doi:10.1007/BF00300234.

34.	Cunningham, F.; Allen, J.E.; Allen, J.; Alvarez-Jarreta, J.; Amode, M.R.; Armean, I.M.; Austine-Orimoloye, O.; Azov, A.G.; Barnes, I.; Bennett, R.; et al. Ensembl 2022. *Nucleic Acids Res* **2022**, *50*, D988-D995, doi:10.1093/nar/gkab1049.

35.	Gregorio, A.C.; Fonseca, N.A.; Moura, V.; Lacerda, M.; Figueiredo, P.; Simoes, S.; Dias, S.; Moreira, J.N. Inoculated Cell Density as a Determinant Factor of the Growth Dynamics and Metastatic Efficiency of a Breast Cancer Murine Model. *PLoS One* **2016**, *11*, e0165817, doi:10.1371/journal.pone.0165817.

36.	Memarnejadian, A.; Meilleur, C.E.; Shaler, C.R.; Khazaie, K.; Bennink, J.R.; Schell, T.D.; Haeryfar, S.M.M. PD-1 Blockade Promotes Epitope Spreading in Anticancer CD8(+) T Cell Responses by Preventing Fratricidal Death of Subdominant Clones To Relieve Immunodomination. *J Immunol* **2017**, *199*, 3348-3359, doi:10.4049/jimmunol.1700643.

37.	Adams, S.; Gatti-Mays, M.E.; Kalinsky, K.; Korde, L.A.; Sharon, E.; Amiri-Kordestani, L.; Bear, H.; McArthur, H.L.; Frank, E.; Perlmutter, J.; et al. Current Landscape of Immunotherapy in Breast Cancer: A Review. *JAMA Oncol* **2019**, *5*, 1205-1214, doi:10.1001/jamaoncol.2018.7147.

38.	Dutoit, V.; Taub, R.N.; Papadopoulos, K.P.; Talbot, S.; Keohan, M.L.; Brehm, M.; Gnjatic, S.; Harris, P.E.; Bisikirska, B.; Guillaume, P.; et al. Multiepitope CD8(+) T cell response to a NY-ESO-1 peptide vaccine results in imprecise tumor targeting. *J Clin Invest* **2002**, *110*, 1813-1822, doi:10.1172/JCI16428.

39.	Fenstermaker, R.A.; Ciesielski, M.J.; Qiu, J.; Yang, N.; Frank, C.L.; Lee, K.P.; Mechtler, L.R.; Belal, A.; Ahluwalia, M.S.; Hutson, A.D. Clinical study of a survivin long peptide vaccine (SurVaxM) in patients with recurrent malignant glioma. *Cancer Immunol Immunother* **2016**, *65*, 1339-1352, doi:10.1007/s00262-016-1890-x.

40. Kameshima, H.; Tsuruma, T.; Kutomi, G.; Shima, H.; Iwayama, Y.; Kimura, Y.; Imamura, M.; Torigoe, T.; Takahashi, A.; Hirohashi, Y.; et al. Immunotherapeutic benefit of alpha-interferon (IFNalpha) in survivin2B-derived peptide vaccination for advanced pancreatic cancer patients. *Cancer Sci* **2013**, *104*, 124-129, doi:10.1111/cas.12046.

41. Kameshima, H.; Tsuruma, T.; Torigoe, T.; Takahashi, A.; Hirohashi, Y.; Tamura, Y.; Tsukahara, T.; Ichimiya, S.; Kanaseki, T.; Iwayama, Y.; et al. Immunogenic enhancement and clinical effect by type-I interferon of anti-apoptotic protein, survivin-derived peptide vaccine, in advanced colorectal cancer patients. *Cancer Sci* **2011**, *102*, 1181-1187, doi:10.1111/j.1349-7006.2011.01918.x.

42. Honma, I.; Kitamura, H.; Torigoe, T.; Takahashi, A.; Tanaka, T.; Sato, E.; Hirohashi, Y.; Masumori, N.; Tsukamoto, T.; Sato, N. Phase I clinical study of anti-apoptosis protein survivin-derived peptide vaccination for patients with advanced or recurrent urothelial cancer. *Cancer Immunol Immunother* **2009**, *58*, 1801-1807, doi:10.1007/s00262-009-0691-x.

43. Nitschke, N.J.; Bjoern, J.; Iversen, T.Z.; Andersen, M.H.; Svane, I.M. Indoleamine 2,3-dioxygenase and survivin peptide vaccine combined with temozolomide in metastatic melanoma. *Stem Cell Investig* **2017**, *4*, 77, doi:10.21037/sci.2017.08.06.

44. Lennerz, V.; Gross, S.; Gallerani, E.; Sessa, C.; Mach, N.; Boehm, S.; Hess, D.; von Boehmer, L.; Knuth, A.; Ochsenbein, A.F.; et al. Immunologic response to the survivin-derived multi-epitope vaccine EMD640744 in patients with advanced solid tumors. *Cancer Immunol Immunother* **2014**, *63*, 381-394, doi:10.1007/s00262-013-1516-5.

45. Becker, J.C.; Andersen, M.H.; Hofmeister-Muller, V.; Wobser, M.; Frey, L.; Sandig, C.; Walter, S.; Singh-Jasuja, H.; Kampgen, E.; Opitz, A.; et al. Survivin-specific T-cell reactivity correlates with tumor response and patient survival: a phase-II peptide vaccination trial in metastatic melanoma. *Cancer Immunol Immunother* **2012**, *61*, 2091-2103, doi:10.1007/s00262-012-1266-9.

46.   Widenmeyer, M.; Griesemann, H.; Stevanovic, S.; Feyerabend, S.; Klein, R.; Attig, S.; Hennenlotter, J.; Wernet, D.; Kuprash, D.V.; Sazykin, A.Y.; et al. Promiscuous survivin peptide induces robust CD4+ T-cell responses in the majority of vaccinated cancer patients. *Int J Cancer* **2012**, *131*, 140-149, doi:10.1002/ijc.26365.

47.   Feyerabend, S.; Stevanovic, S.; Gouttefangeas, C.; Wernet, D.; Hennenlotter, J.; Bedke, J.; Dietz, K.; Pascolo, S.; Kuczyk, M.; Rammensee, H.G.; et al. Novel multi-peptide vaccination in Hla-A2+ hormone sensitive patients with biochemical relapse of prostate cancer. *Prostate* **2009**, *69*, 917-927, doi:10.1002/pros.20941.