Chlamydomonas Mutants Null for Chloroplast Triose Phosphate Transporter3 are Metabolically Compromised and Light Sensitive

Weichao Huang (黄伟超)[1]†*, Anagha Krishnan[2]†, Anastasija Plett[3], Michelle Meagher[4], Nicole Linka[3], Yongsheng Wang[1,5], Bijie Ren[1], Justin Findinier[1], Petra Redekop[1], Neda Fakhimi[1], Rick G. Kim[1], Devin A. Karns[2], Nanette Boyle[4], Matthew C. Posewitz[2], Arthur R. Grossman[1]*

[1]Department of Plant Biology, Carnegie Institution for Science, Stanford, CA 94305, USA

[2]Department of Chemistry, Colorado School of Mines, Golden, CO 80401, USA

[3]Institute of Plant Biochemistry, Heinrich-Heine University Düsseldorf, 40225 Düsseldorf, Germany

[4]Department of Chemical and Biological Engineering, Colorado School of Mines, Golden, CO 80401, USA

[5]School of Life Sciences, Tsinghua University, 100084 Beijing, China

†Authors contributed equally.

*Corresponding Author: Weichao Huang; Address: The Carnegie Institution for Science,

260 Panama St. Bldg. 100; Stanford, CA 94305; email: whuang@carnegiescience.edu

*Corresponding Author: Arthur R. Grossman; Address: The Carnegie Institution for Science,

260 Panama St. Bldg. 100; Stanford, CA 94305; email: agrossman@carnegiescience.edu

Key words: TPT3, reactive oxygen species, CRISPR-Cas9, photosynthetic activity, starch accumulation, redox shuttles

Abstract

Modulation of export of photoassimilates from the chloroplast is essential for controlling the distribution of fixed carbon in the cell and maintaining optimum photosynthetic rates. In this study we identified chloroplast triose phosphate/phosphate translocators 2 and 3 (CreTPT2 and CreTPT3) in the green alga *Chlamydomonas reinhardtii* that exhibited similar substrate specificities but were differentially expressed over the diel cycle. We focused mostly on analyzing CreTPT3 because of its high level of expression and the severe phenotype exhibited by *tpt3* relative to the *tpt2* mutants. Null mutants for CreTPT3 had a pleiotropic phenotype that impacted growth, photosynthetic activities, metabolite profiles, carbon partitioning, and organelle-specific accumulation of H_2O_2. These analyses demonstrated that CreTPT3 is a dominant conduit on the chloroplast envelope for the transport of photoassimilate. In addition, CreTPT3 can serve as a safety valve that moves excess reductant out of the chloroplast and appears to be essential for preventing the cells from experiencing oxidative stress and accumulating of reactive oxygen species, even under low/moderate light intensities. Finally, our studies indicate subfunctionalization of the CreTPT transporters and suggest that there are differences in managing the export of photoassimilates from the chloroplasts of Chlamydomonas and vascular plants.

Introduction

Photosynthetic organisms can absorb excess, potentially damaging levels of light energy during mid-day (when photosynthetic electron transport becomes saturated) and can suffer from extreme damage, especially when experiencing rapid fluctuations in light intensities or when subjected to nutrient limiting conditions and other environmental stresses that impair the productive utilization of excitation energy (Chaux et al., 2017; Saroussi et al., 2017). To cope with excess excitation energy, plants and algae have evolved mechanisms to dissipate this energy through nonphotochemical quenching (NPQ) (Niyogi and Truong, 2013). The excess reductant and energy can also be eliminated by photochemical quenching. While the dominant form of photochemical quenching often involves the use of the electrons/reductant to fix inorganic carbon (Ci), which is directed toward growth or stored in the form of starch and lipids (Krishnan et al., 2015; Ge et al., 2014; Huang et al., 2018), redox equivalents can also be trafficked to other outlets where they are not used for anabolic processes. The main alternative photochemical electron outlets involve the reduction of O_2 to H_2O in H_2O-to-H_2O cycles that include a (*i*) Mehler-type reaction that is noncatalytic and functions on the acceptor side of PSI (Badger et al., 2000), (*ii*) flavodiiron protein reactions (FLVs, NADPH:flavin oxidoreductase) that catalytically reduce O_2 on the acceptor side of PSI without generating reactive oxygen species (ROS) (Helman et al., 2003; Chaux et al., 2017), (*iii*) the plastid terminal oxidase (PTOX, plastoquinol: oxygen oxidoreductase) reaction that can use electrons from the PQ pool to reduce O_2 (Houille-Vernes et al., 2011), and (*iv*) the movement of redox equivalents from the chloroplast to the mitochondrion where they can be used to reduce O_2 through various electron transport activities.

Mechanisms also exist in which fixed carbon and reducing equivalents can be shuttled between the chloroplast and cytoplasm. Studies with *Arabidopsis thaliana* revealed that the malate shuttle, which involves multiple malate dehydrogenases (MDH) and malate/OAA translocators (OMT), functions in the export of reductant from the chloroplast and the management of redox conditions in the chloroplast (Zhao et al., 2020, 2018). In addition to the malate shuttle, the triose phosphate (triose-P)/phosphate (Pi) translocator (TPT) has been proposed to be involved in moving fixed carbon out of the chloroplast, but can also act as a safety valve for eliminating excess reducing power from the chloroplast (Fliege et al., 1978; Flügge et al., 1989; Lee et al., 2017; Stocking and Larson, 1969; Raghavendra and Padmasree, 2003; Johnson and Alric, 2013). The synthesis of triose-Ps during photosynthetic CO_2 fixation by the Calvin-Benson-Bassham Cycle (CBBC) is

supported by the reducing power/energy (NADPH, ATP) derived from photosynthetic electron
transport.

The TPTs reside on the inner chloroplast envelope membrane and can transport triose-Ps
(glyceraldehyde 3-phosphate (GAP), dihydroxyacetone phosphate (DHAP)) and the three carbon
acid 3-phosphoglycerate (3-PGA) in a counter exchange for cytosolic Pi (Fliege et al., 1978; Lee
et al., 2017; Flügge et al., 1989). These transporters belong to a family of plastidic phosphate
translocators (pPTs) that function as antiport systems involved in exchanging Pi with
phosphorylated C3, C5 or C6 compounds (Flügge et al., 2003). Most angiosperms have two *TPT*
genes in their genome, except for various monocots and two dicot families, the Amaranthaceae
and the Brassicaceae, which have a single *TPT* gene (Bockwoldt et al., 2019). In plants, cytosolic
trioses exported from the chloroplast by the TPTs are used for the biosynthesis of sucrose and
other metabolites (Riesmeier et al., 1993) and to drive respiratory activity. In addition to TPTs,
plants harbor the three other pPT subfamilies (Fischer et al., 1997; Kammerer et al., 1998; Eicks
et al., 2002; Lee et al., 2017; Flügge et al., 1989). These include the glucose 6-phosphate (Glc6P)
translocator (GPT), which imports Glc6P into plastids in heterotrophic tissue (Kammerer et al.,
1998), the xylulose phosphate translocator (XPT), which plays a key role in coordinating the
cytosolic and plastidic pentose phosphate pathways (Eicks et al., 2002), and the
phosphoenolpyruvate (PEP) translocator (PPT), which imports PEP into C3 plastids; the PEP can
be used for the synthesis of fatty acids, as substrate for the shikimate pathway (Streatfield et al.,
1999; Prabhakar et al., 2010), and for the export of PEP from plastids in C4 plants (Häusler et al.,
2000).

Over the last few decades, the physiological functions of the TPTs have been examined in some
detail. Various plants do not exhibit a strong phenotype if the chloroplast TPT is either eliminated
or its level is reduced (Häusler et al., 1998; Walters et al., 2004; Schneider et al., 2002; Riesmeier
et al., 1993). A reduction in TPT activity can be compensated for by diverting assimilated carbon
into a transitory starch pool that is subjected to accelerated turnover in the light and/or dark
(Häusler et al., 1998; Walters et al., 2004; Riesmeier et al., 1993), leading to accumulation of starch
degradation products that can be exported from the chloroplast and used in other cellular
compartments. Interestingly, a deficiency of the TPT in rice, a plant that uses sucrose stored in the
leaves as its major transitory form of fixed carbon, led to severe phenotypic consequences; the
plants exhibited reduced photosynthetic rates and decreased levels of both starch and soluble

sugars relative to wild type (WT) plants (Lee et al., 2014).

Microalgae have high photosynthetic conversion efficiencies, can thrive in fresh to hypersaline
waters and can be metabolically versatile. They have attracted considerable interest worldwide
because of their ability to synthesize large quantities of lipids (e.g. for biofuels and food products),
starch, pigments and other bioproducts, and can serve in the remediation of wastewater (Khan et
al., 2018; Bhatt et al., 2022). The transporters used for moving photoassimilate between the
chloroplast and other cellular compartments and the mechanisms and regulation of these
transporters in microalgae have not been extensively explored. Developing a more informed
understanding of central metabolism in microalgae and the movement of metabolites among
compartments can enable additional work on the establishment, regulation, and evolution of
metabolic networks in algae and the ways in which algae can be tailored for production purposes
and for sustained growth under specific environmental conditions.

In this study, we used Chlamydomonas, a well-established model green algal system that has been
extensively used to analyze various physiological processes, to dissect the function of chloroplast
TPTs. Of the four putative Chlamydomonas pPTs, we discovered that at least two are 'genuine'
TPTs (CreTPT2 and CreTPT3) based on yeast liposome transport assays. CreTPT2 was previously
reported to be a PPT based on phylogenetic analysis (Bockwoldt et al., 2019). These two TPTs
exhibited sub-functionalization that is reflected by their expression levels and temporally and
environmentally distinct regulatory patterns. Due to the more severe phenotypes exhibited by *tpt3*
relative to *tpt2* mutants, we focused our analyses on Chlamydomonas TPT3 (CreTPT3) which,
among the four predicted pPTs, is highly expressed in the light and strongly induced by various
environmental stresses. Through a series of physiological analysis, we demonstrated that CreTPT3
is a major conduit on the chloroplast envelope for the trafficking of fixed carbon, sustaining central
carbon metabolism, dissipating excess energy, enabling high rates of photosynthetic electron
transport, preventing intracellular hydrogen peroxide (H_2O_2) accumulation, and balancing redox
conditions at the subcellular level.

Results

The triose-P/Pi translocator (DMT/TPT) family

Candidate genes encoding triose-P/Pi transporters (TPT) of Chlamydomonas were identified by

blasting the Arabidopsis TPT protein (AT5G46110.1) against proteins encoded on the Chlamydomonas genome. The TPT family is the largest within the drug/metabolite transporter (DMT) superfamily in eukaryotes and includes triose-P and sugar-phosphate transporters associated with chloroplasts; many of the family members are still not functionally characterized (Jack et al., 2001; Knappe et al., 2003; Weber et al., 2006). We identified 32 genes encoding potential TPTs in the version v6.1 genome of Chlamydomonas, with four members, CreTPT10 (CreTPT1 in v5.6 genome), CreTPT2 (CreTPT2 in v5.6 genome), CreTPT3 and CGL51 (CreTPT25 in v5.6 genome), predicted to have a transit peptide that would localize the protein to the chloroplast (**Supplementary Table 1**). These four candidates were included in a phylogenetic analysis of plant and algal pPTs that showed that CreTPT3 is a putative TPT, CreTPT2 and CreTPT10 are putative PPTs, and CGL51 is a putative GPT or XPT (Bockwoldt et al., 2019).

To examine the potential substrate specificities of the four putative pPTs, their amino acid sequences were aligned with pPTs from Arabidopsis (AtPTs) (**Supplementary Fig. 1A**). The ability of these transporters to use triose-P/DHAP or 3-PGA as substrate is dependent on five highly conserved amino acid residues (H184, K203, Y338, K359, and R360 in AtTPT1) (Lee et al., 2017; Moog et al., 2020). Of the putative pPTs in Chlamydomonas, only CreTPT3 and CGL51 contain all five of these residues (H170, K189, Y322, K345, and R346 in CreTPT3) (**Supplementary Fig. 1A**). In AtTPT1, residue F262 (F248 in Chlamydomonas in the analogous protein, CreTPT3) is thought to inhibit PEP access to the binding site; this residue is replaced by N in AtPPT1 (PPT) (Lee et al., 2017; Moog et al., 2020). CGL51 has an M at position F248, indicating that unlike CreTPT3, CGL51 might have preference for other substrates. Protein sequence similarity and identity analysis show CreTPT3 and CreTPT2 share the highest and the second highest similarity and identity with AtTPT1, (**Supplementary Fig. 1B**) and are 57% and 49% similar to AtTPT1, respectively. Furthermore, *CreTPT2* and *CreTPT3* have significantly higher transcript accumulation in light relative to Cre*TPT10* and *CGL51* (**Supplementary Fig. 1C, D**). Given the high similarity and high expression levels, CreTPT2 and CreTPT3 were chosen for further study.

CreTPT2 and CreTPT3 transport properties

We defined the subcellular localization and substrate preferences of both CreTPT2 and CreTPT3. As shown in **Fig. 1A**, both CreTPT2 and CreTPT3 fused to VENUS localized to the chloroplast

envelope. To evaluate the substrate specificity of these transporters, each of them was expressed

in *Saccharomyces cerevisiae* (yeast) and the resulting recombinant protein was biochemically

analyzed using a liposome uptake assay. The Cre*TPT2* and Cre*TPT3* genes were fused at their C

termini to a sequence encoding a His tag, codon-optimized, and expressed in yeast (**Fig. 1B**), and

total cell membranes were isolated and reconstituted into liposomes (Loddenkötter et al., 1993;

Linka et al., 2008). Chloroplast phosphate transporters in vascular plants can catalyze a Pi/Pi

homo-exchange in vitro. Both CreTPT2 and CreTPT3 reconstituted in liposomes were able to

catalyze the signature Pi homo-exchange, whereas in the absence of a counter-exchange substrate,

little Pi uptake was detected (**Fig. 1C** and **D**, left). In contrast, very low Pi uptake rates were

observed for liposomes reconstituted with membranes from yeast cells lacking CreTPT2 or

CreTPT3 (**Supplementary Fig. 2**), indicating that the introduced transporters were responsible for

the detected Pi import activity in the yeast liposomes.

To assess the substrate specificity of the CreTPT transporters, the initial rates of Pi uptake into

liposomes preloaded with saturating concentrations (30 mM) of various potential counter-

exchange substrates were determined. As the right panels of **Fig. 1C** and **D** show, both CreTPT2

and CreTPT3 exhibited the highest activity when DHAP was used as the substrate for the yeast

liposome assay. For both transporters, the relative initial velocity for DHAP/Pi exchange was

slightly higher than that of 3-PGA/Pi (the relative 3-PGA/Pi exchange was 75% of DHAP/Pi

exchange) while Pi uptake into liposomes preloaded with PEP was much lower (**Fig. 1C** and **D**,

right panels). Pi import was negligible when CreTPT2 or CreTPT3 liposomes were preloaded with

Glc-6-P, Glc-1-P, Fru-6-P, and Gal-1-P (**Fig. 1C** and **D**, right panels). These results show that

CreTPT2 and CreTPT3 have almost the same substrate preferences, with both specifically

catalyzing the transport of triose-P and 3-PGA across the membrane in exchange for Pi.

Characterizations of the K_M and K_i for CreTPT2 and CreTPT3 were performed to determine the

affinity of these transporters for the various substrates. CreTPT3 has an apparent Michaelis-

Menten constant (K_M) of 1.1 +/- 0.2 mM for Pi (**Fig. 1E**), which is comparable to the value

obtained for the vascular plant TPT ortholog (Fliege et al., 1978). CreTPT2 has a slightly lower

K_M for Pi than CreTPT3 (0.77 +/- 0.05 mM) (**Fig. 1E**). However, while the 3-PGA K_i values were

comparable for the two transporters, DHAP was more effective in inhibiting the CreTPT3-

dependent Pi exchange than CreTPT2-dependent Pi exchange (**Fig. 1E**). These results suggest that

CreTPT3 may have a greater specificity for the transport of DHAP than CreTPT2 and that it may

be more effective in transporting C3 phosphorylated compounds than CreTPT2. In the case of

PEP, no inhibition of the Pi/Pi homo-exchange was observed, even at the non-physiologically high

concentration of 5 mM. Overall, the results of these in vitro assays indicate that both CreTPT2 and

CreTPT3 have a typical plant-TPT substrate spectrum (Fliege et al., 1978) and can transport Pi,

triose-P (DHAP) and 3-PGA in a counter-exchange mode. However PEP might not be a

physiologically relevant substrate for either CreTPT2 or CreTPT3, as was shown for apoplast TPT

homologues in other organisms (Lim et al., 2010; Moog et al., 2020). Furthermore, the values

generated in these analyses may not precisely reflect transport kinetics in vivo since the two

transporters used for these in vitro assays were fused to a His-tag at their N terminus, and the yeast

liposomes would have a different lipid composition than the chloroplast inner envelop membrane,

where these transporters are normally localized.

Isolation of *tpt2* and *tpt3* null mutants of Chlamydomonas and their impacts on cell growth

To explore the role of CreTPT2 and CreTPT3 in trafficking carbon and potentially reductant across

the chloroplast envelope in vivo, CRISPR knockouts of Cre*TPT2* or Cre*TPT3* were generated. We

used the CRISPR-Cas9 editing system to disrupt the Cre*TPT2* or Cre*TPT3* genes while at the same

time integrating the hygromycin resistance marker gene (*AphVII*) into the edited site (**Fig. 2A,**

Supplementary Fig. 3 and **4**). Four independent knockouts of Cre*TPT2* (*t2ko1, t2ko2, t2ko3 and*

4) were obtained, with the marker gene inserted into exon 8 (**Supplementary Fig. 3)**. Three

independent knockouts of Cre*TPT3* were obtained, including *t3ko1*, with the marker gene inserted

into exon 1, and *t3ko2* and *t3ko3*, with the marker gene inserted into exon 7; no CreTPT3 protein

was detected in any of these edited strains (**Fig. 2B**).

To elucidate the physiological roles of CreTPT2 and CreTPT3, we first examined growth of the

parental WT strain and the knockout mutants under either low light (LL, 30 μmol photons m^{-2} s^{-}

^{1}), moderate light (ML, 250-300 μmol photons m^{-2} s^{-1}) or high light (HL, 450 μmol photons m^{-2} s^{-}

^{1}). The design of the experiments in which the cells were transferred from one light condition to

another is shown in **Supplementary Fig. 5**. The four *tpt2* mutants (*t2ko1, t2ko2, t2ko3 and 4*) were

not impacted by growth in LL, but their growth was significantly impaired relative to WT cells

under HL on either photoautotrophic (TP) or heterotrophic (TAP) solid agar medium

(**Supplementary Fig. 3D, Fig. 2C**). In contrast, the Cre*TPT3* mutants (*t3ko1, t3ko2* and *t3ko3*)

exhibited severe growth impairment even in LL, and completely stopped growing under ML and

HL, on either TP or TAP agar medium (**Fig. 2C, D**). Due to the striking growth phenotypes caused

by the loss of CreTPT3, we focused on investigating the physiological functions of TPT3 by

analyzing *tpt3* mutants in more detail in this study. The differences in transcript changes between

Cre*TPT2* and Cre*TPT3* under various environmental conditions were examined and are described

in the 'Discussion' section.

Growth curves were also determined for WT and *tpt3* mutant cells in liquid medium (TP) in LL,

ML and HL (**Fig. 2E** and **2F**); the results were in accord with those observed for the solid medium

growth assays. Additionally, *tpt3* mutant cells in both LL and ML exhibited an increased cell

diameter and formed clusters of cells that appear to be less able to separate following cell division

(**Supplementary Fig. 6A**, **B**). To further confirm that the growth phenotypes are a consequence

of the *tpt3* knockout, we introduced a wild-type copy of Cre*TPT3* fused to VENUS into the mutant

strains (**Fig. 2B**, right). As shown in **Fig. 2D-G**, ectopic expression of the wild-type Cre*TPT3* in

the *t3ko2* mutant (*C_T3KO2*) rescued the reduced growth phenotype of the mutant under all light

conditions tested in this study (LL/ML/HL).

We also analyzed the chlorophyll content of photoautotrophically grown cells after transferring

them from LL to ML and found that *t3ko2* had reduced chlorophyll content relative to WT cells

after 24 h in ML (**Fig. 2F-G** and **Supplementary Fig. 7**). Cell numbers and total chlorophyll were

quantified following the LL to ML transition. The chlorophyll levels per cell declined in all strains

at 24 and 48 h following transfer to ML; however, it was lower by approximately half in *t3ko2*

(0.34 µg/10^6 cells) relative to either WT cells or the *C_T3KO2* rescued strain (both ~0.72 µg/10^6

cells) after 24 h of ML (**Supplementary Fig. 7**), with some additional increase in cell density for

WT and *C_T3KO2* after 48 h of ML (which might result in some shading). Finally, when mutant

cells were transferred to HL for 24 h they became strongly bleached (**Fig. 2F**).

Taken together, these results indicate that the activity of the CreTPT3 transporter is essential for

optimal growth over a range of light intensities (LL/ML/HL).

The *tpt3* mutant exhibits hyper-accumulation of 'storage' carbon

To explore the impact of the loss of CreTPT3 activity on carbon partitioning, we quantified carbon

storage (starch and lipids) following a transition of WT and the *t3ko2* mutant from LL to ML.

Lugol staining showed extensive starch accumulation in the *t3ko2* cells after a 48-h exposure to ML, whereas WT cells were barely stained (**Fig. 3B**). Furthermore, as shown in **Fig. 3A**, there is a ~55-fold difference in the level of starch that accumulated in *t3ko2* relative to WT cells (13.29 compared to 0.24 µg starch/µg chlorophyll, respectively) after 24 h of illumination in ML. The mutant also accumulated ~24-fold more lipid than WT and *C_T3KO2* cells (on a chlorophyll basis), as monitored by Nile Red fluorescence, over the same time period (WT and *t3ko2*: 1038 and 41 Nile Red fluorescence/chlorophyll, respectively) (**Fig. 3C, D**). Since the chlorophyll in the mutant on a per cell basis was approximately 50% relative to that of WT cells after 24 h in ML, the accumulated starch and lipid on a per cell basis would be ~25-fold more starch and ~12-fold more lipid in the mutant relative to the WT strain. Additionally, mutant cells are much larger and tended to exhibit more aggregation than WT cells (**Supplementary Fig. 6A, B**). These results suggest that the inability to transport triose-P between the chloroplast and the cytosol through CreTPT3 resulted in a repartitioning of photosynthetic assimilates (carbon, reductant, and ATP) toward the synthesis of both starch and neutral lipid.

Cre*TPT3* deletion leads to accumulation of CBBC/glycolytic/gluconeogenic intermediates

To understand the metabolic consequences of the loss of CreTPT3 on growth under LL and ML, comparative metabolite analyses of WT and the *t3ko2* mutant were performed on cells grown in LL and after shifting them to ML for both 45 min and 6 h. As mentioned above, the level of starch increased dramatically in the mutant relative to WT cells (**Fig. 3A, B** and **Supplementary Fig. 8A**). Pool sizes of various central carbon metabolites, particularly those of the CBBC/glycolysis/gluconeogenesis pathways and intermediates of the TCA/glyoxylate cycle (**Fig. 3F**), were quantified. Data was normalized to both cell number (**Fig. 3E**) and chlorophyll content (**Supplementary Fig. 8B**). The fold-change in the quantity of each metabolite in the mutant relative to WT under LL or at 45 min and 6 h after the switch to ML is given in **Fig. 3G** and **Supplementary Fig. 8C**.

At the time of shifting cells from LL (**Fig. 3E**, time 0 h) to ML, the *t3ko2* mutant had already accumulated a significantly larger pool (2- to 6-fold) of glycolytic/gluconeogenic intermediates [DHAP, 3-PGA, FBP (all three shared with CBBC), G6P, and PEP] compared to WT cells, suggesting that the loss of CreTPT3 resulted in a back-up of these metabolites within the cell; our hypothesis is that these metabolites are accumulating in the chloroplast stroma due to the reduced

ability of the mutant chloroplast to export fixed carbon, which is supported by the observed

accumulation of starch and TAG in the mutant strain (**Fig. 3A-D** and **Supplementary Fig. 8A**).

While the pool sizes of fumarate and malate (metabolites of the TCA/glyoxylate cycle) were

similar in both strains under LL, following the transition of the cells to ML, WT cells exhibited a

significant increase in the pool sizes of those metabolites while the mutant maintained a lower

level, indicating that the loss of CreTPT3 either directly (by supplying precursors) or indirectly

(metabolic rewiring) impacts their levels.

Thus, a primary function of CreTPT3 appears to be the export of photosynthetically synthesized

reduced carbon, which would drive metabolic processes in the cytoplasm and other cellular

compartments, such as the mitochondrion, while enabling the import of Pi into the chloroplast,

which sustains ATP synthesis.

The *tpt3* mutant has reduced photosynthetic activity when grown in ML

To understand the impact of the loss of CreTPT3 activity on PET, we quantified photosynthetic

activities following a transition from LL to ML (**Supplementary Fig. 5**). Photosynthetic O_2

evolution rates (OERs) were measured for WT, *t3ko2* and *C_T3KO2* over a range of light

intensities. LL acclimated WT, *t3ko2*, and *C_T3KO2* showed comparable OERs at intensities

below 200 μmol photons m^{-2} s^{-1}, while at saturating light intensities of $\geq$600 μmol photons m^{-2} s^{-1},

the OER of *t3ko2* was diminished slightly relative to the WT and the complemented strain (**Fig.**

4A). Upon acclimation of the cells to ML for 24 h, the *t3ko2* mutant consistently displayed lower

OERs than WT cells or *C_T3KO2* under all actinic light intensities used (**Fig. 4A**). Additionally,

while the Fv/Fm in the mutant grown in LL was comparable to that of WT and *C_T3KO2* (**Fig.**

4B), after exposure of the mutant cells to ML for 6 h, the Fv/Fm declined to about 50% of the WT

and *C_T3KO2* levels; the decline in the mutant continued over a period of 24 h in ML, with a 3-

fold decrease in Fv/Fm for *t3ko2* compared to that of WT cells (**Fig. 4B**). These results indicate

that damage to PSII reaction centers occurs following exposure of the mutant to ML.

To examine the redox state of the photosynthetic apparatus, the pool of electron acceptors

downstream of PSII were evaluated. We quantified the photochemical efficiency (qL) of all strains

at various light intensities after growth in LL or shifting them to ML for 6 and 24 h (**Fig. 4C**). The

photosynthetic parameter qL indicates the redox state of Q_A, the primary electron acceptor of the

PSII reaction center. Assuming that Q_A and Q_B are in equilibrium, qL would reflect the redox

status of the PQ pool; a lower qL value indicates a more reduced electron transport chain (Kramer

et al., 2004). 1-qL positively correlates with the PQ pool redox state. In the LL-acclimated cells,

1-qL was significantly higher in *t3ko2* even at low levels of actinic illumination compared to that

of WT cells and the rescued strain. For cells that had been acclimated to ML (6 h and 24 h), 1-qL

was 70% of the near maximum value for *t3ko2*, even under relatively low actinic light conditions

(e.g. 100 µmol photons $m^{-2} s^{-1}$), indicating a highly reduced PQ pool. In contrast, for WT and the

complemented strain at the same light intensity, the 1-qL value attained only ~20% of the

maximum value (**Fig. 4C**). These results suggest that the PQ pool is much more reduced in LL-

acclimated *t3ko2* relative to WT and *C_T3KO2* (**Fig. 4C**), and that there is a much more

pronounced reduction of this pool at all actinic light intensities in ML grown *t3ko2* mutants. These

results suggest a limitation in electron flow downstream of the PQ pool in both LL and ML grown

t3ko2.

Moreover, we analyzed how fast PET becomes restricted following the transfer of LL-grown *t3ko2*,

C_T3KO2 and WT cells to HL (400-450 µmol photons $m^{-2} s^{-1}$) by monitoring changes in the redox

state of the PQ pool. We observed that the PQ pool in *t3ko2* began to be more reduced than that of

WT cells and *C_T3KO2* after 105 s of elevated illumination (**Fig. 4D**), and that this was even more

pronounced at longer times in HL. These results suggest that PET can be rapidly limited by

diminished triose-P export from the chloroplast; elimination of CreTPT3 has a strong impact on

PET activity.

To determine if PSI was also impacted in the mutant cells, we analyzed PSI/P700

oxidation/reduction kinetics following exposure of LL-grown cells to ML in the presence of

DCMU (20 µM) and hydroxylamine (1 mM) to block a contribution of electrons from PSII. Levels

of photo-oxidizable P700 following exposure to actinic light for 5 sec were similar in all LL-

acclimated strains (**Fig. 4E**). However, the level of photo-oxidized P700 in ML-acclimated cells

declined in the mutant to ~40-50% of that in WT and *C_T3KO2* cells (**Fig. 4E**), indicating that

PSI is more reduced in the *t3ko2* mutant after ML exposure. As shown in **Fig. 4F**, the oxidation

rate of P700 following a dark to light transition was much slower in the ML-acclimated *t3ko2*

compared to that of WT. These results suggest that the mutant has a diminished level of available

electron acceptors on the acceptor side of PSI (relative to WT and *C_T3KO2*) after growth in ML.

CreTPT3 inactivation dramatically affected accumulation and distribution of compartmentalized H_2O_2

To investigate the relationship of CreTPT3 activity to oxidative stress, we assayed ROS production in the mutant using the fluorescent probe CM-H2DCFDA, which upon exposure to increasing ROS levels is converted to the green-fluorescent molecule dichlorofluorescein (DCF). DCF fluorescence was visualized by confocal microscopy. As shown in **Fig. 5A** and **B**, ROS levels in the *t3ko2* mutant were markedly increased (~3-fold) after 48 h in ML, while little difference in ROS levels was detected in WT cells.

For an alternative, dynamic method for evaluating redox changes in the chloroplast, we used the redox sensitive green fluorescent protein roGFP2, which was targeted to the chloroplast stroma (**Fig. 5C**, upper panel) and mitochondrion (**Fig. 5C**, lower panel). roGFP2 monitors ratiometric redox changes of glutathione, which reflects cellular ROS levels (Dorion et al., 2021; Vevea et al., 2013). LL-acclimated WT cells and the *t3ko2* mutant exhibited similar levels of chloroplast roGFP2 oxidation (**Fig. 5D**). Upon transfer of these cells to ML, the mutant showed an increase in chloroplast oxidative conditions, with a 5.0-fold increase after 1.5 h, and a 2.0-fold increase after 6 h, which is 1.8- and 1.6-fold higher than the values measured in WT cells (**Fig. 5D**). Additionally, as triose-P is exported to the cytosol by CreTPT3, it could potentially be further metabolized and donate redox equivalents to the mitochondrial electron transport chain and alter mitochondrial ROS production. Therefore, we also measured mitochondrial redox levels at different light intensities in both WT and the *t3ko2* mutant using the roGFP2 sensor targeted to the mitochondrion. Upon a LL to ML shift for 1.5 and 6.0 h, neither WT nor *t3ko2* displayed a significant change in fluorescence from the mitochondrial targeted roGFP (**Fig. 5E**). These results suggest that the mitochondrial redox level is maintained after shifting either LL acclimated WT or *t3ko2* cells to ML. Overall, the inability to export triose-P through CreTPT3 markedly increased the level of oxidative stress in the chloroplast but not in the mitochondrion.

We also determined if the ROS accumulated in *t3ko2* is H_2O_2 and whether this molecule shows differential accumulation in the different subcellular compartments; the analysis was based on the use of a hypersensitive sensor of H_2O_2, roGFP2-Tsa2ΔCR, which was previously used for studies with Chlamydomonas (Niemeyer et al., 2021). In this analysis, we monitored real-time accumulation of H_2O_2 in the stroma, cytosol, mitochondrion, and nucleus (**Fig. 6A**) following a 20

min exposure of the cells to either HL or very low light (**Fig. 6B-E**). The *t3ko2* stromal H_2O_2 level

increased within 2.5 min of the light exposure and attained a 1.4-fold increase after 20 min of

illumination with HL compared to the initial level in LL (**Fig. 6B**). The stromal H_2O_2 levels in WT

cells showed little change after being shifted to HL, however, the level declined when the cells

were shifted to 10 μmol photons m^{-2} s^{-1}, or very low light. Notably, after 20 min in HL, the stroma

of the *t3ko2* mutant accumulated ~1.6-fold more H_2O_2 than that of the WT cells; *t3ko2* mutant cells

in very low light accumulated a similar amount of stromal H_2O_2 as WT cells after exposure to HL

(**Fig. 6B**). The cytosolic probe also responded rapidly, with a 1.2-fold increase for WT and a 1.4-

fold increase for the *t3ko2* mutant (**Fig. 6C**); note that the initial levels of H_2O_2 prior to the transfer

of the cells to HL or very low light were much lower in the WT cells than in the mutant. The *t3ko2*

cytosolic H_2O_2 level was elevated relative to the level in WT cells by 2.1-fold after 20 min of

illumination in HL (**Fig. 6C**). We did not observe a significant change in H_2O_2 levels in the

mitochondrion for either the WT or the *t3ko2* mutant after the cells were shifted to the higher light

intensity (**Fig. 6D**). Finally, the *t3ko2* mutant already accumulated much higher levels of H_2O_2 in

the nucleus in LL compared to that in the WT cells, although both mutant and WT cells did not

show significant changes in nuclear H_2O_2 levels after HL exposure (**Fig. 6E**).

Discussion

Cre*TPT2* and Cre*TPT3* genes exhibit different expression patterns

In this study, we discovered that Chlamydomonas contains at least two TPTs that are located on the chloroplast envelope. An earlier report suggested that CreTPT2 was a plastidic PPT (for the transport of PEP) (Bockwoldt et al., 2019), but based on our results, it appears to be functionally more similar to a triose-P transporter. In vitro assays show that CreTPT2 has almost the same substrate specificity as CreTPT3, although it may be less effective in DHAP/Pi exchange (**Fig. 1C-E**). To determine expression patterns of Cre*TPT2* and Cre*TPT3*, the abundance of the Cre*TPT2* and Cre*TPT3* transcripts were analyzed using RT-qPCR and by mining published RNA-seq data over the diurnal cycle, and during nitrogen/sulfur/iron starvation (Zones et al., 2015; Ngan et al., 2015; González-Ballester et al., 2010; Urzica et al., 2013) (**Supplementary Fig. 1C-D,** and **9**). Cre*TPT3* was highly expressed in the light and dark, with significantly higher transcript accumulation than that of Cre*TPT2* and the other genes (Cre*TPT10* and *CGL51*) potentially encoding chloroplast localized pPTs (**Supplementary Fig. 1C-D**). Cre*TPT2* and Cre*TPT3* also responded differentially to abiotic stresses. Cre*TPT3* was strongly induced by nitrogen, sulfur, and iron starvation and upon exposure to HL, whereas the level of the Cre*TPT2* transcript remained almost unchanged under the same conditions (**Supplementary Fig. 9A-D**), suggesting that CreTPT3 plays a more prominent role in exporting triose-P from the chloroplast than CreTPT2, with potentially increasing export from the plastid under HL and nutrient limitation conditions. This hypothesis is supported by the observation that *tpt3* mutants displayed much more severe growth retardation relative to *tpt2* mutants upon exposure to ML or HL (**Fig. 2C**). There is no evidence showing that the *TPTs* from plants are induced by stress/excess absorbed excitation, and expression of the Arabidopsis *TPT* gene appears to even decrease following HL exposure (Weise et al., 2019).

Cre*TPT2* and Cre*TPT3* exhibited distinct expression patterns over the diurnal cycle; the expression of Cre*TPT2* increased rapidly after transitioning from the dark to the light, with peak accumulation after one hour in the light, when the transcript level of Cre*TPT3* was at its lowest (**Supplementary Fig. 1D**). Continued exposure to light led to a decrease in the level of the Cre*TPT2* transcript to near zero while the transcript from Cre*TPT3* steadily increased in the light, reaching a peak in mid-

day (**Supplementary Fig. 1D**). The Cre*TPT2* expression pattern suggests that it might play a role

in exporting triose-P at the beginning of light period, when the light intensity and photosynthesis

are low and low levels of triose-P would be synthesized in the stroma. As the light intensity

increases over the course of the day, higher levels of triose-P are synthesized and its trafficking

out of the chloroplast for use in other subcellular compartments would likely predominantly

involve the activity of CreTPT3. The greater specificity of CreTPT3 than CreTPT2 for transporting

DHAP (**Fig. 1E**) may make it more effective than CreTPT2 in transporting C3 phosphorylated

compounds. This possibility is congruent with the finding that there is elevated

synthesis/accumulation of CreTPT3 mRNA during the day when the light intensity reaches its

peak (Zones et al. 2015) and there would be rapid synthesis of the C3 phosphorylated compounds.

Overall, the subfunctionalization of the two Chlamydomonas triose-P transporters based on their

expression levels, patterns of RNA accumulation over the course of the day and upon nutrient

deprivation, and their substrate specificities, may help tune the export of triose-P from the

chloroplast with respect to the diurnal cycle and dynamic environmental cues.

CreTPT3 also potentially serves as a redox valve, transferring reductant to the cytoplasm

It was previously proposed that chloroplast TPTs could catalyze two potential reactions in the light

(**Fig. 7**) based on the crystal structure of the red algal TPT (Lee et al., 2017) and in vitro assays

using isolated spinach chloroplasts (Stocking and Larson, 1969); both triose-P/Pi, and triose-P/3-

PGA exchange across the chloroplast inner envelope membrane. The former reaction can route

both carbon skeletons and reductants into the cytoplasm while importing Pi back into the

chloroplast for ATP regeneration. The latter reaction would import 3-PGA into the chloroplast in

exchange for triose-P (DHAP, GAP), which would serve to transfer reductant from the chloroplast

to the cytosol while transferring 3-PGA back into the chloroplast where it can be reduced by the

CBBC and stimulate the regeneration of ribulose 1,5-bisphosphate. Indeed, in vitro, CreTPT3 can

actively transport both triose-P and 3-PGA in exchange for Pi (**Fig. 1**), indicating that this

transporter can serve as both a carbon and 'reductant shuttle' which would help sustain

photosynthetic electron flow.

Studies of photosynthetic activities and growth of WT and *tpt3* mutants (e.g. *t3ko2*) in LL, ML

and HL support the idea that the mutant is highly compromised in its ability to export fixed carbon

and potentially also reductant from the chloroplast. In *t3ko2* exposed to LL (after growth in LL),

growth was slow (**Fig. 2E**) and the PQ pool (**Fig. 4C**) was more reduced than in WT cells, while there was little loss of PSII or PSI activities (**Fig. 4A, B, E, F**). These results suggest that there is a reduced rate of PQH2 oxidation. During ML exposure, the mutant stopped growing, and when placed in HL experienced severe bleaching (**Fig. 2F**). The highly reduced PQ pool and PSI reaction center in ML-acclimated *t3ko2* cells (**Fig. 4C, E**, respectively) reflects hyper reduction of PET and the generation of 'over-flow' electrons. The phenotypes of the *t3ko2* cells, including an elevated 1-qL (**Fig. 4C-D**), slower oxidation rate of PSI in ML-acclimated cells (**Fig. 4F**), accumulation of storage carbon (**Fig. 3A-D**), an increase in intracellular accumulation of triose-P and 3-PGA (**Fig. 3E**), light dependent damage to the photosynthetic apparatus (**Fig. 4B**) and elevated production/accumulation of ROS (**Fig. 5A, B**), especially in the chloroplast (**Fig. 6B**), indicates a block on the acceptor side of PSI, which reflects the function of CreTPT3 and its central role in fixed carbon export from the chloroplast and for fueling central metabolism. Furthermore, the inability to efficiently transport fixed carbon from the chloroplast in ML and HL would result in diminished CBBC activity, diminished photosynthetic electron transport and the hyper-reduction of the stroma, which would also result in diminished ATP synthesis and NADPH production.

The malate-OAA shuttle represents another route that, under high redox stress, might have partially compensated for the loss of the CreTPT3 by transporting reductant from the chloroplast (schematic in **Fig. 7**). Intriguingly, malate levels in *t3ko2* were 4-fold lower than in WT cells (**Fig. 3E**, **G**). Moreover, expression of the plastid localized malate dehydrogenases (Cre*MDH1* and Cre*MDH5*) was 3-fold to 5-fold lower in the mutant than in WT cells following a transition from LL to ML (**Supplementary Fig. 10D-E**), indicating that the malate-OAA shuttle is likely unable to compensate for a loss of CreTPT3. Inactivation of CreTPT3 appears to have a negative impact on the malate-OAA shuttle, potentially because of the compromised physiological state of the *t3ko2* mutant. Furthermore, a previous study of metabolic flux analysis during heterotrophic growth of Chlamydomonas showed that the CreTPT shuttle(s) is almost 10-fold more active than the malate-OAA shuttle (Boyle et al., 2017).

TPT deficiency in Chlamydomonas cannot be compensated for by a day/night regime

TPT deficiency in plants can be almost fully compensated for by the starch-mediated night pathways that elicit the breakdown of starch and the export of the breakdown products via the maltose transporter (MEX1) and glucose translocator (GlcT) (Niittylä et al., 2004; Cho et al., 2011).

Furthermore, in plants, starch turnover may also be occurring in the light, at the same time as starch

is being synthesized (Häusler et al., 1998; Walters et al., 2004). Compared to algae, plant cells

appear to display a high plasticity in their capacity to transport fixed carbon between the

chloroplast and cytosol. This high degree of plasticity in plants is reflected by the findings: i) most

dicots contain a larger number of pPTs (from 5 to 16) (Bockwoldt et al., 2019). For example,

Arabidopsis contains 6 pPTs, which includes one TPT, two GPTs, two PPTs and one XPT

(Bockwoldt et al., 2019), whereas Chlamydomonas harbors four pPTs, which include two TPTs,

one putative PPT (TPT10 in this study) and one putative GPT/XPT (CGL51 in this study); ii) the

plant TPTs play an important role in the export of carbon from the chloroplast during the day.

However, XPT has been shown to transport triose-Ps and partially compensate for the loss of TPTs

under both ML and HL conditions (Eicks et al., 2002; Hilgers et al., 2018b). Hence, it appears that

the paths for fixed carbon export in plants are cooperative, with contributions of transporters that

use various sugars, sugar phosphates and triose phosphates.

Additionally, the elevated starch content in the *t3ko2* mutant during growth in both LL and ML

compared to WT cells suggests that the lesion creates a bottleneck in the export of fixed carbon,

which in part becomes stored as starch and lipids (**Fig. 3A-D, Supplementary Fig. 8A**). The

conversion of triose-P to starch releases Pi within the chloroplast that can at least partially

compensate for the shortage of Pi caused by inactivation of CreTPT3 (Börnke and Sonnewald,

2011). Thus, photosynthetic carbon assimilation can be maintained until the cellular system is

completely compromised. In this study, we found that Chlamydomonas *tpt3* mutants exhibited

severe growth retardation and the accumulation of starch and lipid in either continuous light (CL)

or when experiencing a day/night regime (**Fig. 2D-G** and **8A**). Furthermore, the light-induced

electron transport rate (ETR) through PSII in the *t3ko2* mutant maintained on a diurnal cycle was

similar to that of WT cells exposed to actinic light intensities of up to 200 μmol photons m^{-2} s^{-1},

but was approximately 30% lower than that of WT cells at a light intensity of 400 μmol photons

m^{-2} s^{-1} (**Fig. 8B**). We also observed that diurnally maintained *t3ko2* cells grew slightly better than

when the cells were maintained in CL (**Fig. 8A**), indicating that the loss of CreTPT3 might be

partially compensated for by starch turnover during the night (or allow for some repair of cellular

damage that might accumulate during the day), but to a lesser extent than in plants. We observed

night-time starch degradation in *tpt3* cells, although immediately following the night period a

higher level (~5-fold) of undegraded starch remained relative to that of WT cultures (**Fig. 8C**).

These data suggest that starch mobilization may partially compensate for a CreTPT3 deficiency in
Chlamydomonas.

The mechanism by which starch breakdown products are exported from the chloroplast during the
night in Chlamydomonas remains largely unknown, although it could involve various transporters
including MEX1 (like in plants). Chlamydomonas MEX1 can transport starch break down
products in the form of glucose and/or glucose phosphate, but there is no evidence suggesting that
it can export maltose since a *mex1* mutant did not accumulate maltose or exhibit growth
impairment (Findinier et al., 2017). We speculate that the loss of TPT3 might be partially
compensated for by starch turnover in the dark, with degradation products exported via MEX1 or
other pPTs that are highly expressed in the dark and downregulated in the light, such as CreTPT10
and CGL51 (**Supplementary Fig. 1D**). However, even if hexose-P is exported from the
chloroplast, it may not be readily converted to triose-P (see below), which would fuel the TCA
cycle and respiration and serve to support anabolic processes.

t3ko2 **experiences oxidative stress**

In plants, the export of sugars and other molecules (e.g. redox equivalents, ROS) can serve as
signals that coordinate chloroplast and nuclear gene expression during acclimation to HL (Häusler
et al., 2014; Zirngibl et al., 2022). We probed the impact of impaired triose-P export in *t3ko2* on
ROS production and accumulation in various subcellular compartments following exposure of the
cells to ML or HL. In plant cells, H_2O_2 is produced as a side-product of cellular processes including
PET, mitochondrial respiration, and substrate level oxidation (Smirnoff and Arnaud, 2019; Foyer
and Noctor, 2016). While ROS stability is generally low, it can accumulate in cells experiencing
oxidative stress, with H_2O_2 being the most prevalent species that can function as a redox messenger
(Li and Kim, 2022). Moreover, the trafficking of H_2O_2 into or out of different cellular
compartments can trigger activation of other retrograde and anterograde signals that may
coordinate activities among the compartments, including the nucleus (Exposito-Rodriguez et al.,
2017; Mittler et al., 2022; Shapiguzov et al., 2012). Upon exposure to HL, the *t3ko2* mutant
accumulated more stromal H_2O_2 than WT cells. The cytosolic H_2O_2 levels in the mutant exhibited
a similar increase, which may reflect the ability of this metabolite to rapidly diffuse from the
chloroplast and into the cytoplasm (**Fig. 6B, C**). Furthermore, it is unlikely that the mitochondrion
contributes to an increase in H_2O_2 in *t3ko2* since no (or little) increase in accumulation of

intramitochondrial H_2O_2 was observed in the mutant in either LL or ML (**Fig. 6D**). An increase in H_2O_2 accumulation in the nucleus of *t3ko2* relative to WT cells was also observed, although the light intensity (HL or LL) did not alter these levels in either WT or the mutant (**Fig. 6E**); a previous report showed that the nuclear H_2O_2 level was not significantly affected in WT Chlamydomonas cells following a HL exposure (Niemeyer et al., 2021), which may reflect both the level of H_2O_2 accumulation and barriers that limit its diffusion. The higher levels of H_2O_2 in the nucleus of *t3ko2* cells may trigger retrograde signals that modulate nuclear gene expressions, which in turn could ameliorate some of the negative effects of ROS and elicit repair of any damage experienced by the photosynthetic apparatus. A similar response may be elicited in WT cells at higher intensity actinic light.

Superoxide is predominantly produced by PSI, which can enzymatically be converted into H_2O_2 (Foyer, 2018). Highly reduced PSI in the *tpt3* mutant can lead to an increase in the production of H_2O_2. It was also shown that H_2O_2 can be synthesized in thylakoid membranes as a consequence of the oxidation of plastoquinol (PQH_2), suggesting a positive correlation between the redox state of the PQ pool and the generation of H_2O_2 (Khorobrykh et al., 2015). A similar finding was noted for both *Nicotiana benthamiana* and Chlamydomonas based on the use of hypersensitive H_2O_2 sensors (Exposito-Rodriguez et al., 2017; Niemeyer et al., 2021). These two organisms were shown to accumulate more stromal H_2O_2 in HL, which was dependent on photosynthesis. We observed this positive correlation between PQ pool reduction and the accumulation of H_2O_2 in the *t3ko2* mutant; the PQ pool was more reduced in *t3ko2* relative to the WT cells after exposing the cells to 105 s of HL (**Fig. 4D**). In parallel, there was a marked increase in stromal H_2O_2 following 2.5 min of HL (**Fig. 6B**). Therefore, hyper-reduction of the PQ pool in the mutant likely results in elevated stromal H_2O_2 accumulation, suggesting that CreTPT3 activity and the export of triose-P from the chloroplast is critical for maintaining low level synthesis/accumulation H_2O_2 and sustaining a high rate of PET in HL; the export of fixed carbon relieves the redox pressure and lessens ROS formation. Additionally, Cre*TPT3* is the most nutrient-deprivation responsive/upregulated of the *pPT* family genes; it responds strongly to nitrogen, sulfur, and iron deprivation (**Supplementary Fig. 9 B-D**). These findings are in accord with the hypothesis that the ability to traffic fixed carbon from the chloroplast is important for both the distributing carbon to other cellular compartments and relieving oxidative stress in the organelle.

CreTPT3 is critical for maintaining intracellular partitioning of fixed carbon

Why is the phenotype of the Chlamydomonas *tpt3* mutant so severe? Land plants contain the entire glycolytic pathway in both the chloroplast and cytosol while the pathway is partitioned between two compartments in Chlamydomonas; 90% of the upper activities of the pathways (from F6P→ 3PG) was associated with the plastid while over 95% of the activities of the lower part of the pathway (3PG→Pyruvate) occurred in the cytosol (Klein, 1986; Rochaix et al., 1998). The oxidative pentose phosphate pathway also appears to be in the chloroplast (Klein, 1986). The partitioning of glycolysis between the chloroplast and cytosol is supported by the comparative quantification of the metabolites, with glucose-1-P, fructose-6-P, and fructose-1,6-P2 being exclusively in the chloroplast and 2-phosphoglycerate only in the cytosol (Klöck and Kreuzberg, 1991). Recently, it was suggested that the flux of metabolites through hexose-P is negligible in the Chlamydomonas cytosol (Treves et al., 2022), possibly because of the absence of glycolytic reactions that would facilitate its metabolism. Therefore, even if hexose-P is exported from the chloroplast, it would likely not be rapidly metabolized or maintain rapid cell growth. Overall, the results strongly suggest that triose-P exported from the Chlamydomonas chloroplast is likely the major source of fixed carbon transported into the cytoplasm of the cell, facilitating algal growth in the light.

Thus, we hypothesize that the export of triose-P would drive the cytosolic segment of glycolysis and downstream metabolic pathways. This hypothesis is supported by the metabolite analysis; specifically, a marked increase of most metabolites associated with the upper-glycolytic/gluconeogenic pathways, and a significant decrease of some metabolites of the TCA cycle upon exposure of the mutant cells to either LL or ML (**Fig. 3E-G, Supplementary Fig. 8B, C**). In contrast, elimination of the chloroplast-targeted TPT1 protein of Arabidopsis showed no significant phenotype, although growth was retarded in the TPT/XPT double mutants (Hilgers et al., 2018a). Furthermore, based on Pearson correlation analyses presented in **Supplementary Fig. 11** and **Table. 4**, Cre*TPT3* is co-expressed with many genes involved in respiratory electron transport and the major ATP transporters located on the mitochondria and chloroplast envelope membranes. In addition, transcript levels of some genes involved in starch degradation, glycolysis, the TCA cycle, and malate/OAA shuttle shared a high correlation coefficient with Cre*TPT3*. Together, these data indicate that the export of triose-P from the chloroplast is closely linked to

central energy metabolism in Chlamydomonas, starting with the production of triose-P in the

chloroplast by the CBBC or starch degradation (chloroplast localized reactions of glycolysis),

followed by transport to the cytosol which houses the remaining reactions of glycolysis. The

products of glycolysis can be trafficked to the mitochondria where they can be used to drive the

TCA cycle, respiratory metabolism, and the generation of ATPs (**Supplementary Fig. 11**). Hence,

triose-P is the major photoassimilate routed from the chloroplast, supplying substrates for

downstream metabolic processes.

Summary

As depicted in **Fig. 7**, we propose that various tiers of regulation are responsible for the

physiological responses of the Chlamydomonas *t3ko2* mutant. When *t3ko2* cells are transferred

from LL to ML, the triose-P pool and metabolites derived from that pool accumulate because of

the reduced capacity of the strain to move triose-P out of the chloroplast where it could be further

metabolized. Some compensation may occur through the activity of other transporters, although

expression of Cre*TPT2* is especially low during the day (in the light) and the transport of hexose

phosphate may not compensate for the loss of CreTPT3 because the cytoplasm does not have (or

has little of) the activities of glycolysis that would convert hexose-P to DHAP. The compromised

ability to export fixed carbon from the chloroplast also suppresses CBBC activity, causes hyper-

reduction of PET and the accumulation of ROS (which would inhibit photosynthetic activity). The

highly diminished export of triose-P to the cytoplasm would compromise respiration and

downstream biosynthetic processes. Furthermore, hyper-reduction of PET and accumulation of

carbon metabolites in the stroma would activate AGPase through allosteric regulation and by the

FDX/TRX (ferredoxin/thioredoxin) and NTRC (NADPH-dependent thioredoxin reductase C)

redox systems, which would result in starch hyper-accumulation (Ballicora et al., 2000; Lepistö et

al., 2013) (**Fig. 7**). Increased ROS accumulation in the mutant chloroplasts and an elevated PET

redox state would also elicit the generation of retrograde signals that mediate changes in nuclear

gene expression (Wakao and Niyogi, 2021; Suzuki et al., 2012; Shapiguzov et al., 2012),

stimulating the synthesis of specific activities that may function to ameliorate the impact of the

hyper-reduced state attained in the chloroplast.

Material and methods

Strains and culture conditions

WT Chlamydomonas M10 (CC-4403, isogenic line derived from CC-124) was used as the parental strain for the generation of knockout mutants. Cultures were routinely cultivated in growth chambers (LED-41L2, Percival Scientific, Inc.) at 25°C with continuous shaking on an orbital shaker (VWR OS-500 Shaker) at 120 rpm, in Tris-Acetate-Phosphate (TAP) medium (Harris, 2009). Cultures were illuminated with continuous cool white LEDs (LED-41L2, Percival Scientific, Inc.) at low light (LL, 30 μmol photons m^{-2} s^{-1}). Experiments were mostly performed with cells grown in the photoautotrophic TP medium (TAP medium without acetate) and in some cases, in TAP medium, at 25°C, and sparged with air while being shaken at 120 rpm in a growth chamber (LED-41L2, Percival Scientific, Inc.). For growth assays, cultures were inoculated to a density of 0.02 at $OD_{750 nm}$ (~1×10^5 cells/mL) in TP sparged with air under either LL or moderate light (ML, 250-300 μmol photons m^{-2} s^{-1}) intensities. For spectrophotometric and chlorophyll fluorescence analyses, the experimental design is described in **Supplementary Fig. 5**. Growth assays on solid medium were performed with cultures spotted onto the medium at different dilutions (indicated in the text) and exposed to different light intensities; spot tests for photoautotrophic growth were on TP agar plates and for mixotrophic growth on TAP agar plates. Agar plates were incubated for 7 d under either LL or ML (cool white LED) at 25°C.

Reconstitution into liposomes and transport assays

The procedures for the construction and expression of Cre*TPT2* and Cre*TPT3* in yeast are described in supplemental information (SI). For uptake studies, yeast membranes from cells with and without recombinant His-tagged CreTPT2 or CreTPT3 were enriched and reconstituted into 3% (w/v) L-alpha-phosphatidylcholine using a freeze-thaw-sonication procedure (Linka et al., 2008). The reconstituted liposomes were preloaded with 30 mM Pi or phosphorylated metabolites to be tested as potential transport substrates. As a negative control for antiport activity, liposomes were also generated without metabolite preloading. The external counter-exchange substrate was removed via gel filtration on Sephadex G-25M columns (GE Healthcare). Transport assays were initiated by adding 0.25 mM [alpha-^{32}P]-phosphoric acid (6,000 Ci/mmol) to the medium bathing the liposomes and performed as previously described by Linka et al. (2008). Measurements of Km for Pi and competitive inhibition constants (Ki) are described in supplemental information (SI).

Vector construction, transformation, and subcellular localization

The pRam118_VENUS plasmid, which harbors the VENUS gene and the *AphVII* cassette (confers resistance to hygromycin) (Cre01.g045550)(Kaye et al., 2019), was used to express Cre*TPT2* (Cre06.g263850_4532) and Cre*TPT3* (Cre01.g045550_4532). The step-by-step description is in SI. 2-4 μg of the engineered plasmids was linearized and added to 250 μL of a cell suspension of $\sim3\times10^8$ cells/mL per reaction. The GeneArt MAX Efficiency Transformation Reagent for algae (Invitrogen) was used for introducing the plasmid into the algal cells by electroporation according to the instructions provided by the manufacturer. Transformants were selected on solid TAP medium containing hygromycin (10 μg/mL; Enzo Life).

Drug resistant transformants were visualized for VENUS fluorescence as previously described (Kaye et al., 2019). In brief, transgenic cell lines resistant to hygromycin were screened for VENUS fluorescence using a microplate reader (Infinite M1000; TECAN). Excitation and emission settings were: VENUS, excitation at 515 nm, bandwidth 12 nm and emission at 550 nm, bandwidth 12 nm; chlorophyll excitation was at 440 nm, bandwidth 9 nm and emission was at 680 nm, bandwidth 20 nm. The TCS SP8 confocal laser-scanning microscope (Leica) was used to visualize the VENUS fluorescence signal (Kaye et al., 2019).

CRISPR-CAS9 mediated mutagenesis

The Chlamydomonas WT strain CC-124 was used for mutant generation. WT cells were cultured under CL (50 μmol photons m^{-2} s^{-1}) for 2 d to a density of $3\text{-}5\times10^6$ cells/mL. The cells were then concentrated to 2×10^8 cells/mL in 0.5×TAP medium supplemented with 80 mM sucrose. Two single guide RNAs (sgRNA) were designed by CHOPCHOP and synthesized by Integrated DNA Technologies (IDT). The sequences of the generated sgRNAs are: *TPT2*-sg (5′-AUAAGGGCAAGGACAUGUCAGGG -3) for editing exon 8, *TPT3*-sg1 (5′-CGCUGGGCGTCACUUCCCGGCGG -3′) for editing exon 1, and *TPT3*-sg2 (5′-AAGGCCGCUAUCGCCAACGUGGG -3′) for editing of exon 7. The protocol for disruption of Cre*TPT2* and Cre*TPT3* was adapted from (Findinier et al., 2019) and is described in SI.

Complementation of mutants *tpt3* mutants

Mutant strains were transformed with the linearized pRam118_CreTPT3 plasmid. Transformed cells were selected in ML and screened for VENUS fluorescence. Colonies exhibiting VENUS

fluorescence and an *AphVII* cassette knock-in at the CAS9 target site were examined for
accumulation of the CreTPT3 protein by immunodetection using CreTPT3 antibodies generated
by GenScript USA Inc (Piscataway, USA). Immuno-positive colonies were subjected to growth
assays using spot tests under ML on either solid TAP or TP medium.

P700 activity measurements

Absorbance spectroscopy [JTS-100 spectrophotometer (SpectroLogiX, TN)] to measure P700
activity was performed with dark-adapted liquid cultures (15 μg/mL chlorophyll, in 20 mM
HEPES-KOH, pH 7.2, and 10% ficoll) as previously described (Clowez et al., 2021). Actinic light
was provided by an orange LED (165 μmol photons m^{-2} s^{-1}) for PSI oxidation, followed by a
saturating pulse and dark incubation. 20 μM DCMU and 1 mM hydroxylamine were added to the
cell suspension to inhibit linear electron flow (LEF) prior to the measurement. P700 activity was
measured by monitoring the absorbance at 705 nm (interference filter 6 nm FWHM was used to
create a narrow excitation beam); the absorbance at 740 nm was used to correct for an unspecific
contribution to the 705 nm signal.

Chlorophyll fluorescence analysis

Chlorophyll fluorescence used to evaluate photosynthetic electron transport was monitored with a
DUAL PAM-100 fluorometer. Cells were acclimated in the dark for 20 min prior to illumination
at increasing light intensities (0, 10, 50, 100, 200, 400, 800, 1000, 1200 μmol photons m^{-2} s^{-1}) for
2 min at each intensity, or at a constant intensity of 450 μmol photons m^{-2} s^{-1} for 10 min to evaluate
1-qL. 1 mM CO_2 ($NaHCO_3$) was added to the reaction mix as an electron acceptor for the CBBC.

ROS measurements and roGFP2 imaging analysis

All strains were grown photoautotrophically in LL for 16-24 h. After dilution with fresh medium,
the cultures were transferred to ML and stained with CM-H2DCFDA (Thermo Fisher Scientific)
as described in (Kong et al., 2018) for detecting ROS. A step-by-step protocol is given in SI.

Constructs containing chloroplast or mitochondria targeting sequences (Crozet et al., 2018) fused
to codon-optimized roGFP2 (Vevea et al., 2013) were transformed into WT and mutant strains.
Transgenic cell lines were screened for green roGFP2 fluorescence using a microplate reader
(Infinite M1000; TECAN); excitation was at 488 nm, bandwidth 9 nm and emission at 525 nm,

bandwidth 10 nm. Cells with strong green fluorescence were cultured as depicted in **Supplementary Fig. 5**. Signals from the transformed lines were visualized using a TCS SP8 confocal laser-scanning microscope (Leica). roGFP2 signals were collected and analyzed as previously described (Vevea et al., 2013). Using the sequential setup of the SP8, roGFP2 signals were collected at the emission wavelength of 510-550 nm immediately following excitation at 405 nm and 488 nm (Vevea et al., 2013). The degree of roGFP2 oxidation was analyzed as the ratio of the emission signals after excitation at 405 nm and 488 nm (Vevea et al., 2013).

Plasmids harboring H_2O_2 roGFP2-Tsa2ΔCR sensors (Niemeyer et al., 2021) were obtained from the Chlamydomonas Resource Center (https://www.chlamycollection.org/) (**Supplementary Table. 3)**. Cells in which roGFP2-Tsa2ΔCR was targeted to the stroma, cytosol, mitochondrial matrix, or nucleus were initially grown in LL in TAP to exponential growth phase. Cells were then grown in TP in LL for 2 h and shifted either to HL or very low light (10 μmol photons m^{-2} s^{-1}). Prior to collecting the algal samples, N-Ethylmaleimide (NEM) (in 100-mM MES–Tris buffer, pH 7.0) was added to a centrifuge tube to trap the oxidation state of the sensor. The final concentration of NEM is 10 mM after adding the algae culture. After centrifugation (1459 x g, 3 min), cells were resuspended in 100-mM MES–Tris buffer (pH 7.0) to reach the chlorophyll concentration of 30 μg/mL; and roGFP2 fluorescence was measured in a plate reader (Infinite M1000; TECAN). For calibration and data calculation, fully oxidized sensors in the control samples were prepared by adding H_2O_2 to a final concentration of 5 mM, and fully reduced sensors were prepared by adding DTT to a final concentration of 100 mM. Signals were detected using the excitation wavelengths of 410 and 488 nm and the emission wavelength of 514 nm. The degree of sensor oxidation (OxD) was calculated using the previous equation as described in (Niemeyer et al., 2021).

Photosynthesis-Irradiance curve

Photosynthesis-Irradiance curves were measured using a custom Pt-Ag/AgCl polarographic electrode system (ALGI) with a water jacketed (for temperature-control), 1 mL glass reaction chamber. A step-by-step protocol is provided in SI.

Starch and TAG quantification

Cells were grown under LL to mid-exponential phase, diluted to 0.5 μg/mL chl with fresh medium and transferred to and then grown in ML for 24 h or more, as indicated. Starch was measured

according to (Klein and Betz, 1978), with slight modification (see SI). Triacylglycerol (TAG)

levels were estimated by a fluorometric assay using the dye Nile Red (Thermo Fisher)(Yu et al.,

2009). A Nile Red solution (500 μg/mL in acetone) was added to 1 mL of the cell suspensions to

a final concentration of 0.5 μg/mL. Samples were then incubated at room temperature for 30 min,

and the Nile Red fluorescence emission quantified at 575 nm following excitation at 530 nm using

a microplate reader (Infinite M1000; TECAN).

Metabolic analysis

Cells grown in LL were shifted to ML for 45 min or 6 h, as indicated. 45 mL of culture was rapidly

quenched in cold saline solution (-2 to -3°C), extracted using cold methanol and then analyzed by

LC-MS/MS. Quenching and analysis of metabolites were performed, with modifications,

according to (Sake et al., 2020). A step-by-step protocol is described in SI. Metabolite extracts

were analyzed using LC-MS/MS, as adapted from (Young et al., 2011). A Phenomenex 150 mm

x 2 mm Synergi Hydro-RP column was used on an Agilent 1200 Series HPLC system coupled to

an AB Sciex 5500 QTRAP system. LC was performed with a liquid injection volume of 20 μL and

a gradient elution with 10 mM tributylamine and 15 mM acetic acid (aqueous phase) in acetonitrile

(organic phase) (reagent B) at a constant flow rate of 0.3 mL/min, and a constant temperature of

40°C. The gradient profile of the organic phase is as follows: 0% B (0 min), 8% B (10 min), 16%

B (15 min), 30% B (16.5 min), 30% B (19 min), 90% B (21.5 min), 90% B (28 min), 0% B (28.5

min), and 0% B (35 min). MS analysis was performed in negative mode using a multiple reaction

monitoring (MRM) acquisition method. Data acquisition was performed on the ABSciex Analyst

1.7 software. Absolute quantification of intracellular metabolites was performed using the

quantitation mode on the Analyst software. All chemicals used for metabolite extraction and LC-

MS/MS analysis were Optima grade reagents.

Acknowledgments:

WH thanks Dr. Masayuki Onishi for advice on the generation of CRISPR mutants and Adrien Burlacot for help with PQ pool measurements.

Funding:

This project has supported by DOE award DE-SC0019417 to ARG. WH was supported solely by a grant from the US Department of Energy, Office of Science, Basic Energy Systems (DE-SC0019417). AK and MCP were supported solely by a grant from the US Department of Energy, Office of Science, Basic Energy Systems (DE-SC0019341 to MP). NB and MM were supported by US Department of Energy, Office of Science, Office of Biological and Environmental Research (BER), (DE-SC0018301 to NB). NL and AP were supported by German Research Foundations (DFG Grant LI1781/2-2). BR was supported by National Science Foundation (Award Number 1645164). RGK and JF were supported by the Department of Plant Biology of the Carnegie Institution for Science. PR was supported by Human Frontier Science Program (HFSP) RGP0046/2018 (to ARG).

Author contributions:

WH, AK, ARG and MP conceptualized the study. WH generated the CRISPR mutants as well as the complemented strain, localized the TPT2 and 3 proteins, analyzed the photosynthetic performances and starch changes over diel cycle. AK performed the experiments of photosynthetic O_2 evolution, extraction of samples for LC-MS/MS, data analysis of LC-MS/MS. AP and NL performed the reconstitution into liposomes and transport activity assays. MM and NB performed LC-MS/MS and analyses of metabolite data. WH and YW analyzed the redox status in the cell. BR performed data mining for transcriptome and the Nile red staining. JF and NF analyzed starch changes upon transition from LL to ML. WH, PR and ARG analyzed PSI and PSII activities. JF and BR constructed roGFP2 associated vectors. WH and ARG wrote the manuscript. ARG, MP, AK, NL, NB, AP, MM and all the other authors helped writing and revising the manuscript.

Competing interests:

All authors declare they have no competing interests.

Distribution of materials:

All manuscripts must include the following statement as an unnumbered footnote: "The author(s) responsible for distribution of materials integral to the findings presented in this article in accordance with the policy described in the Instructions for Authors (https://academic.oup.com/plcell/pages/General-Instructions) are: Weichao Huang (whuang@carnegiescience.edu) and Arthur Grossman (agrossman@carnegiescience.edu).

References

Badger, M.R., von Caemmerer, S., Ruuska, S., and Nakano, H. (2000). Electron flow to oxygen in higher plants and algae: rates and control of direct photoreduction (Mehler reaction) and rubisco oxygenase. Philos. Trans. R. Soc. Lond. B Biol. Sci. **355**: 1433–1446.

Ballicora, M.A., Frueauf, J.B., Fu, Y., Schürmann, P., and Preiss, J. (2000). Activation of the potato tuber ADP-glucose pyrophosphorylase by thioredoxin. J. Biol. Chem. **275**: 1315–1320.

Bhatt, P., Bhandari, G., Turco, R.F., Aminikhoei, Z., Bhatt, K., and Simsek, H. (2022). Algae in wastewater treatment, mechanism, and application of biomass for production of value-added product. Environ. Pollut. **309**: 119688.

Bockwoldt, M., Heiland, I., and Fischer, K. (2019). The evolution of the plastid phosphate translocator family. Planta **250**: 245–261.

Börnke, F. and Sonnewald, S. (2011). Biosynthesis and metabolism of starch and sugars. In Plant Metabolism and Biotechnology (John Wiley & Sons, Ltd: Chichester, UK), pp. 1–25.

Boyle, N.R., Sengupta, N., and Morgan, J.A. (2017). Metabolic flux analysis of heterotrophic growth in *Chlamydomonas reinhardtii*. PLoS One **12**: e0177292.

Chaux, F., Burlacot, A., Mekhalfi, M., Auroy, P., Blangy, S., Richaud, P., and Peltier, G. (2017). Flavodiiron proteins promote fast and transient O_2 photoreduction in Chlamydomonas. Plant Physiol. **174**: 1825–1836.

Cho, M.-H., Lim, H., Shin, D.H., Jeon, J.-S., Bhoo, S.H., Park, Y.-I., and Hahn, T.-R. (2011). Role of the plastidic glucose translocator in the export of starch degradation products from the chloroplasts in *Arabidopsis thaliana*. New Phytol. **190**: 101–112.

Clowez, S., Renicke, C., Pringle, J.R., and Grossman, A.R. (2021). Impact of menthol on growth and photosynthetic function of *Breviolum minutum* (Dinoflagellata, Dinophyceae, Symbiodiniaceae) and Interactions with its Aiptasia Host. J. Phycol. **57**: 245–257.

Crozet, P. et al. (2018). Birth of a photosynthetic chassis: A MoClo toolkit enabling synthetic biology in the microalga *Chlamydomonas reinhardtii*. ACS Synth. Biol. **7**: 2074–2086.

Dorion, S., Ouellet, J.C., and Rivoal, J. (2021). Glutathione metabolism in plants under stress: beyond reactive oxygen species detoxification. Metabolites **11**: 641.

Eicks, M., Maurino, V., Knappe, S., Flügge, U.-I., and Fischer, K. (2002). The plastidic pentose phosphate translocator represents a link between the cytosolic and the plastidic pentose phosphate pathways in plants. Plant Physiol. **128**: 512–522.

Exposito-Rodriguez, M., Laissue, P.P., Yvon-Durocher, G., Smirnoff, N., and Mullineaux, P.M. (2017). Photosynthesis-dependent H_2O_2 transfer from chloroplasts to nuclei provides a high-light signalling mechanism. Nat. Commun. **8**: 49.

Findinier, J., Delevoye, C., and Cohen, M.M. (2019). The dynamin-like protein Fzl promotes
thylakoid fusion and resistance to light stress in *Chlamydomonas reinhardtii*. PLoS Genet.
15: e1008047.

Fischer, K., Kammerer, B., Gutensohn, M., Arbinger, B., Weber, A., Hausler, R.E., and
Flugge, U.-I. (1997). A new class of plastidic phosphate translocators: A putative link
between primary and secondary metabolism by the phosphoenolpyruvate/phosphate
antiporter. Plant Cell **9**: 453.

Fliege, R., Flügge, U.-I., Werdan, K., and Heldt, H.W. (1978). Specific transport of inorganic
phosphate, 3-phosphoglycerate and triosephosphates across the inner membrane of the
envelope in spinach chloroplasts. Biochim. Biophys. Acta Bioenerg. **502**: 232–247.

Flügge, U.I., Fischer, K., Gross, A., Sebald, W., Lottspeich, F., and Eckerskorn, C. (1989).
The triose phosphate-3-phosphoglycerate-phosphate translocator from spinach
chloroplasts: nucleotide sequence of a full-length cDNA clone and import of the in vitro
synthesized precursor protein into chloroplasts. EMBO J. **8**: 39–46.

Flügge, U.I., Häusler, R.E., Ludewig, F., and Fischer, K. (2003). Functional genomics of
phosphate antiport systems of plastids. Physiol. Plant. **118**: 475–482.

Foyer, C.H. (2018). Reactive oxygen species, oxidative signaling and the regulation of
photosynthesis. Environ. Exp. Bot. **154**: 134–142.

Foyer, C.H. and Noctor, G. (2016). Stress-triggered redox signalling: what's in pROSpect? Plant
Cell Environ. **39**: 951–964.

Ge, F., Huang, W., Chen, Z., Zhang, C., Xiong, Q., Bowler, C., Yang, J., Xu, J., and Hu, H.
(2014). Methylcrotonyl-CoA carboxylase regulates triacylglycerol accumulation in the
model diatom *Phaeodactylum tricornutum*. Plant Cell **26**: 1681–1697.

Gibson, D.G., Young, L., Chuang, R.-Y., Venter, J.C., Hutchison, C.A., 3rd, and Smith, H.O.
(2009). Enzymatic assembly of DNA molecules up to several hundred kilobases. Nat.
Methods **6**: 343–345.

Gietz, R.D. and Schiestl, R.H. (2007). High-efficiency yeast transformation using the LiAc/SS
carrier DNA/PEG method. Nat. Protoc. **2**: 31–34.

González-Ballester, D., Casero, D., Cokus, S., Pellegrini, M., Merchant, S.S., and Grossman,
A.R. (2010). RNA-seq analysis of sulfur-deprived Chlamydomonas cells reveals aspects
of acclimation critical for cell survival. Plant Cell **22**: 2058–2084.

Harris, E.H. (2009). Chlamydomonas sourcebook: Introduction to Chlamydomonas and its
laboratory use 2nd ed. (Academic Press).

Häusler, R.E., Baur, B., Scharte, J., Teichmann, T., Eicks, M., Fischer, K.L., Flügge, U.I.,
Schubert, S., Weber, A., and Fischer, K. (2000). Plastidic metabolite transporters and

870   their physiological functions in the inducible crassulacean acid metabolism plant
871   Mesembryanthemum crystallinum. Plant J. **24**: 285–296.

872 **Häusler, R.E., Heinrichs, L., Schmitz, J., and Flügge, U.-I.** (2014). How sugars might
873   coordinate chloroplast and nuclear gene expression during acclimation to high light
874   intensities. Mol. Plant **7**: 1121–1137.

875 **Häusler, R.E., Schlieben, N.H., Schulz, B., and Flügge, U.I.** (1998). Compensation of decreased
876   triose phosphate/phosphate translocator activity by accelerated starch turnover and glucose
877   transport in transgenic tobacco. Planta **204**: 366–376.

878 **Helman, Y., Tchernov, D., Reinhold, L., Shibata, M., Ogawa, T., Schwarz, R., Ohad, I., and**
879   **Kaplan, A.** (2003). Genes encoding A-type flavoproteins are essential for photoreduction
880   of O2 in cyanobacteria. Curr. Biol. **13**: 230–235.

881 **Hilgers, E.J.A., Schöttler, M.A., Mettler-Altmann, T., Krueger, S., Dörmann, P., Eicks, M.,**
882   **Flügge, U.-I., and Häusler, R.E.** (2018a). The combined loss of triose phosphate and
883   xylulose 5-Phosphate/Phosphate translocators leads to severe growth retardation and
884   impaired photosynthesis in *Arabidopsis thaliana* tpt/xpt double mutants. Front. Plant Sci.
885   **9**: 1331.

886 **Hilgers, E.J.A., Staehr, P., Flügge, U.-I., and Häusler, R.E.** (2018b). The Xylulose 5-
887   Phosphate/Phosphate Translocator Supports Triose Phosphate, but Not
888   Phosphoenolpyruvate Transport Across the Inner Envelope Membrane of Plastids in
889   Arabidopsis thaliana Mutant Plants. Front. Plant Sci. **9**: 1461.

890 **Houille-Vernes, L., Rappaport, F., Wollman, F.-A., Alric, J., and Johnson, X.** (2011). Plastid
891   terminal oxidase 2 (PTOX2) is the major oxidase involved in chlororespiration in
892   Chlamydomonas. Proc. Natl. Acad. Sci. U. S. A. **108**: 20820–20825.

893 **Huang, W., Haferkamp, I., Lepetit, B., Molchanova, M., Hou, S., Jeblick, W., Río Bártulos,**
894   **C., and Kroth, P.G.** (2018). Reduced vacuolar β-1,3-glucan synthesis affects carbohydrate
895   metabolism as well as plastid homeostasis and structure in *Phaeodactylum tricornutum*.
896   Proc. Natl. Acad. Sci. U. S. A. **115**: 4791–4796.

897 **Jack, D.L., Yang, N.M., and Saier, M.H., Jr** (2001). The drug/metabolite transporter superfamily.
898   Eur. J. Biochem. **268**: 3620–3639.

899 **Johnson, X. and Alric, J.** (2013). Central carbon metabolism and electron transport in
900   Chlamydomonas reinhardtii: metabolic constraints for carbon partitioning between oil and
901   starch. Eukaryot. Cell **12**: 776–793.

902 **Kammerer, B., Fischer, K., Hilpert, B., Schubert, S., Gutensohn, M., Weber, A., and Flügge,**
903   **U.I.** (1998). Molecular characterization of a carbon transporter in plastids from
904   heterotrophic tissues: the glucose 6-phosphate/phosphate antiporter. Plant Cell **10**: 105–
905   117.

Kaye, Y., Huang, W., Clowez, S., Saroussi, S., Idoine, A., Sanz-Luque, E., and Grossman,
A.R. (2019). The mitochondrial alternative oxidase from *Chlamydomonas reinhardtii*
enables survival in high light. J. Biol. Chem. **294**: 1380–1395.

Khan, M.I., Shin, J.H., and Kim, J.D. (2018). The promising future of microalgae: current status,
challenges, and optimization of a sustainable and renewable industry for biofuels, feed, and
other products. Microb. Cell Fact. **17**: 36.

Khorobrykh, S.A., Karonen, M., and Tyystjärvi, E. (2015). Experimental evidence suggesting
that H_2O_2 is produced within the thylakoid membrane in a reaction between plastoquinol
and singlet oxygen. FEBS Lett. **589**: 779–786.

Klein, U. (1986). Compartmentation of glycolysis and of the oxidative pentose-phosphate pathway
in *Chlamydomonas reinhardii*. Planta **167**: 81–86.

Klein, U. and Betz, A. (1978). Fermentative metabolism of hydrogen-evolving *Chlamydomonas*
moewusii. Plant Physiol. **61**: 953–956.

Klöck, G. and Kreuzberg, K. (1991). Compartmented metabolite pools in protoplasts from the
green alga *Chlamydomonas reinhardtii*: changes after transition from aerobiosis to
anaerobiosis in the dark. Biochim. Biophys. Acta **1073**: 410–415.

Knappe, S., Flügge, U.-I., and Fischer, K. (2003). Analysis of the plastidic phosphate
translocator gene family in Arabidopsis and identification of new phosphate translocator-
homologous transporters, classified by their putative substrate-binding site. Plant Physiol.
131: 1178–1190.

Kong, F., Burlacot, A., Liang, Y., Légeret, B., Alseekh, S., Brotman, Y., Fernie, A.R.,
Krieger-Liszkay, A., Beisson, F., Peltier, G., and Li-Beisson, Y. (2018). Interorganelle
communication: peroxisomal MALATE DEHYDROGENASE2 connects lipid catabolism
to photosynthesis through redox coupling in Chlamydomonas. Plant Cell **30**: 1824–1847.

Kramer, D.M., Johnson, G., Kiirats, O., and Edwards, G.E. (2004). New fluorescence
parameters for the determination of QA redox state and excitation energy fluxes.
Photosynth. Res. **79**: 209.

Krishnan, A., Kumaraswamy, G.K., Vinyard, D.J., Gu, H., Ananyev, G., Posewitz, M.C., and
Dismukes, G.C. (2015). Metabolic and photosynthetic consequences of blocking starch
biosynthesis in the green alga *Chlamydomonas reinhardtii sta6* mutant. Plant J. **81**: 947–
960.

Lee, S.-K., Eom, J.-S., Voll, L.M., Prasch, C.M., Park, Y.-I., Hahn, T.-R., Ha, S.-H., An, G.,
and Jeon, J.-S. (2014). Analysis of a triose phosphate/phosphate translocator-deficient
mutant reveals a limited capacity for starch synthesis in rice leaves. Mol. Plant **7**: 1705–
1708.

Lee, Y., Nishizawa, T., Takemoto, M., Kumazaki, K., Yamashita, K., Hirata, K., Minoda, A.,
Nagatoishi, S., Tsumoto, K., Ishitani, R., and Nureki, O. (2017). Structure of the triose-

943   phosphate/phosphate translocator reveals the basis of substrate specificity. Nat Plants **3**:
944   825–832.

945 **Lepistö, A., Pakula, E., Toivola, J., Krieger-Liszkay, A., Vignols, F., and Rintamäki, E.**
946   (2013). Deletion of chloroplast NADPH-dependent thioredoxin reductase results in
947   inability to regulate starch synthesis and causes stunted growth under short-day
948   photoperiods. J. Exp. Bot. **64**: 3843–3854.

949 **Li, M. and Kim, C.** (2022). Chloroplast ROS and stress signaling. Plant Commun **3**: 100264.

950 **Lim, L., Linka, M., Mullin, K.A., Weber, A.P.M., and McFadden, G.I.** (2010). The carbon and
951   energy sources of the non-photosynthetic plastid in the malaria parasite. FEBS Lett. **584**:
952   549–554.

953 **Linka, M., Jamai, A., and Weber, A.P.M.** (2008). Functional characterization of the plastidic
954   phosphate translocator gene family from the thermo-acidophilic red alga *Galdieria*
955   *sulphuraria* reveals specific adaptations of primary carbon partitioning in green plants and
956   red algae. Plant Physiol. **148**: 1487–1496.

957 **Loddenkötter, B., Kammerer, B., Fischer, K., and Flügge, U.I.** (1993). Expression of the
958   functional mature chloroplast triose phosphate translocator in yeast internal membranes
959   and purification of the histidine-tagged protein by a single metal-affinity chromatography
960   step. Proc. Natl. Acad. Sci. U. S. A. **90**: 2155–2159.

961 **Mittler, R., Zandalinas, S.I., Fichman, Y., and Van Breusegem, F.** (2022). Reactive oxygen
962   species signalling in plant stress responses. Nat. Rev. Mol. Cell Biol. **23**: 663–679.

963 **Moog, D., Nozawa, A., Tozawa, Y., and Kamikawa, R.** (2020). Substrate specificity of plastid
964   phosphate transporters in a non-photosynthetic diatom and its implication in evolution of
965   red alga-derived complex plastids. Sci. Rep. **10**: 1167.

966 **Ngan, C.Y. et al.** (2015). Lineage-specific chromatin signatures reveal a regulator of lipid
967   metabolism in microalgae. Nat Plants **1**: 15107.

968 **Niemeyer, J., Scheuring, D., Oestreicher, J., Morgan, B., and Schroda, M.** (2021). Real-time
969   monitoring of subcellular H_2O_2 distribution in *Chlamydomonas reinhardtii*. Plant Cell **33**:
970   2935–2949.

971 **Niittylä, T., Messerli, G., Trevisan, M., Chen, J., Smith, A.M., and Zeeman, S.C.** (2004). A
972   previously unknown maltose transporter essential for starch degradation in leaves. Science
973   **303**: 87–89.

974 **Niyogi, K.K. and Truong, T.B.** (2013). Evolution of flexible non-photochemical quenching
975   mechanisms that regulate light harvesting in oxygenic photosynthesis. Curr. Opin. Plant
976   Biol. **16**: 307–314.

Prabhakar, V. et al. (2010). Phosphoenolpyruvate provision to plastids is essential for
gametophyte and sporophyte development in Arabidopsis thaliana. Plant Cell **22**: 2594–
2617.

Raghavendra, A.S. and Padmasree, K. (2003). Beneficial interactions of mitochondrial
metabolism with photosynthetic carbon assimilation. Trends Plant Sci. **8**: 546–553.

Riesmeier, J.W., Flügge, U.I., Schulz, B., Heineke, D., Heldt, H.W., Willmitzer, L., and
Frommer, W.B. (1993). Antisense repression of the chloroplast triose phosphate
translocator affects carbon partitioning in transgenic potato plants. Proc. Natl. Acad. Sci.
U. S. A. **90**: 6160–6164.

Rochaix, J.-D., Goldschmidt-Clermont, M., and Merchant, S. (1998). The molecular biology
of chloroplasts and mitochondria in Chlamydomonas J.-D. Rochaix, M. Goldschmidt-
Clermont, and S. Merchant, eds (Plenum Publishing Corporation).

Sake, C.L., Newman, D.M., and Boyle, N.R. (2020). Evaluation of quenching methods for
metabolite recovery in photoautotrophic *Synechococcus sp*. PCC 7002. Biotechnol. Prog.
36: e3015.

Saroussi, S., Sanz-Luque, E., Kim, R.G., and Grossman, A.R. (2017). Nutrient scavenging and
energy management: acclimation responses in nitrogen and sulfur deprived
Chlamydomonas. Curr. Opin. Plant Biol. **39**: 114–122.

Schneider, A., Häusler, R.E., Kolukisaoglu, Ü., Kunze, R., Van der Graaff, E., Schwacke, R.,
Catoni, E., Desimone, M., and Flügge, U.I. (2002). An Arabidopsis thaliana knock-out
mutant of the chloroplast triose phosphate/phosphate translocator is severely compromised
only when starch synthesis, but not starch mobilisation is abolished. Plant Journal.

Shapiguzov, A., Vainonen, J.P., Wrzaczek, M., and Kangasjärvi, J. (2012). ROS-talk - how
the apoplast, the chloroplast, and the nucleus get the message through. Front. Plant Sci. **3**:
292.

Smirnoff, N. and Arnaud, D. (2019). Hydrogen peroxide metabolism and functions in plants.
New Phytol. **221**: 1197–1214.

Stocking, C.R. and Larson, S. (1969). A chloroplast cytoplasmic shuttle and the reduction of
extraplastid NAD. Biochem. Biophys. Res. Commun. **37**: 278–282.

Streatfield, S.J., Weber, A., Kinsman, E.A., Hausler, R.E., Li, J., Post-Beittenmiller, D.,
Kaiser, W.M., Pyke, K.A., Flugge, U.-I., and Chory, J. (1999). The
phosphoenolpyruvate/phosphate translocator is required for phenolic metabolism, palisade
cell development, and Plastid-dependent nuclear gene expression. Plant Cell **11**: 1609.

Strenkert, D., Schmollinger, S., Gallaher, S.D., Salomé, P.A., Purvine, S.O., Nicora, C.D.,
Mettler-Altmann, T., Soubeyrand, E., Weber, A.P.M., Lipton, M.S., Basset, G.J., and
Merchant, S.S. (2019). Multiomics resolution of molecular events during a day in the life
of Chlamydomonas. Proc. Natl. Acad. Sci. U. S. A. **116**: 2374–2383.

Suzuki, N., Koussevitzky, S., Mittler, R., and Miller, G. (2012). ROS and redox signalling in
the response of plants to abiotic stress. Plant Cell Environ. **35**: 259–270.

Treves, H., Küken, A., Arrivault, S., Ishihara, H., Hoppe, I., Erban, A., Höhne, M., Moraes,
T.A., Kopka, J., Szymanski, J., Nikoloski, Z., and Stitt, M. (2022). Carbon flux through
photosynthesis and central carbon metabolism show distinct patterns between algae, C3
and C4 plants. Nat Plants **8**: 78–91.

Urzica, E.I., Vieler, A., Hong-Hermesdorf, A., Page, M.D., Casero, D., Gallaher, S.D., Kropat,
J., Pellegrini, M., Benning, C., and Merchant, S.S. (2013). Remodeling of membrane
lipids in iron-starved Chlamydomonas. J. Biol. Chem. **288**: 30246–30258.

Vevea, J.D., Alessi Wolken, D.M., Swayne, T.C., White, A.B., and Pon, L.A. (2013).
Ratiometric biosensors that measure mitochondrial redox state and ATP in living yeast
cells. J. Vis. Exp.: 50633.

Wakao, S. and Niyogi, K.K. (2021). Chlamydomonas as a model for reactive oxygen species
signaling and thiol redox regulation in the green lineage. Plant Physiol. **187**: 687–698.

Walters, R.G., Ibrahim, D.G., Horton, P., and Kruger, N.J. (2004). A mutant of Arabidopsis
lacking the triose-phosphate/phosphate translocator reveals metabolic regulation of starch
breakdown in the light. Plant Physiol. **135**: 891–906.

Weber, A.P.M., Linka, M., and Bhattacharya, D. (2006). Single, ancient origin of a plastid
metabolite translocator family in Plantae from an endomembrane-derived ancestor.
Eukaryot. Cell **5**: 609–612.

Weise, S.E., Liu, T., Childs, K.L., Preiser, A.L., Katulski, H.M., Perrin-Porzondek, C., and
Sharkey, T.D. (2019). Transcriptional regulation of the glucose-6-phosphate/phosphate
translocator 2 is related to carbon exchange across the chloroplast envelope. Front. Plant
Sci. **10**: 827.

Winder, C.L., Dunn, W.B., Schuler, S., Broadhurst, D., Jarvis, R., Stephens, G.M., and
Goodacre, R. (2008). Global metabolic profiling of Escherichia coli cultures: an
evaluation of methods for quenching and extraction of intracellular metabolites. Anal.
Chem. **80**: 2939–2948.

Young, J.D., Shastri, A.A., Stephanopoulos, G., and Morgan, J.A. (2011). Mapping
photoautotrophic metabolism with isotopically nonstationary [13]C flux analysis. Metab.
Eng. **13**: 656–665.

Yu, E.T., Zendejas, F.J., Lane, P.D., Gaucher, S., Simmons, B.A., and Lane, T.W. (2009).
Triacylglycerol accumulation and profiling in the model diatoms *Thalassiosira*
pseudonana and *Phaeodactylum tricornutum* (Baccilariophyceae) during starvation. J.
Appl. Phycol. **21**: 669–681.

Zhao, Y. et al. (2018). Malate transported from chloroplast to mitochondrion triggers production
of ROS and PCD in *Arabidopsis thaliana*. Cell Res. **28**: 448–461.

Zhao, Y., Yu, H., Zhou, J.-M., Smith, S.M., and Li, J. (2020). Malate circulation: linking
chloroplast metabolism to mitochondrial ROS. Trends Plant Sci. **25**: 446–454.

Zirngibl, M.-E., Araguirang, G.E., Kitashova, A., Jahnke, K., Rolka, T., Kühn, C., Nägele,
T., and Richter, A.S. (2022). Triose phosphate export from chloroplasts and cellular sugar
content regulate anthocyanin biosynthesis during high light acclimation. Plant Commun:
100423.

Zones, J.M., Blaby, I.K., Merchant, S.S., and Umen, J.G. (2015). High-Resolution Profiling of
a Synchronized Diurnal Transcriptome from *Chlamydomonas reinhardtii* Reveals
Continuous Cell and Metabolic Differentiation. Plant Cell **27**: 2743–2769.

1060

Figures

Fig. 1: Subcellular localization of CreTPT2 and 3 and in vitro measurements of transport activities. (**A**) Cellular localization of CreTPT2 (upper) and CreTPT3 (lower). The construct for expression of the CreTPT2-3-VENUS fusion protein is shown above the cell images and described in the **Materials and methods**. VENUS fluorescence, green; Chlorophyll (Chl) autofluorescence, magenta. (Scale bar: 5 μm). (**B**) Expression of Cre*TPT2* (1, 2) and Cre*TPT3* (3, 4) in yeast. **Left in each panel**, SDS-PAGE of proteins from total membranes isolated from yeast cells containing the empty vector (1, 3) or expressing His-CreTPT2 (2) or His-CreTPT3 (4). **Right in each panel**, corresponding immunoblot blot detection of recombinant His-CreTPT2 or His-CreTPT3 protein using anti-His antibodies. The calculated molecular masses of the N-terminal His-tagged proteins of CreTPT2 and CreTPT3 were 39, and 37 kDa. (**C**) **Left**, kinetics of Pi exchange by His-CreTPT2 in lipid vesicles. Uptake of Pi (0.25 mM) into liposomes was measured in the presence (■) or absence (□) of Pi (30 mM) loaded into the vesicles. **Right**, analyses of substrate specificity of His-CreTPT2. (**D**) **Left**, kinetics of Pi exchange by His-CreTPT3 in lipid vesicles. Uptake of Pi (0.25 mM) into liposomes was measured in the presence (■) or absence (□) of Pi (30 mM) loaded into the vesicles. **Right**, analyses of substrate specificity of His-CreTPT3. For the left panels of **C** and **D**, the arithmetic mean (±SD) of three independent experiments (each with three technical replicates) was plotted with respect to the time after initiating the assay. For the right panels of **C** and **D**, liposomes in which His-CreTPT2 and His-CreTPT3, respectively, were incorporated, were preloaded with various substrates (30 mM) and the initial Pi uptake rates (at 0.25 mM external concentration) were determined. Relative velocities were compared to the Pi/Pi homo-exchange, which was set to 100%. The data represents the arithmetic mean (±SD) of three independent experiments (each consisting of three technical replicates). (**E**) Comparison of kinetic constants of His-CreTPT2 and His-CreTPT3. The Michaelis-Menten constant (K_M) for Pi uptake was determined using various external Pi concentrations (0.05-5 mM). The competitive inhibition constant (K_i) of Pi uptake (0.25 mM) was measured with increasing competitor concentrations (0.05-5 mM). Liposomes were preloaded with 30 mM Pi as the counter-exchange substrate. The data represent the arithmetic mean +/- SE of three independent experiments. Nd, no competitive inhibitory constant could be measured under the given experimental conditions. DHAP, dihydroxyacetone phosphate; 3-PGA, 3-phosphoglycerate; PEP, phosphoenolpyruvate; Glc6P, glucose 6-P; Glc1P, glucose 1-P; Fru6P, fructose 6-P; Gal1P, galactose 1-P.

Fig. 2: Growth characterization of *tpt* mutants and complemented strains. (**A**) Molecular map of the Cre*TPT2* (upper) and Cre*TPT3* (lower) genes and the positions of the Cas9 targeted sites. Edited site of Cre*TPT2* is located in exon 8, edited sites of Cre*TPT3* are located in exons 1 and 7; *t3ko1* has the marker gene inserted in exon 1 and *t3ko2* and *t3ko3* have the marker gene inserted in exon 7. (**B**) Immunoblot (left panel) of the stained protein profiles (right panel) for wild type (WT), *t3ko1*, *t3ko2*, *t3ko3* and the complemented strains (*C_T3KO1*, *C_T3KO2* and *C_T3KO3*, respectively). (**C**) Growth of WT, *t2ko1*, *t2ko2*, and *t3ko2* at various dilutions (see below) on agar plates incubated under LL and HL for 4 d. (**D**) Growth of WT, *t3ko1*, *t3ko2*, *t3ko3* and the complemented strains (*C_T3KO1*, *C_T3KO2*, and *C_T3KO3*, respectively), at various dilutions (see below) on agar plates incubated under LL and ML for 4 d. (**E**) Growth of the various strains (indicated) in liquid medium under LL (left panel) and ML (right panel). (**F**) Transition of the strains from LL to ML (upper panel) or to HL (lower panel) for the times indicated. (**G**) Cell growth (number) and chlorophyll content of cultures at various times after shifting from LL to ML for up to 48 h. In **C** and **D**, the cells were spotted on agar plates containing TAP or TP medium and maintained under continuous LL, ML or HL conditions. The dilution series used was 1.5, 0.75, 0.375 µg/mL chlorophyll (left to right). For growth in liquid medium, the cells were cultured in TP medium to an initial OD$_{750nm}$ 0.02 in air and under continuous LL before initiating the various growth analyses in LL, ML or HL. Each curve represents the arithmetic mean ($\pm$SD) of three independent experiments.

1063

Fig. 3: Changes in storage carbon and metabolite levels in WT and *t3ko2* after transitioning cells from growth in LL to ML. (**A**) Starch content in indicated strains following 0, 6, and 24 h of ML exposure. (**B**) Lugol staining of starch in WT and *t3ko2* after 48 h of ML. (**C**) TAG (TAG: triacylglycerol) content in the indicated strains following 0, 6, and 24 h of ML exposure. (**D**) Nile Red staining of TAG in WT and *t3ko2* following 48 h of ML exposure. (**E**) Time course of metabolite accumulation at 0 h (LL), 45 min and 6 h following ML exposure. Data was normalized to cell numbers. Each data point shows the mean and standard error; the data represents three biological replicates for each metabolite. ppb: parts per billion. (**F**) Select metabolic pathways in Chlamydomonas adapted from (Johnson and Alric, 2013). (**G**) Bar graph representation of fold-change for metabolites shown in (**E**); calculated by dividing the mean of pool size in *t3ko2* by that of WT at the respective light levels. A fold change of 1 (no change) is indicated with a dashed line. An asterisk indicates statistically significant difference in pool sizes in *t3ko2* compared to WT (from **E**) (* P<0.05, ** P<0.01, *** P<0.001). Abbreviations: G6P, glucose-6-P; FBP, fructose bisphosphate; DHAP, dihydroxyacetone phosphate; 3-PGA, 3-phosphoglycerate; PEP, phosphoenolpyruvate; PYR, pyruvate; CIT, citrate; SUCC, succinate; FUM, fumarate; MAL, malate; E4P, Erythrose 4-P; R5P, Ribose 5-P; RU5P/X5P, ribulose 5-P/xylulose-5-P; OMT, 2-oxoglutarate/malate transporter.

Fig. 4: Photosynthetic activities following a transition from LL to ML or to HL. (A) Light intensity curve of O_2 evolution of LL (left) and ML (right)-acclimated cells for the indicated strains. **(B)** Fv/Fm of WT, *t3ko2* and the rescued strain (*C_T3KO2*) following 0 (LL), 6 and 24 h in ML. **(C)** 1-qL values of LL (left panel) and ML (right panel) acclimated cells exposed to increasing actinic light intensities. **(D)** Kinetics of 1-qL of LL-acclimated cells after illumination at HL (400-450 μmol photons $m^{-2}s^{-1}$) for the times indicated on the x-axis. **(E)** P700 oxidation and reduction kinetics in LL-grown cells and after 6 and 24 h in ML. **(F)** Kinetics of P700 oxidation upon dark-to-light transition. P700 measurements were performed in the presence of DCMU and hydroxylamine, as indicated in the **Materials and methods**. Absorbance differences were monitored at 705 nm during continuous illumination with 150 μmol photons·$m^{-2} s^{-1}$ for 5 sec (white box above), followed by a saturating light pulse at 1,500 μmol photons·$m^{-2} s^{-1}$ (arrow) and a 5 sec dark incubation (black box). For panel **E**, the kinetics was normalized by setting maximum oxidation (after light pulse) of WT to 100% and for panel **F**, by setting maximum oxidation (after light pulse) of individual strain to 100%. The data from all panels represent the arithmetic mean ($\pm$SD) of three independent experiments. For panel **B**, the asterisks represent significant differences determined by ANOVA tests., ****P < 0.0001.

1065

Fig. 5: Measurement of intracellular ROS in *t3ko2* and WT upon transition from LL to ML. (**A**) ROS levels were evaluated by CM-H2DCFDA fluorescence of WT and the *t3ko2* mutant in LL and following exposure to ML for 48 h. Scale bar: 20 μm. (**B**) Quantitation of data in **A**. Shown are mean values from three independent experiments, error bars represent standard deviation. roGFP2 protein was targeted to the chloroplast stroma (**C**, upper panel) and mitochondrion (**C**, lower panel). Scale bar: 10 μm; BF: bright field; roGFP2 fluorescence: green; Chlorophyll (Chl) autofluorescence: red. (**D-E**) Monitoring redox levels in stroma (**D**) and mitochondrion (**E**) after exposure of WT and the *t3ko2* mutant for 1.5 and 6.0 h of ML. Shown are mean values from three independent experiments, 3-4 random figures were analyzed for each experiment, error bars represent standard deviation. For panel **B** and **D**, the asterisks represent significant differences determined by ANOVA tests., **P < 0.005, ***P < 0.001.

1066

Fig. 6: Real-time monitoring of H₂O₂ compartmentalized distribution in *t3ko2* and WT upon transition from LL to HL. (**A**) The hypersensitive H₂O₂ sensor (roGFP2-Tsa2ΔCR) protein was targeted to the chloroplast stroma (St), cytosol (Cyto), mitochondrion (Mito) and nucleus (Nu). Shown are GFP fluorescence, chlorophyll autofluorescence (Chl) and the two signals merged (scale bar: 5 µm). WT and *t3ko2* transformant cells, accumulating roGFP2-Tsa2ΔCR in the stroma (**B**), cytosol (**C**), mitochondrial matrix (**D**), and nucleus (**E**), were acclimated to LL in TP and then were transferred either to HL (450 µmol photons m⁻² s⁻¹, solid line) or to very LL (10 µmol photons m⁻² s⁻¹, dotted line) for 20 min. The oxidation state of the sensor was trapped by the addition of NEM and roGFP2 fluorescence was measured in a plate reader as previously described (Niemeyer et al., 2021). Shown are the mean values from three independent experiments with the error bars representing the standard deviation.

1067

Fig. 7: Impact of eliminating Chlamydomonas TPT3 on the chloroplast metabolic landscape. i. reduced triose-P export leads to accumulation of the sugar-P in CBBC and glycolysis pathways (e.g. triose-P, hexose-P, marked in orange), and an elevated ratio of 3-PGA/Pi (which can allosterically activate AGPase activity) would begin to stimulate the synthesis and accumulation of storage carbon; **ii.** CBBC activity diminishes as a consequence of the hyper-accumulation of the precursors with the fixation of CO_2, leading to elevation of reductant (NADPH) and energy (ATP) within the chloroplast; the elevated NADPH:NADP$^+$ and ATP:ADP ratios, elicit strong feedback on PET causing hyper-reduction of electron carriers (eg: over reduced PQ pool and highly reduced PSI) that slows electron flow across the cytochrome b_6f complex; **iii.** the highly reduced PET system and the elevated NADPH can actively reduce FDX/TRX and the NTRC systems, respectively; reduced TRX and/or NTRC can activate the AGPase (above the activity elicited by an increase in the 3-PGA/Pi ratio), leading to additional starch accumulation; **iv.** hyper-reduction of PET can also lead to stromal ROS accumulation, that can act as a signal that controls expression of chloroplast and nuclear genes; ROS also cause damage to both PSII and I and leads to neutral lipid accumulation as a consequence of activation of diacylglycerol acyltransferases (DGAT) and phospholipid diacylglycerol acyltransferase (PDAT). Many of these metabolic/acclimatory processes both overlap and are interconnected. Under conditions of extended exposure to ML, the damage in Chlamydomonas *tpt3* mutants can accumulate and lead to cell death; cell death occurs rapidly in HL.

1068

Fig. 8: Effects of deletion of CreTPT3 over diurnal cycle. (**A**) Growth of WT, and *t3ko2* strains on TP agar plates for 4 days either under diurnal cycle (DC) (left panel; light : dark/ 12h : 12h) or continuous light (CL) (right panel) at an intensity of 60 µmol photon m⁻²s⁻¹ (upper panel) or HL (lower panel: 450 µmol photon m⁻²s⁻¹). The dilution series are 3, 1.5, 0.75, 0.375 µg/mL chlorophyll. (**B**) Photosynthetic electron transport rate (ETR) of cells grown under a diurnal rythym with the light period at an intensity of 60 µmol photon m⁻²s⁻¹. (**C**) Accumulation of starch during growth under a diurnal rythym at a light intensity of 60 µmol photon m⁻²s⁻¹ .

1069
1070

Supplementary Materials

Materials and methods

Expression of CreTPT2 and CreTPT3 in yeast

The DNA sequence encoding the mature CreTPT2 or CreTPT33 protein was codon-optimized for expression in *Saccharomyces cerevisiae* (GeneART, ThermoFisher Scientific). This coding sequence was inserted in frame with an N-terminal His tag into the yeast vector pYES-NTa (ThermoFisher Scientific) using Gibson cloning (NEB). Briefly, pYES-NTa was linearized with BamHI and the Cre*TPT2* and *TPT3* cDNA amplified with the primer pairs: fwd:5'-gacgataaggtacctagcGCTGCTGCTGTTCCAGCTG-3' and rev: 5'- agaattccaccacactgTTAAGCAGCTTCTGGCTTAACATC-3' (or Cre*TPT2*); fwd: 5' - gacgataaggtacctagcGCTTCTGCTGCTGATGCTC-3' and rev: 5' - agaattccaccacactgTTAAGCGGCAGCTGGAGAAG-3' (for Cre*TPT3*). This cDNA was ligated into the linearized pYES-NTa vector, which was then transformed into the yeast strain INVSc1 (MATa, his3D1, leu2, trp1-289, ura3-52 / MATa, his3D1, leu2, trp1-289, and ura3-52, Thermo Fisher Scientific) using the lithium-acetate/PEG method (Gietz and Schiestl, 2007). Transformed yeast cells were selected on synthetic complete medium containing 2% (w/v) glucose with the uracil auxotrophic marker. Galactose-inducible expression of His-CreTPT2 or CreTPT3 in yeast was performed as described in (Linka et al., 2008). The presence of the His-tagged fusion protein was verified by standard SDS-PAGE and immunoblot analysis using an anti-His-antibody conjugated with horseradish peroxidase (Miltenyi Biotech).

Transport activity assays

The Km for Pi was determined by measuring the initial velocity at each of six external Pi concentrations between 0.05 and 5 mM. To obtain competitive inhibition constants (Ki), the uptake of 0.25 mM Pi into liposomes containing 30 mM Pi was measured over a 4 min period in the presence of increasing external competitor–concentrations (0.05 - 5 mM). Three biological replicates were performed for all described experiments. GraphPad Prism software version 9.3.0 was used for non-linear regression analyses of the kinetic data.

Vector construction

The plasmid of pRam118_VENUS was linearized with HpaI (NEB). Primers: g*TPT2*_pRam118_f, g*TPT2*_pRam118_r and g*TPT3*_pRam118_f, g*TPT3*_pRam118_r (**Supplementary Table 1**) were used to amplify Cre*TPT2* and Cre*TPT3* genomic DNAs containing an overlap with the linearized pRam118_VENUS vector. For generating the plasmids pRam118-CreTPT2&VENUS and pRam118-CreTPT3&VENUS, genomic DNA of Cre*TPT2* or Cre*TPT3* was assembled with pRam118_VENUS plasmid using Gibson assembly (Gibson et al., 2009). The plasmid of pRam118-CreTPT2&VENUS (2-4 µg)

linearized with AseI (NEB), was transferred into the Chlamydomonas M10 strain by
electroporation.

A 1,000 bp sequence upstream of Cre*TPT3* and containing the promoter region of the
gene was amplified using the primers TPT3pro1000_f and TPT3pro1000_r
(**Supplementary Table 2**). The 3' UTR of Cre*TPT3* was amplified using TPT_3UTR_f
and TPT3_3UTR_r (**Supplementary Table 2**). The *PSAD* promoter and the *RBCS2*
3'UTR of pRam118-CreTPT3&VENUS were replaced by the amplified fragments of the
1,000 bp upstream region and the 3' UTR of Cre*TPT3*, respectively. This final vector,
designated pRam118_CreTPT3, contains the original Cre*TPT3* promoter (driving
expression of Cre*TPT3*), the genomic DNA sequence of Cre*TPT3*, VENUS and the
Cre*TPT3* 3' UTR plus the *AphVII* cassette. To locate CreTPT3 and complement the *tpt3*
mutant, the mutant was transformed by electroporation with a total amount of 2-4 μg
(~500 ng/μL) of pRam118_CreTPT3 that was linearized with AseI.

CRISPR-Cas9 mediated mutagenesis

Prior to electroporation, Cas9 (IDT) and sgRNAs were incubated together at 37 °C for
30 min. Approximately 500 ng PCR product of the *AphVII* cassette, which confers
resistance to hygromycin, was added to the RNP (ribonucleoprotein) mixture. 250 μl
aliquots were electroporated using Super Electroporator NEPA21 type II (NEPA GENE).
After 16 h of recovery in very low light (10-15 μmol photons $m^{-2}\,s^{-1}$), cells were plated onto
solid TAP medium containing 10 μg/mL hygromycin. Sense or antisense-oriented knock-
ins of *AphVII* were determined by amplification using primer pairs with one primer
annealing to the genomic sequence and the other to the inserted sequence
(**Supplementary Table 2**). The amplified fragments were sequenced to verify the
insertion sites (ELIM BIOPHARM, Hayward, USA).

ROS measurements and roGFP2 imaging analysis

Briefly, after various treatment (e.g. 48 h in ML), 10 million Chlamydomonas cells were
pelleted and washed once with 1×PBS. The cells were then resuspended in 1×PBS
containing 8 μM CM-H2DCFDA and incubated at room temperature in the dark for 30 min.
Following this incubation, the cells were washed three times with 1×PBS buffer, the
fluorescent signals were either visualized using TCS SP8 confocal laser-scanning
microscope (Leica) or quantified with a microplate reader (Infinite M1000; TECAN).
Excitation and emission settings for the microscope were: 488 nm/510-530 nm HyD SMD
hybrid detector for reactive oxygen species (ROS), and 488 nm/650-700 nm-HyD SMD
hybrid detector for chlorophyll autofluorescence. Excitation and emission settings for the
plate reader were as follows: ROS excitation 488/5 nm and emission 530/12 nm;
chlorophyll excitation 514/5 nm and emission 690/5 nm.

1144

Starch analysis

In brief, cells were collected by centrifugation and pigments extracted with methanol. Dried pellets were resuspended in water and heated at 100°C to break the cells and release the starch. After cooling, amyloglucosidase and α-amylase (2.25 U/mL) were used to hydrolyze the starch and the products of hydrolysis were quantified using a Glucose Colorimetric Detection Kit (Thermo Fisher).

Photosynthesis-Irradiance curve

The YSI 5331A electrodes (Yellow Springs Instruments) were polarized at −0.8 V. The cultures used for the assays were concentrated to 2.5-5 µg mL^{-1} chlorophyll, supplemented with 15 µL of 0.5 M sodium bicarbonate in water and then purged with 1% CO_2/99% He. Using a gas tight syringe, the sample was transferred into the reaction chamber that was also purged with 1% CO_2/99% He. The rate of change in O_2 levels was measured sequentially at the light intensities 100, 200, 600, 1000 µmol photons m^{-2} s^{-1} (photosynthetic active radiation, PAR); each intensity was maintained for 3-5 min followed by a 3-min intervening dark period, and then the light level was raised to the next higher intensity (stepped change) until the full range of intensities was tested. At 1,000 µmol photons m^{-2} s^{-1}, the light was held for 10 min followed by an 8-min dark period. Before the measurement of each experimental series, the electrodes were calibrated with air (~21% O_2) and 1% CO_2/99% He mixture (0% O_2). The initial slope of the response was used to determine the O_2 evolution rate.

Extraction of samples for LC-MS/MS

The quenching solution, filtered saline (9 g/L NaCl), was prechilled to 4°C in a refrigerator. 30 mL of the quenching solution was transferred to a 50 mL conical tube kept in an ice bath mixed with salt to depress the temperature to between -3 °C and -1 °C. 15 mL of the culture was rapidly plunged into the 30 mL of the quenching solution and the samples were centrifuged at 4,000 rpm for 10 min at 4 °C. Cell pellets were washed with fresh prechilled saline solution and centrifuged again at 4,000 rpm for 10 min in a 2 mL centrifuge tube. For each replicate, 45 mL of the culture was sampled and the cell pellets were pooled. Washed pellets were frozen at -80 °C until extraction. Metabolite extraction and further analysis were modified from (Young et al., 2011). To each cell pellet, 500 µL methanol was added along with the internal standards ribitol and PIPES to a final concentration of 2 µM each. Samples were then vortexed for 30 sec, frozen in liquid nitrogen as described in (Winder et al., 2008), and allowed to thaw at 0 °C. This vortex-freeze-thaw cycle was repeated twice more before the samples were centrifuged at 10,000 x g at 1 °C for 5 min. The supernatant was collected, and the remaining pellets were extracted twice more with 500 µL of a 50:50 mixture of methanol and water, with 3 vortex-freeze-thaw cycles done for each extraction. Supernatants from each extraction procedure were pooled and dried on a Thermo Fisher SpeedVac Concentrator. Dried

extracts were then resuspended in 500 µL water and cleaned to remove any residual
large cell debris by filtration, first through a 0.22 µm pore size Spin-X centrifugal tube filter,
followed by 10 K molecular mass cutoff filters, followed finally by 3 K molecular mass
cutoff filters. After each filtration step, filters were rinsed with 50 µL water, which was
added to the total sample volume for subsequent steps. Filtered extracts were dried again
and resuspended in 200 µL water for LC-MS/MS analysis.

Protein extraction and immunoblot analysis

Affinity-purified polyclonal antibodies to CreTPT3 were custom-made by GenScript. The
antigen sequence used for antibody generation was KSWSFGRPVTKQEF.
Chlamydomonas cells were grown in liquid cultures to $2–5 \times 10^6$ cells ml^{-1} and collected
by centrifugation (1459 x g, 5 min). Cells were resuspended in resuspension buffer [(5
mM HEPES, pH 7.5, KOH), 10 mM EDTA (pH 7.5, NaOH), 1x protease inhibitor "complete
EDTA-free" (Roche)]. For the disruption of cells, bead-beating was performed using a
mini beadbeater (Biospec) in two cycles of 30 sec min^{-1}, with a 1 min period of cooling on
ice between cycles. Disrupted cells were centrifuged at 4 °C, 30 min, 14000 x g in a
microfuge. The supernatant was removed, the pellet was resuspended in sample buffer
(resuspension buffer supplemented with 100 mM Na_2CO_3, 100 mM DTT, 2% WT/vol SDS,
and 12% WT /vol sucrose).

SDS-PAGE was performed using a 12% polyacrylamide gel (Bio-Rad), with the
electrophoresis for 90 min at 120 V. The proteins within the gel were transferred onto a
PVDF membrane (Bio-Rad) using Trans-Blot Turbo (Bio-Rad). The detection was
performed using Clarity Max Western ECL Substrate (Bio-Rad).

RT-qPCR

RT-qPCR, was performed as described previously (Kaye et al., 2019). Briefly, total RNA
was isolated using the RNeasy Plant Mini Kit (Qiagen) and treated with DNase I (Qiagen).
First-strand cDNA was generated by reverse transcription of 0.5 µg total RNA using the
iScript cDNA Synthesis Kit (Bio-Rad). Real-time PCR using the Roche Light Cycler 480
was performed with the SensiMix NO-Rox SYBR Green I Kit as described by the
manufacturer (Bioline). Oligos used in this research for analyzing the expression levels
of Cre*TPT10/TPT2/TPT3/CGL51* are listed in **Supplemental Table 1**. *mRNA levels were*
calculated using DeltaC (T) method, CBLP gene was used as an internal control.

Analysis of transcript levels in response to different conditions

In order to analyze the transcript level of Cre*TPT* genes under different conditions, we
obtained RNAseq data published previously in which cells experienced a diurnal cycle,
nitrogen starvation, sulfur starvation and Fe^{2+} starvation (Strenkert et al., 2019; Zones et
al., 2015; Ngan et al., 2015; González-Ballester et al., 2010; Urzica et al., 2013). The log2

fold change of each gene was calculated based on the ratio of transcript abundance at
different time point over transcript level at time point 0 h. The fold change was visualized
by a heatmap and the color bar is the scale of log2 (fold-change).

Correlation analysis based on Cre*TPT3* transcription abundance under different
conditions

The Z score of each gene was calculated as follows: the mean transcript level of each
gene from all of the conditions was subtracted from each gene's transcript level. This
difference was divided by the standard deviation of each gene's transcript level under all
conditions to get Z score numeric values. The whole Z score matrix for every gene under
all conditions was used to calculate the Pearson correlation matrix by using python.
Dataframe.corr() function. The genes highly correlated with TPT3 were selected if their
correlation value is higher than 0.75. The Pearson correlation value of TPT3-highly-
correlated genes were plotted based on their KEGG pathway.

In vivo polymeric carbohydrate staining

Lugol's iodine (aqueous solution of 1.8% iodine and 3.0% potassium iodide) was used to
stain in vivo polymeric carbohydrate. Briefly, 0.5 mL of the cell culture was pelleted,
resuspended in 10 μL of Lugol's iodine and visualized under a bright-field microscope.
Lugol's iodine selectively binds to alpha-1,4 glucans and stains it blue-black.

1238

1239

1240

1241

1242

1243

1244

1245

1246

1247

1248

Supplementary Tables

Cre ID	Name from V6.1 / (name used in this study)	Previous name	Prediction	CS Position
Cre08.g379350_4532	TPT10 / (TPT10)	TPT1	cTP	CS pos: 61-62. TCL-AV. Pr: 0.3994
Cre06.g263850_4532	N / (TPT2)	PPT2; TPT2; TPT20	cTP	CS pos: 70-71. VCQ-AA. Pr: 0.5253
Cre01.g045550_4532	TPT3 / (TPT3)	TPT3	cTP	CS pos: 60-61. VTK-AS. Pr: 0.4178
Cre02.g106200_4532	N	TPT4; TPT5	OTHER	
Cre02.g112900_4532	N	TPT5; TPT6	OTHER	
Cre02.g144300_4532	N	TPT6; TPT7	OTHER	
Cre03.g162000_4532	N	TPT7; TPT8	OTHER	
Cre03.g184850_4532	N	TPT8; TPT9	OTHER	
Cre04.g227450_4532	N	TPT9	OTHER	
Cre07.g330850_4532	N	TPT10; TPT11	OTHER	
Cre08.g363600_4532	N	TPT11; TPT12	OTHER	
Cre09.g408400_4532	N	TPT12; TPT13	OTHER	
Cre09.g413700_4532	N	TPT13; TPT14	OTHER	
Cre08.g382350_4532	EZY14	TPT14; TPT16; EZY14	OTHER	
Cre09.g415900_4532	MOT20	TPT15	OTHER	
Cre10.g452750_4532	N	TPT16; TPT18	OTHER	
Cre11.g479950_4532	CGL7	TPT17	OTHER	
Cre12.g490050_4532	N	TPT18; TPT19; GMT1	OTHER	
Cre12.g490100_4532	TPT2	TPT19;GMT2	OTHER	
Cre12.g501000_4532	N	TPT20; TPT22; PPT1	OTHER	
Cre15.g641266_4532	N	TPT22; TPT23	OTHER	
Cre15.g642950_4532	N	TPT23; TPT24	OTHER	
Cre15.g643385_4532	N	TPT24; TPT26	OTHER	
Cre16.g663800_4532	CGL51 / (CGL51)	TPT25	cTP	CS pos: 70-71. IVA-SS. Pr: 0.4817
Cre16.g666250_4532	N	TPT26; TPT27	OTHER	
Cre17.g702700_4532	N	TPT27; TPT28	OTHER	
Cre17.g703250_4532	SLC35D	TPT28; TPT29	OTHER	
Cre17.g710850_4532	N	TPT29; TPT4	OTHER	
Cre11.g467754_4532	N	UAA6	OTHER	
Cre14.g622700_4532	N	N	OTHER	
Cre18.g748947_4532	N	N	OTHER	
Cre09.g408428_4532	N	N	OTHER	

Supplementary Table 1: Members of the triose phosphate transporter family in *Chlamydomonas reinhardtii* and their predicted subcellular localization based on TargetP-2.0. cTP, chloroplast transit peptide; CS position, cleavage site of transit peptides; N: genes without names in the genome. Cre IDs are from v6.1 genome.

oligo name	sequence
gTPT2_pRam118_f	tactcacaacaagcccagttatggcgtctctgctgggc
gTPT2_pRam118_r	gagccacccagatctccgttggcggcctccggcttgac
gTPT3_pRam118_f	tactcacaacaagcccagttATGGCTTCGCTCGCTCAGCGCGTGAGC
gTPT3_pRam118_r	gagccacccagatctccgttGGCCGCGGCCGGCGACGC
TPT3pro1000_f	gccttttgctcacatgtggcggccgccgccatgctggccaacag
TPT3pro1000_r	taccaatccacaaatgtgggctagcccatgtgaaggcgttgacc
TPT3_3UTR_f	gatgacgacaagtaaagatctTAAGGCGGGCAGCGGC
TPT3_3UTR_r	GCGCAAGAAAGAAGCTTGATATCCCCCTTATAATGTTTACAAGTGGTGCG
hygro_crispr_f2	GGAACTGGCGCAGTTCCTCTG
hygro_crispr_re2	GGCTCGCGTAGGAATCATCCG
TPT2_gRNA1_F	TGGGACGGGAGGTGGAGAGC
TPT3_gRNA1_F	GCTCGGTTGAGCGCACCC
TPT3_gRNA1_R	CAAGGGTCTGGCGAGCACG
TPT3_gRNA2_F	ACCTCCCCATCGCCACGAT
TPT3_gRNA2_R	AGGGGTAAGGACAGCGGTG
Qrt_TPT10_f	CCTGTACCAGCAGGTGTCGTAC
Qrt_TPT10_r	CCAGCGCGTTCTGGAGTGACA
Qrt_TPT2_f	ATGGCGTCTCTGCTGGGC
Qrt_TPT2_r	CTGCTCGCCCATTCTCCTC
Qrt_TPT3_f	GCAAAGCTGGCGCCTACTG
Qrt_TPT3_r	TAGACCAGGCCGACCACCAC
Qrt_CGL51_f	CTGGTTCATCGGGACATGGC
Qrt_CGL51_r	TGGCTGAAGCTGACCACT
Qrt_MDH1_f	GGCTCTGCTGCTGAAGATGAAC
Qrt_MDH1_r	CGGTGTAGCCAGTAACCTTCAC
Qrt_MDH5_f	GGCCCTGAACATGAAGCAGC
Qrt_MDH5_r	GGCGGAAGACTCCATAGGCC

1255

Supplementary Table 2: Oligonucleotide list.

1256

ID	Construct	Restriction enzyme for transformation	Localization
pMBS472	PSADpro::aadA::PSAD-T:::: ARpro::roGFP2-Tsa2ΔCR::MStop::RPL23-T	ECoRV/Afel	cytosol
pMBS652	PSADpro::aadA::PSAD-T:::: ARpro::CDJ1::roGFP2-Tsa2ΔCR::MStop::RPL23-T	ECoRV/Afel	stroma
pMBS657	PSADpro::aadA::PSAD-T:::: ARpro::roGFP2-Tsa2ΔCR::SV40::RPL23-T	ECoRV/Afel	nucleus
pMBS655	PSADpro::aadA::PSAD-T:::: ARpro::HSP70C::roGFP2-Tsa2ΔCR::MStop::RPL2	ECoRV	mitochondrion

1257

Supplementary Table 3: Plasmids of H₂O₂ roGFP2-Tsa2ΔCR sensors (Niemeyer et al., 2021)**.**

1258

Name	Cre number	Gene function	Correlation coefficient	Localization
TPT3	Cre01.g045550	Triose phosphate/phosphate translocator 2	1	chloroplast
AMY	Cre08.g362450	Alpha-amylase	0.881792677	chloroplast
SP	Cre07.g336950	Starch phosphorylase	0.861504451	chloroplast
ISA3	Cre03.g207713	Isoamylase, starch debranching enzyme	0.790501386	chloroplast
PGI1	Cre03.g175400	Phosphoglucose isomerase	0.808137223	chloroplast
PFK2	Cre12.g553250	Phosphofructokinase	0.805294876	chloroplast
PGM1	Cre06.g272050	Phosphoglycerate mutase	0.838232423	other
IDH2	Cre02.g143250	Isocitrate dehydrogenase, NAD-dependent	0.81622058	other
MDH4	Cre12.g483950	Malate dehydrogenase 4	0.80363604	mitochondrion
PDH1	Cre16.g677026	Pyruvate dehydrogenase E1 beta subunit	0.805186998	mitochondrion
ASAP3	Cre07.g338050	Mitochondrial F1F0 ATP synthase associated protein 3	0.853067434	mitochondrion
MPPB	Cre12.g523850	Mitochondrial processing peptidase beta subunit and Complex III Core I subunit	0.844453019	mitochondrion
NUO5	Cre10.g450400	NADH dehydrogenase (ubiquinone) flavoprotein 2,mitochondrial(respiratory Complex I)	0.781322987	mitochondrion
NUO6	Cre10.g422600	NADH:ubiquinone oxidoreductase 51 kDa subunit,mitochondrial, respiratory Complex I	0.770479958	mitochondrion
COX2a	Cre03.g154350	Mitochondrial cytochrome c oxidase subunit II	0.824639841	mitochondrion
COX2	Cre01.g049500	Mitochondrial cytochrome c oxidase subunit II, protein IIb of split subunit	0.814345856	mitochondrion
COX3	Cre04.g221700	Mitochondrial cytochrome c oxidase subunit III	0.813942712	mitochondrion
COX90	Cre16.g691850	Cytochrome c oxidase subunit Cox90, mitochondrial	0.803959451	mitochondrion
COX10	Cre12.g516350	Mitochondrial cytochrome c oxidase assembly protein	0.770332065	mitochondrion
ATP1a	Cre02.g116750	Mitochondrial F1F0 ATP synthase, alpha subunit	0.850554844	mitochondrion
ATP2	Cre17.g698000	Mitochondrial F1F0 ATP synthase, beta subunit	0.839116637	mitochondrion
ATP3	Cre15.g635850	Mitochondrial F1F0 ATP synthase, gamma subunit	0.790390859	mitochondrion
ATP4	Cre11.g467707	Mitochondrial F1F0 ATP synthase, delta subunit	0.782621502	mitochondrion
ASA7	Cre09.g416150	Mitochondrial F1F0 ATP synthase associated protein 7	0.762529504	mitochondrion
ATP5	Cre16.g680000	Mitochondrial ATP synthase subunit 5	0.75995889	mitochondrion
CYC1	Cre15.g638500	Ubiquinol:cytochrome c oxidoreductase cytochrome c1,mitochondrial	0.78087886	mitochondrion
OMT1	Cre17.g713350	Oxoglutarate:malate antiporter1	0.7724107	chloroplast
MDH1	Cre03.g194850	NAD-dependent malate dehydrogenase	0.770420148	chloroplast
ANT1	Cre09.g386650	ADP/ATP carrier protein, mitochondrial	0.841663755	mitochondrion
AAA1	Cre08.g358526	Plastidic ADP/ATP translocase	0.831658774	chloroplast
ATPva3	Cre04.g220350	Vacuolar ATP synthase subunit A	0.806662753	other
VTC1	Cre12.g510250	Vacuolar Transport Chaperone-like protein	0.753558866	other
PST1	Cre07.g331550	Phosphoserine aminotransferase	0.787494263	chloroplast
CPL12	Cre10.g466500	Glyoxalase	0.787085275	chloroplast
NAGSAD	Cre03.g146187	N-acetyl-gamma-glutamyl-phosphate reductase / NAGSA dehydrogenase/oxidoreductase	0.772201917	chloroplast
NAGK1	Cre01.g015000	Acetylglutamate kinase/L-arginine biosynthesis II (acetyl cycle)	0.771816595	chloroplast
PFGS	Cre08.g364800	Phosphoribosylformylglycinamidine synthase /superpathway of purine nucleotides	0.761886415	chloroplast
PPP2r2	Cre01.g055420	serine/threonine-protein phosphatase 2A regulatory subunit B	0.776264136	chloroplast
CLTC	Cre02.g101400	clathrin heavy chain	0.775380317	other
SLC40a1	Cre06.g251000	solute carrier family 40	0.771233669	secretory pathway

1259

1260 **Supplementary Table 4: List of genes with strong Pearson correlation to *CreTPT3*.**

1261
1262

Supplementary Figures

Supplementary Fig. 1: Features of putative plastid phosphate transporter (pPT) in *Arabidopsis thaliana* and *Chlamydomonas reinhardtii*. (**A**) Alignment of sequences of pPTs in Arabidopsis and Chlamydomonas. Plastid putative sugar phosphate transporters from Chlamydomonas are CreTPT10 (Cre08.g379350_4532), CreTPT2 (Cre06.g263850_4532), CreTPT3 (Cre01.g045550_4532), and CGL51 (Cre16.g663800_4532). Sequences from *Arabidopsis thaliana* are AtTPT1 (AT5G46110.1), AtPPT1 (AT5G33320.1), AtGPT1 (AT5G54800.1) and AtXPT1 (AT5G17630.1). Conserved substrate binding pockets are colored red. Phosphate-binding residues are colored blue. The gate capping residues are colored yellow. The positions of residue in green are crucial for the substrate preference of either triose-phosphate or 3-PGA. The purple residues are critical for the specificity of PEP. (**B**) The similarity and identity of the protein sequences between Chlamydomonas pPTs and Arabidopsis TPT1. The similarity and identity were calculated using https://www.bioinformatics.org/sms2/ident_sim.html. (**C**) mRNA levels encoding putative chloroplast envelope localized pPTs under LL during growth in TP medium. (**D**) Expression of pPTs over the diel cycle; data was extracted from Zones et al., 2015. Supports Fig. 1 and 2.

1281

Supplementary Fig. 2: Background activity of Pi uptake with the endogenous yeast transporters reconstituted in into liposomes. Time-dependent uptake of Pi (0.25 mM) into reconstituted liposomes preloaded with 30 mM of Pi (■) or without exchange substrate (●) prepared from yeast cells harboring the empty vector (pYES-NTa). The arithmetic mean (±SD) of three independent experiments (each with three technical replicates) was plotted against time. The observed Pi uptake rates of endogenous yeast carriers reconstituted into liposomes were negligible compared to the rates with His-CreTPT2/3 (20-fold higher, **Fig. 1C, D**), indicating that this expression system is suitable for functional analysis of recombinant phosphate transporters. Supports Fig. 1.

1291

1292

Supplementary Fig. 3: Mutations generated in the Cre*TPT2* gene by CRISPR-Cas9 mediated insertion. (**A**) Presentation of the orientation and position of *AphVII* cassette in the Cre*TPT2* gene in *t2ko1*, *t2ko2*, *t2ko3* and *t2ko4*. Sequencing of genomic DNA fragments across the site of *AphVII* cassette in *t2ko1* (**B**) and *t2ko2, t2ko3* and *t2ko4* (**C**). The mismatched nucleotides are marked in pink. Deletions are shown with dashes. (**D**) Growth of WT, *t2ko1*, *t2ko2*, *t2ko3* and *t2ko4* at various dilutions on agar plates with TP or TAP medium and incubated under LL and HL for 6 d. The dilution series used was 1, 0.5, 0.25, 0.125 µg/mL chlorophyll (left to right). Supports Fig. 2.

1293

1294

1295

1296

1297

1298

1299

1300

1301

1302

Supplementary Fig. 4: Types of mutations generated in the Cre*TPT3* gene by CRISPR-Cas9 mediated insertion with sgRNA1 in exon1 and sgRNA2 in exon7. Sequencing of genomic DNA fragments across the site of insertion of *AphVII* cassette in *t3ko1* (**A**) *t3ko2* (**B**) and *t3ko3* (**C**). The orientation and position of *AphVII* cassette are presented above the alignment. The mismatched nucleotides are marked in pink. Deletions are shown with dashes. Supports Fig. 2.

1310

1311

Supplementary Fig. 5: Experimental set up. All strains were grown photoautotrophically under
low white light (LL, 30 µmol photons m^{-2} s^{-1}) for 16-24 h. After dilution with fresh medium, the
cultures from LL were then either spotted onto solid medium or to liquid medium and maintained
at either LL, moderate light (ML, 250-300 µmol photon m^{-2} s^{-1}) or high light (HL, 450 µmol photon
m^{-2} s^{-1}) for 24 or 48 h. To avoid self-shading, the culture density was adjusted to 1-1.5 µg/mL
chlorophyll before exposure to ML or HL. Supports Fig. 2,3,4,5.

1318

1319

Supplementary Fig. 6: (**A**) Cell size of WT, *t3ko2*, and *C_T3KO2* in TP in both LL and ML. (**B**) WT and *t3ko2* morphology 24 h following a LL to ML transition. Supports Fig. 2.

1322

Supplementary Fig. 7: Quantification of chlorophyll contents in indicated strains after exposure to ML for 24 and 48 h. All strains were grown photoautotrophically in LL for 16-24 h. After dilution with fresh medium, the culture was transferred to ML and the chlorophyll content determined after 24 and 48 h in the ML. The initial inoculum density used for the transition was between 1-1.5 µg/mL chlorophyll in order to avoid self-shading. Supports Fig. 2.

1328

1329

Supplementary Fig. 8: Starch and metabolites in WT and *t3ko2*. (A) Starch content in indicated strains following 0, 6, and 24 h in LL. (**B**) Time course, over a period of 6 h, of changes in metabolite pool sizes measured at 0 h (LL), 45 min and 6 h after shifting LL-grown cells to ML. Data was normalized to chlorophyll contents. (**C**) Schematic representation of the fold-change for the metabolites shown in (**B**), calculated by dividing the pool size in *t3ko2* by that of WT cells under the same conditions. Fold change of 1 (no change) is shown by a dashed line. Each data point is the mean and standard error of three biological replicates. ppb: parts per billion. Asterisk indicates statistically significant differences relative to WT (* P<0.05, ** P<0.01, *** P<0.001). Abbreviations: G6P, glucose-6-phosphate; FBP, fructose bisphosphate; DHAP, dihydroxyacetone phosphate; 3-PGA, 3-phosphoglycerate; PEP, phosphoenolpyruvate; CIT, citrate; SUCC, succinate; FUM, fumarate; MAL, malate; S7P, sedoheptulose-7-phosphate; E4P, Erythrose 4-phosphate; R5P, Ribose 5-phosphate; RU5P/X5P, ribulose 5-phosphate/xylulose-5-phosphate. Supports Fig. 3.

1343

Supplementary Fig. 9: transcript changes of Cre*TPT2* and Cre*TPT3* with various conditions.
(A) Expressions of Cre*TPT2* and Cre*TPT3* following exposure of LL grown cells to HL. Expression
of the putative pPTs under nitrogen starvation **(B)**, sulfur starvation **(C)**, and iron starvation **(D)**
from (Ngan et al., 2015; González-Ballester et al., 2010; Urzica et al., 2013). Supports Fig. 5,6,7.

1344
1345
1346
1347

1348
1349

1350

1351 **Supplementary Fig. 10:** mRNA levels of Cre*TPT10* (**A**), Cre*TPT2* (**B**) and *CGL51* (**C**) in the
1352 indicated strains 0, 2, 6 h after the cells were shifted from LL to ML. (**D, E**) mRNA levels of
1353 plastidial malate dehydrogenases (*MDH*) in the indicated strains 0, 6 h after the cells were shifted
1354 from LL to ML. Supports Fig. 7.

1355

A

B

Supplementary Fig 11: List of genes with strong correlation relative to Cre*TPT3*. (**A**) Correlation coefficients greater than 0.75 were listed. (**B**) Construction of Cre*TPT3* co-expression networks. Supports Fig. 7 and 8.