Title

Triose phosphate translocator drives carbohydrates/energetic shuttling from the chloroplast in

Chlamydomonas reinhardtii

Weichao Huang[1]†*, Anagha Krishnan[2]†, Anastasija Plett[3], Michelle Meagher[4], Nicole Linka[3],

Bijie Ren[1], Justin Findinier[1], Petra Redekop[1], Neda Fakhimi[1], Jin Xu[5], Rick G. Kim[1], Devin A.

Karns[2], Nanette Boyle[4], Matthew C. Posewitz[2], Arthur R. Grossman[1]*

[1]Department of Plant Biology, Carnegie Institution for Science, Stanford, CA 94305, USA

[2]Department of Chemistry, Colorado School of Mines, Golden, CO 80401, USA

[3]Institute of Plant Biochemistry, Heinrich-Heine University Düsseldorf, 40225 Düsseldorf,

Germany

[4]Department of Chemical and Biological Engineering, Colorado School of Mines, Golden, CO

80401, USA

[5]Department of Microbiology and Cell Science, Institute of Food and Agricultural Sciences (IFAS),

University of Florida, Lake Alfred, 33885, FL, USA

†Authors contributed equally.

*Corresponding Author: Weichao Huang; Address: The Carnegie Institution for Science,

260 Panama St. Bldg. 100; Stanford, CA 94305; email: whuang@carnegiescience.edu

*Corresponding Author: Arthur R. Grossman; Address: The Carnegie Institution for Science,

260 Panama St. Bldg. 100; Stanford, CA 94305; email: agrossman@carnegiescience.edu

Abstract

Photosynthetic eukaryotic organisms contain several chloroplast-associated metabolite transporters that enable energetic/metabolic exchange between the chloroplast and other cellular compartments. In this study, we used the model photosynthetic alga *Chlamydomonas reinhardtii* to investigate a highly expressed chloroplast triose phosphate transporter. The chloroplast triose phosphate/phosphate translocator 3 (CreTPT3) showed high DHAP and 3-PGA transport activities in vitro using a yeast liposome uptake system. Null mutants for CreTPT3, generated by CRISPR-Cas9, resulted in a pleiotropic phenotype impacting photosynthetic activity, metabolite pools, carbon partitioning and storage, the redox status of the chloroplast, and the accumulation of reactive oxygen species. The results presented demonstrate that CreTPT3 is the major conduit on the chloroplast envelope for the intracellular distribution of fixed carbon and reductant. Its function is critical for optimizing the use of resources and supporting cell fitness, managing the chloroplast redox status and sustaining high rates of photosynthetic CO_2 fixation.

Introduction

Photosynthetic organisms have evolved numerous strategies to distribute fixed carbon among the different cellular compartments, maintaining a dynamic metabolic/redox balance that fuels growth and biosynthetic processes. However, the extent and characteristics of metabolite and reductant shuttling between the chloroplast and other cellular compartments have not been extensively analyzed despite its importance to cellular metabolism. The chloroplast is delineated by a double envelope membrane that contains several metabolite translocators that are part of the machinery that governs metabolic trafficking among the different cellular compartments[1, 2]. Various mechanisms have been proposed in the shuttling of fixed carbon and reducing equivalents between the chloroplast and cytoplasm. Studies with *Arabidopsis thaliana* revealed that the malate shuttle, which involves multiple malate dehydrogenases (MDH) and malate/OAA translocators (OMT), functions in the export of reductants from the chloroplast and also enables the cells to manage redox conditions in the chloroplast[3, 4].

In addition to the malate shuttle, reduced carbon can be exported from the chloroplast by the triose phosphate (triose-P)/phosphate (Pi) translocator (TPT), which can also serve potentially as a means of exporting carbon from the chloroplast and a safety valve for eliminating excess reducing power from the chloroplast and lessening the potential for ROS accumulation and damage[5, 6]. The TPTs can potentially function as a major path for exchanging metabolites between the chloroplast and cytosol. Triose-Ps are generated during photosynthetic CO_2 fixation by Calvin-Benson-Bassham cycle (CBBC) activity that is supported by the reducing power/energy (NADPH/ATP) derived from photosynthetic electron transport. The TPTs reside on the inner chloroplast envelope membrane and can export triose-Ps, including glyceraldehyde 3-phosphate (G3P/GAP), dihydroxyacetone phosphate (DHAP), and 3-phosphoglycerate (3-PGA) across the chloroplast envelope membrane and into the cytosol in a 1:1 counter exchange for Pi[5–7]. Various plants like tobacco and Arabidopsis, do not exhibit a strong phenotype if TPT is either eliminated or its level is reduced[8–10]. A reduction in TPT activity can be compensated for by moving the assimilated carbon into a transitory starch pool that is subjected to accelerated turnover in the light[8, 9, 11], leading to the production of hexoses that can be exported from the chloroplast. Characterization of a double mutant (*tpt-1/adg1-1*) of Arabidopsis null for TPT (*tpt-1*) and starch synthesis (*adg1-1*) exhibited severely compromised photosynthetic capacity and growth[10, 12].

While there are a few studies that demonstrate the functionalities of TPT proteins in single-celled
photosynthetic organisms (red algae, apicomplexans, cryptophyte) based on the in vitro activity of
these proteins in liposome transport assays(*13–15*), the impact of null mutations in genes encoding
TPT proteins on the physiology of the cells has not been investigated so far. In this study, we used
the unicellular green alga *Chlamydomonas reinhardtii* (Chlamydomonas throughout), a well-
established model organism that has been extensively used to dissect various physiological
processes, including photosynthesis and acclimation to environmental change. Chlamydomonas
contains more than 30 TPT homologs; the informatic analyses of these proteins are still not refined
enough to be certain of their substrate specificities and, in many cases, the transported metabolites
are likely not a triose-P. Here we present work that approaches the questions: *i*) which of the TPTs
is a triose-P/Pi translocator associated with the chloroplast envelope; *ii*) what are the roles of
chloroplast TPTs in partitioning carbon and reductant in cells and sustaining central carbon
metabolism; and finally, *iii*) do the TPTs contribute to dissipating excess reducing equivalents and
balancing cellular redox conditions? Our study demonstrates, for the first time, the physiological
impact of a null mutation of a highly expressed algal TPT gene.

We have focused this analysis on Chlamydomonas TPT3 (CreTPT3), which exhibits the strongest
sequence similarity to plant chloroplast TPTs among the four predicted chloroplast associated
CreTPTs, and is highly expressed over the diel cycle, especially during the light period. Through
a series of in vitro and physiological analyses we demonstrate that CreTPT3 is essential for the
exchange of fixed carbon, sustaining central carbon metabolism and maintaining intracellular
redox balance.

Results

Features of the triose-P/Pi translocator (TPT) family

Candidate genes encoding triose-P/Pi transporters (TPT) in the Chlamydomonas genome were
identified by blasting the Arabidopsis TPT protein to the Chlamydomonas genome. The TPT
family is the largest drug/metabolite transporter (DMT) family in eukaryotes and includes triose-
P and sugar-phosphate permeases that are associated with chloroplasts; many family members are
still functionally uncharacterized(*16–18*). Chlamydomonas harbors ~32 genes encoding proteins
of the TPT family, with four members (CreTPT1, 2, 3 and 25) predicted to have a transit peptide
and be chloroplast-localized (**Supplementary Table 2**). To learn about the potential substrate

specificities of the four putative chloroplast CreTPTs, their amino acid sequences were aligned

with the plastidial phosphate transporters (PT) from Arabidopsis (AtPTs) (**Supplementary Fig.**

2).

Land plants harbor four classes of phosphate translocators on the inner plastid envelope, including

a glucose 6-phosphate transporter (GPT), a xylulose phosphate transporter (XPT), a

phosphoenolpyruvate transporter (PPT), and a triose-P translocator (TPT)(*6*, *19–22*). The ability

of the transporters to use triose-P/DHAP or 3-PGA is dependent on five highly conserved residues

(H184, K203, Y338, K359, and R360 in AtTPT1)(*13*, *22*). Only CreTPT3 and CreTPT25 contain

all five of these residues (H170, K189, Y322, K345, and R346 in CreTPT3) (**Supplementary Fig.**

2, **3**). In AtTPT1, residue F262 (F248 in Chlamydomonas) is thought to inhibit

phosphoenolpyruvate (PEP) access to the binding site; this residue is replaced by N in AtPPT1

(PPT)(*13*, *22*). CreTPT25 has M at position F248, indicating that unlike the other CreTPTs,

CreTPT25 might have higher preference for PEP (**Supplementary Fig. 3**). These findings suggest

that of the four chloroplast TPT family members, CreTPT3 is most likely to function in specifically

shuttling triose-P or 3-PGA across the chloroplast envelope.

Additionally, a phylogenic tree was constructed to help determine the specificities of the putative

CreTPTs. This tree includes all of the CreTPTs and the plastidial phosphate transporters (PT) from

Arabidopsis thaliana (AtPPT1, AtTPT1, AtGPT1 and AtXPT1). The CreTPT1, 2, 3 and 25 amino

acid sequences cluster with the different AtTPTs (**Fig. 1A**, **Supplementary Fig 3**), again

suggesting that these individual Chlamydomonas transporters have different substrate specificities.

Since CreTPT3 is most closely related to AtTPT1 (**Fig. 1A**, **Supplementary Fig. 3**), the major

triose-P permease in Arabidopsis, we considered CreTPT3 as the Chlamydomonas ortholog of

AtTPT1.

TPT3 transport properties

To help establish the physiological (in vivo) function of CreTPT3, its subcellular localization was

first experimentally confirmed in Chlamydomonas. The CreTPT3 coding sequence with its

putative chloroplast targeting sequence was fused with the VENUS tag (CreTPT3-VENUS) and

expressed under the control of its own promoter in the Chlamydomonas mutant null for CreTPT3.

As shown in **Fig. 1B**, CreTPT3-VENUS was localized to the chloroplast envelope.

To evaluate the substrate specificity of CreTPT3, it was heterologously expressed in

Saccharomyces cerevisiae (yeast) cells and the resulting recombinant protein was biochemically

analyzed using a liposome uptake system. The Cre*TPT3* gene was codon-optimized, expressed in

yeast (**Fig. 1C**), total cell membranes were isolated and then CreTPT3 was reconstituted into

liposomes(*15*, *23*). The chloroplast phosphate transporters in vascular plants can catalyze a strict

Pi/Pi homo-exchange in vitro. The reconstituted CreTPT3 in liposomes was able to catalyze the

signature Pi homo-exchange, whereas in the absence of a counter-exchange substrate, little Pi

uptake was detected. This result demonstrates that the recombinant CreTPT3 protein can function

in Pi/Pi homo-exchange in liposomes (**Fig. 1D**). In contrast, very low Pi uptake rates were observed

for liposomes reconstituted with membranes from yeast cells lacking CreTPT3 (**Supplementary**

Fig. 4), indicating that the detected Pi import activity was the result of CreTPT3 expression in

yeast.

To assess the substrate specificity, the initial rates of Pi uptake into liposomes preloaded with

saturating concentrations (30 mM) of various potential counter-exchange substrates were

determined. As **Fig. 1E** shows, CreTPT3 exhibited the highest activity when DHAP was used as

substrate. The relative initial velocity for Pi uptake in exchange for DHAP was slightly higher than

that of 3-PGA (3-PGA exhibited a 75% exchange rate relative to that of Pi/DHAP) while the Pi

uptake into liposomes preloaded with PEP was much less efficient (about 16% relative to the

Pi/DHAP exchange rate) (**Fig. 1E**). Pi import was negligible when CreTPT3 liposomes were

preloaded with Glc-6-P, Glc-1-P, Fru-6-P, and Gal-1-P (**Fig. 1E**).

Characterizations of the K_M and K_i of CreTPT3 were performed to determine the affinity of the

transporter for the various substrates. CreTPT3 has an apparent Michaelis-Menten constant (K_M)

value of 1.1 +/- 0.2 mM for Pi (**Fig. 1F**), which is comparable to the value obtained for the vascular

plant TPT ortholog(*6*). DHAP and 3-PGA competitively inhibited Pi uptake with an inhibition

constant (Ki) of 0.57 +/- 0.01 mM and 0.79 +/- 0.12 mM, respectively (**Fig. 1F**), suggesting that

CreTPT3 has a higher affinity for DHAP and 3-PGA than for Pi. In the case of PEP, no inhibition

of the Pi/Pi homo-exchange was observed, even at the non-physiologically high concentration of

5 mM. Overall, the results of this in vitro assay indicate that CreTPT3 has a typical plant-TPT

substrate spectrum (*6*). It can transport Pi, triose-P (DHAP) and 3-PGA in a counter-exchange

mode, however PEP might not be a physiologically relevant substrate for CreTPT3, as was shown

for apoplast TPT homologues in other organisms(*13*, *24*).

Isolation of *tpt3* null mutants and their impact on cell growth

To explore the role of CreTPT3 in trafficking carbon and reductant across the chloroplast envelope
in vivo, CRISPR knockouts of Cre*TPT3* were generated. We used the CRISPR-Cas9 editing
system to disrupt the Cre*TPT3* gene in exons 1 and 7, while at the same time integrating the
hygromycin marker gene (*AphVII*) into the edited site (**Supplementary Fig. 5A**). Multiple
independent knockouts of Cre*TPT3* were obtained, including *t3ko1*, with the marker gene inserted
into exon 1, and *t3ko2* and *t3ko3*, with the marker gene inserted into exon 7 (**Supplementary Fig.
5**, **6**) using specific sgRNA2 guides.

To elucidate the physiological roles of CreTPT3, we first examined growth of the parental WT
strain and the knockout mutants under either LL or ML. In contrast to WT cells, growth of the
Cre*TPT3* knockouts (*t3ko1*, *t3ko2*, *t3ko3*) was significantly reduced in LL on either
photoautotrophic (TP) or heterotrophic (TAP) agar medium (**Fig. 2A**, **Supplementary Fig. 7A**,
D), while the growth of the mutant in ML was completely blocked in both TP and TAP (**Fig. 2A**,
Supplementary Fig. 7D). Growth curves were also determined for WT and mutant cells in liquid
medium (TP) in both LL and ML (**Fig. 2B**, **C**); the results were in accord with those observed for
the agar medium growth assays. Additionally, *tpt3* mutant cells in both LL and ML exhibited an
increased cell diameter and formed clusters of cells that appear to be less able to separate following
division (**Supplementary Fig. 8A**, **B**). To further confirm that the growth phenotypes are a
consequence of the *tpt3* knockout, we introduced a wild-type copy of Cre*TPT3* fused to VENUS
into the mutant strains (**Supplementary Fig. 5B**). As demonstrated in **Fig. 2B-E**, ectopic
expression of the wild-type Cre*TPT3* in the *t3ko2* mutant (*C_T3KO2*) rescued the reduced growth
phenotype of the mutant in all light conditions tested in this study (LL/ML/HL).

We also analyzed the chlorophyll content of cells grown photoautotrophically upon transfer from
LL to ML (**Supplementary Fig. 1**) and found that *t3ko2* had reduced chlorophyll content relative
to WT cells after 24 h in ML (**Fig. 2D-F** and **Supplementary Fig. 9**). Cell numbers and total
chlorophyll were quantified following the LL to ML transition. The chlorophyll levels per cell
declined in all strains at 24 and 48 h following the transfer to ML; however, it was lower by more
than half in *t3ko2* (0.34 µg/10^6 cells) relative to either WT cells or the *C_T3KO2* rescued strain
(both ~0.72 µg/10^6 cells) after 24 h of ML (**Supplementary Fig. 9**), with some additional increase

for WT and *C_T3KO2* after 48 h of ML illumination (which might reflect some shading caused

by an increase in the cell density). Finally, when mutant cells were transferred to HL for 24 h they

became highly bleached (**Fig. 2E**).

Taken together, these results indicate that the activity of the CreTPT3 shuttle is essential for growth

over a range of light intensities (LL/ML/HL).

The *tpt3* mutant exhibits hyper-accumulation of 'storage' carbon

To explore the impact of the loss of CreTPT3 activity on carbon partitioning, we quantified carbon

storage (starch and lipids) following a transition of WT and the *t3ko2* mutant from LL to ML.

Lugol staining showed extensive starch accumulation in the *t3ko2* cells after a 48-h exposure to

ML, whereas WT cells were barely stained (**Fig. 3A**). Furthermore, as shown in **Fig. 3B**, there is

an ~55-fold (WT and *t3ko2*: 13.29 and 0.24 μg starch/μg chlorophyll, respectively) increase in

starch accumulation (normalized to chlorophyll) in the *t3ko2* mutant after 24 h of illumination in

ML. The mutant also exhibited a ~24-fold increase in lipid accumulation (on a chlorophyll basis),

as monitored by Nile Red fluorescence, relative to WT and *C_T3KO2* over the same time period

(WT and *t3ko2*: 1038.93 and 41.98 Nile Red fluorescence/chlorophyll, respectively) (**Fig. 3C**, **D**).

Since the chlorophyll in the mutant was approximately 50% relative to that of WT cells after 24 h

in ML, the accumulated starch and lipid on a per cell basis would be ~25-fold for starch and ~12-

fold for lipid in the mutant relative to the WT strains. Furthermore, the mutant cells are much

bigger than WT cells (**Supplementary Fig. 8A**, **B**), which may reflect the large increase of storage

carbon that they accumulated. Accumulated starch granules in the stroma of ML-exposed *t3ko2*

cells are readily visualized under bright field light microscopy (**Supplementary Video 1A**, **B**).

These results suggest that the inability to transport triose-P between the chloroplast and the cytosol

through CreTPT3 resulted in a repartitioning of photosynthetic assimilates (carbon, reductant, and

ATP) toward the synthesis of both starch and neutral lipid.

Cre*TPT3* deletion leads to accumulation of CBBC/glycolytic/gluconeogenic intermediates

To understand the metabolic consequences of the loss of CreTPT3 on growth under LL and ML,

comparative metabolic analysis of WT and *t3ko2* mutant was performed on cells grown at LL and

after shifting them to ML for 45 min or 6 h. Pool sizes of various central carbon metabolites,

particularly those of the CBBC/glycolysis/gluconeogenesis pathways and intermediates of the

TCA/glyoxylate cycle (**Fig. 3F**), were quantified. Data was normalized to both chlorophyll content

(**Fig. 3E**) and cell number (**Supplementary Fig. 10B**). The fold-change in the quantity of each

metabolite in the mutant relative to WT under LL or at 45 min and 6 h after the switch to ML is

given in **Fig. 3G** and **Supplementary Fig. 10C**.

Under LL and following transfer to ML, the *t3ko2* mutant maintained a significantly larger pool

(2 to 6-fold) of glycolytic/gluconeogenic intermediates [DHAP, 3-PGA and FBP (all three shared

with CBBC), G6P, PEP], and of some metabolites of the CBBC pathway (E4P, RU5P/X5P, R5P)

compared to WT, suggesting that the loss of CreTPT3 results in a back-up of these metabolites

within the cell; our hypothesis is that they are accumulating inside the chloroplast due to the lack

of CreTPT3 activity. This is supported by the finding of marked accumulation of starch and TAG

in the mutant strain (**Fig. 3A-D**). While the pool sizes of fumarate and malate (metabolites of the

TCA/glyoxylate cycle) were similar in both strains under LL, following the transition of the cells

to ML, WT cells showed a significant increase in the pool sizes of those metabolites while the

mutant maintained a lower level, indicating that the loss of CreTPT3 either directly (by supplying

precursors) or indirectly (metabolic rewiring) impacts their levels.

Thus, the one primary function of CreTPT3 would be the export of photosynthetically-synthesized

reduced carbon, which would drive metabolic processes in the cytoplasm and other cellular

compartments, while importing Pi into the chloroplast to sustain ATP synthesis.

The *tpt3* mutant has reduced PSII activity when grown in moderate light

To understand the impact of the loss of CreTPT3 activity on photosynthetic electron transport

(PET), we quantified photosynthetic activities following a transition from LL to ML, as shown in

Supplementary Fig. 1. Photosynthetic O_2 evolution rates (OERs) were measured for WT, *t3ko2*

and *C_T3KO2* at a range of light intensities. LL acclimated WT, *t3ko2*, and *C_T3KO2* showed

comparable OER at intensities below 200 μmol photons m^{-2} s^{-1} while at saturating light intensities

of ≥600 μmol photons m^{-2} s^{-1}, the OER of *t3ko2* was diminished slightly relative to the WT and

complemented strain (**Fig. 4A**). Upon acclimation of the cells to ML for 24 h, the *t3ko2*

consistently displayed low OER relative to WT and *C_T3KO2* under all actinic light intensities

used (**Fig. 4B**).

The PSII quantum yield, ΦPSII, is maximum after a dark pre-incubation (Fv/Fm). The Fv/Fm in
the mutant grown in LL was similar to that of WT and *C_T3KO2* immediately following the dark
incubation (**Fig. 4C**). The LL-acclimated WT and *t3ko2* cells exhibited comparable ΦPSII values
when the actinic light intensity was below 100 μmol photons m^{-2} s^{-1}, but decreased by about 20-
25% when the actinic light was increased to 400 μmol photons m^{-2} s^{-1}; that decrease was
maintained even when the actinic light was raised to over 1,000 μmol photons m^{-2} s^{-1}
(**Supplementary Fig. 11**). In contrast, dark acclimated mutant cells that were exposed to ML for
6 h exhibited an Fv/Fm that was about 2-fold lower than that of WT and *C_T3KO2* and continued
to decline over a period of 24 h in ML, with a 3-fold decrease in Fv/Fm for *t3ko2* compared to that
of WT (**Fig. 4C**, **Supplementary Fig. 11**).

The oxidation of the PQ pool is the slowest step in the PET chain between PSII and PSI(*25*, *26*).
The redox potential of the PQ pool changes upon increasing the irradiance to which the cells are
exposed. Thus, the redox potential of the PQ pool serves as a biological measure for evaluating
the balance between the photochemistry of PSII and linear photosynthetic electron transport.
Changes of the redox state of the PQ pool (measured as 1-qL)(*27*) in LL and ML acclimated cells
were analyzed at increasing actinic light intensities (**Fig. 4D**, **Supplementary Fig. 12**). In LL-
grown cells, the PQ pool was significantly more reduced in *t3ko2* even at low level illumination
compared to that of WT and the rescued strain. When the actinic light intensity was elevated to
400 μmol photons m^{-2} s^{-1}, the reduction levels of the PQ pool increased for all strains, although it
was still much more reduced in *t3ko2* compared to that of WT and the rescued strains (**Fig. 4D**,
Supplementary Fig. 12).

For cells that had been acclimated to ML (6 h and 24 h), the PQ pool was already very highly
reduced in *t3ko2* following exposure to LL, while it remained highly oxidized for both the WT and
the rescued strain (**Fig. 4D**, **Supplementary Fig. 12**). The level of PQ reduction in ML-acclimated
t3ko2 increased only slightly with an increase in the actinic light intensity (**Fig. 4D**,
Supplementary Fig. 12). There was also an increase in the PQ pool reduction of WT and
C_T3KO2 as the light intensity was raised, however, even at the highest light intensity the WT and
rescued strains maintained a much more oxidized PQ pool than that of the mutant; the highest
levels of 1-qL for WT and *C_T3KO2* were still ~2-fold lower than that of *t3ko2* (**Fig. 4D**,
Supplementary Fig. 12). Overall, inactivation of CreTPT3 resulted in attenuated PSII activities

(Fv/Fm, OER) and hyper-reduction of the PQ pool, suggesting a block in electron flow
downstream of the PQ pool.

Impact of *tpt3* mutant on PSI activity

To determine if PSI activity was also impacted in the mutant cells, we analyzed PSI/P700
oxidation/reduction kinetics following exposure of LL-grown cells to ML. Levels of photo-
oxidizable P700 were similar in all LL-acclimated strains (**Fig. 4E**). However, after 6 h in ML, the
level of photo-oxidizable P700 declined in the mutant to ~40% of that in WT and *C_T3KO2* cells
(**Fig. 4F**), followed by slightly more inhibition of PSI activity after 24 h of exposure to ML (**Fig.**
4G). These deleterious impacts on photosynthetic electron transport likely reflect a defect in
routing carbon/energy out of the chloroplast, which in turn will cause hyper-accumulation of
reductant in the chloroplast, which would potentially result in a backup of PET, elevated ROS
accumulation and damage to the photosynthetic apparatus.

CreTPT3 is crucial for balancing metabolite flux and chloroplast redox

Photosynthetic organisms mainly accumulate ROS in chloroplasts and mitochondria as byproducts
of aerobic metabolism(*28*). To assay ROS production in the mutant, we used the fluorescent probe
CM-H2DCFDA which, upon exposure to increasing ROS levels, is converted to the green-
fluorescent molecule dichlorofluorescein (DCF). DCF fluorescence was visualized by confocal
microscopy. As shown in **Fig. 5A** and **B**, ROS levels in the *t3ko2* mutant were markedly increased
(~3-fold) after 48 h in ML, while no difference in the level was detected in WT cells.

For an alternative, dynamic method for evaluating redox changes in the chloroplast, we used the
redox sensitive green fluorescent protein roGFP2 (**Supplementary Fig. 13**) that was targeted to
the chloroplast stroma. Two cysteine residues were engineered into the roGFP2 to form a
reduction-oxidation-dependent disulfide bond that detects intracellular changes in ROS
accumulation(*29*). roGFP2 was shown to monitor ratiometric redox changes of glutathione(*29*, *30*).
The ascorbate–glutathione (GSH) system is central to ROS detoxification(*31*), with the redox
potential of glutathione reflecting ROS levels. LL-acclimated WT cells and the *t3ko2* mutant
exhibited similar levels of chloroplast roGFP oxidation (**Fig. 5C**). Upon transfer of WT cells to
ML, the oxidation level of roGFP in the chloroplast increased by ~2-fold after 1.5 h, compared to
that of WT cells maintained in LL, followed by a decrease after a 6 h ML exposure (**Fig. 5C**). The

changes of redox in WT upon a LL to ML transition is in accord with previous results(*32*, *33*).

Furthermore, the synthesis of both GSH and ascorbate increases when the redox balance is shifted

to a largely oxidized state after transferring cells to a higher light intensity, and recovers several

hours after the transfer(*32*, *33*) as the cells acclimate to the higher light conditions. A similar redox

pattern determined by chloroplast roGFP2 was observed for the *t3ko2* mutant after it was shifted

to ML, however, the mutant showed a much greater increase in chloroplast oxidative conditions,

with a 5.0-fold increase after 1.5 h, and a 2.0-fold increase after 6 h, or 1.8- and 1.6-fold higher

than the levels measured in WT cells (**Fig. 5C**).

As triose-P is exported to the cytosol by CreTPT3, it could potentially donate redox equivalents to

mitochondrial electron transport and alter mitochondrial ROS production. Therefore, we measured

mitochondrial redox levels at different light intensities in both WT and the *t3ko2* mutant using the

roGFP sensor targeted to the mitochondrion. Upon a LL to ML transition, neither WT nor *t3ko2*

displayed a change in fluorescence from the mitochondrial targeted roGFP (**Fig. 5D**). These results

suggest that the mitochondrial redox level is maintained after shifting either LL acclimated WT or

t3ko2 cells to ML. This is in agreement with the results obtained for diatoms; the roGFP oxidation

level in diatom mitochondria exhibited either no or minor changes in response to various stress

conditions, including nitrogen limitation, HL, and iron limitation(*34*). Overall, these results

suggest that the increase in oxidative conditions caused by a LL to ML transition is primarily

localized to the chloroplast in Chlamydomonas and that the inability to export triose-P through

TPT3 markedly increases the chloroplast oxidative stress level.

Another issue that we addressed was whether CreTPT3 is important for dark metabolism in

Chlamydomonas. We analyzed redox changes in WT and the *t3ko2* mutant upon transferring cells

from LL to the dark. Intriguingly, ROS accumulation in WT was 5 times higher than that of *t3ko2*

after 48 h of darkness (**Fig. 5E**, **F**). The oxidation level of chloroplast roGFP2 in WT was slightly

reduced during the first 3 h of darkness, followed by a >1.5-fold increase after 6 h of darkness

compared to LL-acclimated cells (**Fig. 5G**). This is in keeping with previous observations in

Arabidopsis in which exposing leaves to extended darkness resulted in an initial decrease in the

degree of oxidation of roGFP2 in the chloroplast followed by an increase to the original LL level

when the darkness was prolonged(*35*). Unlike the slight decrease in roGFP2 oxidation in the WT

chloroplast, the oxidation state of the chloroplast in *t3ko2* cells declined by 2.2-fold relative to the

original LL conditions and by 1.77-fold relative to WT after 3 h in the dark (**Fig. 5G**). The chloroplast oxidation level in *t3ko2* did not recover to the level of WT cells after 6 h of darkness but attained its initial LL level after 24 h in the dark (**Fig. 5G**). The mitochondrial oxidation level in *t3ko2* during the dark period was slightly but not strikingly higher than that in WT (**Fig. 5H**). In sum, these results suggest that CreTPT3 also affects cellular redox levels in the dark.

In addition to the light conditions, we analyzed changes in the starch contents over the diel cycles. As shown in **Fig. 5I**, *t3ko2* accumulates 7 times more starch than that of WT cells after 12 h in the light (*t3ko2*, WT: 10.21, 1.403 µg starch/µg chlorophyll). Similar to WT and the rescued strain, *t3ko2* can mobilize 70% of the starch at the end of the light period, indicating that *t3ko2* contains much more carbohydrates in the chloroplast in the dark (**Fig. 5I**).

Discussion

Photoassimilates that serve as substrate for the chloroplast triose-P/Pi translocator (TPT) have been studied in various vascular plants (e,g, tobacco, Arabidopsis, potato, etc). However, surprisingly few studies have characterized chloroplast envelope localized members of the triose-P family in trafficking fixed carbon in single-celled photosynthetic organisms(*13–15*); their physiological role(s) in algae has not been elucidated. The Chlamydomonas genome encodes 4 putative triose-P transporters predicted by TargetP-2.0 to be routed to the chloroplast. Cre*TPT3* is the most highly expressed among the chloroplast associated TPTs (**Supplementary Table 3**, **Supplementary Fig. 14A**). A detailed analysis comparing Chlamydomonas TPT proteins with those of other organisms revealed that CreTPT3 is most similar to the *Arabidopsis thaliana* TPT1 (**Fig. 1A**). In this study, we localized CreTPT3 to the chloroplast envelope. To explore the function of CreTPT3, we expressed it in yeast cells and performed in vitro studies to assay its activity in liposomes (**Fig. 1C-F**) and demonstrated the capacity of this translocator to use Pi, DHAP and 3-PGA as substrates (**Fig. 1D-F**), similar to plant TPT(*6*). Hence, it appears that CreTPT3 can exchange chloroplast/stromal triose-Ps for cytosolic Pi, and may also serve as redox shuttle by catalyzing DHAP/3-PGA exchange. We also generated mutants disrupted for the Cre*TPT3* gene to examine the impact of a loss of function mutant on the physiology of Chlamydomonas.

CreTPT3 is critical for the distribution of photosynthate

An Arabidopsis mutant for the chloroplast TPT was isolated 20 years ago(*10*). The growth and

photosynthesis of this mutant, designated *ape2*, was not affected under LL conditions(*9*), although its starch content was slightly elevated relative to that of WT Arabidopsis(*9*). The absence of AtTPT in LL-grown plants appeared to be fully compensated for by directing more photosynthate toward starch synthesis and turning over the starch rapidly in the light to generate sugars like hexose phosphate (hexose-P) and other metabolites that can be exported from the chloroplast. Furthermore, while *ape2* grew in HL, its growth was attenuated relative to WT Arabidopsis(*9*), although there was no detectable change in the level of starch(*10*). In potato, a 2-3-fold increase in starch was observed in transgenic plants with diminished levels of the chloroplast targeted TPT(*11*). The reduction in TPT activity led to a strong growth retardation during the young stages of development. Similar to Arabidopsis, a tobacco strain with decreased TPT activity was able to compensate for the defect by accelerating the synthesis and turnover of starch and the transport of hexose from the chloroplast(*8*). These observations suggest that various vascular plants respond to a defect in the export of triose-P from chloroplasts by moving the fixed carbon into a transient starch pool that is converted to sucrose/glucose/glucose-P through accelerated hydrolysis. Glucose-P and potentially other metabolites generated from starch degradation can then be exported from the chloroplast and trafficked to sink organs, providing them with carbon skeletons and energy for anabolic processes(*36*). In contrast, a deficiency of TPT in rice, which uses sucrose stored in leaves as its major transitory form of fixed carbon rather than starch, led to reduced photosynthetic rates and about a 2-fold decrease in the level of starch relative to WT plants(*37*).

To probe the in vivo function of CreTPT3, we generated multiple null mutants, designated *t3ko* and characterized one of the mutants (*t3ko2*) and the rescued strain in depth. Our results show that the flux of fixed carbon trafficked through the CreTPT3 is crucial for Chlamydomonas growth under all light intensities tested, and that it becomes essential for survival at the higher intensities (**Fig. 2**, **Supplementary Fig. 7**). Chlamydomonas, unlike plant cells, contains little intracellular sucrose, with starch as its major form of storage carbon(*38*). In addition to a pronounced growth phenotype, the *t3ko2* mutant exhibited a dramatic change in its carbon partitioning. After 24 h in LL, *t3ko2* contained ~16-fold more starch than that of WT cells (**Supplementary Fig. 10A**). After 24 h in ML, the starch and lipid contents of *t3ko2* were 55 and 24-fold higher, respectively, than that of WT cells (**Fig. 3A-D**).

Furthermore, in Arabidopsis, elimination of the chloroplast targeted TPT protein did not

significantly alter the profiles of metabolites of either the CBBC or the glycolytic/gluconeogenic pathways, suggesting the occurrence of compensatory mechanisms that redirect carbon from the chloroplast of the mutant plant(*39*). In contrast, a Chlamydomonas mutant null for Cre*TPT3* exhibited a dramatic increase of most metabolites in the glycolytic/gluconeogenic pathways, suggesting that triose-P exchange catalyzed by CreTPT3 was not readily compensated for by alternative fixed carbon shuttles. CreTPT1, CreTPT2 and CreTPT25, the other shuttles that potentially transport triose-P or hexose-P across the Chlamydomonas chloroplast envelope, are unable to compensate (at RNA level) for the loss of CreTPT3 (**Supplementary Fig. 14B-D**). These results demonstrate the critical role of CreTPT3 in shuttling photoassimilates in Chlamydomonas. The retention/accumulation of photoassimilates in the chloroplast appears to redirect them toward the synthesis of starch and lipid, which would cause a marked deficit of fixed carbon in other cellular compartments, such as the cytoplasm and mitochondrion, and severely compromise physiological processes including cell growth and division; a loss of CreTPT3 impacts carbon partitioning and growth under all light conditions tested (LL/ML) (**Fig. 2**, **3**, **Supplementary Fig. 10**).

Metabolic flux through CreTPT3 is essential for redox homeostasis

Studies of photosynthetic activities of *t3ko2* in ML support our hypothesis that the mutant has a reduced capacity to utilize metabolites for anabolic processes required for growth. The marked alteration in metabolite pools observed in the mutant (**Fig. 3E**) could feedback on PET and potentially elicits photosynthetic control, constricting electron flow at the cyt b_6f complex as a consequence of the elevation of ΔpH across the thylakoid membranes(*40*, *41*). Such control would result in hyper-reduction of the PQ pool (**Fig. 4D**) and reduced electron flow to PSI, which would help protect PSI from photoinhibition. Furthermore, *t3ko2* cells are strongly photoinhibited following their transfer from LL to ML, conditions that cause little change in PET of WT and *C_T3KO2* (**Fig. 4B-C**). These results strongly suggest that CreTPT3 is essential for maintaining the flow of fixed carbon and reductant from the chloroplast to the cytosol where it can be allocated for various anabolic processes that support normal cell growth.

The malate-OAA shunt represents another route that, under high redox stress, might have compensated for the loss of the CreTPT3 shuttle. Intriguingly, malate levels in *t3ko2* were 4-fold lower than in WT cells following a transition from LL to ML (**Fig. 3E**, **G**) suggesting that the

malate-OAA shunt cannot compensate for a loss of CreTPT3 activity in Chlamydomonas. A previous study of metabolic flux analysis during heterotrophic growth of Chlamydomonas showed that the TPT shuttle is almost 10-fold more active than the malate-OAA shunt(*42*).

Cre*TPT3* is the most responsive/upregulated of the four Chlamydomonas, chloroplast localized *TPT* genes following exposure of cells to nitrogen, sulfur or iron deprivation (**Supplementary Fig. 15 A-C**), which reinforces the hypothesis that it plays an important role in controlling chloroplast redox conditions. Deprivation of Chlamydomonas for nitrogen or sulfur caused dramatic attenuation of PET(*43*) as the cells elicited an array of 'energy management' strategies to balance energy generation and utilization(*43*). Elevated NPQ, chlororespiration and lipid/starch accumulation are well recognized mechanisms to dissipate excess light energy and minimize ROS production in Chlamydomonas in response to nutrient deprivation(*43*). In this study, TPT3-dependent chloroplast triose-P/Pi transport appears to be critical for exporting 'overflow' electrons/reductant to other cellular compartments. When this function is compromised, the chloroplast accumulates various metabolites, which results in altered substrate availability for chloroplast-localized metabolic processes. Additionally, hyper-reduction of chloroplast redox components, such as the PQ pool, and elevated production of ROS can lead to dysfunction of the photosynthetic apparatus (and other metabolic activities), almost mirroring the response of the cells to N-stress.

This study also suggests that CreTPT3 impacts the redox levels of the chloroplast in the dark. It was proposed that during an extended period in the dark, Arabidopsis becomes sugar starved which leads to the diminished availability of reductant for scavenging ROS(*35*, *44*). This hypothesis has been supported by the transcriptome analysis of an Arabidopsis starch accumulation-defective mutant (*pgm*) that has a lower sucrose level during the night than the WT plants(*45*). The upregulation in the dark of the majority of the general ROS-induced transcripts in *pgm* is more rapid than that of the WT, suggesting oxidative stress can be related to sugar starvation(*35*). Although our interpretation is speculative, given the highly reducing state of the chloroplast and the elevated accumulation of starch in the chloroplast in *t3ko2*, the mutant likely maintains elevated levels of sugars in the chloroplast during the dark period as it mobilizes the high level of carbon that was stored during the light period. Indeed, as shown in **Fig. 5I**, *t3ko2* accumulates a higher amount of carbohydrates in the dark, potentially leading to a greater capacity to generate reduced

carbons that efficiently scavenge excess ROS in the dark.

CreTPT3 is central to metabolic and redox networks

ADP-glucose pyrophosphorylase (AGPase), a key enzyme in starch synthesis, is known to be
exquisitely sensitive to allosteric regulation; it is activated by 3-PGA and inhibited by Pi(*46*).
When the *t3ko2* mutant was exposed to LL, there was already a dramatic increase of 3-PGA (6-
fold) relative to WT cells (**Fig. 3E**, **G**). The loss of CreTPT3 might also lead to less import of Pi
into chloroplasts resulting in an elevated ratio of 3-PGA/Pi in *tpt3* compared to that of the WT
cells. Thus, even in LL, AGPase in the mutant would become active, which in turn would lead to
starch accumulation (**Supplementary Fig. 10A**).

In chloroplasts, the high reductant levels can also modulate the activity of various redox-sensitive
target proteins via thiol-based redox regulation. Chloroplasts harbor two thiol-based redox systems
that independently regulate plastid metabolism by communicating redox conditions to specific
target enzymes, thereby increasing or decreasing their activities. One system, FDX/TRX reductase,
obtains reducing equivalents from FDX (which is reduced by PET)(*47*), while the other involves
NADPH-TRX reductase (NTRC), which uses NADPH as its substrate(*48*). Both systems can
activate AGPase and starch biosynthesis(*49*, *50*). Elevated reduction of the PQ pool in *t3ko2*
compared to that of WT (**Fig. 4D**, **Supplementary Fig. 12**) could activate the FDX/TRX system
while increased NADPH availability in the mutant might be instrumental in activation of the
NTRC (NADPH-TRX reductase) system.

Additionally, the ROS levels in *t3ko2* (**Fig. 5A-C**) could potentially activate alternative regulatory
pathways. ROS was initially considered harmful, causing oxidative stress, damaging lipids,
proteins, and DNA, eventually resulting in cell death(*51*). Increasing evidence, however, suggests
that ROS can also function in diverse signaling pathways(*51*) that regulate various cellular
processes(*52*, *53*) such as growth, the cell cycle, senescence(*54*), and acclimation to stressful biotic
and abiotic conditions(*55*). ROS can also regulate expression of some chloroplast genes encoding
proteins associated with photosynthesis and modulation of PET(*56*). For example, ROS inhibits
de novo synthesis of the PSII reaction center protein D1, which would lead to photoinhibition
during high light acclimation(*57*, *58*). Moreover, when PSI becomes highly reduced, superoxide
and singlet oxygen both elicit PSI photoinhibition(*59*).

Overall, Chlamydomonas mutants lacking TPT3 develop a hyper-reduced chloroplast landscape

with elevated ROS production (**Fig. 5A-C**). After 6 h or more in ML, PSII and PSI activities in

the mutant become compromised, most likely a consequence of elevated ROS levels (see model

on **Fig. 6**). The chloroplast generated ROS serves as a messenger for reporting redox information

(e.g. light levels, stress conditions) to the regulatory machinery of the cell, which results in

modulation of expression of nuclear genes that would ameliorate the impact and damage associated

with elevated ROS levels(*60–62*). Many studies have shown that stress-induced

lipid/triacylglycerol (TAG) accumulation also correlates with an increase in ROS production(*63*);

ROS may activate diacylglycerol acyltransferases (DGAT) and phospholipid diacylglycerol

acyltransferase activities (PDAT) thereby stimulating lipid accumulation(*64*).

As depicted in **Fig. 6**, we propose that various tiers of redox regulation are potentially responsible

for the differential physiological responses of Chlamydomonas *tpt3* knockout mutants. When

t3ko2 cells are initially transferred from LL to ML, the triose-P pool and metabolites derived from

that pool accumulate in the mutant because of its reduced capacity to move triose-P out of the

chloroplast, which also causes reduced CBBC activity, hyper-reduction of PET and the

accumulation of ROS (which would inhibit photosynthetic activity). The hyper-reduction of PET

and accumulation of carbon metabolites in the chloroplast of the *tpt3* mutants would activate

AGPase through allosteric regulation and by the FDX/TRX and NTRC redox systems, which

would result in starch hyper-accumulation (**Fig. 6**). Furthermore, increased ROS/RS accumulation

and the elevated redox state of PET would elicit retrograde signals that mediate changes in

expression of both chloroplast and nuclear genes, marshaling the synthesis of specific activities

that function to ameliorate the impact of the hyper-reduced state of the chloroplast resulting from

the compromised ability of the cells to export reductant and fixed carbon from the organelle (**Fig.**

6).

Material and methods

Strains and culture conditions

WT Chlamydomonas M10 (CC-4403, isogenic line derived from CC-124) was used as the parental

strain for the generation of knockout mutants. Cultures were routinely cultivated in growth

chambers (LED-41L2, Percival Scientific, Inc.) at 25°C with continuous shaking on an orbital

shaker (VWR OS-500 Shaker) at 120 rpm, in Tris-acetate-phosphate (TAP) medium(*65*). The

cultures were illuminated with continuous cool white LEDs at low light (LL, 30 µmol photons m^-

^{2} s^{-1}). Experiments were mostly performed with cells grown in the photoautotrophic medium TP

(TAP medium without acetate), and in TAP medium in some cases, at 25°C, and sparged with air

while being shaken at 120 rpm in a growth chamber (LED-41L2, Percival Scientific, Inc.). For

growth assays, cultures were inoculated at a density of 0.02 at $OD_{750\ nm}$ (~1×10^5 cells/mL) in TP

sparged with air under either LL or moderate light (ML, 250-300 µmol photons m^{-2} s^{-1}) intensities.

For spectrophotometric and chlorophyll fluorescence analyses, precultures were grown in TAP in

LL to mid-exponential phase, the cells were pelleted by centrifugation at 800xg, for 3 min at room

temperature, and the pellets were washed once with TP and then resuspended in TP at a density of

1-1.5 µg/mL chlorophyll. These cultures were grown in TP sparged with air and after 16-24 h in

LL, were transferred to ML. To avoid self-shading, the culture density was adjusted to 1-1.5 µg/mL

chlorophyll before ML exposure.

Cell samples exposed to the different light intensities were diluted to perform spot tests for

photoautotrophic growth on TP agar plates and for mixotrophic growth on TAP agar plates. Agar

plates were incubated for 7 d under either LL or ML (cool white LED) at 25°C.

Phylogenetic analysis

Protein sequences were aligned using MUSCLE(*66*). The best-fit nucleotide substitution model

was generated based on the Akaike information criterion (AIC) using MEGA7(*67*). Maximum

likelihood phylogenetic trees were constructed based on the best-fit model GTR (general time

reversible) + G (gamma distribution) + I (invariable sites) with 100 bootstrap replicates using

MEGA7(*67*).

Expression of CreTPT3 in yeast

The DNA sequence encoding the mature CreTPT3 protein was codon-optimized for expression in

Saccharomyces cerevisiae (GeneART, ThermoFisher Scientific). This coding sequence was

inserted in frame with an N-terminal His tag into the yeast vector pYES-NTa (ThermoFisher

Scientific) using Gibson cloning (NEB). Briefly, pYES-NTa was linearized with BamHI and the

CreTPT3   cDNA   amplified   with   the   primer   pair   fwd:5'-

gacgataaggtacctagcGCTTCTGCTGCTGATGCTC-3'   and   rev:5'-

agaattccaccacactgTTAAGCGGCAGCTGGAGAAG-3'. This cDNA was ligated into the

linearized pYES-NTa vector, which was then transformed into the yeast strain INVSc1 (MATa,

his3D1, leu2, trp1-289, ura3-52 / MATa, his3D1, leu2, trp1-289, and ura3-52, Thermo Fisher

Scientific) using the lithium-acetate/PEG method(*68*). Transformed yeast cells were selected on

synthetic complete medium containing 2% (w/v) glucose with the uracil auxotrophic marker.

Galactose-inducible expression of His-CreTPT3 in yeast was performed as described in(*15*). The

presence of the His-tagged fusion protein was verified by standard SDS-PAGE and immunoblot

analysis using an anti-His-antibody conjugated with horseradish peroxidase (Miltenyi Biotech).

Reconstitution into liposomes and transport activity assays

For uptake studies, yeast membranes from cells with and without recombinant His-CreTPT3 were

enriched and reconstituted into 3% (w/v) L-alpha-phosphatidylcholine using a freeze-thaw-

sonication procedure(*15*). The reconstituted liposomes were preloaded with 30 mM Pi or the

indicated phosphorylated metabolites that were tested as potential transport substrates. As a

negative control for antiport activity, liposomes were also generated without preloading. The

external counter-exchange substrate was removed via gel filtration on Sephadex G-25M columns

(GE Healthcare). Transport assays were initiated by adding 0.25 mM [alpha-^{32}P]-phosphoric acid

(6,000 Ci/mmol) and performed as previously described in Linka et al. (2008). The Km for Pi was

determined by measuring the initial velocity at each of six external Pi concentrations between 0.05

and 5 mM. To obtain competitive inhibition constants (Ki), the uptake of 0.25 mM Pi into

liposomes containing 30 mM Pi was measured over a 4 min period in the presence of increasing

external inhibitor concentrations (0.05 - 5 mM). Three biological replicates were performed for all

described experiments. GraphPad Prism software version 9.3.0 was used for non-linear regression

analyses of the kinetic data.

Vector construction, transformation and complementation

The pRam118_VENUS plasmid which harbors the VENUS gene and the *AphVII* cassette (confers

resistance to hygromycin) (Cre01.g045550)(*69*), was used to express Cre*TPT3* (Cre01.g045550).

This plasmid was linearized with HpaI (NEB). Primers: g*TPT3*_pRam118_f and

g*TPT3*_pRam118_r (**Supplementary Table 1**) were used to amplify Cre*TPT3* genomic DNA

containing an overlap with the linearized pRam118_VENUS vector. For generating the plasmid

pRam118-CreTPT3&VENUS, genomic DNA of Cre*TPT3* was assembled with

pRam118_VENUS plasmid using Gibson assembly(*70*). A 1,000 bp sequence upstream of

Cre*TPT3* and containing the promoter region of the gene was amplified using the primers

TPT3pro1000_f and TPT3pro1000_r (**Supplementary Table 1**). The 3' UTR of Cre*TPT3* was

amplified using TPT_3UTR_f and TPT3_3UTR_r (**Supplementary Table 1**). The *PSAD*

promoter and the *RBCS2* 3'UTR of pRam118-CreTPT3&VENUS were replaced by the amplified

fragments of the 1,000 bp upstream region and the 3' UTR of Cre*TPT3*, respectively. This final

vector, designated pRam118_CreTPT3, contains the original Cre*TPT3* promoter (driving

expression of Cre*TPT3*), the genomic DNA sequence of Cre*TPT3*, VENUS and the Cre*TPT3* 3'

UTR plus the *AphVII* cassette. The Chlamydomonas *tpt3* mutant was transformed by

electroporation with a total amount of 2 μg (~500 ng/μL) of pRam118_CreTPT3 that was

linearized with AseI. Next, the linearized plasmids were added into 250 μL of cell suspension at

the concentration of ~3×10^8 cells/mL per reaction. The GeneArt MAX Efficiency Transformation

Reagent for algae (Invitrogen) was used for introducing the plasmid by electroporation following

the instructions provided by the manufacturer. Transformants were selected on solid TAP medium

containing hygromycin (10 μg/mL; Enzo Life).

Subcellular localization

Drug resistant transformants were visualized for VENUS fluorescence as previously described(*69*).

In brief, transgenic cell lines resistant to hygromycin were screened for VENUS fluorescence using

a microplate reader (Infinite M1000; TECAN). Excitation and emission settings were as follows:

VENUS, excitation 515/12 nm and emission 550/12 nm; chlorophyll excitation 440/9 nm and

emission 680/20 nm. The TCS SP8 confocal laser-scanning microscope (Leica) was used to

visualize the VENUS fluorescence signal. Detailed methods for visualization can be found in(*69*).

CRISPR-Cas9 mediated mutagenesis

The protocol for disruption of Cre*TPT3* (Cre01.g045550) was adapted from(*71*). Chlamydomonas

WT strain CC-124 was used for mutant generation. WT cells were cultured under continuous light

(50 μmol photons m^{-2} s^{-1}) for 2 d to a density of $3\text{-}5 \times 10^6$ cells/mL. The cells were then concentrated

to 2×10^8 cells/mL in 0.5×TAP supplemented with 80 mM sucrose. Two single guide RNAs

(sgRNA) were designed by CHOPCHOP and synthesized by Integrated DNA Technologies (IDT).

The sequences of the generated sgRNA are: *TPT3*-sg1 (5′-

CGCUGGGCGTCACUUCCCGGCGG-3′) for editing of exon 1 and *TPT3*-sg2 (5′-

AAGGCCGCUAUCGCCAACGUGGG-3′) for editing of exon 7. Prior to electroporation, Cas9

(IDT) and sgRNAs were incubated together at 37 °C for 30 min. Approximately 500 ng PCR

product of the *AphVII* cassette, which confers resistance to hygromycin, was added to the RNP

(ribonucleoprotein) mixture. 250 µl aliquots were electroporated using Super Electroporator

NEPA21 type II (NEPA GENE). After 16 h of recovery in very low light (10-15 µmol photons m⁻

² s⁻¹), cells were plated onto solid TAP medium containing 10 µg/mL hygromycin. Sense or

antisense-oriented knock-ins of *AphVII* were determined by amplification using primer pairs with

one primer annealing to the genomic sequence and the other to the inserted sequence

(**Supplementary Table 1**). The amplified fragments were sequenced to verify the insertion sites

(ELIM BIOPHARM, Hayward, USA).

Complementation of mutants of Cre*TPT3*

Mutant strains were transformed with the linearized plasmid pRam118_CreTPT3 plasmid. After

transformation, transformants were selected in ML and then screened for VENUS fluorescence.

Colonies containing VENUS and an *AphVII* cassette knock-in at the Cas9 targeted site were

examined for accumulation of the CreTPT3 protein by immunoblots using CreTPT3 antibody

generated by GenScript USA Inc (Piscataway, USA). Colonies of immuno-positive cells were

subjected to growth complementation assays using spot tests under ML on solid TAP or TP

medium.

P700 activity measurements

Absorbance spectroscopy for P700 analyses was performed in dark adapted liquid cultures (15

µg/mL chlorophyll, in 20 mM HEPES-KOH, pH 7.2, and 10% ficoll) using a JTS-100

spectrophotometer (SpectroLogiX, TN). Actinic light was provided by an orange LED (165 µmol

photons m⁻² s⁻¹) for PSI oxidation, followed by a saturating pulse and a dark incubation. 20 µM

DCMU and 1 mM hydroxylamine were added to the cell suspension to inhibit linear electron flow

(LEF) prior to the measurement. P_{700} activity was measured by monitoring the absorbance at 705

nm (interference filter 6 nm FWHM was used to create a narrow excitation beam); the absorbance

at 740 nm was used to correct for an unspecific contribution to the 705 nm signal(*72*).

Chlorophyll fluorescence analysis

630 Chlorophyll fluorescence, used to evaluate photosynthetic electron transport, was measured using

631 a DUAL PAM-100 fluorometer. Cells were acclimated in the dark for 20 min. Light curves were

632 generated by illuminating cells with increasing light intensities (0, 10, 50, 100, 200, 400, 800, 1000,

633 1200 μmol photons m^{-2} s^{-1}) for 2 min at each intensity. 1 mM CO_2 ($NaHCO_3$) was added to the

634 reaction mix as an electron receptor for the CBBC.

ROS measurements and roGFP2 imaging analysis

All strains were grown photoautotrophically under LL for 16-24 h. After dilution with fresh

medium, the cultures were transferred to ML or the dark for >48 h and then stained with CM-

H2DCFDA (Thermo Fisher Scientific) as described in([73]) for detecting ROS. Briefly, after the

treatment (e.g. 48 h in ML), 10 million Chlamydomonas cells (treated in various ways) were

pelleted and washed once with 1×PBS. The cells were then resuspended in 1×PBS containing 8

μM CM-H2DCFDA and incubated at room temperature in the dark for 30 min. Following this

incubation, the cells were washed three times with 1×PBS buffer, the fluorescent signals were

either visualized using TCS SP8 confocal laser-scanning microscope (Leica) or quantified with a

microplate reader (Infinite M1000; TECAN). Excitation and emission settings for the microscope

were: 488 nm/510-530 nm HyD SMD hybrid detector for ROS, and 488 nm/650-700 nm-HyD

SMD hybrid detector for chlorophyll autofluorescence. Excitation and emission settings for the

plate reader were as follows: ROS excitation 488/5 nm and emission 530/12 nm; chlorophyll

excitation 514/5 nm and emission 690/5 nm.

Constructs containing chloroplast or mitochondria targeting sequences([74]) fused to codon-

optimized roGFP2([29]) were transformed into WT and mutant strains. Cells were cultured as shown

in **Supplementary Fig. 1**. Signals from the transformed lines were visualized using a TCS SP8

confocal laser-scanning microscope (Leica). Using the sequential setup of the SP8, roGFP2 signals

were collected at the emission wavelength of 510-550 nm immediately following excitation of

roGFP2 at 405 nm and 488 nm([29]). The degree of roGFP2 oxidation was analyzed as the ratio of

the emission signals after excitation at 405 nm and 488 nm([29]).

Photosynthesis-Irradiance curve

Photosynthesis-Irradiance curves were measured using a custom Pt-Ag/AgCl polarographic

electrode system (ALGI) with a water jacketed (for temperature-control), 1 mL glass reaction

chamber. The YSI 5331A electrodes (Yellow Springs Instruments) were polarized at −0.8 V. The

cultures used for the assays were concentrated to 2.5-5 μg mL^{-1} chlorophyll, supplemented with

15 µL of 0.5 M sodium bicarbonate in water and then purged with 1% CO_2/99% He. Using a gas

tight syringe, the sample was transferred into the reaction chamber that was also purged with 1%

CO_2/99% He. The rate of change in O_2 levels was measured sequentially at the light intensities

100, 200, 600, 1000 µmol photons m^{-2} s^{-1} (photosynthetic active radiation, PAR); each intensity

was maintained for 3-5 min followed by a 3-min intervening dark period, and then the light level

was raised to the next higher intensity (stepped change) until the full range of intensities was tested.

At 1,000 µmol photons m^{-2} s^{-1}, the light was held for 10 min followed by an 8-min dark period.

Before the measurement of each experimental series, the electrodes were calibrated with air (~21%

O_2) and 1% CO_2/99% He mixture (0% O_2). The initial slope of the response was used to determine

the O_2 evolution rate.

Starch and TAG quantification

Cells were grown under LL to mid-exponential phase. Cultures were then diluted to 0.5 µg/mL

chlorophyll with fresh medium and transferred to and then grown in ML for 24 h or more, as

indicated. Starch was measured using a modified method(75). In brief, cells were collected by

centrifugation and pigments extracted with methanol. Dried pellets were resuspended in water and

heated at 100°C to break the cells and release the starch. After cooling, amyloglucosidase and α-

amylase (2.25 U/mL) were used to hydrolyze the starch and the products of hydrolysis were

quantified using a Glucose Colorimetric Detection Kit (Thermo Fisher).

Intracellular triacylglycerol (TAG) levels were estimated by a fluorometric assay using the dye

Nile Red (Thermo Fisher)(76). A Nile Red solution (500 µg/mL in acetone) was added to 1 mL of

cell suspensions to a final concentration of 0.5 µg/mL. The samples were then incubated at room

temperature for 30 min, and the Nile Red fluorescence emission quantified at 575 nm after

excitation at 530 nm using a microplate reader (Infinite M1000; TECAN).

Metabolic analysis

Cells grown at LL were shifted to ML for 45 min or 6 h, as indicated. 45 mL of culture was rapidly

quenched in cold saline solution (-2 to -3°C), extracted using cold methanol and then analyzed by

LC-MS/MS. Quenching and analysis of metabolites were performed according to(77) with

modifications. The quenching solution, filtered saline (9 g/L NaCl), was prechilled to 4°C in a

refrigerator. 30 mL of the quenching solution was transferred to a 50 mL conical tube kept in an

ice bath mixed with salt to depress the temperature to between -3ºC and -1ºC. 15 mL of the culture

was rapidly plunged into the 30 mL of the quenching solution and the samples were centrifuged at

4,000 rpm for 10 min at 4ºC. Cell pellets were washed with fresh prechilled saline solution and

centrifuged again at 4,000 rpm for 10 min in a 2 mL centrifuge tube. For each replicate, 45 mL of

the culture was sampled and the cell pellets were pooled. Washed pellets were frozen at -80ºC

until extraction. Metabolite extraction and further analysis were modified from(*78*). A step-by-

step protocol is described in supplemental information (SI). All chemicals used for metabolite

extraction and LC-MS/MS analysis were Optima grade reagents.

References

1. F. Facchinelli, A. P. M. Weber, The metabolite transporters of the plastid envelope: an update. *Front. Plant Sci.* **2**, 50 (2011).

2. I. Haferkamp, S. Schmitz-Esser, The plant mitochondrial carrier family: functional and evolutionary aspects. *Front. Plant Sci.* **3**, 2 (2012).

3. Y. Zhao, H. Yu, J.-M. Zhou, S. M. Smith, J. Li, Malate Circulation: Linking Chloroplast Metabolism to Mitochondrial ROS. *Trends Plant Sci.* **25**, 446–454 (2020).

4. Y. Zhao, L. Luo, J. Xu, P. Xin, H. Guo, J. Wu, L. Bai, G. Wang, J. Chu, J. Zuo, H. Yu, X. Huang, J. Li, Malate transported from chloroplast to mitochondrion triggers production of ROS and PCD in *Arabidopsis thaliana. Cell Res.* **28**, 448–461 (2018).

5. R. Fliege, U.-I. Flügge, K. Werdan, H. W. Heldt, Specific transport of inorganic phosphate, 3-phosphoglycerate and triosephosphates across the inner membrane of the envelope in spinach chloroplasts. *Biochim. Biophys. Acta Bioenerg.* **502**, 232–247 (1978).

6. U. I. Flügge, K. Fischer, A. Gross, W. Sebald, F. Lottspeich, C. Eckerskorn, The triose phosphate-3-phosphoglycerate-phosphate translocator from spinach chloroplasts: nucleotide sequence of a full-length cDNA clone and import of the in vitro synthesized precursor protein into chloroplasts. *EMBO J.* **8**, 39–46 (1989).

7. Y. Lee, T. Nishizawa, M. Takemoto, K. Kumazaki, K. Yamashita, K. Hirata, A. Minoda, S. Nagatoishi, K. Tsumoto, R. Ishitani, O. Nureki, Crystal structure of the triose-phosphate/phosphate translocator in complex with inorganic phosphate (2017), , doi:10.2210/pdb5y78/pdb.

8. R. E. Häusler, N. H. Schlieben, B. Schulz, U. I. Flügge, Compensation of decreased triose phosphate/phosphate translocator activity by accelerated starch turnover and glucose transport in transgenic tobacco. *Planta.* **204**, 366–376 (1998).

9. R. G. Walters, D. G. Ibrahim, P. Horton, N. J. Kruger, A mutant of Arabidopsis lacking the triose-phosphate/phosphate translocator reveals metabolic regulation of starch breakdown in the light. *Plant Physiol.* **135**, 891–906 (2004).

10. A. Schneider, R. E. Häusler, Ü. Kolukisaoglu, R. Kunze, E. Van der Graaff, R. Schwacke, E. Catoni, M. Desimone, U. I. Flügge, An *Arabidopsis thaliana* knock-out mutant of the chloroplast triose phosphate/phosphate translocator is severely compromised only when starch synthesis, but not starch mobilisation is abolished. *Plant Journal* (2002), doi:10.1046/j.1365-313X.2002.01460.x.

11. J. W. Riesmeier, U. I. Flügge, B. Schulz, D. Heineke, H. W. Heldt, L. Willmitzer, W. B. Frommer, Antisense repression of the chloroplast triose phosphate translocator affects carbon partitioning in transgenic potato plants. *Proc. Natl. Acad. Sci. U. S. A.* **90**, 6160–6164 (1993).

12. L. Heinrichs, J. Schmitz, U.-I. Flügge, R. E. Häusler, The Mysterious Rescue of adg1-1/tpt-
2 - an Arabidopsis thaliana Double Mutant Impaired in Acclimation to High Light - by
Exogenously Supplied Sugars. *Front. Plant Sci.* **3**, 265 (2012).

13. D. Moog, A. Nozawa, Y. Tozawa, R. Kamikawa, Substrate specificity of plastid phosphate
transporters in a non-photosynthetic diatom and its implication in evolution of red alga-
derived complex plastids. *Sci. Rep.* **10**, 1167 (2020).

14. I. Haferkamp, P. Deschamps, M. Ast, W. Jeblick, U. Maier, S. Ball, H. E. Neuhaus, Molecular
and biochemical analysis of periplastidial starch metabolism in the cryptophyte *Guillardia*
theta. *Eukaryot. Cell*. **5**, 964–971 (2006).

15. M. Linka, A. Jamai, A. P. M. Weber, Functional characterization of the plastidic phosphate
translocator gene family from the thermo-acidophilic red alga *Galdieria sulphuraria* reveals
specific adaptations of primary carbon partitioning in green plants and red algae. *Plant*
Physiol. **148**, 1487–1496 (2008).

16. D. L. Jack, N. M. Yang, M. H. Saier Jr, The drug/metabolite transporter superfamily. *Eur. J.*
Biochem. **268**, 3620–3639 (2001).

17. S. Knappe, U.-I. Flügge, K. Fischer, Analysis of the plastidic phosphate translocator gene
family in Arabidopsis and identification of new phosphate translocator-homologous
transporters, classified by their putative substrate-binding site. *Plant Physiol.* **131**, 1178–1190
(2003).

18. A. P. M. Weber, M. Linka, D. Bhattacharya, Single, ancient origin of a plastid metabolite
translocator family in Plantae from an endomembrane-derived ancestor. *Eukaryot. Cell*. **5**,
609–612 (2006).

19. K. Fischer, B. Kammerer, M. Gutensohn, B. Arbinger, A. Weber, R. E. Hausler, U.-I. Flugge,
A new class of plastidic phosphate translocators: A putative link between primary and
secondary metabolism by the phosphoenolpyruvate/phosphate antiporter. *Plant Cell*. **9**, 453
(1997).

20. B. Kammerer, K. Fischer, B. Hilpert, S. Schubert, M. Gutensohn, A. Weber, U. I. Flügge,
Molecular characterization of a carbon transporter in plastids from heterotrophic tissues: the
glucose 6-phosphate/phosphate antiporter. *Plant Cell*. **10**, 105–117 (1998).

21. M. Eicks, V. Maurino, S. Knappe, U.-I. Flügge, K. Fischer, The plastidic pentose phosphate
translocator represents a link between the cytosolic and the plastidic pentose phosphate
pathways in plants. *Plant Physiol.* **128**, 512–522 (2002).

22. Y. Lee, T. Nishizawa, M. Takemoto, K. Kumazaki, K. Yamashita, K. Hirata, A. Minoda, S.
Nagatoishi, K. Tsumoto, R. Ishitani, O. Nureki, Structure of the triose-phosphate/phosphate
translocator reveals the basis of substrate specificity. *Nat Plants*. **3**, 825–832 (2017).

23. B. Loddenkötter, B. Kammerer, K. Fischer, U. I. Flügge, Expression of the functional mature
chloroplast triose phosphate translocator in yeast internal membranes and purification of the

histidine-tagged protein by a single metal-affinity chromatography step. *Proc. Natl. Acad. Sci.*
U. S. A. **90**, 2155–2159 (1993).

24. L. Lim, M. Linka, K. A. Mullin, A. P. M. Weber, G. I. McFadden, The carbon and energy
sources of the non-photosynthetic plastid in the malaria parasite. *FEBS Lett.* **584**, 549–554
(2010).

25. H. H. Stiehl, H. T. Witt, Quantitative Treatment of the Function of Plastoquinone in
Photosynthesis. *Zeitschrift für Naturforschung B*. **24**, 1588–1598 (1969).

26. J. M. Escoubas, M. Lomas, J. LaRoche, P. G. Falkowski, Light intensity regulation of cab
gene transcription is signaled by the redox state of the plastoquinone pool. *Proc. Natl. Acad.*
Sci. U. S. A. **92**, 10237–10241 (1995).

27. S. Saroussi, D. A. J. Karns, D. C. Thomas, C. Bloszies, O. Fiehn, M. C. Posewitz, A. R.
Grossman, Alternative outlets for sustaining photosynthetic electron transport during dark-
to-light transitions. *Proc. Natl. Acad. Sci. U. S. A.* **116**, 11518–11527 (2019).

28. B. Halliwell, Reactive species and antioxidants. Redox biology is a fundamental theme of
aerobic life. *Plant Physiol.* **141**, 312–322 (2006).

29. J. D. Vevea, D. M. Alessi Wolken, T. C. Swayne, A. B. White, L. A. Pon, Ratiometric
biosensors that measure mitochondrial redox state and ATP in living yeast cells. *J. Vis. Exp.*,
50633 (2013).

30. S. Dorion, J. C. Ouellet, J. Rivoal, Glutathione Metabolism in Plants under Stress: Beyond
Reactive Oxygen Species Detoxification. *Metabolites*. **11** (2021), p. 641.

31. C. H. Foyer, G. Noctor, Ascorbate and glutathione: the heart of the redox hub. *Plant Physiol.*
155, 2–18 (2011).

32. S.-T. Lin, C.-W. Chiou, Y.-L. Chu, Y. Hsiao, Y.-F. Tseng, Y.-C. Chen, H.-J. Chen, H.-Y.
Chang, T.-M. Lee, Enhanced Ascorbate Regeneration Via Dehydroascorbate Reductase
Confers Tolerance to Photo-Oxidative Stress in *Chlamydomonas reinhardtii*. *Plant Cell*
Physiol. **57**, 2104–2121 (2016).

33. A. Vidal-Meireles, J. Neupert, L. Zsigmond, L. Rosado-Souza, L. Kovács, V. Nagy, A.
Galambos, A. R. Fernie, R. Bock, S. Z. Tóth, Regulation of ascorbate biosynthesis in green
algae has evolved to enable rapid stress-induced response via the *VTC2* gene encoding GDP-
l-galactose phosphorylase. *New Phytol.* **214**, 668–681 (2017).

34. S. G. van Creveld, S. Rosenwasser, D. Schatz, I. Koren, A. Vardi, Early perturbation in
mitochondria redox homeostasis in response to environmental stress predicts cell fate in
diatoms. *ISME J.* **9**, 385–395 (2015).

35. S. Rosenwasser, I. Rot, E. Sollner, A. J. Meyer, Y. Smith, N. Leviatan, R. Fluhr, H. Friedman,
Organelles contribute differentially to reactive oxygen species-related events during extended
darkness. *Plant Physiol.* **156**, 185–201 (2011).

36.   H. Winter, S. C. Huber, Regulation of sucrose metabolism in higher plants: localization and
regulation of activity of key enzymes. *Crit. Rev. Biochem. Mol. Biol.* **35**, 253–289 (2000).

37.   S.-K. Lee, J.-S. Eom, L. M. Voll, C. M. Prasch, Y.-I. Park, T.-R. Hahn, S.-H. Ha, G. An, J.-
S. Jeon, Analysis of a triose phosphate/phosphate translocator-deficient mutant reveals a
limited capacity for starch synthesis in rice leaves. *Mol. Plant.* **7**, 1705–1708 (2014).

38.   U. Klein, Intracellular Carbon Partitioning in *Chlamydomonas reinhardtii*. *Plant Physiol.* **85**,
892–897 (1987).

39.   E. J. A. Hilgers, M. A. Schöttler, T. Mettler-Altmann, S. Krueger, P. Dörmann, M. Eicks, U.-
I. Flügge, R. E. Häusler, The Combined Loss of Triose Phosphate and Xylulose 5-
Phosphate/Phosphate Translocators Leads to Severe Growth Retardation and Impaired
Photosynthesis in *Arabidopsis thaliana* tpt/xpt Double Mutants. *Front. Plant Sci.* **9**, 1331
(2018).

40.   P. Joliot, G. N. Johnson, Regulation of cyclic and linear electron flow in higher plants. *Proc.*
Natl. Acad. Sci. U. S. A. **108**, 13317–13322 (2011).

41.   J. E. Johnson, J. A. Berry, The role of Cytochrome b_6f in the control of steady-state
photosynthesis: a conceptual and quantitative model. *Photosynth. Res.* **148**, 101–136 (2021).

42.   N. R. Boyle, N. Sengupta, J. A. Morgan, Metabolic flux analysis of heterotrophic growth in
Chlamydomonas reinhardtii. *PLoS One.* **12**, e0177292 (2017).

43.   S. Saroussi, E. Sanz-Luque, R. G. Kim, A. R. Grossman, Nutrient scavenging and energy
management: acclimation responses in nitrogen and sulfur deprived Chlamydomonas. *Curr.*
Opin. Plant Biol. **39**, 114–122 (2017).

44.   B. Usadel, O. E. Bläsing, Y. Gibon, K. Retzlaff, M. Höhne, M. Günther, M. Stitt, Global
transcript levels respond to small changes of the carbon status during progressive exhaustion
of carbohydrates in *Arabidopsis rosettes*. *Plant Physiol.* **146**, 1834–1861 (2008).

45.   O. E. Bläsing, Y. Gibon, M. Günther, M. Höhne, R. Morcuende, D. Osuna, O. Thimm, B.
Usadel, W.-R. Scheible, M. Stitt, Sugars and circadian regulation make major contributions
to the global regulation of diurnal gene expression in Arabidopsis. *Plant Cell.* **17**, 3257–3281
(2005).

46.   J. Preiss, in *The Biochemistry of Plants* (Elsevier, 1988; http://dx.doi.org/10.1016/b978-0-08-
092615-5.50012-6), pp. 181–254.

47.   Y. Meyer, W. Siala, T. Bashandy, C. Riondet, F. Vignols, J. P. Reichheld, Glutaredoxins and
thioredoxins in plants. *Biochim. Biophys. Acta.* **1783**, 589–600 (2008).

48.   J. M. Pérez-Ruiz, F. J. Cejudo, A proposed reaction mechanism for rice NADPH thioredoxin
reductase C, an enzyme with protein disulfide reductase activity. *FEBS Lett.* **583**, 1399–1402
(2009).

49. M. A. Ballicora, J. B. Frueauf, Y. Fu, P. Schürmann, J. Preiss, Activation of the potato tuber
ADP-glucose pyrophosphorylase by thioredoxin. *J. Biol. Chem.* **275**, 1315–1320 (2000).

50. A. Lepistö, E. Pakula, J. Toivola, A. Krieger-Liszkay, F. Vignols, E. Rintamäki, Deletion of
chloroplast NADPH-dependent thioredoxin reductase results in inability to regulate starch
synthesis and causes stunted growth under short-day photoperiods. *J. Exp. Bot.* **64**, 3843–
3854 (2013).

51. M. Schieber, N. S. Chandel, ROS function in redox signaling and oxidative stress. *Curr. Biol.*
24, R453-62 (2014).

52. C. H. Foyer, G. Noctor, Redox Homeostasis and Antioxidant Signaling: A Metabolic
Interface between Stress Perception and Physiological Responses. *Plant Cell*. **17**, 1866–1875
(2005).

53. D. Wagner, D. Przybyla, R. op den Camp, C. Kim, F. Landgraf, K. P. Lee, M. Würsch, C.
Laloi, M. Nater, E. Hideg, K. Apel, The Genetic Basis of Singlet Oxygen–Induced Stress
Responses of *Arabidopsis thaliana*. *Science* (2004), doi:10.1126/science.1103178.

54. P. Zimmermann, U. Zentgraf, The correlation between oxidative stress and leaf senescence
during plant development. *Cell. Mol. Biol. Lett.* **10**, 515–534 (2005).

55. G. Miller, V. Shulaev, R. Mittler, Reactive oxygen signaling and abiotic stress. *Physiol. Plant.*
133, 481–489 (2008).

56. J. Yu, Y. Li, Z. Qin, S. Guo, Y. Li, Y. Miao, C. Song, S. Chen, S. Dai, Plant chloroplast stress
response: Insights from thiol redox proteomics. *Antioxid. Redox Signal.* **33**, 35–57 (2020).

57. Y. Nishiyama, H. Yamamoto, S. I. Allakhverdiev, M. Inaba, A. Yokota, N. Murata, Oxidative
stress inhibits the repair of photodamage to the photosynthetic machinery. *EMBO J.* **20**,
5587–5594 (2001).

58. K. Kojima, M. Oshita, Y. Nanjo, K. Kasai, Y. Tozawa, H. Hayashi, Y. Nishiyama, Oxidation
of elongation factor G inhibits the synthesis of the D1 protein of photosystem II. *Mol.*
Microbiol. **65**, 936–947 (2007).

59. D. Takagi, S. Takumi, M. Hashiguchi, T. Sejima, C. Miyake, Superoxide and singlet oxygen
produced within the thylakoid membranes both cause photosystem I photoinhibition. *Plant*
Physiol. **171**, 1626–1634 (2016).

60. M. Exposito-Rodriguez, P. P. Laissue, G. Yvon-Durocher, N. Smirnoff, P. M. Mullineaux,
Photosynthesis-dependent H_2O_2 transfer from chloroplasts to nuclei provides a high-light
signalling mechanism. *Nat. Commun.* **8** (2017), doi:10.1038/s41467-017-00074-w.

61. C. H. Foyer, Reactive oxygen species, oxidative signaling and the regulation of
photosynthesis. *Environ. Exp. Bot.* **154**, 134–142 (2018).

62. S. Wakao, K. K. Niyogi, Chlamydomonas as a model for reactive oxygen species signaling
and thiol redox regulation in the green lineage. *Plant Physiol.* **187**, 687–698 (2021).

63. K. Shi, Z. Gao, T.-Q. Shi, P. Song, L.-J. Ren, H. Huang, X.-J. Ji, Reactive oxygen species-
mediated cellular stress response and lipid accumulation in oleaginous microorganisms: The
state of the art and future perspectives. *Front. Microbiol.* **8**, 793 (2017).

64. N. Chouhan, E. Devadasu, R. M. Yadav, R. Subramanyam, Autophagy induced accumulation
of lipids in pgrl1 and pgr5 of *Chlamydomonas reinhardtii* under high light. *Front. Plant Sci.*
12, 752634 (2021).

65. E. H. Harris, *Chlamydomonas sourcebook: Introduction to Chlamydomonas and its*
laboratory use (Academic Press, ed. 2, 2009), *Chlamydomonas Sourcebook*.

66. R. C. Edgar, MUSCLE: multiple sequence alignment with high accuracy and high throughput.
Nucleic Acids Res. **32**, 1792–1797 (2004).

67. S. Kumar, G. Stecher, K. Tamura, MEGA7: Molecular Evolutionary Genetics Analysis
version 7.0 for bigger datasets. *Mol. Biol. Evol.* **33**, 1870–1874 (2016).

68. R. D. Gietz, R. H. Schiestl, High-efficiency yeast transformation using the LiAc/SS carrier
DNA/PEG method. *Nat. Protoc.* **2**, 31–34 (2007).

69. Y. Kaye, W. Huang, S. Clowez, S. Saroussi, A. Idoine, E. Sanz-Luque, A. R. Grossman, The
mitochondrial alternative oxidase from *Chlamydomonas reinhardtii* enables survival in high
light. *J. Biol. Chem.* **294**, 1380–1395 (2019).

70. D. G. Gibson, L. Young, R.-Y. Chuang, J. C. Venter, C. A. Hutchison 3rd, H. O. Smith,
Enzymatic assembly of DNA molecules up to several hundred kilobases. *Nat. Methods*. **6**,
343–345 (2009).

71. J. Findinier, C. Delevoye, M. M. Cohen, The dynamin-like protein Fzl promotes thylakoid
fusion and resistance to light stress in *Chlamydomonas reinhardtii*. *PLoS Genet.* **15**,
e1008047 (2019).

72. S. Clowez, C. Renicke, J. R. Pringle, A. R. Grossman, Impact of Menthol on Growth and
Photosynthetic Function of *Breviolum Minutum* (Dinoflagellata, Dinophyceae,
Symbiodiniaceae) and Interactions with its Aiptasia Host. *J. Phycol.* **57**, 245–257 (2021).

73. F. Kong, A. Burlacot, Y. Liang, B. Légeret, S. Alseekh, Y. Brotman, A. R. Fernie, A. Krieger-
Liszkay, F. Beisson, G. Peltier, Y. Li-Beisson, Interorganelle Communication: Peroxisomal
MALATE DEHYDROGENASE2 Connects Lipid Catabolism to Photosynthesis through
Redox Coupling in Chlamydomonas. *Plant Cell*. **30**, 1824–1847 (2018).

74. P. Crozet, F. J. Navarro, F. Willmund, P. Mehrshahi, K. Bakowski, K. J. Lauersen, M.-E.
Pérez-Pérez, P. Auroy, A. Gorchs Rovira, S. Sauret-Gueto, J. Niemeyer, B. Spaniol, J. Theis,
R. Trösch, L.-D. Westrich, K. Vavitsas, T. Baier, W. Hübner, F. de Carpentier, M. Cassarini,
A. Danon, J. Henri, C. H. Marchand, M. de Mia, K. Sarkissian, D. C. Baulcombe, G. Peltier,

J.-L. Crespo, O. Kruse, P.-E. Jensen, M. Schroda, A. G. Smith, S. D. Lemaire, Birth of a Photosynthetic Chassis: A MoClo Toolkit Enabling Synthetic Biology in the Microalga *Chlamydomonas reinhardtii*. *ACS Synth. Biol.* **7**, 2074–2086 (2018).

75. U. Klein, A. Betz, Fermentative metabolism of hydrogen-evolving *Chlamydomonas moewusii*. *Plant Physiol.* **61**, 953–956 (1978).

76. E. T. Yu, F. J. Zendejas, P. D. Lane, S. Gaucher, B. A. Simmons, T. W. Lane, Triacylglycerol accumulation and profiling in the model diatoms *Thalassiosira pseudonana* and *Phaeodactylum tricornutum* (Baccilariophyceae) during starvation. *J. Appl. Phycol.* **21**, 669–681 (2009).

77. C. L. Sake, D. M. Newman, N. R. Boyle, Evaluation of quenching methods for metabolite recovery in photoautotrophic *Synechococcus sp.* PCC 7002. *Biotechnol. Prog.* **36**, e3015 (2020).

78. J. D. Young, A. A. Shastri, G. Stephanopoulos, J. A. Morgan, Mapping photoautotrophic metabolism with isotopically nonstationary (13)C flux analysis. *Metab. Eng.* **13**, 656–665 (2011).

79. X. Johnson, J. Alric, Central carbon metabolism and electron transport in *Chlamydomonas reinhardtii*: metabolic constraints for carbon partitioning between oil and starch. *Eukaryot. Cell.* **12**, 776–793 (2013).

80. C. L. Winder, W. B. Dunn, S. Schuler, D. Broadhurst, R. Jarvis, G. M. Stephens, R. Goodacre, Global metabolic profiling of *Escherichia coli* cultures: an evaluation of methods for quenching and extraction of intracellular metabolites. *Anal. Chem.* **80**, 2939–2948 (2008).

81. D. Strenkert, S. Schmollinger, S. D. Gallaher, P. A. Salomé, S. O. Purvine, C. D. Nicora, T. Mettler-Altmann, E. Soubeyrand, A. P. M. Weber, M. S. Lipton, G. J. Basset, S. S. Merchant, Multiomics resolution of molecular events during a day in the life of Chlamydomonas. *Proc. Natl. Acad. Sci. U. S. A.* **116**, 2374–2383 (2019).

82. J. M. Zones, I. K. Blaby, S. S. Merchant, J. G. Umen, High-Resolution Profiling of a Synchronized Diurnal Transcriptome from *Chlamydomonas reinhardtii* Reveals Continuous Cell and Metabolic Differentiation. *Plant Cell.* **27**, 2743–2769 (2015).

83. C. Y. Ngan, C.-H. Wong, C. Choi, Y. Yoshinaga, K. Louie, J. Jia, C. Chen, B. Bowen, H. Cheng, L. Leonelli, R. Kuo, R. Baran, J. G. García-Cerdán, A. Pratap, M. Wang, J. Lim, H. Tice, C. Daum, J. Xu, T. Northen, A. Visel, J. Bristow, K. K. Niyogi, C.-L. Wei, Lineage-specific chromatin signatures reveal a regulator of lipid metabolism in microalgae. *Nat Plants.* **1**, 15107 (2015).

84. D. González-Ballester, D. Casero, S. Cokus, M. Pellegrini, S. S. Merchant, A. R. Grossman, RNA-seq analysis of sulfur-deprived Chlamydomonas cells reveals aspects of acclimation critical for cell survival. *Plant Cell.* **22**, 2058–2084 (2010).

948 85. E. I. Urzica, A. Vieler, A. Hong-Hermesdorf, M. D. Page, D. Casero, S. D. Gallaher, J. Kropat,
949   M. Pellegrini, C. Benning, S. S. Merchant, Remodeling of membrane lipids in iron-starved
950   Chlamydomonas. *J. Biol. Chem.* **288**, 30246–30258 (2013).

Acknowledgments:

WH thanks M. Onishi for advice on generation of CRISPR mutants and A. Burlacot for help with PQ pool measurements. WH also would like to thank N. Linka, N. Boyle, A. Plett, and M. Meagher for the contributions to the in vitro test and metabolomic analyses.

Funding:

This project has received funding from DOE award DE-SC0019417. WH was supported solely by a grant from the US Department of Energy, Office of Science, Basic Energy Systems (DE-SC0019417). AK and MCP were supported solely by a grant from the US Department of Energy, Office of Science, Basic Energy Systems (DE-SC0019341). NB and MM were supported by US Department of Energy, Office of Science, Office of Biological and Environmental Research (BER), (DE-SC0018301). NL and AP were supported by German Research Foundations (DFG Grant LI1781/2-2). BR was supported by National Science Foundation (Award Number: 1645164). RGK and JF were supported by the Department of Plant Biology of the Carnegie Institution for Science. PR was supported by Human Frontier Science Program (HFSP) RGP0046/2018 (to ARG).

Author contributions:

WH, AK, ARG and MP conceptualized the study. WH generated the CRISPR mutants as well as the complemented strain, localized the TPT3 protein, analyzed the redox status in the cell, performed respiration measurements, measured the reduction level of PQ pool, analyzed changes of starch over the diel cycle. AK performed the experiments of photosynthetic O_2 evolution, extraction of samples for LC-MS/MS, data analysis of LC-MS/MS. AP and NL performed the reconstitution into liposomes and transport activity assays. MM and NB performed LC-MS/MS and analyses of metabolite data. BR performed data mining for transcriptome and the Nile red staining. JF and NF analyzed starch changes upon transition from LL to ML. WH, PR and ARG analyzed PSI and PSII activities. JF and BR constructed roGFP2 associated vectors. JX and WH performed the phylogenetic tree. WH drafted the manuscript. ARG, MP, AK, NL, NB, AP, MM and all the other authors helped with revising and finalizing the manuscript.

Competing interests:

All other authors declare they have no competing interests.

Data and materials availability:

All data are available in the main text or the supplementary materials.

 Figures and Tables

Fig. 1: Chloroplast TPT phylogenies, subcellular localization of CreTPT3 and in vitro measurements of CreTPT3 transport activities. (**A**) Truncated phylogenetic tree of potential chloroplast CreTPTs. (**B**) Cellular localization of CreTPT3. The construct for expression of the CreTPT3-VENUS fusion protein is shown above the cell images and described in the Materials and Methods. VENUS fluorescence, green; Chlorophyll (Chl) autofluorescence, magenta. (Scale bar: 5 μm). (**C**) Expression of Cre*TPT3* in yeast. Left, SDS-PAGE gel of proteins from total membranes isolated from yeast cells containing the empty vector (lane 1) or expressing His-CreTPT3 (lane 2). Right, corresponding immunoblot blot detection of recombinant His-CreTPT3 protein using anti-His antibodies. The calculated molecular mass of the N-terminal His-tagged protein was 37 kDa. (**D**) The kinetics of Pi exchange by His-CreTPT3 in lipid vesicles. Uptake of Pi (0.25 mM) into liposomes was measured in the presence (■) or absence (□) of Pi (30 mM) loaded within the vesicles. The arithmetic mean (±SD) of three independent experiments (each consisting of three technical replicates) was plotted with respect to the time after initiating the assay. (**E**) Analyses of substrate specificity of His-CreTPT3. Liposomes in which His-CreTPT3 was incorporated were preloaded with various substrates (30 mM) and the initial Pi uptake rates (at 0.25 mM external concentration) were determined. Relative velocities were compared to the Pi/Pi homo-exchange, which was set to 100%. The data represents the arithmetic mean (±SD) of three independent experiments (each consisting of three technical replicates). (**F**) Comparison of kinetic constants of CreTPT3. The Michaelis-Menten constant (K_M) for Pi uptake was determined using various external Pi concentrations (0.05-5 mM). The competitive inhibition constant (K_i) of Pi uptake (0.25 mM) was measured with increasing inhibitor concentrations (0.05-5 mM). Liposomes were preloaded with 30 mM Pi as the counter-exchange substrate. The data represent the arithmetic mean +/- SE of three independent experiments. ND, no competitive inhibitory constant could be measured under the given experimental conditions. DHAP, dihydroxyacetone phosphate; 3-PGA, 3-phosphoglycerate; PEP, phosphoenolpyruvate; Glc6P, glucose 6-P; Glc1P, glucose 1-P; Fru6P, fructose 6-P; Gal1P, galactose 1-P.

1013

Fig. 2: Growth characterization of *tpt3* mutants and complemented strains. (**A**) Growth of WT, *t3ko1*, *t3ko2* and the complemented strains (*C_T3KO1* and *C_T3KO2*, respectively), at various dilutions (see below) on agar plates incubated under LL and ML for 4 d. Growth of respective strains in liquid medium under LL (**B**) and ML (**C**). The data represents the arithmetic mean (±SD) of three independent experiments. Transition of the strains from LL to ML (**D**) or to high light (HL, 450 μmol photons m^{-2} s^{-1}) (**E**) for the times indicated. (**F**) Cell growth (number) and chlorophyll content of the indicated cultures at various times following exposure to ML (from LL), for up to 48 h. In (**A**) the cells were spotted on agar plates containing TAP or TP medium under continuous LL or ML conditions. The dilution series used was 1.5, 0.75, 0.375 μg/mL chlorophyll (left to right). For growth in liquid medium, the cells were cultured in TP medium to an initial OD$_{750nm}$ 0.02 in air and under continuous LL before initiating the various growth analyses in LL, ML or HL.

1026

1027

Fig. 3: Changes in storage carbon and metabolite levels in WT and *t3ko2* after transitioning the cells from growth in LL to ML. (A) Lugol staining of starch in WT and *t3ko2* after 48 h of ML. (**B**) Starch content in indicated strains following 0, 6, and 24 h of ML exposure. (**C**) Nile Red staining of TAG in WT and *t3ko2* following 48 h of ML exposure. (**D**) TAG content in the indicated strains following 0, 6, and 24 h of ML exposure. (**E**) Time course of metabolite accumulation at 0 h (LL), 45 min and 6 h following ML exposure. (**F**) Select metabolic pathways in Chlamydomonas adapted from Johnson and Alric, 2013(*79*). (**G**) Bar graph representation of fold-change for metabolites shown in (**E**), calculated by dividing the pool size in *t3ko2* by that of WT at the respective light levels. A fold change of 1 (no change) is shown by a dashed line. Each data point shows the mean and standard error; the data represents three biological replicates for each metabolite. An asterisk indicates statistically significant difference compared to WT (* P<0.05, ** P<0.01, *** P<0.001). Abbreviations: G6P, glucose-6-P; FBP, fructose bisphosphate; DHAP, dihydroxyacetone phosphate; 3-PGA, 3-phosphoglycerate; PEP, phosphoenolpyruvate; PYR, pyruvate; CIT, citrate; SUCC, succinate; FUM, fumarate; MAL, malate; E4P, Erythrose 4-P; R5P, Ribose 5-P; RU5P/X5P, ribulose 5-P/xylulose-5-P.

1043

Fig. 4: Photosynthetic activities followed LL to ML transition. (**A**) O_2 evolution of LL-acclimated cells in the indicated strains. (**B**) O_2 evolution of ML-acclimated (24 h) cells. (**C**) Fv/Fm of WT, *t3ko2* and the rescued strain (*C_T3KO2*) following 6 and 24 h in ML. (**D**) 1-qL values (redox state of PQ pool) of LL- and ML-acclimated WT, *t3ko2* and the rescued strain. The acclimated cells were exposed to 50 and 400 μmol photons m^{-2} s^{-1}, as indicated, for 2 mins. (**E-G**) PSI activities following a LL to ML transition. P700 oxidation and reduction kinetics in LL-grown cells (**E**) and after 6 (**F**) and 24 h (**G**) in ML. Absorbance differences were monitored at 705 nm during continuous illumination with 150 μmol photons·m^{-2} s^{-1} for 5 sec (white box above), followed by a saturating light pulse at 1,500 μmol photons·m^{-2} s^{-1} (arrow) and a 5 sec dark incubation (black box). The data represent the arithmetic mean (±SD) of three independent experiments. Kinetics were normalized by setting maximum oxidation (after light pulse) for WT to 100%. For panels **c** and **d**, the asterisks represent significant differences determined by ANOVA tests. ****P < 0.005, ****P < 0.0001.

1058

Fig. 5: Measurement of intracellular ROS. (A) ROS levels were evaluated by CM-H2DCFDA fluorescence of WT and the *t3ko2* mutant in LL and following exposure to ML for 48 h. **(B)** Quantitation of data in **a**. **(C-D)** Monitoring redox levels in chloroplasts **(C)** and mitochondria **(D)** after exposure of WT and the *t3ko2* mutant for 1.5 and 6.0 h of ML. **(E)** ROS levels monitored by CM-H2DCFDA staining in WT and *t3ko2* after the cells are transitioned from LL to darkness in TP medium. **(F)** Quantification of the data presented in **e**. Monitoring redox levels in chloroplasts **(G)** and mitochondria **(H)** in the dark with roGFP2 (targeted to chloroplast and mitochondria) in WT and the *t3ko2* mutant after various times in the dark. **(I)** Starch contents during diel cycle in indicated strains. The intensity of the light phase was 60 μmol photons m^{-2} s^{-1}. Values are means ± SD (n ≥ 3). The asterisks represent a significant difference determined by an ANOVA test. **P < 0.005, ***P < 0.001. Strains were grown photoautotrophically under LL for 16-24 h. After dilution with fresh medium, cultures were transferred to ML or dark for the times indicated.

1071

1072

Fig. 6: Impact of eliminating Chlamydomonas TPT3 on the chloroplast metabolic landscape.
i. reduced triose-P export leads to accumulation of the sugar-P in CBBC and glycolysis pathways (e.g. triose-P, hexose-P, marked in orange), and an elevated ratio of 3-PGA/Pi (which can allosterically activate AGPase activity) would begin to stimulate production of storage carbon; **ii.** CBBC activity diminishes as a consequence of the hyper-accumulation of the precursors along with the fixation of CO_2, leading to elevation of reductant (NADPH) and energy (ATP) within the chloroplast; the elevated NADPH:NADP$^+$ and ATP:ADP ratios, elicit strong feedback on PET causing hyper-reduction of electron carriers (eg: over reduced PQ pool and highly reduced PSI) that slows electron flow across the cytochrome b_6f complex; **iii.** the highly reduced PET system and the elevated NADPH can actively reduce FDX/TRX and the NTRC, respectively; reduced TRX and/or NTRC can activate the AGPase (above the activity elicited by an increase in the 3-PGA/Pi ratio), leading to additional starch accumulation; **iv.** hyper-reduction of PET can also lead to ROS accumulation, that can act as a signal controlling expression of chloroplast and nuclear genes; ROS also cause damage to both PSII and I and lead to neutral lipid (TAG: triacylglycerol) accumulation as a consequence of activation of diacylglycerol acyltransferases (DGAT) and phospholipid diacylglycerol acyltransferase (PDAT). Many of these metabolic/acclimatory processes both overlap and are interconnected. Under conditions of extended exposure to moderate light, the damage in the Chlamydomonas *tpt3* mutant can accumulate and lead to cell death; cell death occurs rapidly in high light.

1099 **Supplementary Materials**
1100

Materials and methods

Extraction of samples for LC-MS/MS

To each cell pellet, 500 µL methanol was added along with the internal standards ribitol and PIPES
to a final concentration of 2 µM each. Samples were then vortexed for 30 sec, frozen in liquid
nitrogen as described in(*80*), and allowed to thaw at 0°C. This vortex-freeze-thaw cycle was
repeated twice more before the samples were centrifuged at 10,000 x g at 1°C for 5 min. The
supernatant was collected, and the remaining pellets were extracted twice more with 500 µL of a
50:50 mixture of methanol and water, with 3 vortex-freeze-thaw cycles done for each extraction.
Supernatants from each extraction procedure were pooled and dried on a Thermo Fisher SpeedVac
Concentrator. Dried extracts were then resuspended in 500 µL water and cleaned to remove any
residual large cell debris by filtration, first through a 0.22 µm pore size Spin-X centrifugal tube
filter, followed by 10 K molecular mass cutoff filters, followed finally by 3 K molecular mass
cutoff filters. After each filtration step, filters were rinsed with 50 µL water, which was added to
the total sample volume for subsequent steps. Filtered extracts were dried again and resuspended
in 200 µL water for LC-MS/MS analysis.

Analysis of metabolite extracts

Metabolite extracts were analyzed using an LC-MS/MS method adapted from(*78*). A Phenomenex
150 mm x 2 mm Synergi Hydro-RP column was used on an Agilent 1200 Series HPLC system
coupled to an AB Sciex 5500 QTRAP system. LC was performed with a liquid injection volume
of 20 µL, using gradient elution of 10 mM tributylamine and 15 mM acetic acid (aqueous phase)
and acetonitrile (organic phase) at a constant flow rate of 0.3 mL/min, and a constant temperature
of 40°C. The gradient profile of the organic phase is as follows: 0% B (0 min), 8% B (10 min),
16% B (15 min), 30%B (16.5 min), 30% B (19 min), 90% B (21.5 min), 90% B (28 min), 0% B
(28.5 min), and 0% B (35 min). MS analysis was performed in negative mode using a multiple
reaction monitoring (MRM) acquisition method. Data acquisition was performed on the ABSciex
Analyst 1.7 software. Absolute quantification of intracellular metabolites was performed using the
quantitation mode on the Analyst software.

Protein extraction and immunoblot analysis

Affinity-purified polyclonal antibodies to CreTPT3 were custom-made by GenScript. The antigen
sequence used for antibody generation was KSWSFGRPVTKQEF. Chlamydomonas cells were
grown in liquid cultures to 2–5 x10^6 cells ml^{-1} and collected by centrifugation (2700 rpm, 5 min).
Cells were resuspended in resuspension buffer [(5 mM HEPES, pH 7.5, KOH), 10 mM EDTA (pH
7.5, NaOH), 1x protease inhibitor "complete EDTA-free" (Roche)]. For the disruption of cells,
bead-beating was performed using a mini beadbeater (Biospec) in two cycles of 30 sec min^{-1}, with
a 1 min period of cooling on ice between cycles. Disrupted cells were centrifuged at 4°C, 30 min,
14000 rpm in a microfuge. The supernatant was removed, the pellet was resuspended in sample
buffer (resuspension buffer supplemented with 100 mM Na$_2$CO$_3$, 100 mM DTT, 2% WT/vol SDS,
and 12% WT /vol sucrose).

SDS-PAGE was performed using a 12% polyacrylamide gel (Bio-Rad), with the electrophoresis
for 90 min at 120 V. Next, proteins were transferred onto a PVDF membrane (Bio-Rad) using
Trans-Blot Turbo (Bio-Rad). The detection was performed using Clarity Max Western ECL
Substrate (Bio-Rad).

RT-qPCR

RT-qPCR, was performed as described previously(*69*). Briefly, total RNA was isolated using the
RNeasy Plant Mini Kit (Qiagen) and treated with DNase I (Qiagen). First-strand cDNA was
generated by reverse transcription of 0.5 μg total RNA using the iScript cDNA Synthesis Kit (Bio-
Rad). Real-time PCR using the Roche Light Cycler 480 was performed with the SensiMix NO-
Rox SYBR Green I Kit as described by the manufacturer (Bioline). Oligos used in this research
for analyzing the expression levels of Cre*TPT1/2/3/25* are listed in **Supplemental Table 1**.

Analysis of transcript levels in response to different conditions

In order to analyze the transcript level of Cre*TPT* genes under different conditions, we obtained
RNAseq data published previously in which cells experienced a diurnal cycle, nitrogen starvation,
sulfur starvation and Fe^{2+} starvation(*81–85*). The log2 fold change of each gene was calculated
based on the ratio of transcript abundance at different time point over transcript level at time point
0 h. The fold change was visualized by a heatmap and the color bar is the scale of log2 (fold-
change).

In vivo polymeric carbohydrate staining

Lugol's iodine (aqueous solution of 1.8% iodine and 3.0% potassium iodide) was used to stain in
vivo polymeric carbohydrate. Briefly, 0.5 mL of the cell culture was pelleted, resuspended in 10
μL of Lugol's iodine and visualized under a bright-field microscope. Lugol's iodine selectively
binds to alpha-1,4 glucans and stains it blue-black.

1162

1163

1164

oligo name	sequence
gTPT3_pRam118_f	tactcacaacaagcccagttATGGCTTCGCTCGCTCAGCGCGTGAGC
gTPT3_pRam118_r	agcggggatccttatccgttTTAGGCCGCGGCCGGCGACGC
TPT3pro1000_f	gccttttgctcacatgtggcggccgccgccatgctggccaacag
TPT3pro1000_r	taccaatccacaaatgtgggctagcccatgtgaaggcgttgacc
TPT3_3UTR_f	gatgacgacaagtaaagatctTAAGGCGGGCAGCGGC
TPT3_3UTR_r	GCGCAAGAAAGAAGCTTGATATCCCCCTTATAATGTTTACAAGTGGTGCG
hygro_crispr_f2	GGAACTGGCGCAGTTCCTCTG
hygro_crispr_re2	GGCTCGCGTAGGAATCATCCG
TPT3_gRNA1_F	GCTCGGTTGAGCGCACCC
TPT3_gRNA1_R	CAAGGGTCTGGCGAGCACG
TPT3_gRNA2_F	ACCTCCCCATCGCCACGAT
TPT3_gRNA2_R	AGGGGTAAGGACAGCGGTG
Qrt_TPT1_f2	CCTGTACCAGCAGGTGTCGTAC
Qrt_TPT1_r2	CCAGCGCGTTCTGGAGTGACA
Qrt_TPT2_f2	ATGGCGTCTCTGCTGGGC
Qrt_TPT2_r2	CTGCTCGCCCATTCTCCTC
Qrt_TPT3_f2	GCAAAGCTGGCGCCTACTG
Qrt_TPT3_r2	TAGACCAGGCCGACCACCAC
Qrt_TPT25_fl	CTGGTTCATCGGGACATGGC
Qrt_TPT25_rl	TGGCTGAAGCTGACCACT

1165

1166 **Supplementary Table 1: Oligonucleotide list.**

Cre ID	Name	Prediction	CS Position
Cre08.g379350.t1.1	CreTPT1	cTP	CS pos: 61-62. TCL-AV.
Cre06.g263850.t1.1	CreTPT2	cTP	CS pos: 70-71. VCQ-AA.
Cre01.g045550.t1.2	CreTPT3	cTP	CS pos: 60-61. VTK-AS.
Cre02.g106200.t1.2	CreTPT4	OTHER	
Cre02.g112900.t1.2	CreTPT5	OTHER	
Cre02.g144300.t1.2	CreTPT6	OTHER	
Cre03.g162000.t1.2	CreTPT7	OTHER	
Cre03.g184850.t1.1	CreTPT8	OTHER	
Cre04.g227450.t1.2	CreTPT9	OTHER	
Cre07.g330850.t1.1	CreTPT10	OTHER	
Cre08.g363600.t1.2	CreTPT11	OTHER	
Cre09.g408400.t1.1	CreTPT12	OTHER	
Cre09.g413700.t1.1	CreTPT13	OTHER	
Cre08.g382350.t1.2	CreTPT14	OTHER	
Cre09.g415900.t1.2	CreTPT15	OTHER	
Cre10.g452750.t1.2	CreTPT16	OTHER	
Cre11.g479950.t1.2	CreTPT17	OTHER	
Cre12.g490050.t1.2	CreTPT18	OTHER	
Cre12.g490100.t1.2	CreTPT19	OTHER	
Cre12.g501000.t1.1	CreTPT20	OTHER	
Cre15.g641266.t1.1	CreTPT22	OTHER	
Cre15.g642950.t1.2	CreTPT23	OTHER	
Cre15.g643385.t1.1	CreTPT24	OTHER	
Cre16.g663800.t1.2	CreTPT25	cTP	CS pos: 70-71. IVA-SS.
Cre16.g666250.t1.2	CreTPT26	OTHER	
Cre17.g702700.t1.1	CreTPT27	OTHER	
Cre17.g703250.t1.2	CreTPT28	OTHER	
Cre17.g710850.t1.2	CreTPT29	OTHER	
Cre11.g467754.t1.1	CreTPT30	OTHER	
Cre14.g622700.t1.2	CreTPT31	OTHER	
Cre18.g748947.t1.1	CreTPT32	OTHER	
Cre09.g408428.t1.2	CreTPT33	OTHER	

1167

Supplementary Table 2: Members of the triose phosphate transporter family in *Chlamydomonas reinhardtii* and their predicted subcellular localization based on TargetP-2.0. cTP, chloroplast transit peptide; CS position, transit peptides cleavage site.

1171

Name	D_11h	D_9h	D_7h	D_5h	D_3h	D_1h	D_0.5h	D_0h	L_0.5h	L_1h	L_3h	L_5h	L_7h	L_9h	L_11h	D_13h
TPT1	5.2883	10.2702	9.45732	8.85549	7.70518	6.18466	7.5509	9.07019	0.511726	0.198621	0.329098	1.99425	2.93498	2.13736	1.02291	9.56813
TPT2	6.46543	9.45107	9.77461	10.5338	12.8513	8.72194	9.51834	10.275	59.1606	62.7836	33.5533	12.891	5.69013	5.94982	12.1866	3.6131
TPT3	233.805	337.793	247.287	181.496	131.602	97.1854	105.57	98.3637	71.9013	43.1935	218.026	280.947	185.973	179.459	157.828	312.135
TPT25	38.9867	62.7835	45.3561	32.7872	22.8554	14.8691	15.5851	13.8916	19.6257	26.8661	10.2213	7.48595	8.11458	19.7874	32.0979	53.4145

1172

Supplemental Table 3: Expression levels of TPTs over the diel cycle in Chlamydomonas. Data were extracted from(*81*).

1175
Supplementary Fig. 1: Experimental set up. All strains were grown photoautotrophically under
low white light (LL, ~30 μmol photons m^{-2} s^{-1}) for 16-24 h. After dilution with fresh medium, the
cultures were transferred to moderate light (ML, 250-300 μmol photon m^{-2} s^{-1}). On occasion, the
cultures were also transferred to the dark or high light (~450 μmol photon m^{-2}s^{-1}) for 24 or 48 h.
To avoid self-shading, the culture density was adjusted to 1-1.5 μg/mL chlorophyll before
exposure to ML or HL.

1182

1194

Supplementary Fig. 3: Phylogeny of TPT homologues in Chlamydomonas. Amino acid residues at the positions responsible for substrate specificity in TPT homologues are listed. The numbers above the amino acids are the position of each residue in CreTPT3. The residues marked with green, blue and purple colors are crucial for the preference of triose-phosphate or 3-PGA, for the specificity of PEP and for the binding of phosphate, respectively. Plastid PPT, XPT, and TPT from *Arabidopsis thaliana* are included in the analysis.

1201

1202

Supplementary Fig. 4: Background activity of Pi uptake derived from the endogenous yeast
transporter reconstituted into liposomes. Time-dependent uptake of Pi (0.25 mM) into
reconstituted liposomes preloaded with 30 mM of Pi (■) or without exchange substrate (●)
prepared from yeast cells harboring the empty vector (pYES-NTa). The arithmetic mean (±SD) of
three independent experiments (each with three technical replicates) was plotted against time. The
observed Pi uptake rates of endogenous yeast carrier reconstituted into liposomes were negligible
compared to the rates of His-CreTPT3 (20-fold higher, **Fig. 1D**), indicating that this expression
system is suitable for functional analysis of recombinant phosphate transporters.

1211

Supplementary Fig. 5: Inactivation of Cre*TPT3* by CRISPR-Cas9. (**A**) Molecular map of Cre*TPT3* gene and positions of the Cas9 targeted sites. The edited sites are located in exons 1 and 7; *t3ko1* has the marker gene inserted in exon 1 and *t3ko2* and *t3ko3* has the marker gene inserted in exon 7. (**B**) Immunoblot (left panel) and loading control (right panel) of wild type, *t3ko1*, *t3ko2*, *t3ko3* and the complemented strains (*C_T3KO1*, *C_T3KO2* and *C_T3KO3*, respectively).

1219

1220 **Supplementary**
1221 the drug resistant
1222 *tpt3* mutants (*t3k*

1223

1224

1225

Supplementary Fig. 7: Growth of the CRISPR-Cas9 generated Cre*tpt3* mutants on solid
medium. (**A**) The various mutant strains (*t3ko1*, *t3ko2*, *t3ko3*, *t3ko4*) and WT were spotted on
plates containing TP medium under continuous LL or (**B**) ML. (**C**) Cells were spotted on agar
plates containing TAP medium under continuous ML. The dilution series from (**A,B,C**) were 2.0,
0.75, 0.375 μg/mL chlorophyll. (**D**) Growth of wild type (WT), *t3ko1*, *t3ko2*, *t3ko3* and the
complemented strains on agar plates containing TAP or TP medium under LL or ML for 4 d. The
dilution series used was 1.5, 0.75, 0.375 μg/mL chlorophyll.

1233

a

b

1234

1235 **Supplementary Fig. 8: (A)** Cell size of WT, *t3ko2*, and *C_T3KO2* in TP under both LL and ML.
1236 **(B)** WT and *t3ko2* morphology following a LL to ML transition.

1237

1238

Supplementary Fig. 9: Quantification of chlorophyll contents in indicated strains after exposure to ML for 24 and 48 h. All strains were grown photoautotrophically under LL for 16-24 h. After dilution with fresh medium, the culture was transferred to ML for over 24 h. The initial inoculum density used for the transition was between 1-1.5 μg/mL chlorophyll in order to avoid self-shading.

1244

1245

1246
Supplementary Fig. 10: Starch and metabolites in WT and *t3ko2*. (**A**) Starch contents in
indicated strains following 0, 6, and 24 h of LL exposure in TP. (**B**) Time course of change in the
metabolite pool sizes at 0 h (LL), 45 min and 6 h after shifting LL-grown cells to ML. Data was
normalized to cell numbers. (**C**) Schematic representation of the fold-change for the metabolites
shown in (**B**), calculated by dividing the pool size in *t3ko2* by that of WT cells under the same
conditions. Fold change of 1 (no change) is shown by a dashed line. Each data point is the mean
and standard error of three biological replicates. Asterisk indicates statistically significant
differences relative to WT (* P<0.05, ** P<0.01, *** P<0.001). Abbreviations: G6P, glucose-6-
phosphate; FBP, fructose bisphosphate; DHAP, dihydroxyacetone phosphate; 3-PGA, 3-
phosphoglycerate; PEP, phosphoenolpyruvate; CIT, citrate; SUCC, succinate; FUM, fumarate;
MAL, malate; S7P, sedoheptulose-7-phosphate; E4P, Erythrose 4-phosphate; R5P, Ribose 5-
phosphate; RU5P/X5P, ribulose 5-phosphate/xylulose-5-phosphate;

1259

Supplementary Fig. 11: Quantum yield of PSII (ΦPSII) grown in LL (0 h: left panel) and after 6 and 24 h of growth in ML (right panel) following exposure to increasing actinic light intensities.

1262

Supplementary Fig. 12: Redox state of the PQ pool following a LL to ML transition at different actinic light intensities. 1-qL values of LL (left panel) and ML (right panel) acclimated cells exposed to increasing light intensities.

1268

1269

1270   **Supplementary Fig. 13:** Redox-sensitive GFP (roGFP) protein was targeted to the chloroplast (**A**)
1271   and mitochondrion (**B**). BF: bright field; roGFP2 fluorescence: green; Chlorophyll (Chl)
1272   autofluorescence: red.

1273

1274   **Supplementary Fig. 14:** (**A**) mRNA levels of the putative chloroplast envelope localized TPTs
1275   under LL during growth in TP medium. mRNA levels of CreTPT1 (**B**), CreTPT2 (**C**) and
1276   CreTPT25 (**D**) in the indicated strains 0, 2, 4 h after the cells were shifted from LL to ML.

1277

Supplementary Fig. 15: Expression of the putative chloroplast envelope localized TPTs under nitrogen starvation **(A)**, sulfur starvation **(B)**, and **(C)** iron starvation(*83–85*).

1280

1281