Title

Carbon partitioning and metabolic control by the TPT3 triose phosphate translocator of *Chlamydomonas*

reinhardtii

Weichao Huang[1]†*, Anagha Krishnan[2]†, Anastasija Plett[3], Michelle Meagher[4], Nicole Linka[3],

Yongsheng Wang[5], Bijie Ren[1], Justin Findinier[1], Petra Redekop[1], Neda Fakhimi[1], Rick G. Kim[1],

Devin A. Karns[2], Nanette Boyle[4], Matthew C. Posewitz[2], Arthur R. Grossman[1]*

[1]Department of Plant Biology, Carnegie Institution for Science, Stanford, CA 94305, USA

[2]Department of Chemistry, Colorado School of Mines, Golden, CO 80401, USA

[3]Institute of Plant Biochemistry, Heinrich-Heine University Düsseldorf, 40225 Düsseldorf,

Germany

[4]Department of Chemical and Biological Engineering, Colorado School of Mines, Golden, CO

80401, USA

[5]School of Life Sciences, Tsinghua University, 100084 Beijing, China

†Authors contributed equally.

*Corresponding Author: Weichao Huang; Address: The Carnegie Institution for Science,

260 Panama St. Bldg. 100; Stanford, CA 94305; email: whuang@carnegiescience.edu

*Corresponding Author: Arthur R. Grossman; Address: The Carnegie Institution for Science,

260 Panama St. Bldg. 100; Stanford, CA 94305; email: agrossman@carnegiescience.edu

 Running title

Triose phosphate/phosphate translocator: a dominant factor in regulating compartmentalized

metabolism and photosynthetic activity in Chlamydomonas

Abstract

Modulation of the rates of export of photoassimilates from the chloroplast is essential for

maintaining optimum photosynthetic rates. Additionally, the export of carbon metabolites from

the chloroplast can serve in the transduction of chloroplast signals to the nucleus in response to

environmental stresses. In this study we characterized a major chloroplast translocator of fixed

carbon, the triose phosphate/phosphate translocator 3 (CreTPT3), of the green alga

Chlamydomonas reinhardtii. This translocator is expressed at high levels, was shown to be on the

chloroplast envelope membrane and preferentially exchanges DHAP and 3-PGA for Pi based on

an in vitro yeast liposome uptake assay. Null mutants for CreTPT3, generated by CRISPR-Cas9,

exhibited a pleiotropic phenotype that impacted the growth, the rate of photosynthetic electron

transport, metabolite profiles, carbon partitioning, and organelle-specific accumulation of H_2O_2.

The severe phenotype of the *tpt3* mutant demonstrates that this transporter is a dominant conduit

on the chloroplast envelope for the transport of photosynthetically fixed carbon. In addition,

CreTPT3 can serve as a safety valve that moves excess reductant out of the chloroplast which helps

stabilize the redox state of the plastid. Finally, we demonstrated that CreTPT3 is essential for

preventing the cells from experiencing oxidative stress by sustaining a high rate of photosynthetic

electron transport as the cells acclimate to high light. Our studies reveal important features about

the localization and activity of CreTPT3, its importance in controlling metabolite partitioning, and

suggest differences in the export of photoassimilates from the chloroplasts of Chlamydomonas and

vascular plants.

Introduction

Photosynthetic organisms can absorb excess, potentially damaging levels of light energy during mid-day (when photosynthetic electron transport becomes saturated) and can suffer from extreme damage, especially when experiencing rapid fluctuations in light intensities or are subjected to nutrient limiting conditions and other environmental stresses that impair the productive utilization of excitation energy (Chaux et al., 2017; Saroussi et al., 2017). To cope with excess excitation energy, plants and algae have evolved mechanisms to dissipate this energy through nonphotochemical quenching (NPQ). The excess reductant and energy can also be eliminated by photochemical quenching. While the dominant form of photochemical quenching often involves the use of the electrons/reductant to fix inorganic carbon (Ci), which is used for growth or stored in the form of starch and lipids (Ge et al., 2014; Huang et al., 2018; Krishnan et al., 2015), redox equivalents can also be trafficked to other outlets where they are not used for anabolic processes. The main alternative photochemical electron outlets involve the reduction of O_2 to H_2O in H_2O-to-H_2O cycles that include a (*i*) Mehler-type reaction that is noncatalytic and functions on the acceptor side of PSI, (*ii*) flavodiiron protein reactions (Jokel et al., 2018) (FLVs, NADPH:flavin oxidoreductase) that catalytically reduces O_2 on the acceptor side of PSI without generating reactive oxygen species (ROS), (*iii*) the plastid terminal oxidase (Houille-Vernes et al., 2011) (PTOX, plastoquinol: oxygen oxidoreductase) reaction that can use electrons from the PQ pool to reduce O_2, and (*iv*) the movement of redox equivalents from the chloroplast to the mitochondrion where they can be used to reduce O_2 through various electron transport activities.

Mechanisms also exist in which fixed carbon and reducing equivalents can be shuttled between the chloroplast and cytoplasm. Studies with *Arabidopsis thaliana* revealed that the malate shuttle, which involves multiple malate dehydrogenases (MDH) and malate/OAA translocators (OMT), functions in the export of reductant from the chloroplast and enables plant cells to manage redox conditions in the chloroplast (Zhao et al., 2018, 2020). In addition to the malate shuttle, the triose phosphate (triose-P)/phosphate (Pi) translocator (TPT) has been proposed to be involved in moving fixed carbon out of the chloroplast, but can also act as a potential safety valve for eliminating excess reducing power from the chloroplast (Fliege et al., 1978; Flügge et al., 1989; Johnson and Alric, 2013; Lee et al., 2017a; Raghavendra and Padmasree, 2003; Stocking and Larson, 1969). The synthesis of triose-Ps during photosynthetic CO_2 fixation by the Calvin-Benson-Bassham Cycle (CBBC) is supported by the reducing power/energy (NADPH, ATP)

derived from photosynthetic electron transport. The TPTs reside on the inner chloroplast envelope

membrane and can transport triose-Ps (glyceraldehyde 3-phosphate (GAP), dihydroxyacetone

phosphate (DHAP)) and the three carbon acid 3-phosphoglycerate (3-PGA) in a counter exchange

for cytosolic Pi (Fliege et al., 1978; Flügge et al., 1989; Lee et al., 2017b).

During the last decades, physiological functions of the TPTs have been examined. Various plants

do not exhibit a strong phenotype if the chloroplast TPT is either eliminated or its level is reduced

(Häusler et al., 1998; Riesmeier et al., 1993; Schneider et al., 2002; Walters et al., 2004). A

reduction in TPT activity can be compensated for by diverting assimilated carbon into a transitory

starch pool that is subjected to accelerated turnover in the light and/or dark (Häusler et al., 1998;

Riesmeier et al., 1993; Walters et al., 2004), leading to accumulation of starch degradation

products that can be exported from the chloroplast and used in other cellular compartments.

Interestingly, a deficiency of the TPT in rice, which uses sucrose stored in leaves as its major

transitory form of fixed carbon rather than starch, led to severe phenotypic consequences; the

plants exhibited reduced photosynthetic rates and decreased levels of both starch and soluble

sugars relative to WT plants (Lee et al., 2014). In addition to the TPTs, plant chloroplasts possess

a maltose transporter (MEX1) and glucose translocator (GlcT), both of which are important for the

export of starch degradation products (Cho et al., 2011; Niittylä et al., 2004). Growth of

Arabidopsis thaliana mutants lacking the GlcT (*pglct-1* and *pglct-2*) was not significantly

impacted. In contrast, an Arabidopsis mutant defective for the maltose transporter (*mex1*)

displayed severe growth retardation (Niittylä et al., 2004). To further investigate the functions of

these sugar transporters in plant growth and development, double mutants of Arabidopsis lacking

MEX1 and either TPT or GlcT (Cho et al., 2011) were generated. The *pglct-1/mex1* and *tpt-2/mex1*

double mutants displayed strong growth inhibition, even relative to the *mex1* single mutant, with

the *pglct-1/mex1* strain exhibiting the most severe growth phenotype. In addition, *pglct-1/mex1*

displayed severely reduced photosynthetic activities and extreme aberrations in chloroplast

morphology. These studies demonstrated that in plants, the starch-mediated pathway for the export

of photoassimilate via GlcT and MEX1 is essential for Arabidopsis growth and development (Cho

et al., 2011).

Microalgae have high photosynthetic conversion efficiencies, can thrive in fresh to hypersaline

waters and are among the most metabolically versatile organisms known. They have also attracted

considerable interest worldwide because of their ability to synthesis large quantities of lipids (e.g.

for biofuels and food products), starch, pigments and other bioproduction, and can serve in the
remediation of waste water (Bhatt et al., 2022; Khan et al., 2018). Furthermore, the transporters
used for moving photoassimilate between the chloroplast and other cellular compartments and the
mechanisms and regulation of these transporters in microalgae have not been extensively explored.
Understanding the activities that comprise metabolite transport into and out of chloroplasts is
linked to an understanding of the metabolisms associated with the chloroplast, which can differ in
plants and algae; this is exemplified by the compartmentalization of the glycolytic and oxidative
pentose-phosphate pathways (OPPP). Plants appear to have duplicated the entire glycolytic and
OPPP pathways, with these pathways localized to both the chloroplast and cytosol (Plaxton, 1996).
In green microalgae such as the unicellular green alga *Chlamydomonas reinhardtii*
(Chlamydomonas throughout), the glycolytic pathway is not duplicated but is highly
compartmentalized, with the "upper half" of glycolysis (from glucose to GAP) exclusively in the
chloroplast and the "lower half" (from 3-PGA to pyruvate) exclusively in the cytosol (Klein, 1986).
Such metabolic differences suggest that metabolite exchange between the chloroplast and cytosol
is likely to differ between land plants and Chlamydomonas. Developing a more informed
understanding of central metabolism in microalgae can enable additional work on the
establishment, regulation and evolution of metabolic networks in algae and the ways in which
algae can be tailored for production purposes and for sustained growth under specific
environmental conditions.

One group of highly expressed translocators associated with movement of triose-P between the
chloroplast and cytosol are in the TPT family, a subgroup in the drug/metabolite transporter (DMT)
superfamily. There are still relatively few studies that demonstrate the functionalities of TPT
translocators in single-celled photosynthetic organisms (red algae, green algae, cryptophyte) based
on in vitro activities of these proteins in liposome transport assays (Haferkamp et al., 2006; Linka
et al., 2008; Moog et al., 2020). Active transport of triose phosphates has been noted multiple times
in Chlamydomonas (Boschetti and Schmid, 1998; Klöck and Kreuzberg, 1991), although, the
physiological fuctions associated with the various TPTs encoded on the genome of this alga have
not been characterized.

In this study, we used Chlamydomonas, the well-established model green algal system that has
been extensively used to analyze various physiological processes and, more specifically, to dissect
the function of its highly expressed TPT3 translocator. Chlamydomonas is predicted to contain

more than 30 TPT homologs; the informatic analyses of these proteins are still not refined enough to be certain of their substrate specificities and, in many cases, the transported metabolites are likely not triose-Ps. Here we present work that approaches the questions: *i*) what are the roles of chloroplast TPTs in partitioning fixed carbon and sustaining central carbon metabolism? *ii*) how do changes in the activities of these transporters, and in particular TPT3, impact photosynthetic electron transport (PET)? *iii*) what is the relationship between the transport of triose-P from chloroplasts and the production of ROS? Our analyses have been focused on Chlamydomonas TPT3 (CreTPT3) which, among the four predicted chloroplast associated TPT-like transporters, is the one most similar to plant chloroplast TPTs, and is highly expressed in the light. Through a series of in vitro and physiological analyses, we demonstrated that CreTPT3 is a major conduit on the chloroplast envelope for the trafficking of fixed carbon, sustaining central carbon metabolism, dissipating excess energy, enabling high rates of photosynthetic electron transport, preventing intracellular hydrogen peroxide (H_2O_2) accumulation, and balancing redox conditions at the subcellular level.

Results

Features of the triose-P/Pi translocator (DMT/TPT) family

Candidate genes encoding triose-P/Pi transporters (TPT) of Chlamydomonas were identified by blasting the Arabidopsis TPT protein (AT5G46110.1) against proteins encoded on the Chlamydomonas genome. The TPT family is the largest within the DMT superfamily in eukaryotes and includes triose-P and sugar-phosphate transporters associated with chloroplasts; many the family members are still not functionally characterized (Jack et al., 2001; Knappe et al., 2003; Weber et al., 2006). We have identified 32 genes encoding potential TPTs in the version v6.1 genome of Chlamydomonas, with four members, CreTPT10 (CreTPT1 in v5.6 genome), CrePPT2 (CreTPT2 in v5.6 genome), CreTPT3 and CGL51 (CreTPT25 in v5.6 genome), predicted to have a chloroplast-localized transit peptide (**Supplementary Table 3**).

Land plants harbor four classes of phosphate translocators on the inner plastid envelope that import and/or export sugars, including glucose 6-phosphate (GPT, transports glucose 6-phosphate), xylulose phosphate (XPT, transports xylulose 6-phosphate), phosphoenolpyruvate (PPT), and triose-P (various TPTs) (Eicks et al., 2002; Fischer et al., 1997; Flügge et al., 1989; Kammerer et al., 1998; Lee et al., 2017a). To explore the potential substrate specificities of the four putative

chloroplast-localized, Chlamydomonas TPT family transporters, their amino acid sequences were aligned with the plastidial phosphate transporters (PT) from Arabidopsis (AtPTs) (**Supplementary Fig. 2**). The ability of these transporters to use triose-P/DHAP or 3-PGA as substrate is dependent on five highly conserved residues (H184, K203, Y338, K359, and R360 in AtTPT1) (Lee et al., 2017a; Moog et al., 2020). Of the four putative chloroplast-targeted TPT family members in Chlamydomonas, only CreTPT3 and CGL51 contain all five of these residues (H170, K189, Y322, K345, and R346 in CreTPT3) (**Supplementary Fig. 2**). In AtTPT1, residue F262 (F248 in Chlamydomonas in the analogous protein CreTPT3) is thought to inhibit phosphoenolpyruvate (PEP) access to the binding site; this residue is replaced by N in AtPPT1 (PPT) (Lee et al., 2017a; Moog et al., 2020). CGL51 has M at position F248, indicating that unlike the other Chlamydomonas putative chloroplast sugar phosphate transporters, CGL51 might have greater preference for PEP or other substrates. These findings suggest that of the four chloroplast TPT family members, CreTPT3 most likely functions in specifically shuttling triose-P or 3-PGA across the chloroplast envelope.

TPT3 transport properties

To help establish the physiological (in vivo) function of CreTPT3, its deduced subcellular localization was experimentally determined. The CreTPT3 coding sequence with its putative targeting sequence was fused with VENUS (CreTPT3-VENUS) and expressed under the control of its own promoter in a Chlamydomonas mutant null for CreTPT3 (*t3ko2*, see below). As shown in **Fig. 1A**, CreTPT3-VENUS localized to the chloroplast envelope.

To evaluate the substrate specificity of CreTPT3, it was expressed in *Saccharomyces cerevisiae* (yeast) and the resulting recombinant protein was biochemically analyzed using a liposome uptake assay. The Cre*TPT3* gene was codon-optimized, expressed in yeast (**Fig. 1B**), and total cell membranes were isolated and reconstituted into liposomes (Linka et al., 2008; Loddenkötter et al., 1993). Chloroplast phosphate transporters in vascular plants can catalyze a Pi/Pi homo-exchange in vitro. The reconstituted CreTPT3 in liposomes was able to catalyze the signature Pi homo-exchange, whereas in the absence of a counter-exchange substrate, little Pi uptake was detected (**Fig. 1C**). In contrast, very low Pi uptake rates were observed for liposomes reconstituted with membranes from yeast cells lacking CreTPT3 (**Supplementary Fig. 3**), indicating that the detected Pi import activity was the result of CreTPT3 expression in yeast.

To assess the substrate specificity, the initial rates of Pi uptake into liposomes preloaded with
saturating concentrations (30 mM) of various potential counter-exchange substrates were
determined. As **Fig. 1D** shows, CreTPT3 exhibited the highest activity when DHAP was used as
substrate. The relative initial velocity for Pi uptake in exchange for DHAP was slightly higher than
that of 3-PGA (3-PGA exhibited a 75% exchange rate relative to that of Pi/DHAP) while the Pi
uptake into liposomes preloaded with PEP was much less efficient (about 16% relative to the
Pi/DHAP exchange rate) (**Fig. 1D**). Pi import was negligible when CreTPT3 liposomes were
preloaded with Glc-6-P, Glc-1-P, Fru-6-P, and Gal-1-P (**Fig. 1D**).

Characterizations of the K_M and K_i of CreTPT3 were performed to determine the affinity of the
transporter for the various substrates. CreTPT3 has an apparent Michaelis-Menten constant (K_M)
value of 1.1 +/- 0.2 mM for Pi (**Fig. 1E**), which is comparable to the value obtained for the vascular
plant TPT ortholog (Flügge et al., 1989). DHAP and 3-PGA competitively inhibited Pi uptake with
an inhibition constant (K_i) of 0.57 +/- 0.01 mM and 0.79 +/- 0.12 mM, respectively (**Fig. 1E**),
suggesting that CreTPT3 has a higher affinity for DHAP and 3-PGA than for Pi. In the case of
PEP, no inhibition of the Pi/Pi homo-exchange was observed, even at the non-physiologically high
concentration of 5 mM. Overall, the results of these in vitro assays indicate that CreTPT3 has a
typical plant-TPT substrate spectrum (Flügge et al., 1989). It can transport Pi, triose-P (DHAP)
and 3-PGA in a counter-exchange mode, however PEP might not be a physiologically relevant
substrate for CreTPT3, as was shown for apoplast TPT homologues in other organisms (Lim et al.,
2010; Moog et al., 2020).

Isolation of *tpt3* null mutants and their impact on cell growth

To explore the role of CreTPT3 in trafficking carbon and potentially reductant across the
chloroplast envelope in vivo, CRISPR knockouts of Cre*TPT3* were generated. We used the
CRISPR-Cas9 editing system to disrupt the Cre*TPT3* gene in exons 1 and 7 while at the same time
integrating the hygromycin marker gene (*AphVII*) into the edited site (**Fig. 2A**). Multiple
independent knockouts of Cre*TPT3* were obtained, including *t3ko1*, with the marker gene inserted
into exon 1, and *t3ko2* and *t3ko3*, with the marker gene inserted into exon 7; no CreTPT3 protein
was detected in any of the edited strains (**Fig. 2B, Supplementary Fig. 4**).

To elucidate the physiological roles of CreTPT3, we first examined growth of the parental WT
strain and the knockout mutants under either low light (LL, 30 μmol photons m^{-2} s^{-1}) or moderate

light (ML, 250-300 μmol photons m^{-2} s^{-1}). In contrast to WT cells, growth of all three of the CreTPT3 knockouts (*t3ko1*, *t3ko2*, *t3ko3*) was significantly reduced in LL on either photoautotrophic (TP) or heterotrophic (TAP) solid agar medium (**Fig. 2C**), while the growth of the mutant in ML was completely blocked on both TP and TAP (**Fig. 2C**). Growth curves were also determined for WT and mutant cells in liquid medium (TP) in both LL and ML (**Fig. 2D**); the results were in accord with those observed for the solid medium growth assays. Additionally, *tpt3* mutant cells in both LL and ML exhibited an increased cell diameter and formed clusters of cells that appear to be less able to separate following cell division (**Supplementary Fig. 5A**, **B**). To further confirm that the growth phenotypes are a consequence of the *tpt3* knockout, we introduced a wild-type copy of CreTPT3 fused to VENUS into the mutant strains (**Fig. 2B**). As shown in **Fig. 2C-F**, ectopic expression of the wild-type CreTPT3 in the *t3ko2* mutant (*C_T3KO2*) rescued the reduced growth phenotype of the mutant under all light conditions tested in this study (LL/ML/HL).

We also analyzed the chlorophyll content of photoautotrophically grown cells after transferring them from LL to ML (**Supplementary Fig. 1**) and found that *t3ko2* had reduced chlorophyll content relative to WT cells after 24 h in ML (**Fig. 2E-F** and **Supplementary Fig. 6**). Cell numbers and total chlorophyll were quantified following the LL to ML transition. The chlorophyll levels per cell declined in all strains at 24 and 48 h following transfer to ML; however, it was lower by more than half in *t3ko2* (0.34 μg/10^6 cells) relative to either WT cells or the *C_T3KO2* rescued strain (both ~0.72 μg/10^6 cells) after 24 h of ML (**Supplementary Fig. 6**), with some additional increase for WT and *C_T3KO2* after 48 h of ML (which might reflect some shading caused by an increase in the cell density). Finally, when mutant cells were transferred to HL for 24 h they became highly bleached (**Fig. 2E**).

Taken together, these results indicate that the activity of the CreTPT3 transporter is essential for growth over a range of light intensities (LL/ML/HL).

The *tpt3* mutant exhibits hyper-accumulation of 'storage' carbon

To explore the impact of the loss of CreTPT3 activity on carbon partitioning, we quantified carbon storage (starch and lipids) following a transition of WT and the *t3ko2* mutant from LL to ML. Lugol staining showed extensive starch accumulation in the *t3ko2* cells after a 48-h exposure to ML, whereas WT cells were barely stained (**Fig. 3B**). Furthermore, as shown in **Fig. 3A**, there is a ~55-fold difference in the level starch that accumulated in *t3ko2* relative to WT cells (13.29

compared to 0.24 µg starch/µg chlorophyll, respectively) after 24 h of illumination in ML. The

mutant also accumulated ~24-fold more lipid than WT and *C_T3KO2* cells (on a chlorophyll basis),

as monitored by Nile Red fluorescence, over the same time period (WT and *t3ko2*: 1038 and 41

Nile Red fluorescence/chlorophyll, respectively) (**Fig. 3C**, **D**). Since the chlorophyll in the mutant

was approximately 50% relative to that of WT cells after 24 h in ML, the accumulated starch and

lipid on a per cell basis would be ~25-fold more starch and ~12-fold more lipid in the mutant

relative to the WT strain. Furthermore, mutant cells are much bigger and tend to show more

aggregation than WT cells (**Supplementary Fig. 5A**, **B**), which may reflect the large increase of

storage carbon that they accumulate. These results suggest that the inability to transport triose-P

between the chloroplast and the cytosol through CreTPT3 resulted in a repartitioning of

photosynthetic assimilates (carbon, reductant, and ATP) toward the synthesis of both starch and

neutral lipid.

Cre*TPT3* deletion leads to accumulation of CBBC/glycolytic/gluconeogenic intermediates

To understand the metabolic consequences of the loss of CreTPT3 on growth under LL and ML,

comparative metabolite analyses of WT and the *t3ko2* mutant were performed on cells grown at

LL and after shifting them to ML for both 45 min and 6 h. Pool sizes of various central carbon

metabolites, particularly those of the CBBC/glycolysis/gluconeogenesis pathways and

intermediates of the TCA/glyoxylate cycle (**Fig. 3F**), were quantified. Data was normalized to both

chlorophyll content (**Fig. 3E**) and cell number (**Supplementary Fig. 7B**). The fold-change in the

quantity of each metabolite in the mutant relative to WT under LL or at 45 min and 6 h after the

switch to ML is given in **Fig. 3G** and **Supplementary Fig. 7C**.

Under LL (**Fig. 3E**, time 0 h) and following transfer to ML, the *t3ko2* mutant maintained a

significantly larger pool (2 to 6-fold) of glycolytic/gluconeogenic intermediates [DHAP, 3-PGA

and FBP (all three shared with CBBC), G6P, PEP], and of some metabolites of the CBBC pathway

(E4P, RU5P/X5P, R5P) compared to WT cells, suggesting that the loss of CreTPT3 results in a

back-up of these metabolites within the cell; our hypothesis is that they are accumulating inside

the chloroplast due to the reduced ability to export carbon in the *t3ko2* mutant. This is supported

by the finding of marked accumulation of starch and TAG in the mutant strain (**Fig. 3A-D**). While

the pool sizes of fumarate and malate (metabolites of the TCA/glyoxylate cycle) were similar in

both strains under LL, following the transition of the cells to ML, WT cells exhibited a significant

increase in the pool sizes of those metabolites while the mutant maintained a lower level, indicating

that the loss of CreTPT3 either directly (by supplying precursors) or indirectly (metabolic rewiring)

impacts their levels.

Thus, a primary function of CreTPT3 appears to be the export of photosynthetically-synthesized

reduced carbon, which would drive metabolic processes in the cytoplasm and other cellular

compartments, while importing Pi into the chloroplast to sustain ATP synthesis.

The *tpt3* mutant has reduced photosynthetic activity when grown in ML

To understand the impact of the loss of CreTPT3 activity on PET, we quantified photosynthetic

activities following a transition from LL to ML (**Supplementary Fig. 1**). Photosynthetic O_2

evolution rates (OERs) were measured for WT, *t3ko2* and *C_T3KO2* over a range of light

intensities. LL acclimated WT, *t3ko2*, and *C_T3KO2* showed comparable OER at intensities below

200 μmol photons m^{-2} s^{-1} while at saturating light intensities of $\geq$600 μmol photons m^{-2} s^{-1}, the

OER of *t3ko2* was diminished slightly relative to the WT and the complemented strain (**Fig. 4A**).

Upon acclimation of the cells to ML for 24 h, the *t3ko2* mutant consistently displayed lower OERs

than WT or *C_T3KO2* under all actinic light intensities used (**Fig. 4A**). Additionally, the Fv/Fm

in the mutant grown in LL was similar to that of WT and *C_T3KO2* (**Fig. 4B**). In contrast, after

exposure of the mutant cells to ML for 6 h, the Fv/Fm declined to about 50% of the WT and

C_T3KO2 values; the decline in the mutant continued over a period of 24 h in ML, with a 3-fold

decrease in Fv/Fm for *t3ko2* compared to that of WT (**Fig. 4B**). These results indicate that there is

damage of PSII reaction centers following exposure of the mutant to the ML exposure.

To examine the redox state of the photosynthetic apparatus, the pool of electron acceptors

downstream of PSII were evaluated. We quantified the photochemical efficiency (qL) of all strains

in LL and ML acclimated cells (**Fig. 4C**). The photosynthetic parameter qL indicates the redox

state of Q_A, the primary electron acceptor of the PSII reaction center. Assuming that Q_A and Q_B

are in equilibrium, qL would reflect the redox status of the PQ pool; a lower qL value indicates a

more reduced electron transport chain (Kramer et al., 2004). 1-qL positively correlates with the

PQ pool redox state. In LL-grown cells, 1-qL was significantly higher in *t3ko2* even at low level

illumination compared to that of WT and the rescued strain. For cells that had been acclimated to

ML (6 h and 24 h), 1-qL was 70% to near maximal for *t3ko2* even under low actinic light conditions

(e.g. 100 μmol photons $m^{-2}\,s^{-1}$), indicating a highly reduced PQ pool. In contrast, for WT and the

complemented strain, at the same light intensity (100 μmol photons $m^{-2}\,s^{-1}$), the 1-qL value attained

only ~20% of the maximum value (**Fig. 4C**). These results suggest that the PQ pool is more

reduced in the LL-acclimated *t3ko2* relative to the WT and *C_T3KO2* (**Fig. 4C**) and that there is a

much more pronounced reduction of this pool at all actinic light intensities in ML grown *t3ko2*.

These results suggest a limitation in electron flow downstream of the PQ pool in both LL and ML

grown *t3ko2*.

Moreover, we analyzed how fast PET becomes restricted following the transfer of LL-grown *t3ko2*,

C_T3KO2 and WT cells to HL (400-450 μmol photons $m^{-2}\,s^{-1}$) by monitoring changes in the redox

state of the PQ pool. We observed that the PQ pool in *t3ko2* began to be more reduced than that of

WT cells and *C_T3KO2* immediately after 105 s of elevated illumination (**Fig. 4D**), which

becomes much more pronounced at longer times in the HL. These results suggest that PET can be

rapidly limited by diminished triose-P export from the chloroplast; elimination of CreTPT3 has a

strong impact on PET activity.

To determine if PSI was also impacted in the mutant cells, we analyzed PSI/P700

oxidation/reduction kinetics following exposure of LL-grown cells to ML in the presence of

DCMU (20 μM) and hydroxylamine (1 mM) to block a contribution of electrons from PSII. Levels

of photo-oxidized P700 following exposure to actinic light for 5 sec were similar in all LL-

acclimated strains (**Fig. 4E**). However, the level of photo-oxidized P700 in ML-acclimated cells

declined in the mutant to ~40-50% of that in WT and *C_T3KO2* cells (**Fig. 4E**), indicating that

PSI is more reduced in *t3ko2* after ML exposure. As shown in **Fig. 4F**, the oxidation rate of P700

following a dark to light transition was much slower in the ML-acclimated *t3ko2* compared to that

of WT. These results suggest that the mutant has a diminished level of available electron acceptors

on the acceptor side of PSI (relative to WT and *C_T3KO2*) after growth in ML.

Inactivation of CreTPT3 dramatically affected cellular H_2O_2 accumulation and distribution

To investigate the function of CreTPT3 in reducing or preventing the generation of oxidative stress,

we assayed ROS production in the mutant using the fluorescent probe CM-H2DCFDA which,

upon exposure to increasing ROS levels, is converted to the green-fluorescent molecule

dichlorofluorescein (DCF). DCF fluorescence was visualized by confocal microscopy. As shown

in **Fig. 5A** and **B**, ROS levels in the *t3ko2* mutant were markedly increased (~3-fold) after 48 h in

ML, while no difference in the ROS level was detected in WT cells.

For an alternative, dynamic method for evaluating redox changes in the chloroplast, we used the

redox sensitive green fluorescent protein roGFP2, which was targeted to the chloroplast stroma

(**Fig. 5C**, upper panel) and mitochondria (**Fig. 5C**, lower panel). roGFP2 monitors ratiometric

redox changes of glutathione, which reflects cellular ROS levels (Dorion et al., 2021; Vevea et al.,

2013). LL-acclimated WT cells and the *t3ko2* mutant exhibited similar levels of chloroplast roGFP

oxidation (**Fig. 5D**). Upon transfer of these cells to ML, the mutant showed a much greater increase

in chloroplast oxidative conditions, with a 5.0-fold increase after 1.5 h, and a 2.0-fold increase

after 6 h, which is 1.8- and 1.6-fold higher than the values measured in WT cells (**Fig. 5D**). As

triose-P is exported to the cytosol by CreTPT3, it could potentially be further metabolized and

donate redox equivalents to the mitochondrial electron transport chain and alter mitochondrial

ROS production. Therefore, we measured mitochondrial redox levels at different light intensities

in both WT and the *t3ko2* mutant using the roGFP2 sensor targeted to the mitochondrion. Upon a

LL to ML transition, neither WT nor *t3ko2* displayed a significant change in fluorescence from the

mitochondrial targeted roGFP (**Fig. 5E**). These results suggest that the mitochondrial redox level

is maintained after shifting either LL acclimated WT or *t3ko2* cells to ML. Overall, the inability

to export triose-P through CreTPT3 markedly increased the level of oxidative stress in the

chloroplast but not in the mitochondrion.

We also determined if the ROS accumulated in *t3ko2* is H_2O_2 and whether this molecule shows

differential accumulation in the different subcellular compartments; the analysis was based on the

use of a hypersensitive sensor of H_2O_2, roGFP2-Tsa2ΔCR, which was previously used for studies

of Chlamydomonas (Niemeyer et al., 2021). In this study, we monitored real-time accumulation

of the H_2O_2 in the stroma, cytosol, mitochondrion and nucleus following a 20 min exposure either

to HL or very low light (**Fig. 6A**). The *t3ko2* stromal H_2O_2 level increased rapidly within 2.5 min

and attained a 1.4-fold increase after 20 min of illumination with HL compared to the initial level

in LL (**Fig. 6B**). The stromal H_2O_2 levels in WT cells showed little change after being shifted to

HL, however, the level declined when the cells were shifted to 10 μmol photons m^{-2} s^{-1}, or very

low light. Notably, after 20 min in HL, the stroma of the *t3ko2* mutant accumulated ~1.6-fold more

H_2O_2 than that of the WT cells; *t3ko2* mutant cells in very low light accumulated a similar amount

of stromal H_2O_2 as WT cells after being exposed to HL (**Fig. 6B**). The cytosolic probe also

responded rapidly, with a 1.2-fold increase for WT and a 1.4-fold increase for the *t3ko2* mutant

(**Fig. 6C**); note that the initial levels of H_2O_2 prior to the transfer of the cells to HL or very low

light were much lower in the WT cells than in the mutant. The *t3ko2* cytosolic H_2O_2 level was

elevated relative to the level in WT cells by 2.1-fold after 20 min of illumination in HL (**Fig. 6C**).

We did not observe a significant change in H_2O_2 levels in the mitochondrion for either the WT or

the *t3ko2*mutant after the cells were shifted to the higher light intensity (**Fig. 6D**). Finally, the

t3ko2 mutant already accumulated much higher levels of H_2O_2 in the nucleus in LL compared to

that in the WT cells, although both the mutant and WT did not show significant changes of H_2O_2

levels after HL exposure (**Fig. 6E**).

Discussion

CreTPT3 is critical for maintaining intracellular partitioning of fixed carbon

An Arabidopsis mutant for the chloroplast TPT was isolated 20 years ago (Schneider et al., 2002).

The growth of this mutant, which was designated *ape2*, was little affected under LL conditions

although its starch content was slightly elevated relative to that of WT Arabidopsis (Walters et al.,

2004). While *ape2* grew in HL, its growth was attenuated relative to WT Arabidopsis, although

there was no detectable change in starch accumulation (Walters et al., 2004). In potato, diminished

levels of chloroplast targeted TPT led to a 2-3-fold increase in starch accumulation; young

transgenic plants had a dwarf phenotype (Riesmeier et al., 1993). In comparison, the flux of fixed

carbon trafficked through CreTPT3 is absolutely critical for the growth of Chlamydomonas under

all light intensities tested, and it becomes essential for survival at the higher intensities (**Fig. 2**). In

addition to severe growth retardation, the *t3ko2* mutant exhibited a dramatic change in its carbon

partitioning. The starch content in *t3ko2* was markedly elevated in the mutant in both LL and ML

compared to WT cells, suggesting bottleneck in the export of fixed carbon in mutant cells, leading

to the storage of this carbon as starch and lipid (**Fig. 3A-D, Supplementary Fig. 7A**). In

Arabidopsis, elimination of the chloroplast targeted TPT1 protein did not significantly alter the

profiles of metabolites of either the CBBC or the glycolytic/gluconeogenic pathways (Hilgers et

al., 2018). In contrast, a Chlamydomonas mutant null for Cre*TPT3* exhibited a marked increase of

most metabolites associated with the CBBC and starch synthesis after exposure to either LL or

ML, indicating that triose-P exchange catalyzed by CreTPT3 was not readily compensated for by

alternative fixed carbon shuttles. CreTPT10, CrePPT2, CGL51, other translocators on the

chloroplast envelope that potentially use sugar-P molecules as substrates are unable to compensate

for the loss of CreTPT3 (**Supplementary Fig. 8B-D**).

In plants, the products of starch break-down can be routed out of chloroplasts by the MEX1 and

GlcT proteins. These translocators can compensate for the loss of TPT (Cho et al., 2011; Niittylä

et al., 2004; Walters et al., 2004). The Chlamydomonas chloroplast also likely contains MEX1,

which exhibited differential transport activities relative to that in vascular plants. Chlamydomonas

MEX1 can transport glucose and/or glucose phosphate across the envelope membrane, but there

is no strong evidence to show that it can export maltose as a *mex1* mutant did not accumulate

maltose or exhibit growth impairment (Findinier et al., 2017). Although the MEX1 transporter

cannot compensate for the loss of CreTPT3 in Chlamydomonas, we cannot rule out the possibility

that part of the accumulated starch in the Chlamydomonas *t3ko2* mutant can be mobilized in light,

with degradation products exported via MEX1.

The loss of chloroplast TPT function appears to have a more severe impact on Chlamydomonas

than on vascular plants. Land plants contain the entire glycolytic pathway in both the chloroplast

and cytosol while the pathway is partitioned between two compartments in Chlamydomonas; 90%

of the upper activities of the pathways(from F6P→ 3PG) was found to be associated with the

plastid while over 95% of the activities associated with the lower part of the pathway

(3PG→Pyruvate) appeared to be present in the cytosol (Klein, 1986; Rochaix et al., 1998)

Furthermore, for intact purified chloroplasts, almost 38% of the exported metabolites consist of 3-

PGA and DHAP (Rochaix et al., 1998). This compartmentalization likely requires CreTPT3,

which serves as the major carbon exporter from the chloroplast in Chlamydomonas in the light.

Investigation of the fluxes of specific carbon metabolites from the chloroplast under various

conditions in WT and the *tpt3* mutant is part of our current focus.

CreTPT3, a safety valve allowing the acclimation of PET to light stress

It was previously proposed that chloroplast TPTs could catalyze two potential reactions in the light

(**Fig. 7**) based on crystal structure analyses of TPT in the red algae (Lee et al., 2017a) and in vitro

assays using isolated spinach chloroplasts (Stocking and Larson, 1969): *i*) triose-P/Pi or *ii*) triose-

P/3-PGA exchange across the chloroplast inner envelope membrane. The former reaction can route

both carbon skeletons and reductants into the cytoplasm while importing Pi back into the

chloroplast for ATP regeneration. The latter reaction can import 3-PGA into the chloroplast in

exchange for triose-P (DHAP, GAP), which can serve as a substrate for the CBBC and allow for

ribulose 1,5-bisphosphate regeneration when it becomes depleted. The latter reaction also results

in the shuttling reductant (NAD(P)H) to the cytosol with zero net-carbon export. Indeed, in vitro,

TPT3 can actively transport both triose-P and 3-PGA in exchange for Pi (**Fig. 1**), suggesting that

TPT3 can potentially catalyze these two reactions and serve as a 'reductant shuttle' in addition to

a carbon shuttle. This exchange would eliminate excess reductant from the chloroplast, which

would help sustain photosynthetic electron flow.

Studies of photosynthetic activities of WT and *t3ko2* in both LL and ML support the idea that the

mutant is aberrant in its export of fixed carbon and reductant from the chloroplast. For LL-

acclimated *t3ko2* cells, the PQ pool (**Fig. 4C**) is much more reduced than in WT cells. However,

there is no loss of functions of either PSII or PSI activities (**Fig. 4A, B, E**) with the redox level and

oxidation rate of PSI comparable to that of the WT (**Fig. 4E, F**), suggesting that there is a reduced

rate of oxidation of the PQ pool. During ML exposure, the mutant stopped growing and, if placed

in HL, experienced severe bleaching (**Fig. 2E**). Furthermore, the highly reduced PQ pool and PSI

in ML-acclimated *t3ko2* (**Fig. 4C, E**) reflects a highly reduced PET system and the generation of

'over-flow' electrons. The slower oxidation rate of PSI in ML-acclimated *t3ko2* cells following

the dark to light transition during P700 measurement (**Fig. 4F**) along with the intracellular

accumulation of triose-P and 3-PGA indicates a block on the acceptor side of PSI. A defect in

routing carbon/reductants out of the chloroplast through CreTPT3 would cause hyper-

accumulation of reductant in the chloroplast, which would potentially result in a backup of PET.

The excess absorbed light energy by photosynthetic reaction centers can be dissipated either by

the export of fixed carbon (reaction: triose-P/Pi), which is coupled with reductant, or by shuttling

reductant out of chloroplasts that results in zero-net carbon export (reaction: malate/oxaloacetate

or triose-P/3-PGA exchange) (**Fig. 7**).

The malate-OAA shuttle represents another route that, under high redox stress, might have

partially compensated for the loss of the CreTPT3 by shuttling reductant from the chloroplast.

Intriguingly, malate levels in *t3ko2* were 4-fold lower than in WT cells (**Fig. 3E**, **G**). Moreover,

the expression of the plastidial malate dehydrogenases (Cre*MDH1* and Cre*MDH5*) were 3- to 6-

fold lower than that of the WT cells following a transition from LL to ML (**Supplementary Fig.**

8E-F), suggesting that the malate-OAA shuttle cannot compensate for a loss of CreTPT3 activity

in Chlamydomonas. Inactivation of triose-P/reductant export through CreTPT3 might have a

negative impact on the malate-OAA shuttle through metabolic rewiring. A previous study of

metabolic flux analysis during heterotrophic growth of Chlamydomonas showed that the TPT

shuttle is almost 10-fold more active than the malate-OAA shuttle (Boyle et al., 2017).

tpt3 **mutant experiences oxidative stress**

In plants, the export of sugars and other molecules (redox/ROS, specific sugar levels) can serve as

signals that coordinate chloroplast and nuclear gene expression during acclimation to HL (Häusler

et al., 2014; Zirngibl et al., 2022). In our study, we probed the impact of impaired triose-P export

(*tpt3* mutant, *t3ko2*) on ROS production and accumulation in the different subcellular

compartments following exposure of the cells to ML or HL. In plant cells, H_2O_2 is produced as a

side-product of cellular processes including PET, mitochondrial respiration, and substrate level

oxidation (Cheeseman; Foyer and Noctor, 2016). While ROS stability is generally low, it can

accumulate in cells experiencing oxidative stress, with H_2O_2 being the most prevalent species that

functions as an important redox messenger (Li and Kim, 2022). Moreover, the trafficking of H_2O_2

into or out of different cellular compartments can trigger activation of other retrograde and

anterograde signals that may coordinate the activities among the compartments, including the

nucleus (Exposito-Rodriguez et al., 2017; Mittler et al., 2022; Shapiguzov et al., 2012). Upon

exposure to HL, the *t3ko2* mutant in Chlamydomonas accumulated more stromal H_2O_2 which

could diffuse rapidly out of the chloroplast and to the cytoplasm (**Fig. 6B, C**). Both stromal and

cytosolic H_2O_2 levels exhibited a similar increase, although it is still uncertain whether or not this

increase is solely dependent on H_2O_2 diffusion from the chloroplast stroma. It is unlikely that

mitochondria contribute to the increase in H_2O_2 in the *t3ko2* since exposing the mutant to elevated

light levels did not lead to accumulation of intramitochondrial H_2O_2 (**Fig. 6D**). However, an

increase in cytosolic H_2O_2 in the *t3ko2* mutant when exposed to either high or low light intensities

appears to influence H_2O_2 accumulation in the nucleus (**Fig. 6E**); the level of H_2O_2 in the cytoplasm

or nucleus of WT cells was not strongly impacted by the light conditions used in the experiments.

Such observations are consistent with previous reports demonstrating that the nuclear H_2O_2 level

was not significantly increased in WT Chlamydomonas cells exposed to HL (Niemeyer et al.,

2021), which suggest that the cytosol or the other organelles, such as the peroxisomes, might act

as a barrier that limits the intracellular diffusion of H_2O_2 into the nucleus. H_2O_2 transfer from the

chloroplast to the cytosol in the *t3ko2* mutant can potentially trigger retrograde signals that

modulate nuclear gene expressions, which in turn could ameliorate some of the negative effects of

ROS and elicit repair of the photosynthetic apparatus. A similar response may be elicited in WT

cells if the intensity of actinic light is increased to higher levels.

It has been shown that H_2O_2 can be synthesized in thylakoid membranes through the oxidation of

plastoquinol (PQH_2), suggesting a positive correlation between the redox state of the PQ pool and

the generation of H_2O_2 (Khorobrykh et al., 2015). This has become evident in both *Nicotiana*

benthamiana and Chlamydomonas based on the use of different H_2O_2 hypersensitive sensors

(Exposito-Rodriguez et al., 2017; Niemeyer et al., 2021). These two organisms have also been

shown to accumulate more stromal H_2O_2 in HL, and that this HL response is dependent on

photosynthesis. We observed this positive correlation between PQ pool reduction and the

generation of H_2O_2 in the *t3ko2* mutant. A more reduced PQ pool was observed in *t3ko2* relative

to the WT cells after a HL exposure of 105 s (**Fig. 4D)**. In parallel, there was a marked increase in

the stromal H_2O_2 level following 2.5 min of HL exposure (**Fig. 6B**). Therefore, hyper-reduction of

the PQ pool in the mutant likely results in elevated production of stromal H_2O_2, suggesting that

CreTPT3 can diminish H_2O_2 synthesis/accumulation and sustain a high rate of PET during

acclimation of the cells to HL; the protection by Cre*TPT3* likely reflects its ability to export fixed

carbon, relieving the redox pressure and formation of ROS in the chloroplast. Cre*TPT3* is the most

responsive/upregulated of the four Chlamydomonas, chloroplast localized *TPT* family genes

following exposure of the cells to nitrogen, sulfur or iron deprivation (**Supplementary Fig. 9 A-**

C), which reinforces the hypothesis that the ability to traffic fixed carbon from the chloroplast is

important not only for distributing carbon to other compartments of the cell, but potentially also

for relieving oxidative stress in the organelle. The isolated *tpt3* mutant represent an important

resource for investigating the dynamics of H_2O_2 synthesis and accumulation, as well as its role in

eliciting transcriptional and post-transcriptional control of gene expression.

As depicted in **Fig. 7**, we propose that various tiers of redox regulation are potentially responsible

for the differential physiological responses of Chlamydomonas *tpt3* knockout mutants. When

t3ko2 cells are initially transferred from LL to ML, the triose-P pool and metabolites derived from

that pool accumulate in the mutant because of its reduced capacity to move triose-P out of the

chloroplast, which also causes reduced CBBC activity, hyper-reduction of PET and accumulation

of ROS (which would inhibit photosynthetic activity). The hyper-reduction of PET and

accumulation of carbon metabolites in the chloroplast of the mutants would activate AGPase

through allosteric regulation and by the Fdx/Trx (ferredoxin/thioredoxin) and NTRC (NADPH-

dependent thioredoxin reductase C) redox systems, which would result in starch hyper-

accumulation (Ballicora et al., 2000; Lepistö et al., 2013) (**Fig. 7**). Furthermore, increased ROS

accumulation in chloroplasts and the elevated redox state of PET would elicit the generation of

retrograde signals that mediate changes in expression of nuclear genes (Shapiguzov et al., 2012;

Suzuki et al., 2012; Wakao and Niyogi, 2021) and stimulate the synthesis of specific activities that

function to ameliorate the impact of the hyper-reduced state of the chloroplast resulting from a

compromised ability to export reductant and fixed carbon from the organelle (**Fig. 7**).

Material and methods

Strains and culture conditions

WT Chlamydomonas M10 (CC-4403, isogenic line derived from CC-124) was used as the parental strain for the generation of knockout mutants. Cultures were routinely cultivated in growth chambers (LED-41L2, Percival Scientific, Inc.) at 25°C with continuous shaking on an orbital shaker (VWR OS-500 Shaker) at 120 rpm, in Tris-Acetate-Phosphate (TAP) medium (Harris, 2009). Cultures were illuminated with continuous cool white LEDs (LED-41L2, Percival Scientific, Inc.) at low light (LL, 30 μmol photons m^{-2} s^{-1}). Experiments were mostly performed with cells grown in the photoautotrophic medium TP (which is TAP medium without acetate), and in TAP medium in some cases, at 25°C, and sparged with air while being shaken at 120 rpm in a growth chamber (LED-41L2, Percival Scientific, Inc.). For growth assays, cultures were inoculated to a density of 0.02 at $OD_{750\ nm}$ (~1×10^5 cells/mL) in TP sparged with air under either LL or moderate light (ML, 250-300 μmol photons m^{-2} s^{-1}) intensities. For spectrophotometric and chlorophyll fluorescence analyses, the experimental design is described in **Supplementary Fig. 1**. Growth assays on solid medium were performed with cultures spotted onto the medium at different dilutions (as indicated in the text) and exposed to different light intensities; spot tests for photoautotrophic growth were on TP agar plates and for mixotrophic growth on TAP agar plates. Agar plates were incubated for 7 d under either LL or ML (cool white LED) at 25°C.

Reconstitution into liposomes and transport assays

The procedures for the construction and expression of CreTPT3 in yeast are described in supplemental information (SI). For uptake studies, yeast membranes from cells with and without recombinant His-tagged CreTPT3 were enriched and reconstituted into 3% (w/v) L-alpha-phosphatidylcholine using a freeze-thaw-sonication procedure (Linka et al., 2008). The reconstituted liposomes were preloaded with 30 mM Pi or phosphorylated metabolites to be tested as potential transport substrates. As a negative control for antiport activity, liposomes were also generated without metabolite preloading. The external counter-exchange substrate was removed via gel filtration on Sephadex G-25M columns (GE Healthcare). Transport assays were initiated by adding 0.25 mM [alpha-^{32}P]-phosphoric acid (6,000 Ci/mmol) to the medium bathing the liposomes and performed as previously described by Linka et al. (2008). Measurements of Km for Pi and competitive inhibition constants (Ki) are described in supplemental information (SI)

Vector construction, transformation and subcellular localization

The pRam118_VENUS plasmid, which harbors the VENUS gene and the *AphVII* cassette (confers resistance to hygromycin) (Cre01.g045550)(Kaye et al., 2019), was used to express Cre*TPT3* (Cre01.g045550). The step-by-step description is in supplemental information (SI). Following construction, 2-4 µg of the plasmid was linearized and added to 250 µL of a cell suspension at a concentration of ~3×10^8 cells/mL per reaction. The GeneArt MAX Efficiency Transformation Reagent for algae (Invitrogen) was used for introducing the plasmid into the algal cells by electroporation according to the instructions provided by the manufacturer. Transformants were selected on solid TAP medium containing hygromycin (10 µg/mL; Enzo Life).

Drug resistant transformants were visualized for VENUS fluorescence as previously described (Kaye et al., 2019). In brief, transgenic cell lines resistant to hygromycin were screened for VENUS fluorescence using a microplate reader (Infinite M1000; TECAN). Excitation and emission settings were as follows: VENUS, excitation 515nm, bandwidth 12 nm and emission 550 nm, bandwidth 12 nm; chlorophyll excitation 440 nm, bandwidth 9 nm and emission 680 nm, bandwidth 20 nm. The TCS SP8 confocal laser-scanning microscope (Leica) was used to visualize the VENUS fluorescence signal (Kaye et al., 2019).

CRISPR-Cas9 mediated mutagenesis

Chlamydomonas reinhardtii WT strain CC-124 was used for mutant generation. WT cells were cultured under continuous light (50 µmol photons m^{-2} s^{-1}) for 2 d to a density of $3\text{-}5\times10^6$ cells/mL. The cells were then concentrated to 2×10^8 cells/mL in 0.5×TAP supplemented with 80 mM sucrose. Two single guide RNAs (sgRNA) were designed by CHOPCHOP and synthesized by Integrated DNA Technologies (IDT). The sequences of the generated sgRNA are: *TPT3*-sg1 (5′- CGCUGGGCGUCACUUCCCGGCGG-3′) for editing exon 1 and *TPT3*-sg2 (5′- AAGGCCGCUAUCGCCAACGUGGG-3′) for editing of exon 7. The protocol for disruption of Cre*TPT3* (Cre01.g045550) was adapted from (Findinier et al., 2019) and is described in (SI).

Complementation of mutants *tpt3* mutants

Mutant strains were transformed with the linearized plasmid pRam118_CreTPT3 plasmid. Transformed cells were selected in ML and then screened for VENUS fluorescence. Colonies containing VENUS and an *AphVII* cassette knock-in at the Cas9 targeted site were examined for

accumulation of the CreTPT3 protein by immunoblots using CreTPT3 antibody generated by

GenScript USA Inc (Piscataway, USA). Colonies of immuno-positive cells were subjected to

growth complementation assays using spot tests under ML on either solid TAP or TP medium.

P700 activity measurements

Absorbance spectroscopy to measure P700 activity was performed in dark-adapted liquid cultures

(15 µg/mL chlorophyll, in 20 mM HEPES-KOH, pH 7.2, and 10% ficoll) using a JTS-100

spectrophotometer (SpectroLogiX, TN) as previously described (Clowez et al., 2021). Actinic light

was provided by an orange LED (165 µmol photons m^{-2} s^{-1}) for PSI oxidation, followed by a

saturating pulse and dark incubation. 20 µM DCMU and 1 mM hydroxylamine were added to the

cell suspension to inhibit linear electron flow (LEF) prior to the measurement. P700 activity was

measured by monitoring the absorbance at 705 nm (interference filter 6 nm FWHM was used to

create a narrow excitation beam); the absorbance at 740 nm was used to correct for an unspecific

contribution to the 705 nm signal.

Chlorophyll fluorescence analysis

Chlorophyll fluorescence, used to evaluate photosynthetic electron transport, was measured with

a DUAL PAM-100 fluorometer. Cells were acclimated in the dark for 20 min prior to illumination

at increasing light intensities (0, 10, 50, 100, 200, 400, 800, 1000, 1200 µmol photons m^{-2} s^{-1}) for

2 min at each intensity, or at a constant intensity of 450 µmol photons m^{-2} s^{-1} for 10 min to evaluate

1-qL. 1 mM CO_2 ($NaHCO_3$) was added to the reaction mix as an electron acceptor for the CBBC.

ROS measurements and roGFP2 imaging analysis

All strains were grown photoautotrophically under LL for 16-24 h. After dilution with fresh

medium, the cultures were transferred to ML and then stained with CM-H2DCFDA (Thermo

Fisher Scientific) as described in (Kong et al., 2018) for detecting ROS. A step-by-step protocol

is given in (SI).

Constructs containing chloroplast or mitochondria targeting sequences (Crozet et al., 2018) fused

to codon-optimized roGFP2 (Vevea et al., 2013) were transformed into WT and mutant strains.

Cells were cultured as shown in **Supplementary Fig. 1**. Signals from the transformed lines were

visualized using a TCS SP8 confocal laser-scanning microscope (Leica). roGFP2 signals were

collected and analyzed as previously described (Vevea et al., 2013). Using the sequential setup of
the SP8, roGFP2 signals were collected at the emission wavelength of 510-550 nm immediately
following excitation at 405 nm and 488 nm (Vevea et al., 2013). The degree of roGFP2 oxidation
was analyzed as the ratio of the emission signals after excitation at 405 nm and 488 nm (Vevea et
al., 2013).

Plasmids harboring H_2O_2 roGFP2-Tsa2ΔCR sensors (Niemeyer et al., 2021) were obtained from
the Chlamydomonas Resource Center (https://www.chlamycollection.org/) (**Supplementary**
Table. 2). Cells in which roGFP2-Tsa2ΔCR was targeted to the stroma, cytosol, mitochondrial
matrix, or nucleus were initially grown in LL in TAP to exponential growth phase. Cells were then
grown in TP in LL for 2 h and shifted either to HL or very low light (10 μmol photons m^{-2} s^{-1}).
The oxidation state of the sensor was trapped by the addition of N-Ethylmaleimide (NEM) (Sigma
Aldrich) and roGFP2 fluorescence was measured in a plate reader (Infinite M1000; TECAN).
Signals were detected using the excitation wavelengths of 410 and 488 nm, and the emission
wavelength of 514 nm. The degree of sensor oxidation (OxD) was calculated as described in
(Niemeyer et al., 2021).

Photosynthesis-Irradiance curve

Photosynthesis-Irradiance curves were measured using a custom Pt-Ag/AgCl polarographic
electrode system (ALGI) with a water jacketed (for temperature-control), 1 mL glass reaction
chamber. A step-by-step protocol is provided in SI.

Starch and TAG quantification

Cells were grown under LL to mid-exponential phase, diluted to 0.5 μg/mL chlorophyll with fresh
medium and transferred to and then grown in ML for 24 h or more, as indicated. Starch was
measured according to (Klein and Betz, 1978), , with slight modification (and see SI).
Triacylglycerol (TAG) levels were estimated by a fluorometric assay using the dye Nile Red
(Thermo Fisher)(Yu et al., 2009). A Nile Red solution (500 μg/mL in acetone) was added to 1 mL
of the cell suspensions to a final concentration of 0.5 μg/mL. Samples were then incubated at room
temperature for 30 min, and the Nile Red fluorescence emission quantified at 575 nm following
excitation at 530 nm using a microplate reader (Infinite M1000; TECAN).

Metabolic analysis

Cells grown in LL were shifted to ML for 45 min or 6 h, as indicated. 45 mL of culture was rapidly quenched in cold saline solution (-2 to -3°C), extracted using cold methanol and then analyzed by LC-MS/MS. Quenching and analysis of metabolites were performed, with modifications, according to (Sake et al., 2020). A step-by-step protocol is described in (SI). Metabolite extracts were analyzed using LC-MS/MS adapted from (Young et al., 2011). A Phenomenex 150 mm x 2 mm Synergi Hydro-RP column was used on an Agilent 1200 Series HPLC system coupled to an AB Sciex 5500 QTRAP system. LC was performed with a liquid injection volume of 20 μL and a gradient elution with 10 mM tributylamine and 15 mM acetic acid (aqueous phase) in acetonitrile (organic phase) (reagent B) at a constant flow rate of 0.3 mL/min, and a constant temperature of 40°C. The gradient profile of the organic phase is as follows: 0% B (0 min), 8% B (10 min), 16% B (15 min), 30%B (16.5 min), 30% B (19 min), 90% B (21.5 min), 90% B (28 min), 0% B (28.5 min), and 0% B (35 min). MS analysis was performed in negative mode using a multiple reaction monitoring (MRM) acquisition method. Data acquisition was performed on the ABSciex Analyst 1.7 software. Absolute quantification of intracellular metabolites was performed using the quantitation mode on the Analyst software. All chemicals used for metabolite extraction and LC-MS/MS analysis were Optima grade reagents.

Acknowledgments:

WH thanks Dr. Masayuki Onishi for advice on the generation of CRISPR mutants and
Adrien Burlacot for help with PQ pool measurements.

Funding:

This project has supported by DOE award DE-SC0019417 to ARG. WH was supported
solely by a grant from the US Department of Energy, Office of Science, Basic Energy
Systems (DE-SC0019417). AK and MCP were supported solely by a grant from the US
Department of Energy, Office of Science, Basic Energy Systems (DE-SC0019341 to MP).
NB and MM were supported by US Department of Energy, Office of Science, Office of
Biological and Environmental Research (BER), (DE-SC0018301 to NB). NL and AP were
supported by German Research Foundations (DFG Grant LI1781/2-2). BR was supported
by National Science Foundation (Award Number : 1645164). RGK and JF were supported
by the Department of Plant Biology of the Carnegie Institution for Science. PR was
supported by Human Frontier Science Program (HFSP) RGP0046/2018 (to ARG).

Author contributions:

WH, AK, ARG and MP conceptualized the study. WH generated the CRISPR mutants as
well as the complemented strain, localized the TPT3 protein, performed respiration
measurements, measured the reduction level of PQ pool. AK performed the experiments of
photosynthetic O_2 evolution, extraction of samples for LC-MS/MS, data analysis of LC-
MS/MS. AP and NL performed the reconstitution into liposomes and transport activity
assays. MM and NB performed LC-MS/MS and analyses of metabolite data. WH and YW
analyzed the redox status in the cell. BR performed data mining for transcriptome and the
Nile red staining. JF and NF analyzed starch changes upon transition from LL to ML. WH,
PR and ARG analyzed PSI and PSII activities. JF and BR constructed roGFP2 associated
vectors. ARG, MP, AK, NL, NB, AP, MM and all the other authors helped with revising
and finalizing the manuscript.

Competing interests:

All other authors declare they have no competing interests.

Data and materials availability:

All data are available in the main text or the supplementary materials.

References

Ballicora, M.A., Frueauf, J.B., Fu, Y., Schürmann, P., and Preiss, J. (2000). Activation of the potato tuber ADP-glucose pyrophosphorylase by thioredoxin. J. Biol. Chem. *275*, 1315–1320. https://doi.org/10.1074/jbc.275.2.1315.

Bhatt, P., Bhandari, G., Turco, R.F., Aminikhoei, Z., Bhatt, K., and Simsek, H. (2022). Algae in wastewater treatment, mechanism, and application of biomass for production of value-added product. Environ. Pollut. *309*, 119688. https://doi.org/10.1016/j.envpol.2022.119688.

Boschetti, A., and Schmid, K. (1998). Energy Supply for ATP-Synthase Deficient Chloroplasts of *Chlamydomonas reinhardii*. Plant Cell Physiol. *39*, 160–168. https://doi.org/10.1093/oxfordjournals.pcp.a029353.

Boyle, N.R., Sengupta, N., and Morgan, J.A. (2017). Metabolic flux analysis of heterotrophic growth in *Chlamydomonas reinhardtii*. PLoS One *12*, e0177292. https://doi.org/10.1371/journal.pone.0177292.

Chaux, F., Burlacot, A., Mekhalfi, M., Auroy, P., Blangy, S., Richaud, P., and Peltier, G. (2017). Flavodiiron proteins promote fast and transient O_2 photoreduction in Chlamydomonas. Plant Physiol. *174*, 1825–1836. https://doi.org/10.1104/pp.17.00421.

Cheeseman, J.M. Hydrogen peroxide and plant stress: A challenging relationship.

Cho, M.-H., Lim, H., Shin, D.H., Jeon, J.-S., Bhoo, S.H., Park, Y.-I., and Hahn, T.-R. (2011). Role of the plastidic glucose translocator in the export of starch degradation products from the chloroplasts in *Arabidopsis thaliana*. New Phytol. *190*, 101–112. https://doi.org/10.1111/j.1469-8137.2010.03580.x.

Clowez, S., Renicke, C., Pringle, J.R., and Grossman, A.R. (2021). Impact of menthol on growth and photosynthetic function of *Breviolum minutum* (Dinoflagellata, Dinophyceae, Symbiodiniaceae) and Interactions with its Aiptasia Host. J. Phycol. *57*, 245–257. https://doi.org/10.1111/jpy.13081.

Crozet, P., Navarro, F.J., Willmund, F., Mehrshahi, P., Bakowski, K., Lauersen, K.J., Pérez-Pérez, M.-E., Auroy, P., Gorchs Rovira, A., Sauret-Gueto, S., et al. (2018). Birth of a photosynthetic chassis: A MoClo toolkit enabling synthetic biology in the microalga *Chlamydomonas reinhardtii*. ACS Synth. Biol. *7*, 2074–2086. https://doi.org/10.1021/acssynbio.8b00251.

Dorion, S., Ouellet, J.C., and Rivoal, J. (2021). Glutathione metabolism in plants under stress: beyond reactive oxygen species detoxification. Metabolites *11*, 641. https://doi.org/10.3390/metabo11090641.

Eicks, M., Maurino, V., Knappe, S., Flügge, U.-I., and Fischer, K. (2002). The plastidic pentose phosphate translocator represents a link between the cytosolic and the plastidic pentose phosphate pathways in plants. Plant Physiol. *128*, 512–522. https://doi.org/10.1104/pp.010576.

Exposito-Rodriguez, M., Laissue, P.P., Yvon-Durocher, G., Smirnoff, N., and Mullineaux, P.M.
(2017). Photosynthesis-dependent H_2O_2 transfer from chloroplasts to nuclei provides a high-light
signalling mechanism. Nat. Commun. *8*, 49. https://doi.org/10.1038/s41467-017-00074-w.

Findinier, J., Tunçay, H., Schulz-Raffelt, M., Deschamps, P., Spriet, C., Lacroix, J.-M., Duchêne,
T., Szydlowski, N., Li-Beisson, Y., Peltier, G., et al. (2017). The Chlamydomonas *mex1* mutant
shows impaired starch mobilization without maltose accumulation. J. Exp. Bot. *68*, 5177–5189.
https://doi.org/10.1093/jxb/erx343.

Findinier, J., Delevoye, C., and Cohen, M.M. (2019). The dynamin-like protein Fzl promotes
thylakoid fusion and resistance to light stress in *Chlamydomonas reinhardtii*. PLoS Genet. *15*,
e1008047. https://doi.org/10.1371/journal.pgen.1008047.

Fischer, K., Kammerer, B., Gutensohn, M., Arbinger, B., Weber, A., Hausler, R.E., and Flugge,
U.-I. (1997). A new class of plastidic phosphate translocators: A putative link between primary
and secondary metabolism by the phosphoenolpyruvate/phosphate antiporter. Plant Cell *9*, 453.
https://doi.org/10.2307/3870494.

Fliege, R., Flügge, U.-I., Werdan, K., and Heldt, H.W. (1978). Specific transport of inorganic
phosphate, 3-phosphoglycerate and triosephosphates across the inner membrane of the envelope
in spinach chloroplasts. Biochim. Biophys. Acta Bioenerg. *502*, 232–247.
https://doi.org/10.1016/0005-2728(78)90045-2.

Flügge, U.I., Fischer, K., Gross, A., Sebald, W., Lottspeich, F., and Eckerskorn, C. (1989). The
triose phosphate-3-phosphoglycerate-phosphate translocator from spinach chloroplasts:
nucleotide sequence of a full-length cDNA clone and import of the in vitro synthesized precursor
protein into chloroplasts. EMBO J. *8*, 39–46. .

Foyer, C.H., and Noctor, G. (2016). Stress-triggered redox signalling: what's in pROSpect? Plant
Cell Environ. *39*, 951–964. https://doi.org/10.1111/pce.12621.

Ge, F., Huang, W., Chen, Z., Zhang, C., Xiong, Q., Bowler, C., Yang, J., Xu, J., and Hu, H. (2014).
Methylcrotonyl-CoA carboxylase regulates triacylglycerol accumulation in the model diatom
Phaeodactylum tricornutum. Plant Cell *26*, 1681–1697. https://doi.org/10.1105/tpc.114.124982.

Gibson, D.G., Young, L., Chuang, R.-Y., Venter, J.C., Hutchison, C.A., 3rd, and Smith, H.O.
(2009). Enzymatic assembly of DNA molecules up to several hundred kilobases. Nat. Methods *6*,
343–345. https://doi.org/10.1038/nmeth.1318.

Gietz, R.D., and Schiestl, R.H. (2007). High-efficiency yeast transformation using the LiAc/SS
carrier DNA/PEG method. Nat. Protoc. *2*, 31–34. https://doi.org/10.1038/nprot.2007.13.

González-Ballester, D., Casero, D., Cokus, S., Pellegrini, M., Merchant, S.S., and Grossman, A.R.
(2010). RNA-seq analysis of sulfur-deprived Chlamydomonas cells reveals aspects of acclimation
critical for cell survival. Plant Cell *22*, 2058–2084. https://doi.org/10.1105/tpc.109.071167.

Haferkamp, I., Deschamps, P., Ast, M., Jeblick, W., Maier, U., Ball, S., and Neuhaus, H.E. (2006).
Molecular and biochemical analysis of periplastidial starch metabolism in the cryptophyte
Guillardia theta. Eukaryot. Cell *5*, 964–971. https://doi.org/10.1128/EC.00381-05.

Harris, E.H. (2009). Chlamydomonas sourcebook: Introduction to Chlamydomonas and its
laboratory use (Academic Press).

Häusler, R.E., Schlieben, N.H., Schulz, B., and Flügge, U.I. (1998). Compensation of decreased
triose phosphate/phosphate translocator activity by accelerated starch turnover and glucose
transport in transgenic tobacco. Planta *204*, 366–376. https://doi.org/10.1007/s004250050268.

Häusler, R.E., Heinrichs, L., Schmitz, J., and Flügge, U.-I. (2014). How sugars might coordinate
chloroplast and nuclear gene expression during acclimation to high light intensities. Mol. Plant *7*,
1121–1137. https://doi.org/10.1093/mp/ssu064.

Hilgers, E.J.A., Schöttler, M.A., Mettler-Altmann, T., Krueger, S., Dörmann, P., Eicks, M., Flügge,
U.-I., and Häusler, R.E. (2018). The combined loss of triose phosphate and xylulose 5-
Phosphate/Phosphate translocators leads to severe growth retardation and impaired photosynthesis
in *Arabidopsis thaliana* tpt/xpt double mutants. Front. Plant Sci. *9*, 1331.
https://doi.org/10.3389/fpls.2018.01331.

Houille-Vernes, L., Rappaport, F., Wollman, F.-A., Alric, J., and Johnson, X. (2011). Plastid
terminal oxidase 2 (PTOX2) is the major oxidase involved in chlororespiration in Chlamydomonas.
Proc. Natl. Acad. Sci. U. S. A. *108*, 20820–20825. https://doi.org/10.1073/pnas.1110518109.

Huang, W., Haferkamp, I., Lepetit, B., Molchanova, M., Hou, S., Jeblick, W., Río Bártulos, C.,
and Kroth, P.G. (2018). Reduced vacuolar β-1,3-glucan synthesis affects carbohydrate metabolism
as well as plastid homeostasis and structure in *Phaeodactylum tricornutum*. Proc. Natl. Acad. Sci.
U. S. A. *115*, 4791–4796. https://doi.org/10.1073/pnas.1719274115.

Jack, D.L., Yang, N.M., and Saier, M.H., Jr (2001). The drug/metabolite transporter superfamily.
Eur. J. Biochem. *268*, 3620–3639. https://doi.org/10.1046/j.1432-1327.2001.02265.x.

Johnson, X., and Alric, J. (2013). Central carbon metabolism and electron transport in
Chlamydomonas reinhardtii: metabolic constraints for carbon partitioning between oil and starch.
Eukaryot. Cell *12*, 776–793. https://doi.org/10.1128/EC.00318-12.

Jokel, M., Johnson, X., Peltier, G., Aro, E.-M., and Allahverdiyeva, Y. (2018). Hunting the main
player enabling *Chlamydomonas reinhardtii* growth under fluctuating light. Plant J. *94*, 822–835.
https://doi.org/10.1111/tpj.13897.

Kammerer, B., Fischer, K., Hilpert, B., Schubert, S., Gutensohn, M., Weber, A., and Flügge, U.I.
(1998). Molecular characterization of a carbon transporter in plastids from heterotrophic tissues:
the glucose 6-phosphate/phosphate antiporter. Plant Cell *10*, 105–117.
https://doi.org/10.1105/tpc.10.1.105.

Kaye, Y., Huang, W., Clowez, S., Saroussi, S., Idoine, A., Sanz-Luque, E., and Grossman, A.R.
(2019). The mitochondrial alternative oxidase from *Chlamydomonas reinhardtii* enables survival
in high light. J. Biol. Chem. *294*, 1380–1395. https://doi.org/10.1074/jbc.RA118.004667.

Khan, M.I., Shin, J.H., and Kim, J.D. (2018). The promising future of microalgae: current status,
challenges, and optimization of a sustainable and renewable industry for biofuels, feed, and other
products. Microb. Cell Fact. *17*, 36. https://doi.org/10.1186/s12934-018-0879-x.

Khorobrykh, S.A., Karonen, M., and Tyystjärvi, E. (2015). Experimental evidence suggesting that
H_2O_2 is produced within the thylakoid membrane in a reaction between plastoquinol and singlet
oxygen. FEBS Lett. *589*, 779–786. https://doi.org/10.1016/j.febslet.2015.02.011.

Klein, U. (1986). Compartmentation of glycolysis and of the oxidative pentose-phosphate pathway
in *Chlamydomonas reinhardii*. Planta *167*, 81–86. https://doi.org/10.1007/BF00446372.

Klein, U., and Betz, A. (1978). Fermentative metabolism of hydrogen-evolving *Chlamydomonas*
moewusii. Plant Physiol. *61*, 953–956. https://doi.org/10.1104/pp.61.6.953.

Klöck, G., and Kreuzberg, K. (1991). Compartmented metabolite pools in protoplasts from the
green alga *Chlamydomonas reinhardtii*: changes after transition from aerobiosis to anaerobiosis in
the dark. Biochim. Biophys. Acta *1073*, 410–415. https://doi.org/10.1016/0304-4165(91)90150-f.

Knappe, S., Flügge, U.-I., and Fischer, K. (2003). Analysis of the plastidic phosphate translocator
gene family in Arabidopsis and identification of new phosphate translocator-homologous
transporters, classified by their putative substrate-binding site. Plant Physiol. *131*, 1178–1190.
https://doi.org/10.1104/pp.016519.

Kong, F., Burlacot, A., Liang, Y., Légeret, B., Alseekh, S., Brotman, Y., Fernie, A.R., Krieger-
Liszkay, A., Beisson, F., Peltier, G., et al. (2018). Interorganelle communication: peroxisomal
MALATE DEHYDROGENASE2 connects lipid catabolism to photosynthesis through redox
coupling in Chlamydomonas. Plant Cell *30*, 1824–1847. https://doi.org/10.1105/tpc.18.00361.

Kramer, D.M., Johnson, G., Kiirats, O., and Edwards, G.E. (2004). New fluorescence parameters
for the determination of QA redox state and excitation energy fluxes. Photosynth. Res. *79*, 209.
https://doi.org/10.1023/B:PRES.0000015391.99477.0d.

Krishnan, A., Kumaraswamy, G.K., Vinyard, D.J., Gu, H., Ananyev, G., Posewitz, M.C., and
Dismukes, G.C. (2015). Metabolic and photosynthetic consequences of blocking starch
biosynthesis in the green alga *Chlamydomonas reinhardtii sta6* mutant. Plant J. *81*, 947–960.
https://doi.org/10.1111/tpj.12783.

Lee, S.-K., Eom, J.-S., Voll, L.M., Prasch, C.M., Park, Y.-I., Hahn, T.-R., Ha, S.-H., An, G., and
Jeon, J.-S. (2014). Analysis of a triose phosphate/phosphate translocator-deficient mutant reveals
a limited capacity for starch synthesis in rice leaves. Mol. Plant *7*, 1705–1708.
https://doi.org/10.1093/mp/ssu082.

Lee, Y., Nishizawa, T., Takemoto, M., Kumazaki, K., Yamashita, K., Hirata, K., Minoda, A.,
Nagatoishi, S., Tsumoto, K., Ishitani, R., et al. (2017a). Structure of the triose-

phosphate/phosphate translocator reveals the basis of substrate specificity. Nat Plants *3*, 825–832.
https://doi.org/10.1038/s41477-017-0022-8.

Lee, Y., Nishizawa, T., Takemoto, M., Kumazaki, K., Yamashita, K., Hirata, K., Minoda, A.,
Nagatoishi, S., Tsumoto, K., Ishitani, R., et al. (2017b). Crystal structure of the triose-
phosphate/phosphate translocator in complex with inorganic phosphate.
https://doi.org/10.2210/pdb5y78/pdb.

Lepistö, A., Pakula, E., Toivola, J., Krieger-Liszkay, A., Vignols, F., and Rintamäki, E. (2013).
Deletion of chloroplast NADPH-dependent thioredoxin reductase results in inability to regulate
starch synthesis and causes stunted growth under short-day photoperiods. J. Exp. Bot. *64*, 3843–
3854. https://doi.org/10.1093/jxb/ert216.

Li, M., and Kim, C. (2022). Chloroplast ROS and stress signaling. Plant Commun *3*, 100264.
https://doi.org/10.1016/j.xplc.2021.100264.

Lim, L., Linka, M., Mullin, K.A., Weber, A.P.M., and McFadden, G.I. (2010). The carbon and
energy sources of the non-photosynthetic plastid in the malaria parasite. FEBS Lett. *584*, 549–554.
https://doi.org/10.1016/j.febslet.2009.11.097.

Linka, M., Jamai, A., and Weber, A.P.M. (2008). Functional characterization of the plastidic
phosphate translocator gene family from the thermo-acidophilic red alga *Galdieria sulphuraria*
reveals specific adaptations of primary carbon partitioning in green plants and red algae. Plant
Physiol. *148*, 1487–1496. https://doi.org/10.1104/pp.108.129478.

Loddenkötter, B., Kammerer, B., Fischer, K., and Flügge, U.I. (1993). Expression of the functional
mature chloroplast triose phosphate translocator in yeast internal membranes and purification of
the histidine-tagged protein by a single metal-affinity chromatography step. Proc. Natl. Acad. Sci.
U. S. A. *90*, 2155–2159. https://doi.org/10.1073/pnas.90.6.2155.

Mittler, R., Zandalinas, S.I., Fichman, Y., and Van Breusegem, F. (2022). Reactive oxygen species
signalling in plant stress responses. Nat. Rev. Mol. Cell Biol. *23*, 663–679.
https://doi.org/10.1038/s41580-022-00499-2.

Moog, D., Nozawa, A., Tozawa, Y., and Kamikawa, R. (2020). Substrate specificity of plastid
phosphate transporters in a non-photosynthetic diatom and its implication in evolution of red alga-
derived complex plastids. Sci. Rep. *10*, 1167. https://doi.org/10.1038/s41598-020-58082-8.

Ngan, C.Y., Wong, C.-H., Choi, C., Yoshinaga, Y., Louie, K., Jia, J., Chen, C., Bowen, B., Cheng,
H., Leonelli, L., et al. (2015). Lineage-specific chromatin signatures reveal a regulator of lipid
metabolism in microalgae. Nat Plants *1*, 15107. https://doi.org/10.1038/nplants.2015.107.

Niemeyer, J., Scheuring, D., Oestreicher, J., Morgan, B., and Schroda, M. (2021). Real-time
monitoring of subcellular H_2O_2 distribution in *Chlamydomonas reinhardtii*. Plant Cell *33*, 2935–
2949. https://doi.org/10.1093/plcell/koab176.

Niittylä, T., Messerli, G., Trevisan, M., Chen, J., Smith, A.M., and Zeeman, S.C. (2004). A
previously unknown maltose transporter essential for starch degradation in leaves. Science *303*,
87–89. https://doi.org/10.1126/science.1091811.

Plaxton, W.C. (1996). The organization and regulation of plant glycolysis. Annu. Rev. Plant
Physiol. Plant Mol. Biol. *47*, 185–214. https://doi.org/10.1146/annurev.arplant.47.1.185.

Raghavendra, A.S., and Padmasree, K. (2003). Beneficial interactions of mitochondrial
metabolism with photosynthetic carbon assimilation. Trends Plant Sci. *8*, 546–553.
https://doi.org/10.1016/j.tplants.2003.09.015.

Riesmeier, J.W., Flügge, U.I., Schulz, B., Heineke, D., Heldt, H.W., Willmitzer, L., and Frommer,
W.B. (1993). Antisense repression of the chloroplast triose phosphate translocator affects carbon
partitioning in transgenic potato plants. Proc. Natl. Acad. Sci. U. S. A. *90*, 6160–6164.
https://doi.org/10.1073/pnas.90.13.6160.

Rochaix, J.-D., Goldschmidt-Clermont, M., and Merchant, S. (1998). The molecular biology of
chloroplasts and mitochondria in Chlamydomonas (Plenum Publishing Corporation).

Sake, C.L., Newman, D.M., and Boyle, N.R. (2020). Evaluation of quenching methods for
metabolite recovery in photoautotrophic *Synechococcus sp.* PCC 7002. Biotechnol. Prog. *36*,
e3015. https://doi.org/10.1002/btpr.3015.

Saroussi, S., Sanz-Luque, E., Kim, R.G., and Grossman, A.R. (2017). Nutrient scavenging and
energy management: acclimation responses in nitrogen and sulfur deprived Chlamydomonas. Curr.
Opin. Plant Biol. *39*, 114–122. https://doi.org/10.1016/j.pbi.2017.06.002.

Schneider, A., Häusler, R.E., Kolukisaoglu, Ü., Kunze, R., Van der Graaff, E., Schwacke, R.,
Catoni, E., Desimone, M., and Flügge, U.I. (2002). An Arabidopsis thaliana knock-out mutant of
the chloroplast triose phosphate/phosphate translocator is severely compromised only when starch
synthesis, but not starch mobilisation is abolished. Plant Journal https://doi.org/10.1046/j.1365-
313X.2002.01460.x.

Shapiguzov, A., Vainonen, J.P., Wrzaczek, M., and Kangasjärvi, J. (2012). ROS-talk - how the
apoplast, the chloroplast, and the nucleus get the message through. Front. Plant Sci. *3*, 292.
https://doi.org/10.3389/fpls.2012.00292.

Stocking, C.R., and Larson, S. (1969). A chloroplast cytoplasmic shuttle and the reduction of
extraplastid NAD. Biochem. Biophys. Res. Commun. *37*, 278–282. https://doi.org/10.1016/0006-
291x(69)90731-1.

Strenkert, D., Schmollinger, S., Gallaher, S.D., Salomé, P.A., Purvine, S.O., Nicora, C.D., Mettler-
Altmann, T., Soubeyrand, E., Weber, A.P.M., Lipton, M.S., et al. (2019). Multiomics resolution
of molecular events during a day in the life of Chlamydomonas. Proc. Natl. Acad. Sci. U. S. A.
116, 2374–2383. https://doi.org/10.1073/pnas.1815238116.

Suzuki, N., Koussevitzky, S., Mittler, R., and Miller, G. (2012). ROS and redox signalling in the
response of plants to abiotic stress. Plant Cell Environ. *35*, 259–270.
https://doi.org/10.1111/j.1365-3040.2011.02336.x.

Urzica, E.I., Vieler, A., Hong-Hermesdorf, A., Page, M.D., Casero, D., Gallaher, S.D., Kropat, J.,
Pellegrini, M., Benning, C., and Merchant, S.S. (2013). Remodeling of membrane lipids in iron-
starved Chlamydomonas. J. Biol. Chem. *288*, 30246–30258.
https://doi.org/10.1074/jbc.M113.490425.

Vevea, J.D., Alessi Wolken, D.M., Swayne, T.C., White, A.B., and Pon, L.A. (2013). Ratiometric
biosensors that measure mitochondrial redox state and ATP in living yeast cells. J. Vis. Exp. 50633.
https://doi.org/10.3791/50633.

Wakao, S., and Niyogi, K.K. (2021). Chlamydomonas as a model for reactive oxygen species
signaling and thiol redox regulation in the green lineage. Plant Physiol. *187*, 687–698.
https://doi.org/10.1093/plphys/kiab355.

Walters, R.G., Ibrahim, D.G., Horton, P., and Kruger, N.J. (2004). A mutant of Arabidopsis
lacking the triose-phosphate/phosphate translocator reveals metabolic regulation of starch
breakdown in the light. Plant Physiol. *135*, 891–906. https://doi.org/10.1104/pp.104.040469.

Weber, A.P.M., Linka, M., and Bhattacharya, D. (2006). Single, ancient origin of a plastid
metabolite translocator family in Plantae from an endomembrane-derived ancestor. Eukaryot. Cell
5, 609–612. https://doi.org/10.1128/EC.5.3.609-612.2006.

Winder, C.L., Dunn, W.B., Schuler, S., Broadhurst, D., Jarvis, R., Stephens, G.M., and Goodacre,
R. (2008). Global metabolic profiling of *Escherichia coli* cultures: an evaluation of methods for
quenching and extraction of intracellular metabolites. Anal. Chem. *80*, 2939–2948.
https://doi.org/10.1021/ac7023409.

Young, J.D., Shastri, A.A., Stephanopoulos, G., and Morgan, J.A. (2011). Mapping
photoautotrophic metabolism with isotopically nonstationary [13]C flux analysis. Metab. Eng. *13*,
656–665. https://doi.org/10.1016/j.ymben.2011.08.002.

Yu, E.T., Zendejas, F.J., Lane, P.D., Gaucher, S., Simmons, B.A., and Lane, T.W. (2009).
Triacylglycerol accumulation and profiling in the model diatoms *Thalassiosira pseudonana* and
Phaeodactylum tricornutum (Baccilariophyceae) during starvation. J. Appl. Phycol. *21*, 669–681.
https://doi.org/10.1007/s10811-008-9400-y.

Zhao, Y., Luo, L., Xu, J., Xin, P., Guo, H., Wu, J., Bai, L., Wang, G., Chu, J., Zuo, J., et al. (2018).
Malate transported from chloroplast to mitochondrion triggers production of ROS and PCD in
Arabidopsis thaliana. Cell Res. *28*, 448–461. https://doi.org/10.1038/s41422-018-0024-8.

Zhao, Y., Yu, H., Zhou, J.-M., Smith, S.M., and Li, J. (2020). Malate circulation: linking
chloroplast metabolism to mitochondrial ROS. Trends Plant Sci. *25*, 446–454.
https://doi.org/10.1016/j.tplants.2020.01.010.

Zirngibl, M.-E., Araguirang, G.E., Kitashova, A., Jahnke, K., Rolka, T., Kühn, C., Nägele, T., and
Richter, A.S. (2022). Triose phosphate export from chloroplasts and cellular sugar content regulate
anthocyanin biosynthesis during high light acclimation. Plant Commun 100423.
https://doi.org/10.1016/j.xplc.2022.100423.

Zones, J.M., Blaby, I.K., Merchant, S.S., and Umen, J.G. (2015). High-Resolution Profiling of a
Synchronized Diurnal Transcriptome from *Chlamydomonas reinhardtii* Reveals Continuous Cell
and Metabolic Differentiation. Plant Cell *27*, 2743–2769. https://doi.org/10.1105/tpc.15.00498.

Legends

Fig. 1: Subcellular localization of CreTPT3 and in vitro measurements of CreTPT3 transport
activities. (**A**) Cellular localization of CreTPT3. The construct for expression of the CreTPT3-VENUS
fusion protein is shown above the cell images and described in the **Materials and methods**. VENUS
fluorescence, green; Chlorophyll (Chl) autofluorescence, magenta. (Scale bar: 5 μm). (**B**) Expression of
Cre*TPT3* in yeast. Left, SDS-PAGE of proteins from total membranes isolated from yeast cells containing
the empty vector (lane 1) or expressing His-CreTPT3 (lane 2). Right, corresponding immunoblot blot
detection of recombinant His-CreTPT3 protein using anti-His antibodies. The calculated molecular mass
of the N-terminal His-tagged protein was 37 kDa. (**C**) The kinetics of Pi exchange by His-CreTPT3 in lipid
vesicles. Uptake of Pi (0.25 mM) into liposomes was measured in the presence (■) or absence (□) of Pi
(30 mM) loaded within the vesicles. The arithmetic mean (±SD) of three independent experiments (each
with three technical replicates) was plotted with respect to the time after initiating the assay. (**D**) Analyses
of substrate specificity of His-CreTPT3. Liposomes in which His-CreTPT3 was incorporated were
preloaded with various substrates (30 mM) and the initial Pi uptake rates (at 0.25 mM external concentration)
were determined. Relative velocities were compared to the Pi/Pi homo-exchange, which was set to 100%.
The data represents the arithmetic mean (±SD) of three independent experiments (each consisting of three
technical replicates). (**E**) Comparison of kinetic constants of CreTPT3. The Michaelis-Menten constant (K_M)
for Pi uptake was determined using various external Pi concentrations (0.05-5 mM). The competitive
inhibition constant (K_i) of Pi uptake (0.25 mM) was measured with increasing competitor concentrations
(0.05-5 mM). Liposomes were preloaded with 30 mM Pi as the counter-exchange substrate. The data
represent the arithmetic mean +/- SE of three independent experiments. ND, no competitive inhibitory
constant could be measured under the given experimental conditions. DHAP, dihydroxyacetone phosphate;
3-PGA, 3-phosphoglycerate; PEP, phosphoenolpyruvate; Glc6P, glucose 6-P; Glc1P, glucose 1-P; Fru6P,
fructose 6-P; Gal1P, galactose 1-P.

Fig. 2: Growth characterization of *tpt3* mutants and complemented strains. (**A-B**) Cre*TPT3* and the
positions edited using CRISPR-Cas9. Molecular map of Cre*TPT3* gene and positions of the Cas9 targeted
sites (**A**) Edited sites are located in exons 1 and 7; *t3ko1* has the marker gene inserted in exon 1 and *t3ko2*
and *t3ko3* has the marker gene inserted in exon 7. (**B**) Immunoblot (left panel) of the stained protein profiles
(right panel) for wild type, *t3ko1*, *t3ko2*, *t3ko3* and the complemented strains (*C_T3KO1*, *C_T3KO2* and
C_T3KO3, respectively). (**C**) Growth of WT, *t3ko1*, *t3ko2*, *t3ko3* and the complemented strains (*C_T3KO1*,
C_T3KO2, and *C_T3KO3*, respectively), at various dilutions (see below) on agar plates incubated under
LL and ML for 4 d. (**D**) Growth of the various strains (indicated) in liquid medium under LL (left panel)
and ML (right panel). (**E**) Transition of the strains from LL to ML (upper panel) or to high light (HL, 450
μmol photons m^{-2} s^{-1}) (lower panel) for the times indicated. (**F**) Cell growth (number) and chlorophyll
content of cultures at various times after shifting from LL to ML for up to 48 h. In (**C**) the cells were spotted
on agar plates containing TAP or TP medium and maintained under continuous LL or ML conditions. The
dilution series used was 1.5, 0.75, 0.375 μg/mL chlorophyll (left to right). For growth in liquid medium,
the cells were cultured in TP medium to an initial OD_{750nm} 0.02 in air and under continuous LL before
initiating the various growth analyses in LL, ML or HL. Each curve represents the arithmetic mean (±SD)
of three independent experiments.
1003

Fig. 3: Changes in storage carbon and metabolite levels in WT and *t3ko2* after transitioning cells from growth in LL to ML. (**A**) Starch content in indicated strains following 0, 6, and 24 h of ML exposure. (**B**) Lugol staining of starch in WT and *t3ko2* after 48 h of ML. (**C**) TAG (TAG: triacylglycerol) content in the indicated strains following 0, 6, and 24 h of ML exposure. (**D**) Nile Red staining of TAG in WT and *t3ko2* following 48 h of ML exposure. (**E**) Time course of metabolite accumulation at 0 h (LL), 45 min and 6 h following ML exposure. (**F**) Select metabolic pathways in Chlamydomonas adapted from (Johnson and Alric, 2013). (**G**) Bar graph representation of fold-change for metabolites shown in (**E**); calculated by dividing the pool size in *t3ko2* by that of WT at the respective light levels. A fold change of 1 (no change) is indicated with a dashed line. Each data point shows the mean and standard error; the data represents three biological replicates for each metabolite. An asterisk indicates statistically significant differences compared to WT (* P<0.05, ** P<0.01, *** P<0.001). Abbreviations: G6P, glucose-6-P; FBP, fructose bisphosphate; DHAP, dihydroxyacetone phosphate; 3-PGA, 3-phosphoglycerate; PEP, phosphoenolpyruvate; PYR, pyruvate; CIT, citrate; SUCC, succinate; FUM, fumarate; MAL, malate; E4P, Erythrose 4-P; R5P, Ribose 5-P; RU5P/X5P, ribulose 5-P/xylulose-5-P; OMT, 2-oxoglutarate/malate transporter.

Fig. 4: Photosynthetic activities followed LL to ML or HL transitions. (**A**) O_2 evolution of LL (left) and ML (right)-acclimated cells for the indicated strains. (**B**) Fv/Fm of WT, *t3ko2* and the rescued strain (*C_T3KO2*) following 0 (LL), 6 and 24 h in ML. (**C**) 1-qL values of LL (left panel) and ML (right panel) acclimated cells exposed to increasing actinic light intensities. (**D**) Kinetics of 1-qL of LL-acclimated cells after illumination at HL (400-450 µmol photons m^{-2} s^{-1}) actinic light for the times indicated on the x-axis. (**E**) P700 oxidation and reduction kinetics in LL-grown cells and after 6 and 24 h in ML. (**F**) Kinetics of P700 oxidation upon dark-to-light transition**.** P700 measurements were made in the presence of DCMU and hydroxylamine, as indicated in the **Materials and methods**. Absorbance differences were monitored at 705 nm during continuous illumination with 150 µmol photons·m^{-2} s^{-1} for 5 sec (white box above), followed by a saturating light pulse at 1,500 µmol photons·m^{-2} s^{-1} (arrow) and a 5 sec dark incubation (black box). For panel E, kinetics was normalized by setting maximum oxidation (after light pulse) of WT to 100% and for panel F, by setting maximum oxidation (after light pulse) of individual strain to 100%. The data from all panels represent the arithmetic mean (±SD) of three independent experiments. For panel **B**, the asterisks represent significant differences determined by ANOVA tests., ****P < 0.0001.

Fig. 5: Measurement of intracellular ROS in *t3ko2* and WT upon transition from LL to ML. (**A**) ROS levels were evaluated by CM-H2DCFDA fluorescence of WT and the *t3ko2* mutant in LL and following exposure to ML for 48 h. (**B**) Quantitation of data in **A**. roGFP2 protein was targeted to the chloroplast stroma (**C**, upper panel) and mitochondrion (**C**, lower panel). BF: bright field; roGFP2 fluorescence: green; Chlorophyll (Chl) autofluorescence: red. (**D-E**) Monitoring redox levels in stroma (**D**) and mitochondrion (**E**) after exposure of WT and the *t3ko2* mutant for 1.5 and 6.0 h of ML. Shown are mean values from three independent experiments, error bars represent standard deviation. For panel **B** and **D**, the asterisks represent significant differences determined by ANOVA tests., **P < 0.005, ***P < 0.001.

Fig. 6: Real-time monitor of H₂O₂ distribution in *t3ko2* and WT upon transition from LL to HL. (A) The hypersensitive H_2O_2 sensor (roGFP2-Tsa2ΔCR) protein was targeted to the chloroplast stroma (St), cytosol (Cyto), mitochondrion (Mito) and nucleus (Nu). Shown are GFP fluorescence, chlorophyll autofluorescence (Chl) and the two signals merged (scale bar: 5 μm). WT and *t3ko2* transformant cells, accumulating roGFP2-Tsa2ΔCR in the stroma (**B**), cytosol (**C**), mitochondrial matrix (**D**), and nucleus (**E**), were acclimated to LL in TP and then were transferred either to HL (450 μmol photons m^{-2} s^{-1}, solid line) or to very LL (10 μmol photons m^{-2} s^{-1}, dotted line). The oxidation state of the sensor was trapped by the addition of NEM and roGFP2 fluorescence was measured in a plate reader as previously described (Niemeyer et al., 2021). Shown are mean values from three independent experiments with the error bars representing the standard deviation.

Fig. 7: Impact of eliminating Chlamydomonas TPT3 on the chloroplast metabolic landscape. i. reduced triose-P export leads to accumulation of the sugar-P in CBBC and glycolysis pathways (e.g. triose-P, hexose-P, marked in orange), and an elevated ratio of 3-PGA/Pi (which can allosterically activate AGPase activity) would begin to stimulate production of storage carbon; **ii.** CBBC activity diminishes as a consequence of the hyper-accumulation of the precursors along with the fixation of CO_2, leading to elevation of reductant (NADPH) and energy (ATP) within the chloroplast; the elevated $NADPH:NADP^+$ and ATP:ADP ratios, elicit strong feedback on PET causing hyper-reduction of electron carriers (eg: over reduced PQ pool and highly reduced PSI) that slows electron flow across the cytochrome b_6f complex; **iii.** the highly reduced PET system and the elevated NADPH can actively reduce FDX/TRX and the NTRC systems, respectively; reduced TRX and/or NTRC can activate the AGPase (above the activity elicited by an increase in the 3-PGA/Pi ratio), leading to additional starch accumulation; **iv.** hyper-reduction of PET can also lead to stromal ROS accumulation, that can act as a signal that controls expression of chloroplast and nuclear genes; ROS also cause damage to both PSII and I and leads to neutral lipid (TAG: triacylglycerol) accumulation as a consequence of activation of diacylglycerol acyltransferases (DGAT) and phospholipid diacylglycerol acyltransferase (PDAT). Many of these metabolic/acclimatory processes both overlap and are interconnected. Under conditions of extended exposure to ML, the damage in Chlamydomonas *tpt3* mutants can accumulate and lead to cell death; cell death occurs rapidly in high light.

Figures and Tables

1070

Fig. 1: Subcellular localization of CreTPT3 and in vitro measurements of CreTPT3 transport activities. (**A**) Cellular localization of CreTPT3. The construct for expression of the CreTPT3-VENUS fusion protein is shown above the cell images and described in the **Materials and methods**. VENUS fluorescence, green; Chlorophyll (Chl) autofluorescence, magenta. (Scale bar: 5 μm). (**B**) Expression of Cre*TPT3* in yeast. Left, SDS-PAGE of proteins from total membranes isolated from yeast cells containing the empty vector (lane 1) or expressing His-CreTPT3 (lane 2). Right, corresponding immunoblot blot detection of recombinant His-CreTPT3 protein using anti-His antibodies. The calculated molecular mass of the N-terminal His-tagged protein was 37 kDa. (**C**) The kinetics of Pi exchange by His-CreTPT3 in lipid vesicles. Uptake of Pi (0.25 mM) into liposomes was measured in the presence (■) or absence (□) of Pi (30 mM) loaded within the vesicles. The arithmetic mean (±SD) of three independent experiments (each with three technical replicates) was plotted with respect to the time after initiating the assay. (**D**) Analyses of substrate specificity of His-CreTPT3. Liposomes in which His-CreTPT3 was incorporated were preloaded with various substrates (30 mM) and the initial Pi uptake rates (at 0.25 mM external concentration) were determined. Relative velocities were compared to the Pi/Pi homo-exchange, which was set to 100%. The data represents the arithmetic mean (±SD) of three independent experiments (each consisting of three technical replicates). (E) Comparison of kinetic constants of CreTPT3. The Michaelis-Menten constant (K_M) for Pi uptake was determined using various external Pi concentrations (0.05-5 mM). The competitive inhibition constant (K_i) of Pi uptake (0.25 mM) was measured with increasing competitor concentrations (0.05-5 mM). Liposomes were preloaded with 30 mM Pi as the counter-exchange substrate. The data represent the arithmetic mean +/- SE of three independent experiments. ND, no competitive inhibitory constant could be measured under the given experimental conditions. DHAP, dihydroxyacetone phosphate; 3-PGA, 3-phosphoglycerate; PEP, phosphoenolpyruvate; Glc6P, glucose 6-P; Glc1P, glucose 1-P; Fru6P, fructose 6-P; Gal1P, galactose 1-P.

1094

1095

Fig. 2: Growth ━━━━━━━━ **d strains.** (**A-B**) Cre*TPT3* and the positions edited using CRISPR-Cas9. Molecular map of Cre*TPT3* gene and positions of the Cas9 targeted sites (**A**) Edited sites are located in exons 1 and 7; *t3ko1* has the marker gene inserted in exon 1 and *t3ko2* and *t3ko3* has the marker gene inserted in exon 7. (**B**) Immunoblot (left panel) of the stained protein profiles (right panel) for wild type, *t3ko1*, *t3ko2*, *t3ko3* and the complemented strains (*C_T3KO1*, *C_T3KO2* and *C_T3KO3*, respectively). (**C**) Growth of WT, *t3ko1*, *t3ko2*, *t3ko3* and the complemented strains (*C_T3KO1*, *C_T3KO2*, and *C_T3KO3*, respectively), at various dilutions (see below) on agar plates incubated under LL and ML for 4 d. (**D**) Growth of the various strains (indicated) in liquid medium under LL (left panel) and ML (right panel). (**E**) Transition of the strains from LL to ML (upper panel) or to high light (HL, 450 μmol photons m^{-2} s^{-1}) (lower panel) for the times indicated. (**F**) Cell growth (number) and chlorophyll content of cultures at various times after shifting from LL to ML for up to 48 h. In (**C**) the cells were spotted on agar plates containing TAP or TP medium and maintained under continuous LL or ML conditions. The dilution series used was 1.5, 0.75, 0.375 μg/mL chlorophyll (left to right). For growth in liquid medium, the cells were cultured in TP medium to an initial OD$_{750nm}$ 0.02 in air and under continuous LL before initiating the various growth analyses in LL, ML or HL. Each curve represents the arithmetic mean (±SD) of three independent experiments.

1115

1116

Fig. 3: Changes in storage carbon and metabolite levels in WT and *t3ko2* after transitioning cells from growth in LL to ML. (**A**) Starch content in indicated strains following 0, 6, and 24 h of ML exposure. (**B**) Lugol staining of starch in WT and *t3ko2* after 48 h of ML. (**C**) TAG (TAG: triacylglycerol) content in the indicated strains following 0, 6, and 24 h of ML exposure. (**D**) Nile Red staining of TAG in WT and *t3ko2* following 48 h of ML exposure. (**E**) Time course of metabolite accumulation at 0 h (LL), 45 min and 6 h following ML exposure. (**F**) Select metabolic pathways in Chlamydomonas adapted from (Johnson and Alric, 2013). (**G**) Bar graph representation of fold-change for metabolites shown in (**E**); calculated by dividing the pool size in *t3ko2* by that of WT at the respective light levels. A fold change of 1 (no change) is indicated with a dashed line. Each data point shows the mean and standard error; the data represents three biological replicates for each metabolite. An asterisk indicates statistically significant differences compared to WT (* P<0.05, ** P<0.01, *** P<0.001). Abbreviations: G6P, glucose-6-P; FBP, fructose bisphosphate; DHAP, dihydroxyacetone phosphate; 3-PGA, 3-phosphoglycerate; PEP, phosphoenolpyruvate; PYR, pyruvate; CIT, citrate; SUCC, succinate; FUM, fumarate; MAL, malate; E4P, Erythrose 4-P; R5P, Ribose 5-P; RU5P/X5P, ribulose 5-P/xylulose-5-P; OMT, 2-oxoglutarate/malate transporter.

1131

Fig. 4: Photosynthetic activities followed LL to ML or HL transitions. (**A**) O_2 evolution of LL (left) and ML (right)-acclimated cells for the indicated strains. (**B**) Fv/Fm of WT, *t3ko2* and the rescued strain (*C_T3KO2*) following 0 (LL), 6 and 24 h in ML. (**C**) 1-qL values of LL (left panel) and ML (right panel) acclimated cells exposed to increasing actinic light intensities. (**D**) Kinetics of 1-qL of LL-acclimated cells after illumination at HL (400-450 µmol photons $m^{-2} s^{-1}$) actinic light for the times indicated on the x-axis. (**E**) P700 oxidation and reduction kinetics in LL-grown cells and after 6 and 24 h in ML. (**F**) Kinetics of P700 oxidation upon dark-to-light transition**.** P700 measurements were made in the presence of DCMU and hydroxylamine, as indicated in the **Materials and methods**. Absorbance differences were monitored at 705 nm during continuous illumination with 150 µmol photons·$m^{-2} s^{-1}$ for 5 sec (white box above), followed by a saturating light pulse at 1,500 µmol photons·$m^{-2} s^{-1}$ (arrow) and a 5 sec dark incubation (black box). For panel E, kinetics was normalized by setting maximum oxidation (after light pulse) of WT to 100% and for panel F, by setting maximum oxidation (after light pulse) of individual strain to 100%. The data from all panels represent the arithmetic mean (±SD) of three independent experiments. For panel **B**, the asterisks represent significant differences determined by ANOVA tests., ****P < 0.0001.

Fig. 5: Measurement of intracellular ROS in *t3ko2* and WT upon transition from LL to ML. (**A**) ROS
levels were evaluated by CM-H2DCFDA fluorescence of WT and the *t3ko2* mutant in LL and following
exposure to ML for 48 h. (**B**) Quantitation of data in **A**. roGFP2 protein was targeted to the chloroplast
stroma (**C**, upper panel) and mitochondrion (**C**, lower panel). BF: bright field; roGFP2 fluorescence: green;
Chlorophyll (Chl) autofluorescence: red. (**D-E**) Monitoring redox levels in stroma (**D**) and mitochondrion
(**E**) after exposure of WT and the *t3ko2* mutant for 1.5 and 6.0 h of ML. Shown are mean values from three
independent experiments, error bars represent standard deviation. For panel **B** and **D**, the asterisks represent
significant differences determined by ANOVA tests., $**P < 0.005$, $***P < 0.001$.

1156

Fig. 6: Real-time monitor of H$_2$O$_2$ compartmentalized distribution in *t3ko2* and WT upon transition from LL to HL. (**A**) The hypersensitive H$_2$O$_2$ sensor (roGFP2-Tsa2ΔCR) protein was targeted to the chloroplast stroma (St), cytosol (Cyto), mitochondrion (Mito) and nucleus (Nu). Shown are GFP fluorescence, chlorophyll autofluorescence (Chl) and the two signals merged (scale bar: 5 μm). WT and *t3ko2* transformant cells, accumulating roGFP2-Tsa2ΔCR in the stroma (**B**), cytosol (**C**), mitochondrial matrix (**D**), and nucleus (**E**), were acclimated to LL in TP and then were transferred either to HL (450 μmol photons m^{-2} s^{-1}, solid line) or to very LL (10 μmol photons m^{-2} s^{-1}, dotted line). The oxidation state of the sensor was trapped by the addition of NEM and roGFP2 fluorescence was measured in a plate reader as previously described (Niemeyer et al., 2021). Shown are mean values from three independent experiments with the error bars representing the standard deviation.

1167

1168

Fig. 7: Impact of eliminating Chlamydomonas TPT3 on the chloroplast metabolic landscape. i.
reduced triose-P export leads to accumulation of the sugar-P in CBBC and glycolysis pathways (e.g. triose-
P, hexose-P, marked in orange), and an elevated ratio of 3-PGA/Pi (which can allosterically activate
AGPase activity) would begin to stimulate production of storage carbon; **ii.** CBBC activity diminishes as a
consequence of the hyper-accumulation of the precursors along with the fixation of CO_2, leading to
elevation of reductant (NADPH) and energy (ATP) within the chloroplast; the elevated NADPH:NADP+
and ATP:ADP ratios, elicit strong feedback on PET causing hyper-reduction of electron carriers (eg: over
reduced PQ pool and highly reduced PSI) that slows electron flow across the cytochrome b_6f complex; **iii.**
the highly reduced PET system and the elevated NADPH can actively reduce FDX/TRX and the NTRC
systems, respectively; reduced TRX and/or NTRC can activate the AGPase (above the activity elicited by
an increase in the 3-PGA/Pi ratio), leading to additional starch accumulation; **iv.** hyper-reduction of PET
can also lead to stromal ROS accumulation, that can act as a signal that controls expression of chloroplast
and nuclear genes; ROS also cause damage to both PSII and I and leads to neutral lipid accumulation as a
consequence of activation of diacylglycerol acyltransferases (DGAT) and phospholipid diacylglycerol
acyltransferase (PDAT). Many of these metabolic/acclimatory processes both overlap and are
interconnected. Under conditions of extended exposure to ML, the damage in Chlamydomonas *tpt3* mutants
can accumulate and lead to cell death; cell death occurs rapidly in high light.

1186

Supplementary Materials

Materials and methods

Expression of CreTPT3 in yeast

The DNA sequence encoding the mature CreTPT3 protein was codon-optimized for expression in *Saccharomyces cerevisiae* (GeneART, ThermoFisher Scientific). This coding sequence was inserted in frame with an N-terminal His tag into the yeast vector pYES-NTa (ThermoFisher Scientific) using Gibson cloning (NEB). Briefly, pYES-NTa was linearized with BamHI and the CreTPT3 cDNA amplified with the primer pair fwd:5'- gacgataaggtacctagcGCTTCTGCTGCTGATGCTC-3' and rev:5'- agaattccaccacactgTTAAGCGGCAGCTGGAGAAG-3'. This cDNA was ligated into the linearized pYES-NTa vector, which was then transformed into the yeast strain INVSc1 (MATa, his3D1, leu2, trp1-289, ura3-52 / MATa, his3D1, leu2, trp1-289, and ura3-52, Thermo Fisher Scientific) using the lithium-acetate/PEG method (Gietz and Schiestl, 2007). Transformed yeast cells were selected on synthetic complete medium containing 2% (w/v) glucose with the uracil auxotrophic marker. Galactose-inducible expression of His-CreTPT3 in yeast was performed as described in (Linka et al., 2008). The presence of the His-tagged fusion protein was verified by standard SDS-PAGE and immunoblot analysis using an anti-His-antibody conjugated with horseradish peroxidase (Miltenyi Biotech).

Transport activity assays

The Km for Pi was determined by measuring the initial velocity at each of six external Pi concentrations between 0.05 and 5 mM. To obtain competitive inhibition constants (Ki), the uptake of 0.25 mM Pi into liposomes containing 30 mM Pi was measured over a 4 min period in the presence of increasing external competitor concentrations (0.05 - 5 mM). Three biological replicates were performed for all described experiments. GraphPad Prism software version 9.3.0 was used for non-linear regression analyses of the kinetic data.

Vector construction

This plasmid was linearized with HpaI (NEB). Primers: g*TPT3*_pRam118_f and g*TPT3*_pRam118_r (**Supplementary Table 1**) were used to amplify Cre*TPT3* genomic DNA containing an overlap with the linearized pRam118_VENUS vector. For generating the plasmid pRam118-CreTPT3&VENUS, genomic DNA of Cre*TPT3* was assembled with pRam118_VENUS plasmid using Gibson assembly(Gibson et al., 2009). A 1,000 bp sequence upstream of Cre*TPT3* and containing the promoter region of the gene was amplified using the primers TPT3pro1000_f and TPT3pro1000_r (**Supplementary Table 1**). The 3' UTR of Cre*TPT3* was amplified using TPT_3UTR_f and TPT3_3UTR_r (**Supplementary Table 1**). The *PSAD* promoter and the *RBCS2* 3'UTR of pRam118-CreTPT3&VENUS were replaced by the amplified fragments of the 1,000 bp upstream region and the 3' UTR of Cre*TPT3*, respectively. This final vector, designated pRam118_CreTPT3, contains the original Cre*TPT3* promoter (driving expression of Cre*TPT3*), the genomic DNA sequence of Cre*TPT3*, VENUS and the Cre*TPT3* 3' UTR plus the *AphVII* cassette. The Chlamydomonas *tpt3* mutant was transformed by electroporation with a total amount of 2 μg (~500 ng/μL) of pRam118_CreTPT3 that was linearized with AseI.

CRISPR-Cas9 mediated mutagenesis

Prior to electroporation, Cas9 (IDT) and sgRNAs were incubated together at 37 °C for 30 min. Approximately 500 ng PCR product of the *AphVII* cassette, which confers resistance to hygromycin, was added to the RNP (ribonucleoprotein) mixture. 250 µl aliquots were electroporated using Super Electroporator NEPA21 type II (NEPA GENE). After 16 h of recovery in very low light (10-15 µmol photons m^{-2} s^{-1}), cells were plated onto solid TAP medium containing 10 µg/mL hygromycin. Sense or antisense-oriented knock-ins of *AphVII* were determined by amplification using primer pairs with one primer annealing to the genomic sequence and the other to the inserted sequence (**Supplementary Table 1**). The amplified fragments were sequenced to verify the insertion sites (ELIM BIOPHARM, Hayward, USA).

ROS measurements and roGFP2 imaging analysis

Briefly, after various treatment (e.g. 48 h in ML), 10 million Chlamydomonas cells were pelleted and washed once with 1×PBS. The cells were then resuspended in 1×PBS containing 8 µM CM-H2DCFDA and incubated at room temperature in the dark for 30 min. Following this incubation, the cells were washed three times with 1×PBS buffer, the fluorescent signals were either visualized using TCS SP8 confocal laser-scanning microscope (Leica) or quantified with a microplate reader (Infinite M1000; TECAN). Excitation and emission settings for the microscope were: 488 nm/510-530 nm HyD SMD hybrid detector for reactive oxygen species (ROS), and 488 nm/650-700 nm-HyD SMD hybrid detector for chlorophyll autofluorescence. Excitation and emission settings for the plate reader were as follows: ROS excitation 488/5 nm and emission 530/12 nm; chlorophyll excitation 514/5 nm and emission 690/5 nm.

Starch analysis

In brief, cells were collected by centrifugation and pigments extracted with methanol. Dried pellets were resuspended in water and heated at 100 °C to break the cells and release the starch. After cooling, amyloglucosidase and α-amylase (2.25 U/mL) were used to hydrolyze the starch and the products of hydrolysis were quantified using a Glucose Colorimetric Detection Kit (Thermo Fisher).

Photosynthesis-Irradiance curve

The YSI 5331A electrodes (Yellow Springs Instruments) were polarized at −0.8 V. The cultures used for the assays were concentrated to 2.5-5 µg mL^{-1} chlorophyll, supplemented with 15 µL of 0.5 M sodium bicarbonate in water and then purged with 1% CO_2/99% He. Using a gas tight syringe, the sample was transferred into the reaction chamber that was also purged with 1% CO_2/99% He. The rate of change in O_2 levels was measured sequentially at the light intensities 100, 200, 600, 1000 µmol photons m^{-2} s^{-1} (photosynthetic active radiation, PAR); each intensity was maintained for 3-5 min followed by a 3-min intervening dark period, and then the light level was raised to the next higher intensity (stepped change) until the full range of intensities was tested. At 1,000 µmol photons m^{-2} s^{-1}, the light was held for 10 min followed by an 8-min dark period. Before the measurement of each experimental series, the electrodes were calibrated with air (∼21% O_2) and 1% CO_2/99% He mixture (0% O_2). The initial slope of the response was used to determine the O_2 evolution rate.

Extraction of samples for LC-MS/MS

The quenching solution, filtered saline (9 g/L NaCl), was prechilled to 4°C in a refrigerator. 30 mL of the quenching solution was transferred to a 50 mL conical tube kept in an ice bath mixed with salt to depress the temperature to between -3°C and -1°C. 15 mL of the culture was rapidly plunged into the 30 mL of the quenching solution and the samples were centrifuged at 4,000 rpm for 10 min at 4°C. Cell pellets were washed with fresh prechilled saline solution and centrifuged again at 4,000 rpm for 10 min in a 2 mL centrifuge tube. For each replicate, 45 mL of the culture was sampled and the cell pellets were pooled. Washed pellets were frozen at -80°C until extraction. Metabolite extraction and further analysis were modified from(Young et al., 2011). To each cell pellet, 500 μL methanol was added along with the internal standards ribitol and PIPES to a final concentration of 2 μM each. Samples were then vortexed for 30 sec, frozen in liquid nitrogen as described in(Winder et al., 2008), and allowed to thaw at 0°C. This vortex-freeze-thaw cycle was repeated twice more before the samples were centrifuged at 10,000 x g at 1°C for 5 min. The supernatant was collected, and the remaining pellets were extracted twice more with 500 μL of a 50:50 mixture of methanol and water, with 3 vortex-freeze-thaw cycles done for each extraction. Supernatants from each extraction procedure were pooled and dried on a Thermo Fisher SpeedVac Concentrator. Dried extracts were then resuspended in 500 μL water and cleaned to remove any residual large cell debris by filtration, first through a 0.22 μm pore size Spin-X centrifugal tube filter, followed by 10 K molecular mass cutoff filters, followed finally by 3 K molecular mass cutoff filters. After each filtration step, filters were rinsed with 50 μL water, which was added to the total sample volume for subsequent steps. Filtered extracts were dried again and resuspended in 200 μL water for LC-MS/MS analysis.

Protein extraction and immunoblot analysis

Affinity-purified polyclonal antibodies to CreTPT3 were custom-made by GenScript. The antigen sequence used for antibody generation was KSWSFGRPVTKQEF. Chlamydomonas cells were grown in liquid cultures to 2–5 x10^6 cells ml^{-1} and collected by centrifugation (1459 x g, 5 min). Cells were resuspended in resuspension buffer [(5 mM HEPES, pH 7.5, KOH), 10 mM EDTA (pH 7.5, NaOH), 1x protease inhibitor "complete EDTA-free" (Roche)]. For the disruption of cells, bead-beating was performed using a mini beadbeater (Biospec) in two cycles of 30 sec min^{-1}, with a 1 min period of cooling on ice between cycles. Disrupted cells were centrifuged at 4°C, 30 min, 14000 x g in a microfuge. The supernatant was removed, the pellet was resuspended in sample buffer (resuspension buffer supplemented with 100 mM Na_2CO_3, 100 mM DTT, 2% WT/vol SDS, and 12% WT /vol sucrose).

SDS-PAGE was performed using a 12% polyacrylamide gel (Bio-Rad), with the electrophoresis for 90 min at 120 V. Next, proteins were transferred onto a PVDF membrane (Bio-Rad) using Trans-Blot Turbo (Bio-Rad). The detection was performed using Clarity Max Western ECL Substrate (Bio-Rad).

RT-qPCR

RT-qPCR, was performed as described previously(Kaye et al., 2019). Briefly, total RNA was isolated using the RNeasy Plant Mini Kit (Qiagen) and treated with DNase I (Qiagen). First-strand cDNA was generated by reverse transcription of 0.5 µg total RNA using the iScript cDNA Synthesis Kit (Bio-Rad). Real-time PCR using the Roche Light Cycler 480 was performed with the SensiMix NO-Rox SYBR Green I Kit as described by the manufacturer (Bioline). Oligos used in this research for analyzing the expression levels of Cre*TPT10/PPT2/TPT3/CGL51* are listed in **Supplemental Table 1**.

Analysis of transcript levels in response to different conditions

In order to analyze the transcript level of Cre*TPT* genes under different conditions, we obtained RNAseq data published previously in which cells experienced a diurnal cycle, nitrogen starvation, sulfur starvation and Fe^{2+} starvation(González-Ballester et al., 2010; Ngan et al., 2015; Strenkert et al., 2019; Urzica et al., 2013; Zones et al., 2015). The log2 fold change of each gene was calculated based on the ratio of transcript abundance at different time point over transcript level at time point 0 h. The fold change was visualized by a heatmap and the color bar is the scale of log2 (fold-change).

In vivo polymeric carbohydrate staining

Lugol's iodine (aqueous solution of 1.8% iodine and 3.0% potassium iodide) was used to stain in vivo polymeric carbohydrate. Briefly, 0.5 mL of the cell culture was pelleted, resuspended in 10 µL of Lugol's iodine and visualized under a bright-field microscope. Lugol's iodine selectively binds to alpha-1,4 glucans and stains it blue-black.

oligo name	sequence
gTPT3_pRam118_f	tactcacaacaagcccagttATGGCTTCGCTCGCTCAGCGCGTGAGC
gTPT3_pRam118_r	agcggggatcctatccgttTTAGGCCGCGGCCGGCGACGC
TPT3pro1000_f	gccttttgctcacatgtggcggccgccgccatgctggccaacag
TPT3pro1000_r	taccaatccacaaatgtgggctagcccatgtgaaggcgttgacc
TPT3_3UTR_f	gatgacgacaagtaaagatctTAAGGCGGGCAGCGGC
TPT3_3UTR_r	GCGCAAGAAAGAAGCTTGATATCCCCCTTATAATGTTTACAAGTGGTGCG
hygro_crispr_f2	GGAACTGGCGCAGTTCCTCTG
hygro_crispr_re2	GGCTCGCGTAGGAATCATCCG
TPT3_gRNA1_F	GCTCGGTTGAGCGCACCC
TPT3_gRNA1_R	CAAGGGTCTGGCGAGCACG
TPT3_gRNA2_F	ACCTCCCCATCGCCACGAT
TPT3_gRNA2_R	AGGGGTAAGGACAGCGGTG
Qrt_TPT10_f2	CCTGTACCAGCAGGTGTCGTAC
Qrt_TPT10_r2	CCAGCGCGTTCTGGAGTGACA
Qrt_PPT2_f2	ATGGCGTCTCTGCTGGGC
Qrt_PPT2_r2	CTGCTCGCCCATTCTCCTC
Qrt_TPT3_f2	GCAAAGCTGGCGCCTACTG
Qrt_TPT3_r2	TAGACCAGGCCGACCACCAC
Qrt_CGL51_f1	CTGGTTCATCGGGACATGGC
Qrt_CGL51_r1	TGGCTGAAGCTGACCACT

1318

Supplementary Table 1: Oligonucleotide list.

1320

ID	Construct	Restriction enzyme for transformation	Localization
pMBS472	PSADpro::aadA::PSAD-T:::: ARpro::roGFP2-Tsa2ΔCR::MStop::RPL23-T	ECoRV/Afel	cytosol
pMBS652	PSADpro::aadA::PSAD-T:::: ARpro::CDJ1::roGFP2-Tsa2ΔCR::MStop::RPL23-T	ECoRV/Afel	stroma
pMBS657	PSADpro::aadA::PSAD-T:::: ARpro::roGFP2-Tsa2ΔCR::SV40::RPL23-T	ECoRV/Afel	nucleus
pMBS655	PSADpro::aadA::PSAD-T:::: ARpro::HSP70C::roGFP2-Tsa2ΔCR::MStop::RPL2	ECoRV	mitochondrion

1321

Supplementary Table 2: Plasmids of H$_2$O$_2$ roGFP2-Tsa2ΔCR sensors (Niemeyer et al., 2021)**.**

1323

Cre ID	Name from V6.1 / (name used in this study)	Previous name	Prediction	CS Position
Cre08.g379350_4532	TPT10 / (TPT10)	TPT1	cTP	CS pos: 61-62. TCL-AV. Pr: 0.3994
Cre06.g263850_4532	N / (PPT2)	PPT2; TPT2; TPT20	cTP	CS pos: 70-71. VCQ-AA. Pr: 0.5253
Cre01.g045550_4532	TPT3 / (TPT3)	TPT3	cTP	CS pos: 60-61. VTK-AS. Pr: 0.4178
Cre02.g106200_4532	N	TPT4; TPT5	OTHER	
Cre02.g112900_4532	N	TPT5; TPT6	OTHER	
Cre02.g144300_4532	N	TPT6; TPT7	OTHER	
Cre03.g162000_4532	N	TPT7; TPT8	OTHER	
Cre03.g184850_4532	N	TPT8; TPT9	OTHER	
Cre04.g227450_4532	N	TPT9	OTHER	
Cre07.g330850_4532	N	TPT10; TPT11	OTHER	
Cre08.g363600_4532	N	TPT11; TPT12	OTHER	
Cre09.g408400_4532	N	TPT12; TPT13	OTHER	
Cre09.g413700_4532	N	TPT13; TPT14	OTHER	
Cre08.g382350_4532	EZY14	TPT14; TPT16; EZY14	OTHER	
Cre09.g415900_4532	MOT20	TPT15	OTHER	
Cre10.g452750_4532	N	TPT16; TPT18	OTHER	
Cre11.g479950_4532	CGL7	TPT17	OTHER	
Cre12.g490050_4532	N	TPT18; TPT19; GMT1	OTHER	
Cre12.g490100_4532	TPT2	TPT19;GMT2	OTHER	
Cre12.g501000_4532	N	TPT20; TPT22; PPT1	OTHER	
Cre15.g641266_4532	N	TPT22; TPT23	OTHER	
Cre15.g642950_4532	N	TPT23; TPT24	OTHER	
Cre15.g643385_4532	N	TPT24; TPT26	OTHER	
Cre16.g663800_4532	CGL51 / (CGL51)	TPT25	cTP	CS pos: 70-71. IVA-SS. Pr: 0.4817
Cre16.g666250_4532	N	TPT26; TPT27	OTHER	
Cre17.g702700_4532	N	TPT27; TPT28	OTHER	
Cre17.g703250_4532	SLC35D	TPT28; TPT29	OTHER	
Cre17.g710850_4532	N	TPT29; TPT4	OTHER	
Cre11.g467754_4532	N	UAA6	OTHER	
Cre14.g622700_4532	N	N	OTHER	
Cre18.g748947_4532	N	N	OTHER	
Cre09.g408428_4532	N	N	OTHER	

1324

1325 **Supplementary Table 3: Members of the triose phosphate transporter family in *Chlamydomonas***
1326 ***reinhardtii* and their predicted subcellular localization based on TargetP-2.0.** cTP, chloroplast transit
1327 peptide; CS position, transit peptides cleavage site; N: genes without names in the genome. Cre IDs are
1328 from v6.1 genome.

1329

1330 **Supplementary Fig. 1: Experimental set up.** All strains were grown photoautotrophically under low
1331 white light (LL, 30 µmol photons m^{-2} s^{-1}) for 16-24 h. After dilution with fresh medium, the cultures were
1332 transferred to moderate light (ML, 250-300 µmol photon m^{-2} s^{-1}) for 24 or 48 h. On occasion, the cultures
1333 were also transferred to the high light (~450 µmol photon m^{-2} s^{-1}). To avoid self-shading, the culture density
1334 was adjusted to 1-1.5 µg/mL chlorophyll before exposure to ML or HL.

1335

1336

Supplementary Fig. 2: Alignment of putative plastid phosphate transporter sequences of *Arabidopsis*
thaliana* and *Chlamydomonas reinhardtii. Plastid putative sugar phosphate transporters from
Chlamydomonas are CreTPT10 (Cre08.g379350_4532), CrePPT2 (Cre06.g263850_4532), CreTPT3
(Cre01.g045550_4532), and CGL51 (Cre16.g663800_4532). Sequences from *Arabidopsis thaliana* are
AtTPT1 (AT5G46110.1), AtPPT1 (AT5G33320.1), AtGPT1 (AT5G54800.1) and AtXPT1
(AT5G17630.1). Conserved substrate binding pockets are colored red. Phosphate-binding residues are
colored blue. The gate capping residues are colored yellow. The positions of residue in green are crucial
for the preference of triose-phosphate or 3-PGA as substrate. The purple residues are critical for the
specificity of PEP.

1346

1347

Supplementary Fig. 3: Background activity of Pi uptake derived from the endogenous yeast
transporters reconstituted into liposomes. Time-dependent uptake of Pi (0.25 mM) into reconstituted
liposomes preloaded with 30 mM of Pi (■) or without exchange substrate (●) prepared from yeast cells
harboring the empty vector (pYES-NTa). The arithmetic mean (±SD) of three independent experiments
(each with three technical replicates) was plotted against time. The observed Pi uptake rates of endogenous
yeast carriers reconstituted into liposomes were negligible compared to the rates of His-CreTPT3 (20-fold
higher, **Fig. 1C**), indicating that this expression system is suitable for functional analysis of recombinant
phosphate transporters.

1356

1357

1358 **Supplementary F**

1359 resistant marker (*A*

1360 (*t3ko1*, *t3ko2*, *t3ko*

1361

1362

Supplementary Fig. 5: (**A**) Cell size of WT, *t3ko2*, and *C_T3KO2* in TP in both LL and ML. (**B**) WT and *t3ko2* morphology 24 h following a LL to ML transition.

1365

Supplementary Fig. 6: Quantification of chlorophyll contents in indicated strains after exposure to ML for 24 and 48 h. All strains were grown photoautotrophically in LL for 16-24 h. After dilution with fresh medium, the culture was transferred to ML and the chlorophyll content determined after 24 and 48 h in the ML. The initial inoculum density used for the transition was between 1-1.5 µg/mL chlorophyll in order to avoid self-shading.

1371

1372

Supplementary Fig. 7: Starch and metabolites in WT and *t3ko2*. (**A**) Starch content in indicated strains following 0, 6, and 24 h in LL. (**B**) Time course, over a period of 6 sec, of change in metabolite pool sizes at 0 h (LL), 45 min and 6 h after shifting LL-grown cells to ML. Data was normalized to cell numbers. (**C**) Schematic representation of the fold-change for the metabolites shown in (**B**), calculated by dividing the pool size in *t3ko2* by that of WT cells under the same conditions. Fold change of 1 (no change) is shown by a dashed line. Each data point is the mean and standard error of three biological replicates. Asterisk indicates statistically significant differences relative to WT (* $P<0.05$, ** $P<0.01$, *** $P<0.001$). Abbreviations: G6P, glucose-6-phosphate; FBP, fructose bisphosphate; DHAP, dihydroxyacetone phosphate; 3-PGA, 3-phosphoglycerate; PEP, phosphoenolpyruvate; CIT, citrate; SUCC, succinate; FUM, fumarate; MAL, malate; S7P, sedoheptulose-7-phosphate; E4P, Erythrose 4-phosphate; R5P, Ribose 5-phosphate; RU5P/X5P, ribulose 5-phosphate/xylulose-5-phosphate.

1384

1385

Supplementary Fig. 8: (**A**) mRNA levels encoding putative chloroplast envelope localized TPTs under LL during growth in TP medium. mRNA levels of CreTPT10 (**B**), CrePPT2 (**C**) and CGL51 (**D**) in the indicated strains 0, 2, 6 h after the cells were shifted from LL to ML. (**E, F**) mRNA levels of plastidial malate dehydrogenases (MDH) in the indicated strains 0, 6 h after the cells were shifted from LL to ML.

1386

1387
1388
1389
1390

1391

1392

Supplementary Fig. 9: Expression of the putative chloroplast envelope localized TPTs under nitrogen
starvation **(A)**, sulfur starvation **(B)**, and iron starvation **(C)** (González-Ballester et al., 2010; Ngan et al.,
2015; Urzica et al., 2013).