Chlamydomonas reinhardtii triose-phosphate/phosphate translocator3 (TPT3): A major

chloroplast shunt for the export of fixed carbon and reductant

Weichao Huang[a1+], Anagha Krishnan[b1], Anastasija Plett[c], Michelle Meagher[d], Nicole Linka[c], Bijie

Ren[a], Justin Findinier[a], Petra Redekop[a], Neda Fakhimi[a], Jin Xu[e], Rick G. Kim[a], Devin A. Karns[b],

Nanette Boyle[d], Matthew C. Posewitz[b], Arthur R. Grossman[a+]

[a]Department of Plant Biology, Carnegie Institution for Science, Stanford, CA 94305, USA

[b]Department of Chemistry and Geochemistry, Colorado School of Mines, Golden, CO 80401,

USA

[c]Institute of Plant Biochemistry, Heinrich-Heine University Düsseldorf, 40225 Düsseldorf,

Germany

[d]Department of Chemical and Biological Engineering, Colorado School of Mines, Golden, CO

80401, USA

[e]Department of Microbiology and Cell Science, Institute of Food and Agricultural Sciences (IFAS),

University of Florida, Lake Alfred, 33885, FL, USA

[1]Authors contributed equally.

[+]Corresponding Author: Weichao Huang; Address: The Carnegie Institution for Science,

260 Panama St. Bldg. 100; Stanford, CA 94305; email: whuang@carnegiescience.edu

[+]Corresponding Author: Arthur R. Grossman; Address: The Carnegie Institution for Science,

260 Panama St. Bldg. 100; Stanford, CA 94305; email: agrossman@carnegiescience.edu

Abstract

Photosynthetic eukaryotic organisms contain several chloroplast-associated metabolite

transporters that enable energetic/metabolic exchange between the chloroplast and other cellular

compartments. In this study, we used the model photosynthetic alga *Chlamydomonas reinhardtii*

to investigate a highly expressed chloroplast triose phosphate transporter. The triose

phosphate/phosphate translocator 3 (CreTPT3), located on the Chlamydomonas chloroplast

envelope, was found to be highly expressed under both non-stressed/stressed conditions (RNA

level) and was characterized for substrate specificity in vitro using a yeast liposome uptake system.

The CreTPT3 transporter showed high DHAP and 3-PGA transport activities, but little activity

with PEP. Null mutants for CreTPT3, generated by CRISPR-Cas9 editing of the Cre*TPT*3 gene,

resulted in a pleiotropic phenotype impacting photosynthetic activity, metabolite pools, carbon

partitioning and storage, the redox status of the chloroplast, and the accumulation of reactive

oxygen species. The results presented demonstrate that CreTPT3 is a major conduit on the

chloroplast envelope for the intracellular distribution of fixed carbon and reductant generated by

photosynthetic electron transport. Its function is critical for optimizing the use of resources

supporting cell fitness, especially as light intensities increase, the rate of photosynthetic CO_2

fixation is elevated and the chloroplast environment becomes highly reducing.

Introduction

Photosynthetic organisms have evolved numerous strategies to distribute fixed carbon among the different cellular compartments, maintaining a dynamic metabolic/redox balance that fuels growth and biosynthetic processes. However, photosynthetic organisms can also absorb excess, potentially damaging levels of light energy during mid-day (when photosynthesis becomes saturated) and can suffer from extreme damage, especially when they experience rapid fluctuations in light intensities or are subjected to nutrient limiting conditions and other environmental stresses that impair the productive utilization of excitation energy (Chaux et al., 2017; Saroussi et al., 2017). To cope with excess excitation energy, plants and algae have evolved mechanisms to dissipate this energy through nonphotochemical quenching (NPQ), which involves both energy dependent quenching (or qE) that dissipates the energy as heat within light harvesting complexes of photosystem II (PSII), and a state transition (or qT), which results in the movement of the light harvesting antennae complexes from PSII to photosystem I (PSI) where the energy can be dissipated within the PSI reaction center. Excess reductant can also be eliminated by photochemical quenching. While the dominant form of photochemical quenching often involves the use of the electrons/reductant to fix inorganic carbon (Ci), which is used for growth or stored in the form of starch or lipids (Ge et al., 2014; Huang et al., 2018; Krishnan et al., 2015), redox equivalents can also be trafficked to other outlets where they are involved in 'nonproductive' photochemistry. The main alternative photochemical electron outlets involve the reduction of O_2 in H_2O-to-H_2O cycles that include a (*i*) Mehler-type reaction (Badger et al., 2000) that is noncatalytic and functions on the acceptor side of PSI, (*ii*) flavodiiron protein reactions (Jokel et al., 2018) (FLVs, NADPH:flavin oxidoreductase) that catalytically reduces O_2 on the acceptor side of PSI without generating reactive oxygen species (ROS), (*iii*) the plastid terminal oxidase (Houille-Vernes et al., 2011) (PTOX, plastoquinol:oxygen oxidoreductase) reaction that can use electrons of the plastoquinone (PQ) pool to reduce O_2 (thought to maintain/tune the redox state of the photosynthetic electron transport (PET) chain), and (*iv*) the movement of redox equivalents from the chloroplast to the mitochondrion where they can be used to reduce O_2 through various electron transport activities.

The extent and characteristics of metabolite and reductant shuttling between the chloroplast and other cellular compartments have not been extensively characterized despite its importance to cellular metabolism. The chloroplast is delineated by a double envelope membrane that contains

several metabolite translocators that are part of the machinery that governs metabolic trafficking

among the different cellular compartments (Facchinelli and Weber, 2011; Haferkamp and

Schmitz-Esser, 2012). Various mechanisms have been proposed in the shuttling of fixed carbon

and reducing equivalents between the chloroplast and cytoplasm. Studies with *Arabidopsis*

thaliana revealed that the malate shuttle, which involves multiple malate dehydrogenases (MDH)

and malate/OAA translocators (OMT), functions in the export of reductants from the chloroplast

and also enables the cells to manage redox conditions in the chloroplast (Zhao et al., 2018, 2020).

The exported malate can be used for anabolic processes or to generate NADH (during the

conversion of malate to oxaloacetate), which can be delivered to the mitochondrion to drive

oxidative phosphorylation and ATP synthesis (Zhao et al., 2020) or diverted to the mitochondrial

alternative oxidase pathway where it reduces O_2 to H_2O without (or with little) generation of ATP

(Kaye et al., 2020). The buildup of reducing equivalents in the mitochondrion or chloroplast can

trigger the production of ROS, which potentially activates signaling systems that modulate energy

metabolism and, under extreme conditions, could elicit programmed cell death (PCD) (Dietz et al.,

2016; Kuźniak and Kopczewski, 2020; Li and Kim, 2022; Zhao et al., 2018).

In addition to the malate shuttle, reduced carbon can be exported from the chloroplast by the triose

phosphate (triose-P)/phosphate (Pi) translocator (TPT), which can also serve potentially as a means

of exporting carbon from the chloroplast and a safety valve for eliminating excess reducing power

from the chloroplast and lessening the potential for ROS accumulation and damage (Fliege et al.,

1978; Flügge et al., 1989). The TPTs can potentially function as a major path for exchanging

metabolites between the chloroplast and cytosol. Triose-Ps are generated during photosynthetic

CO_2 fixation by Calvin-Benson-Bassham cycle (CBBC) activity that is supported by the reducing

power/energy (NADPH, ATP) derived from photosynthetic electron transport. The TPTs reside

on the inner chloroplast envelope membrane and can export triose-Ps, including glyceraldehyde

3-phosphate (G3P/GAP), dihydroxyacetone phosphate (DHAP), and 3-phosphoglycerate (3-PGA)

across the chloroplast envelope membrane and into the cytosol in a 1:1 counter exchange for Pi

(Fliege et al., 1978; Flügge et al., 1989; Lee et al., 2017a). Various plants like tobacco and

Arabidopsis, do not exhibit a strong phenotype if TPT is either eliminated or its level is reduced

(Häusler et al., 1998; Schneider et al., 2002; Walters et al., 2004). This lack of TPT activity can be

compensated for by moving the assimilated carbon into a transitory starch pool that is subjected to

accelerated turnover in the light (Häusler et al., 1998; Riesmeier et al., 1993; Walters et al., 2004),

leading to the production of hexoses that can be exported from the chloroplast. Characterization of a double mutant (*tpt-1/adg1-1*) of Arabidopsis null for TPT (*tpt-1*) and starch synthesis (*adg1-1*) exhibited severely compromised photosynthetic capacity and growth (Heinrichs et al., 2012; Schneider et al., 2002). However, there are still few studies that demonstrate the functionalities of TPT proteins in single-celled photosynthetic organisms (red algae, apicomplexans, cryptophyte) by analyzing the activity of these proteins using liposome transport assays (Haferkamp et al., 2006; Linka et al., 2008; Moog et al., 2020), the impact of null mutations in the gene encoding the TPT on the physiology of the cells has not been investigated so far.

In this study, we used the unicellular green alga *Chlamydomonas reinhardtii* (Chlamydomonas throughout), a well-established model organism that has been extensively used to dissect various physiological processes, including photosynthesis and acclimation to environmental change. Chlamydomonas contains more than 30 TPT homologs; the informatic analyses of these proteins are still not refined enough to be certain of their substrate specificities and, in many cases, the transported metabolite is likely not a triose-P. Here we present work that approaches the questions: *i*) which of the TPTs is a triose-P/Pi translocator associated with the chloroplast envelope; *ii*) what are the roles of chloroplast TPTs in partitioning carbon and reductant in cells and sustaining central carbon metabolism; and finally, *iii*) do the TPTs contribute to dissipating excess reducing equivalents and balancing cellular redox conditions? Our study is showing for the first time the impact of *tpt* null mutations on the physiology of algal cells.

We have focused our analyses on Chlamydomonas TPT3 (CreTPT3) which, of the four predicted chloroplast-associated CreTPTs, exhibits the strongest sequence similarity to plant chloroplast TPTs and is highly expressed over the diel cycle, especially during the light period. We demonstrate that CreTPT3 is located on the chloroplast envelope. Expressing CreTPT3 in yeast and reconstituting it into liposomes allowed us to show that this transporter has a strong preference for DHAP (dihydroxyacetone phosphate) and 3-PGA (3-phosphoglycerate) in Pi exchange assays. We generated null mutants for the Cre*TPT3* gene, which exhibited growth attenuation even in low light and, over the long term, were unable to survive the transition to moderate/high light. Growth, along with the various other mutant phenotypes, were rescued by ectopic expression of the Cre*TPT3* gene in *tpt3* mutant cells. We also examined photosynthetic performance, performed metabolite analyses, measured accumulation of carbon storage products (starch and lipids), and analyzed subcellular redox changes and ROS accumulation in wild type (WT) cells and the

137   Chlamydomonas *tpt3* mutant. Our results demonstrated that CreTPT3 is essential for the exchange

138   of fixed carbon, sustaining central carbon metabolism and maintaining intracellular redox balance.

Material and methods

Strains and culture conditions

WT Chlamydomonas M10 (CC-4403, isogenic line derived from CC-124) was used as the parental

strain for the generation of knockout mutants. Cultures were routinely cultivated in growth

chambers (LED-41L2, Percival Scientific, Inc.) at 25°C with continuous shaking on an orbital

shaker (VWR OS-500 Shaker) at 120 rpm, in Tris-acetate-phosphate (TAP) medium (Harris, 2009).

The cultures were illuminated with continuous cool white LEDs at low light (30 μmol photons m^{-2}

s^{-1}). Experiments were mostly performed with cells grown in the photoautotrophic medium TP

(TAP medium without acetate), and in TAP medium in some cases, at 25°C, and sparged with air

while being shaken at 120 rpm in a growth chamber (LED-41L2, Percival Scientific, Inc.). For

growth assays, cultures were inoculated at a density of 0.02 at OD$_{750\,nm}$ (~1×10^5 cells/mL) in TP

sparged with air under either LL or moderate light (ML, 250-300 μmol photons m^{-2} s^{-1}) intensities.

For spectrophotometric and chlorophyll fluorescence analyses, precultures were grown in TAP in

LL to mid-exponential phase, the cells were pelleted by centrifugation at 800x g, for 3 min at room

temperature, and the pellets were washed once with TP and then resuspended in TP at a density of

1-1.5 μg/mL chlorophyll. These cultures were grown in TP sparged with air and after 16-24 h in

LL, were transferred to ML. To avoid self-shading, the culture density was adjusted to 1-1.5 μg/mL

chlorophyll before ML exposure.

Cell samples exposed to the different light intensities were diluted to perform spot tests for

photoautotrophic growth on TP agar plates and for mixotrophic growth on TAP agar plates. Agar

plates were incubated for 7 d under either LL or ML (cool white LED) at 25°C.

Phylogenetic analysis

161   Protein sequences were aligned using MUSCLE (Edgar, 2004). The best-fit nucleotide substitution

162   model was generated based on the Akaike information criterion (AIC) using MEGA7 (Kumar et

163   al., 2016). Maximum likelihood phylogenetic trees were constructed based on the best-fit model

164   GTR (general time reversible) + G (gamma distribution) + I (invariable sites) with 100 bootstrap

165   replicates using MEGA7 (Kumar et al., 2016).

Expression of CreTPT3 in yeast

The DNA sequence encoding the mature CreTPT3 protein was codon-optimized for expression in *Saccharomyces cerevisiae* (GeneART, ThermoFisher Scientific). This coding sequence was inserted in frame with an N-terminal His tag into the yeast vector pYES-NTa (ThermoFisher Scientific) using Gibson cloning (NEB). Briefly, pYES-NTa was linearized with BamHI and the CreTPT3 cDNA amplified with the primer pair fwd:5'-gacgataaggtacctagcGCTTCTGCTGCTGATGCTC-3' and rev:5'-agaattccaccacactgTTAAGCGGCAGCTGGAGAAG-3'. This cDNA was ligated into the linearized pYESS-NTa vector, which was then transformed into the yeast strain INVSc1 (MATa, his3D1, leu2, trp1-289, ura3-52 / MATα, his3D1, leu2, trp1-289, and ura3-52, ThermoFisher Scientific) using the lithium-acetate/PEG method (Gietz and Schiestl, 2007). Transformed yeast cells were selected on synthetic complete medium containing 2% (w/v) glucose with the uracil auxotrophic marker. Galactose-inducible expression of His-CreTPT3 in yeast was performed as described in (Linka et al., 2008). The presence of the His-tagged fusion protein was verified by standard SDS-PAGE and immunoblot analysis using an anti-His-antibody conjugated with horseradish peroxidase (Miltenyi Biotech).

Reconstitution into liposomes and transport activity assays

For uptake studies, yeast membranes from cells with and without recombinant His-CreTPT3 were enriched and reconstituted into 3% (w/v) L-alpha-phosphatidylcholine using a freeze-thaw-sonication procedure (Linka et al., 2008). The reconstituted liposomes were preloaded with 30 mM Pi or the indicated phosphorylated metabolites that were tested as potential transport substrates. As a negative control for antiport activity, liposomes were also generated without preloading. The external counter-exchange substrate was removed via gel filtration on Sephadex G-25M columns (GE Healthcare). Transport assays were initiated by adding 0.25 mM [alpha-^{32}P]-phosphoric acid (6,000 Ci/mmol) and performed as previously described in Linka et al. (2008). The Km for Pi was determined by measuring the initial velocity at each of six external Pi concentrations between 0.05 and 5 mM. To obtain competitive inhibition constants (Ki), the uptake of 0.25 mM Pi into liposomes containing 30 mM Pi was measured over a 4 min period in the presence of increasing external inhibitor concentrations (0.05 - 5 mM). Three biological replicates were performed for all

196   described experiments. GraphPad Prism software version 9.3.0 was used for non-linear regression
197   analyses of the kinetic data.

Vector construction, transformation and complementation

The pRam118_VENUS plasmid which harbors the VENUS gene and the *AphVII* cassette (confers
resistance to hygromycin) (Cre01.g045550) (Kaye et al., 2019), was used to express Cre*TPT3*
(Cre01.g045550). This plasmid was linearized with HpaI (NEB). Primers: g*TPT3*_pRam118_f and
g*TPT3*_pRam118_r (**Supplementary Table 1**) were used to amplify Cre*TPT3* genomic DNA
containing an overlap with the linearized pRam118_VENUS vector. For generating the plasmid
pRam118-CreTPT3&VENUS, genomic DNA of Cre*TPT3* was assembled with
pRam118_VENUS plasmid using Gibson assembly (Gibson et al., 2009). A 1,000 bp sequence
upstream of Cre*TPT3* and containing the promoter region of the gene was amplified using the
primers TPT3pro1000_f and TPT3pro1000_r (**Supplementary Table 1**). The 3' UTR of Cre*TPT3*
was amplified using TPT_3UTR_f and TPT3_3UTR_r (**Supplementary Table 1**). The *PSAD*
promoter and the *RBCS2* 3'UTR of pRam118-CreTPT3&VENUS were replaced by the amplified
fragments of the 1,000 bp upstream region and the 3' UTR of Cre*TPT3*, respectively. This final
vector, designated pRam118_CreTPT3, contains the original Cre*TPT3* promoter (driving
expression of Cre*TPT3*), the genomic DNA sequence of Cre*TPT3*, VENUS and the Cre*TPT3* 3'
UTR plus the *AphVII* cassette. The Chlamydomonas *tpt3* mutant was transformed by
electroporation with 2 μg of pRam118_CreTPT3 that was linearized with AseI. The GeneArt MAX
Efficiency Transformation Reagent for algae (Invitrogen) was used for introducing the plasmid by
electroporation following the instructions provided by the manufacturer. Transformants were
selected on solid TAP medium containing hygromycin (10 μg/mL; Enzo Life).

Subcellular localization

Drug resistant transformants were visualized for VENUS fluorescence as previously described
(Kaye et al., 2019). In brief, transgenic cell lines resistant to hygromycin were screened for
VENUS fluorescence using a microplate reader (Infinite M1000; TECAN). Excitation and
emission settings were as follows: VENUS, excitation 515/12 nm and emission 550/12 nm;
chlorophyll excitation 440/9 nm and emission 680/20 nm. The TCS SP8 confocal laser-scanning
microscope (Leica) was used to visualize the VENUS fluorescence signal. Detailed methods for
visualization can be found in (Kaye et al., 2019).

CRISPR-Cas9 mediated mutagenesis

The protocol for disruption of Cre*TPT3* (Cre01.g045550) was adapted from (Findinier et al., 2019). Chlamydomonas WT strain CC-124 was used for mutant generation. WT cells were cultured under continuous light (50 µmol photons m^{-2} s^{-1}) for 2 d to a density of 3-5×10^6 cells/mL. The cells were then concentrated to 2×10^8 cells/mL in 0.5×TAP supplemented with 80 mM sucrose. Two single guide RNAs (sgRNA) were designed by CHOPCHOP and synthesized by Integrated DNA Technologies (IDT). The sequences of the generated sgRNA are: *TPT3*-sg1 (5′-CGCUGGGCGTCACUUCCCGGCGG-3′) for editing of exon 1 and *TPT3*-sg2 (5′-AAGGCCGCUAUCGCCAACGUGGG-3′) for editing of exon 7. Prior to electroporation, Cas9 (IDT) and sgRNAs were incubated together at 37 °C for 30 min. Approximately 500 ng PCR product of the *AphVII* cassette, which confers resistance to hygromycin, was added to the RNP (ribonucleoprotein) mixture. 250 µl aliquots were electroporated using Super Electroporator NEPA21 type II (NEPA GENE). After 16 h of recovery in very low light (10-15 µmol photons m^{-2} s^{-1}), cells were plated onto solid TAP medium containing 10 µg/mL hygromycin. Sense or antisense-oriented knock-ins of *AphVII* were determined by amplification using primer pairs with one primer annealing to the genomic sequence and the other to the inserted sequence (**Supplementary Table 1**). Amplified fragments were verified by sequencing (ELIM BIOPHARM, Hayward, USA).

Complementation of mutants of Cre*TPT3*

Mutant strains were transformed with the linearized plasmid pRam118_CreTPT3 plasmid. After transformation, transformants were selected in ML and then screened for VENUS fluorescence. Colonies containing VENUS and an *AphVII* cassette knock-in at the Cas9 targeted site were examined for accumulation of the CreTPT3 protein by western blots using CreTPT3 antibody generated by GenScript USA Inc (Piscataway, USA). Clones of Western-positive cells were subjected to growth complementation assays using spot tests under ML on solid TAP or TP medium.

P700 activity measurements

Absorbance spectroscopy for P700 analyses was performed in dark adapted liquid cultures (15 µg/mL chlorophyll, in 20 mM HEPES-KOH, pH 7.2, and 10% ficoll) using a JTS-100

spectrophotometer (SpectroLogiX, TN). Actinic light was provided by an orange LED (165 μmol

photons m^{-2} s^{-1}) for PSI oxidation, followed by a saturating pulse and a dark incubation. 20 μM

DCMU and 1 mM hydroxylamine were added to the cell suspension to inhibit linear electron flow

(LEF) prior to the measurement. P$_{700}$ activity was measured by monitoring the absorbance at 705

nm (interference filter 6 nm FWHM was used to create a narrow excitation beam); the absorbance

at 740 nm was used to correct for an unspecific contribution to the 705 nm signal.

Chlorophyll fluorescence analysis

262   Chlorophyll fluorescence, used to evaluate photosynthetic electron transport, was measured using

263   a DUAL PAM-100 fluorometer. Cells were adapted in the dark for 20 min. Light curves were

264   generated by illuminating cells with increasing light intensities (0, 10, 50, 100, 200, 400, 800, 1000,

265   1200 μmol photons m^{-2} s^{-1}) for 2 min at each intensity. 1 mM CO_2 (NaHCO$_3$) was added to the

266   reaction mix as an electron receptor for the CBBC.

ROS measurements and roGFP2 imaging analysis

All strains were grown photoautotrophically under LL for 16-24 h. After dilution with fresh

medium, the cultures were transferred to ML or the dark for >48 h and then stained with CM-

H2DCFDA (Thermo Fisher Scientific) as described in (Kong et al., 2018) for detecting ROS.

Briefly, after the treatment (e.g. 48 h in ML), approximately 10 million Chlamydomonas cells

(treated in various ways) were pelleted and washed once with 1×PBS. The cells were then

resuspended in 1×PBS containing 8 μM CM-H2DCFDA and incubated at room temperature in the

dark for 30 min. Following this incubation, the cells were washed three times with 1×PBS buffer,

the fluorescent signals were either visualized using TCS SP8 confocal laser-scanning microscope

(Leica) or quantified with a microplate reader (Infinite M1000; TECAN). Excitation and emission

settings for the microscope were: 488 nm/510-530 nm HyD SMD hybrid detector for ROS, and

488 nm/650-700 nm-HyD SMD hybrid detector for chlorophyll autofluorescence. Excitation and

emission settings for the plate reader were as follows: ROS excitation 488/5 nm and emission

530/12 nm; chlorophyll excitation 514/5 nm and emission 690/5 nm.

Constructs containing chloroplast or mitochondria targeting sequences (Crozet et al., 2018) fused

to codon-optimized roGFP2 (Vevea et al., 2013) were transformed into WT and mutant strains.

Cells were cultured as shown in **Supplementary Fig. 1**. Signals from the transformed lines were

visualized using a TCS SP8 confocal laser-scanning microscope (Leica). Using the sequential

setup of the SP8, roGFP2 signals were collected at the emission wavelength of 510-550 nm upon

excitation of roGFP2 at 405 nm and 488 nm (Vevea et al., 2013). The degree of roGFP2 oxidation

was analyzed as the ratio of the emission signals after excitation at 405 nm and 488 nm (Vevea et

al., 2013).

Photosynthesis-Irradiance curve

Photosynthesis-Irradiance curves were measured using a custom Pt-Ag/AgCl polarographic

electrode system (ALGI) with a water jacketed (for temperature-control), 1 mL glass reaction

chamber. The YSI 5331A electrodes (Yellow Springs Instruments) were polarized at -0.8 V. The

cultures used for the assays were concentrated to 2.5-5 µg mL^{-1} chlorophyll, supplemented with

15 µL of 0.5 M sodium bicarbonate in water and then purged with 1% CO_2/99% He. Using a gas

tight syringe, the sample was transferred into the reaction chamber that was also purged with 1%

CO_2/99% He. The rate of change in O_2 levels was measured sequentially at the light intensities

100, 200, 600, 1000 µmol photons m^{-2} s^{-1} (photosynthetic active radiation, PAR); each intensity

was maintained for 3-5 min followed by a 3-min intervening dark period, and then the light level

was raised to the next higher intensity (stepped change) until the full range of intensities was tested.

At 1,000 µmol photons m^{-2} s^{-1}, the light was held for 10 min followed by an 8-min dark period.

Before the measurement of each experimental series, the electrodes were calibrated with air ($\sim$21%

O_2) and 1% CO_2/99% He mixture (0% O_2). The initial slope of the response was used to determine

the O_2 evolution rate.

Starch and TAG quantification

Cells were grown under LL to mid-exponential phase. Cultures were then diluted to 0.5 µg/mL

chlorophyll with fresh medium and transferred to and then grown in ML for 24 h or more. Starch

was measured using a modified method from (Klein and Betz, 1978). In brief, cells were collected

by centrifugation and bleached with methanol. Dried pellets were dissolved in water and heated at

100 $^{\circ}$C to break the cells and release the starch. After cooling, amyloglucosidase and α-amylase

(2.25 U/mL) were used to hydrolyze the starch and the products of hydrolysis were quantified

using a Glucose Colorimetric Detection Kit (Thermo Fisher).

Intracellular TAG levels were estimated by a fluorometric assay using the dye Nile Red (Thermo
Fisher) (Yu et al., 2009). A Nile Red solution (500 μg/mL in acetone) was added to 1 mL of cell
suspensions at a final concentration of 0.5 μg/mL. The samples were then incubated at room
temperature for 30 min, and the Nile Red fluorescence emission quantified at 575 nm after
excitation at 530 nm using a microplate reader (Infinite M1000; TECAN).

Metabolic analysis

Cells grown at 30 μmol photons m^{-2} s^{-1} (LL) were shifted to 250-300 μmol photons m^{-2} s^{-1} (ML)
for 45 min or 6 h, as indicated. 45 mL of culture was rapidly quenched in cold saline solution (-2
to -3°C), extracted using cold methanol and then analyzed by LC-MS/MS. Quenching and analysis
of metabolites were performed according to (Sake et al., 2020) with modifications. The quenching
solution, filtered saline (9 g/L NaCl), was prechilled to 4°C in a refrigerator. 30 mL of the
quenching solution was transferred to a 50 mL conical tube kept in an ice bath mixed with salt to
depress the temperature to between -3°C and -1°C. 15 mL of the culture was rapidly plunged into
the 30 mL of the quenching solution and the samples were centrifuged at 4,000 rpm for 10 min at
4°C. Cell pellets were washed with fresh prechilled saline solution and centrifuged again at 4,000
rpm for 10 min in a 2 mL centrifuge tube. For each replicate, 45 mL of the culture was sampled
and the cell pellets were pooled. Washed pellets were frozen at -80°C until extraction. Metabolite
extraction and further analysis were modified from (Young et al., 2011). A step-by-step protocol
is described in supplemental information (SI). All chemicals used for metabolite extraction and
LC-MS/MS analysis were Optima grade reagents.

Results

Features of the triose-P/Pi translocator (TPT) family

Candidate genes encoding triose-P/Pi transporters (TPT) in the Chlamydomonas genome were
identified by blasting the Arabidopsis TPT protein to the Chlamydomonas genome. The TPT
family is the largest drug/metabolite transporter (DMT) family in eukaryotes and includes triose-
P and sugar-phosphate permeases that are associated with chloroplasts; many family members are
still functionally uncharacterized (Jack et al., 2001; Knappe et al., 2003; Weber et al., 2006).
Chlamydomonas harbors ~32 genes encoding proteins of the TPT family, with four members
(CreTPT1, 2, 3 and 25) predicted to have a transit peptide and be chloroplast-localized

(**Supplementary Table 2**). To learn about the potential substrate specificities of the four putative chloroplast CreTPTs, their amino acid sequences were aligned with the plastidial phosphate transporters (PT) from Arabidopsis (AtPTs) (**Supplementary Fig. 2**).

Land plants harbor four classes of phosphate translocators on the inner plastid envelope, including a glucose 6-phosphate transporter (GPT), a xylulose phosphate transporter (XPT), a phosphoenolpyruvate transporter (PPT), and a triose-P translocator (TPT) (Eicks et al., 2002; Fischer et al., 1997; Flügge et al., 1989; Kammerer et al., 1998; Lee et al., 2017b). The ability of the transporters to use triose-P/DHAP or 3-PGA is dependent on five highly conserved residues (H184, K203, Y338, K359, and R360 in AtTPT1) (Lee et al., 2017b; Moog et al., 2020). Only CreTPT3 and CreTPT25 contain all five of these residues (H170, K189, Y322, K345, and R346 in CreTPT3) (**Supplementary Fig. 2**, **3**). In CreTPT2 and TPT1, H170 is replaced by N, which might result in distinct substrate preferences (**Supplementary Fig. 2**, **3**). In AtTPT1, the residue F262 (F248 in Chlamydomonas) is thought to inhibit phosphoenolpyruvate (PEP) access to the binding site; this residue is replaced by N in AtPPT1 (PPT) (Lee et al., 2017b; Moog et al., 2020). CreTPT25 has M at position F248, indicating that unlike the other CreTPTs, CreTPT25 might have higher preference for PEP (**Supplementary Fig. 3**). Additionally, these findings suggest that of the four chloroplast TPT family members, CreTPT3 is most likely to function in specifically shuttling triose-P or 3-PGA across the chloroplast envelope.

Figure 1: Chloroplast TPT phylogenies, subcellular localization of CreTPT3 and in vitro measurements of CreTPT3 transport activities. (**a**) Truncated phylogenetic tree of potential chloroplast CreTPTs. (**b**) Cellular localization of CreTPT3. The construct for expression of the CreTPT3-VENUS fusion protein is shown above the cell images and explained in the Materials and Methods. VENUS fluorescence, green; Chlorophyll (Chl) autofluorescence, magenta. (Scale bar: 5 µm). (**c**) Expression of CreTPT3 in yeast. Left, SDS-PAGE gel of proteins from total membranes isolated from yeast cells containing the empty vector (lane 1) or expressing His-CreTPT3 (lane 2). Right, corresponding immunoblot blot detection of recombinant His-CreTPT3 protein using anti-His antibodies. The calculated molecular mass of the N-terminal His-tagged protein was 37 kDa. (**d**) The kinetics of Pi exchange by His-CreTPT3 in lipid vesicles. Uptake of Pi (0.25 mM) into liposomes was measured in the presence (■) or absence (□) of Pi (30 mM) loaded within the vesicles. The arithmetic mean (±SD) of three independent experiments (each consisting of three technical replicates) was plotted with respect to the time after initiating the assay. (**e**) Analyses of substrate specificity of His-CreTPT3. Liposomes in which His-CreTPT3 was incorporated were preloaded with various substrates (30 mM) and the initial Pi uptake rates (at 0.25 mM external concentration) were determined. Relative velocities were compared to the Pi/Pi homo-exchange, which was set to 100%. The data represents the arithmetic mean (±SD) of three independent experiments (each consisting of three technical replicates). (**f**) Comparison of kinetic constants of CreTPT3. The Michaelis-Menten constant (K_M) for Pi uptake was determined using various external Pi concentrations (0.05-5 mM). The competitive inhibition constant (K_i) of Pi uptake (0.25 mM) was measured with increasing inhibitor concentrations (0.05-5 mM). Liposomes were preloaded with 30 mM Pi as the counter-exchange substrate. The data represent the arithmetic mean +/- SE of three independent experiments. ND, no competitive inhibitory constant could be measured under the given experimental conditions. DHAP, dihydroxyacetone phosphate; 3-PGA, 3-phosphoglycerate; PEP, phosphoenolpyruvate; Glc6P, glucose 6-P; Glc1P, glucose 1-P; Fru6P, fructose 6-P; Gal1P, galactose 1-P.

Additionally, a phylogenic tree was constructed to help determine the specificities of the putative

CreTPTs. This tree includes all of the CreTPTs and the plastidial phosphate transporters (PT) from

Arabidopsis thaliana (AtPPT1, AtTPT1, AtGPT1 and AtXPT1). The CreTPT1, 2, 3 and 25 protein

sequences cluster with the different AtTPTs (**Fig. 1a**, **Supplementary Fig 3**), again suggesting

that these individual Chlamydomonas transporters have different substrate specificities. Since

CreTPT3 is most closely related to AtTPT1 (**Fig. 1a**, **Supplementary Fig. 3**), the major triose-P

permease in Arabidopsis, we considered CreTPT3 as the Chlamydomonas ortholog of AtTPT1.

TPT3 transport properties

To help establish the physiological (in vivo) function of CreTPT3, its subcellular localization was

first experimentally confirmed in Chlamydomonas. The CreTPT3 coding sequence with its

putative chloroplast targeting sequence was fused with the VENUS tag (CreTPT3-VENUS) and

expressed under the control of its own promoter in the Chlamydomonas mutant null for CreTPT3.

As shown in **Fig. 1b**, CreTPT3-VENUS was localized to the chloroplast envelope.

To evaluate the substrate specificity of CreTPT3, it was heterologously expressed in

Saccharomyces cerevisiae (yeast) cells and the resulting recombinant protein was biochemically

analyzed using a liposome uptake system. The Cre*TPT3* gene was codon-optimized, expressed in

yeast (**Fig. 1c**), total cell membranes were isolated and then CreTPT3 was reconstituted into

liposomes (Linka et al., 2008; Loddenkötter et al., 1993). The chloroplast phosphate transporters

in vascular plants can catalyze a strict Pi/Pi homo-exchange in vitro. The reconstituted CreTPT3

in liposomes was able to catalyze the signature Pi homo-exchange, whereas in the absence of a

counter-exchange substrate, little Pi uptake was detected. This result demonstrates that the

recombinant CreTPT3 protein can function in Pi/Pi homo-exchange in liposomes (**Fig. 1d**). In

contrast, very low Pi uptake rates were observed for liposomes reconstituted with membranes from

yeast cells lacking CreTPT3 (**Supplementary Fig. 4**), indicating that the detected Pi import

activity was the result of CreTPT3 expression in yeast.

To assess the substrate specificity, the initial rates of Pi uptake into liposomes preloaded with

saturating concentrations (30 mM) of various putative counter-exchange substrates were

determined. As **Fig. 1e** shows, CreTPT3 exhibited the highest activity when DHAP was used as

substrate. The relative initial velocity for Pi uptake in exchange for DHAP was slightly higher than

that of 3-PGA (3-PGA exhibited a 75% exchange rate relative to that of Pi/DHAP) while the Pi

uptake into liposomes preloaded with PEP was much less efficient (about 16% relative to the Pi/DHAP exchange rate) (**Fig. 1e**). Pi import was negligible when CreTPT3 liposomes were preloaded with Glc-6-P, Glc-1-P, Fru-6-P, and Gal-1-P (**Fig. 1e**).

Characterizations of the K_M and K_i of CreTPT3 were performed to determine the affinity of the transporter for the various substrates. CreTPT3 has an apparent Michaelis-Menten constant (K_M) value of 1.1 +/- 0.2 mM for Pi (**Fig. 1f**), which is comparable to the value obtained for the vascular plant TPT ortholog (Flügge et al., 1989). DHAP and 3-PGA competitively inhibited Pi uptake with an inhibition constant (Ki) of 0.57 +/- 0.01 mM and 0.79 +/- 0.12 mM, respectively (**Fig. 1f**), suggesting that CreTPT3 has a higher affinity for DHAP and 3-PGA than for Pi. In the case of PEP, no inhibition of the Pi/Pi homo-exchange was observed, even at the non-physiologically high concentration of 5 mM. Overall, the results of this in vitro assay indicate that CreTPT3 has a typical plant-TPT substrate spectrum (Flügge et al., 1989). It can transport Pi, triose-P (DHAP) and 3-PGA in a counter-exchange mode, however PEP might not be a physiologically relevant substrate for CreTPT3, as was shown for apoplast TPT homologues in other organisms (Lim et al., 2010; Moog et al., 2020).

Isolation of *tpt3* null mutants and their impact on cell growth

To explore the role of CreTPT3 in trafficking carbon and reductant across the chloroplast envelope in vivo, CRISPR knockouts of Cre*TPT3* were generated. We used the CRISPR-Cas9 editing system to disrupt the Cre*TPT3* gene in exons 1 and 7, while at the same time integrating the hygromycin marker gene (*AphVII*) into the edited site (**Supplementary Fig. 5a**). Multiple independent knockouts of Cre*TPT3* were obtained, including *t3ko1*, with the marker gene inserted into exon 1, and *t3ko2* and *t3ko3*, with the marker gene inserted into exon 7 (**Supplementary Fig. 5**, **6**) using specific sgRNA2 guides. It was previously proposed that chloroplast TPT could catalyze two potential reactions based on TPT structure analysis in red algae (Lee et al., 2017b) and on in vitro assays using isolated spinach chloroplasts (Stocking and Larson, 1969): *i*) triose-P (DHAP, G3P)/Pi or *ii*) triose-P /3-PGA exchange across the chloroplast inner envelope membrane. The former reaction can route both carbon skeletons and reductants into the cytoplasm while importing Pi back into the chloroplast for ATP regeneration. The latter reaction imports 3-PGA into the chloroplast in exchange for triose-P, which allows for the export of reductant from the chloroplast to the cytoplasm (DHAP or G3P), with the imported 3-PGA serving as substrate for

448 the CBBC, allowing for regeneration of ribulose 1,5-bisphosphate when it becomes depleted.

Figure 2: Growth characterization of *tpt3* mutants and complemented strains. (a) Growth of
WT, *t3ko1*, *t3ko2* and the complemented strains (*C_T3KO1* and *C_T3KO2*, respectively), at
various dilutions (see below) on agar plates incubated under low light (LL, ~30 μmol photons m^{-2}
s^{-1}) and moderate light (ML, 250~300 μmol photons m^{-2} s^{-1}) for 4 days. Growth of respective
strains in liquid medium under LL (**b**) and ML (**c**). The data represents the arithmetic mean (±SD)
of three independent experiments. Transition of the strains from LL to ML (**d**) or to high light (HL,
450 μmol photons m^{-2} s^{-1}) (**e**) for the times indicated. (**f**) Cell growth (number) and chlorophyll
content of the indicated cultures at various times following exposure to ML (from LL), for up to
48 h. In (**a**) the cells were spotted on agar plates containing TAP or TP medium under continuous

LL or ML conditions. The dilution series used was 1.5, 0.75, 0.375 μg/mL chlorophyll (left to

right). For growth in liquid medium, the cells were cultured in TP medium to an initial OD_{750nm}

0.02, in air and under continuous LL, before initiating the various growth analyses in LL, ML or

HL.

To elucidate the physiological roles of CreTPT3, we first examined growth of the parental WT

strain and the knockout mutants under either LL or ML. In contrast to WT cells, growth of the

CreTPT3 knockouts (*t3ko1*, *t3ko2*, *t3ko3*) was significantly reduced in LL on either

photoautotrophic (TP) or heterotrophic (TAP) agar medium (**Fig. 2a**, **Supplementary Fig. 7**),

while the growth of the mutant in ML was completely blocked in both TP and TAP (**Fig. 2a**).

Growth curves were also determined for WT and mutant cells in liquid medium (TP) in both LL

and ML (**Fig. 2b**, **c**); the results were in accord with those observed for the agar medium growth

assays. Additionally, *tpt3* mutant cells in both LL and ML exhibited an increased cell diameter and

formed clusters of cells that appear to be less able to separate following division (**Supplementary**

Fig. 8a, **b**). To further confirm that the growth phenotypes are a consequence of the *tpt3* knockout,

we introduced a wild-type copy of CreTPT3 fused to VENUS into the mutant strains

(**Supplementary Fig. 5b**). As demonstrated in **Fig. 2b** and **2c**, ectopic expression of the wild-type

CreTPT3 in the *t3ko2* mutant (*C_T3KO2*) rescued the reduced growth phenotype of the mutant in

both LL and ML.

We also analyzed the chlorophyll content of cells grown photoautotrophically upon transfer from

LL to ML (**Supplementary Fig. 1**) and found that *t3ko2* had reduced chlorophyll content relative

to WT cells after 24 h in ML (**Fig. 2d-f** and **Supplementary Fig. 9**). Cell numbers and total

chlorophyll were quantified following the LL to ML transition. The chlorophyll levels per cell

declined in all strains at 24 and 48 h following the transfer to ML; however, it was lower by more

than half in *t3ko2* (0.34 μg/10^6 cells) relative to either WT cells or the *C_T3KO2* rescued strain

(both ~0.72 μg/10^6 cells) after 24 h of ML (**Supplementary Fig. 9**), with some additional increase

for WT and *C_T3KO2* after 48 h of ML illumination (which might reflect some shading caused

by an increase in the cell density). Finally, when mutant cells were transferred HL for 24 h they

became highly bleached (**Fig. 2e**).

Taken together, these results indicate that the activity of the CreTPT3 shuttle is essential for growth

over a range of light intensities (LL/ML/HL).

The *tpt3* mutant exhibits hyper-accumulation of 'storage' carbon

To explore the impact of the loss of CreTPT3 activity on carbon partitioning, we quantified carbon storage (starch and lipids) following a transition of WT and the *t3ko2* mutant from LL to ML. Lugol staining showed extensive starch accumulation in the *t3ko2* cells after a 48-h exposure in the ML, whereas, the WT cells were barely stained **(Fig. 3a)**. Furthermore, as shown in **Fig. 3b**, there is an ~55-fold (WT and *t3ko2*: 13.29 and 0.24 µg starch/µg chlorophyll, respectively) increase in starch accumulation (normalized to chlorophyll) in the *t3ko2* mutant after 24 h of illumination in ML. The mutant also exhibited a ~24-fold (WT and *t3ko2*: 1038.93 and 41.98 Nile Red fluorescence/chlorophyll, respectively) increase in lipid accumulation, as monitored by Nile Red fluorescence, relative to WT and *C_T3KO2* over the same time period (**Fig. 3c, d**) (on a chlorophyll basis). Since the chlorophyll in the mutant was approximately 50% relative to that of WT cells after 24 h in ML, the accumulated starch and lipid on a per cell basis would be ~25-fold for starch and ~12-fold for lipid in the mutant relative to the WT strains. Furthermore, the mutant cells are much bigger (**Supplementary Fig. 8a, b**), which may reflect the large increase in storage carbon that they accumulated in the mutant cells. Accumulated starch granules in the stroma of ML-exposed *t3ko2* cells are readily visualized under bright field light microscopy (**Supplementary Video 1a, b**). These results suggest that the inability to transport triose-P between the chloroplast and the cytosol through CreTPT3 resulted in a repartitioning of photosynthetic assimilates (carbon, reductant, and ATP) toward the synthesis of both starch and TAG.

Figure 3: Changes in storage carbon and metabolite levels in WT and *t3ko2* after transitioning the cells from growth in LL to ML. (a) Lugol staining of starch in WT and *t3ko2* after 48 h of ML. **(b)** Starch content in indicated strains following 0, 6, and 24 h of ML exposure. **(c)** Nile Red staining of TAG in WT and *t3ko2* following 48 h of ML exposure. **(d)** TAG content in strains following 0, 6, and 24 h of ML exposure. **(e)** Time course of metabolite accumulation at 0 h (LL), 45 min and 6 h following ML exposure. **(f)** Select metabolic pathways in Chlamydomonas; adapted from (Johnson and Alric, 2013). **(g)** Bar graph representation of fold-change for metabolites shown in **(e)**, calculated by dividing the pool size in *t3ko2* by that of WT

at the respective light levels. A fold change of 1 (no change) is shown by a dashed line. Each data
point shows the mean and standard error; the data represents three biological replicates for each
metabolite. An asterisk indicates statistically significant difference compared to WT (* P<0.05, **
P<0.01, *** P<0.001). Abbreviations: G6P, glucose-6-P; FBP, fructose bisphosphate; DHAP,
dihydroxyacetone phosphate; 3-PGA, 3-phosphoglycerate; PEP, phosphoenolpyruvate; PYR,
pyruvate; CIT, citrate; SUCC, succinate; FUM, fumarate; MAL, malate; E4P, Erythrose 4-P; R5P,
Ribose 5-P; RU5P/X5P, ribulose 5-P/xylulose-5-P.

Cre*TPT3* deletion leads to accumulation of CBBC/glycolytic/gluconeogenic intermediates

To understand the metabolic consequences of the loss of CreTPT3 on growth under LL and ML,
comparative metabolic analysis of WT and *t3ko2* mutant was performed on cells grown at LL and
after shifting them to ML for 45 min or 6 h. Pool sizes of various central carbon metabolites,
particularly those of the CBBC/glycolysis/gluconeogenesis pathways and intermediates of the
TCA/glyoxylate cycle (**Fig. 3f**), were quantified. Data was normalized to both chlorophyll content
(**Fig. 3e**) and cell number (**Supplementary Fig. 10b**). The fold-change in the quantity of each
metabolite in the mutant relative to WT under LL or at 45 min and 6 h after the switch to ML is
given in **Fig. 3g** and **Supplementary Fig. 10c**.

Under LL and upon transfer to ML, the *t3ko2* mutant maintained a significantly larger pool (2 to
6-fold) of glycolytic/gluconeogenic intermediates [DHAP, 3-PGA and FBP (all three shared with
CBBC), G6P, PEP], and of some metabolites of the CBBC pathway (E4P, RU5P/X5P, R5P)
compared to WT, suggesting that the loss of CreTPT3 results in a back-up of these metabolites
within the cell; our hypothesis is that they are accumulating inside the chloroplast due to the lack
of CreTPT3 activity. This is supported by the finding of a marked accumulation of starch and TAG
in the mutant strain (**Fig. 3a-d**). While the pool sizes of fumarate and malate (metabolites of the
TCA/glyoxylate cycle) were similar in both strains under LL, following the transition of the cells
to ML, WT cells showed a significant increase in the pool sizes of those metabolites while the
mutant maintained a lower level, indicating that the loss of CreTPT3 either directly (by supplying
precursors) or indirectly (metabolic rewiring) impacts their levels.

Thus, the one primary function of CreTPT3 would be the export of photosynthetically-synthesized
reduced carbon, which would drive metabolic processes in the cytoplasm and other cellular
compartments, while importing Pi into the chloroplast to sustain ATP synthesis.

The *tpt3* mutant has reduced PSII activity when grown in moderate light

To understand the impact of the loss of CreTPT3 activity on photosynthetic electron transport (PET), we quantified photosynthetic activities following a transition from LL to ML. After growing all strains photoautotrophically in LL for 16-24 h, the cultures were diluted with fresh medium to a concentration of 1-1.5 µg/mL chlorophyll and then transferred to ML for 24 h (**Supplementary Fig. 1**). Photosynthetic O_2 evolution rates (OERs) were measured for WT, *t3ko2* and *C_T3KO2* at a range of light intensities. LL acclimated WT, *t3ko2*, and *C_T3KO2* showed comparable OER at intensities below 200 µmol photons m^{-2} s^{-1} while at saturating light intensities of ≥600 µmol photons m^{-2} s^{-1}, the OER of *t3ko2* was diminished relative to the WT and complemented strain (**Fig. 4a**). Upon acclimation of the cells to ML for 24 h, the *t3ko2* consistently displayed low OER relative to WT and *C_T3KO2* under all actinic light intensities used (**Fig. 4b**).

The PSII quantum yield, ΦPSII, is maximum after a dark pre-incubation (Fv/Fm). The Fv/Fm in the mutant grown in LL was similar to that of WT and *C_T3KO2* immediately following the dark incubation (**Fig. 4c**). The LL-acclimated WT and *t3ko2* cells exhibited comparable ΦPSII values when the actinic light intensity was below 100 µmol photons m^{-2} s^{-1}, but decreased by about 20-25% when the actinic light was increased to 400 µmol photons m^{-2} s^{-1}; that decrease was maintained even when the actinic light was raised to over 1,000 µmol photons m^{-2} s^{-1} (**Supplementary Fig. 11**). In contrast, dark acclimated mutant cells that had been exposed to ML for 6 h exhibited an Fv/Fm that was about 2-fold lower than that of WT and *C_T3KO2* and continued to decline over a period of 24 h in ML, with a 3-fold decrease in Fv/Fm for *t3ko2* compared to that of WT (**Fig. 4c** and **Supplementary Fig. 11**).

The oxidation of the PQ pool is the slowest step in the PET chain between PSII and PSI (Escoubas et al., 1995; Stiehl and Witt, 1969). The redox potential of the PQ pool changes upon increasing the irradiance to which the cells are exposed. Thus, the redox potential of the PQ pool serves as a biological measure for evaluating the balance between the photochemistry of PSII and linear photosynthetic electron transport. Changes of the redox state of the PQ pool in LL and ML acclimated cells were analyzed at increasing light intensities (**Fig. 4d, Supplementary Fig. 12**). In LL-grown cells, the PQ pool was significantly more reduced in *t3ko2* even at low level illumination (measured as 1-qL) (Saroussi et al., 2019) compared to that of WT and the rescued strain. When the actinic light intensity was elevated to 400 µmol photons m^{-2} s^{-1}, the reduction

levels of the PQ pool increased for all strains, although it was still much more reduced in *t3ko2*

compared to that of WT and the rescued strains (**Fig. 4d, Supplementary Fig. 12**).

For cells that had been acclimated to ML (6 h and 24 h), the PQ pool was already very highly

reduced in *t3ko2* following exposure to LL, while it remained highly oxidized for both the WT and

the rescued strain (**Fig. 4d, Supplementary Fig. 12**). The level of PQ reduction in ML-acclimated

t3ko2 increased only slightly with an increase in the actinic light intensity (**Fig. 4d**,

Supplementary Fig. 12). There was also an increase in the PQ pool reduction of WT and

C_T3KO2 as the light intensity was raised, however, even at the highest light intensity the WT and

rescued strains maintained a much more oxidized PQ pool than that of the mutant; the highest

levels of 1-qL for WT and *C_T3KO2* were still ~2-fold lower than that of *t3ko2* (**Fig. 4d,**

Supplementary **Fig. 12**). Overall, inactivation of CreTPT3 resulted in attenuated PSII activities

(Fv/Fm, OER) and hyper-reduction of the PQ pool, suggesting a block in electron flow

downstream of PQ pool.

Figure 4: Photosynthetic activities followed LL to ML transition. (**a**) O_2 evolution of LL-

acclimated cells in the indicated strains. (**b**) O_2 evolution of ML-acclimated (24 h) cells. (**c**) Fv/Fm

of WT, *t3ko2* and the rescued strain following 6 and 24 h in ML. (**d**) 1-qL values (redox state of

PQ pool) of LL- and ML-acclimated WT, *t3ko2* and the rescued strain. The acclimated cells were

exposed to 50 and 400 µmol photons $m^{-2} s^{-1}$, as indicated, for 2 mins. (**e-g**) PSI activities following

a LL to ML transition. P700 oxidation and reduction kinetics in LL-grown cells (**e**) and after 6 (**f**) and 24 h (**g**) in ML. Absorbance differences were monitored at 705 nm during continuous illumination with 150 μmol photons·m^{-2} s^{-1} for 5 sec (white box above e), followed by a saturating light pulse at 1,500 μmol photons·m^{-2} s^{-1} (arrow) and a 5 sec dark incubation (black box). The data represent the arithmetic mean (±SD) of three independent experiments. Kinetics were normalized by setting maximum oxidation (after light pulse) for WT to 100%. For panels c and d, the asterisks represent significant differences determined by ANOVA tests. **P < 0.005, ****P < 0.0001.

Impact of *tpt3* mutant on PSI activity

To determine if PSI activity was also impacted in the mutant cells, we analyzed PSI/P700 oxidation/reduction kinetics following exposure of LL-grown cells to ML. Levels of photo-oxidizable P700 were similar in all LL-acclimated strains (**Fig. 4e**). However, after 6 h in ML, the level of photo-oxidizable P700 declined in the mutant to ~40% of that in WT and *C_T3KO2* cells (**Fig. 4f**), followed by slightly more inhibition of PSI activity after 24 h of exposure to ML (**Fig. 4g**). These deleterious impacts on photosynthetic electron transport likely reflect a defect in routing carbon/energy out of the chloroplast, which in turn will cause hyper-accumulation of reductant in the chloroplast, which would potentially result a backup of PET, elevated ROS accumulation and damage to the photosynthetic apparatus.

CreTPT3 is crucial for balancing metabolite flux and chloroplast redox

Photosynthetic organisms mainly accumulate ROS in the chloroplast and mitochondria as byproducts of aerobic metabolism (Halliwell, 2006). To assay ROS production in the mutant, we used the fluorescent probe CM-H2DCFDA which, upon exposure to increasing ROS levels, is converted to the green fluorescent molecule dichlorofluorescein (DCF). DCF fluorescence was visualized by confocal microscopy. As shown in **Fig. 5a** and **b**, ROS levels in the *t3ko2* mutant were markedly increased (~3-fold) after 48 h in ML, while no difference in the level was detected in WT cells.

For an alternative, dynamic method for evaluating redox changes in the chloroplast, we used the redox sensitive green fluorescent protein roGFP2 (**Supplementary Fig. S13**) that was targeted to the chloroplast stroma. Two cysteine residues were engineered into the roGFP2 to form a reduction-oxidation-dependent disulfide bond that detects intracellular changes in ROS accumulation (Vevea et al., 2013). roGFP2 was shown to monitor ratiometric redox changes of glutathione (Dorion et al., 2021; Vevea et al., 2013). The ascorbate–glutathione (GSH) system is

central to ROS detoxification (Foyer and Noctor, 2011), with the redox potential of glutathione
reflecting ROS levels. LL-acclimated WT cells and the *t3ko2* mutant exhibited similar levels of
chloroplast roGFP oxidation (**Fig. 5c**). Upon transfer of WT cells to ML, the oxidation level of
roGFP in the chloroplast increased by ~2-fold after 1.5 h, compared to that of WT cells maintained
in LL, followed by a decrease after a 6 h ML exposure (**Fig. 5c**). The changes of redox in WT
upon a LL to ML transition is in accord with previous results (Lin et al., 2016; Vidal-Meireles et
al., 2017). Furthermore, the synthesis of both GSH and ascorbate increases when the redox balance
is shifted to a largely oxidized state after transferring cells to a higher light intensity, and recovers
several hours after the transfer (Lin et al., 2016; Vidal-Meireles et al., 2017) as the cells acclimate
to the higher light conditions. A similar redox pattern determined by chloroplast roGFP2 was
observed for the *t3ko2* mutant after it was shifted to ML, however, the mutant showed a much
greater increase in chloroplast oxidative conditions, with a 5.0-fold increase after 1.5 h, and a 2.0-
fold increase after 6 h, or 1.8- and 1.6-fold higher than the levels measured in WT cells (**Fig. 5c**).

Figure 5: Measurement of intracellular reactive ROS. (**a**) ROS levels were evaluated by CM-
H2DCFDA fluorescence of WT and the *t3ko2* mutant in LL and following exposure to ML for 48
h. (**b**) Quantitation of data in **a**. (**c-d**) Monitoring redox levels in the chloroplast (**c**) and
mitochondria (**d**) after ML exposure of WT and the *t3ko2* mutant for 1.5 and 6.0 h. Values are
means ± SD (n ≥ 3). The asterisks represent a significant difference determined by an ANOVA
test. **$P < 0.005$, ***$P < 0.001$. Strains were grown photoautotrophically under LL for 16-24 h.
After dilution with fresh medium, cultures were transferred to ML for the times indicated.

As triose-P is exported to the cytosol by CreTPT3, it could potentially donate redox equivalents to

mitochondrial electron transport and alter mitochondrial ROS production. Therefore, we measured

mitochondrial redox levels at different light intensities in both WT and the *t3ko2* mutant using the

roGFP sensor targeted to the mitochondrion. Upon a LL to ML transition, neither WT nor *t3ko2*

displayed a change in fluorescence from the mitochondrial targeted roGFP (**Fig. 5d**). These results

suggest that the mitochondrial redox level is maintained after shifting either LL acclimated WT or

t3ko2 cells to ML. This is in agreement with the results obtained for diatoms; the roGFP oxidation

level in diatom mitochondria exhibited either no or minor changes in response to various stress

conditions, including nitrogen limitation, HL, and iron limitation (van Creveld et al., 2015).

Overall, these results suggest that the increase in oxidative conditions caused by a LL to ML

transition is primarily localized to the chloroplast in Chlamydomonas and that the inability to

export triose-P through TPT3 markedly increases the chloroplast oxidative stress level.

Export of triose-P is essential for maintaining chloroplast redox levels in the dark

Another issue that we addressed was whether CreTPT3 is important for dark metabolism in

Chlamydomonas. We analyzed redox changes in WT and the *t3ko2* mutant upon transferring cells

from LL to the dark. Intriguingly, ROS accumulation in WT was 5 times higher than that of *t3ko2*

after 48 h of darkness (**Fig. 6a, b**). RoGFP2 was used to determine the cellular compartment

associated with ROS accumulated. The oxidation level of chloroplast roGFP2 in WT was slightly

reduced during the first 3 h of darkness, followed by a >1.5-fold increase after 6 h of darkness

compared to LL-acclimated cells (**Fig. 6c**). This is in keeping with previous observations in

Arabidopsis in which exposing leaves to extended darkness resulted in an initial decrease in the

degree of oxidation of roGFP2 in the chloroplast followed by an increase to the original LL level

when the darkness was prolonged (Rosenwasser et al., 2011). Unlike the slight decrease in roGFP2

oxidation in the WT chloroplast, the oxidation state of the chloroplast in *t3ko2* cells declined by

2.2-fold relative to the original LL conditions and by 1.77-fold relative to WT after 3 h in the dark

(**Fig. 6c**). The chloroplast oxidation level in *t3ko2* did not recover to the level of WT cells after 6

h of darkness but attained its initial LL level after 24 h in the dark (**Fig. 6c**) suggesting that the

diminished export of triose-P transport and the consequent accumulation of storage carbon resulted

in a decline in the chloroplast redox potential in dark.

The initial decline in the oxidation level of the probe upon a light to dark transition indicates that

chloroplast ROS generated in the light immediately decreases after the cells are shifted to the dark.

It was proposed that during an extended period in the dark, Arabidopsis becomes sugar starved

which leads to the diminished availability of reductant for scavenging ROS (Rosenwasser et al.,

2011; Usadel et al., 2008). Although our interpretation is speculative, given the highly reducing

state of the chloroplast and the elevated accumulation of starch in the chloroplast in *t3ko2*, the

mutant will likely maintain elevated levels of sugars in the chloroplast during the dark period as it

mobilizes the high level of carbon stored during the light period. Indeed, as shown in **Fig. 6e**, *t3ko2*

accumulates 7 times (*t3ko2*, WT: 10.21, 1.403 µg starch/µg chlorophyll) more starch than that of

WT cells after 12 h in the light. Similar to WT and the rescued strain, *t3ko2* can mobilize 70% of

the starch at the end of the light period (**Fig. 6e**). Thus, as the cells transition from the light to the

dark, the chloroplast of *t3ko2* has approximately 7-fold more carbohydrates than that of WT cells

and the rescued strain potentially giving it a greater capacity to generate metabolites that efficiently

scavenge excess ROS.

Figure 6: Metabolic and redox changes in *tpt3* over a diel cycle or extended darkness. (a)

ROS levels monitored by CM-H2DCFDA staining in WT and *t3ko2* after the cells are transitioned

from LL to darkness in TP medium. (**b**) Quantification of the data presented in **a**. Monitoring redox

levels in the chloroplast (**c**) and mitochondrion (**d**) in the dark with roGFP2 (targeted to chloroplast

and mitochondria) in WT and the *t3ko2* mutant after various times in the dark. (**e**) Starch content

over the diel cycle in indicated strains. The intensity of the light phase was 60 µmol photons m^{-2}

s^{-1}. (**f**) Respiration during the dark phase of the diel cycle with/without *n*-propyl gallate (PG:
inhibitor of alternative respiration AOX+PTOX). (**g**) Growth of WT, *t3ko2* and the complemented
strains on agar plates in continuous light (CL) (~60 μmol photons m^{-2} s^{-1}) or on a diel cycle for 6
d. The cells were plated at 3, 1.5, 0.75, 0.375, and 0.1875 μg/mL chlorophyll. Values (from b,c,d,f)
are means ± SD (n ≥ 3). The asterisks represent significant difference determined by ANOVA. *P
< 0.05, **P < 0.005, ****P < 0.0001.

Mitochondrial dark respiration relieves redox stress in *tpt3* mutant

The mitochondrial oxidation level of *t3ko2* during the dark period was generally slightly higher
than that in WT (**Fig. 6d**). The respiration rate of *t3ko2* in the dark increased 2-fold compared to
that of WT cells irrespective of whether the mitochondrial alternative oxidase was blocked (**Fig.
6f**), suggesting that there is more substrate available for fueling mitochondrial electron flow and
oxidative phosphorylation, which in turn could fuel growth. These results imply that during the
dark period of the diel cycle, mitochondrial respiration could dissipate a proportion of the reductant
that accumulated in the *t3ko2* mutant during the light period. Indeed, the growth of *t3ko2* during
diel cycle growth (relative to continuous light) is partially rescued in TP and almost fully recovered
in TAP (**Fig. 6g**). In addition to growth, the morphology of *t3ko2* mutant observed over an
extended dark period is recovered. As shown in **Supplementary Fig. 14**, some of *t3ko2* cells are
aggregated under LL, however, an extended dark period results in the complete separation of *t3ko2*
cells. These results indicate that CreTPT3 might play only a minor role during darkness, or that
the loss of CreTPT3 in the dark can be compensated for by the other sugar-P transporters, such as
the TPT2, the putative glucose-P transporter (CreTPT25), or the maltose exporter (Findinier et al.,
2017). Indeed, the Cre*TPT25* and *TPT1* genes, which potentially encode the homologues of the
Arabidopsis glucose 6-P and phosphoenolpyruvate transporters, respectively, are upregulated in
darkness and downregulated in light (**Supplementary Table S3**, **Supplementary Fig. 16a**).

Discussion

CreTPT3 is critical for the distribution of photosynthate

Photoassimilates that serve as substrate for the chloroplast triose-P/Pi translocator (TPT) have been
studied in various vascular plants (e,g, tobacco, *Arabidopsis*, potato, etc). However, surprisingly
few studies have characterized chloroplast envelope localized members of the triose-P family in
trafficking fixed carbon in single-celled photosynthetic organisms (Haferkamp et al., 2006; Linka
et al., 2008; Moog et al., 2020); their physiological role(s) in algae has not been elucidated. In this

work, we hypothesized that the highly expressed Chlamydomonas chloroplast envelope triose-P/Pi

translocator3 (CreTPT3) functions as a 'genuine' TPT. To explore the function of CreTPT3, we

expressed it in yeast cells and performed in vitro studies to assay its activity in liposomes (**Fig. 1c-**

f) and also generated mutants disrupted for the Cre*TPT3* gene to examine the impact of a loss of

function mutant on the physiology of Chlamydomonas.

An Arabidopsis mutant for the chloroplast TPT was isolated 20 years ago (Schneider et al., 2002).

The growth and photosynthesis of this mutant, designated *ape2*, was not affected under LL

conditions (Walters et al., 2004), although its starch content was slightly elevated relative to that

of WT Arabidopsis (Walters et al., 2004). The absence of AtTPT in LL-grown plants appeared to

be fully compensated for by directing more photosynthate toward starch synthesis and turning over

the starch rapidly in the light to generate sugars like hexose phosphate (hexose-P) and other

metabolites that can be exported from the chloroplast. Furthermore, while *ape2* grew in HL, its

growth was attenuated relative to WT Arabidopsis (Walters et al., 2004), although there was no

detectable change in the level of starch (Schneider et al., 2002). In potato, a 2-3-fold increase in

starch was observed in transgenic plants with diminished levels of the chloroplast targeted TPT

(Riesmeier et al., 1993). The reduction in TPT activity led to a strong growth retardation during

the young stages of development. Similar to Arabidopsis, a tobacco strain with decreased TPT

activity was able to compensate for the defect by accelerating the synthesis and turnover of starch

and the transport of hexose from the chloroplast (Häusler et al., 1998). These observations suggest

that various vascular plants respond to a defect in the export of triose-P from chloroplasts by

moving the fixed carbon into a transient starch pool that is converted to sucrose/glucose/glucose-

P through accelerated hydrolysis. Glucose-P and potentially other metabolites generated from

starch degradation can then be exported from the chloroplast and trafficked to sink organs,

providing them with carbon skeletons and energy for anabolic processes (Winter and Huber, 2000).

In contrast, a deficiency of TPT in rice, which uses sucrose stored in leaves as its major transitory

form of fixed carbon rather than starch, led to reduced photosynthetic rates and about a 2-fold

decrease in the level of starch relative to WT plants (Lee et al., 2014).

The Chlamydomonas genome encodes 4 putative triose-P transporters predicted by TargetP-2.0 to

be routed to the chloroplast. Cre*TPT3* is the most highly expressed among the chloroplast

associated TPTs (**Supplementary Table S3** and **Supplementary Fig. 15a**). A detailed analysis

comparing Chlamydomonas TPT proteins with those of other organisms revealed that CreTPT3 is

most similar to the *Arabidopsis thaliana* TPT1 (**Fig. 1a**). In this study we localized CreTPT3 to

the chloroplast envelope. We also performed in vitro transport studies with liposomes reconstituted

with CreTPT3 expressed in yeast and demonstrated the capacity of this translocator to use Pi,

DHAP and 3-PGA as substrates (**Fig. 1d-f**), similar to the plant TPT (Flügge et al., 1989). Hence,

it appears that CreTPT3 can exchange chloroplast/stromal triose-Ps for cytosolic Pi, and may also

serve as redox shuttle by catalyzing DHAP/3-PGA exchange.

To probe the in vivo function of CreTPT3, we generated multiple null mutants, designated *t3ko*

and characterized one of the mutants (*t3ko2)* in depth. Under LL, growth of the *t3ko2* knockout

mutant was markedly inhibited (**Fig. 2a**, **b**, **Supplementary Fig. 7a, d**), a phenotype exacerbated

upon transferring the cells from LL to either ML or HL (**Fig. 2a**, **c**, **d**, **e**, **Supplementary Fig. 7b**,

c, **d**). Upon exposure to ML, *t3ko2* cells stopped growing, and when placed in HL, they exhibited

severe bleaching (**Fig. 2e**); this phenotype can be rescued by ectopic expression of Cre*TPT3* in the

mutant background. These results indicate that the flux of fixed carbon trafficked through the

CreTPT3 is crucial for Chlamydomonas growth under all light intensities tested, and that it

becomes essential for survival at the higher intensities.

Chlamydomonas, unlike plant cells, contains little intracellular sucrose, with starch as its major

form of storage carbon (Klein, 1987). In addition to a pronounced growth phenotype, the *t3ko2*

mutant exhibited a dramatic change in its carbon partitioning. After 24 h of LL, *t3ko2* contained

~16-fold more starch than that of WT cells (**Supplementary Fig. 10a**). After 24 h in ML, the

starch and lipid contents of *t3ko2* were 55 and 24-fold higher than that of WT cells, respectively

(**Fig. 3a-d**); these results were based on normalization to chlorophyll, which was reduced by ~50%

in the ML-grown mutants based on the number of cells.

In Arabidopsis, elimination of the chloroplast targeted TPT protein did not significantly alter the

profiles of metabolite of either the CBBC, or the glycolytic/gluconeogenic pathways, suggesting

the occurrence of compensatory mechanisms that redirect carbon from the chloroplast of the

mutant plant (Hilgers et al., 2018). In contrast, a Chlamydomonas mutant null for Cre*TPT3*

exhibited a dramatic increase of most metabolites in the glycolytic/gluconeogenic pathways,

suggesting that triose-P exchange catalyzed by CreTPT3 was not readily compensated for by

alternative fixed carbon shuttles. CreTPT1, CreTPT2 and CreTPT25, the other shuttles that

potentially transport triose-P or hexose-P across the Chlamydomonas chloroplast envelope, are

unable to fully compensate for the loss of CreTPT3 (**Supplementary Fig. 15b-d**). These results

demonstrate the critical role of CreTPT3 in shuttling photoassimilates in Chlamydomonas. The

retention/accumulation of photoassimilates in the chloroplast appears to redirect them toward the

synthesis of starch and lipid, which would cause a marked deficit of fixed carbon in other cellular

compartments, such as the cytoplasm and mitochondrion, and severely compromise physiological

processes including cell growth and division; a loss of CreTPT3 even impacts carbon partitioning

and growth under various light conditions (LL/ML) as well as in the dark (**Fig. 2**, **3**, **6,**

Supplementary Fig. 10, **14**).

CreTPT3 is essential for controlling 'overflow' electrons/reductant in the chloroplasts

Disruption of triose-P export from the chloroplast through disruption of Cre*TPT3* also resulted in

inhibition of PET (**Fig. 4**). When WT, *t3ko2*, and the complemented strain *C_T3KO2* were grown

in LL, they exhibited a similar Fv/Fm and photosynthetic OER when exposed to low intensity

actinic light (**Fig. 4a**, **c**). However, even after exposure to LL, the PQ pool became more highly

reduced in *t3ko2* cells than in either WT or *C_T3KO2* cells (**Fig. 4d**). Hence, PSII is functioning

properly in LL-acclimated *t3ko2* and is capable of normal electron flow and reduction of

plastoquinone (PQ) to plastoquinol (plastohydroquinone, PQH2). However, reoxidation of PQH2

to PQ in *t3ko2* is slowed in LL-acclimated mutant cultures, which results in a decline in the rate

of O_2 evolution as the actinic light intensity is increased. PSI activity also appears to be unaffected

in LL-acclimated mutant cultures (**Fig. 4e**) which, in combination with measurements of PSII

activity, suggest that hyper-reduction of the PQ pool in LL-acclimated cells is caused by a block

in the utilization of electrons downstream of PSI. A similar hyper-reduction of the PQ pool occurs

when Chlamydomonas cells are unable to synthesize starch (Krishnan et al., 2015).

Disruption of the Cre*TPT3* gene impacted PET more acutely when the cells were transferred to

ML (**Fig. 4**). ML-acclimated *t3ko2* cells exhibited diminished Fv/Fm and photosynthetic OER

relative to WT and C_*T3KO2*. The PQ pool of ML-acclimated *t3ko2* cells was completely reduced

(**Fig. 4d**). Additionally, PSI activity was depressed in *t3ko2* cells transferred to ML (**Fig. 4e**).

These results together with the altered metabolite pool sizes and carbon partitioning in the mutant

(**Fig. 3**) suggest that an inability to efficiently export the terminal products of carbon fixation

results in hyper-reduction of the photosynthetic electron transport chain that in turn impacts the

activity of the photosynthetic reaction centers (PSII and PSI). The diminished Fv/Fm value in cells

maintained in ML clearly indicates a loss of active PSII reaction center proteins; further studies

will be needed to elucidate the effect of the altered metabolism on light harvesting function and

composition. Lower net OER may also reflect an increase in O_2 photoreduction in the light (H_2O-

to-H_2O cycle) in *t3ko2* mutant relative to WT cells. Future studies will focus on elucidating the

dynamics of O_2 evolution in this mutant using MIMS, which will help determine the importance

of H_2O-to-H_2O cycle in *t3ko* mutants relative to the WT cells.

Studies of photosynthetic activities of *t3ko2* in ML support our hypothesis that the mutant has a

reduced capacity to utilize reductant for anabolic processes required for growth. The ML-incubated

t3ko2 mutant accumulated high levels of carbon metabolites that serve as precursors for the

biosynthesis of starch, lipids, and amino acids, suggesting a systems-level consequence of

accumulating high levels of reduced carbon in the chloroplast (**Fig. 3e**) and slowing the

regeneration of $NADP^+$ and ADP from NADPH and ATP, respectively. The marked alteration in

metabolite pools observed in the mutant could feedback on PET and potentially elicits

photosynthetic control, constricting electron flow at the cyt b_6f complex as a consequence of the

generation of a large ΔpH across the thylakoid membranes (Joliot and Johnson, 2011; Johnson and

Berry, 2021). Such control would result in hyper-reduction of the PQ pool and reduced electron

flow to PSI, which would help protect PSI from photoinhibition. Furthermore, *t3ko2* cells are

strongly photoinhibited following their transfer from LL to ML, conditions that cause little change

in PET of WT and *C_T3KO2*. These results strongly suggest that CreTPT3 is essential for

maintaining the flow of fixed carbon and reductant from the chloroplast to the cytosol where it can

be allocated for various anabolic processes that support normal cell growth.

The malate-OAA shunt was shown to play a role in exporting reductant from chloroplasts and

linking chloroplast redox stress, which can be elicited by high light and various conditions that

limit growth and/or block photosynthetic activities, to mitochondrial metabolism (Zhao et al.,

2020). The malate-OAA shunt represents another activity that, under high redox stress, might have

compensated for the loss of the CreTPT3 shuttle. Intriguingly, malate levels in *t3ko2* were 4-fold

lower than in WT cells following a transition from LL to ML (**Fig. 3e**, **g**). Furthermore, a previous

study of metabolic flux analysis during heterotrophic growth of Chlamydomonas showed that the

TPT shuttle is almost 10-fold more active than the malate-OAA shunt (Boyle et al., 2017). These

results together with the physiological phenotypes of the *tpt3ko* mutant suggest that the malate-
OAA shunt cannot compensate for a loss of CreTPT3 activity. Furthermore, Cre*TPT3* is the most
responsive/upregulated of the four Chlamydomonas, chloroplast-localized *TPT* genes following
exposure of cells to nitrogen, sulfur or iron deprivation (**Supplementary Fig. 16 b-d**), suggesting
that it plays an important role in controlling chloroplast redox conditions. Deprivation of
Chlamydomonas for nitrogen or sulfur caused dramatic attenuation of PET (Saroussi et al., 2017)
as the cells elicited an array of 'energy management' strategies to balance energy generation and
utilization (Saroussi et al., 2017). Elevated NPQ, chlororespiration and lipid/starch accumulation
are well recognized mechanisms to dissipate excess light energy and minimize ROS production in
Chlamydomonas in response to nutrient deprivation (Saroussi et al., 2017). In this study, TPT3-
dependent chloroplast triose-P/Pi transport appears to be critical for distributing fixed carbon and
reductant to other cellular compartments. When this function is compromised, the chloroplast
accumulates various metabolites and altered substrate availability for chloroplast-localized
metabolic processes; elevated NADPH/NADP$^+$ and ATP/ADP$^+$ ratios, hyper-reduction of
chloroplast redox components, such as the PQ pool, and elevated production of ROS can lead to
dysfunction of the photosynthetic apparatus (and other metabolic activities).

CreTPT3 is central to metabolic and redox networks

AGPase, a key enzyme in starch synthesis, is exquisitely sensitive to allosteric regulation; it is
activated by 3-PGA and inhibited by Pi (Preiss, 1988). When the *t3ko2* mutant was exposed to LL,
there was already a dramatic increase of 3-PGA (6-fold) relative to WT cells (**Fig. 3e**, **g**). The loss
of CreTPT3 might also lead to less import of Pi into chloroplasts. Therefore, the ratio of 3-PGA/Pi
would be markedly elevated in the *tpt3* mutant compared to that of the WT cells and the
complemented strain. Thus, even in LL, AGPase in the mutant would become active, which in turn
would lead to starch accumulation (**Supplementary Fig. 10a**).

In chloroplasts, reductant generated by photosynthetic electron transport can modulate the activity
of various redox-sensitive target proteins via thiol-based redox regulation. Key metabolic
pathways both within and external to the chloroplast must be coordinated with the rate at which
the photosynthetic light reactions produce reductant and ATP (Cejudo et al., 2021; Yoshida and
Hisabori, 2018; Yu et al., 2020). Chloroplasts harbor two thiol-based redox systems that
independently regulate plastid metabolism by communicating redox conditions to specific target

enzymes, thereby increasing or decreasing their activities. One system, FDX/TRX reductase, obtains reducing equivalents from FDX (which is reduced by PET) (Meyer et al., 2008) while the other involves NADPH-TRX reductase (NTRC), which uses NADPH as its substrate (Pérez-Ruiz and Cejudo, 2009). Both systems can activate ADP-glucose pyrophosphorylase (AGPase) and starch biosynthesis (Ballicora et al., 2000; Lepistö et al., 2013). There was a dramatic impact of the loss of CreTPT3 on the redox state of PET components. Illumination of LL-acclimated *t3ko2* mutant cells with 50 μmol photons m^{-2} s^{-1} actinic light resulted in elevated reduction of the PQ pool compared to that of WT (**Fig. 4d**) while exposure of the mutant to ML for 6 h led to near complete reduction of the PQ pool (**Fig. 4d**, **Supplementary Fig. 12**), which could activate the FDX/TRX system and alter the activities of its various target enzymes, including increasing AGPase activity, which is needed for the synthesis of starch. Furthermore, an increase in the levels of phosphorylated intermediates of the CBBC (**Fig. 3e**, **g**) typically leads to a reduced rate of cycling NADPH to NADP$^+$, resulting in a higher NADPH:NADP$^+$ ratio and availability of reductant for activation of the NTRC (NADPH-TRX reductase) system, which would also contribute to AGPase activation and starch hyperaccumulation.

In sum, we show that the loss of a chloroplast envelope localized triose-P/Pi exchange protein results in elevated 3-PGA levels and slow turnover of carbon precursors and reducing equivalents, leading to PET hyper-reduction. This could possibly result in an increase in the NADPH:NADP$^+$ ratio. These metabolic alterations would lead to both allosteric (high 3-PGA) and redox regulated (Trx or NTRC systems) activation of AGPase, even in LL. Hence, starch overaccumulation in *t3ko2* would initially reflect metabolic control, with subsequent regulation through the FDX/TRX or NTRC redox system (see model on **Fig. 7**).

Additionally, the highly compromised transport of triose-P out of the chloroplast in *t3ko2* would lead to hyper-reduction of photosynthetic electron carriers, which is reflected by a highly reduced PQ pool, the redox state of the chloroplast and mitochondrion obtained by roGFP analyses, and elevated ROS production/accumulation (**Fig. 5a-c**). ROS was initially considered harmful, causing oxidative stress, damaging lipids, proteins, and DNA, eventually resulting in cell death (Schieber and Chandel, 2014). Increasing evidence, however, suggests that low levels of ROS can function in diverse signaling pathways (Schieber and Chandel, 2014) that regulate various cellular processes (Foyer and Noctor, 2005; Wagner et al., 2004) such as growth, the cell cycle, senescence

(Zimmermann and Zentgraf, 2005), and acclimation to stressful biotic and abiotic conditions
(Miller et al., 2008). ROS can also regulate expression of some chloroplast genes encoding proteins
associated with photosynthesis and modulation of PET (Yu et al., 2020). For example, ROS
inhibits *de novo* synthesis of the PSII reaction center protein D1, which would lead to
photoinhibition during high light acclimation (Kojima et al., 2007; Nishiyama et al., 2001).
Moreover, when PSI becomes highly reduced, superoxide and singlet oxygen both elicit PSI
photoinhibition (Takagi et al., 2016).

Overall, Chlamydomonas mutants lacking TPT3 develop a hyper-reduced chloroplast landscape
with elevated ROS production (**Fig. 5c**). After 6 h or more in ML, PSII and PSI activities in the
mutant become compromised, most likely a consequence of elevated ROS levels (see model on
Fig. 7). The chloroplast generated ROS serves as a messenger for reporting redox information (e.g.
light levels, stress conditions) to the regulatory machinery of the cell, which results in modulation
of expression of nuclear genes that would ameliorate the impact and damage associated with
elevated ROS levels (Exposito-Rodriguez et al., 2017; Foyer, 2018; Wakao and Niyogi, 2021).
Many studies have shown that stress-induced lipid/triacylglycerol (TAG) accumulation also
correlates with an increase in ROS production (Shi et al., 2017); ROS may activate diacylglycerol
acyltransferases (DGAT) and phospholipid diacylglycerol acyltransferase activities (PDAT)
thereby stimulating lipid accumulation (Chouhan et al., 2021).

As depicted in **Fig. 7**, we propose that various tiers of redox regulation are responsible for the
differential physiological responses of the Chlamydomonas *tpt3* knockout mutants relative to WT
cells and the rescued strain. When *t3ko2* cells are initially transferred from LL to ML, the triose-P
pool and metabolites derived from that pool accumulate in the mutant because of its reduced
capacity to move triose-P out of the chloroplast, which also causes reduced CBBC activity, hyper-
reduction of PET and the accumulation of ROS (which would inhibit photosynthetic activity). The
hyper-reduction and accumulation of fixed carbon in the chloroplast of the *tpt3* mutants would
activate AGPase through allosteric regulation and the FDX/TRX and the NTRC redox systems,
leading to starch hyper-accumulation (**Fig. 7**). Furthermore, increased ROS/RS accumulation and
the elevated redox state of PET would elicit retrograde signals that mediate changes in expression
of both chloroplast and nuclear genes, marshaling the synthesis of specific activities that function
to ameliorate the impact of the hyper-reduced state of the chloroplast resulting from the

compromised ability of the cells to export reductant and fixed carbon from the organelle (**Fig. 7**).

Figure 7: Impact of eliminating Chlamydomonas TPT3 on the chloroplast metabolic
landscape. i. reduced triose-P export leads to accumulation of sugar-P (e.g. triose-P, hexose-P,
marked in pink), and an elevated ratio of 3-PGA/Pi (which can activate AGPase activity) would
stimulate production of storage carbon; CBBC activity diminishes along with the fixation of CO_2,
leading to hyperaccumulation of reductant (NADPH) and energy (ATP) within the chloroplast; **ii.**
the elevated NADPH:NADP$^+$ and ATP:ADP ratios, elicit strong feedback on PET causing hyper-
reduction of electron carriers (eg: over reduced PQ pool and highly reduced PSI) that slows
electron flow across the cytochrome b_6f complex; **iii.** the highly reduced PET system and the
elevated NADPH can actively reduce FDX/TRX and the NTRC, respectively; **iv.** reduced TRX
and/or NTRC can activate the AGPase (above the activity elicited by an increase in the 3-PGA/Pi
ratio), leading to additional starch accumulation; **v.** hyper-reduction of PET can also lead to ROS
accumulation; **vi.** ROS can act as a signal that controls expression of chloroplast and nuclear genes;
vii. ROS also cause damage to both PSII and I and lead to lipid accumulation as a consequence of
activation of diacylglycerol acyltransferases (DGAT) and phospholipid diacylglycerol
acyltransferase (PDAT). Many of these metabolic/acclimatory processes both overlap and are
interconnected. Under conditions of extended exposure to moderate light, the damage in the
Chlamydomonas *tpt3* mutant can accumulate and lead to cell death; cell death occurs rapidly in
high light.
962

Acknowledgments:

WH thanks M. Onishi for advice on generation of CRISPR mutants and A. Burlacot for help with
PQ pool measurements.

Fundings:

This project has received funding from DOE award DE-SC0019417. RGK and JF were supported
by the Department of Plant Biology of the Carnegie Institution for Science. PR was supported by
Human Frontier Science Program (HFSP) RGP0046/2018 (to ARG).

Author contributions

WH, AK, ARG and MP conceptualized the study. WH generated the CRISPR mutants as well as
the complemented strain, localized the TPT3 protein, analyzed the redox status in the cell,
performed respiration measurements, measured the reduction level of PQ pool, analyzed changes
of starch over the diel cycle. AK performed the experiments of photosynthetic O_2 evolution,
extraction of samples for LC-MS/MS, data analysis of LC-MS/MS. AP and NL performed the
reconstitution into liposomes and transport activity assays. MM and NB performed LC-MS/MS
and analyses of metabolite data. BR performed data mining for transcriptome and the Nile red
staining. JF and NF analyzed starch changes upon transition from LL to ML. WH, PR and ARG
analyzed PSI and PSII activities. JF and BR constructed roGFP2 associated vectors. JX and WH
performed the phylogenetic tree. WH drafted the manuscript. ARG, MP, AK, NL, NB, AP, MM
and all the other authors helped with revising and finalizing the manuscript.

References

Badger, M.R., von Caemmerer, S., Ruuska, S., and Nakano, H. (2000). Electron flow to oxygen
in higher plants and algae: rates and control of direct photoreduction (Mehler reaction) and rubisco
oxygenase. Philos. Trans. R. Soc. Lond. B Biol. Sci. *355*, 1433–1446.
https://doi.org/10.1098/rstb.2000.0704.

Ballicora, M.A., Frueauf, J.B., Fu, Y., Schürmann, P., and Preiss, J. (2000). Activation of the
potato tuber ADP-glucose pyrophosphorylase by thioredoxin. J. Biol. Chem. *275*, 1315–1320.
https://doi.org/10.1074/jbc.275.2.1315.

Boyle, N.R., Sengupta, N., and Morgan, J.A. (2017). Metabolic flux analysis of heterotrophic
growth in *Chlamydomonas reinhardtii*. PLoS One *12*, e0177292.
https://doi.org/10.1371/journal.pone.0177292.

Cejudo, F.J., González, M.-C., and Pérez-Ruiz, J.M. (2021). Redox regulation of chloroplast
metabolism. Plant Physiol. *186*, 9–21. https://doi.org/10.1093/plphys/kiaa062.

Chaux, F., Burlacot, A., Mekhalfi, M., Auroy, P., Blangy, S., Richaud, P., and Peltier, G. (2017).
Flavodiiron Proteins Promote Fast and Transient O_2 Photoreduction in Chlamydomonas. Plant
Physiol. *174*, 1825–1836. https://doi.org/10.1104/pp.17.00421.

Chouhan, N., Devadasu, E., Yadav, R.M., and Subramanyam, R. (2021). Autophagy induced
accumulation of lipids in *pgrl1* and *pgr5* of *Chlamydomonas reinhardtii* under high light. Front.
Plant Sci. *12*, 752634. https://doi.org/10.3389/fpls.2021.752634.

van Creveld, S.G., Rosenwasser, S., Schatz, D., Koren, I., and Vardi, A. (2015). Early perturbation
in mitochondria redox homeostasis in response to environmental stress predicts cell fate in diatoms.
ISME J. *9*, 385–395. https://doi.org/10.1038/ismej.2014.136.

Crozet, P., Navarro, F.J., Willmund, F., Mehrshahi, P., Bakowski, K., Lauersen, K.J., Pérez-Pérez,
M.-E., Auroy, P., Gorchs Rovira, A., Sauret-Gueto, S., et al. (2018). Birth of a Photosynthetic
Chassis: A MoClo Toolkit Enabling Synthetic Biology in the Microalga *Chlamydomonas*
reinhardtii. ACS Synth. Biol. *7*, 2074–2086. https://doi.org/10.1021/acssynbio.8b00251.

Dietz, K.-J., Turkan, I., and Krieger-Liszkay, A. (2016). Redox- and Reactive Oxygen Species-
Dependent Signaling into and out of the Photosynthesizing Chloroplast. Plant Physiol. *171*, 1541–
1550. https://doi.org/10.1104/pp.16.00375.

Dorion, S., Ouellet, J.C., and Rivoal, J. (2021). Glutathione Metabolism in Plants under Stress:
Beyond Reactive Oxygen Species Detoxification. Metabolites *11*, 641.
https://doi.org/10.3390/metabo11090641.

Edgar, R.C. (2004). MUSCLE: multiple sequence alignment with high accuracy and high
throughput. Nucleic Acids Res. *32*, 1792–1797. https://doi.org/10.1093/nar/gkh340.

Eicks, M., Maurino, V., Knappe, S., Flügge, U.-I., and Fischer, K. (2002). The plastidic pentose
phosphate translocator represents a link between the cytosolic and the plastidic pentose phosphate

pathways in plants. Plant Physiol. *128*, 512–522. https://doi.org/10.1104/pp.010576.

Escoubas, J.M., Lomas, M., LaRoche, J., and Falkowski, P.G. (1995). Light intensity regulation
of cab gene transcription is signaled by the redox state of the plastoquinone pool. Proc. Natl. Acad.
Sci. U. S. A. *92*, 10237–10241. https://doi.org/10.1073/pnas.92.22.10237.

Exposito-Rodriguez, M., Laissue, P.P., Yvon-Durocher, G., Smirnoff, N., and Mullineaux, P.M.
(2017). Photosynthesis-dependent H_2O_2 transfer from chloroplasts to nuclei provides a high-light
signalling mechanism. Nat. Commun. *8*. https://doi.org/10.1038/s41467-017-00074-w.

Facchinelli, F., and Weber, A.P.M. (2011). The metabolite transporters of the plastid envelope: an
update. Front. Plant Sci. *2*, 50. https://doi.org/10.3389/fpls.2011.00050.

Findinier, J., Tunçay, H., Schulz-Raffelt, M., Deschamps, P., Spriet, C., Lacroix, J.-M., Duchêne,
T., Szydlowski, N., Li-Beisson, Y., Peltier, G., et al. (2017). The Chlamydomonas *mex1* mutant
shows impaired starch mobilization without maltose accumulation. J. Exp. Bot. *68*, 5177–5189.
https://doi.org/10.1093/jxb/erx343.

Findinier, J., Delevoye, C., and Cohen, M.M. (2019). The dynamin-like protein Fzl promotes
thylakoid fusion and resistance to light stress in *Chlamydomonas reinhardtii*. PLoS Genet. *15*,
e1008047. https://doi.org/10.1371/journal.pgen.1008047.

Fischer, K., Kammerer, B., Gutensohn, M., Arbinger, B., Weber, A., Hausler, R.E., and Flugge,
U.-I. (1997). A new class of plastidic phosphate translocators: A putative link between primary
and secondary metabolism by the phosphoenolpyruvate/phosphate antiporter. Plant Cell *9*, 453.
https://doi.org/10.2307/3870494.

Fliege, R., Flügge, U.-I., Werdan, K., and Heldt, H.W. (1978). Specific transport of inorganic
phosphate, 3-phosphoglycerate and triosephosphates across the inner membrane of the envelope
in   spinach   chloroplasts.   Biochim.   Biophys.   Acta   Bioenerg.   *502*,   232–247.
https://doi.org/10.1016/0005-2728(78)90045-2.

Flügge, U.I., Fischer, K., Gross, A., Sebald, W., Lottspeich, F., and Eckerskorn, C. (1989). The
triose   phosphate-3-phosphoglycerate-phosphate   translocator   from   spinach   chloroplasts:
nucleotide sequence of a full-length cDNA clone and import of the in vitro synthesized precursor
protein into chloroplasts. EMBO J. *8*, 39–46. .

Foyer, C.H. (2018). Reactive oxygen species, oxidative signaling and the regulation of
photosynthesis.   Environ.   Exp.   Bot.   *154*,   134–142.
https://doi.org/10.1016/j.envexpbot.2018.05.003.

Foyer, C.H., and Noctor, G. (2005). Redox Homeostasis and Antioxidant Signaling: A Metabolic
Interface between Stress Perception and Physiological Responses. Plant Cell *17*, 1866–1875.
https://doi.org/10.1105/tpc.105.033589.

Foyer, C.H., and Noctor, G. (2011). Ascorbate and glutathione: the heart of the redox hub. Plant
Physiol. *155*, 2–18. https://doi.org/10.1104/pp.110.167569.

Ge, F., Huang, W., Chen, Z., Zhang, C., Xiong, Q., Bowler, C., Yang, J., Xu, J., and Hu, H. (2014). Methylcrotonyl-CoA Carboxylase Regulates Triacylglycerol Accumulation in the Model Diatom *Phaeodactylum tricornutum*. Plant Cell *26*, 1681–1697. https://doi.org/10.1105/tpc.114.124982.

Gibson, D.G., Young, L., Chuang, R.-Y., Venter, J.C., Hutchison, C.A., 3rd, and Smith, H.O. (2009). Enzymatic assembly of DNA molecules up to several hundred kilobases. Nat. Methods *6*, 343–345. https://doi.org/10.1038/nmeth.1318.

Gietz, R.D., and Schiestl, R.H. (2007). High-efficiency yeast transformation using the LiAc/SS carrier DNA/PEG method. Nat. Protoc. *2*, 31–34. https://doi.org/10.1038/nprot.2007.13.

Haferkamp, I., and Schmitz-Esser, S. (2012). The plant mitochondrial carrier family: functional and evolutionary aspects. Front. Plant Sci. *3*, 2. https://doi.org/10.3389/fpls.2012.00002.

Haferkamp, I., Deschamps, P., Ast, M., Jeblick, W., Maier, U., Ball, S., and Neuhaus, H.E. (2006). Molecular and biochemical analysis of periplastidial starch metabolism in the cryptophyte Guillardia theta. Eukaryot. Cell *5*, 964–971. https://doi.org/10.1128/EC.00381-05.

Halliwell, B. (2006). Reactive species and antioxidants. Redox biology is a fundamental theme of aerobic life. Plant Physiol. *141*, 312–322. https://doi.org/10.1104/pp.106.077073.

Harris, E.H. (2009). Chlamydomonas sourcebook: Introduction to Chlamydomonas and its laboratory use (Academic Press).

Häusler, R.E., Schlieben, N.H., Schulz, B., and Flügge, U.I. (1998). Compensation of decreased triose phosphate/phosphate translocator activity by accelerated starch turnover and glucose transport in transgenic tobacco. Planta *204*, 366–376. https://doi.org/10.1007/s004250050268.

Heinrichs, L., Schmitz, J., Flügge, U.-I., and Häusler, R.E. (2012). The Mysterious Rescue of adg1-1/tpt-2 - an Arabidopsis thaliana Double Mutant Impaired in Acclimation to High Light - by Exogenously Supplied Sugars. Front. Plant Sci. *3*, 265. https://doi.org/10.3389/fpls.2012.00265.

Hilgers, E.J.A., Schöttler, M.A., Mettler-Altmann, T., Krueger, S., Dörmann, P., Eicks, M., Flügge, U.-I., and Häusler, R.E. (2018). The Combined Loss of Triose Phosphate and Xylulose 5-Phosphate/Phosphate Translocators Leads to Severe Growth Retardation and Impaired Photosynthesis in *Arabidopsis thaliana* tpt/xpt Double Mutants. Front. Plant Sci. *9*, 1331. https://doi.org/10.3389/fpls.2018.01331.

Houille-Vernes, L., Rappaport, F., Wollman, F.-A., Alric, J., and Johnson, X. (2011). Plastid terminal oxidase 2 (PTOX2) is the major oxidase involved in chlororespiration in Chlamydomonas. Proc. Natl. Acad. Sci. U. S. A. *108*, 20820–20825. https://doi.org/10.1073/pnas.1110518109.

Huang, W., Haferkamp, I., Lepetit, B., Molchanova, M., Hou, S., Jeblick, W., Río Bártulos, C., and Kroth, P.G. (2018). Reduced vacuolar β-1,3-glucan synthesis affects carbohydrate metabolism as well as plastid homeostasis and structure in *Phaeodactylum tricornutum*. Proc. Natl. Acad. Sci. U. S. A. *115*, 4791–4796. https://doi.org/10.1073/pnas.1719274115.

Jack, D.L., Yang, N.M., and Saier, M.H., Jr (2001). The drug/metabolite transporter superfamily.

Eur. J. Biochem. *268*, 3620–3639. https://doi.org/10.1046/j.1432-1327.2001.02265.x.

Johnson, X., and Alric, J. (2013). Central carbon metabolism and electron transport in
Chlamydomonas reinhardtii: metabolic constraints for carbon partitioning between oil and starch.
Eukaryot. Cell *12*, 776–793. https://doi.org/10.1128/EC.00318-12.

Jokel, M., Johnson, X., Peltier, G., Aro, E.-M., and Allahverdiyeva, Y. (2018). Hunting the main
player enabling *Chlamydomonas reinhardtii* growth under fluctuating light. Plant J. *94*, 822–835.
https://doi.org/10.1111/tpj.13897.

Kammerer, B., Fischer, K., Hilpert, B., Schubert, S., Gutensohn, M., Weber, A., and Flügge, U.I.
(1998). Molecular characterization of a carbon transporter in plastids from heterotrophic tissues:
the glucose 6-phosphate/phosphate antiporter. Plant Cell *10*, 105–117.
https://doi.org/10.1105/tpc.10.1.105.

Kaye, Y., Huang, W., Clowez, S., Saroussi, S., Idoine, A., Sanz-Luque, E., and Grossman, A.R.
(2019). The mitochondrial alternative oxidase from *Chlamydomonas reinhardtii* enables survival
in high light. J. Biol. Chem. *294*, 1380–1395. https://doi.org/10.1074/jbc.RA118.004667.

Klein, U. (1987). Intracellular Carbon Partitioning in *Chlamydomonas reinhardtii*. Plant Physiol.
85, 892–897. https://doi.org/10.1104/pp.85.4.892.

Klein, U., and Betz, A. (1978). Fermentative metabolism of hydrogen-evolving *Chlamydomonas*
moewusii. Plant Physiol. *61*, 953–956. https://doi.org/10.1104/pp.61.6.953.

Knappe, S., Flügge, U.-I., and Fischer, K. (2003). Analysis of the plastidic phosphate translocator
gene family in Arabidopsis and identification of new phosphate translocator-homologous
transporters, classified by their putative substrate-binding site. Plant Physiol. *131*, 1178–1190.
https://doi.org/10.1104/pp.016519.

Kojima, K., Oshita, M., Nanjo, Y., Kasai, K., Tozawa, Y., Hayashi, H., and Nishiyama, Y. (2007).
Oxidation of elongation factor G inhibits the synthesis of the D1 protein of photosystem II. Mol.
Microbiol. *65*, 936–947. https://doi.org/10.1111/j.1365-2958.2007.05836.x.

Kong, F., Burlacot, A., Liang, Y., Légeret, B., Alseekh, S., Brotman, Y., Fernie, A.R., Krieger-
Liszkay, A., Beisson, F., Peltier, G., et al. (2018). Interorganelle Communication: Peroxisomal
MALATE DEHYDROGENASE2 Connects Lipid Catabolism to Photosynthesis through Redox
Coupling in Chlamydomonas. Plant Cell *30*, 1824–1847. https://doi.org/10.1105/tpc.18.00361.

Krishnan, A., Kumaraswamy, G.K., Vinyard, D.J., Gu, H., Ananyev, G., Posewitz, M.C., and
Dismukes, G.C. (2015). Metabolic and photosynthetic consequences of blocking starch
biosynthesis in the green alga *Chlamydomonas reinhardtii* sta6 mutant. Plant J. *81*, 947–960.
https://doi.org/10.1111/tpj.12783.

Kumar, S., Stecher, G., and Tamura, K. (2016). MEGA7: Molecular Evolutionary Genetics
Analysis version 7.0 for bigger datasets. Mol. Biol. Evol. *33*, 1870–1874.
https://doi.org/10.1093/molbev/msw054.

Kuźniak, E., and Kopczewski, T. (2020). The chloroplast reactive oxygen species-redox system in
plant immunity and disease. Front. Plant Sci. *11*, 572686.
https://doi.org/10.3389/fpls.2020.572686.

Lee, S.-K., Eom, J.-S., Voll, L.M., Prasch, C.M., Park, Y.-I., Hahn, T.-R., Ha, S.-H., An, G., and
Jeon, J.-S. (2014). Analysis of a triose phosphate/phosphate translocator-deficient mutant reveals
a limited capacity for starch synthesis in rice leaves. Mol. Plant *7*, 1705–1708.
https://doi.org/10.1093/mp/ssu082.

Lee, Y., Nishizawa, T., Takemoto, M., Kumazaki, K., Yamashita, K., Hirata, K., Minoda, A.,
Nagatoishi, S., Tsumoto, K., Ishitani, R., et al. (2017a). Crystal structure of the triose-
phosphate/phosphate translocator in complex with inorganic phosphate.
https://doi.org/10.2210/pdb5y78/pdb.

Lee, Y., Nishizawa, T., Takemoto, M., Kumazaki, K., Yamashita, K., Hirata, K., Minoda, A.,
Nagatoishi, S., Tsumoto, K., Ishitani, R., et al. (2017b). Structure of the triose-
phosphate/phosphate translocator reveals the basis of substrate specificity. Nat Plants *3*, 825–832.
https://doi.org/10.1038/s41477-017-0022-8.

Lepistö, A., Pakula, E., Toivola, J., Krieger-Liszkay, A., Vignols, F., and Rintamäki, E. (2013).
Deletion of chloroplast NADPH-dependent thioredoxin reductase results in inability to regulate
starch synthesis and causes stunted growth under short-day photoperiods. J. Exp. Bot. *64*, 3843–
3854. https://doi.org/10.1093/jxb/ert216.

Li, M., and Kim, C. (2022). Chloroplast ROS and stress signaling. Plant Commun *3*, 100264.
https://doi.org/10.1016/j.xplc.2021.100264.

Lim, L., Linka, M., Mullin, K.A., Weber, A.P.M., and McFadden, G.I. (2010). The carbon and
energy sources of the non-photosynthetic plastid in the malaria parasite. FEBS Lett. *584*, 549–554.
https://doi.org/10.1016/j.febslet.2009.11.097.

Lin, S.-T., Chiou, C.-W., Chu, Y.-L., Hsiao, Y., Tseng, Y.-F., Chen, Y.-C., Chen, H.-J., Chang,
H.-Y., and Lee, T.-M. (2016). Enhanced Ascorbate Regeneration Via Dehydroascorbate Reductase
Confers Tolerance to Photo-Oxidative Stress in *Chlamydomonas reinhardtii*. Plant Cell Physiol.
57, 2104–2121. https://doi.org/10.1093/pcp/pcw129.

Linka, M., Jamai, A., and Weber, A.P.M. (2008). Functional characterization of the plastidic
phosphate translocator gene family from the thermo-acidophilic red alga *Galdieria sulphuraria*
reveals specific adaptations of primary carbon partitioning in green plants and red algae. Plant
Physiol. *148*, 1487–1496. https://doi.org/10.1104/pp.108.129478.

Loddenkötter, B., Kammerer, B., Fischer, K., and Flügge, U.I. (1993). Expression of the functional
mature chloroplast triose phosphate translocator in yeast internal membranes and purification of
the histidine-tagged protein by a single metal-affinity chromatography step. Proc. Natl. Acad. Sci.
U. S. A. *90*, 2155–2159. https://doi.org/10.1073/pnas.90.6.2155.

Meyer, Y., Siala, W., Bashandy, T., Riondet, C., Vignols, F., and Reichheld, J.P. (2008).
Glutaredoxins and thioredoxins in plants. Biochim. Biophys. Acta *1783*, 589–600.

https://doi.org/10.1016/j.bbamcr.2007.10.017.

Miller, G., Shulaev, V., and Mittler, R. (2008). Reactive oxygen signaling and abiotic stress.
Physiol. Plant. *133*, 481–489. https://doi.org/10.1111/j.1399-3054.2008.01090.x.

Moog, D., Nozawa, A., Tozawa, Y., and Kamikawa, R. (2020). Substrate specificity of plastid
phosphate transporters in a non-photosynthetic diatom and its implication in evolution of red alga-
derived complex plastids. Sci. Rep. *10*, 1167. https://doi.org/10.1038/s41598-020-58082-8.

Nishiyama, Y., Yamamoto, H., Allakhverdiev, S.I., Inaba, M., Yokota, A., and Murata, N. (2001).
Oxidative stress inhibits the repair of photodamage to the photosynthetic machinery. EMBO J. *20*,
5587–5594. https://doi.org/10.1093/emboj/20.20.5587.

Pérez-Ruiz, J.M., and Cejudo, F.J. (2009). A proposed reaction mechanism for rice NADPH
thioredoxin reductase C, an enzyme with protein disulfide reductase activity. FEBS Lett. *583*,
1399–1402. https://doi.org/10.1016/j.febslet.2009.03.067.

Preiss, J. (1988). Biosynthesis of starch and its regulation. In The Biochemistry of Plants,
(Elsevier), pp. 181–254.

Riesmeier, J.W., Flügge, U.I., Schulz, B., Heineke, D., Heldt, H.W., Willmitzer, L., and Frommer,
W.B. (1993). Antisense repression of the chloroplast triose phosphate translocator affects carbon
partitioning in transgenic potato plants. Proc. Natl. Acad. Sci. U. S. A. *90*, 6160–6164.
https://doi.org/10.1073/pnas.90.13.6160.

Rosenwasser, S., Rot, I., Sollner, E., Meyer, A.J., Smith, Y., Leviatan, N., Fluhr, R., and Friedman,
H. (2011). Organelles contribute differentially to reactive oxygen species-related events during
extended darkness. Plant Physiol. *156*, 185–201. https://doi.org/10.1104/pp.110.169797.

Sake, C.L., Newman, D.M., and Boyle, N.R. (2020). Evaluation of quenching methods for
metabolite recovery in photoautotrophic *Synechococcus sp.* PCC 7002. Biotechnol. Prog. *36*,
e3015. https://doi.org/10.1002/btpr.3015.

Saroussi, S., Sanz-Luque, E., Kim, R.G., and Grossman, A.R. (2017). Nutrient scavenging and
energy management: acclimation responses in nitrogen and sulfur deprived Chlamydomonas. Curr.
Opin. Plant Biol. *39*, 114–122. https://doi.org/10.1016/j.pbi.2017.06.002.

Schieber, M., and Chandel, N.S. (2014). ROS function in redox signaling and oxidative stress.
Curr. Biol. *24*, R453-62. https://doi.org/10.1016/j.cub.2014.03.034.

Schneider, A., Häusler, R.E., Kolukisaoglu, Ü., Kunze, R., Van der Graaff, E., Schwacke, R.,
Catoni, E., Desimone, M., and Flügge, U.I. (2002). An Arabidopsis thaliana knock-out mutant of
the chloroplast triose phosphate/phosphate translocator is severely compromised only when starch
synthesis, but not starch mobilisation is abolished. Plant Journal https://doi.org/10.1046/j.1365-
313X.2002.01460.x.

Shi, K., Gao, Z., Shi, T.-Q., Song, P., Ren, L.-J., Huang, H., and Ji, X.-J. (2017). Reactive oxygen
species-mediated cellular stress response and lipid accumulation in oleaginous microorganisms:

The state of the art and future perspectives. Front. Microbiol. *8*, 793. https://doi.org/10.3389/fmicb.2017.00793.

Stiehl, H.H., and Witt, H.T. (1969). Quantitative Treatment of the Function of Plastoquinone in Photosynthesis. Zeitschrift Für Naturforschung B *24*, 1588–1598. https://doi.org/10.1515/znb-1969-1219.

Stocking, C.R., and Larson, S. (1969). A chloroplast cytoplasmic shuttle and the reduction of extraplastid NAD. Biochem. Biophys. Res. Commun. *37*, 278–282. https://doi.org/10.1016/0006-291x(69)90731-1.

Takagi, D., Takumi, S., Hashiguchi, M., Sejima, T., and Miyake, C. (2016). Superoxide and singlet oxygen produced within the thylakoid membranes both cause photosystem I photoinhibition. Plant Physiol. *171*, 1626–1634. https://doi.org/10.1104/pp.16.00246.

Usadel, B., Bläsing, O.E., Gibon, Y., Retzlaff, K., Höhne, M., Günther, M., and Stitt, M. (2008). Global transcript levels respond to small changes of the carbon status during progressive exhaustion of carbohydrates in Arabidopsis rosettes. Plant Physiol. *146*, 1834–1861. https://doi.org/10.1104/pp.107.115592.

Vevea, J.D., Alessi Wolken, D.M., Swayne, T.C., White, A.B., and Pon, L.A. (2013). Ratiometric biosensors that measure mitochondrial redox state and ATP in living yeast cells. J. Vis. Exp. 50633. https://doi.org/10.3791/50633.

Vidal-Meireles, A., Neupert, J., Zsigmond, L., Rosado-Souza, L., Kovács, L., Nagy, V., Galambos, A., Fernie, A.R., Bock, R., and Tóth, S.Z. (2017). Regulation of ascorbate biosynthesis in green algae has evolved to enable rapid stress-induced response via the VTC2 gene encoding GDP-l-galactose phosphorylase. New Phytol. *214*, 668–681. https://doi.org/10.1111/nph.14425.

Wagner, D., Przybyla, D., den Camp, R. op, Kim, C., Landgraf, F., Lee, K.P., Würsch, M., Laloi, C., Nater, M., Hideg, E., et al. (2004). The Genetic Basis of Singlet Oxygen–Induced Stress Responses of *Arabidopsis thaliana*. Science https://doi.org/10.1126/science.1103178.

Wakao, S., and Niyogi, K.K. (2021). Chlamydomonas as a model for reactive oxygen species signaling and thiol redox regulation in the green lineage. Plant Physiol. *187*, 687–698. https://doi.org/10.1093/plphys/kiab355.

Walters, R.G., Ibrahim, D.G., Horton, P., and Kruger, N.J. (2004). A mutant of Arabidopsis lacking the triose-phosphate/phosphate translocator reveals metabolic regulation of starch breakdown in the light. Plant Physiol. *135*, 891–906. https://doi.org/10.1104/pp.104.040469.

Weber, A.P.M., Linka, M., and Bhattacharya, D. (2006). Single, ancient origin of a plastid metabolite translocator family in Plantae from an endomembrane-derived ancestor. Eukaryot. Cell *5*, 609–612. https://doi.org/10.1128/EC.5.3.609-612.2006.

Winter, H., and Huber, S.C. (2000). Regulation of sucrose metabolism in higher plants: localization and regulation of activity of key enzymes. Crit. Rev. Biochem. Mol. Biol. *35*, 253–289. https://doi.org/10.1080/10409230008984165.

Yoshida, K., and Hisabori, T. (2018). Determining the Rate-Limiting Step for Light-Responsive
Redox Regulation in Chloroplasts. Antioxidants (Basel) *7*. https://doi.org/10.3390/antiox7110153.

Young, J.D., Shastri, A.A., Stephanopoulos, G., and Morgan, J.A. (2011). Mapping
photoautotrophic metabolism with isotopically nonstationary (13)C flux analysis. Metab. Eng. *13*,
656–665. https://doi.org/10.1016/j.ymben.2011.08.002.

Yu, E.T., Zendejas, F.J., Lane, P.D., Gaucher, S., Simmons, B.A., and Lane, T.W. (2009).
Triacylglycerol accumulation and profiling in the model diatoms *Thalassiosira pseudonana* and
Phaeodactylum tricornutum (Baccilariophyceae) during starvation. J. Appl. Phycol. *21*, 669–681.
https://doi.org/10.1007/s10811-008-9400-y.

Yu, J., Li, Y., Qin, Z., Guo, S., Li, Y., Miao, Y., Song, C., Chen, S., and Dai, S. (2020). Plant
chloroplast stress response: Insights from thiol redox proteomics. Antioxid. Redox Signal. *33*, 35–
57. https://doi.org/10.1089/ars.2019.7823.

Zhao, Y., Luo, L., Xu, J., Xin, P., Guo, H., Wu, J., Bai, L., Wang, G., Chu, J., Zuo, J., et al. (2018).
Malate transported from chloroplast to mitochondrion triggers production of ROS and PCD in
Arabidopsis thaliana. Cell Res. *28*, 448–461. https://doi.org/10.1038/s41422-018-0024-8.

Zhao, Y., Yu, H., Zhou, J.-M., Smith, S.M., and Li, J. (2020). Malate Circulation: Linking
Chloroplast Metabolism to Mitochondrial ROS. Trends Plant Sci. *25*, 446–454.
https://doi.org/10.1016/j.tplants.2020.01.010.

Zimmermann, P., and Zentgraf, U. (2005). The correlation between oxidative stress and leaf
senescence during plant development. Cell. Mol. Biol. Lett. *10*, 515–534. .

1258

1259

1260

1261

1262

1263

1264

1265

1266

1267

1268

Materials and methods

Extraction of samples for LC-MS/MS

To each cell pellet, 500 μL methanol was added along with internal standards ribitol and PIPES to a final concentration of 2 μM each. Samples were then vortexed for 30 sec, frozen in liquid nitrogen as described in (Winder et al. 2008), and allowed to thaw at 0°C. This vortex-freeze-thaw cycle was repeated twice more before the samples were centrifuged at 10,000 x g at 1°C for 5 minutes. The supernatant was collected, and the remaining pellets were extracted twice more with 500 μL of a 50:50 mixture of methanol and water, with 3 vortex-freeze-thaw cycles done for each extraction. Supernatants from each extraction procedure were pooled and dried on a Thermo Fisher SpeedVac Concentrator. Dried extracts were then resuspended in 500 μL water and cleaned to remove any residual large cell debris by filtration, first through a 0.22 μm pore size Spin-X centrifugal tube filter, followed by 10 K molecular mass cutoff filters, followed finally by 3 K molecular mass cutoff filters. After each filtration step, filters were rinsed with 50 μL water, which was added to the total sample volume for subsequent steps. Filtered extracts were dried again and resuspended in 200 μL water for LC-MS/MS analysis.

Analysis of metabolite extracts

Metabolite extracts were analyzed using an LC-MS/MS method adapted from (Young et al. 2011). A Phenomenex 150 mm x 2 mm Synergi Hydro-RP column was used on an Agilent 1200 Series HPLC system coupled to an AB Sciex 5500 QTRAP system. LC was performed with a liquid injection volume of 20 μL, using gradient elution of 10 mM tributylamine and 15 mM acetic acid (aqueous phase) and acetonitrile (organic phase) at a constant flow rate of 0.3 mL/min, and a constant temperature of 40°C. The gradient profile of the organic phase is as follows: 0% B (0 min), 8% B (10 min), 16% B (15 min), 30%B (16.5 min), 30% B (19 min), 90% B (21.5 min), 90% B (28 min), 0% B (28.5 min), and 0% B (35 min). MS analysis was performed in negative mode using a multiple reaction monitoring (MRM) acquisition method. Data acquisition was performed on the ABSciex Analyst 1.7 software. Absolute quantification of intracellular metabolites was performed using the quantitation mode on the Analyst software.

Protein extraction and immunoblot analysis

Affinity-purified polyclonal antibodies to CreTPT3 were custom-made by GenScript. The antigen sequence used for antibody generation was KSWSFGRPVTKQEF. Chlamydomonas cells were grown in liquid cultures to 2–5 x10^6 cells ml^{-1} and collected by centrifugation (2700 rpm, 5 min). Cells were resuspended in resuspension buffer [(5 mM HEPES, pH 7.5, KOH), 10 mM EDTA (pH 7.5, NaOH), 1x protease inhibitor "complete EDTA-free" (Roche)]. For the disruption of cells, bead-beating was performed using a mini beadbeater (Biospec) in two cycles of 30 sec min^{-1}, with a 1 min period of cooling on ice between cycles. Disrupted cells were centrifuged at 4°C, 30 min, 14000 rpm in a microfuge. The supernatant was removed, the pellet was resuspended in sample buffer (resuspension buffer supplemented with 100 mM Na$_2$CO$_3$, 100 mM DTT, 2% WT/vol SDS, and 12% WT /vol sucrose).

SDS-PAGE was performed using a 12% polyacrylamide gel (Bio-Rad), with the electrophoresis
for 90 min at 120 V. Next, proteins were transferred onto a PVDF membrane (Bio-Rad) using
Trans-Blot Turbo (Bio-Rad). The detection was performed using Clarity Max Western ECL
Substrate (Bio-Rad).

RT-qPCR

RT-qPCR, was performed as described previously (Kaye et al. 2019). Briefly, total RNA was
isolated using the RNeasy Plant Mini Kit (Qiagen) and treated with DNase I (Qiagen). First-strand
cDNA was generated by reverse transcription of 0.5 µg total RNA using the iScript cDNA
Synthesis Kit (Bio-Rad). Real-time PCR using the Roche Light Cycler 480 was performed with
the SensiMix NO-Rox SYBR Green I Kit as described by the manufacturer (Bioline). Oligos used
in this research for analyzing the expression levels of *CreTPT1/2/3/25* are listed in **Supplemental**
Table 1.

Analysis of transcript levels in response to different conditions

In order to analyze the transcript level of Cre*TPT* genes under different conditions, we obtained
RNAseq data published previously in which cells experienced a diurnal cycle, nitrogen starvation,
sulfur starvation and Fe^{2+} starvation (Strenkert et al. 2019; Zones et al. 2015; Ngan et al. 2015;
González-Ballester et al. 2010; Urzica et al. 2013). The log2 fold change of each gene was
calculated based on the ratio of transcript abundance at different time point over transcript level at
time point 0 h. The fold change was visualized by a heatmap and the color bar is the scale of log2
(fold-change).

In vivo polymeric carbohydrate staining

Lugol's iodine (aqueous solution of 1.8% iodine and 3.0% potassium iodide) was used to stain in
vivo polymeric carbohydrate. Briefly, 0.5 mL of the cell culture was pelleted, resuspended in 10
µL of Lugol's iodine and visualized under a bright-field microscope. Lugol's iodine selectively
binds to alpha-1,4 glucans and stains it blue-black.

References

González-Ballester, David, David Casero, Shawn Cokus, Matteo Pellegrini, Sabeeha S. Merchant,
and Arthur R. Grossman. 2010. "RNA-Seq Analysis of Sulfur-Deprived Chlamydomonas Cells
Reveals Aspects of Acclimation Critical for Cell Survival." *The Plant Cell* 22 (6): 2058–84.

Kaye, Yuval, Weichao Huang, Sophie Clowez, Shai Saroussi, Adam Idoine, Emanuel Sanz-Luque,
and Arthur R. Grossman. 2019. "The Mitochondrial Alternative Oxidase from *Chlamydomonas*
reinhardtii Enables Survival in High Light." *The Journal of Biological Chemistry* 294 (4): 1380–
95.

Ngan, Chew Yee, Chee-Hong Wong, Cindy Choi, Yuko Yoshinaga, Katherine Louie, Jing Jia,
Cindy Chen, et al. 2015. "Lineage-Specific Chromatin Signatures Reveal a Regulator of Lipid
Metabolism in Microalgae." *Nature Plants* 1 (8): 15107.

Strenkert, Daniela, Stefan Schmollinger, Sean D. Gallaher, Patrice A. Salomé, Samuel O. Purvine,

Carrie D. Nicora, Tabea Mettler-Altmann, et al. 2019. "Multiomics Resolution of Molecular
Events during a Day in the Life of Chlamydomonas." *Proceedings of the National Academy of*
Sciences of the United States of America 116 (6): 2374–83.

Urzica, Eugen I., Astrid Vieler, Anne Hong-Hermesdorf, M. Dudley Page, David Casero, Sean D.
Gallaher, Janette Kropat, Matteo Pellegrini, Christoph Benning, and Sabeeha S. Merchant. 2013.
"Remodeling of Membrane Lipids in Iron-Starved Chlamydomonas." *The Journal of Biological*
Chemistry 288 (42): 30246–58.

Winder, Catherine L., Warwick B. Dunn, Stephanie Schuler, David Broadhurst, Roger Jarvis,
Gillian M. Stephens, and Royston Goodacre. 2008. "Global Metabolic Profiling of Escherichia
Coli Cultures: An Evaluation of Methods for Quenching and Extraction of Intracellular
Metabolites." *Analytical Chemistry* 80 (8): 2939–48.

Young, Jamey D., Avantika A. Shastri, Gregory Stephanopoulos, and John A. Morgan. 2011.
"Mapping Photoautotrophic Metabolism with Isotopically Nonstationary (13)C Flux Analysis."
Metabolic Engineering 13 (6): 656–65.

Zones, James Matt, Ian K. Blaby, Sabeeha S. Merchant, and James G. Umen. 2015. "High-
Resolution Profiling of a Synchronized Diurnal Transcriptome from *Chlamydomonas reinhardtii*
Reveals Continuous Cell and Metabolic Differentiation." *The Plant Cell* 27 (10): 2743–69.

1363

1364

1365

1366

1367

1368

1369

1370

1371

1372

1373

1374

1375

1376

oligo name	sequence
gTPT3_pRam118_f	tactcacaacaagcccagttATGGCTTCGCTCGCTCAGCGCGTGAGC
gTPT3_pRam118_r	agcggggatccttatccgttTTAGGCCGCGGCCGGCGACGC
TPT3pro1000_f	gccttttgctcacatgtggcggccgccgccatgctggccaacag
TPT3pro1000_r	taccaatccacaaatgtgggctagcccatgtgaaggcgttgacc
TPT3_3UTR_f	gatgacgacaagtaaagatctTAAGGCGGGCAGCGGC
TPT3_3UTR_r	GCGCAAGAAAGAAGCTTGATATCCCCCTTATAATGTTTACAAGTGGTGCG
hygro_crispr_f2	GGAACTGGCGCAGTTCCTCTG
hygro_crispr_re2	GGCTCGCGTAGGAATCATCCG
TPT3_gRNA1_F	GCTCGGTTGAGCGCACCC
TPT3_gRNA1_R	CAAGGGTCTGGCGAGCACG
TPT3_gRNA2_F	ACCTCCCCATCGCCACGAT
TPT3_gRNA2_R	AGGGGTAAGGACAGCGGTG
Qrt_TPT1_f2	CCTGTACCAGCAGGTGTCGTAC
Qrt_TPT1_r2	CCAGCGCGTTCTGGAGTGACA
Qrt_TPT2_f2	ATGGCGTCTCTGCTGGGC
Qrt_TPT2_r2	CTGCTCGCCCATTCTCCTC
Qrt_TPT3_f2	GCAAAGCTGGCGCCTACTG
Qrt_TPT3_r2	TAGACCAGGCCGACCACCAC
Qrt_TPT25_fl	CTGGTTCATCGGGACATGGC
Qrt_TPT25_r1	TGGCTGAAGCTGACCACT

1377

1378 **Supplementary Table 1: oligo list.**

1379

Cre ID	Name	Prediction	CS Position
Cre08.g379350.t1.1	CreTPT1	cTP	CS pos: 61-62. TCL-AV.
Cre06.g263850.t1.1	CreTPT2	cTP	CS pos: 70-71. VCQ-AA.
Cre01.g045550.t1.2	CreTPT3	cTP	CS pos: 60-61. VTK-AS.
Cre02.g106200.t1.2	CreTPT4	OTHER	
Cre02.g112900.t1.2	CreTPT5	OTHER	
Cre02.g144300.t1.2	CreTPT6	OTHER	
Cre03.g162000.t1.2	CreTPT7	OTHER	
Cre03.g184850.t1.1	CreTPT8	OTHER	
Cre04.g227450.t1.2	CreTPT9	OTHER	
Cre07.g330850.t1.1	CreTPT10	OTHER	
Cre08.g363600.t1.2	CreTPT11	OTHER	
Cre09.g408400.t1.1	CreTPT12	OTHER	
Cre09.g413700.t1.1	CreTPT13	OTHER	
Cre08.g382350.t1.2	CreTPT14	OTHER	
Cre09.g415900.t1.2	CreTPT15	OTHER	
Cre10.g452750.t1.2	CreTPT16	OTHER	
Cre11.g479950.t1.2	CreTPT17	OTHER	
Cre12.g490050.t1.2	CreTPT18	OTHER	
Cre12.g490100.t1.2	CreTPT19	OTHER	
Cre12.g501000.t1.1	CreTPT20	OTHER	
Cre15.g641266.t1.1	CreTPT22	OTHER	
Cre15.g642950.t1.2	CreTPT23	OTHER	
Cre15.g643385.t1.1	CreTPT24	OTHER	
Cre16.g663800.t1.2	CreTPT25	cTP	CS pos: 70-71. IVA-SS.
Cre16.g666250.t1.2	CreTPT26	OTHER	
Cre17.g702700.t1.1	CreTPT27	OTHER	
Cre17.g703250.t1.2	CreTPT28	OTHER	
Cre17.g710850.t1.2	CreTPT29	OTHER	
Cre11.g467754.t1.1	CreTPT30	OTHER	
Cre14.g622700.t1.2	CreTPT31	OTHER	
Cre18.g748947.t1.1	CreTPT32	OTHER	
Cre09.g408428.t1.2	CreTPT33	OTHER	

Supplementary Table 2: Members of the triose phosphate transporter family in *Chlamydomonas reinhardtii* and their predicted subcellular localization based on TargetP-2.0. cTP, chloroplast transit peptide; CS position, transit peptides cleavage site.

Name	D_11h	D_9h	D_7h	D_5h	D_3h	D_1h	D_0.5h	D_0h	L_0.5h	L_1h	L_3h	L_5h	L_7h	L_9h	L_11h	D_13h
TPT1	5.2883	10.2702	9.45732	8.85549	7.70518	6.18466	7.5509	9.07019	0.511726	0.198621	0.329098	1.99425	2.93498	2.13736	1.02291	9.56813
TPT2	6.46543	9.45107	9.77461	10.5338	12.8513	8.72194	9.51834	10.275	59.1606	62.7836	33.5533	12.891	5.69013	5.94982	12.1866	3.6131
TPT3	233.805	337.793	247.287	181.496	131.602	97.1854	105.57	98.3637	71.9013	43.1935	218.026	280.947	185.973	179.459	157.828	312.135
TPT25	38.9867	62.7835	45.3561	32.7872	22.8554	14.8691	15.5851	13.8916	19.6257	26.8661	10.2213	7.48595	8.11458	19.7874	32.0979	53.4145

Supplemental Table S3: Expression levels of TPTs predicted on the chloroplast envelope over the diel cycle in Chlamydomonas. Data were extracted from (Strenkert et al. 2019).

1388
Supplementary Fig. 1: Experimental set up. All strains were grown photoautotrophically under
low white light ($\sim$30 µmol photons m^{-2} s^{-1}, LL) for 16-24 h. After dilution with fresh medium, the
cultures were transferred to moderate light (250-300 µmol photon m^{-2} s^{-1}, ML). On occasion, the
cultures were also transferred to the dark or high light ($\sim$450 µmol photon m^{-2}s^{-1}) for 24 or 48 h.
To avoid self-shading, the culture density was adjusted to 1-1.5 µg/mL chlorophyll before
exposure to ML or HL.

1395

1396

Supplementary Fig. 2: Sequence alignment of the plastid phosphate transporter sequences
of *Arabidopsis thaliana* and *Chlamydomonas reinhardtii*. Plastid TPTs from Chlamydomonas
are CreTPT1 (Cre08.g379350.t1.1), CreTPT2 (Cre06.g263850.t1.1), CreTPT3
(Cre01.g045550.t1.2), and CreTPT25 (Cre16.g663800.t1.2). Sequences from *Arabidopsis*
thaliana are AtTPT1 (AT5G46110.1), AtPPT1 (AT5G33320.1), AtGPT1 (AT5G54800.1) and
AtXPT1 (AT5G17630.1). Conserved substrate binding pockets are colored red. Phosphate-binding
residues are highlighted with blue. The gate capping residues are marked with yellow. The
positions of residue in green are crucial for the preference of triose-phosphate or 3-PGA. The
positions of purple residues are critical for the specificity of PEP.

1406

1407

Supplementary Fig. 3: Phylogeny of TPT homologues in Chlamydomonas. Amino acid residues at the positions responsible for substrate specificity in TPT homologues are listed. The numbers above the amino acids are the position of each residue in CreTPT3. The residues marked with green, blue and purple colors are crucial for the preference of triose-phosphate or 3-PGA, for the specificity of PEP and for the binding of phosphate, respectively. Plastid PPT, XPT, and TPT from *Arabidopsis thaliana* are included in the analysis.

1414

1415

Supplementary Fig. 4: Background activity of Pi uptake derived from endogenous yeast
transporter reconstituted into liposomes. Time-dependent uptake of Pi (0.25 mM) into
reconstituted liposomes preloaded with 30 mM of Pi (■) or without exchange substrate (●)
prepared from yeast cells harboring the empty vector (pYES-NTa). The arithmetic mean (±SD) of
three independent experiments (each with three technical replicates) was plotted against time. The
observed Pi uptake rates of endogenous yeast carrier reconstituted into liposomes were negligible
compared to the rates of His-CreTPT3 (20-fold higher, **Fig. 1d**), indicating that this expression
system is suitable for functional analysis of recombinant phosphate transporters.

1424

Supplementary Fig. 5: Inactivation of Cre*TPT3* by CRISPR-Cas9. (a) Molecular map of Cre*TPT3* gene and positions of the Cas9 targeted sites. The edited sites are located in exons 1 and 7; *t3ko1* has the marker gene inserted in exon 1 and *t3ko2* and *t3ko3* has the marker gene inserted in exon 7. **(b)** Western blot (left) and loading control (right) of wild type, *t3ko1*, *t3ko2*, *t3ko3* and the complemented strains (C_T3KO1, C_T3KO2 and C_T3KO3, respectively).

1432

1433 **Supplementary**

1434 the drug resistant

1435 *tpt3* mutants (*t3k*

1436

1437

1438

Supplementary Fig. 7: Growth of the CRISPR-Cas9 generated Cre*tpt3* mutants on solid medium. (**a**) Cells were spotted on plates containing TP medium under continuous LL or (**b**) ML. (**c**) Cells were spotted on agar plates containing TAP medium under continuous ML. The dilution series from (**a,b,c**) were 2.0, 0.75, 0.375 µg/mL chlorophyll. (**d**) Growth of wild type (WT), *t3ko1*, *t3ko2*, *t3ko3* and the complemented strains on agar plates containing TAP or TP medium under LL or ML for 4 d. The dilution series used was 1.5, 0.75, 0.375 µg/mL chlorophyll.

1445

1446

Supplementary Fig. 8: (a) Cell size of WT, *t3ko2*, and C_T3KO2 in TP under both LL and ML.
(b) WT and *t3ko2* morphology changes following a LL to ML transition.

1449

1450

1451   **Supplementary Fig. 9: Loss of chlorophyll. Quantification of chlorophyll contents in**
1452 **indicated strains after exposure to ML for 24 and 48 h.** All strains were grown
1453 photoautotrophically under LL (~30 μmol photons m^{-2} s^{-1}) for 16-24 h. After dilution with fresh
1454 medium, the culture was transferred to ML for over 24 h. The initial inoculum density used for the
1455 transition was between 1-1.5 μg/mL chlorophyll in order to avoid self-shading.

1456

1457

1458
Supplementary Fig. 10: (**a**) Starch contents in indicated strains following 0, 6, and 24 h of LL
exposure in TP. (**b**) Time course of change in the metabolite pool sizes at 0 h (LL), 45 min and 6
h after shifting LL-grown cells to ML. Data was normalized to cell numbers. (**c**) Schematic
representation of the fold-change for the metabolites shown in (**b**), calculated by dividing the pool
size in *t3ko2* by that of WT cells under the same conditions. Fold change of 1 (no change) is shown
by a dashed line. Each data point is the mean and standard error of three biological replicates.
Asterisk indicates statistically significant differences relative to WT (* P<0.05, ** P<0.01, ***
P<0.001). Abbreviations: G6P, glucose-6-phosphate; FBP, fructose bisphosphate; DHAP,
dihydroxyacetone phosphate; 3-PGA, 3-phosphoglycerate; PEP, phosphoenolpyruvate; CIT,
citrate; SUCC, succinate; FUM, fumarate; MAL, malate; S7P, sedoheptulose-7-phosphate; E4P,
Erythrose 4-phosphate; R5P, Ribose 5-phosphate; RU5P/X5P, ribulose 5-phosphate/xylulose-5-
phosphate;

1471

Supplementary Fig. 11: Quantum yield of PSII (ΦPSII) grown in LL (0 h: left) and after 6 and
24 h in ML (right) following exposure to increasing actinic light intensities.

1474

Supplementary Fig. 12: Redox state of the PQ pool followed by LL to ML transition. 1-qL
values of LL (left) and ML (right) acclimated cells exposed to increasing light intensities.

1477

1478

1479

| | BF | roGFP2 | Chl | Chl&roGFP2 |

a RoGFP2 in chloroplast

b roGFP2 in mitochondria

1480

1481 **Supplementary Fig. 13:** Redox-sensitive GFP (roGFP) protein was targeted to the chloroplast (**a**)
1482 and mitochondrion (**b**). BF: bright field; roGFP2 fluorescence: green; Chlorophyll (Chl)
1483 autofluorescence: red.

1484

1485 **Supplementary Fig. 14:** WT and *t3ko2* morphology changes followed by a LL (~30 μmol photons
1486 $m^{-2} s^{-1}$) to dark transition.

1487

1488

1489

Supplementary Fig. 15: (**a**) mRNA levels of TPTs predicted to be on the chloroplast envelope under LL during growth in TP medium. mRNA levels of CreTPT1 (**b**), CreTPT2 (**c**) and CreTPT25 (**d**) in the indicated strains 0, 2, 4 h after the cells were shifted from LL to ML.

1493

1494

Supplementary Fig. 16: Expression of chloroplast Chlamydomonas TPTs under diurnal cycle (**a**), nitrogen starvation (**b**), sulfur starvation (**c**), and (**d**) iron starvation (Zones et al. 2015; Ngan et al. 2015; González-Ballester et al. 2010; Urzica et al. 2013).

1498