<u>Title page</u>

Title

naRNA is a canonical neutrophil extracellular trap (NET) component

and novel inflammation-amplifying composite DAMP

Running title

naRNA – canonical NET component and composite DAMP

Authors

Francesca Bork[1], Carsten L. Greve[1], Christine Youn[2], Sirui Chen[1], Yu Wang[2], Masoud Nasri[3], Jule
Focken[4], Jasmin Scheurer[4], Pujan Engels[1], Marissa Dubbelaar[1,5], Katharina Hipp[6], Birgit Schittek[4,7,8],
Stefanie Bugl[1], Markus W. Löffler[1,7,9], Julia Skokowa[3,7], Nathan K. Archer[2] (ORCID: 0000-0002-8212-
8985), Alexander N.R. Weber[1,7,8]* (ORCID: 0000-0002-8627-7056)

Author information

[1]Interfaculty Institute for Cell Biology, Department of Immunology, University of Tübingen, Auf der
Morgenstelle 15, 72076 Tübingen, Germany

[2]Department of Dermatology, Johns Hopkins University School of Medicine, Baltimore, MD 21231,
USA

[3]Division of Translational Oncology, Department of Oncology, Hematology, Clinical Immunology and
Rheumatology, University Hospital Tübingen, Otfried-Müller Str. 10, 72076 Tübingen, Germany

[4]Division of Dermatooncology, Department of Dermatology, University Hospital Tübingen,
Liebermeisterstr. 25, 72076 Tübingen, Germany

[5]Quantitative Biology Center (QBiC), University of Tübingen, Auf der Morgenstelle 10, 72076
Tübingen, Germany

[6]Electron Microscopy Facility, Max Planck Institute for Biology Tübingen, Max-Planck-Ring 5, 72076
Tübingen, Germany

[7]iFIT – Cluster of Excellence (EXC 2180) "Image-Guided and Functionally Instructed Tumor Therapies",
University of Tübingen, Germany

[8]CMFI – Cluster of Excellence (EXC 2124) "Controlling microbes to fight infection", University of

Tübingen, Germany.

[9]Department of General, Visceral and Transplant Surgery, University Hospital Tübingen, Hoppe-

Seyler-Str. 3, 72076 Tübingen, Germany and Department of Clinical Pharmacology, University

Hospital Tübingen, Auf der Morgenstelle 8, 72076 Tübingen, Germany

Corresponding Author

Alexander N. R. Weber, Interfaculty Institute for Cell Biology, Department of Immunology, University

of Tübingen, Auf der Morgenstelle 15, 72076 Tübingen, Germany. Tel.: +49 7071 29 87623. Fax: +49

7071 29 4579. Email: alexander.weber@uni-tuebingen.de; ORCID: 0000-0002-8627-7056, Twitter

@aweberlab

Abstract

Neutrophil extracellular traps (NETs) have emerged as a key feature of cellular innate immunity mediated by polymorphonuclear neutrophils (PMNs), the primary leukocyte population in humans. Forming web-like structures composed of DNA, histones, and antimicrobial proteins, NETs trap and kill microbial invaders and thus enhance host defense. However, they have also been linked to inflammatory states, e.g., in atherosclerosis or psoriasis. Whilst DNA has been in focus as a primary structural component of NETs, we here characterize naRNA (NET-associated RNA), as a new canonical, abundant, and largely unexplored NET component. naRNA decorated all types of NETs in complex with the antimicrobial peptide LL37. In fact, naRNA was pre-associated with LL37 intracellularly as a 'composite' danger-associated molecular pattern (DAMP) prior to neutrophil activation. Externalized, naRNA propagated NET formation in naïve PMN, dependent on TLR8 in humans and Tlr13 in mice, in vitro and in vivo. TLR8/Tlr13 signaling contributed significantly to the highly sensitive pro-inflammatory response to naRNA, involving tissue cells, like keratinocytes, and other immune cell types, such as macrophages. Those responses could be blocked by inhibition and genetic ablation of RNA receptors or RNase treatment. Importantly, in vivo naRNA strongly drove skin inflammation whereas genetic ablation of RNA sensing drastically ameliorated skin inflammation in the imiquimod psoriasis model. Our data highlight naRNA as a novel composite DAMP signaling and amplifying neutrophil activation. Moreover, naRNA emerges as the likely driver of inflammation in conditions previously linked to NETs and extracellular RNA, suggesting blockade of TLR-mediated RNA sensing as potential new intervention target.

Graphical abstract

Created   with   biorender.com

Introduction

The formation of Neutrophil Extracellular Traps (NETs), since its discovery in 2004 [1], has emerged as a

fascinating phenomenon of host defense. Hereby neutrophils, the primary leukocyte population,

extrude their genomic DNA to form web-like structures that, similar to barbed wire roadblocks, trap

and kill microbial invaders such as bacteria or fungi [2]. DNA is thus a defining structural and functional

feature of NETs. Additionally, DNA-associated proteins, histones and HMGB-1, antimicrobial peptides

like LL37 and enzymes such as myeloperoxidase (MPO) are released during the formation of NETs

and contribute to their antimicrobial function [2]. More than 1,000 publications on NETs have sought

to detail the multi-faceted phenomenon and the processes that lead to its execution [3]. However,

with the main focus devoted to DNA and protein components, another primary cellular biomolecule,

RNA, has so far received very little attention in vertebrates. Interestingly, a study in insects showed

that hemocytes (macrophage-like immune cells) can release both DNA and RNA in NET-like

structures during microbe-triggered clotting reactions, and in response to extracellular RNA or DNA [4].

Our lab recently provided evidence of a responsiveness of human polymorphonuclear neutrophils

(PMNs) to RNA, but not DNA, in combination with the antimicrobial peptide LL37, fueling neutrophil

activation [5]. In addition, earlier work described a role for DNA- and RNA-LL37 complexes in the

activation of plasmacytoid dendritic cells [6, 7]. This has given rise to the notion that under certain

conditions tolerance to self-DNA and RNA can be broken[8], involving the nucleic acid-sensing TLR7,

TLR8 (ssRNA) and TLR9 (DNA) in humans, and Tlr7, Tlr13 and Tlr9 in mice [9, 10]. In general, many roles

have been ascribed to so-called extracellular RNA (exRNA), for example, macrophage polarization,

recruitment of leukocytes to the site of inflammation, leukocyte rolling at the vascular endothelium,

as well as integrin-mediated firm adhesion of immune cells and promotion of thrombosis [11].

Additionally, exRNA is considered as a reliable biomarker for various diseases such as cancer or

cardiovascular pathologies [12, 13]. However, RNA being considered a short-lived biomolecule,

physiological sources of exRNA have been somewhat unclear. Under sterile conditions, vascular

injury, tissue damage, or ischemia have been suggested to trigger release of exRNA along with other

cellular material [11]it remains unclear it remains unclear whether these relatively slow processes could

amount to physiologically detectable exRNA amounts in the serum of patients, or whether a rapid,

as-yet unidentified process of RNA extrusion would have to be postulated.

We speculated that RNA contained in NETs, so-called NET-associated RNA or naRNA, might lead to a

feed-forward loop of rapidly expanding NET release amongst neutrophils which might also engulf

additional tissue or immune cells able to respond to exRNA. We show here that naRNA drives such a

self-amplifying loop not only engaging PMNs but also macrophages, and keratinocytes. naRNA

responsiveness was dependent on *TLR8* and *Tlr13* in human and murine myeloid immune cells,

respectively. Interestingly, naRNA appears pre-complexed to LL37 as a novel composite DAMP.

Notably, in mice naRNA caused considerable skin inflammation in a Tlr13-dependent manner.

Additionally, in a well-established model of psoriatic skin inflammation, genetic ablation of RNA

sensing strongly ameliorated skin inflammation, highlighting naRNA as a potent damage associate

molecular pattern (DAMP) amplifying PMN activation and alerting tissue and other immune cells.

Results

naRNA is a common component of NETs

To first investigate whether naRNA was a common component of NETs, we compared NETs from

human PMNs formed in response to the well-defined molecular agonists PMA, different complexes

of LL37 with purified RNA (synthetic, as well as total fungal and bacterial RNA from *S. aureus* and *C.*

albicans, respectively), nigericin and the live pathogen *C. albicans*. Confocal microscopy analysis

using the well-characterized rRNA-specific antibody, Y10b [14], revealed the presence of RNA in all

corresponding NETs, independent of stimulus and whether NET formation was 'suicidal' or not [3] (Fig.

1A, quantified in B; control in Fig. S1A): the RNA signal was readily detectable along the web-like DNA

threads in human NETs. Similar results were obtained in NETs released by murine bone marrow-

derived PMNs (BM-PMNs, Fig. 1C; control Fig. S1B). To avoid the use of staining reagents, we also

metabolically labeled primary human hematopoietic stem cells (HSCs) with 5-ethynyluridine (5-EU), a

nucleotide that can be incorporated in cellular RNA but not DNA, and that is amenable to labeling by

click chemistry [15]. These HSCs were then differentiated to neutrophils [16] and the differentiation

validated by microscopy and flow cytometric analysis (Fig. S1C, D). NETs released from HSC-derived

PMNs could indeed be labeled with click reagents conferring a fluorescent label only when grown in

5-EU-containing medium (Fig. 1D). Moreover, the overlap between click-label and rRNA confirmed

the specificity of the staining with anti-rRNA antibodies and indicated that naRNA contained rRNA

but also other types of cellular RNAs. This experiment also unequivocally confirmed that upon PMN

stimulation, cellular RNA is turned into a component of the NET, wherein it decorates the fine, web-

like DNA structures. The localization of naRNA on DNA strands was further confirmed by high-

resolution fluorescence microscopy and 3D analysis (AiryScan, Fig. 1E and Supplemental Movie S1) as

well as scanning electron microscopy (SEM, Fig. 1F; controls Fig. S1E). Extraction of naRNA from PMA-

induced NETs (Fig. 1G) and subsequent RNA-sequencing analysis confirmed that purified naRNA

contained multiple RNA types, with noticeably more non-ribosomal RNA than the corresponding

total cellular RNA of PMNs (Fig. 1H). Collectively, these results indicate that naRNA is a canonical

component of NETs released upon multiple stimuli from both human and murine neutrophils.

naRNA is a potent 'composite' DAMP propagating NET formation in primary PMNs

A primary function of NETs is host defense by trapping and killing bacteria. Therefore, we first

explored if naRNA participated in this process [1]. However, the antibacterial effect of NETs on live

Staphylococcus aureus (evidenced by lower CFUs compared to resting PMNs) was similar to that of

RNase-treated NETs, whereas DNase digestion reduced the drop in CFU discernibly but non-

significantly (Fig. S2A). Consequently, we turned our attention to a possible role of naRNA as a

DAMP, since primary PMNs can respond to synthetic RNA-LL37 complexes with NET formation, and

our earlier analysis indicated that NETs also contain LL37 [5]. To prepare a naRNA-containing stimulant,

PMA-induced NETs were prepared as described in Methods and the original stimulus, PMA, removed

by extensive washing before harvesting. Interestingly, these 'PMA NETs', when applied to naïve

PMNs from another donor, potently induced new NETs (Fig. 2B, quantified in C, controls in Fig. S2B).

Treatment with an RNase inhibitor enhanced the stimulatory effect of PMA NETs rendering a 1:500

dilution of PMA-NETs more effective than a 1:50 dilution of non-treated PMA-NETs (Fig. 2B, C,

controls Fig. S2B). Interestingly, similar NET preparations from mock-treated PMNs (here referred to

as 'mock NETs') did not stimulate NET formation under the same experimental conditions. To rule

out the possibility that residual PMA in PMA NETs could be responsible for the observed stimulatory

effects, RNase treatment was also performed which strongly reduced NET formation (Fig. 2D,

quantified in E; controls Fig. S2C).. The destruction of naRNA in NETs was further confirmed by rapid

loss of the RNA signal (using the RNA-selective dye, SYTO RNAselect [5] in time-lapse digestion analysis

(Supplemental Movie S2). Bearing in mind that PMNs do not respond to DNA or DNA-LL37 complexes

[5], the opposing effects of RNase inhibitor and RNase treatments thus clearly indicated naRNA (and

not DNA) to be the relevant immunostimulatory component for NETs to drive the activation of naïve

PMNs. In line with our previous work showing that synthetic RNA or LL37 alone cannot trigger PMN

activation [5], purified 'naked' naRNA isolated from PMA NETs (*cf.* Fig. 1G) was unable to activate NET

formation; however, re-complexing with exogenous LL37 was sufficient to restore NET formation

(Fig. 2F, controls in Fig. S2D). We next investigated whether in NETs, naRNA co-localized with LL37.

Strong co-localization was evident in confocal microscopy analysis (Fig. 2G) and confirmed by

Pearson's colocalization analysis (Fig. 2H). Additional line plot analysis of PMA-stimulated PMN

revealed that RNA and LL37 did in fact co-localize in NETs, i.e., after extrusion (Fig. S2E). Surprisingly,

in unstimulated PMNs naRNA and LL37 showed even greater colocalization (Fig. 2H) and shared co-

localization maxima, suggesting pre-association before extrusion (Fig. 2I, Fig. S2E). rRNA and LL37

double staining on ultrathin (50-60 nm) sections of resting PMN confirmed the reported localization

of LL37 to neutrophil granules [17], but also the unexpected concomitant presence of naRNA in the

same structures (Fig. 2J). Transmission electron microscopy combined with double staining of rRNA

Y10b and LL37 using two different immunogold sizes corroborated this further at even higher

magnification (Fig. 2K, Fig. S2F). The concept of pre-packaging of RNA with LL37 in specific

compartments also fits with the RNAseq data showing that certain RNA types are lost upon release

(*cf.* Fig. 1H), probably because they are not bound to LL37. Collectively, naRNA appears to be LL37-

pre-associated, and hence a 'composite' DAMP, enabling PMN activation and NET propagation.

naRNA DAMP activity is dependent on RNA sensors TLR8/Tlr13

To validate naRNA as an immunostimulatory RNA component from the receptor side, we turned to

HEK293T cells which do not respond to RNA, unless transfected with plasmids encoding the human

ssRNA sensors TLR7 or TLR8 [18]. NF-κB reporter assays revealed that TLR7 or TLR8- (both ssRNA

sensors) but not TLR9 (human DNA sensor)-transfected HEK293T cells stimulated with PMA NETs or

mock NETs showed robust NF-κB activity only for PMA NETs (Fig. 3A). R848 (TLR7 and TLR8 agonist)

and TL8 or ssRNA+DOTAP (both TLR8 agonists) served as controls. Of note, compared to its cognate

agonist, CpG, the DNA sensor TLR9 was only poorly activated by PMA NETs, indicating that naRNA is a

superior immune stimulant compared to NET DNA in this system. Indeed, the observed NET response

of primary PMNs (Fig. 3B, quantified in C, and Fig. S3A) could be completely blocked by the TLR8-

specific inhibitor, CU-CPT9a [19], similar to the effects of the non-specific inhibitor, Cl-amidine, a PAD4

inhibitor which also blocked the response to PMA (Fig. 3C and S3A). This revealed the RNA sensor

TLR8 [20] to be the specific naRNA receptor in primary human PMNs. We further genetically validated

the involvement of RNA sensing in naRNA-mediated NET propagation using BM-PMNs from *Unc93b1-*

or *Tlr13*-defective mice: whereas in *Unc93b1*-defective mice signaling of all endosomal TLRs is

abrogated [21], mice lacking *Tlr13,* the murine equivalent of TLR8 [9], specifically lack ssRNA sensing [22].

Confocal microscopy analysis showed that WT BM-PMNs responded readily to both PMA (used as

control) and PMA NETs, whereas *Unc93b1* (Fig. S3B) or *Tlr13* KO PMNs (Fig. 3D, quantified in E) only

responded to the TLR-independent PMA control but not PMA NETs. Additionally, naïve BM-PMNs

from *Tlr9* KO mice, which lack DNA sensing, responded to both PMA and PMA NETs with NET

formation just like WT BM-PMNs (Fig. S3C), emphasizing the redundancy of DNA vs the key role of

naRNA in the self-propagating properties of NETs. This receptor analysis unequivocally confirmed

naRNA to be a primary immune stimulant in NETs.

PMN-derived naRNA triggers broader immune cell and keratinocyte activation

Given the potent effect of naRNA on naïve PMNs (*cf.* Fig. 3B-E) and the ability of human

macrophages to sense RNA via TLR8 [23], we hypothesized that naRNA might also directly activate

macrophages, which are rapidly recruited at the sites of PMN activation in peripheral tissues [24],

contributing to an inflammatory process in vivo. We thus assessed the effects of naRNA on

genetically modified macrophage-like cell lines: BlaER1 macrophages [25] responded to PMA NETs but

not mock NETs with IL-8 release, and this was TLR8-dependent as evidenced by comparisons of WT

and *TLR8*-edited BlaER1 cells (Fig. 3F). Use of a naRNA-stabilizing RNase inhibitor during NET

preparation (*cf.* Fig. 2B) drastically increased the potency of PMA NETs on these macrophages.

Additionally, TLR8 and TLR7-edited THP-1 macrophage-like cells [26] responded to PMA NETs with

lower relative IL-8 release than WT THP-1 cells (Fig. S4A), confirming RNA as the active agent in

monocyte/macrophage activation. Human peripheral blood mononuclear cells (PBMCs) were also

stimulated with different NET preparations. RNA-stabilized PMA NETs consistently elicited higher

TNF, IL-6 and IL-8 levels than non-stabilized PMA- or mock NETs (Fig. S4B-D). However, this effect was

not sensitive to the TLR8 inhibitor Cu-CPT9a, which is consistent with the ability of TLR7 to also sense

naRNA (*cf.* Fig. 3A) and possibly due to a mixture of cells being present. Further analysis revealed that

the natural killer (NK) cell line NK-92 MI showed low IFNγ release in response to RNA-stabilized PMA

NETs (Fig. S4E). Collectively, our results show that the RNA component of NETs, naRNA, can engage

not only bystander neutrophils in a feed-forward inflammatory response but also other myeloid

(monocytes/macrophages) and lymphoid (NK cells) cellular innate immunity. To investigate whether

non-hematopoietic cells with immune functions, e.g. tissue cells like keratinocytes, could also

respond to RNA-stabilized PMA NETs, N/TERT-1 keratinocytes [27] in monolayers were exposed to

PMA-NETs (Fig. 3G). This triggered a dose-dependent IL-8 release comparable to synthetic RNA+LL37,

especially under RNA stabilization by RNase inhibitors. Similar results were obtained using primary

normal human epidermal keratinocyte (NHEK) monolayers (Fig. 3H). Furthermore, in an 'in vivo'-like

human skin equivalent 3D model in which natural keratinocyte differentiation was recapitulated by

NHEK cells [28], PMA but not mock NETs significantly induced *IL8* mRNA and protein (Fig. 3I, J). Thus,

naRNA can not only activate primary neutrophils in vitro but also trigger broad immune activation in

other immune and tissue cells/equivalents.

naRNA displays DAMP activity in vivo in dependence on RNA sensing

Moreover, to gain an insight into whether naRNA could trigger inflammation in vivo in an RNA

receptor-dependent manner, we intradermally injected RNase inhibitor-stabilized mock or PMA NETs

in the ears of C57BL6 mice. RNA-stabilized PMA NETs were almost as potent to induce ear swelling as

synthetic RNA+LL37 (Fig. 4A), but even non-stabilized PMA NETs induced a response significantly

higher than mock NETs. The same stimulants were also injected into the ear of *LysM*[EGFP/+] mice, in

which myeloid cells are GFP-positive, thus enabling in vivo monitoring of cellular influx during skin

inflammation. Here, RNA-stabilized PMA NETs elicited even greater cellular influx than synthetic

RNA+LL37 and mock NETs (Fig. 4B). Importantly, a comparison of WT and *Tlr13*-deficient animals

showed that the ear swelling reaction resulting in vivo was naRNA-dependent because RNA-

stabilized PMA-NETs were significantly less effective in *Tlr13*-deficient animals (Fig. 4C).

RNA recognition contributes to progressive skin inflammation in experimental psoriasis

Finally, we sought to explore if naRNA was relevant in an animal disease model. The most frequently
used murine model of psoriasis uses the TLR7 ligand imiquimod (IMQ) to trigger increasing skin
inflammation characterized by epidermal thickening and immune cell infiltration [29]. We previously
showed that IMQ treatment also leads to the occurrence of NETs in the tissue [5]. We therefore
speculated that naRNA might be involved in disease progression, a hypothesis that could be probed
by genetically ablating naRNA sensing and comparing WT and *Tlr13* KO mice. Although in the early
induction phase, WT and *Tlr13* KO animals showed a similar increase in ear thickness, the groups
diverged from day 3 onwards, after which *Tlr13*-deficient animals were significantly protected from
skin inflammation compared to WT mice (Fig. 4D). Additionally, the characteristic epidermal
thickening upon IMQ treatment was significantly lower in *Tlr13*-deficient mice compared to WT mice
(Fig. 4E, representative image Fig. 4F). The most plausible explanation is that IMQ-initiated NET
formation [5] amplifies skin inflammation in WT animals via naRNA, whereas this is prevented in *Tlr13*
KO animals. This shows that naRNA can significantly contribute to inflammation in a well-established
disease model and can act as a potent inflammation-amplifying endogenous DAMP.

Discussion

At first sight, it may not seem surprising that the release of NETs, a process that churns up the most
critical compartment of a cell, e.g., the nucleus, also inevitably leads to the release of another
primary cellular biomolecule, namely RNA. However, independently of whether the process is
regulated or not, the release of RNA by NETting PMNs appears to have physiological importance as
we have demonstrated here. Rather than acting directly in antimicrobial defense (*cf.* Fig. S2A),
naRNA appears to be an LL37-precomplexed DAMP of PMN origin that can be released in the NETting
process, and then activates both PMN and other immune and tissue cells in an RNA sensor-
dependent manner (Fig. S5). Several further findings warrant further discussion:

Firstly, our data revise previous concepts of self-RNA and LL37 in the 'breaking' of innate immune
tolerance in psoriasis and the physiological role of NETs in general. naRNA when complexed with
LL37, gains stimulatory properties on neutrophils, in line with the existing notion that LL37 may
'break' immune tolerance by acting as a physiological 'transfection reagent' for uptake into immune
cells, and by shielding it from RNase degradation. However, RNA and LL37 do not appear to
aggregate and co-localize 'accidentally' in NETs: quite unexpectedly, they were rather found in even
greater co-localization in resting neutrophils, indicating a kind of 'pre-association' and intracellular
storage of a pre-defined naRNA-LL37 'composite' DAMP before extrusion during NET formation. It
will be interesting to study the process for pre-association for this first 'composite DAMP' and
elucidate the precise nature of the RNAs assembled with LL37 in the future Importantly, the
identification of such pre-association challenges existing concepts of LL37-RNA complexes as

'accidentally assembled' tolerance breakers in certain diseases [7]. Rather, it suggests naRNA-LL37

complexes act as purposefully pre-associated DAMPs that arm neutrophils at steady state.

Intracellular (rather than accidental/extracellular) pre-association points to a specific role of the

naRNA-LL37 DAMP in NET biology and may indicate the NET response to be a DAMP response, rather

than an antimicrobial reponse, especially in sterile conditions. Beyond simply amplifying

inflammation, we speculate that the physiological relevance of deploying this composite DAMP upon

NET formation could be to tag fresh NETs with a timed molecular beacon: Freshly extruded NETs

decorated with the naRNA-LL37 DAMP would likely highlight the lingering presence of a pathogenic

microorganism or acute tissue damage to other cells not directly engaged by the threat. The naRNA-

LL37 DAMP would thus label 'fresh NETs' as 'requiring attention' and triggering further immune

activation. Over time, whilst the DNA-related structural and antimicrobial properties of NETs may

remain longer, inevitable RNA degradation would lead to deactivation of the composite DAMP,

rendering 'old NETs' less immunostimulatory. Further work outside the scope of the present study

may explore such a role of naRNA as time-restricted molecular label for NETs.

Moreover, our data pertain to the origin and effects of 'extracellular RNA' (exRNA). exRNA is a

generic term to indicate a heterogeneous group of RNA molecules which are actively or passively

released during sterile inflammation or infectious processes. exRNA can be released in a 'free' state,

bound to proteins or phospholipids, in association with extracellular vesicles (EVs) or apoptotic

bodies [11]. In all these forms, exRNA may function as a DAMP but also as e.g., procoagulant or

regenerative factor [11]. Our data identify naRNA as the first type of exRNA for which a clear

physiological origin (NETs) is provided. We speculate that our findings will help trace multiple

descriptions of exRNA, on the one side, to neutrophil traps on the other. For example, exRNA has

emerged as disease-relevant in atherosclerosis, where it was described to act as a proinflammatory

mediator enhancing the recruitment of leukocytes to the site of atherosclerotic lesions as shown in a

mouse model of accelerated atherosclerosis [30]. At the same time, NETs have been ascribed a role in

amplifying sterile inflammatory responses in an independent mouse model of atherosclerosis by

priming macrophages [31]. However, never have these two independent strains of research been

connected. By showing NETs to release naRNA, a DAMP type of exRNA, our work connects both lines

of enquiry. Likewise, for rheumatoid arthritis (RA), our work makes a plausible link between exRNA in

synovial fluid contributing to joint inflammation [32], the emerging role of NETs in RA [33] and even a

hitherto enigmatic but therapeutically relevant role of TLR8 [34, 35].

Moreover, NETs have been of broad interest focus in COVID-19-related research regarding thrombo-

inflammatory states, like sepsis, thrombosis, and respiratory failure, , have been found in plasma of

hospitalized COVID-19 patients, and linked to thrombus formation by several research groups [36] [37] [38]

[39, 40]. Furthermore, exRNAs have been in focus as procoagulant cofactors in blood coagulation [41] [42]. A

link between NET-related and RNA-induced thrombus formation in COVID-19 has never been made
but highlightss naRNA as a potential contributing factor of NET-induced thrombo-inflammatory
states in COVID-19 disease pathogenesis. Collectively, we suspect similar links between NETs and
exRNA via naRNA to emerge for cardiovascular diseases and cancer if the role of naRNA is thoroughly
assessed. Furthermore, the broad sensitivity of immune and tissue cells to naRNA observed by us
makes sense of how exRNA may act pathophysiologically.

Although we believe PMNs to be the primary trap forming human leukocyte population and hence
naRNA the most significant "trap-associated RNA", it will be interesting to explore whether mast cell
(MCETs) [43] or macrophage extracellular traps (METs) [44] contain RNA. However, unlike PMN, the latter
immune cells are not primary sources of LL37 [17], so that RNA associated with MCETs or METs would
be of lesser physiological relevance as a DAMP than naRNA due to a lack of LL37. Therefore,
translational approaches, e.g., to restrict trap-RNA mediated amplification of inflammation, should
probably center on PMN-derived naRNA. The use of PAD4 inhibitors has already been investigated in
animal models to treat cancer or atherosclerotic lesions [45 46] and would represent one potential way
of eliminating NETs and hence naRNA-mediated inflammation. However, this would also prevent the
effects of NETs that are beneficial for host defense (e.g. physical trapping via DNA and antimicrobial
enzymatic activities) and may render treated patients vulnerable to infections. From a translational
perspective, our in vivo data indicate that blockade of RNA sensing might be more advantageous,
restricting only naRNA mediated responses. Although probably not applicable to patients, it is
underscored by evidence that neuroinflammation upon subarachnoid hemorrhage, which is
characterized by a NET pathology, is strongly ameliorated by RNase treatment in vivo [47]. More
applicable to patients may be the targeting of TLR8 by small molecular antagonists [48] or inhibitory
oligonucleotides, which are able to block PMN responsiveness to synthetic RNA in vitro [5]. Exciting is
thus the observed efficacy of TLR inhibitory oligoribonucleotides in a psoriasis mouse model [49] and
even a first clinical trial in psoriasis patients [50]. We could imagine the blockade of naRNA effects via
TLRs highlighted here to emerge as an effective novel strategy to target multiple exRNA or NET-
related inflammatory responses.

<u>Figure legends</u>

Fig. 1: **naRNA is a canonical component of NETs.**

(A, B) Confocal microscopy of primary human PMNs stimulated as indicated for 3 h and stained for

naRNA (anti-rRNA Y10b, magenta) and DNA (Hoechst 33342, white, n = 3, scale bar: 10 μm,

arrowheads point to selected NET strands; representative images in **A**) were quantified in **B** (each dot

represents one image, *p<0.05 according to one-way ANOVA). **(C)** Confocal microscopy of primary

murine BM-PMNs stimulated as indicated for 16 h and stained as in **A** (n = 3, representative images,

scale bar: 10 μm). **(D)** Confocal microscopy of primary human stem cells differentiated *in vitro*

with/without 100 μM 5-ethynyluridine (5-EU), click-labeled with a fluorescent dye (yellow, total

RNA), anti-rRNA and Hoechst 33342 as in **A** (n = 3, representative images, scale bar: 10 μm, 2 μm in

cropped image). **(E)** as in **A** but showing 3D image reconstruction of z-stacks (n = 3, representative

images, scale bar: 10 μm). **(F)** Scanning electron microscopy of human primary PMNs treated as

indicated and using anti-rRNA primary and immunogold (white arrow)-labeled secondary antibodies

and silver enhancement (n = 3, representative images, scale bars as indicated; the two rightmost

images show composite images with signals from secondary electron and backscattered electron

detectors for topography and additional material information, respectively). **(G)** Agilent TapeStation

quantification of naRNA isolated from mock or PMA NETs (from n = 4-6 donors, combined data,

mean+SD, each dot represents one biological replicate/donor, *p<0.05 according to Mann-Whitney

test). **(H)** RNAseq of PMA NET naRNA (n = 4) and whole PMN RNA (n = 1) (combined data).

Fig. 2. naRNA is a composite DAMP driving NET propagation in human PMN

(A) Workflow for NET content preparation from one donor and transfer to naïve human primary

PMNs from a second donor. **(B)** Confocal microscopy of primary human PMNs stimulated with NET

content harvested with/without RNase inhibitor and diluted 1:50 or 1:500, and then stained for

NETs/DNA (Hoechst 33342, n = 9, representative images, scale bar: 10 μm). **(C)** Quantification of **B**

using DNA (Hoechst 33342) signal to quantify NET formation (n = 3, combined data, mean+SD, each

dot represents one image, *p<0.05 according to one-way ANOVA). **(D)** as in **B** but with/without pre-

digestion of NET content with RNase A (n = 3, representative images, scale bar: 10 μm). **(E)**

Quantification of **D** (n = 3, combined data, mean+SD, each dot represents the number of NET-positive

tiles in one image quantified from three images/condition, *p<0.05 according to one-way ANOVA).

(F) As in **B** but using purified naRNA (*cf.* Fig. 1G) alone or in complex with exogenously added LL37 (n

= 3, representative images, scale bar: 10 μm). **(G)** Confocal microscopy of primary human PMNs

stimulated as indicated for 3 h and stained for naRNA (anti-rRNA Y10b, magenta), LL37 (anti-hLL37-

DyLight550, yellow), and DNA (Hoechst 33342, white, n = 3, representative images, scale bar 10 μm).

(H) Pearson's correlation coefficient (colocalization) analysis of **G** (n = 3, combined data, mean+SD,

each dot represents one image, three images/condition, *p<0.05 according to one-way ANOVA). **(I)**

Line plot analysis of LL37, RNA and DNA staining of unstimulated primary human PMNs of **G** was

performed using ZenBlue3 software (n=3, representative graph). **(J)** As in **G/I** but on ultrathin

sections of unstimulated PMNs (n=3, representative image, scale bar 2 µm). Line plot analysis was

performed using ImageJ-Win64 software (n=3, representative graph). **(K)** Transmission electron

microscopy of unstimulated human primary PMNs using anti-rRNA and anti-hLL-37 primary and

immunogold (6 nm (black arrow) and 12 nm (white arrow) respectively)-labeled secondary

antibodies (n = 3, representative images, scale bars as indicated).

Fig. 3. naRNA activity depends on RNA sensors

(A) NF-κB dual luciferase assay in HEK293T cell transiently transfected and stimulated as indicated

(eV = empty vector, n=3-5, combined data, mean+SD, *p<0.05 according to one-way ANOVA). **(B)**

Confocal microscopy of human primary PMNs stimulated as indicated in the presence or absence of

CU-CPT9a (100 nM) or Cl-amidine (200 µM, not shown) and stained for NETs/DNA (Hoechst 33342,

white, n = 3, representative images, scale bar: 10 µm). **(C)** Quantification of **B** using DNA (Hoechst

33342) signal to quantify NET formation (n = 3, combined data, mean+SD, each dot represents the

number of NET-positive tiles in one image, three images/condition, *p<0.05 according to non-

parametric one-way ANOVA). **(D)** Confocal microscopy analysis of primary C57BL/6 WT or *Tlr13*$^{-/-}$

murine BM-PMNs stimulated as indicated and stained for NETs/DNA (Hoechst 33342, white, n = 3,

representative images, scale bar: 10 µm). **(E)** Quantification of **D** as in **C** (n = 3, *p<0.05 according to

one-way ANOVA). **(F)** Triplicate IL-8 ELISA from WT, *TLR8*$^{-/-}$ and *UNC93B1*$^{-/-}$ BlaER1 macrophage-like

cells stimulated as indicated for 18 h (n = 3-4, combined data, mean+SD, each dot represents one

biological replicate, *p<0.05 according to one-way ANOVA). **(G)** Triplicate IL-8 ELISA from N/TERT-1

keratinocytes stimulated as indicated for 24 h (n = 3, combined data, mean+SD, each dot represents

one biological replicate, *p<0.05 according to one-way ANOVA). **(H)** As in **G** but with primary normal

human epidermal keratinocytes (NHEK) (n = 3, combined data, mean+SD, each dot represents one

biological replicate, *p<0.05 according to one-way ANOVA). **(I, J)** Triplicate relative (to unstimulated)

IL8 mRNA qPCR or IL-8 ELISA from NHEK 3D human skin equivalent constructs stimulated as indicated

for 24 h (n = 3, representative of one biological replicate is shown, mean+SD, each dot represents

one technical replicate, *p<0.05 according to one-way ANOVA).

Fig. 4. naRNA is a driver of NET-associated *in vivo* inflammation

(A) Ear thickness quantified daily in WT C57BL6 mice injected intradermally in vivo on day 0 as

indicated (n = 5 per group, combined data, mean+SD, *p<0.05 according to two-way ANOVA). **(B)**

Fluorescence imaging monitored hourly in *LysM*$^{EGFP/+}$ mice injected intradermally in vivo at t=0 as

indicated (n = 10 per group, combined data, mean+SD, *p<0.05 according to two-way ANOVA). **(C)** As

in **A** but also using *Tlr13*$^{-/-}$ mice (n = 7, combined data, mean+SD, *p<0.05 according to two-way

ANOVA). **(D)** As in **C** but instead of intradermal injection with topical imiquimod application on day 0–

4 (C57BL/6 n = 8, *Tlr13*$^{-/-}$ n = 10, combined data, mean+SD, *p<0.05 according to two-way ANOVA).

(E) Measurement of epidermal thickness of H&E-stained ear skin specimens of **D** (C57BL/6 n = 8,

Tlr13$^{-/-}$ n = 10, combined data, mean+SD, *p<0.05 according to Student's *t*-test). **(F)** Histological

analysis of H&E-stained ear skin specimens of **E** (C57BL/6 n = 8, *Tlr13*$^{-/-}$ n = 10, representative of one

biological replicate is shown, scale bar: 100 µm).

Fig. S1. Controls of IF microscopy, stem cell-derived PMNs, and electron microscopy

(A) Confocal microscopy of unstimulated primary human PMNs (control to Fig. 1A) after 3 h and

stained for naRNA (anti-rRNA Y10b, magenta) and DNA (Hoechst 33342, white, n = 3, representative

images, scale bar: 10 µm). **(B)** Confocal microscopy of unstimulated primary murine BM-PMNs

(control to Fig. 1C) after 16 h and stained as in **A** (n = 3, representative images, scale bar: 10 µm). **(C)**

Brightfield microscopy analysis of cytospun control primary human stem cell-derived PMNs (control

to Fig. 1D) differentiated *in vitro* with/without 100 µM 5-Ethynyluridine (5-EU, n = 3, representative

images, scale bar: 20 µm). **(D)** FACS analysis for cells shown in **C** and Fig. 1D (n=3, representative

data). **(E)** Scanning electron microscopy of PMA-treated human primary PMNs showing only

secondary antibody staining (no primary antibody) control of Fig. 1F (n = 1, representative data; the

image on the right is a composite image with signals from secondary electron and backscattered

electron detectors for topography and additional material information, respectively).

Fig. S2. Antibacterial effect of NETs on live *S. aureus*, controls of IF microscopy and line plot
analysis of naRNA and LL37

(A) Extracellular bactericidal activity of human PMNs/NETs after infection with *S. aureus* and

treatment with RNase A and DNase I during or after formation of PMA-induced NETs (n = 8,

combined data, mean+SD, *p<0.05 according to one-way ANOVA). **(B)** Confocal microscopy of

unstimulated or PMA-stimulated primary human PMNs (control to Fig. 2B) after 3 h, stained DNA

(Hoechst 33342, white, n = 9, representative images, scale bar: 10 µm). **(C)** as in **B** but controls of Fig.

2D (n = 3). **(D)** As in **B** but controls of Fig. 2F (n = 3). **(E)** Confocal microscopy of primary human PMNs

stimulated as indicated for 3 h and stained for naRNA (anti-rRNA Y10b, magenta), LL37 (anti-hLL37-

DyLight550, yellow) and DNA (Hoechst 33342, white, n = 3, representative images). The line plot

analysis of LL37, RNA and DNA staining was performed using ZenBlue3 software. Two to three

different line plots from the same representative image are shown. **(F)** Transmission electron

microscopy of unstimulated human primary PMNs using anti-rRNA and anti-hLL-37 primary and

immunogold (6 nm (black arrow) and 12 nm (white arrow) respectively)-labeled secondary

antibodies (n = 3, representative images, scale bars as indicated).

Fig. S3. Inhibition of PAD4 in human PMNs during NET formation assay and *Unc93b1*[-/-] and *Tlr9*[-/-]
BM-PMN stimulation with human NETs

(A) Confocal microscopy of stimulated primary human PMNs after 3 h and stained for DNA (Hoechst
33342, white) in the presence or absence of the PAD4-inhibitor Cl-amidine (200 µM, n = 3,
representative images; scale bar 10 µm). **(B)** Confocal microscopy of primary C57BL/6 WT or
Unc93b1[-/-] murine BM-PMNs stimulated as indicated for 16 h as in **A** (n = 3 WT, n = 1 *Unc93b1*[-/-],
representative images, scale bar: 10 µm). **(C)** Confocal microscopy of primary C57BL/6 WT or *Tlr9*[-/-]
murine BM-PMNs stimulated as indicated for 16 h as in **A** (n = 3, representative images, scale bar: 10
µm).

Fig. S4. Immune responses of PBMCs, macrophages, and NK-cells to NETs and in vivo fluorescence
imaging

(A) Triplicate IL-8 ELISA from WT, *TLR8*[-/-] and *TLR7*[-/-] THP-1 cells stimulated as indicated for 18 h,
normalized to PMA+ionomycin control (n = 4, combined data, mean+SD, each dot represents one
biological replicate, *$p < 0.05$ according to one-way ANOVA). **(B-D)** Triplicate ELISA for TNF (**A**, n = 4),
IL-6 (**B**, n = 3), and IL-8 (**C**, n = 3) from primary human PBMCs stimulated as indicated with/without
CU-CPT9a for 24 h (combined data, mean+SD, each dot represents one biological replicate, *$p < 0.05$
according to one-way ANOVA). **(E)** Triplicate IFN-γ ELISA from NK-92 MI cells stimulated as indicated
for 24 h (n = 3, combined data, mean+SD, each dot represents one biological replicate, *$p < 0.05$
according to one-way ANOVA).

Fig. S5. Graphical abstract

Supplemental movie S1: 3D reconstruction of NET DNA network decorated with naRNA

Confocal microscopy of primary human PMNs stimulated with PMA for 3 h, stained for naRNA (anti-
rRNA Y10b, magenta) and DNA (Hoechst 33342, white, n = 3) and 3D analysis and animation were
performed using ZenBlue3 software.

Supplemental movie S2: Digest of naRNA in PMA-induced NETs

Confocal live microscopy of RNase digestion of NETs obtained from primary human PMNs stimulated
with PMA for 3 h, stained for naRNA (SYTO RNAselect, magenta) and DNA (Hoechst 33342, white, n =
3). RNase A was added between time point 0 and 5 min (n = 3, scale bar 20 µm). Video analysis and
animation were performed using ImageJ-Win64 software.

<u>References</u>

462   1.   Brinkmann, V. *et al.* Neutrophil extracellular traps kill bacteria. *Science* **303**, 1532-1535
463   (2004).
464

465   2.   Kruger, P. *et al.* Neutrophils: Between host defence, immune modulation, and tissue injury.
466   *PLoS Pathog* **11**, e1004651 (2015).
467

468   3.   Boeltz, S. *et al.* To NET or not to NET:current opinions and state of the science regarding the
469   formation of neutrophil extracellular traps. *Cell Death Differ* **26**, 395-408 (2019).
470

471   4.   Altincicek, B., Stotzel, S., Wygrecka, M., Preissner, K.T. & Vilcinskas, A. Host-derived
472   extracellular nucleic acids enhance innate immune responses, induce coagulation, and
473   prolong survival upon infection in insects. *J Immunol* **181**, 2705-2712 (2008).
474

475   5.   Herster, F. *et al.* Neutrophil extracellular trap-associated RNA and LL37 enable self-amplifying
476   inflammation in psoriasis. *Nat Commun* **11**, 105 (2020).
477

478   6.   Lande, R. *et al.* Plasmacytoid dendritic cells sense self-DNA coupled with antimicrobial
479   peptide. *Nature* **449**, 564-569 (2007).
480

481   7.   Ganguly, D. *et al.* Self-RNA-antimicrobial peptide complexes activate human dendritic cells
482   through TLR7 and TLR8. *J Exp Med* **206**, 1983-1994 (2009).
483

484   8.   Lande, R. *et al.* Cationic antimicrobial peptides in psoriatic skin cooperate to break innate
485   tolerance to self-DNA. *Eur J Immunol* **45**, 203-213 (2015).
486

487   9.   Eigenbrod, T. & Dalpke, A.H. Bacterial RNA: An Underestimated Stimulus for Innate Immune
488   Responses. *J Immunol* **195**, 411-418 (2015).
489

490   10. Lind, N.A., Rael, V.E., Pestal, K., Liu, B. & Barton, G.M. Regulation of the nucleic acid-sensing
491   Toll-like receptors. *Nat Rev Immunol* **22**, 224-235 (2022).
492

493   11. Preissner, K.T., Fischer, S. & Deindl, E. Extracellular RNA as a Versatile DAMP and Alarm Signal
494   That Influences Leukocyte Recruitment in Inflammation and Infection. *Front Cell Dev Biol* **8**,
495   619221 (2020).
496

497   12. Schmidt, B. *et al.* Quantification of free RNA in serum and bronchial lavage: a new diagnostic
498   tool in lung cancer detection? *Lung Cancer* **48**, 145-147 (2005).
499

500   13. Zernecke, A. & Preissner, K.T. Extracellular Ribonucleic Acids (RNA) Enter the Stage in
501   Cardiovascular Disease. *Circ Res* **118**, 469-479 (2016).
502

503   14. Lerner, E.A., Lerner, M.R., Janeway, C.A., Jr. & Steitz, J.A. Monoclonal antibodies to nucleic
504   acid-containing cellular constituents: probes for molecular biology and autoimmune disease.
505   *Proc Natl Acad Sci U S A* **78**, 2737-2741 (1981).
506

507   15. Presolski, S.I., Hong, V.P. & Finn, M.G. Copper-Catalyzed Azide-Alkyne Click Chemistry for
508   Bioconjugation. *Curr Protoc Chem Biol* **3**, 153-162 (2011).
509

510   16. Sioud, M. RNA Interference and CRISPR Technologies Technical Advances and New
511   Therapeutic Opportunities Preface. *Methods Mol Biol* **2115**, V-Vii (2020).
512

513   17.   Sorensen, O., Arnljots, K., Cowland, J.B., Bainton, D.F. & Borregaard, N. The human
514   antibacterial cathelicidin, hCAP-18, is synthesized in myelocytes and metamyelocytes and
515   localized to specific granules in neutrophils. *Blood* **90**, 2796-2803 (1997).
516
517   18.   Colak, E. *et al.* RNA and imidazoquinolines are sensed by distinct TLR7/8 ectodomain sites
518   resulting in functionally disparate signaling events. *J Immunol* **192**, 5963-5973 (2014).
519
520   19.   Zhang, S. *et al.* Small-molecule inhibition of TLR8 through stabilization of its resting state. *Nat*
521   *Chem Biol* **14**, 58-64 (2018).
522
523   20.   Heil, F. *et al.* Species-specific recognition of single-stranded RNA via toll-like receptor 7 and 8.
524   *Science* **303**, 1526-1529 (2004).
525
526   21.   Tabeta, K. *et al.* The Unc93b1 mutation 3d disrupts exogenous antigen presentation and
527   signaling via Toll-like receptors 3, 7 and 9. *Nat Immunol* **7**, 156-164 (2006).
528
529   22.   Li, X.D. & Chen, Z.J. Sequence specific detection of bacterial 23S ribosomal RNA by TLR13.
530   *Elife* **1**, e00102 (2012).
531
532   23.   Ishii, N., Funami, K., Tatematsu, M., Seya, T. & Matsumoto, M. Endosomal localization of
533   TLR8 confers distinctive proteolytic processing on human myeloid cells. *J Immunol* **193**, 5118-
534   5128 (2014).
535
536   24.   Mahdavian Delavary, B., van der Veer, W.M., van Egmond, M., Niessen, F.B. & Beelen, R.H.
537   Macrophages in skin injury and repair. *Immunobiology* **216**, 753-762 (2011).
538
539   25.   Vierbuchen, T., Bang, C., Rosigkeit, H., Schmitz, R.A. & Heine, H. The Human-Associated
540   Archaeon Methanosphaera stadtmanae Is Recognized through Its RNA and Induces TLR8-
541   Dependent NLRP3 Inflammasome Activation. *Front Immunol* **8**, 1535 (2017).
542
543   26.   Coch, C. *et al.* Human TLR8 Senses RNA From Plasmodium falciparum-Infected Red Blood
544   Cells Which Is Uniquely Required for the IFN-gamma Response in NK Cells. *Front Immunol* **10**,
545   371 (2019).
546
547   27.   Dickson, M.A. *et al.* Human keratinocytes that express hTERT and also bypass a p16(INK4a)-
548   enforced mechanism that limits life span become immortal yet retain normal growth and
549   differentiation characteristics. *Mol Cell Biol* **20**, 1436-1447 (2000).
550
551   28.   Bitschar, K. *et al.* Lugdunin amplifies innate immune responses in the skin in synergy with
552   host- and microbiota-derived factors. *Nat Commun* **10**, 2730 (2019).
553
554   29.   Gilliet, M. *et al.* Psoriasis triggered by toll-like receptor 7 agonist imiquimod in the presence
555   of dermal plasmacytoid dendritic cell precursors. *Arch Dermatol* **140**, 1490-1495 (2004).
556
557   30.   Simsekyilmaz, S. *et al.* Role of extracellular RNA in atherosclerotic plaque formation in mice.
558   *Circulation* **129**, 598-606 (2014).
559
560   31.   Warnatsch, A., Ioannou, M., Wang, Q. & Papayannopoulos, V. Inflammation. Neutrophil
561   extracellular traps license macrophages for cytokine production in atherosclerosis. *Science*
562   **349**, 316-320 (2015).
563
564   32.   Neumann, E., Hasseli, R., Lange, U. & Muller-Ladner, U. The Role of Extracellular Nucleic
565   Acids in Rheumatoid Arthritis. *Curr Pharm Biotechnol* **19**, 1182-1188 (2018).

566
567   33.   Song, W., Ye, J., Pan, N., Tan, C. & Herrmann, M. Neutrophil Extracellular Traps Tied to
568   Rheumatoid Arthritis: Points to Ponder. *Front Immunol* **11**, 578129 (2020).
569
570   34.   Sacre, S. *et al.* Oligodeoxynucleotide inhibition of Toll-like receptors 3, 7, 8, and 9 suppresses
571   cytokine production in a human rheumatoid arthritis model. *Eur J Immunol* **46**, 772-781
572   (2016).
573
574   35.   Sacre, S.M. *et al.* Inhibitors of TLR8 reduce TNF production from human rheumatoid synovial
575   membrane cultures. *J Immunol* **181**, 8002-8009 (2008).
576
577   36.   Zuo, Y. *et al.* Neutrophil extracellular traps in COVID-19. *JCI Insight* **5** (2020).
578
579   37.   Yaqinuddin, A. & Kashir, J. Novel therapeutic targets for SARS-CoV-2-induced acute lung
580   injury: Targeting a potential IL-1beta/neutrophil extracellular traps feedback loop. *Med
581   Hypotheses* **143**, 109906 (2020).
582
583   38.   Cicco, S., Cicco, G., Racanelli, V. & Vacca, A. Neutrophil Extracellular Traps (NETs) and
584   Damage-Associated Molecular Patterns (DAMPs): Two Potential Targets for COVID-19
585   Treatment. *Mediators Inflamm* **2020**, 7527953 (2020).
586
587   39.   Barnes, B.J. *et al.* Targeting potential drivers of COVID-19: Neutrophil extracellular traps. *J
588   Exp Med* **217** (2020).
589
590   40.   Golonka, R.M. *et al.* Harnessing innate immunity to eliminate SARS-CoV-2 and ameliorate
591   COVID-19 disease. *Physiol Genomics* **52**, 217-221 (2020).
592
593   41.   Kannemeier, C. *et al.* Extracellular RNA constitutes a natural procoagulant cofactor in blood
594   coagulation. *Proc Natl Acad Sci U S A* **104**, 6388-6393 (2007).
595
596   42.   Sahu, A. *et al.* MicroRNA-145 Impedes Thrombus Formation via Targeting Tissue Factor in
597   Venous Thrombosis. *EBioMedicine* **26**, 175-186 (2017).
598
599   43.   Mollerherm, H., von Kockritz-Blickwede, M. & Branitzki-Heinemann, K. Antimicrobial Activity
600   of Mast Cells: Role and Relevance of Extracellular DNA Traps. *Front Immunol* **7**, 265 (2016).
601
602   44.   Doster, R.S., Rogers, L.M., Gaddy, J.A. & Aronoff, D.M. Macrophage Extracellular Traps: A
603   Scoping Review. *J Innate Immun* **10**, 3-13 (2018).
604
605   45.   Li, M. *et al.* A Novel Peptidylarginine Deiminase 4 (PAD4) Inhibitor BMS-P5 Blocks Formation
606   of Neutrophil Extracellular Traps and Delays Progression of Multiple Myeloma. *Mol Cancer
607   Ther* **19**, 1530-1538 (2020).
608
609   46.   Knight, J.S. *et al.* Peptidylarginine deiminase inhibition reduces vascular damage and
610   modulates innate immune responses in murine models of atherosclerosis. *Circ Res* **114**, 947-
611   956 (2014).
612
613   47.   Fruh, A. *et al.* RNase A Inhibits Formation of Neutrophil Extracellular Traps in Subarachnoid
614   Hemorrhage. *Front Physiol* **12**, 724611 (2021).
615
616   48.   Vlach, J. *et al.* Discovery of M5049: A Novel Selective Toll-Like Receptor 7/8 Inhibitor for
617   Treatment of Autoimmunity. *J Pharmacol Exp Ther* **376**, 397-409 (2021).
618

619 49. Jiang, W. *et al.* A Toll-like receptor 7, 8, and 9 antagonist inhibits Th1 and Th17 responses and
620   inflammasome activation in a model of IL-23-induced psoriasis. *J Invest Dermatol* **133**, 1777-
621   1784 (2013).
622
623 50. Balak, D.M. *et al.* IMO-8400, a toll-like receptor 7, 8, and 9 antagonist, demonstrates clinical
624   activity in a phase 2a, randomized, placebo-controlled trial in patients with moderate-to-
625   severe plaque psoriasis. *Clin Immunol* **174**, 63-72 (2017).
626
627 51. Chang, T.-H. *et al.* Transkingdom mechanism of MAMP generation by chitotriosidase (CHIT1)
628   feeds oligomeric chitin from fungal pathogens and allergens into TLR2-mediated innate
629   immune sensing. *bioRxiv*, 2022.2002.2017.479713 (2022).
630
631 52. Faust, N., Varas, F., Kelly, L.M., Heck, S. & Graf, T. Insertion of enhanced green fluorescent
632   protein into the lysozyme gene creates mice with green fluorescent granulocytes and
633   macrophages. *Blood* **96**, 719-726 (2000).
634
635 53. Bitschar, K. *et al.* Staphylococcus aureus Skin Colonization Is Enhanced by the Interaction of
636   Neutrophil Extracellular Traps with Keratinocytes. *J Invest Dermatol* **140**, 1054-1065 e1054
637   (2020).
638
639 54. Zack, G.W., Rogers, W.E. & Latt, S.A. Automatic-Measurement of Sister Chromatid Exchange
640   Frequency. *Journal of Histochemistry & Cytochemistry* **25**, 741-753 (1977).
641
642

Additional information

644 **Author contributions:** according to Credit guidelines:

	FB	CGL	CY	SC	YW	MN	JF	JS	PE	MD	KH	BS	SB	MWL	YS
Conceptualization	x														
Data curation															
Formal analysis	x	x	x	x		x	x	x	x		x				
Funding acquisition												x			
Investigation	x	x	x	x		x	x	x	x	x	X				
Methodology											X				x
Project administration					x									X	
Resources													x	X	
Software															
Supervision	X											x			x
Validation	X														
Visualization	x	x	x								X				
Writing: original draft	x														
Writing: review and editing	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x

Conflict of interest statement

NKA has received previous grant support from Pfizer and Boehringer Ingelheim and was a paid

consultant for Janssen Pharmaceuticals. All other authors declare no competing interests. S.D.G.

Acknowledgments

We gratefully acknowledge Jim Rheinwald, Holger Heine, Austin Chang, Thomas Zillinger for the

provision of reagents, respectively, and Jon Kagan and Libera Lo Presti for helpful scientific and

editorial comments. We further acknowledge Mark Helm and Martina Christina Schmidt-Dengler for

very helpful technical and scientific support. We thank all voluntary healthy donors of biomaterials

for participating in the study. The study was supported by the Deutsche Forschungsgemeinschaft

(German Research Foundation, DFG) grants CRC TR156 "The skin as an immune sensor and effector

organ – Orchestrating local and systemic immunity" (to FB, CG, JF, JS, BS and ANRW), NIH grants

R01AI146177, R01AR073665, and R01AR069502 (to NKA). NKA. has received previous grant support

from Pfizer and Boehringer Ingelheim and was a paid consultant for Janssen Pharmaceuticals.

Infrastructural funding was provided by the University of Tübingen, the University Hospital Tübingen

and the DFG Clusters of Excellence "iFIT – Image-Guided and Functionally Instructed Tumor

Therapies" (EXC 2180, to AW, PE, BS and MWL), "CMFI – Controlling Microbes to Fight Infection (EXC

2124 to AW and BS). Gefördert durch die Deutsche Forschungsgemeinschaft (DFG) im Rahmen der

Exzellenzstrategie des Bundes und der Länder - EXC 2180 and EXC 2124. The authors declare no

competing financial interests.

Abbreviations

AF – Alexa Fluor; bRNA – bacterial ribonucleic acid; DAMP – damage-associated molecular pattern; 5-

EU – 5-Ethynyluridine; FACS – Fluorescence Activated Cell Sorting; fRNA – fungal ribonucleic acid;

H&E - Hematoxylin and Eosin; HEK – Human embryonic kidney; HMGB-1 – High-Mobility-Group-

Protein B1; HSC – hematopoietic stem cell; IFN – Interferon; IL – Interleukin; KO – knockout; MCET –

Mast cell extracellular trap; MET – Macrophage extracellular trap; MPO – Myeloperoxidase; naRNA –

Neutrophil extracellular trap-associated RNA; NET – Neutrophil extracellular trap; NK – natural killer;

PAD4 - Peptidyl arginine deiminase 4; PBMC - Peripheral Blood Mononuclear Cell; PKC- Protein kinase

C; PMN – Polymorphonuclear leukocytes; RT – room temperature; SEM – scanning electron

microscopy;   TLR   –   Toll-like   receptor;   TNF   –   Tumor   necrosis   factor

Materials and Methods

Reagents

PMA (tlrl-pma), Nigericin (tlrl-nig), LL37 (tlrl-l37), as well as the PRR ligands LPS (tlrl-peklps), R848 (tlrl-r848), TL8 (tlrl-tl8506), and the TLR8-inhibitor CU-CPT9a (inh-cc9a) were from Invivogen, Ionomycin was acquired from Sigma (I0634-1MG). RNase inhibitor (N2615) was from Promega, RNase A (EN0531), DNase I (EN0521) and DNase inhibitor (EN0521) were from Thermo Fisher. The PAD4-inhibitor Cl-amidine (506282) was from Merck Millipore. DOTAP (L787.2) was from Roth (see Supplementary Table S1), ssRNA40 was from Eurogentec and CpG PTO 2006 from TIB Molbiol (see Supplementary Table S2). Bacterial RNA isolated from *S. aureus* was prepared as described [5]. Fungal RNA from *C. albicans* strain SC5314 was isolated as described below, as well as naRNA isolated from PMA NETs. For complex formation to stimulate cells in a volume of 500 µL, 5.8 µM ssRNA40 (~ 34.4 µg/mL), fungal RNA (125 ng/mL), bacterial RNA (10 µg/mL) or PMA NET derived naRNA (~ 600 ng/mL) was mixed with 10 µg LL37 and left for 1 h at room temperature (RT). For a smaller volume of cells, complexes were used in the according fractional amount. For RNA-only or LL37-only controls, the same amounts and volumes were used replacing one of the components with sterile, endotoxin-free H_2O. For complex formation with DOTAP, the according RNA or CpG was incubated with the transfection reagent for 10 min at RT prior to stimulation of the cells. NET content for stimulation was prepared as described below. Antibodies used for fluorescence microscopy, as well as click chemistry reagents are listed in Supplementary Tables S3 and S4. Constructs used for transfection of HEK293T cells are listed in Supplementary Table S5.

Preparation of fungal RNA from *C. albicans*

C. albicans SC5314 (kindly cultured and prepared by Tzu-Hsuan Chang, Tübingen) was plated in a slant tube containing YPD agar and grown overnight at 30 °C as described in [51]. One colony was picked from the slant tube and resuspended in 500 µL YPD medium, centrifuged at 10,000 rpm for 1 min and washed with phosphate-buffered saline (PBS). Afterwards, the pellet was resuspended in 200 µL RLT buffer (derived from RNeasy Mini Kit, Qiagen, #74106) and transferred into a 2 mL tube containing 0.5 mm diameter ceramic beads. The tube was filled up to 1 mL with RLT buffer and the fungi were subsequently homogenized by using a microtube homogenizer (BeadBug™, Merck) with an interval of seven times 2 min shaking at 2800 rpm and 1 min cooling break on ice. The supernatant was transferred into a new tube containing 1 mL 75% ethanol (VWR, 20821.330). The further RNA isolation was performed according to the manufacturer's instructions using the Qiagen RNeasy Mini Kit for purification of Total RNA from Animal Tissues (RNeasy Mini Kit, Qiagen, 74106). The RNA was eluted in 30 µL RNase DNase-free H_2O and the concentration was determined with a Nanodrop Spectrophotometer.

Mice

Unc93b1[3d/3d] [21], *Tlr13*[-/-] [22] (both kindly provided by Tatjana Eigenbrod, Heidelberg and on C57BL/6 background) and WT C57BL/6 mice between 8 and 20 weeks of age were used in accordance with local institutional guidelines on animal experiments, regular hygiene monitoring, and specific locally approved protocols compliant with the German regulations of the Gesellschaft für Versuchstierkunde/Society for Laboratory Animal Science (GV-SOLAS) and the European Health Law of the Federation of Laboratory Animal Science Associations (FELASA) for sacrificing and *in vivo* work. *Unc93b1*[3d/3d], *Tlr13*[-/-] and WT C57BL/6 control mice were housed in controlled specific-pathogen-free animal facilities at the Interfaculty Institute of Cell Biology, Tübingen. Local federal authority for the approval of experimental protocols was the Regierungspräsidium Tübingen. *Tlr9*[-/-] mice and matched WT C57BL/6 control mice were a kind gift from Birgit Schittek, Tübingen. LysM[EGFP/+] [52], *Tlr13*[-/-] mice [22] (kindly provided by James Chen, Houston and David Nemazee, La Jolla) and WT (all on a C57BL/6 background) mouse strains were bred and maintained under the specific pathogen-free conditions, with air isolated cages at an American Association for the Accreditation of Laboratory Animal Care

(AAALAC)-accredited animal facility at Johns Hopkins University and handled according to procedures
described in the Guide for the Care and Use of Laboratory Animals as well as Johns Hopkins
University's policies and procedures as set forth in the Johns Hopkins University Animal Care and Use
Training Manual, and all animal experiments were approved by the Johns Hopkins University Animal
Care and Use Committee (MO21M378). Gender-and age-matched 6–8-week-old mice were used for
each experiment.

Isolation and stimulation of primary bone-marrow-derived polymorphonuclear neutrophils (BM-
PMNs)

Bone-marrow (BM)-PMNs were isolated from the bone marrow as described [5]. In brief, bones were
isolated from the respective mice and the bone marrow was flushed out. Afterwards, neutrophils
were isolated using magnetic separation (mouse Neutrophil isolation kit, Miltenyi Biotec, 130-097-
658) following the manufacturer's instructions. In total, 1×10^5 cells/well PMNs were seeded in a
24-well plate, and stimulation was carried out with PMA (600 nM), ssRNA+LL37 complex (as
previously described), nigericin (50 µM), live *C. albicans* (MOI1) or human NET content (mock control
and PMA NETs, 1:50 dilution) for 16 h at 37°C and 5% CO2. Subsequently, cells were stained for
immunofluorescence.

Study participants and human blood or tissue sample acquisition

All healthy donors included in this study provided their written informed consent before
participation. Approval for use of biomaterials was obtained for this project by the local ethics
committee of the Medical Faculty Tübingen in accordance with the principles laid down in the
Declaration of Helsinki as well as applicable laws and regulations.

Primary human neutrophil isolation and stimulation

Neutrophils of healthy human donors were isolated as described [5]. In brief, EDTA-anticoagulated
whole blood was diluted in PBS (Thermo Fisher, 14190-169), loaded on Ficoll (1.077 g/mL, Sigma,
10771) and centrifuged for 25 min at 509 x g at RT without brake. Afterwards, all layers except the
erythrocyte-granulocyte layer were discarded and erythrocyte lysis was performed twice (for 20 and
for 10 min) using 1x ammonium chloride erythrocyte lysis buffer (see Supplementary Table S6) at 4
°C on roller shaker. The remaining cells were resuspended in culture medium (RPMI culture medium
(Sigma-Aldrich, R8758) + 10% FBS (heat-inactivated, sterile filtered, Th. Geyer, 11682258)) to a
concentration of 1.6×10^6 cells/mL. 500 µL of cells were seeded in a 24-well plate for
immunofluorescence microscopy or 8 mL of 5×10^6 cells/mL in 10 cm uncoated dishes for NET
preparation and naRNA isolation/isolation of whole PMN RNA. After seeding, the cells were rested
for 30 min at 37°C and 5% CO$_2$, followed by 3 h stimulation with PMA (600 nM), nigericin (50 µM),
live *C. albicans* (MOI2), RNA+LL37 complexes (as previously described) or NET content at indicated
dilutions for IF or 4 h stimulation with PMA (600 nM) for NET preparation. Where indicated, the cells
were incubated with 100 nM TLR8-inhibitor CU-CPT9a or 200 µM PAD4-inhibitor Cl-amidine 30 min
before stimulation and medium was not replaced during incubation with the respective stimuli.

Primary peripheral blood mononuclear cell (PBMC) isolation and stimulation

PBMCs were isolated from whole blood or buffy coats as described [5]. In brief, EDTA-anticoagulated
blood was diluted in PBS and density gradient separation was performed as described above. The
PBMC layer was then carefully transferred into another reaction tube and diluted in PBS (1:1). The
cell suspension was spun down at 645 × g for 8 min and the cells were washed two more times in
PBS and resuspended in culture medium (RPMI + 10% FBS (heat inactivated) +1% L-glutamine) at a
density of 1×10^6 cells/mL. Afterwards, 200 µL of PBMCs were seeded in a 96-well plate and
stimulated with LPS (100 ng/mL), R848 (5 µg/mL), TL8 (100 ng/mL), ssRNA (1.6 µg/mL) + DOTAP (50
µg/mL), ssRNA+LL37 complex (as described above), and respective NET content (1:20 dilution) for 24
h at 37°C and 5% CO$_2$. Where indicated, the TLR8 inhibitor CU-CPT9a was added to the cells at a
concentration of 1 µM 2 h before stimulation and was not removed for the incubation of the cells

with the respective stimuli. After the stimulation, the plate was centrifuged for 5 min at 1500 rpm
and the supernatant was collected and stored at -80 °C until the ELISA was performed.

Preparation of NETs and isolation of naRNA/whole PMN RNA

NETs were prepared by 600 nM PMA treatment for 4 h at 37°C and 5% CO_2, or cells were left
untreated during the incubation as the mock control. After incubation, the neutrophils were gently
washed three times with PBS to get rid of PMA as the stimulus for NET formation, any cytokines
released by the cells and unstimulated PMNs, as those do not adhere to the uncoated petri dish used
here. In some conditions (as indicated) during the NET preparation process and for storage, naRNA
was protected by addition of 10 U/µL RNase inhibitor (= mock or PMA NETs + RNase Inhibitor). For
digest of NETs, the NET content was incubated for 20 min at 37 °C with RNase A (Thermo Fisher,
EN0531; 100 µg/mL, EDTA for DNase inhibition added) at 37 °C. For isolation of naRNA, RNase
inhibitor was added to the PMNs during NET formation. After the above-described washing process,
PMA or mock NETs were resuspended in 300 µL of ML buffer and RNA isolation was performed
according to the manufacturer's instructions (NucleoSpin miRNA isolation kit, Macherey-Nagel,
740971.50). The RNA was eluted in 50 µL RNase DNase-free H_2O and the concentration was
determined with a Nanodrop Spectrophotometer. For isolation of whole PMN RNA, untreated PMNs
were directly lysed, and the RNA was isolated from the cells according to the manufacturer's
instructions.

Preparation of human primary stem cell-derived PMNs

Stem cells derived from human healthy donors were prepared and differentiated as described [16]. In
brief, bone marrow was diluted in PBS, carefully layered on Ficoll-Paque medium (density: 1.077
g/mL) and centrifuged at 500 x g for 25 min at RT without brakes. The interphase layer containing the
mononuclear cell fraction was transferred to a new tube and washed twice with 30 mL ice-cold PBS
by centrifugation at 300 x g for 8 min at 8 °C. Further, the cells were resuspended, counted, and
isolated using Human CD34 MicroBead Kit (Miltenyi) for magnetic beads-based isolation of $CD34^+$
cells from BM-MNCs. Afterwards, the number of $CD34^+$ HPSCs was determined and cultured in $CD34^+$
culture medium (Stemline II Hematopoietic Stem Cell Expansion medium supplemented with 10%
FCS, 1% L-glutamine, 1% penicillin/streptomycin, and a human recombinant cytokine cocktail
consisting of 20 ng/mL IL-3, 20 ng/mL IL-6, 20 ng/mL TPO, 50 ng/mL SCF, and 50 ng/mL FLT-3L) at a
density of 2 x 10^5/mL at 37 °C and 5% CO_2. The medium was replaced every second day and the cells
were cultured for 14 days. During the differentiation process, the cells were treated with 100 µM 5-
ethynyluridine for 14 days for subsequent click chemistry labeling of endogenous RNA or were left
untreated as negative controls. To verify differentiation, cell morphology was assessed using
Cytospin assay. In brief, the Cytoclip™ slide clips were loaded by fitting the filter card, the sample
chamber, and the glass slide. An assemble Cytoclip™ slide clip was then placed in the slide clip
support plate of the cytospin centrifuge. 2 x 10^4 cells from liquid culture differentiation were
pipetted into Cytofunnel™ and centrifuged for 3 min at 200 x g. The cytospin slides were stained for 5
min in May-Grünwald stain, rinsed shortly with ddH_2O, and then stained for 10 min in Wright-Giemsa
stain. Afterwards, the slides were rinsed shortly with ddH_2O, and cell morphology was determined
using a microscope. To further verify differentiation, flow cytometric analysis was performed using
antibodies specific for the following hematopoietic/myeloid markers: CD45 (leukocyte marker), CD34
(HSPC marker), CD33 (promyelocyte marker), CD11b (myeloid cell marker), CD14 (monocyte marker),
and CD15 and CD16 (neutrophil markers). Neutrophil percentage was determined by gating on
neutrophils as follows: $CD45^+CD11b^+CD15^+$, or $CD45^+CD11b^+CD16^+$, or $CD45^+CD15^+CD16^+$ cells

BlaER1 cell culture, transdifferentiation and stimulation

BlaER1 cells (WT, *Unc93b*$^{-/-}$ and *Tlr8*$^{-/-}$), a kind gift of Holger Heine, Borstel, Germany [25], were
cultured, transdifferentiated, and stimulated for 18 h with the respective stimuli as described [5]. In
brief, 1 x 10^6 cells/well were seeded in a 6-well plate and differentiated with 10 ng/mL hIL-3, 10
ng/mL hMCSF and 150 nM β-estradiol in complete RPMI-1640 (PANBiotech, P04-18525) for 7 days.
Afterwards 5 x 10^4 differentiated cells were reseeded in a 96-well plate, followed by 1 h resting. Cells

were treated with the respective stimuli (LPS at 0.1 µg/mL, R848 at 5 µg/mL, TL8 at 100 ng/mL or
ssRNA+LL37 complex as described above) and mock or PMA NETs with or without RNase inhibitor) in
complete medium in a total volume of 125 µL/well for 18 h. After stimulation, the cells were
centrifuged for 5 min at 1200 rpm, the supernatant was transferred into a new plate and stored at -
80°C until the ELISA was performed.

THP-1 cell culture, differentiation and stimulation

THP-1 cells (THP-1 cells were a kind gift from Thomas Zillinger, Bonn, Germany [26]), were cultured in
complete RPMI-1640 (Sigma, R8758-24X500ML) medium. For differentiation into macrophage-like
cells, 5 x 10^4 cells/well were seeded in a 96-well plate, treated with 300 ng/mL PMA and incubated
for 16 h at 37°C and 5% CO_2. The next day, the cells were washed three times with PBS and fresh
medium was added, followed by 48 h of resting. Subsequently, the medium was removed, exchanged
by medium containing 200 U/mL IFN-γ (Sigma, I-3265), and the cells were incubated for 6 h. After
repeated washing and medium exchange, cells were treated with the respective stimuli (PMA (25
µg/mL) + Ionomycin (0.375 µg/mL), LPS (0.1 µg/mL), R848 (5 µg/mL), TL8 (40 ng/mL), ssRNA+LL37
complex (as described above), mock NETs + RNase inhibitor (1:50 dilution), PMA NETs (1:50 dilution)
and PMA NETs + RNase inhibitor (1:50 dilution)) in complete medium in a total volume of 125 µL/well
for 18 h. After stimulation, the cells were centrifuged for 5 min at 1200 rpm, the supernatant was
transferred into a new plate and stored at -80°C until the ELISA was performed.

N/TERT-1 keratinocyte cell culture and stimulation

N/TERT-1 cells (a kind gift from Prof. James Rheinwald [27]) were cultured for less than ten passages in
complete CnT-07 medium (CELLnTEC, CnT-07). Two days prior to stimulation, a total amount of 2 x
10^4 cells/well was seeded in a 96-well plate and incubated at 37°C and 5% CO_2. The medium was
renewed, and the cells were stimulated in a total volume of 125 µL/well of the respective stimuli
diluted in medium (PMA (25 µg/mL) + Ionomycin (0.375 µg/mL), TL8 (200 ng/mL), ssRNA+LL37
complex (as previously described), as well as mock and PMA NETs with and without RNase inhibitor
at indicated dilutions). After 24 h stimulation, the cells were centrifuged at 1200 rpm for 5 min and
the supernatant was stored in a new plate at -80°C until further usage.

Primary normal human epidermal keratinocyte (NHEK) cell culture and stimulation

NHEK cells (Normal Human Epidermal Keratinocytes (NHEK) single juvenile donor, proliferating,
PromoCell, C-12002) were grown and stimulated in Keratinocyte Growth Medium 2 (PromoCell, C-
20111). Two days prior to stimulation, a total amount of 2 x 10^4 cells/well was seeded in a 96-well
plate and incubated at 37°C and 5% CO_2. For stimulation, the medium was renewed with basal
medium (PromoCell, C-20211) containing 1.7 mM $CaCl_2$ (Roth, CN93.1) and stimuli were added and
diluted in medium in a final volume of 125 µL (R848 (20 µg/mL), TL8 (200 ng/mL), ssRNA+LL37
complex, as well as mock NETs + RNase inhibitor (1:25 dilution) and PMA NETs with and without
RNase inhibitor (1:25 dilution)). After 24 h stimulation, the cells were centrifuged at 1200 rpm for 5
min and the supernatant was stored in a new plate at -80°C until further usage.

Preparation of 3D human skin equivalent

The 3D human skin equivalent was prepared as described [53]. Briefly, primary fibroblasts were seeded
on collagen and incubated in FF medium for five days. Subsequently, primary keratinocytes were
added to the wells and airlifting was performed on day 12. On day 22, the 3D skin model was
stimulated with NET content (25 µL/well for one 3D construct grown in a 12-well chamber) or
respective water control for 24 h. Afterwards, supernatant was harvested for ELISA, RNA was isolated
for qPCR analysis and H&E staining was performed.

NK-92 MI cell culture and stimulation

NK-92 MI cells (kindly provided by Melanie Märklin, University Hospital Tübingen) were cultured in
IMDM-Medium (Lonza, 12-722F). For stimulation, a total amount of 1 x 10^5 cells/well were seeded in
a volume of 200 µL and rested for 2 h at 37°C and 5% CO_2. The cells were stimulated with LPS (100

ng/mL), CpG (2.5 µM) + DOTAP (25 µg/mL), R848 (5 µg/mL), TL8 (100 ng/mL), ssRNA (1.6 µg/mL) +
DOTAP (50 µg/mL), ssRNA+LL37 complex (as described above), or NET content (1:100 dilution) for 24
h and afterwards centrifuged for 5 min at 1500 rpm. Subsequently, the supernatant was transferred
into a new plate and stored at -80°C until further usage.

Flow cytometry

After PMN isolation, the purity and activation status of the cells was determined by flow cytometry
as described [5]. In brief, 200 µL of cells were transferred into a 96-well plate (U-bottom) and
centrifuged for 5 min at 448 x g. Afterwards, blocking was performed using pooled human serum
diluted 1:10 in PBS for 15 min at 4°C. After washing, the samples were stained for 20 min at RT in
the dark and fixed (4% PFA in PBS) after repeated washing for 10 min at RT in the dark. After an
additional washing step, the cell pellets were resuspended in 300 µL PBS and measurements were
performed on a FACS Canto II (BD Bioscience, Diva software). Analysis was performed using FlowJo
V10 analysis software.

RNA sequencing analysis of naRNA

Isolated naRNA was analyzed for quality control using the Agilent 4200 TapeStation system.
Subsequently, the RNA was sequenced according to NEBNext® Ultra™ II Directional RNA Library Prep
Kit for Illumina® using the protocol for use with rRNA Depleted FFPE RNA. The data was quantified
using Salmon Version1.5.0 and tximport was used to obtain the transcript-level quantification. For
transcript classification, GENECODE annotation was performed.

Fluorescence microscopy of fixed human or murine primary neutrophils

500 µL of 1.6 x 10^6 cells/mL of human blood PMNs, and 2 x 10^6 cells/mL murine BM-PMNs were
seeded in a 24-well plate containing poly-L-lysine-coated glass coverslips (Electron Microscopy
Sciences, 72292-04) and rested for 30 min before stimulation at 37°C and 5% CO_2 with the according
stimuli (as described above) for 3 h (human) or 16 h (murine), respectively, as adapted from
Brinkmann *et al.*, 2004 [1]. After stimulation, the cells were carefully washed with PBS and fixed with
fixation buffer (Biolegend, 420801) for 10 min at RT in the dark. Afterwards, the cells were blocked
with PBS containing 0.1% heat-inactivated diethylpyrocarbonate (DEPC) (Roth, K028.2), 10% chicken
serum (Normal Chicken Serum Blocking Solution S-3000, Biozol/Vectorlabs, VEC-S-3000-20), 0.1%
saponin (Applichem, A4518.0100), as well as 10 U/µL RNase inhibitor for 2 h at RT. The primary
antibodies (rRNA Y10b, hLL37, see Supplementary Table S3) were diluted 1:50 in blocking buffer and
subsequently incubated for 2 h at RT. Afterwards, the cells were washed three times with PBS
containing 0.1% heat inactivated DEPC and incubated with the secondary antibodies (see
Supplementary Table S3) in a 1:500 dilution in blocking buffer for 1 h. After repeated washing, the
cells were incubated with Hoechst 33342 (Thermo Fisher; 1 µg/mL) for 5 min to stain nuclear DNA.
Secondary antibodies alone did not yield any significant staining under identical staining and
acquisition conditions. The coverslips were mounted (ProLong™ Diamond Antifade Mountant,
Thermo Fisher, P36961) on glass slides and left to dry overnight at RT in the dark. Subsequently, the
samples were stored at 4°C before microscopy using a Zeiss LSM800 Confocal microscope (40x or 63x
magnification with z-stack acquisition, AiryScan mode) and image analysis using ImageJ-Win64 and
Zen Blue3 software was performed. For IF of ultrathin sections, coverslips were coated with a carbon
layer, glow-discharged and UV-treated overnight before cells were seeded. After incubation, the cells
were fixed in 2.5% glutaraldehyde in PBS for 2 h at room temperature followed by incubation at 4°C.
Samples were prepared by the progressive lowering of temperature method (PLT). Therefore, the
fixed cells were incubated in ethanol for infiltration of Lowicryl HM20, followed by UV polymerization
and subsequent sectioning into 50-60 nm ultrathin slices. Immunolabeling was performed by
incubation with anti-rRNA Y10b and anti-LL37 primary antibodies for 1 h at RT. Afterwards, the cells
were incubated with goat anti-mouse Cy3 and goat anti-rabbit AF488 in blocking buffer for 1 h at RT.
Following immunolabeling, the samples were treated with 1% uranyl acetate for 5 min at RT before
analysis using a Zeiss Axioplan microscope (63x objective, Olympus SIS cell software) and image/line
plot analysis was performed using ImageJ-Win64.

Quantification of NET formation

To quantify the formation of NETs, microscopy images were obtained using a Zeiss LSM800 Confocal microscope with a 40x objective and 3x3 tiles acquisition. Three images per sample of three biological replicates were taken. To quantify NET formation by using NET-related signal dispersion of rRNA and DNA signal, ImageJ-Win64 software was used, and a threshold (Triangle threshold) was applied as originally described [54]. Particles were analyzed with a ROI (region of interest) manager (size (micron^2): 100-infinity (pixel units); circularity 0.00-1.00) and average size and number of particles (ROIs) were assessed. In NETs, RNA and DNA signals showed up in a greater number and smaller size, making the usage of the ratio suitable as a measurement of NET-related signal dispersion. In the case of using the DNA signal only to assess NET formation, ImageJ-Win64 software was used to create a PNG image and a grid with 8x8 tiles was manually applied to the images. Tiles containing extracellular DNA structures were manually counted in a blinded manner as NET-positive tiles.

Live fluorescence microscopy of enzymatic digest of human NETs

500 µL of 1.6 x 10^6 cells/mL of human blood PMNs were seeded in a 4-well glass bottom microscopy cell culture dish (Greiner, 627871) and rested for 30 min before stimulation at 37°C and 5% CO_2 with PMA (600 nM) for 3 h. After stimulation, the medium was carefully removed and the cells were washed with PBS before adding fresh culture medium (RPMI culture medium without phenol red (Sigma-Aldrich, R7509) + 10% FBS (heat inactivated, sterile filtered, TH Geyer, 11682258)). Hoechst 33342 (Thermo Fisher, 1 µg/mL) to stain nuclear DNA and SYTO RNAselect Green fluorescent dye (Thermo Fisher, 50 µM) to stain naRNA was added to the cells, as well as RNase A (Thermo Fisher, EN0531; 100 µg/mL) was added between time point 0 and 5 min. Live cell imaging was performed using a Zeiss LSM800 Confocal microscope with a 63x objective and z-stack acquisition, taking an image every 5 min for 30-60 min respectively. Image analysis and video creation was performed using ImageJ-Win64.

Click chemistry of primary, stem cell-derived PMNs and fluorescence microscopy

500 µL of 1.6 x 10^6 cells/mL of human stem cell-derived PMNs treated with 5-ethynyluridine or left untreated were seeded in a 24-well plate containing poly-L-lysine-coated glass coverslips and rested for 30 min before stimulation with PMA (600 nM) at 37°C and 5% CO_2 for 12 h. After stimulation, the cells were washed and permeabilized with ice-cold acetone (Applichem, A1582.2500PE) 1:1 methanol (Honeywell, 32213-2.5L) for 5 min at RT. Subsequently, the click chemistry (reagents see Supplementary Table S4) labeling of endogenous RNA was performed as described [15]. Briefly, in a total volume of 500 µL, 2 µL of AF546-Azide, a pre-mixture of 1 mM $CuSO_4$ and 1.25 mM THPTA, 5 mM aminoguanidine-hydrochloride and 5 mM Na-ascorbate in PBS were added to the cells in a 24-well plate. The wells were sealed with plastic foil and incubated for 1 h at RT while shaking in the dark. For negative controls, 5-ethynyluridine untreated cells incubated with complete click chemistry reagents and 5-ethynyluridine treated cells incubated with PBS and AF546-Azide only were used. No significant signals were observed. After the incubation, the cells were washed three times for 5 min with PBS and counterstained with rRNA Y10b-AF647 (see Supplementary Table S3) at 1:50 in PBS for 2 h at RT in the dark. After repeated washing, the cells were incubated with Hoechst 33342 to stain nuclear DNA and mounted as described above. Imaging and analysis were performed as previously described.

Electron microscopy

For electron microscopy, 500 µL of 1.6 x 10^6 cells/mL of human blood-derived PMNs were seeded in a 24-well plate containing coverslips which were pre-coated with 0.01 % poly-L-lysine (Sigma, A-005-C) for 15 min at 37°C for scanning electron microscopy (SEM). For transmission electron microscopy (TEM), coverslips were coated with a carbon layer, glow-discharged and UV-treated overnight before cells were seeded. Cells were rested for 30 min at 37°C and 5% CO_2 and subsequently stimulated with 1200 nM PMA or left untreated for 3 h. Afterwards, the cells were fixed in 2.5% glutaraldehyde in

PBS for 2 h at room temperature followed by incubation at 4°C. For SEM, samples were post-fixed
with 1% osmium tetroxide for 1 h on ice. Subsequently, samples were dehydrated in a graded
ethanol series followed by critical point drying (CPD300, Leica Microsystems) with CO_2. Finally, the
cells were sputter-coated with a 4 nm thick layer of platinum (CCU-010, Safematic) and examined
with a field emission scanning electron microscope (Regulus 8230, Hitachi High Technologies) at an
accelerating voltage of 3 kV. For antibody labeling, cells treated as described above were fixed in 4%
formaldehyde in PBS for 1-2 hours at room temperature and 4°C overnight. After washing and
blocking (0.2% gelatin in PBS) samples were incubated for 1 h at room temperature with rRNA Y10b
as the primary antibody in blocking buffer and for 1 h at RT with goat anti-mouse antibodies coupled
to 6 nm gold in blocking buffer (Jackson ImmunoResearch, code number 115-195-166). Samples
labeled with 6 nm gold were further silver enhanced. Following immunolabeling, the samples were
treated with 1% uranyl acetate for 5 min at RT, dehydrated and critical point dried as before.
Samples were sputter-coated with a 5 nm thick layer of carbon (CCU-010, Safematic) and analyzed in
the SEM with an accelerating voltage of 5 kV. For TEM, samples were prepared by the PLT method as
described above for IF of ultrathin sections. Samples were placed on grids (Cu, Ni) and subsequently
sectioned into 50-60 nm ultrathin slices. Immunolabeling was performed by incubation with anti-
rRNA Y10b as the first primary antibody in blocking buffer at 4°C overnight, followed by incubation
with anti-LL37 as the second primary antibody for 1 h at RT. Afterwards, the cells were incubated
with goat anti-mouse coupled to 6 nm gold and goat anti-rabbit 12 nm gold secondary antibodies in
blocking buffer for 1 h at RT. Following immunolabeling, the samples were treated with 1% uranyl
acetate for 5 min at RT before analysis in the TEM.

Transient transfection of HEK293T cells

HEK293T cells were transiently transfected with the respective TLR8, TLR7, TLR9 and NF-κB reporter
plasmids as described [18] (see Supplementary Table S5 for plasmids) using X-tremeGENE™ HP DNA
Transfection Reagent (Merck, 6366236001). A total amount of 5×10^4 cells/well were seeded in a 24-
well plate one day prior to transfection. For the transfection of one well, 100 ng of the according TLR
plasmid, 100 ng of the firefly luciferase NF-κB reporter and 10 ng *Renilla* luciferase control reporter
was mixed in Opti-MEM™ Reduced Serum Medium (Thermo Fisher, 31985062) in a total volume of
50 µL. After 15 min incubation at RT, the transfection mix was added to the cells and the cells were
incubated for 48 h. Prior to subsequent stimulation, the medium was changed to complete DMEM
medium (Sigma, D5796-24X500ML), and the cells were incubated with the respective NET content
stimuli and controls (R848 (2.5 µg/mL), TL8 (100 ng/mL), CpG (1.25 µM) + DOTAP (25 µg/mL), ssRNA
(0.6 µg/mL) + DOTAP (20 µg/mL), mock NETs + RNase inhibitor (1:50 dilution), PMA NETs (1:50
dilution), PMA NETs + RNase inhibitor (1:50)) for 18 h at 37°C and 5% CO_2. Supernatants were then
removed, and the cells frozen briefly at -80 °C. Subsequently they were used for dual luciferase assay.

Dual luciferase reporter assay

The dual luciferase reporter assay for detection NF-κB activation after TLR transfection and
subsequent stimulation was performed as described [5]. In brief, supernatants were removed from the
cells after stimulation and 60 µL/well of 1x passive lysis buffer (E194A, Promega) was added. The
plate was then incubated for 15 min at RT on the plate shaker and subsequently stored at −80°C for
at least 15 min to facilitate complete cell lysis. After thawing, the cell solution (60 µL) was
transferred into a V-bottom 96-well plate and centrifuged for 10 min at 2500 rpm and 4°C to pellet
cell debris. Ten microliters of supernatant were then transferred into a white microplate and each
condition was measured in triplicates using the FLUOstar OPTIMA device (BMG Labtech). Firefly and
Renilla luciferase activity were determined using the Promega Dual luciferase kit. Both enzyme
activities were measured for 12.5 s with 24 intervals of 0.5 s, respectively. The data was analyzed by
calculating the ratio of the two measured signals, thereby normalizing each firefly luciferase signal to
its corresponding Renilla luciferase signal. The ratios were represented as the relative light units
(RLU) of NF-κB activation.

Extracellular bacterial killing of S. aureus

The killing assay of *S. aureus* with human PMNs was performed according to Brinkmann *et al.*, 2004 [1]. In brief, PMNs were seeded at a density of 2×10^6 cells/mL and incubated with PMA (600 nM) for 2 h at 37°C. Afterwards, the medium was carefully replaced with serum-free culture medium, containing 2% heat-inactivated pooled human serum with cytochalasin D (10 µg/mL) and incubated further for 15 min before infection with bacteria. Cytochalasin D treatment did not affect NETs and this concentration was effective in blocking bacterial phagocytosis. To investigate whether naRNA of NETs was important in extracellular killing, samples were either treated with RNase A (Thermo Fisher, EN0531; 100 µg/mL) or DNase I (Thermo Fisher, EN0521; 1 U/10 µL) for 2 h during NET formation, or after NET formation during the 30 min bacterial killing process. Samples were centrifuged at 700 x g for 10 min and incubated at 37°C and 5% CO_2 for 30 minutes. Bacterial killing was measured as percentages of control values (bacteria incubated alone in media without neutrophils).

In vivo analysis of naRNA DAMP effects

To investigate the effect of NETs with or without RNase inhibitors and the respective TLR signaling, 20 µL of NET content and the respective controls were injected into the ears of C57BL/6 and *Tlr13*[-/-] mice intradermally on day 0. Afterwards, as a measure of inflammation, the ear thickness was assessed using a manual caliper (0.01–10 mm, Peacock, Tokyo, Japan) until day 4.

Neutrophil infiltration *in vivo* fluorescence imaging

The *in vivo* experiment for investigating neutrophil infiltration was performed as described [5]. Briefly, LysM[EGFP/+] mice were injected intradermally with 20 µL of PMA or mock NET content or respective controls. LysM[EGFP/+] mice were then anesthetized with inhalation isoflurane and *in vivo* fluorescence imaging was performed using the IVIS Lumina II imaging system (Caliper). EGFP fluorescence was measured using excitation (465 nm), emission (515–575 nm), and exposure time (0.5 s). Data are quantified as total radiant efficiency ($[photons/s]/[\mu W/cm^2]$) within a circular region of interest using Living Image software (Caliper).

Imiquimod model of psoriatic skin inflammation

To analyze the effect of RNA signaling in an *in vivo* model for psoriasis, the well-established imiquimod mouse model was used [29]. C57BL/6 and *Tlr13*[-/-] mice were used and, briefly, 70 µL (62.5 mg) of imiquimod (5%, Taro Pharmaceuticals Industries, Hawthorne, NY) was applied daily to both sides of a mouse ear for 5 consecutive days (day 0 to 4). Ear thickness was measured with a manual caliper (0.01–10 mm, Peacock, Tokyo, Japan) before imiquimod application. A day after the last application of imiquimod (day 5), full-thickness ear skin was excised with surgical scissors for histologic analysis.

Histologic analysis of mouse ear skin

Ear skin specimens were collected by excising the entire ear with surgical scissors, fixed in formalin (10%), and embedded in paraffin. 4 µm thickness sections were mounted onto glass slides and stained with H&E by the Johns Hopkins Reference Histology Laboratory. To measure epidermal thickness, 100 epidermal thickness measurements per mouse were averaged from images taken at ×200 magnification (Hamamatsu Nanozoomer) using ImageJ software (NIH).

ELISA

To measure cytokine release of BlaER-1, THP-1, N/TERT-1, NHEK, NK-92 MI, PBMCs and 3D human skin equivalent after stimulation with NET content and respective controls, ELISA Kits for hIL-8 (ELISA MAX™ Deluxe Set Human IL-8, Biolegend, 431504), hIL-6 (ELISA MAX™ Deluxe Set Human IL-6, Biolegend, 430504), IFN-γ (ELISA MAX™ Deluxe Set Human IFN-γ, Biolegend, 430104), and TNF (TNF alpha Human Uncoated ELISA Kit, Invitrogen, 88-7346-88) were used according to the manufacturer's instructions. Samples were assessed in triplicates.

qPCR analysis of IL-8 expression of 3D human skin equivalent

To investigate IL-8 expression of 3D human skin equivalent after stimulation with NET contents, qPCR analysis was performed. First, total RNA was isolated using the RNeasy Mini Kit (Qiagen, 74106) for animal tissues and cells. For cDNA preparation, the High-Capacity RNA-to-cDNA-Kit (Thermo Fisher, 4387406) was used according to the manufacturer's instructions. For the qPCR, the TaqMan™ system was used. Briefly, a master mix of TaqMan™ Universal Mastermix II (Thermo Fisher, 4440040) and TaqMan™ Gene Expression Assay (Thermo Fisher, 4448892) was prepared according to the manufacturer's instructions. For one reaction, 5.5 μL master mix and 4.5 μL of the respective cDNA (IL-8 and TBP) were mixed and the qPCR was run. Analysis in triplicates was performed using QuantStudio Real-Time-PCR software version 1.3 (Thermo Fisher).

Statistics

Experimental data were analyzed using Excel 2019 (Microsoft) and/or GraphPad Prism 8, microscopy data with ImageJ-Win64 or ZenBlue3 software, flow cytometry data with FlowJo V10. Normal distribution in each group was always tested using the Shapiro–Wilk test first for the subsequent choice of a parametric (ANOVA, Student's t-test for normally distributed data) or non-parametric (Mann–Whitney U) test as indicated in figure legelnds. p-values ($\alpha = 0.05$) were then calculated, and multiple testing was corrected for in Prism, as indicated in the figure legends. Values < 0.05 were generally considered statistically as significant and denoted by * throughout even if considerably lower. Comparisons made to unstimulated control, unless indicated otherwise, were denoted by brackets.

1086   **Supplementary Tables**

1087   **Supplementary Table 1: commercial TLR ligands and inhibitors; enzymes**

Component	Company	Product no.
Cl-amidine	Merck	506282
CpG PTO 2006	TIB Molbiol	n/a, see Table S2
CU-CPT9a	Invivogen	inh-cc9a
DNase I	Thermo Fisher	EN0521
DNase inhibitor 50 mM EDTA	Thermo Fisher	EN0521
DOTAP	Roth	L787.2
Ionomycin	Sigma	I0634-1MG
LL37	Invivogen	tlrl-l37
LPS-EK (ultrapure)	Invivogen	tlrl-peklps
Nigericin	Invivogen	tlrl-nig
PMA	Invivogen	tlrl-pma
R848 (Resiquimod)	Invivogen	tlrl-r848
RNase A	Thermo Fisher	EN0531
RNase inhibitor	Promega	N2615
TL8-506	Invivogen	tlrl-tl8506

1088

1089   **Supplementary Table 2: Nucleic acid TLR agonists**

Component	Sequence	Backbone	Company
RNA40	5'GsCsCsCsGsUsCsUsGsUsUsGsUsGsUsGsAsCsUsC3'	Phosphorothioate	Eurogentec
CpG PTO 2006	5'TsCsGsTsCsGsTsTsTsTsGsTsCsGsTsTsTsTsGsTsCsGsTsT3'	Phosphorothioate	TIB

1090

1091   **Supplementary Table 3: Antibodies and conjugation kit**

Item	Fluorophore	Species	Isotype	Company	Product no.
Anti-hLL37	DyLight 550	rabbit	IgG	LSBio	LS-B6696-500
DyLight 550 Conjugation Kit (Fast)	DyLight 550	-	-	Abcam	ab201800

Anti-rRNA (Y10b)	unconjugated	mouse	IgG$_{2a}$ κ	Santa Cruz Biotechnology	sc-33678
Anti-rRNA (Y10b) Alexa Fluor® 647	AF647	mouse	IgG$_{2a}$ κ	Santa Cruz Biotechnology	sc-33678 AF647
Anti-mouse IgG	AF647	chicken	IgY	Thermo Fisher	A-21463
Hoechst 33342	-	-	-	Thermo Fisher	H21492
SYTO RNAselect Green fluorescent dye	-	-	-	Thermo Fisher	S32703
Anti-mouse Cy3	Cy3	goat	IgG	Jackson ImmunoResearch	115-165-146
Anti-rabbit AF488	AF488	goat	IgG	Molecular Probes	A-11008
Anti-mouse 6 nm gold	-	goat	IgG	Jackson ImmunoResearch	115-195-166
Anti-rabbit 12 nm gold	-	goat	IgG	Jackson ImmunoResearch	111-205-144

1092

Supplementary Table 4: Click chemistry reagents

Reagent	Company	Product no.
AF546-Azide	Jena Bioscience	CLK-1283-1
Aminoguanidine-Hydrochloride	Merck	396494-25G
CuSO$_4$-click chemistry grade	Jena Bioscience	CLK-MI004-50.1
5-Ethynyluridine	Jena Bioscience	CLK-N002-10
Na-Ascorbate-click chemistry grade	Jena Bioscience	CLK-MI005-1G
THPTA (Tris((1-hydroxy-propyl-1H-1,2,3-triazol-4-yl)methyl)amine)	Jena Bioscience	CLK-1010-25

1094

Supplementary Table 5: Plasmids used for HEK293T transfection

Plasmid name insert	Vector backbone	Insert
NF-κB reporter	pGL3	6x NF-κB response element
Renilla	pRL-TK	*Renilla*
hTLR7	pcDNA3.1 (+)	hTLR7
hTLR8	pcDNA3.1 (+)	hTLR8
hTLR9	pSEM-hTLR9	hTLR9

1096

Supplementary Table 6: primers used for qPCR

Gene	Assay number
IL-8	Hs00174103_m1
TBP (housekeeper)	HS00427620_m1

1098

1099   **Supplementary Table 7: 10x Ammonium chloride erythrocyte lysis buffer**

Compound	Company	Product no.
1.54 M NH$_4$Cl	Roth	5470.1
100 mM KHCO$_3$	Fluka	60220
1 mM EDTA; pH 8 dissolved in Ampuwa water	Thermo Fisher Fresenius Kabi	15575020 1833
pH adjusted to 7.3, sterile filtered (0.22 µm)		

1100

A human PMNs

PMA | Nigericin | live *C. albicans*

synthetic RNA+LL37 | bacterial RNA+LL37 | fungal RNA+LL37

Hoechst rRNA Y10b

B

human PMNs — DNA — RNA

NET related signal dispersion (rel. ROI number/ROI average size)

Unstim. | PMA | ssRNA+LL37 | bRNA+LL37 | fRNA+LL37 | Nigericin | live *C.albicans*

C murine BM-PMNs

PMA | synthetic RNA+LL37

Nigericin | live *C. albicans*

Hoechst rRNA Y10b

D stem cell-derived PMNs

Unstim. | PMA | PMA crop

− 5-EU

+ 5-EU

Hoechst rRNA Y10b 5-EU

10 µm | 2 µm

E human PMNs

Hoechst rRNA Y10b

10 µm | 10 µm | 1 µm

G

NET derived RNA

naRNA (ng/µL)

Mock | PMA

H

RNA-Seq

RNA types (%)

whole RNA | naRNA

- unknown
- rRNA
- ncRNA
- nonsense mediated decay
- altern. spliced transcript
- pseudogene
- mRNA

F human PMNs

Unstim. | PMA

200 µm | 20 µm | 200 µm | 5 µm | 1 µm

A

PMNs from HD1 +/- RNase inh.

PMA

unstimulated

wash, add new medium, scrape; +/- RNase inh.

PMA NETs

Mock NETs

PMNs from HD2

+ PMA NETs

+ Mock NETs

B human PMNs

Mock NETs+RNase inh. | PMA NETs | PMA NETs+RNase inh.

1:50

1:500

C human PMNs

NET related signal dispersion (rel. ROI number/ROI average size)

RNase inh.

Unstim. PMA Mock PMA PMA Mock PMA PMA

1:50 1:500

NETs

D human PMNs

Mock NETs | Mock NETs RNase treated

PMA NETs | PMA NETs RNase treated

E human PMNs

number of tiles with NETs

Unstim. PMA Mock PMA RNase
Mock PMA NETs

F human PMNs

PMA NET naRNA | PMA NET naRNA+LL37

G human PMNs

Unstim. | Unstim. crop

PMA | PMA crop

Hoechst rRNA Y10b LL37

H human PMNs

coloc. with LL37 (PCC)

DNA RNA

Unstim. PMA

I human PMNs

Intensity

Shared co-localization maxima

Unstim.

rRNA
LL37
DNA

Distance (μm)

J human PMNs

K human PMNs

200 nm

5 μm

Shared co-localization maxima

Gray value

Unstim.

rRNA
LL37
DNA

Distance (pixels)