PROTEIN CODING VARIATION IN OUTBRED LABORATORY

MOUSE STOCKS PROVIDES A MOLECULAR BASIS FOR

DISTINCT RESEARCH APPLICATIONS.

Belinda Cornes[1], Carolyn Paisie[1], Emily Swanzey[1], Andrew Schile[1], Kelly Brackett[1], Laura

Reinholdt*[1], Anuj Srivastava*[1]

1. The Jackson Laboratory, 600 Main Street, Bar Harbor, ME 04609

*co-corresponding

laura.reinholdt@jax.org

anuj.srivastava@jax.org

26 **keywords:** outbred stocks, laboratory mice, reproducibility, exome, genetic quality control

28 **running title:** Exome sequencing of outbred laboratory mouse stocks

ABSTRACT

Outbred laboratory mice (*Mus musculus*) are readily available and have high fecundity, making them a popular choice for biomedical research, especially toxicological and pharmacological applications. Direct high throughput genome sequencing (HTS) of these widely used research animals is an important genetic quality control measure that ensures research reproducibility. HTS data have been used to confirm the common origin of outbred stocks and to molecularly define distinct outbred populations. But these data have also revealed unexpected population structure and homozygosity in some populations; genetic features that emerge when outbred stocks are not properly maintained. We used exome sequencing to discover and interrogate protein coding variation in a newly established population of Swiss-derived outbred stock (J:ARC) that is closely related to other, commonly used CD-1 outbred populations. We used these data to describe the genetic architecture of the J:ARC population including heterozygosity, minor allele frequency, LD decay, and we defined the novel, protein-coding sequence variation. These data reveal the expected genetic architecture for a properly maintained outbred stock and provide a basis for on-going genetic quality control. We also compared these data to protein coding variation found in a multiparent outbred stock, the Diversity Outbred (J:DO). We found that the more recently derived, multiparent outbred stock has significantly higher interindividual variability, greater overall genetic variation, higher heterozygosity, and fewer novel variants than the Swiss derived J:ARC stock. However, among the novel variants found in the J:DO stock, significantly more are predicted to be protein-damaging. That individuals from this population can tolerate a higher load of potentially damaging variants highlights the buffering effects of allelic diversity and the differing selective pressures in these stocks. While both outbred stocks offer significant individual heterozygosity, our data provide a molecular basis for their intended applications, where the J:DO are best suited for studies requiring maximum, population-level genetic diversity and power for mapping, while the J:ARC are best suited as a general-purpose outbred stock with robust fecundity, relatively low allelic diversity, and less potential for extreme phenotypic variability.

INTRODUCTION

There are 1,374 inbred strains and outbred laboratory mouse stocks cited in the biomedical research literature (BULT *et al.* 2019). Genome sequencing of inbred strains has confirmed the expected homozygosity and intrastrain genetic divergence that are the results of decades of inbreeding and reproductive isolation (MOUSE GENOME SEQUENCING *et al.* 2002; KEANE *et al.* 2011; YALCIN *et al.* 2011; CHANG *et al.* 2017; SRIVASTAVA *et al.* 2017; LILUE *et al.* 2018). Because inbred mice are genetically identical, interindividual phenotypic variability can be attributed to experimental variables and to a lesser extent, epigenetic variation (WEICHMAN AND CHAILLET 1997). These unique features make inbred mouse strains a popular choice for biomedical researchers who endeavor to minimize genetic variability and to quantify experimental (extrinsic, non-genetic) variability. However, inbred strains do not recapitulate the extent of interindividual variation found in a human cohort, patient group, or population making them inappropriate for studies that seek to model such variation.

In contrast to inbred strains, outbred stocks exhibit interindividual variation and heterozygosity, coupled with higher fecundity. The four most widely cited outbred stocks are CF-1, Swiss Webster, NMRI, and CD-1 (BULT *et al.* 2019), but all trace their origins to a colony at The Rockefeller Institute that was established with 2 male and 7 female 'Swiss' mice imported from the Pasteur Institute in 1926 (CHIA *et al.* 2005). Over the ensuing decades, mice from these stocks were imported by commercial breeders, NIH, and academic institutions where colonies were ideally maintained with at least 25 breeding pairs to maintain a coefficient of inbreeding (F) of less than 1% per generation with selection for maximum fecundity (FESTING *et al.* 1972) (CHIA *et al.* 2005). A variety of approaches have been used to examine the genetic variation resident in these outbred stocks including alloenzyme analysis and other biochemical approaches (GROEN AND LAGERWERF 1979; RICE AND O'BRIEN 1980; CUI *et al.* 1993), high-density SNP panels (ALDINGER *et al.* 2009; YALCIN *et al.* 2010), and most recently low coverage genome sequencing (YALCIN *et al.* 2010; NICOD *et al.* 2016). Data from these studies provide estimates of minor allele frequency, heterozygosity, and coefficients of inbreeding. Specifically, low coverage sequencing of these stocks has revealed 1) relatively low allelic diversity given their common origin, 2) appreciable levels of heterozygosity of common variants (0.20-0.35) in properly maintained populations, and 3) relatively few novel variants (<5%) relative to the

C57BL/6J reference genome and other common inbred laboratory strains (YALCIN *et al.* 2010;

NICOD *et al.* 2016).

The JAX Swiss Outbred stock was imported to The Jackson Laboratory (JAX) from The Animal

Resources Centre (ARC) in Canning Vale, Western Australia in 2020 as a group of 128

individuals (64 male and 64 female). To fully characterize the extent of protein coding variation

in this new outbred population, we sequenced the exomes of the 32 male and 32 female G3

animals that were used to initiate new breeding funnels at JAX (J:ARC). We used these data to

estimate inbreeding, heterozygosity, and interindividual variation, and to molecularly define the

colony. These metrics confirmed the outbred nature of this new colony of CD-1 stock, and

additionally reveal 5,105 (3.8%) novel protein coding and splice site variants, which falls in the

range of the proportion of novel variants found in other outbred colonies and stocks (YALCIN *et*

al. 2010; NICOD *et al.* 2016). To provide a molecular genetic quality control standard for outbred

stocks, we compared the J:ARC exome call sets to exome variant call sets from very different

outbred stock, derived from a multiparent outbred population, the Diversity Outbred (J:DO). We

found that the J:DO have higher population level genetic variation (~3.6X), higher interindividual

variation, higher heterozygosity, and higher allelic diversity due to their multi-substrain parental

origins. We found fewer novel variants in the J:DO, yet these novel variants are predicted to

have higher impact on protein function. These common and distinct genomic features of the

J:ARC and J:DO can serve as the basis for estimating sample size, where fewer J:ARC mice

are needed to capture the full range of segregating variation in the population, but a similarly

sized cohort of J:DO mice offers more interindividual genetic variation and higher population

level genetic variation. Both types of outbred stocks can be used for genetic mapping,

however, their distinct genetic architectures must be accounted for in study design as

previously shown (GATTI *et al.* 2014; NICOD *et al.* 2016), and should be regularly monitored to

ensure reproducibility.

MATERIALS & METHODS

Samples

J:ARC(S), JAX Swiss Outbred. The JAX Swiss Outbred stock was imported to The Jackson

Laboratory from The Animal Resources Centre (ARC), Canning Vale, Western Australia in 2020.

The imported mice (G0) were paired into 64 breeding units, and sperm and eggs were

harvested from the G1 offspring. To establish each of the 32 breeding funnels for live colony

maintenance in the JAX vivarium barrier, two distinct units were selected for reciprocal *in vitro*

fertilization and IVF-generated embryos were pooled. The resulting live-born animals from each

funnel were designated G2 (Figure 1). These mice were subsequently bred according to the

Poiley rotational breeding scheme to produce 32 breeder pair units / funnels at G3(POILEY

1960). Spleen samples were collected the 64 G3 J:ARC (J:ARC(S), RRID:IMSR_JAX:034608,

JAX stock number 034608) (32 females and 32 males) that populated each unit (Supplementary

Data 1).

J:DO, JAX Diversity Outbred. We collected spleen from 20 J:DO, Diversity Outbred

(RRID:IMSR_JAX:009376, JAX stock #9376) mice from The Jackson Laboratory, 10 female and

10 male, non-sibling mice, breeding generation 43 (Supplementary Data 1).

DNA Isolation, Exome Library Preparation, and Sequencing

DNA isolation, exome library preparation, and sequencing were performed by Genome

Technologies at The Jackson Laboratory. DNA was isolated from spleen using the NucleoMag

Tissue Kit (Machery-Nagel) according to the manufacturer's protocol. DNA concentration and

quality were assessed using the Nanodrop 8000 spectrophotometer (Thermo Scientific), the

Qubit Flex dsDNA BR Assay (Thermo Scientific), and the Genomic DNA ScreenTape Analysis

Assay (Agilent Technologies). Mouse exome libraries were constructed using the KAPA

HyperPrep Kit (Roche Sequencing and Life Science) and SureSelect XT Community Design

Mouse All Exon v2 Target Enrichment System (Agilent Technologies), according to the

manufactures' protocols. Briefly, the protocol entails shearing the DNA using the E220

Focused-ultrasonicator (Covaris), size selection targeting 400 bp, ligating Illumina specific

adapters, and PCR amplification. The amplified DNA libraries are then hybridized to the Mouse

All Exon probes (Agilent Technologies) and amplified using indexed primers. The quality and

concentration of the libraries were assessed using the High Sensitivity D1000 ScreenTape

(Agilent Technologies) and KAPA Library Quantification Kit (Roche Sequencing and Life

Science), respectively, according to the manufacturers' instructions. Libraries were sequenced

150 bp paired-end on an Illumina NovaSeq 6000 using the S4 Reagent Kit v1.5.

Sequence Data Analysis

An overview of the sequence data analysis pipeline is shown in Supplementary Figure 1. All

reads were subjected to quality control using an in-house QC script. Samples with base

qualities greater ≥ 30 over 70 % of read length were used in the downstream analysis. High-

quality reads were mapped to the mouse genome (build-mm10) using BWA-mem (bwa-0.7.9a)

at default parameters (ALDINGER *et al.* 2009). The resulting alignment was sorted by coordinates

and converted to binary alignment map (BAM) format by Picard v 2.8.1- SortSam utility

(http://picard.sourceforge.net). Picard-MarkDuplicates module was used to remove the

duplicates from data. The Genome Analysis Toolkit (GATK v4) (MCKENNA *et al.* 2010; DEPRISTO

et al. 2011) module BaseRecalibrator was used to pre-process the alignments. Target capture

efficiency was determined using Picard-HsMetrics (1.95). The recalibrated bam alignment file

was used to input GATK-Haplotype Caller at parameters -stand_call_conf 50.0, -

stand_emit_conf 30.0 and variants calls were restricted to the target region (Mouse All Exon v2).

Finally, raw variants calls were soft filtered using GATK VariantFiltration (DP < 25, QD < 1.5 and

FS > 60), annotated by snpEff 3.6.c (CINGOLANI *et al.* 2012) and the highest impact variant was

reported by GATK VariantAnnotator. All variants were further annotated with mouse dbSNP

v150.

GATK-HaplotypeCaller in GVCF mode was used for joint genotyping of J;DO, J:ARC, and

combined (J:DO and J:ARC) samples. The CombineGVCF utility was used to gather all the

samples and then executed the GenotypeGVCF command. All analyses were performed on

jointly called variants and only included variants found in both "J:ARC only" and "J:DO only"

called datasets (N: 120,864). This effectively revealed the differences between the two datasets

and directly provided the data needed to resolve genotypes.

Heterozygosity

To observe the distribution of mean heterozygosity across all samples, Plink was used

(PURCELL *et al.* 2007). The `--het` option in Plink was used to create heterozygous information

for each sample which included the observed number of homozygous genotypes `[O(Hom)]`

and the number of non-missing genotypes `[N(NM)]`. This information was then used to

calculate the observed heterozygosity rate per individual using the formula `(N(NM) -

O(Hom))/N(NM)`.

To calculate per variant heterozygosity for variants shared between J:DO and J:ARC, the

number of samples in which a shared variant was heterozygous was divided by the total

number of genotyped samples. The inbreeding coefficient (F_H), was calculated

using *VCFtools* (–het) as previously described (Stoffel *et al.* 2016; Foster *et al.* 2021).

PLINK was used to calculate r^2 for all pairs of autosomal SNPs called from joint genotyping in

the J:ARC samples (425,409) and the J:DO samples (117,429). SNPs that were missing in

more than 5% of the samples and that were monomorphic were removed. The 95th percentile of

r^2 values for SNPs spaced up to 1 Mb apart were selected.

Allele Frequency

194   To observe differences between allele frequencies across the genome between J:DO and

195   J:ARC samples, the minor allele frequency (MAF) was calculated separately for J:DO and

196   J:ARC samples at each variant site using the `--freq` option in Plink.

Clustering/Dendrogram

Principal Component Analysis (PCA) plots were performed in Plink. PCA is a multivariate

statistical method used to produce any number of uncorrelated variables (or principal

components) from a data matrix containing observations across a number of potentially

correlated variables. The principal components are calculated so that the first principal

component accounts for as much variation in the data as possible in a single axis of variation

(component), followed by additional components. Here the variants called in both J:ARC and

J:DO datasets were used to observe the similarities within and between datasets.

To show similarity groups among the J:ARC samples, a dendrogram was performed using the

ggdendro package in R-3.6.2 (https://cran.r-project.org). A dendrogram is a tree diagram

showing hierarchical clustering that represents the relationships of similarity among a group of

entities.

Novel Variants

The joint genotyping variant call file (vcf) of J:ARC and J:DO were flagged for known variants in

dbSNP150 (Sherry *et al.* 2001), European Variation Archive (EVA) (Cezard *et al.* 2022) , Sanger

mouses genome project (KEANE *et al.* 2011) (ftp://ftp-mouse.sanger.ac.uk//REL-2004-v7-

SNPs_Indels/mgp_REL2005_snps_indels.vcf.gz), mouse Collaborative Cross genome

(ftp://ftp.sra.ebi.ac.uk/vol1/ERZ460/ERZ460702/Merged_69_flagged.tab.vcf.gz) (SRIVASTAVA *et*

al. 2017), and Diversity Outbred low pass sequencing (in-house resource) using vcftools 'vcf-

annotate' (KEANE *et al.* 2011) and snpsift_4.2 (CINGOLANI *et al.* 2012). We also flagged J:ARC

variants for their presence in J:DO joint and single sample variant call sets and did a similar

analysis for J:DO variants. The variants not present in any of these resources were considered

novel and were further annotated by snpeff v4.3 (CINGOLANI *et al.* 2012) using the mouse

GRCm38.75 snpeff database. Finally, highest effect variants are selected by gatk-3.6

VariantAnnotator (MCKENNA *et al.* 2010). Functional gene and pathway annotations (including

KEGG, GO, UP, BIOCARTA, REACTOME) for the genes harboring novel high impact variants

were compiled using DAVID Bioinformatics tools. DAVID Bioinformatics tools were then used

to identify clusters of genes with related functional annotations using a medium classification

stringency and a Benjamini score of >0.05 as the significance threshold for enrichment

(SHERMAN *et al.* 2022)

229   Results and Discussion

The J:ARC outbred stock currently maintained at The Jackson Laboratory was

established through the importation of mice from the Australian Animal Resource Center (ARC)

in 2020. The origins of this stock are CD-1 mice that were acquired by ARC from Charles River

Laboratories in 1991 and 2005. Like most outbred laboratory mice, CD-1 mice trace their

origins to 2 male and 7 female "Swiss" mice that were obtained by The Rockefeller Institute in

1926 (CHIA *et al.* 2005). The live J:ARC colony was initiated with founder animals (G3) in 32

breeding funnels. We sought to determine the level of inbreeding and relatedness in the

founder animals of this colony and to catalog the coding variants segregating in each funnel for

future use in genetic quality control monitoring (STROBEL *et al.* 2015). Since some outbred

stocks differ quite dramatically, we sought to compare the genetic architecture of the outbred

J:ARC to a multiparent outbred population (J:DO) to demonstrate the differing genetic

architectures of these stocks and to establish a reference set of genetic variants for genetic

monitoring. The J:DO mice are derived from eight founder inbred strains representing the three

major subspecies of laboratory mice, *M.m.musculus, M.m.domestics*, and *M.m.castaneous*.

This population captures 90% of the genetic variation in laboratory mice and with each

generation of breeding, sufficient accumulated recombination to ensure high LD decay, and

thereby maximum power for genetic mapping (GATTI *et al.* 2014; SAUL *et al.* 2019). This

population is maintained at the Jackson Laboratory through 175 breeding funnels using

MateSel(KINGHORN AND KINGHORN 2021) to minimize inadvertent phenotypic selection, to

optimize diversity, and to prevent intracolony structure in each generation based on pedigreed

population history.

Genetic variation. We generated exome data from 64 J:ARC mice (32 male and 32 female)

representing 32 breeding funnels at G3 and from 20 J:DO mice (10 male and 10 females) at

generation 43. We generated ~203M high-quality reads on average, and after alignment to the

reference genome, the mean target coverage was 162X (91% target covered at 30X) for J:ARC

samples and 155X (90% target covered at 30X) for J:DO samples. Overall, 90% of the target

exome regions were covered by 30 or more reads in both sets of samples (see Supplementary

Figure 2).

Through joint genotyping, we identified 478,011 and 135,825 variants in J:DO and

J:ARC samples, respectively, where a variant is defined as a SNP/INDEL present in a

sequenced sample when compared to the C57BL/6J, inbred mouse reference genome (mm10,

GRCm38). To adjust for the difference in sample size between the J:DO and J:ARC datasets

and any potential influence this could have on overall discovery rates, we randomly subset the

J:ARC samples to a set of twenty from joint genotyping results and then recorded the total

number of variants present in the subset. We repeated this analysis over 100 iterations and

found from this bootstrapping method that the J:DO harbor, on average, 3.6X more variants

overall than J:ARC. This higher allelic diversity is expected in the J:DO since the stock is a

multiparent population derived from eight diverse founder inbred strains.

Heterozygosity and inbreeding. To evaluate heterozygosity in these two outbred populations,

we estimated sample level heterozygosity for the entire set of variant calls (Figure 2A). The

average, individual heterozygosity in J:ARC is 24%, which is consistent with previously

estimates from low pass WGS for CFW outbred populations (NICOD *et al.* 2016) (YALCIN *et al.*

2010). In contrast, sample level heterozygosity in the J:DO samples was notably higher (nearly

40%) than J:ARC. Given that previous analyses have estimated J:DO haplotype-level

heterozygosity at around 80% (https://www.jax.org/-/media/jaxweb/files/jax-mice-and-

), it is important to note that we used reference-based

genotype calls and we calculated heterozygosity across the entire call set. Previously,

heterozygosity for J:DO mice was calculated on a limited set of variants based on genotype

probabilities of eight parental genotypes called from known polymorphic SNPs from the Mouse

Universal Genotyping Array (MUGA) series (MORGAN *et al.* 2015; SIGMON *et al.* 2020). To

recapitulate this haplotype-based approach using our exome variant calls, we used 3007 calls

at markers from GigaMUGA genotyping array (MORGAN *et al.* 2015) and for this subset of

variants, we found heterozygosity of ~82% consistent with previous array-based estimates

(see Supplementary Figure 3).

To compare the variant level heterozygosity between the two populations, we identified

the variants that are common between J:ARC and J:DO and then for each variant determined

the number of samples with heterozygous genotypes (Figure 2B). We found that a lower

proportion of J:ARC samples are heterozygous for any given variant than the J:DO, and that

there is greater intraindividual variability in the proportion of heterozygous variants in the J:DO

population. To further explore these differences in the distribution of heterozygosity, we

calculated the minor allele frequencies (MAF) for each of the common variants across the

genome (Figure 3A-C). We found that for each common variant, the MAF in the J:ARC was

consistently lower (mean 0.18, median 0.13) than the J:DO (mean 0.28, median 0.28) and

similar to previously published MAF for variants segregating in properly maintained outbred

CD-1 mouse populations (YALCIN *et al.* 2010).

Variant data can also be used to estimate the coefficient of inbreeding (F), which is a

metric that describes the distribution of variants in a population. According to the Hardy-

Weinberg (HW) principle, F=0 for a given variant indicates that it is in HW equilibrium. By this

measure, a fully inbred strain has an inbreeding coefficient of F=1.0 (100%). When calculated

across all J:ARC variants and samples, the average level of inbreeding is -.033 (-3.3%), which

falls within expectations for an outbred population (YALCIN *et al.* 2010). Though fewer samples

were used to generate the J:DO dataset, the inbreeding coefficient calculated from the J:DO

was closer to HW equilibrium at -0.007 (-0.7%). These F coefficients indicate that both

populations are maximally outbred. The negative values are the result of heterozygous calls

that exceed HW expectations which in these datasets could be due to erroneous genotype

calls or low sampling of the populations. While the coefficients of inbreeding for these

populations are low, there are regions of the genome where shared variants are fixed. These

regions of homozygosity are distributed through the J:ARC and J:DO genomes (Figure 4). It is

likely that some homozygous variant calls (regions of homozygosity, ROH) are the result of

heterozygous deletions. Because our methodology doesn't distinguish between potential

hemizygous and true homozygous variant calls, we may be overestimating the extent of ROH

in these genomes.

Interindividual variability. An important consideration in selecting an outbred laboratory mouse

strain and sample size in experimental design is interindividual genetic variability. Taking

advantage of our variant data, we used principal component analysis to assess interindividual

and interstrain variation in the J:ARC and J:DO. Using all jointly called variants, we found the

expected interstrain variation (PC1) but more interindividual variability in J:DO samples than

J:ARC (PC2) (Figure 5A). These data show that greater genetic diversity can be achieved with

fewer samples in the J:DO by at least an order of magnitude. Unbiased clustering of the J:ARC

variant data recapitulated the known pedigree and sibling relationships of the J:ARC samples

(Figure 5B), highlighting the utility of variant data for pedigree reconstruction in outbred

populations.

LD Decay. Linkage disequilibrium (LD) describes the degree to which variants co-segregate in

a population. Meiotic recombination leads to new haplotypes and is the molecular mechanism

that drives LD decay. Pairwise analysis of linked markers in an interval is used to assess LD

decay in a population and these data are useful for inferring swept radius for marker spacing in

genetic association studies. Higher LD decay in a mapping population provides higher

resolution for genetic mapping. We examined this facet of genetic architecture in the J:ARC

and J:DO populations and while our sample size was low, we found that both populations offer

LD decay sufficient for genetic mapping, with higher LD decay in the J:DO ($LD_{1/2}$ = ~0.1 Mb)

(Figure 5C). Our estimate of LD decay is higher than previous estimates for J:DO, but this is

consistent with the more advanced breeding generation used in our study (GATTI *et al.* 2014).

LD decay in the J:ARC is consistent with previous estimates for commonly used laboratory

outbred populations ($LD_{1/2}$ = ~0.3 Mb) *(YALCIN et al. 2010).*

Novel Variants. To identify the novel variant calls in the J:ARC and J:DO, we compared the

calls to all published variants from all sequenced laboratory mouse strains in dbSNP, EVA, the

Sanger mouse genomes project (LILUE *et al.* 2018), Mouse Collaborative Cross Genome

(SRIVASTAVA *et al.* 2017), and to in house J:DO genome variation call sets (unpublished). Of the

135,825 and 478,011 variants found in J:ARC and J:DO respectively, 96.24 and 99.08% are

variants that have been identified in other laboratory mouse strains. We identified 5,105 [2,191

> AF 0.2] and 4,308 [1,888 > AF 0.2] novel variants respectively (Table 1). The larger number of

novel variants in J:ARC mice is attributable to the relative lack of published variant data from

Swiss derived outbred stocks compared to the complete catalog of variation available from the

eight founder inbred strains of the Collaborative Cross recombinant inbred panel from which

the J:DO were derived. Functional annotation of the novel variants revealed that J:DO have

nearly twice the number of protein-damaging variants (including exon deletions, frameshift,

stop gain / loss, splice acceptor / donor sites, start loss) ((J:DO:797 [329 > AF 0.2, HIGH],

J:ARC: 408 [151 > AF 0.2, HIGH]) (Figure 6). The details of each variant are provided in

Supplementary Data 2. The genes harboring these high impact variants in J:DO show

functional enrichment for post-translational modifications (Ubl conjugation, isopepeptide

bond), DNA repair, and translational regulation / RNA binding; while the genes harboring high

impact variants in J:ARC do not show functional enrichment. The novel protein damaging

variants in J:DO are variants that have naturally arisen in the population but have not been lost

through purifying selection or through artificial selection. These genetic differences are

attributable not only to the distinct origins and ages of these outbred stocks, but to the

different breeding paradigms and selective pressures that have occurred during their

maintenance. For example, many outbred mouse stocks and populations have been subject to

artificial selection for desirable laboratory traits (fecundity, docility, size, etc.) leading to lack of

variability for some phenotypes. One example is body weight where variation is less desirable

because it complicates pharmacological applications, i.e. dosing studies. As a result, body

weight variation in CD-1 and related outbred like J:ARC is minimized, however this is not the

case in J:DO where a larger range of body weights is observed (Supplementary Figure 4).

369   CONCLUSIONS

We used exome sequencing to define protein coding variation in two outbred populations of

mice with distinct origins. We found that the multiparent, J:DO outbred population has more

than 3X population-level variation than the CD-1 derived J:ARC outbred population. Both

populations have coefficients of inbreeding that are consistent with previous estimates,

confirming that both are maximally outbred. Our analysis of variants that are shared between

the J:ARC and J:DO shows that individual heterozygosity in both populations is high, but allelic

diversity, intraindividual variability, and minor allele frequency are higher in the J:DO

population. For research applications this means that similarly sized cohorts of J:DO mice will

provide more genetic diversity than J:ARC, making the former population the better choice for

studies that endeavor to maximize genetic diversity. Our data and multiple previous studies

show that both outbred populations are useful for genetic mapping. But the higher genome-

wide LD decay in J:DO will confer higher mapping resolution.

Overall, we found more novel variants in J:ARC compared to J:DO and this is likely due

to the relative paucity of published / accessible variant data for commonly used outbred

strains, especially CD-1 from which the J:ARC population was derived. While there were fewer

novel variants in the J:DO, more of these were variants that are predicted to be protein

damaging and potentially more likely to reduce survival and/or fecundity. While this study

doesn't allow us to make any inferences about the load of deleterious alleles and their

respective contributions to fecundity, litter size is a trait that is frequently used to estimate

fecundity and breeding data from The Jackson Laboratory show that J:DO females have litter

sizes that are comparable to many common laboratory inbred strains (mean litter size = 9.9,

https://www.jax.org/strain/009376). The J:ARC population is derived from an outbred stock

that has been subject to nearly a century of artificial selection for high fecundity, docility, and

other desirable laboratory traits including consistent body weight which is preferred for dosing

studies. In contrast, the J:DO are derived from phenotypically diverse recombinant inbred lines

(Collaborative Cross) and artificial selection for any phenotype is actively avoided through

randomized breeding that optimized for genetic diversity. These differences in selective

pressure may also explain the apparent loss of deleterious alleles through drift in the J:ARC.

That deleterious alleles have not been subject to purifying selection in the J:DO, also highlights

the potential buffering effects of high allelic diversity and heterozygosity.

Our data, together with previously published studies, reinforce the fact that not all

outbred stocks have comparable genetic and phenotypic diversity. Direct sequencing of

outbred stocks provides molecular basis for genetic quality control, breeding paradigms, and

optimization of experimental design for effective deployment these strains in biomedical

research.

Study limitations. Our exome sequencing data provide just a subset of the overall genetic

variation segregating in these outbred populations, specifically SNPs and small insertions /

deletions in coding sequence. Structural variation (SV), which we have not profiled here, is also

a significant source of genetic variation. SV have the potential to impact multiple genes with

potentially large phenotypic effects. While certain types of structural variants (deletions) can be

robustly detected using short read sequencing data, long read sequencing and *de novo*

assembly are the gold standard for genome-wide detection of SV. Moreover, outside of splice

junction sites and UTRs, our data do not capture non-coding variation, yet non-coding

variation is also a recognized source of phenotypic variation, especially for complex traits.

Finally, while our sequencing depth per sample is high, our samples sizes are low, making our

study underpowered for detection of ultra-rare SNPs/INDELs.

Tables

Stock	Total Variants	Mouse Collaborative Cross	Diversity Outbred Mouse Genomes	SangerSNP_Indel_May_2020	dbSNP	EVA	Novel	High Impact Novel Variants
J:DO	478011	468487	427929	452989	439682	439642	4308	797
J:ARC	135825	116849	109808	121395	115557	115561	5105	408

419
420
Table 1. Total variants (SNPs/INDELs) and total novel variants in J:DO and J:ARC stocks.

Comparison of the total variants to all variants in the major, publicly available mouse variant

resource revealed that 0.9% and 3.75% of the total variants found in J:DO and J:ARC

respectively are novel.

Figure Legends

Figure 1. The Poiley method used for the importation and breeding of the JAX Swiss Outbred

(J:ARC) population. G0 were live animals from The Animal Resources Centre (ARC) in Canning

Vale that were subsequently bred and rederived through IVF to create 32 distinct breeding lines

at The Jackson Laboratory which continue to be maintained according to the Poiley method to

minimize inbreeding.

Figure 2. Heterozygosity in J:ARC (green) and J:DO (blue) datasets. The proportion (**A**) of

heterozygous calls per samples and distribution of heterozygous calls (**B**) for J:ARC (green) and

J:DO (blue) datasets. The mean heterozygosity for J:ARC samples was 0.25 while in J:DO it

was 0.39. Samples from both datasets were joint called and this analysis included only those

variants that are were called in both datasets (120,864)

Figure 3. Allele Frequency in J:ARC (green) and J:DO (blue) datasets. Samples from both

datasets were joint called and this analysis was restricted to those variants found in keeping

both datasets (120,864). Allele frequency was calculated using Plink. (**A**) MAF distribution for

120,864 SNPs for J:ARC (green) and J:DO (blue). Monomorphic variants are not removed. (**B**)

Allele frequency at the variant level for J:ARC samples (mean 0.18, median 0.13). (**C**) Allele

Frequency for J:DO samples at the variant level (mean 0.28, median 0.28).

Figure 4. Consensus segments of homozygosity for J:ARC (green) and J:DO (blue). The

number of consensus segments of homozygosity in J:ARC samples was higher (**A**) than in

J:DO samples (**B**). Plink's default parameters were used. Only runs of homozygosity containing

at least 100 SNPs, and of total length ≥ 1000 kilobases, are noted.

Figure 5. Interindividual variability, relatedness, and LD decay. (**A**) PCA plot of J:DO (blue) and

J:ARC (green) joint variant calls. (**B**) A dendrogram of the J:ARC samples (using all called

variants in the dataset) showing the samples clustering in known sibling relationships. (**C**)

Comparison of LD decay J:ARC samples (green) and J:DO samples (blue). Each curve was

plotted using the 95th percentile of r2 values for SNPs spaced up to 1Mb apart. Plink was used

to calculate r^2 for all pairs of autosomal SNPs called from joint genotyping in the J:ARC

samples (425,409) and the J:DO samples (117,429). SNPs that were missing in more than 5%

of the samples and that were monomorphic were removed.

Figure 6. Novel variant consequences. (**A**) Showing the SnpEff effect categories for novel

variants for J:DO (on the left) and J:ARC (on the right) (**B**) High-impact (SnpEff effect

categories) novel variants are enriched in J:DO (on the left) in comparison to J:ARC (on the

right)

Supplementary Material

Supplementary Data 1. J:DO and J:ARC samples used for the study.

Supplementary Data 2. All novel, protein coding variants in J:ARC and J:DO with functional annotation and allele frequency. Functional annotation (DAVID Bioinformatics Resources) clustering of genes harboring novel variants in J:ARC and J:DO.

Figure S1. The Jackson Laboratory (JAX) Nexflow pipeline for laboratory mouse whole exome sequencing.

Figure S2. Exome sequencing coverage statistics. **(A)** The mean target coverage versus high quality filtered reads showing mean target coverage was 162X (91% target covered at 30X) for J:ARC samples (green) and 155X (90% target covered at 30X) for J:DO (blue) samples. **(B)** Percent target bases at 30X versus high quality filtered reads showing 90% of the target exome regions were covered by 30 or more reads in both sets of samples.

Figure S3. The proportion of heterozygous variants based on the subset of variant calls for known SNPs within the Mouse Universal Genotyping Array probe set. There were 3007 exome variant calls at positions that are included in the GigaMUGA probe set. These were used to estimate the proportion of heterozygous calls per J:DO using the R package, R/qtl2. Estimates are based on autosomes at the haplotype level, using genotype probabilities of the 8 J:DO parental inbred strain genotypes.

Figure S4. Distribution of body weight in J:ARC and J:DO mice. Jackson Laboratory data (unpublished, Schile 34608) and publicly available datasets from the Mouse Phenome Database ((Shorter8, Chesler4 (LOGAN *et al.* 2013; RECLA *et al.* 2014), https://phenome.jax.org/)and Wright et al. (WRIGHT *et al.* 2022) were used to assess the distribution of body weight in J:DO and J:ARC mice). Data include both males and females and sub-sets of 60 were randomly sampled from larger datasets for comparison. Overall body weight in the J:ARC is higher than J:DO (ANOVA, adjusted P value <0.0001), while the distribution of body weight (range) is greater in J:DO.

Acknowledgements

This work was partially supported by The Jackson Laboratory and the Mouse Mutant Resource and Research Center at The Jackson Laboratory (NIH U42 OD010921, LGR). We gratefully acknowledge the contribution of the Genome Technologies Service at The Jackson Laboratory for expert assistance with exome library preparation and sequencing.

Author Contributions

Conceptualization   K.B, L.R., A.Sr.

Methodology   A.S., B.C.

Formal analysis   A.S., B.C.

Investigation   A.S., B.C.

Resources   C.P., E.S., A. Sh.

Data Curation   A.S., B.C.

Writing, Review & Editing   L.R., A.Sr., B.C., K.B.

Visualization   B.C., A.Sr., L.R.

Supervision   A.Sr., L.R.

Funding acquisition   L.R., K.B., A.Sr.

Data availability

All the sequence data used in this study has been submitted to NCBI Sequence Read Archive under the bioproject PRJNA835415 and study SUB11284953.

LITERATURE CITED

Aldinger, K. A., G. Sokoloff, D. M. Rosenberg, A. A. Palmer and K. J. Millen, 2009 Genetic variation and population substructure in outbred CD-1 mice: implications for genome-wide association studies. PLoS One 4**:** e4729.

Bult, C. J., J. A. Blake, C. L. Smith, J. A. Kadin, J. E. Richardson *et al.*, 2019 Mouse Genome Database (MGD) 2019. Nucleic Acids Res 47**:** D801-D806.

525 Cezard, T., F. Cunningham, S. E. Hunt, B. Koylass, N. Kumar *et al.*, 2022 The European
526   Variation Archive: a FAIR resource of genomic variation for all species. Nucleic Acids
527   Res 50**:** D1216-D1220.

528 Chang, P. L., E. Kopania, S. Keeble, B. A. J. Sarver, E. Larson *et al.*, 2017 Whole exome
529   sequencing of wild-derived inbred strains of mice improves power to link phenotype and
530   genotype. Mamm Genome 28**:** 416-425.

531 Chia, R., F. Achilli, M. F. Festing and E. M. Fisher, 2005 The origins and uses of mouse outbred
532   stocks. Nat Genet 37**:** 1181-1186.

533 Cingolani, P., A. Platts, L. Wang le, M. Coon, T. Nguyen *et al.*, 2012 A program for annotating
534   and predicting the effects of single nucleotide polymorphisms, SnpEff: SNPs in the
535   genome of Drosophila melanogaster strain w1118; iso-2; iso-3. Fly (Austin) 6**:** 80-92.

536 Cui, S., C. Chesson and R. Hope, 1993 Genetic variation within and between strains of outbred
537   Swiss mice. Lab Anim 27**:** 116-123.

538 DePristo, M. A., E. Banks, R. Poplin, K. V. Garimella, J. R. Maguire *et al.*, 2011 A framework
539   for variation discovery and genotyping using next-generation DNA sequencing data. Nat
540   Genet 43**:** 491-498.

541 Festing, M., K. Kondo, R. Loosli, S. M. Poiley and A. Spiegel, 1972 International standardized
542   nomenclature for outbred stocks of laboratory animals. Z Versuchstierkd 14**:** 215-224.

543 Foster, Y., L. Dutoit, S. Grosser, N. Dussex, B. J. Foster *et al.*, 2021 Genomic signatures of
544   inbreeding in a critically endangered parrot, the kakapo. G3 (Bethesda) 11.

545 Gatti, D. M., K. L. Svenson, A. Shabalin, L. Y. Wu, W. Valdar *et al.*, 2014 Quantitative trait
546   locus mapping methods for diversity outbred mice. G3 (Bethesda) 4**:** 1623-1633.

547 Groen, A., and A. J. Lagerwerf, 1979 Genic heterogeneity and genetic monitoring of mouse
548   outbred stocks. Lab Anim 13**:** 81-85.

549 Keane, T. M., L. Goodstadt, P. Danecek, M. A. White, K. Wong *et al.*, 2011 Mouse genomic
550   variation and its effect on phenotypes and gene regulation. Nature 477**:** 289-294.

551 Kinghorn, B., and A. Kinghorn, 2021 Management of inbreeding and coancestry to target short-
552   term and long-term genetic gains. Proc. Assoc. Advmt. Anim. Breed. Genet. 24**:** 191-194.

553 Lilue, J., A. G. Doran, I. T. Fiddes, M. Abrudan, J. Armstrong *et al.*, 2018 Sixteen diverse
554   laboratory mouse reference genomes define strain-specific haplotypes and novel
555   functional loci. Nat Genet 50**:** 1574-1583.

Logan, R. W., R. F. Robledo, J. M. Recla, V. M. Philip, J. A. Bubier *et al.*, 2013 High-precision
genetic mapping of behavioral traits in the diversity outbred mouse population. Genes
Brain Behav 12**:** 424-437.

McKenna, A., M. Hanna, E. Banks, A. Sivachenko, K. Cibulskis *et al.*, 2010 The Genome
Analysis Toolkit: a MapReduce framework for analyzing next-generation DNA
sequencing data. Genome Res 20**:** 1297-1303.

Morgan, A. P., C. P. Fu, C. Y. Kao, C. E. Welsh, J. P. Didion *et al.*, 2015 The Mouse Universal
Genotyping Array: From Substrains to Subspecies. G3 (Bethesda) 6**:** 263-279.

Mouse Genome Sequencing, C., R. H. Waterston, K. Lindblad-Toh, E. Birney, J. Rogers *et al.*,
2002 Initial sequencing and comparative analysis of the mouse genome. Nature 420**:** 520-
562.

Nicod, J., R. W. Davies, N. Cai, C. Hassett, L. Goodstadt *et al.*, 2016 Genome-wide association
of multiple complex traits in outbred mice by ultra-low-coverage sequencing. Nat Genet
48**:** 912-918.

Poiley, S. M., 1960 A systematic method of breeder rotation for non-inbred laboratory animal
colonies. Proc. Anim. Care Panel 10**:** 159-166.

Purcell, S., B. Neale, K. Todd-Brown, L. Thomas, M. A. Ferreira *et al.*, 2007 PLINK: a tool set
for whole-genome association and population-based linkage analyses. Am J Hum Genet
81**:** 559-575.

Recla, J. M., R. F. Robledo, D. M. Gatti, C. J. Bult, G. A. Churchill *et al.*, 2014 Precise genetic
mapping and integrative bioinformatics in Diversity Outbred mice reveals Hydin as a
novel pain gene. Mamm Genome 25**:** 211-222.

Rice, M. C., and S. J. O'Brien, 1980 Genetic variance of laboratory outbred Swiss mice. Nature
283**:** 157-161.

Saul, M. C., V. M. Philip, L. G. Reinholdt, A. Center for Systems Neurogenetics of and E. J.
Chesler, 2019 High-Diversity Mouse Populations for Complex Traits. Trends Genet 35**:**
501-514.

Sherman, B. T., M. Hao, J. Qiu, X. Jiao, M. W. Baseler *et al.*, 2022 DAVID: a web server for
functional enrichment analysis and functional annotation of gene lists (2021 update).
Nucleic Acids Res.

Sherry, S. T., M. H. Ward, M. Kholodov, J. Baker, L. Phan *et al.*, 2001 dbSNP: the NCBI
database of genetic variation. Nucleic Acids Res 29**:** 308-311.

Sigmon, J. S., M. W. Blanchard, R. S. Baric, T. A. Bell, J. Brennan *et al.*, 2020 Content and
Performance of the MiniMUGA Genotyping Array: A New Tool To Improve Rigor and
Reproducibility in Mouse Research. Genetics 216**:** 905-930.

Srivastava, A., A. P. Morgan, M. L. Najarian, V. K. Sarsani, J. S. Sigmon *et al.*, 2017 Genomes
of the Mouse Collaborative Cross. Genetics 206**:** 537-556.

Stoffel, M., M. Esser, M. Kardo, E. Humble, H. Nichols *et al.*, 2016 inbreedR: an R package for
the analysis of inbreeding based on genetic markers. Methods in Ecology and Evolution
7**:** 1331-1339.

Strobel, M. C., L. G. Reinholdt, R. D. Malcolm and K. Pritchett-Corning, 2015 Genetic
Monitoring of Laboratory Rats and Mice, pp. 1403-1416 in *Laboratory Animal Medicine,*
3rd Edition, edited by J. G. Fox, L. C. Anderson, G. Otto, K. R. Pritchett-Corning and M.
T. Whary. Elsevier, Boston.

Weichman, K., and J. R. Chaillet, 1997 Phenotypic variation in a genetically identical population
of mice. Mol Cell Biol 17**:** 5269-5274.

Wright, K. M., A. G. Deighan, A. Di Francesco, A. Freund, V. Jojic *et al.*, 2022 Age and diet
shape the genetic architecture of body weight in diversity outbred mice. Elife 11.

Yalcin, B., J. Nicod, A. Bhomra, S. Davidson, J. Cleak *et al.*, 2010 Commercially available
outbred mice for genome-wide association studies. PLoS Genet 6**:** e1001085.

Yalcin, B., K. Wong, A. Agam, M. Goodson, T. M. Keane *et al.*, 2011 Sequence-based
characterization of structural variation in the mouse genome. Nature 477**:** 326-329.

FIGURE 1

FIGURE 2

FIGURE 3

FIGURE 4

FIGURE 5

SUPPLEMENTARY FIGURE S1

A

B

SUPPLEMENTARY FIGURE S3

SUPPLEMENTARY FIGURE S4