Emergent dynamics of adult stem cell lineages from single nucleus and single cell RNA-

Seq of *Drosophila* testes

Amelie A. Raz[1#], Gabriela S. Vida[2#], Sarah R. Stern[3#], Sharvani Mahadevaraju[4#], Jaclyn M.

Fingerhut[1#], Jennifer M. Viveiros[5#], Soumitra Pal[6#], Jasmine R. Grey[5#], Mara R. Grace[5#],

Cameron W. Berry[3], Hongjie Li[7], Jasper Janssens[8], Wouter Saelens[9], Zhantao Shao[10], Chun

Hu[10], Yukiko M. Yamashita[1], Teresa M. Przytycka[6], Brian Oliver[4], Julie A. Brill[11,12,13], Henry M.

Krause[10,12], Erika L. Matunis[5], Helen White-Cooper[14], Stephen DiNardo[2*], and Margaret T.

Fuller[3,15*]

[1]Whitehead Institute for Biomedical Research and Department of Biology, Massachusetts

Institute of Technology; Howard Hughes Medical Institute, Cambridge, MA 02139, USA

[2]Department of Cell and Developmental Biology, The Perelman School of Medicine and The

Penn Institute for Regenerative Medicine, Philadelphia, PA 19104, USA.

[3]Department of Developmental Biology, Stanford University School of Medicine, Stanford, CA

94305 USA.

[4]National Institute of Diabetes and Digestive and Kidney Diseases, National Institutes of Health,

Bethesda, MD 20814, USA.

[5]Department of Cell Biology, Johns Hopkins University School of Medicine, 725 N. Wolfe Street,

Baltimore, MD 21205, USA.

[6]National Center for Biotechnology Information, National Library of Medicine, National Institutes

of Health, Bethesda, MD 20894, USA.

[7]Huffington Center on Aging and Department of Molecular and Human Genetics, Baylor College

of Medicine, Houston, TX 77030, USA.

[8]JVIB Center for Brain & Disease Research, and the Department of Human Genetics, KU

Leuven, Leuven, Belgium.

[9]Data Mining and Modeling for Biomedicine, VIB Center for Inflammation Research, and

Department of Applied Mathematics, Computer Science and Statistics, Ghent University, Ghent,

Belgium.

[10]Donnelly Centre for Cellular and Biomolecular Research, University of Toronto, Toronto,

Ontario M5S 3E1, Canada.

[11]Cell Biology Program, The Hospital for Sick Children, Toronto, Ontario M5G 0A4, Canada.

[12]Department of Molecular Genetics, University of Toronto, Toronto, Ontario M5S 1A8, Canada.

[13]Institute of Medical Science, University of Toronto, Toronto, Ontario M5S 1A8, Canada.

[14]School of Biosciences, Cardiff University, Museum Avenue, Cardiff, CF10 3AT, UK.

[15]Department of Genetics, Stanford University School of Medicine, Stanford, CA 94305 USA.

[*]Correspondence to:

sdinardo@pennmedicine.upenn.edu   ORC ID: 0000-0003-4131-5511

and

mtfuller@stanford.edu   ORC ID: 0000-0002-3804-4987

[#]'contributed equally' justification: AAR, GSV, SRS, SM, JMF, JMV, SP, JRG, and MRG each

took the lead on a particular aspect of the coding, analysis, documentation, construction of at

least one specific Figure and or its supplement, and/or assembly of supporting source data, and

also re-iteratively drafted sections of the narrative reporting that work.

Running title: adult fly testis atlas

Key words: stem cell / lineage / spermatogenesis / snRNA-seq / scRNA-seq

Abstract

Proper differentiation of sperm from germline stem cells, essential for production of the next

generation, requires dramatic changes in gene expression that drive remodeling of almost all

cellular components, from chromatin to organelles to cell shape itself. Here we provide a single

nucleus and single cell RNA-seq resource covering all of spermatogenesis in *Drosophila*

starting from in-depth analysis of adult testis single nucleus RNA-seq (snRNA-seq) data from

the Fly Cell Atlas (FCA) study (Li *et al.*, 2022). With over 44,000 nuclei and 6,000 cells

analyzed, the data provide identification of rare cell types, mapping of intermediate steps in

differentiation, and the potential to identify new factors impacting fertility or controlling

differentiation of germline and supporting somatic cells. We justify assignment of key germline

and somatic cell types using combinations of known markers, *in situ* hybridization, and analysis

of extant protein traps. Comparison of single cell and single nucleus datasets proved

particularly revealing of dynamic developmental transitions in germline differentiation. To

complement the web-based portals for data analysis hosted by the FCA, we provide datasets

compatible with commonly used software such as Seurat and Monocle. The foundation

provided here will enable communities studying spermatogenesis to interrogate the datasets to

identify candidate genes to test for function *in vivo.*

Introduction

Single cell RNA-seq (scRNA-seq) of developing tissues can reveal new cell types as

well as previously unknown steps in the differentiation of lineages underlying tissue

homeostasis and repair. In fact, high-resolution expression maps are being created for entire

organisms, from *C. elegans*, planaria, and schistosomes to *Drosophila* and mouse (Cao et al.,

2017, 2019; Fincher et al., 2018; Li et al., 2022; Plass et al., 2018; Schaum et al., 2018; Sebé-

Pedrós et al., 2018; Siebert et al., 2019; Wendt et al., 2020), with such atlases providing a

foundational reference for several important model organisms. In particular, for tissues

maintained by stem cell lineages, scRNA-seq can identify the developmental trajectories that

lead from dedicated tissue stem cell to terminally differentiated cell types, an important resource

for understanding tissue maintenance, repair, and the origins of cancer.

The testis harbors a highly active, unipotent adult stem cell lineage that must produce

sperm throughout reproductive life. Spermatogenesis relies on self-renewing germline stem

cells, the progeny of which differentiate into one of the most highly specialized cell types in the

body. Production of functional sperm requires intimate interactions between germ cells and

somatic support cells, with defects at almost any step compromising fertility. Interest in

spermatogenesis has motivated scRNA-seq analyses of testes from a variety of organisms,

including mouse (Cao et al., 2021; Chen et al., 2018; Green et al., 2018; Guo et al., 2020; Law

et al., 2019) and *Drosophila* (Li et al., 2022; Mahadevaraju et al., 2021; Witt et al., 2019).

Notably, the testis of *Drosophila* has the highest complexity in terms of mRNAs expressed of

any tissue in the fly, likely reflecting the dramatic differentiation events required (Li et al., 2022).

Many aspects of spermatogenesis are conserved from *Drosophila* to mammals. One

striking difference, however, is that spermatogenesis in *Drosophila* relies on not one but two

adult stem cell lineages. The co-differentiating germ cells and their closely associated somatic

support cells descend from distinct stem cell populations, housed together in a well-defined

niche (Fuller, 1998). Additionally, the many mutations affecting male fertility, plus powerful

genetic tools for cell type specific functional analysis, have allowed identification of stage-

specific regulatory factors underlying niche function in stem cell maintenance, control of

proliferation, and soma-germline feedback circuits that act during co-differentiation of these two

lineages. This comprehensive knowledge of spermatogenesis offers a rich biological foundation

for interpreting single nucleus and single cell RNA-seq data.

Here we present an in-depth analysis of the testis subset of the Fly Cell Atlas (FCA)

single nucleus RNA-Seq (snRNA-seq) data. We supplement this with scRNA-seq from the

same tissue, together providing a foundational reference for the field. While several recent

RNA-seq analyses of *Drosophila* testes have been illuminating, they generally focused on

particular stages (Gan et al., 2010; Hof-Michel and Bökel, 2020; Lu et al., 2020; Mahadevaraju

et al., 2021; Shi et al., 2020; Vedelek et al., 2018; Witt et al., 2019). In contrast, the scale and

comprehensive nature of the FCA dataset allowed us to profile rare cell types, such as the stem

cell niche, and to follow spermatogenesis from early spermatogonia to late spermatids, a

remarkable conversion of precursors to highly elongated, specialized cells.   We present

supporting data for assignment of key cell types, both germline and somatic, and show how

progression through two distinct, yet intimately interacting, stem cell-based lineages emerges

from the changes in gene expression.

The data confirm and extend known features of the male germ line transcription

program, including cell type specific expression of many genes in spermatocytes,

downregulation of X-linked genes in later spermatocytes, and repression of most transcription in

early spermatids. At the same time, surprising new features emerged, including unexpected

complexity in the somatic support cell lineage. In addition, comparison of single nucleus with

single cell sequencing data significantly expanded our understanding of gene expression

dynamics in spermatocytes and spermatids as they mature. In particular, these data showed

how dynamic changes in active transcription reflected in the snRNA-seq can be obscured by the

endowment of mRNAs stored in the cytoplasm. This is especially clear in early haploid

spermatids, which have little transcriptional activity but contain many mRNAs transcribed in

spermatocytes and stored to be translated later to support spermatid morphogenesis. With a

gene expression framework for the two testis stem cell lineages now in place, mining the

snRNA-seq data for changes in gene expression as one cluster advances to the next should

identify new sub-stage-specific markers, thereby opening the way for tests of function for such

newly identified genes in male germ cell differentiation.

Results

Clustering by gene expression signature reveals progression of differentiation in two

stem cell lineages

Spermatogenesis in *Drosophila* involves obligate, intimate interactions between cells

differentiating in two adult stem cell-founded lineages. Male germline stem cells (GSCs) and

their partners, the somatic cyst stem cells (CySCs), are both physically anchored to a small

cluster of somatic cells termed the apical hub (Figure 1A), which provides short-range niche

signals important for maintenance of the two stem cell states. The interleaved arrangement of

GSCs and CySCs ensures that their immediate daughters are positioned to interact. Two

postmitotic early cyst cells enclose each gonialblast (immediate GSC daughter), forming a two-

cell squamous epithelium that later seals the progeny of the gonialblast off from the rest of the

testis (Fairchild et al., 2015). The gonialblast initiates four rounds of spermatogonial transit

amplifying divisions with incomplete cytokinesis, producing 16 interconnected germ cells (Figure

1A). After the fourth mitosis, the germ cells undergo premeiotic S-phase and enter an extended

G2 cell cycle phase termed meiotic prophase. Over the next three and a half days the 16

primary spermatocytes increase 25-fold in volume, engaging in a robust transcription program in

preparation for the meiotic divisions and the extensive elongation and remodeling of the

resulting 64 haploid spermatids into mature sperm. Although they do not divide, the two

somatic cyst cells co-differentiate with the germ cells they enclose (Gönczy et al., 1992),

eventually taking on different identities as head and tail cyst cells. The head cyst cell cups the

nuclear end of elongating spermatid bundles and eventually inserts into the terminal epithelium

at the base of the testis, while the tail cyst cell elongates extensively to cover the rest of the

spermatid bundle (Tokuyasu et al., 1972). All these cell types, as well as somatic structural cells

of the testis sheath (muscle and pigment cells) and cells of the seminal vesicle are represented

in the FCA testis dataset.

The relative similarity and differences in gene expression for 44,621 single nuclei from

triplicate 10X snRNA-seq runs from adult testis plus seminal vesicle (See Materials and

Methods) can be visualized in a Uniform Manifold Approximation and Projection (UMAP)-based

dimensionality reduction plot (Figure 1B). The geography of the UMAP is dominated by the

dynamic sequences of differentiating states in the germline (blue) and somatic cyst cell (yellow)

lineages. Each lineage manifests as an emergent trajectory of nuclei with continuously

progressing gene expression profiles, unlike the discrete clusters characteristic of most

terminally differentiated cell types. Despite their physical proximity and cooperation *in vivo* (blue

and yellow in Figure 1A), the germ line and cyst cell lineages mapped to largely non-overlapping

formations in gene expression space represented in the UMAP (blue and yellow in Figure 1B),

consistent with their different embryological origin, cell biology, and known roles.

From the perspective in Figure 1B, the spatial arrangement of nuclei in the UMAP

whimsically resembles a hammerhead shark (blue - germ line) playing a saxophone (yellow -

cyst cell lineage) watched over by a mermaid (several somatic epithelial-based structural

elements, including the seminal vesicle (sv, dark gray) and terminal epithelial cells at the testis

base (te, pink)). One notable cluster located near the mermaid head is the hub (hb, light green),

the niche that supports the two stem cell lineages. Other clusters on the UMAP contain

differentiated cell types that contribute to organ structure, including muscle (mc) and pigment

cells (pc) of the testis sheath (Figure 1B). Additionally, sample dissection carried over small

numbers of non-testis cells, including tracheal (tr) and fat body (ft) cells, hemocytes (hm),

neurons (nr), and male reproductive tract secretory cells (sr).

Identity of key clusters was assigned based on expression of known markers from the

literature (citations for all published markers employed given in Table 1). Expression of *vasa*

(vas) identified early germ line nuclei while expression of *traffic jam (tj)* identified nuclei from

early stages in the somatic cyst cell lineage (Figure 1C,D). A small proportion of nuclei in other

tissues reported *vas* expression at a low level, perhaps due to ambient RNA in the nuclear

preparation resulting from lysed cells rather than reflecting *vas* expression outside the germ line

(Figure 1 - figure supplement 1A). However, it is also worth noting that Vasa has been shown to

be expressed in somatic cells of the gonad (Renault, 2012). Expression of the cdc25

phosphatase *string (stg)*, required for the G2/M transition in mitotic cells (Alphey et al., 1992;

Edgar and O'Farrell, 1990), and *escargot (esg)*, a gene expressed in diploid proliferative cells

(Fuse et al., 1994), marked CySCs in the cyst cell lineage and proliferating GSCs and

spermatogonia in the germ line lineage (Figure 1E,F). Expression of *esg* also marked the hub,

as expected from prior studies (Voog et al., 2014) (Figure 1F). Together, these markers

established that the germ line lineage begins at the tail end of the "shark" with germ line stem

cells (GSCs) and proliferating spermatogonia at the tapered point. The somatic cyst cell lineage

begins at the mouthpiece of the "saxophone" at the UMAP center, with early cyst cell nuclei

extending down and leftward in a thin line. In addition, analysis by fluorescent *in situ*

hybridization (FISH), and the average number of unique transcripts (Unique Molecular Identifier

- UMI) expressed helped assign identity. For example, spermatocytes are highly

transcriptionally active, whereas *Drosophila* early spermatids are nearly quiescent. While

clustering was carried out using the Leiden algorithm at increasing levels of resolution, we

settled on Leiden 6.0 as providing optimal granularity along both somatic and germline

differentiation trajectories. We assigned 43 clusters as germline and 22 clusters as likely cyst

cell lineage, with many inferred from the UMAP geography as representing putative

intermediate cell types in the respective lineages (Figures 1G, 2A, 6A, and Figure 2 – figure

supplement 1A).

Table 1: List of identifying genes

Gene symbol	Gene_name	FBgn	Reference	DOI
aly	always early	FBgn0004372	White-Cooper et al., 2000	10.1242/dev.127.24.5463
aub	aubergine	FBgn0000146	Nishida et al., 2007	10.1261/rna.744307
bam	bag of marbles	FBgn0000158	Schulz et al., 2004	10.1534/genetics.103.023184
CadN	Cadherin-N	FBgn0015609	Boyle et al., 2007	10.1016/j.stem.2007.08.002
can	cannonball	FBgn0011569	Hiller et al., 2001	10.1101/gad.869101
cher	cheerio	FBgn0014141	Tanentzapf et al., 2007	10.1038/ncb1660
CycB	Cyclin B	FBgn0000405	White-Cooper et al., 1998	10.1242/dev.125.1.125
Dic61B	Dynein intermediate chain at 61B	FBgn0263988	Lu et al., 2020	10.1101/gad.335331.119
dlg1	discs large 1	FBgn0001624	Papagianoulli and Mechler, 2009	10.1038/cr.2009.71
esg	escargot	FBgn0287768	Kiger et al., 2000	10.1038/35037606
eya	eyes absent	FBgn0000320	Fabrizio et al., 2003	10.1016/s0012-1606(03)00127-1
f-cup	flyers-cup	FBgn0028487	Barreau et al., 2008	10.1242/dev.021949
Fas3	Fasciclin III	FBgn0000636	Brower et al., 1981	10.1242/dev.63.1.233
fzo	fuzzy onions	FBgn0011596	Hwa et al., 2002	10.1016/s0925-4773(02)00141-7
hh	hedgehog	FBgn0004644	Michel et al., 2012	10.1242/dev.075242
Hml	Hemolectin	FBgn0029167	Li et al., 2022	10.1126/science.abk2432
kl-2	male fertility factor kl2	FBgn0001313	Carvalho et al, 2000	10.1073/pnas.230438397
kl-3	male fertility factor kl3	FBgn0267432	Carvalho et al, 2001	10.1073/pnas.230438398
kl-3	male fertility factor kl3	FBgn0267432	Fingerhut et al., 2019	10.1371/journal.pgen.1008028
kl-5	male fertility factor kl5	FBgn0267433	Gepner and Hays, 1993	10.1073/pnas.90.23.11132
kl-5	male fertility factor kl5	FBgn0267433	Fingerhut et al., 2019	10.1371/journal.pgen.1008028
kmg	kumgang	FBgn0032473	Kim et al., 2017	10.1126/science.aal3096
Mst77F	Male-specific transcript 77F	FBgn0086915	Barckmann et al., 2013	10.1016/j.ydbio.2013.02.018
Mst84Db	Male-specific RNA 84Db	FBgn0004173	Kuhn et al., 1991	10.1016/0925-4773(91)90064-d
Mst84Dc	Male-specific RNA 84Dc	FBgn0004174	Kuhn et al., 1991	10.1016/0925-4773(91)90064-d
Mst87F	Male-specific RNA 87F	FBgn0002862	Kuhn et al., 1991	10.1016/0925-4773(91)90064-d
MtnA	Metallothionein A	FBgn0002868	Zhao et al., 2010	10.1093/nar/gkp1006

Nep5	Neprilysin 5	FBgn0039478	Sitnik et al., 2014	10.1534/genetics.113.160945
p53	p53	FBgn0039044	Monk et al., 2012	10.1007/s00441-012-1479-4
p-cup	presidents-cup	FBgn0030840	Barreau et al., 2008	10.1242/dev.021949
piwi	P-element induced wimpy testis	FBgn0004872	Gonzalez et al., 2015	10.1016/j.celrep.2015.06.004
Rbp4	RNA-binding protein 4	FBgn0010258	Baker et al., 2015	10.1242/dev.122341
sa	spermatocyte arrest	FBgn0002842	Hiller et al., 2004	10.1242/dev.01314
shg	shotgun	FBgn0003391	Voog et al., 2008	10.1038/nature07173
so	sine oculis	FBgn0003460	Fabrizio et al., 2003	10.1016/s0012-1606(03)00127-1
soti	scotti	FBgn0038225	Barreau et al., 2008	10.1242/dev.021949
stg	string	FBgn0003525	Alphey et al., 1992	10.1016/0092-8674(92)90616-K
Syt1	synaptotagmin 1	FBgn0004242	Li et al., 2022	10.1126/science.abk2432
tj	traffic jam	FBgn0000964	Li et al., 2003	10.1038/ncb1058
tomboy20	tomboy20	FBgn0037828	Hwa et al., 2004	10.1016/j.febslet.2004.07.025
upd1	unpaired 1	FBgn0004956	Tulina and Matunis 2001	10.1126/science.1066700
vas	vasa	FBgn0283442	Hay et al., 1988	10.1016/0092-8674(88)90216-4
wa-cup	walker cup	FBgn0037502	Barreau et al., 2008	10.1242/dev.021949
zpg	zero population growth	FBgn0024177	Tazuke et al., 2002	10.1242/dev.129.10.2529

Progression of differentiation in the male germ line stem cell lineage

Figure 2A shows the UMAP for the germ line stem cell lineage with Leiden 6.0 clusters labeled. Expression of the germ cell-specific gap junction gene *zero population growth* (*zpg),* required for survival of early spermatogonia (Tazuke et al., 2002) (Figure 2D, L), along with *vasa, stg* and *esg* (Figure 1C, E and F), further established nuclei at the pointed tip of the shark tail (clusters 25 and 22) as GSCs and spermatogonia. *In vivo*, GSCs are distinguished from gonialblasts and transit amplifying spermatogonia cytologically, by attachment to the apical hub and cell biological characteristics such as oriented centrosomes and spindles, and functionally, by lineage analysis. However, mRNA markers restricted to GSCs have not yet been identified, preventing us from determining what percent of these early nuclei are GSCs. Many nuclei in cluster 22 express *bag-of-marbles (bam)* (Figure 2L) but lack known spermatocyte markers, suggesting that these nuclei represent mid-to-late spermatogonia or germ cells undergoing premeiotic S phase. Moving rightward, several known early spermatocyte markers such as *kumgang* (*kmg)* and *RNA-binding protein 4* (*Rbp4*) began to be expressed (clusters 5, 78, and 40) (Figure 2F-G, L). Transcripts from *aubergine* (*aub),* a piRNA binding protein, were detected in the spermatogonial region (clusters 25, 22) and overlapping with early spermatocyte markers (clusters 5, 78, 40 and 41) (Figure 2E and L). Fluorescent *in situ* hybridization (FISH) confirmed *aub* transcripts present in GSCs around the hub, spermatogonia, and extending into early spermatocyte cysts, with their characteristic larger nuclei (Figure 2B). FISH also confirmed expression of *kmg* mRNA starting in early spermatocytes (Figure 2C), with early spermatocytes showing both *aub* and *kmg* transcripts, consistent with the snRNA-seq data.

Progressively maturing spermatocytes along the bottom right of the germline UMAP expressed later markers, including mRNAs for the spermatocyte-specific tMAC subunit *always early* (*aly)* and the testis-specific TAFs (tTAFs) *spermatocyte arrest (sa)* and *cannonball* (*can*) (clusters 41, 51, 35, and onward; Figure 2H,L). Expression of *fuzzy onions* (*fzo)* and *Dynein intermediate chain 61B* (*Dic61B*) was detected later, as the germ cell clusters curved upward (clusters 33, 48,105, 45, 13, 56; Figure 2J,L), consistent with the dependence of *fzo* and *Dic61B* transcription on *aly* (Hwa et al., 2002; Lu et al., 2020). Correlating *in vivo* morphology with gene

expression space (visualized in the UMAP), *fzo* transcripts were not detected by FISH in the

young spermatocytes near the spermatogonial region but were strongly detected in more

mature spermatocytes further away from the testis apical tip (Figure 2B). The G2/M cell cycle

regulator *CyclinB* (*CycB*) is transcribed from one promoter in mitotic spermatogonia, silenced,

then re-expressed from an alternate promoter in later spermatocytes, dependent on *aly* function

(Lu et al., 2020; White-Cooper et al., 1998). These two distinct stages of *CycB* transcript

expression are clearly visible in the snRNA-seq data (Figure 2I,L). Maturing spermatocytes,

marked by expression of Y-linked genes (Figure 3D), lie toward the top of the upward curve

where the tail meets the torso of the shark. The progression of germ cell differentiation

continues with early stage spermatids along the upper torso and head of the shark (marked by

low UMI - see below). Mid-to-late elongation stage spermatids, marked by expression of *p-cup*

mRNA, lie in the blunt projection toward the upper right of the UMAP (clusters 66, 10, 46; Figure

2K,L).

The order of clusters in expression space reflects differentiation in the lineage, as

indicated by plotting the expression of known germline markers in each UMAP cluster (Figure

2L). Notably, using the published marker genes scored here, sequential cluster identities (e.g.,

25, 22, 5, 78, 40) were not each delimited by unique marker genes. Instead, graded expression

of the markers examined extended across boundaries between clusters, as expected for a

continuous differentiation process. This was also observed in a UMAP with just nine clusters

created at lower resolution of Leiden 0.8 (Figure 2 – figure supplement 1B,C). In principle, there

may also be genes where expression is delimited by cluster boundaries, and these may identify

novel markers demarcating germline transition stages. Four genes were selected to test using

RNA FISH as potentially meeting this criterion. Two represented the earliest cluster (GSCs and

early spermatogonia, *Pxt* and *Drep2*) and two represented a mid-trajectory cluster (mid-

spermatocytes, *CG43317* and *Vha16-5*). As predicted, *Pxt* and *Drep2 expression was*

restricted to early germine stages, while *CG43317* and *Vha16-5* were robustly expressed in

spermatocytes and not detectable at early stages (Figure 2 – figure supplement 1 D-E, genes

marked with solid squares in Figure 4G-H). The results validate use of these genes as new

markers for their respective stages, and demonstrate the general utility of the dataset for

discovery of novel markers.

The geography of the UMAP is reminiscent of the spatio-temporal organization in the

testis itself, with stages laid out from GSCs through transit amplifying spermatogonia, and then

through young, mid, and late spermatocyte stages. However, a UMAP displays changes in

gene expression space rather than physical space. As a consequence, some surprisingly long

stretches of the germline testis UMAP represent what are known to be short periods in

developmental time. For example, the short, young spermatocyte stage is represented by a

long stretch along the UMAP (clusters 5, 78, 40, 41, 51). This is underscored by the long gap in

detection of *CycB* mRNA representing these clusters in the UMAP (Figure 2I, L). However, in

terms of physical space, *in situ* hybridization for *CycB* revealed only a relatively narrow gap near

the boundary between spermatogonia and spermatocytes (White-Cooper et al., 1998).   The

UMAP territory for young spermatocytes may be stretched in gene expression space because

this stage is a time of extensive and dynamic changes in gene expression, with many genes

dramatically upregulated as the spermatocyte expression program initiates, and many genes

downregulated from the spermatogonial phase (Figure 2,L; see also Shi et al., 2020).

Meanwhile, the young spermatocytes do not take as much physical space as the more mature

stages, since they are smaller in size and fewer in number (2462 nuclei in clusters 5, 78, 40, 41,

51, compared to ~4100 nuclei in clusters 35, 33, 48, 105, 64, 45, 13, 56,16).

The spermatocyte transcription program

The spermatocyte period features onset of dramatic transcriptional changes. Many

genes expressed in spermatocytes are transcribed in few or no other known cell types, including

the markers *kmg, Rbp4, fzo, can, sa* (see references in Table 1).   This robust onset of cell type-

specific transcription appears as an increase in the number of different genes detected per

nucleus (Figure 3A,C), leading to a substantial increase in transcriptome complexity. Coincident

with this was a large increase in the number of unique molecular identifiers (UMI) scored,

peaking in mid-to-late spermatocyte nuclei (clusters 35 and 33), with average UMI per cluster

increasing from <5000 to >30,000 UMI per nucleus as spermatogonia differentiated to late

spermatocytes (Figure 3B,C). These observed increases in the number of genes and UMIs

detected were independently observed in a single cell testis dataset to be introduced below

(Figure 3 – figure supplement 1B,C). The FCA paper noted that testis, heart, fat body,

Malpighian tubules, and male reproductive glands had relatively high RNA levels and number of

genes expressed compared to other tissues (Li et al., 2022). Reanalysis showed that mid-to-

late spermatocyte nuclei exhibited the highest complexity of all, with average expressed gene

(6,000 compared to 2,000) and UMI (30,000 compared to <20,000) numbers higher than for any

cluster in the tissues noted by the FCA paper (Figure 3 - figure supplement 1A;). High

transcriptome complexity has also been noted in mammalian spermatocytes (Soumillon et al.,

2013).

After peaking in clusters 35 and 33, UMI values per nucleus decreased through clusters

48, 105, 56, 106, 21, where the "tail" meets the "torso" of the shark (Figure 3B,C - See Figure

2A for positions of numbered clusters), consistent with the observed lower expression of

spermatocyte marker genes (Figure 2L). In the shark's upper torso and head, many clusters

had very low UMI (Figure 3C), making developmental order difficult to assign. This is reflected

in the UMAP shape, with clusters grouped rather than extended along a string as in early germ

cell stages. We surmise these nuclei represent early spermatids, as classic studies showed

that transcription falls dramatically from shortly before onset of the meiotic divisions, with no

bulk incorporation of radioactive uridine detected in haploid round and early elongating

spermatid nuclei (Gould-Somero and Holland, 1974; Olivieri and Olivieri, 1965). Although UMI

counts and average number of genes expressed in the snRNA-seq were low in post-meiotic

clusters (Figure 3C), nonetheless, post-meiotic transcription appeared more extensive than

previously appreciated, with transcripts from approximately 1000 genes detected.

Spermatocytes showed sex chromosome specific trends in gene expression changes.

Overall, Y-linked transcripts were strongly upregulated in spermatocytes (Figure 3D), primarily

driven by the robust expression of 8 of the 12 single copy genes. For example, transcription of

the Y-linked fertility factors *kl-3* and *kl-5*, which encode flagellar dyneins expressed only in male

germ cells, was massively upregulated (125 and 275 fold respectively, with similarly large

differences in absolute expression). As 10X sequencing utilizes oligo(dT) primers, the late

appearance of reads from the Y-linked fertility factors in spermatocytes may reflect the very long

time required to complete synthesis of the mature transcripts, which have extremely large

introns (Fingerhut et al., 2019).   For X-linked genes, analysis of the snRNA-seq data showed a

similar level of expression relative to a control set of genes from all chromosomes in

spermatogonia and early spermatocytes. However mid-to-late spermatocytes featured a

transition to reduced expression of X-linked genes relative to the control set (Figure 3E),

consistent with the roughly 2 fold lower expression of X linked genes compared to generally

expressed autosomal genes observed previously (Mahadevaraju et al., 2021).

One surprise that emerged from the UMAP geography was that later stage

spermatocytes split into three parallel streams, all expressing spermatocyte specific markers.

(Figure 2A). Strikingly, nuclei in the leftmost and middle streams (clusters 64 and 45,

respectively) had considerably lower UMI count than in the robust mainstream (cluster 35;

Figure 3B,C). The cause underlying such different UMI levels among late spermatocytes is not

known, but could suggest a stochastic component to meiotic chromosome condensation and the

attendant chromosome-wide downregulation in gene expression.

A second notable feature was the expression in mid-to-late spermatocytes of many

markers classically associated with other cell types. Notably, markers for hemocytes

(*Hemolectin – Hml*), muscle *(Myosin heavy chain - Mhc)*, neurons (*Synaptotagmin - Syt1*), and

epithelial cells (*grainy head - grh*) selected as identifiers of these cell types in the FCA study of

adult *Drosophila* tissues (Li et al., 2022), were upregulated in late spermatocytes (Figure 3F,G,I,

Figure 3 – figure supplement 2A,B). Similar upregulation of *Mhc*, *Hml*, *grh,* and *Syt1* in

spermatocytes was independently observed in a single cell testis dataset introduced below

(Figure 3 - figure supplement 2E), so is not likely to be an artifact of isolation of nuclei. Other

mRNAs normally thought of as markers of somatic cells were revealed to be upregulated in mid-

to-late spermatocytes in the snRNA-seq dataset, such as *eyes absent* (*eya*) and *unpaired 1*

(*upd*; Figure 3 - figure supplement 2C,D).

Expression in spermatocytes was confirmed by in situ analysis for *Hml* in wild-type

testes and under mutant conditions where spermatocytes accumulate (Figure 3H, Figure 3 -

figure supplement 2F). Additionally, the mRNAs for the late cyst cell markers *geko* and *Visual*

system homeobox 1 (*Vsx1*) and the cyst cell marker *eya* were directly detected in

spermatocytes by FISH (Figure 3 - figure supplement 2G, H, I; and see below)**.** Note that

expression of the somatic marker mRNAs was generally lower in spermatocytes than in the

marker somatic tissue (Figure 3I), as observed in *Vsx1* in situs, for example (Figure 3 – figure

supplement 2H,I). It will be fruitful to investigate the cause of the seemingly promiscuous

expression in spermatocytes of certain somatic marker genes, whether their encoded proteins

accumulate, and what role these genes may have in spermatocyte biology.

Although spermatocytes feature an overall increase in UMIs per cell, spermatocytes do

not appear to simply be 'permissive' for general expression. For example, mRNA for the cyst

marker *tj* was not detected in spermatocytes (Figure 1D). This selectivity argues against the

possibility that observing 'somatic markers' in spermatocytes represents some technical artifact.

In addition, some transcription factors that mark specific somatic tissues were detected as

upregulated in spermatocytes compared to spermatogonia, while others were not. Across all

adult fly tissues, the FCA project identified 496 transcription factors predicted to have a high

tissue specificity score (Li et al., 2022). That analysis predicted 351 of these as expressed in

certain somatic cell types but not in germ line (Table S3 from Li et al., 2022). Our analysis of

the testis plus seminal vesicle portion of the FCA data showed that about a fifth (76) of these

'somatic' factors were upregulated in spermatocytes compared to spermatogonia, but most

(275) exhibited no upregulation (Figure 3 - source data 1).

sn vs. scRNA-seq: dynamics of active transcription vs. stored RNAs

One of the regulatory mechanisms used to control cell differentiation in tissues is the

programmed storage of mRNAs to be used later, long after the genes encoding these mRNAs

are silenced. Spermatogenesis in *Drosophila* is especially conducive to the study of this

phenomena because early spermatids, although largely transcriptionally silent, carry numerous

cytoplasmic transcripts, many of which are recruited to be translated for temporal control of

protein expression during spermatid morphogenesis (Schäfer et al., 1990, 1995). To identify

transcripts under such control, we took advantage of the fact that single nucleus RNA

sequencing detects recently transcribed or nuclear resident transcripts, whereas single cell RNA

sequencing (scRNA-seq) samples a largely different RNA population. Therefore, we generated

single cell RNA-seq of adult testes (without seminal vesicle), producing data for 6438 germ cells

after quality control steps (Materials and Methods).

The UMAP geography for both the snRNA-seq and scRNA-seq datasets showed progression

from spermatogonia to spermatids, with germline differentiation classes present in sequential

order (Figure 4A-D, Figure 4 - figure supplement 1). In the scRNA-seq UMAP, as for snRNA-

seq, expression of *zpg* marked a small number of spermatogonia located at the bottom tip, *aub*

marked those same plus additional cells, presumably early spermatocytes, *fzo* marked largely

differentiating spermatocytes with expression abruptly ending in early spermatids (yellow

arrowheads in Figure 4C,E), and *presidents-cup* (*p-cup*) marked later elongating spermatids in

an arm extending from the top of the UMAP (Figure 4E). Corroborating these expression

patterns, FISH to whole mount testes clearly showed *aub* expression in spermatogonia and

early spermatocytes, and *fzo* expression beginning in spermatocytes (Figures 2B,4F). Notably,

fzo mRNA was abruptly downregulated in early round spermatids soon after the second meiotic

division (Figure 4F, yellow arrowhead).

Trajectory inference can assign a differentiation distance parameter to cells inferred from

transcriptional differences, with distance noted as 'pseudotime' (Trapnell et al., 2014). Applying

trajectory inference independently to the snRNA-seq and scRNA-seq germ line datasets

produced contiguous trajectories. Using Monocle3, 99.9% (21,061/21,091) of the snRNA-seq

germline nuclei were connected (Figure 4B). Notably, unlike prior trajectory analysis using

Slingshot (Li et al., 2022), the inferred trajectory was contiguous, connecting cells of all

differentiation points from early spermatogonia to late spermatids. Likewise, Monocle3 analysis

of the 6438 germline cells from the scRNA-seq also produced a contiguous trajectory from

spermatogonia to elongating spermatids (Figure 4D), although it did include a late bifurcation,

the explanation of which may be technical or biological. For both datasets, pseudotime staging

paralleled the ordered trajectory deduced from marker gene expression in UMAP clusters.

(Figure 2, Figure 4A-E).

Plotting normalized gene expression across pseudotime in the sn- and scRNA-seq

datasets revealed both shared and contrasting dynamics. In both datasets, the same set of

genes were expressed in early germ cells, with expression diminishing over pseudotime (Fig

4G,H; red/orange in the top color bar, ending at the point indicated by an arrow, corresponding

to the colors and arrow in Figure 4A,C). Similarly, some genes, including *fzo* and *CycB* (Figure

4G,H - black arrowheads on side) reached peak expression in spermatocytes and dropped to

low levels by early spermatid stages in the scRNA-seq (Figure 4E,H, white vertical line marking

fzo drop to z-score 0), consistent with *in situ* hybridization data (Figure 4F and published in

White-Cooper et al., 1998, respectively).

Interestingly, in the snRNA-seq dataset, a group of over 200 genes in the middle region

of the heat map reached peak expression in the mid-spermatocyte stages (green-aqua hues in

the top color bar) then dropped in expression, falling halfway to their nadir at a point similar to

the drop in *fzo* and *cycB* expression (vertical white line) (Figure 4G). In the scRNA-seq dataset,

however, these same genes were still at or near peak mRNA accumulation in the same cells in

which expression of *fzo* and *cycB* had already dropped (Figure 4H, white line). Thus, it is the

comparison of the datasets that is most revealing: these genes are transcribed in

spermatocytes, then transcription halts (inferred from the snRNA-seq dataset), but their mRNAs

remain high in spermatids (inferred from the scRNA-seq dataset) well past the stage when *fzo*

and *cycB* mRNAs disappear (inferred from both datasets). Several genes from this set,

including *Male-specific transcript 77F* (*Mst77F*) and *tomboy20* (asterisks), have been previously

demonstrated by *in situ* hybridization to maintain abundant transcripts in both spermatocytes

and elongating spermatids (Barckmann et al., 2013; Hwa et al., 2004). Comparison of the

snRNA-seq and scRNA-seq datasets suggests that such a pattern, previously described for a

small number of transcripts, is shared by hundreds of genes.

We chose one such novel predicted perdurant transcript, *l(2)41Ab* (Figure 4G-H, marked

with a circle), proposed to be involved in axoneme assembly in spermatids (Zur Lage et al.,

2019), to validate *in vivo*. Signal from RNA FISH probes designed against a *l(2)41Ab* intron was

clearly present in nuclear foci in spermatocytes, but absent from spermatid nuclei, suggesting

that active *l(2)41Ab* transcription occurs in meiotic prophase (G2) and has ended by the early

round spermatid stage (Figure 4I). Signal from probes designed against a *l(2)41Ab* exon, by

contrast, was visible in the cytoplasm of spermatocytes and even more strongly present in early

spermatids and the tails of elongating spermatids (Figure 4I). Such a pattern suggests that

comparison of snRNA-seq and scRNA-seq datasets can correctly uncover novel genes that

encode transcripts that remain in the cytoplasm for days after transcription has ceased.

Comparison of single cell and single nucleus data revealed several distinct classes of

transcript behaviors in spermatids, each worthy of targeted follow-up study. Notably, perduring

transcripts from the class of genes described above showed two types of behavior. For over a

hundred genes, transcripts disappeared sharply in later elongating spermatids (7 genes with this

pattern highlighted in pink box in Figure 4G,H). To plot transcript levels in the sn- and scRNA-

seq datasets on the same X axes, a common 'warped' time scale was derived for the datasets

(see Materials and Methods). First, for all genes upregulated in spermatocytes, onset of

transcription in the nuclear transcriptome was followed (with delay) by upregulation in the whole-

cell transcriptome, generally dominated by cytoplasmic transcripts (Figure 4I,J). As seen in the

heat map (Figure 4H), the graphs for the pink box genes show that the mRNAs remained at

peak levels considerably later in the scRNA-seq than in the snRNA-seq data, but the transcripts

were eventually strongly downregulated by late spermatid stages (Figure 4J: blue lines). This

suggests complexity in mRNA regulation in the cytoplasm: stable maintenance in early

spermatids and abrupt degradation in later spermatids, perhaps once transcripts have been

translated. Interestingly, the protein products of several of these genes are present and

functional in late spermatids and sperm (Raja and Renkawitz-Pohl, 2006), suggesting these

proteins are actively maintained in the absence of new translation.

A second type of behavior was noted for a much smaller group of genes (green box in

Figure 4G,H, graphed in K) where transcripts perdured even longer, remaining high through the

latest stages assessed by scRNA-seq (blue lines in Figure 4K). As differentiation of late

spermatocytes to late spermatids takes days (Chandley and Bateman, 1962), these remarkable

transcripts maintained high levels of cytoplasmic abundance, with almost no sign of

degradation, even days after active transcription had dropped off. This suggests exceptional

stability, likely provided by specialized RNA-binding proteins. Some such transcripts, encoded

by *Mst84Db*, *Mst84Dc*, and *Mst87F*, have long been recognized to be translationally regulated,

with perdurance in the cytoplasm for up to three days (Kuhn et al., 1991; White-Cooper et al.,

1998). Others, including *CG30430, CG9016, MCU*, and *CG17666* have not been previously

reported to undergo translational regulation. The differences in degradation timing revealed by

scRNA-seq (Figure 4G-H, pink vs green boxes) may hint that distinct groups of RNAs, and thus

their protein products, are engaged at different stages of spermatid morphogenesis.

Another compelling example of the utility of comparing snRNA-seq and scRNA-seq data

is highlighted by the group of genes outlined in black (Figure 4G,H). In the snRNA-seq dataset,

these genes are expressed in spermatocytes but transcription shuts down in early spermatids

and remains off for a considerable period before expression is activated again in mid-to late

elongation stages. Thus, few transcripts bridge the gap between late spermatocytes and mid-

stage elongating spermatids, as if the two stages were disconnected. In contrast, in the scRNA-

seq dataset, many of these same genes showed continued high transcript levels throughout the

spermatid stages, presumably representing storage of mRNAs in the cytoplasm. In

consequence, the mature spermatocyte to elongating spermatid stage transcriptomes were well

connected through a smooth gradient of transcript levels in scRNA-seq data (Figure 4H, black

box).

Reactivation of transcription in mid-to-late elongating spermatids

The ability of snRNA-seq to highlight dynamic transcriptional changes during cellular

differentiation revealed striking transcriptional (re)activation of a subset of 162 genes in mid-to-

late elongating spermatids, a phenomenon previously described for only 24 genes, called "post-

meiotic transcripts" (Barreau et al., 2008). Analysis for genes specifically enriched in late

pseudotime identified a list of 162, here termed spermatid transcribed genes (Figure 4G, black

box, and Figure 5 - source data 1). These included 18 of the previously identified 24.   FISH

revealed *flyers-cup* (*f-cup*) RNA at the distal end of elongated spermatid bundles, as expected

(Barreau et al., 2008, Figure 5A,B; Figure 5 - figure supplement 1A). RNA from *loquacious*

(loqs), a newly identified spermatid transcribed gene, was similarly localized (Figure 5C; Figure

5 - figure supplement 1B). Transcripts from *walker cup* (*wa-cup*) and *scotti* (*soti*) also localized

to the distal ends of elongating spermatids as expected (Barreau et al., 2008, Figure 5D-F;

Figure 5 - figure supplement 1C, D). Analysis of earlier elongating spermatid cysts by single

molecule RNA FISH (smFISH) supported active transcription of *wa-cup* and *soti* in spermatids:

smFISH revealed foci in spermatid nuclei, suggesting nascent, post-meiotic transcription (Figure

5E,F, arrows), as well as perinuclear granules (Figure 5E,F, arrowheads), which could represent

newly synthesized RNAs being trafficked toward the distal ends of the spermatids.

Analysis of the snRNA-seq data showed that many newly identified spermatid

transcribed genes, including *Pp2C1* and *CG6701*, were initially expressed in spermatocytes or

spermatogonia, downregulated in early spermatids, and later reactivated during mid-to-late

elongation (Figure 4G-upper half of black box; Figure 5G,H). Other newly identified spermatid

transcribed genes, including *Parp16* and *Glut3,* were weakly expressed in spermatocytes but

robustly transcribed in elongating spermatids (Figure 4F - lower half of black box; Figure 5G,H).

Both patterns are consistent with RT-qPCR and RNA *in situ* hybridisation results for the 24 post-

meiotic transcripts previously identified (Barreau et al., 2008). Together the results show two

sources of RNAs in elongating spermatids: cytoplasmic perdurance of RNAs transcribed in

spermatocytes (Figure 4H), and *de novo* post-meiotic (re)activation of transcription of certain

genes (Figure 4G, Figure 5G,H).

The majority of the spermatid transcribed genes remain functionally uncharacterized,

and await investigation. GO term analysis showed no significant enrichment for any single

biological process or pathway, although several functional classes were represented (Figure 5 -

source data 1). Additionally, genes in this set did not appear to be coordinately reactivated, as

by a single regulatory circuit. Rather, the likelihood of genes to be (re)activated concordantly

was weakly correlated with their expression level, with a few outliers (Figure 5 - figure

supplement 1E).

Progression of differentiation in the somatic cyst cell lineage

Somatic cyst cells govern many germline transitions, from stem cell behavior through

sperm maturation and release (Figure 1A; Figure 6O). The snRNA-seq approach may be

especially useful for characterizing cyst cell transcriptomes across differentiation stages

because the long, thin, extended shape of many cyst cell types may make isolation of intact

cells difficult.

Cyst lineage identity was assigned by expression of three transcriptional regulators,

traffic jam (*tj*), *eya* and *sine oculis* (*so*) (Figure 1B,D; Figure 6A,B,U; Figure 6 - figure

supplement 1C,D). While both very early clusters (62, 36, 58) and late clusters (79, 68 and 80,

84, 72) were simple to describe (see below), the middle region of the 2-D UMAP presented a

tangle. However, re-projecting the lineage to preserve a third dimension clarified the

assignment of cluster order through the middle region (Figure 6C; see Materials and Methods;

rotatable 3D representation available at https://doi.org/10.5061/dryad.m63xsj454). This,

combined with pseudotemporal ordering (Figure 6 - figure supplement 1A,B) and marker

analysis, enabled assignment of specific clusters to the cyst cells associated with stem cell,

spermatogonial, spermatocyte, spermatid or sperm release stages of germline development.

Perhaps notably, the few clusters difficult to assign had relatively low UMI count (Figure 6E;

clusters 81, 94, 95, 67, 60).

The cyst lineage begins with proliferative *stg*-expressing CySCs (Cluster 62, Figure 1E;

Figure 6A,D).   Expression of *tj* suggested that clusters 36 and 58 represent early-stage cyst

cells that enclose transit amplifying spermatogonia (Figure 6B,U) labeled SgCC in Figure 6D, as

Tj protein marks the nucleus of these early cyst cells but was not detected by

immunofluorescence staining in spermatocyte-associated cyst cells (ScCC) (Li *et al.,* 2003;

Zoller and Schulz, 2012). Complementing this, *CG3902* mRNA was also detected in the same

clusters as *tj*, and a CG3902 protein trap line revealed cytoplasmic protein accumulation up to

early ScCC and no detectable protein thereafter (Figure 6U; Figure 6 - figure supplement 1E,F).

P-element induced wimpy testis (*piwi*) mRNA was enriched in the same clusters as *tj and*

CG3902, and also was detected at lower levels in nuclei of subsequent clusters (for example,

clusters 47 and 77; Figure 6D SgCC and ScCC, F,U). FISH and analysis of a Piwi protein trap

confirmed expression in early cyst cells associated with spermatogonia at the testis apex, as

well as in cyst cells enclosing growing spermatocyte cysts (Figure 6G-I, arrowhead and dashed

arrow, respectively). Interestingly, *piwi* transcripts were also detected in more mature cyst cells

associated with elongated and polarized spermatids (Figure 6G, solid arrow), highlighting the

differences between active transcription detected by snRNA-seq and perdurance of cytoplasmic

RNA detected by FISH.

Both *tj* and *eya* mRNA expression was detected in cluster 58, but onward, *tj* mRNA was

abruptly down regulated while *eya* transcript expression increased. We surmise this marks the

transition to cyst cells associated with spermatocytes, labeled ScCC in Figure 6D (Figure

6A,D,J,U), since Eya protein is known to accumulate in cyst cell nuclei from late stage

spermatogonia to spermatocytes (Fabrizio *et al.*, 2003). FISH confirmed the cyst cell *eya*

pattern (Figure 6K, arrowhead and dashed arrow), while also revealing accumulation of *eya*

transcript in cyst cells associated with post-meiotic spermatids (Figure 6L, solid arrow).

Additionally, sparser but detectable *eya* transcript was found in spermatocytes themselves,

corroborating our observation that it is one of several somatic markers expressed in

spermatocytes (Figure 3I; Figure 6K, white outline).

Intriguingly, the cyst lineage bifurcates after cluster 58, with clusters 77, 65, and 98

successively in one arm and 47, 104, and 88 in the other (Figure 6A,C,D). This might be due to

onset of differentiation of head versus tail cyst cells, and would represent the first hint at when

this occurs in the lineage. Identification and characterization of genes differentially expressed

within the split could reveal whether cyst cells specific to a given arm of the lineage govern

different properties of spermatocyte-containing cysts.

Expression of *Amphiphysin* (*Amph)* supported the conclusion that the bifurcation in the

cyst cell lineage after cluster 58 represents cyst cells associated with spermatocytes. *Amph*

mRNA was high before the bifurcation and persisted at lower levels in the two arms of the split,

dropping substantially by cluster 74 (Figure 6M,U). Expression of a protein trap confirmed that

Amph protein levels were high in SgCCs and ScCCs (Figure 6N, arrowhead and dashed arrow,

respectively) and declined significantly in cysts containing early spermatids (data not shown). It

is intriguing that the snRNA-seq and protein trap indicate that Amph expression is strongest in

early cyst cells, even though it encodes a BAR domain protein required to form the actomyosin

clamp that maintains head cyst cell membrane integrity as these cells wrap around spermatid

heads late in spermatogenesis (Kapoor *et al.,* 2021).

Branches of the cyst lineage rejoin at cluster 74 implying a transition to a common

transcriptional state. Interestingly, comparison of *Akr1B* vs. *Amph* suggests that cluster 74

contains cyst cells associated with early or elongating spermatids. Akr1B transcripts were

elevated in both arms of the split through to cluster 74 (Figure 6 - figure supplement 1G) while

Amph expression had decreased by cluster 74 (Figure 6M,U). Analysis of a *Akr1B::GFP* protein

trap confirmed its expression in cyst cells associated with elongating spermatid cysts (SpCC)

(Figure 6 - figure supplement 1H,I, arrows). Thus, it appears that, with respect to the

transcriptome, a developmental transition occurs within the cyst lineage as these cells mature

from support of spermatocytes to early spermatid cysts.

After cluster 74, the lineage again splits, with marker analysis suggesting that this
correlates with differentiation of late stage head cyst cells (HCC, clusters 69, 79, 68) versus
putative tail cyst cells (TCC, clusters 57, 80, 84, 72; Figure 6A,C,D). Enrichment of *eya* was
dramatically reduced in the clusters along either late cyst cell branch compared to those of
earlier lineage nuclei (Figure 6J,U). However, *Neprilysin-4* (*Nep4*), a metalloprotease involved
in male fertility (Sitnik *et al.,* 2014), was upregulated in both late branches (Figure 6P,U). FISH-
IF confirmed *Nep4* mRNA expression in late HCCs and TCCs associated with fully coiled
spermatids (Figure 6Q, arrow and arrowheads, respectively) with Coracle staining defining the
septate junctions between HCCs and TCCs. Intriguingly, the snRNA-seq data showed that
geko, an olfactory gene not studied in the testis (Shiraiwa *et al.,* 2000), was upregulated both in
ScCCs and in part of cluster 68 (Figure 6A,R,U). In fact, higher resolution (Leiden 8.0) analysis
divided cluster 68 into nuclei either enriched or not for *geko* (data not shown). FISH for *geko*
revealed expression in the testis basal region with high expression in HCCs (Figure 6T, arrow)
and lower expression in TCCs (Figure 6T, arrowhead) as predicted by the UMAP (Figure 6R,U).
FISH signal was also observed in cyst cells associated with spermatocytes and elongating
spermatid cysts throughout the testis (Figure 6S, dashed arrow). While TCC-specific transcripts
remain elusive, exploring the putative HCC and TCC clusters further will not only test our
tentative assignments, but possibly reveal unique roles and processes carried out by these cells
during spermatid retraction and coiling (Figure 6O).

The hub: architectural organizer and key signaling center

The hub is a small group of somatic, epithelial-like cells at the testis apex that acts as a
niche, providing signals that maintain GSC and CySC fate (Hardy *et al.,* 1979; Kawase *et al.,*
2004; Tulina and Matunis 2001; Kiger *et al.,* 2001; Leatherman and DiNardo 2010; Shivdasani
and Ingham 2003). Initial marker analysis suggested that the hub maps to cluster 90. However,
only 79 of these 120 nuclei were near to each other in 2D UMAP space, while other members of
cluster 90 were dispersed (Figure 7 - figure supplement 1A). The 41 outcast nuclei either
expressed germline genes such as *Rbp4, zpg, p53*, or *vas* (Figure 7 - figure supplement 1B-E)
or rarely or only inconsistently expressed signature genes known to be enriched in hub cells
(Figure 7A-D, dashed red circles). This strongly suggests that they differ transcriptionally from

the 79 tightly clustered nuclei. This conclusion was further supported by additional subclustering

of the 120 nuclei by two independent methods (Figure 7 - figure supplement 2A-B).

Consequently, cluster 90 was pared down to 79 definitive hub nuclei.

The snRNA-seq data show that hub nuclei express genes involved in signaling as well

as markers common in epithelial cells. For example, *upd1* and *hedgehog* (*hh)* are upregulated,

consistent with the hub's role in stem cell maintenance (Figure 7A,B,K) (Tulina and Matunis

2001; Kiger *et al.,* 2001; Michel *et al,.* 2012; Amoyel *et al.,* 2013). Additionally, the definitive hub

cell nuclei showed enriched expression of mRNAs encoding proteins implicated in cell-cell

adhesion/junctional complexes, including Fasciclin III (Fas3), E-Cadherin (Shg), N-Cadherin

(CadN), Discs large (Dlg1), and Cheerio (Cher; an orthologue of Filamin), all of which proteins

have been shown to mark hub cells (Brower *et al.,* 1981; Le Bras and Van Doren 2006; Boyle *et*

al., 2007; Tanentzapf *et al.,* 2007; Papagiannouli and Mechler 2009) (Figure 7C,D,F,K).

The definitive identification of hub nuclei allowed analysis for upregulated genes ($\log_2$FC

≥ 1, compared to the full testis plus seminal vesicle dataset; Figure 7 - source data 1) as new

candidate hub markers. One such encodes the transcription factor Tailup (Tup, also known as

Islet (Thor and Thomas 1997; She *et al.,* 2021; Boukhatmi *et al.,* 2012) (Figure 7E). Indeed, a

Tup::GFP transgene showed strong protein expression marking hub nuclei at the apex of adult

testes (Figure 7F-F'''). This is consistent with recent evidence that Tup is expressed in and

required for niche organization in the male embryonic gonad (Anllo *et al.,* 2019; Anllo and

DiNardo 2022). The expression of Tup in the 79 "true hub" nuclei of the snRNA-seq dataset but

not the other nuclei of the original 120 further validates the exclusion of the 41 dispersed nuclei

previously included in cluster 90.

Previous lineage-tracing showed that hub cells and CySCs derive from a common pool

of gonadal cells during embryogenesis (DeFalco, 2008; Le Bras and Van Doren *et al.,* 2006;

DiNardo *et al.,* 2011). Comparison of genes up-regulated in hub cells with those up-regulated in

CySCs strongly reflects this developmental relatedness. Hub and CySC (cluster 62) nuclei

shared 27% of their up-regulated genes, likely reflecting their embryonic co-origin (Figure 7G,

Figure 7 - source data 2). In contrast, the fraction of shared upregulated genes was much lower

between the hub and several lineally and functionally distinct cell types, including

spermatogonia, hemocytes and terminal epithelia (Figure 7 H-J). Supporting this, several genes

up-regulated in hub nuclei were also highly expressed in the CySCs and the cyst lineage, but

much lower in spermatogonia and hemocytes (Figure 7K). The transcriptional similarity

observed in the adult cell types could account for the ability of hub cells to replenish CySCs

after drastic injury and explain the shift of one lineage toward the other when specific gene

functions are compromised (Greenspan *et al.,* 2022; Herrera *et al.,* 2021; Hétié *et al.,* 2014;

DiNardo *et al.,* 2011; Voog *et al.,* 2014).

Epithelial Cells of the Testis Organ

A key role of the terminal epithelial cells (TE) is to anchor the head end of elongated

spermatid bundles at the base of the testis during individualization and coiling (Figure 6O) so

that the sperm are positioned for release into the seminal vesicle (SV). Marker gene analysis

suggested that both the TE and SV reside in the "mermaid" of the UMAP (Figure 1B, 8A).

snRNA-seq showed enrichment for *hh* in two broad areas of the UMAP "mermaid" (Figure 8B;

Figure 8 - figure supplement 1A), while FISH showed *hh* RNA in TE and SV cells (Figure 8C,

arrowhead and arrow, respectively). *Metallothionein A* (*MtnA*) was most highly upregulated in

the lower clusters enriched for *hh* (Figure 8A,B,D; Figure 8 - figure supplement 1B; clusters 42,

20), with colorimetric ISH confirming strong expression of *MntA* in the TE (Figure 8E, arrow).

The SV marker *Neprilysin-5* (*Nep5;* Sitnik et al., 2014), confirmed by FISH to be expressed in

SV, was upregulated in the uppermost clusters (mermaid head) (Figure 8A,F,G, arrow; Figure 8

- figure supplement 1C; clusters 1, 4, 7, 9, 29, 32, 34, 93). Although they encode nearly

identical proteins, *Nep4* and *Nep5* mRNAs were expressed in different cell types of the adult

testis (HCC/TCC and SV, respectively; Figure 8H). Note that while the TE and SV each map

as a single-block cluster at lower clustering resolution (Figure 2 - figure supplement 1B), the

appearance of multiple clusters for each cell type at higher resolution suggests notable

transcriptional heterogeneity within both cell types (Figure 8A). Thus, epithelial cell types in

general represent an underexplored area in testis biology.

Discussion

Study of the *Drosophila* testis has had great impact on reproductive biology and on

understanding stem cell-niche interactions, developmentally regulated cell cycles and cell type

specific cellular morphogenesis. Against this foundation, examining single cell and single

nuclear RNA-seq has revealed several notable features. First, in gene expression space, the

stem cell-based lineages for the germ line and the supporting somatic cyst cells were

geographically isolated from each other, reflecting their distinct lineage identity. Second, the

individual cells within each stem cell lineage were essentially arranged in strings, due to

progressive changes in gene expression during differentiation. Third, a UMAP for each lineage

revealed complexities along its trajectory that reflect both known intricacies of development

within each lineage, as well as previously unappreciated gene expression dynamics. Fourth, in

contrast to the two stem cell lineages, terminally differentiated cells generally clustered into

discrete groups, as expected. Finally, the comparison of single cell and single nucleus datasets

proved particularly revealing of dynamic developmental transitions in lineage differentiation.

Germline and soma map to distinct gene expression domains

While *Drosophila* spermatogenesis relies on two separate stem cell lineages, the

progeny of each lineage associate intimately with each other, cooperating at multiple points

during differentiation to produce functional sperm. Despite their physical association, in gene

expression space the two lineages lie well-separated. This is not a surprise, since the lineages

are specified independently during embryogenesis in space and time, as well as by different

gene regulatory circuits. Such separation between germline and supporting soma is also

observed in the scRNA-seq from the *Drosophila* larval testis, the ovary and murine testis

(Mahadevaraju et al, 2021; Green et al., 2018; Rust et al., 2020).

Accurate pairing of each germline cluster with a somatic cluster representing its

interacting partner could facilitate identification of the underlying cell signaling circuits that form

the basis for cooperation between the lineages at various points along their differentiation

trajectory. We are able to highlight some likely pairing assignments due to the extensive

knowledge of *Drosophila* spermatogenesis and testis biology. For example, the CySC cluster

(62) likely associates directly with germ cells in cluster 25, while cyst cell lineage clusters 36 and

58 likely associate with later spermatogonia (cluster 22). Other associations are also suggested

by the data (Figure 6D) with more remaining to be defined. The transcriptome in these pairings

can be mined to identify candidate signaling pathways by recently developed tools, such as

FlyPhone (Liu et al., 2022).

Stem cell lineages appear as strings along their differentiation path

The data highlight how tissues maintained by stem cell lineages display a characteristic

geography in gene expression space. The UMAP reveals these lineages to be arranged largely

in strings, due to progressive changes in gene expression. The linear arrangement is strongly

supported by trajectory inference for both the sc- and sn-RNA-seq datasets (see also Li et al.,

2022). This arrangement might be diagnostic of at least some stem cell lineages, as it is

observed to a degree in germline data from murine testes, in murine small intestine (Green et

al., 2018; Haber et al., 2017), and in the follicle cell lineage in the *Drosophila* ovary (Li et al.,

2022; Rust et al., 2020). While in some tissues maintained by stem cells, such an arrangement

is made apparent only by trajectory inference, in the *Drosophila* testis it is apparent in the UMAP

representing gene expression space.

Additionally, even with progressive changes in gene expression along their respective

trajectories (Figures 2L, 6U), cells in these strings nevertheless can be sorted into clusters. The

extensive study of germline and somatic testis biology provided excellent markers, which

allowed assignment of cell stage identity with high confidence for many of these clusters. These

assignments provide abundant opportunities to identify new, stage-specific markers to test for

function in each stem cell lineage *in vivo*.

Each stem cell lineage exhibited complexities, revealing known and previously

unappreciated expression dynamics

In the testis, differentiation proceeds stepwise from the apical tip of the tubule. It is

tempting to equate this progression with the arrangement of cell states revealed in the UMAP.

However, the UMAP represents gene expression space and not physical space. This means

that a stage characterized by dramatic changes in transcription in a differentiating lineage, while

localized to a very small region within the organ, might be spread out in a thin stream in the

UMAP. An example is seen in early spermatocytes, from completion of premeiotic S phase

through to apolar spermatocytes (Fuller, 1993). Physically located in a narrow band next to the

spermatogonial region in the testes, nuclei at these stages extend across much of the lower

region of the germ cell UMAP (Figure 2A). A reciprocal case is highlighted by early spermatids,

which have very low levels of transcription and thus show very low UMI values. These cells are

clustered together in the UMAP, even though the cells are undergoing dramatic changes in

morphology easily visible by microscopy (Fuller, 1993).

For the soma, the complexity observed within portions of the cyst lineage was a surprise.

Head and tail cyst cells execute very different roles for the spermatid bundles with which they

associate. Not surprisingly, the distinction between head and tail cyst cells is clear late in the

differentiation process in the UMAP, when the two cell types have very different morphologies.

Our analysis confirmed that both head and tail cyst cells are derived from the same progenitor

population (Figure 6D). However, the somatic clusters show complex intertwining before they

eventually resolve to generate the two lineages. Interestingly, this complex tangling seems to

coincide with the stages of spermatid development where there already is clear polarization in

the architecture of the bundle (Figure 6D). Only by reprojecting this lineage, purposefully

preserving an extra dimension, could we infer the trajectory properly, discerning a split and then

a merge in expression profiles that tracked with cell identity. The initial split in the cyst lineage,

as well as the subsequent and transient merge followed by a new split, suggests an interesting

sequence of transcriptional cell states within this key supporting lineage.

Differentiated cells map as discrete groups

Whereas the two stem cell lineages each appear as strings reflecting progression in their

gene expression patterns, terminally differentiated cell types appear as more discrete patches in

the UMAP (Figure 1G). This was the case for differentiated cell types that are an integral part of

the testis, such as the hub and terminal epithelia, as well as for cell types that associate with the

testis but are structurally distinct, such as the seminal vesicle. Even within a patch, however,

increasing cluster resolution reveals complexity in cell type identity. For example, the snRNA-

seq identifies a 'muscle' group (Figure 1B, 'mc') that is composed of several different Leiden 6.0

clusters (Figure 2 - figure supplement 1, compare A with B). Perhaps the clusters reflect

different muscle cell types, for example those covering the testis tubule vs. the seminal vesicle

(Susic-Jung et al., 2012). Likewise, different clusters that comprise 'seminal vesicle' may

represent distinct portions of the structure, such as entry and exit points (Figure 8A).

The value of comparing sn- versus sc-RNA-seq

When carried out on the same tissue, single nucleus RNA-seq is typically comparable to

single cell RNA-seq, with very high percent similarity in gene identification (McLaughlin et al.,

2021). In the *Drosophila* testis, this was largely the case for germ cells early in the differentiation

lineage (Figure 4 G and H). In contrast, comparing these two approaches for later stage germ

cells revealed a striking difference, reflecting an important aspect of testis biology. Our data

show that directly comparing sn- to scRNA-seq can highlight cases where mRNAs are

expressed at an early developmental stage (e.g., in spermatocytes) then stored for later use

(e.g., in spermatids, where mRNAs may be used temporally). This phenomenon was previously

described for a relatively small set of transcripts that persist in the cytoplasm for days following

the cessation of transcriptional activity during *Drosophila* male meiosis (Fingerhut et al., 2019;

Kuhn et al., 1991; White-Cooper et al., 1998). Our results demonstrate that such perdurance is

likely a feature of hundreds of transcripts.

Key to this finding was the use of sn- and scRNA-seq datasets in tandem, rather than

individually. snRNA-seq likely reports actively transcribed mRNAs at a given developmental

time and therefore may be more sensitive to dynamic changes in transcription than scRNA-seq.

Thus snRNA-seq may be a better approach for mapping changes in cell state, for example,

during embryonic development or differentiation in stem cell lineages. Data from scRNA-seq,

by contrast, may be strongly influenced by mRNAs perduring in the cytoplasm from earlier time

points, as well as mRNAs no longer being actively synthesized but purposefully stored for later

usage. Thus, while snRNA-seq reveals gene expression dynamics through *active* transcriptional

changes, scRNA-seq can capture post-transcriptional gene regulation, as is required by

transcriptionally silent cells like early spermatids. Together, these datasets allow for inference

of the full lifetime and velocity of all RNAs, from nuclear transcription to cytoplasmic degradation

and/or protection.

Dynamic inference of transcript state by separately assaying nuclear and cytoplasmic

RNAs, as used here, could prove a relevant measurement in many other differentiation systems

as well. Cytoplasmic storage of transcripts expressed at an earlier stage to be used later is a

widespread differentiation strategy and is especially predominant during oogenesis and

spermatogenesis. Many mRNAs expressed during oogenesis are stored in the cytoplasm in a

translationally silent state, to be recruited for translation in the early embryo (Jenkins et al.,

1978). This is especially important in organisms with large, yolk-rich eggs, in which transcription

from the zygotic genome is delayed until after several rounds of mitotic divisions. In the male

germ line as well, where transcription ceases during spermiogenesis as the nucleus compacts,

many mRNAs expressed at earlier stages are stored in the cytoplasm, initially translationally

repressed, then recruited for translation during spermatid morphogenesis. In both cases,

perdurance of mRNAs in the cytoplasm after transcription has shut down is important to allow

subsequent stages of development and differentiation to take place, and recruitment of specific

mRNAs for translation at different times may play an important role in temporal control of

morphogenetic events.

It is also notable that just 18/162 of the spermatid transcribed genes expressed late in

spermiogenesis encode proteins detected in the mature sperm proteome (Wasbrough et al.,

2010). While this might be due to limited sensitivity in proteome detection, alternatively, many

may play roles in spermatid development but not mature sperm function. Examples could

include regulating or mediating spermatid elongation, the histone-to-protamine transition, or

individualization, as is the case for *soti* (Barreau et al., 2008; Kaplan et al., 2010). A further

twenty of the 22 cytoskeletal or motor-related genes transcribed in spermatids (see Figure 5 -

source data 1 for functional classes of the spermatid transcribed genes) have predicted

functions in microtubule assembly, flagellar axoneme assembly, axonemal dynein regulation or

microtubule transport, consistent with roles in assembling and elongating the 1.8mm sperm

flagellum and transporting cargos within this very long cell (Ghosh-Roy et al., 2004; Noguchi et

al., 2011; Tokuyasu, 1975). The set of late upregulated genes identified as involved in lipid

synthesis, lipid transfer and membrane trafficking could contribute to membrane addition at the

distal (growing) ends of spermatid flagella to facilitate cell elongation (Ghosh-Roy et al., 2004).

Finally, genes encoding RNA binding proteins could regulate transcript localization or translation

in the increasingly long spermatid cells, as polarized mRNA localization by RNA binding

proteins such as Orb2 (localizing *aPKC* and *orb2* mRNA) and Reptin and Pontin (localizing

axonemal dynein mRNA) has been observed and shown to be important for sperm maturation

(Fingerhut and Yamashita, 2020, Xue et al., 2014). Whether and how many other mRNAs are

localized to the growing flagellar tip in spermatids remains to be studied.

Use as a resource

With the transcriptional profiling of over 44,000 nuclei isolated from testis and associated

supporting tissues, we have connected differentiation events throughout the germline and

somatic lineages, capitalizing on the extensive literature on *Drosophila* testis biology. The

expectation is that this will be a foundational resource for the field. Several other *Drosophila*

RNA-seq and scRNA-seq efforts have been reported (Gan et al., 2010; Hof-Michel and Bökel,

2020; Mahadevaraju et al., 2021; Shi et al., 2020; Vedelek et al., 2018; Witt et al., 2019). Since

each approach has different relative strengths and limitations, the foundation we have laid with

the scRNA-seq and snRNA-seq datasets described here should assist others in comparisons

with more stage-restricted transcriptome analyses. More broadly, the data presented here, in

their easily shared formats, should enable a deeper exploration of the conserved aspects of

germline and support cell biology during *Drosophila* and mammalian spermatogenesis.

Materials and Methods

Key Resources Table

Reagent type (species)	Designation	Source or reference	Identifiers	Additional information
genetic reagent (D. melanogaster; male)	w1118	BDSC		
genetic reagent (D. melanogaster; male)	Oregon-R	BDSC		
genetic reagent (D. melanogaster; male)	y1w1	BDSC		

				y w; PBac{y[+mDint2] w[+mC]=tup-GFP.FPTB}VK00031/TM6C, Sb
genetic reagent (D. melanogaster; male)	tup::GFP	BDSC	RRID:BDSC_81278	
genetic reagent (D. melanogaster; male)	Amph::GFP	Kyoto Stock Center	CPTI-002789	
genetic reagent (D. melanogaster; male)	piwi::GFP	Kyoto Stock Center	CPTI-003588	
genetic reagent (D. melanogaster; male)	Akr1B::GFP	Kyoto Stock Center	CPTI-002728	
genetic reagent (D. melanogaster; male)	CG3902::GFP	Kyoto Stock Center	CPTI-100001	
genetic reagent (D. melanogaster; male)	*comr z2-1340*	White-Cooper	FBal0144389	
antibody	anti Fasciclin III, mouse monoclonal	Developmental Studies Hybridoma Bank	DSHB:7G10; RRID:AB_528238	1:50
antibody	Chick polyclonal anti GFP	Abcam	ab13970	1:10,000
antibody	goat anti chicken Alexafluor 488	Invitrogen	A11039	1:200'
antibody	goat anti mouse Alexafluor 568	Invitrogen	A11004	1:100'
antibody	anti-FITC, horseradish peroxidase	Roche	11426338910	1:2,000
antibody	anti-Digoxygenin, horseradish peroxidase	Roche	11207733910	1:1,500
antibody	anti-digoxygenin, alkaline phosophatase	Roche	11093274910	1:2,000
antibody	coracle	Developmental Studies Hybridoma Bank	DSHB: C615.16	1:20
recombinant DNA reagent	piwi (plasmid containing cDNA)	BDGP DGC library	GM05853	FBcl0142639
recombinant DNA reagent	eya (plasmid containing cDNA)	BDGP DGC library	GH05272	FBcl0108545

recombinant DNA reagent	f-cup (plasmid containing cDNA)	BDGP DGC library	GH09045	FBcl0128895
recombinant DNA reagent	loqs (plasmid containing cDNA)	BDGP DGC library	RE14437	FBcl0204474
recombinant DNA reagent	geko (plasmid containing cDNA)	BDGP DGC library	RE30284	FBcl0192532
recombinant DNA reagent	Vsx1 (plasmid containing cDNA)	BDGP DGC library	SD01032	FBcl0286608
chemical compound, drug	DAPI	Millipore Sigma	1023627600	1:1,000 (of 2-5 mg/ml)
chemical compound, drug	Vectashield	Vector Labs	H-1000	
chemical compound, drug	GMM tissue clearing reagent	Gary Struhl		2g Canada Balsam in 1ml methyl salicylate
chemical compound, drug	rhodamine tyramide	Thermo Scientific	46406	1:1,000
chemical compound, drug	fluorescein tyramide	Thermo Scientific	46410	1:1,500
commercial assay or kit	Stellaris RNA FISH Probe Designer	Biosearch Technologies Inc.		
commercial assay or kit	10X Genomics Chromium Controller; Next GEM Single Cell3' v3.1	https://www.10xgenomics.com/product-catalog	1000269	
software, algorithm	Seurat 4.0.5	Satija et al., 2015		
software, algorithm	Monocle3	https://cole-trapnell-lab.github.io/monocle3/		
software, algorithm	Venny 2.1	https://bioinfogp.cnb.csic.es/tools.html		
microscopy, lenses	Olympus Bx50 microscope	Olympus		
microscopy, lenses	20x/0.60 NA objective	Olympus		
microscopy, lenses	10x/0.30 NA objective	Olympus		

	40x/0.75 NA			
microscopy, lenses	objective	Olympus		
microscopy, lenses	KY F75U Camera	JVC		
microscopy, lenses	Orca-05G Camera	Hamamatsuu		
microscopy, lenses	63x/1.4 NA objective	Zeiss		
microscopy, lenses	LSM800 confocal	Zeiss		
microscopy, lenses	63x/1.4 NA objective	Leica		
microscopy, lenses	Stellaris Confocal	Leica		

Drosophila lines

For snRNA-seq, testes and attached seminal vesicles were dissected from 0-1 day old w^{1118}

males and processed as described in (Li et al., 2022). For scRNA-seq, testes alone were

dissected from 1-5 day old Oregon-R males. Oregon-R testes were also used for *in situ*

hybridization of *aub* and *fzo* (Figure 2). $y^1 w^1$ flies were used for smFISH in Figure 5.

Amph::GFP (CPTI-002789), piwi::GFP (CPTI-003588), Akr1B::GFP (CPTI-002728),

CG3902::GFP (CPTI-100001) and Tup::GFP (BDSC line 81278) were from CPTI and Bac

collections (Kudron et al., 2018; Lowe et al., 2014). *comr$^{Z2-1340}$* homozygotes and w^{1118} were

used for colorimetric *in situ* hybridization.

Testis squashes and analysis of expression of fluorescent fusion proteins

All tissue analyses included samples from multiple males of each genotype (5-10 pairs of testes

per experiment), typically processed as a single batch. Testes from transgenic flies of the YFP

CPTI collections were dissected in testis buffer (183mM KCl, 47mM NaCl, 10mM Tris pH6.8),

cut open using tweezers and gently squashed on a glass slide by application of a coverslip.

Testis squashed preparations were imaged live in sequentially captured images by phase

contrast and epifluorescence microscopy using an Olympus Bx50 microscope, with 20x, 0.60

NA, 10x, 0.30 NA, or 40x, 0.75 NA UPlanFl objectives and either a JVC KY F75U or a

Hamamatsu Orca-05G camera.

For Tup::GFP analysis, testes were dissected from BAC transgenic flies, fixed in 4%

paraformaldehyde for twenty minutes, and blocked in 3% bovine serum albumin, 0.02% sodium

azide, 0.1% Triton X-100 in phosphate-buffered saline for one hour. After a one hour wash with

PBX (0.1% Triton X-100 in 1X phosphate-buffered saline, pH 7.4), testes were incubated

overnight at 4°C with antibodies to GFP (Abcam 13970; 1:10,000) and Fasciclin-III

(Developmental Studies Hybridoma Bank 7G10; 1:50). After washing with PBX testes were

treated with goat anti-chicken AlexaFluor 488 (Invitrogen A, 11039; 1:200), goat anti-mouse

AlexaFluor 568 (Invitrogen A11004; 1:100), and DAPI (Millipore Sigma 1023627600; 1:1,000)

for two hours. After final washes in PBX, testes were mounted in VectaShield (Vector Labs H-

1000), and images were captured at 63X, NA 1.4, on a Zeiss LSM800 confocal microscope.

RNA in situ Hybridization

For *in situs* presented in Figures 2 and 4:

aub (forward primer 5'-CCTGGGCGGCTACATCTT-3'; reverse primer 5'-

GCGCAGATTTCGACTCGG'-3), *kmg (*forward primer 5'-TGCCTCTATGCCTCACGC-3';

reverse primer 5'- GCGCCTACCGGTCTCATC-3''*), fzo* (forward primer 5'-

GGCATCCAAACTCTCGCG-3';

reverse primer 5'-TGTCGCAACTGGAGCTCA-3') were amplified by PCR on cDNA from

Oregon-R Drosophila Testes. Using TA cloning (Promega, "Easy-T" Cloning), the resulting

amplicons were cloned into th pGEM vector (Promega). Subsequent PCR added a T7 binding

site (5'-GAAGTAATACGACTCACTATAGGGAGAGGG-3') upstream of the amplicon. The

resulting plasmids then served as templates for *in vitro* transcription with Digoxigenin (DIG)- and

fluorescein isothiocynate (FITC)-labeled ribonucleotides to generate labeled single-stranded

antisense riboprobes.

Testes were isolated and fixed in 4% formaldehyde for 30-60 minutes, dehydrated into methanol

and stored at -20°C for up to 1 month. After rehydration in PBS + 0.1% Triton-X100, testes

were permeabilized with 4 ug/ml Proteinase K for 6 minutes and washed with Pre-Hybridization

solution (50% deionized formamide, 5x SSC, 1mg/ml yeast RNA, 1% Tween-20) for up to 2

hours at 56°C. Testes were incubated overnight at 56°C with probes diluted 1:800 in

Hybridization solution (50% deionized formamide, 5x SSC, 1 mg/mL yeast RNA, 1% Tween-20,

and 5% Dextran Sulfate). After washes in Pre-Hybridization solution, 2x SSC, and 0.2x SSC,

then PBS+0.1% Triton-X100, samples were blocked for 30 minutes in 1% Roche Western

Blocking Reagent prior to incubating overnight at 4°C with either anti-FITC with horseradish

peroxidase conjugate (Roche) at a 1:2000 concentration or anti-DIG- with horseradish

peroxidase conjugate (Roche) at a 1:1500 concentration in 1% Roche Western Blocking

Reagent. Fluorescent tyramide development and amplification were performed by first placing

the testes for 5 min in borate buffer (0.1M boric acid, 2M NaCl, pH 8.5), followed by 10 min in

borate buffer with rhodamine (1:1000) or fluorescein (1:1500) tyramide, and 0.0003% hydrogen

peroxide. After development, peroxidase activation was performed in a 1% sodium azide

solution for at least 1h, followed by antibody labeling for the second probe. Coverslips were

mounted in Vectashield with DAPI (Vector Labs). Fluorescence image acquisition was

performed on a Leica Stellaris Confocal microscope using a 63X oil-immersion objective (NA =

1.4).

For _in situs_ presented in Figures 5, 6, 8, and Figure 3 – figure supplement 2:

Fixation and hybridization for FISH was as described for Figures 2 and 4, with minor

modifications as described (Wilk et al., 2017). Briefly, these include using a cold acetone

permeabilization step and 0.3% Triton X-100 instead of Proteinase K for improved tissue

permeabilization, and DIG-labeled probe detection via tyramide amplification. DAPI (4" ,6-

diamidino-2-phenylindole; Sigma, cat. no. D-9542) was used at 1 ug/ml to reveal nuclei. For

detection of _piwi_, _eya_, _f-cup_, _loqs_, _geko_ and _Vsx1_ transcripts, RNA probes were transcribed from

the BDGP DGC library plasmid clones GM05853, GH05272, GH09045, RE14437, RE30284

and SD01032, respectively using T7 RNA polymerase. Testes were incubated with an antibody

to Coracle (Developmental Studies Hybridoma Bank C615.16; 1:20) for detection of Coracle

protein in Figure 3 – figure supplement 2. For _hh_ and _Nep5_, templates were made by PCR from

genomic DNA using the following T7 and T3 promoter-containing primers:

_Hh_ last exon:

Forward: GTAATACGACTCACTATAGGGAGACCACTGCCGATTGATTTTCTCAGG

Reverse: AATTAACCCTCACTAAAGGGTTGTGGAGATCGTGTTTTGAGCAT

Nep5 exon 6:

Forward: GTAATACGACTCACTATAGGGAGACCACGGGGAAATCCGATAAAGCTC

Reverse: AATTAACCCTCACTAAAGGGTTGTATCTGCAGAACCAAACTGAC

For colorimetric *in situ* hybridisation in Figure 3, 8 and Figure 3 - figure supplement 2

Probe preparation and *in situ* hybridisation were performed as described (Morris et al., 2009).

Primers Hml-F ATTTAGGTGACACTATAGAATAAGTGGACCCATGCCAAG and Hml-R

TAACCCTCACTAAAGGGTGACCATCATCGCAAATC and. primers MtnA-F

ATTTAGGTGACACTATAGAAGCGGTAAGTTCGCAGTC and MtnA-R

TAACCCTCACTAAAGGGACATTTATTGCAGGGTGTG were used to amplify 628bp and 443bp

fragments respectively from cDNA generated from w^{1118} testes. After re-amplification using

primers 5'-SP6 ACGGCAATTTAGGTGACACTATAGAA and 3'-T3

GCAACGAATTAACCCTCACTAAAGGG, and the products served as templates for T3 RNA

polymerase to generate dig-labeled single stranded, antisense RNA probes. The probes were

hydrolysed for 15 minutes, precipitated and resuspended in 200 µl water. Testes were dissected

from young w^{1118} males, fixed in 4% paraformaldehyde for 20-60 minutes, washed in PBS,

permeabilised with 50 µg/ml proteinase K for 5 minutes, washed in PBS, then hybridisation

buffer (HB: 50% Formamide, 5× SSC, 100 µg/ml denatured sonicated salmon sperm DNA, 50

µg/ml heparin, 0.1% Tween 20, 100 mM citric acid). Probes were diluted 1:100 in HB and testes

were hybridized for 16 hr at 65°C. Testes were washed 6x30 min at 65°C in HB, followed by 15

min each step at room temp in 4:1, 3:2, 2:3 1:4 HB:PBST, 2x15 min PBST, then incubated

overnight at 4°C in alkaline phosphatase-conjugated anti-digoxygenin antibody diluted 1:2,000

in PBST. Testes were washed 4x15 min in PBST and finally 3x 5 min HP (100 mM NaCl, 100

mM Tris (pH 9.5), 50 mM MgCl2, 0.1% Tween 20). NBT and BCIP diluted in HP were added as

a colorimetric substrate for alkaline phosphatase and color allowed to develop in the dark at

room temperature. The testes were washed in PBST, dehydrated through an ethanol series,

mounted in GMM, and imaged using DIC microscopy (10x objective) with a JVC KY F75U

camera mounted on an Olympus BX50 microscope.

For single molecule FISH presented in Figure 4

Testes were dissected in 1X PBS and fixed in 4% formaldehyde in 1X PBS for 20 minutes.

Testes were then washed twice for five minutes each in 1X PBS with 0.1% Tween 20 followed

by permeabilization in 1X PBS with 0.1% Triton-X 100 for two hours. Next, the testes were

washed twice with 5X SSCT (0.1% Tween 20) for five minutes each. Probes were added to pre-

warmed Probe Hybridization Buffer (Molecular Instruments) to a final concentration of 16nM.

100uL of hybridization solution was added to each sample, pipetted to mix, and allowed to

hybridize overnight at 37C. Samples were then washed four times with Probe Wash Buffer

(Molecular Instruments) for 15 minutes each at 37C. During the first wash, amplifier hairpins

(Molecular Instruments) were individually incubated at 95C for 90 seconds and allowed to slowly

cool to room temperature. Hairpins were then added to Amplification Buffer (Molecular

Instruments) to a concentration of 60nM. After washing the samples twice with 5C SSCT for five

minutes each, 100uL of hairpin solution was added and the samples were allowed to incubate at

room temperature overnight. The testes were then washed twice with 5X SSCT for 30 minutes

each at room temperature, mounted in Vectashield with DAPI, and imaged on a Leica Stellaris 8

confocal microscope with a 63X oil immersion objective lens (NA = 1.4) and processed using

ImageJ and Adobe Photoshop software. Probes (sequences below) and amplifier hairpins

designed by Molecular Instruments (https://www.molecularinstruments.com).

l(2)41Ab exon probe-binding sequences	l(2)41Ab intron probe-binding sequences
CATgAAAgAATggCAgAACTAggggACATgCAA AgCTTAggAAgTAACATCA	CTCTCgCTgTCTgCAAgCgAATAAAAACgATT ACTCTATAACATgCATgTgA
CTATCAATgCTTgAggTTgAAAAACTTAgATgC gTAgTTTTTAATgAAAgAg	TCgACggACATTCCAgCATCTATCATCTTCTT TACTgATAgCAAATTgTgCg
ggAgAATTgCAAgTTCgTgAACATCAgCAgTTT TggATAATgCTgCAgAACC	gCCAACCTCTCATgCCACAAgCgATACTCTgC ATCATTTgATATTgCAgATg
CgAAAACAACTATCCATTATTAggCAAAAgTC ATTAAAAAgTgACCAAgAgC	AgCgTAATCggACCgCATACATCCgATTgAAT TACAAATAgTgggCgTTCAA

AgAgACgCTCAgTTgCAAgACCAgCTATCACAggAAAAACgAAAATCAgATA	TTAgTgCTTATACTgTAgTTgTgggTATTAggAACCgCTAAATgATAgCTgT
ACACAggCATggAACgAggTTATTATAgCCATTCAgCTgAgAgATAAAATTT	CgACCAgTATAgTTTTgATCgATTTCTCgTCAAATTTTCgCTTTCgAACTTT
TTTCAAgTACAAgACgCAgCAAgAAAACACAgggAggCAATgAAACACAATA	TCATAAATTTTCTCgAgTTCTCATTCCACAACTTATAgCCATTTggCTCATA
AAATTggATTCCTTTAAgCAACgACTggCAgATgAAACTAgTTCCTTAAgAA	CATTTgAAgTCACCgATTTCTTgTTTATTTCATCCACCACTACATCTCTAgC
gAATgTCAggATgAAAgCgAATCTCTTTTgAAgATgCAAgCAAAgCTAgATg	TTAgAgAATCACTgTATTggCCAAAAAAATCATgTTTTTCCACAgAgTCATT
ggAAgTCATCTATTCACAgAAgTTCgAAAACTAgACATCgCCTTggATAAAT	TCgTAgATTCACAATTTTCCTgCAATggAAAATTCgACTCTTCATTCTCTTT
TTAACgCAAAAgTTCCgCACTgAggAAACAATTCTACggTTAgCgCAggATC	TgTATTCTCgAgACACgTggAATTTTCATTTggATTTTCATCAgACTTCgAA
TTAATAACAgACAAggCTACCgCTTACCgTTTgAAACTATTAggTAAATCTC	gATTTTAATggCATCCTCCCACAAggTTCTgTCTTTACgAgTTTTAATCTCT
gTAgAggAgACTgAgTTCgACCACgAACAggCTgAgCAAACTCAAAAACTAA	gATCggCTCgAAgTgAgTCgCgAgAAACgAAAgTgTCTAgCCAgCATATTTg
ATTgCggAgCTggAAATACTTAAATTggAAggAAAAATTgTTgAgCTACAAA	CgACTTAAgATACTTATCgATgTACTTAAgAggAggAgTCCCCCAACTggAT
CgCTCTgTAAAAggTgAAATCggTCTTATgAAAACTgAAATACACCgCATgA	ACTTTCAAATTCCTTTTAAATgATAACCTTAAATTgACAgTTgCCTTTAgAT
CATTgCgTTATgCATCgTgAACAAATCTTTgTCAATgCATCggTCAAggAgC	TATTCAgCTTCggTAgATgAAgCAgCAACCgATTTTTgTTTAATAgTACTCC
CgTAgAATTCAgTCAgATCTTAACgATATCATTAAggAggACgCCAATATTC	gTTggTAgTggCTTTgTCAAAATATTAgAAAATTgTAAgTCTgTggAAAgAT
gATCgAATAgAAgAAggAATATTAgCTAAgCATgCCAACTTggAACAAATAA	ATCATTCAAgATTCggTTAgCCCCAgTTggTCTCgCAgTgCAgTgAATCTgg

CAAgTAATACTCCgTTCAgAgTTgACAgTTCAACAgCTATCTCAAAAgCAAA	ATTTCTTTACAATACAACACgTTgTgTTATgTAATgTTgCTTCCAgAATggT
gAgCTAAATTATgATCTCACAgAAATAgCgAAggCTCTTTTACTAgACTATC	AgTTTAACTTgTCCTTATgCAgTAgCATAAACAAATTCTCCACgCTTTgCAA

For single molecule FISH presented in Figure 5

smFISH was performed as previously described (Fingerhut et al., 2019). Briefly, testes were

dissected in 1xPBS (Invitrogen, AM9624) and fixed in 4% formaldehyde (Polysciences, Inc.,

18814-10) in 1xPBS for 30 minutes, washed twice in 1xPBS for 5 minutes each, and

permeabilized in 70% ethanol overnight at 4°C. Testes were washed with wash buffer (2x

saline-sodium citrate [SSC, Invitrogen, AM9770], 10% formamide [Fisher Scientific, BP227]),

and then hybridized overnight at 37°C in hybridization buffer (2xSSC, 10% dextran sulfate

[Sigma-Aldrich, D8906], 1mg/mL yeast tRNA [Sigma-Aldrich, R8759], 2mM Vanadyl

Ribonucleoside complex [New England Biolabs, S1402S], 0.5% BSA [Invitrogen, Am2616], 10%

formamide). Following hybridization, samples were washed twice with wash buffer for 30

minutes each at 37°C and mounted in Vectashield with DAPI (Vector Laboratory, H-1200,

Burlingame, USA). Images were acquired using an upright Leica Stellaris 8 confocal microscope

with a 63X oil immersion objective lens (NA = 1.4) and processed using ImageJ and Adobe

Photoshop software (National Institutes of Health, Bethesda, USA).

Fluorescently labeled probes were added to the hybridization buffer to a final concentration of

100nM. Probes against *soti* and *wa-cup* transcripts were designed using the Stellaris® RNA

FISH Probe Designer (Biosearch Technologies, Inc., Novato, USA) available online at

www.biosearchtech.com/stellarisdesigner.

Probe Target	Fluorophore	5'-Sequence-3'	

wa-cup	Quasar® 670	cttccgataagctcatttgg, ccgctgttgggtgaaaaaga,
		ccagagcgctgttgaaatac, gagctctttcattgaacgga,
		atcgatcttttcagctgact, ctttcggctcatcaacagat,
		gactgacattgggattggtt, tacagagcatcgcagacttc,
		gaatcttccaagcgattgga, ccggagctaaatcgctttaa,
		tgagacggtcgagaacagga, gacatggtggtatcatctga,
		aactagccatcatgcgattg, gtatccttaatatccttggg,
		gttttccattatgctaacca, agtggggacaaatgggtttc,
		ttaaaatgcctttttcgcct, ccagcatttgttcagatacg,
		tcgcagtgtcttcagaaagg, gacttttttgcaatgcttggg,
		catcagtttgcccaaatact, tgcgttccgctttataattg,
		taaggcgtacatgggacttc, ctcgttgggatatttctgtt,
		ctcccggtgcttattataaa, gtctggtactgaatgcgata,
		atgctgcgcagaatcttgaa, gttctccaactcgaattagc,
		cccatgacttcctcaataaa, cgtcttgatggtgacatagt,
		aaatttccacggcattacgc, aatgcgagctaaacccaagt,
		agacagtcatattgctggga, gtgttagcagacgttgtttg, actcgttcgttttgtctttg,
		gtagccgatctggttatatt, ttaaaatgctccgctttggg,
		atacgattttccagtcggac, gttcaatgtgatactcggca,
		aattcgtgcagtagataggc, acaattcagatgctcttggg,
		ctccatataacactcttgca, gccttgcatataaccatgag,
		taagcacaggtcaaggttct, ctcctccgcattaactttaa,
		tgagccaaacttttgtctct, ctgatcgttgctttggaaca,
		taatttggttgcgatcctca

soti	Quasar® 570	tctcgacgaggtaatttg, tccgtgtagtacgtccat, gctcatcgtacagatcgt, ccgactcgatcgattagc, atcttcattcaccgcgtc, tgtccaagtcatcgccag, tgctgtccatcctccaat, tgacgattgactcccagg, gtccaggagtatgtccat, caacggtggctcttgagg, ctccttgcgccggaaaaa, acgtggtggtccatttgg, aacttcgtttcttccgcc, ggagtgggtttggtcata, ctcctgactttggcatgg, ttaggaggcacatctccg, attgccctcgtgacactg, atcctcgcgaacgtgacg, caaagtactcgcctcgct, gggtagttctgactggtc, tggcagacccataccatt, agaactgaccccaatgct

Preparation of samples for snRNA-seq

1000 FACS-sorted nuclei were obtained from hand dissected, 0-1 day-old adult testes (plus seminal

1001 vesicles) and processed as described by the Fly Cell Atlas project (Li et al., 2022). Data from

1002 44,621 sequenced nuclei passed quality control metrics. The raw data are publicly available

1003 (https://www.ebi.ac.uk/arrayexpress/experiments/E-MTAB-10519/).

1004

Importing FCA data into Seurat

The "Testis, 10X, Relaxed" loom file (https://flycellatlas.org/; Li et al., 2022) was imported into

Seurat 4.0.5, and a standard pipeline run on the resulting Seurat R object to normalize and

scale the data (NormalizeData, FindVariableFeatures, and ScaleData). The loom file had

already been filtered with nuclei expressing less than 200 genes or exceeding 15%

mitochondrial content removed and genes not expressed in at least 3 nuclei removed. We

chose to use the relaxed rather than a '10X, Stringent' testis loom file due to the particular

biology of the testis. Testis germline cells can express 'somatic' genes (see Results), including

the hub cell marker *upd1*. As a consequence, filtering algorithms that generated the stringent

dataset led to loss of a documented somatic cell type (the hub), as well as a large number of

late spermatocytes. The Fly Cell Atlas website provides links and tutorials to ASAP and SCope,

two web-based pipeline and visualization portals where users can examine the data or re-run

analyses (Davie, 2018; Gardeux, 2017). Within the SCope interface, select the 10X > Testis >

Relaxed dataset, and Settings > HVG UMAP. The analysis in this paper complements SCope

and ASAP with Seurat and Monocle3, two R programming-based tools for single cell analysis

(Satija et al., 2015; Trapnell et al., 2014, Tools for Single Cell Genomics • Seurat; https://cole-
trapnell-lab.github.io/monocle3/).

1022

The cluster information contained in the original loom file was preserved in the Seurat object,

with clusterings available at increasing levels of resolution (Leiden algorithm, 0.4 to 10). The

level of granularity provided by resolution 6.0 was deemed most revealing and is used for most

analysis here. The full Seurat Object, FCAloomToSeurat2TFP_Annotations.rds, is available for

download at https://doi.org/10.5061/dryad.m63xsj454, as is the script used to produce it from

the loom file (convert_loom_to_seurat_normScale_adjHub.R).

1029

1030

1031   Original cluster 90 was manually split into two in the Seurat Object,

1032   FCAloomToSeurat2TFP_Annotations.rds, with one resultant cluster of 79 nuclei definitively

1033   established as representing the hub (retaining its cluster number, #90), and the remaining nuclei

1034   placed into a new cluster, #111. Venn diagrams were created using Venny 2.1 (Oliveros, J.C.

1035   2007-2015).

1036

In some cases, data were extracted from subsets of the original Seurat object: a germline only

subset (Figures 2 through 5), a somatic cyst cell lineage subset (Figure 6), and a subset

representing several epithelial cell types of the testis as well as specific additional epithelial cell

types (Figure 8). All UMAPs were generated within Seurat using the "DimPlot" function. UMAPs

that highlighted particular genes of interest were generated using the "FeaturePlot" function

(Figures 1, 2, 3, 5, 6, 7, 8). Dotplots were generated using the "DotPlot" function (Figures 2

through 7).

1044

The testis 10X Relaxed loom file only contains a UMAP reduction projected to two dimensions.

To inspect a 3D UMAP representation, the appropriate lineage was isolated and reprojected,

passing the argument 'n.components =3L' to the Seurat function 'RunUMAP' (see

cystlineage3Dcode.R code; (Qadir et al., 2020); 10.5281/zenodo.348317). An html version of

the resulting 3D representation is available for download at

https://doi.org/10.5061/dryad.m63xsj454.

1051

1052   For analysis of heart, Malpghian tubule and male reproductive gland (Figure 3 - figure

1053   supplement 1A), the appropriate stringent 10X loom file from Fly Cell Atlas was imported into

1054   Seurat. Means were calculated for nCount (UMI) and nFeature (gene). Cluster numbers were

1055   assigned using the "Annotation" metadata field in each object, and a plot produced for average

1056   UMI by cluster number, with dot size reflecting average gene number per cluster.

1057

Graphs generated directly from ASAP were used to produce Figures 3D-E (Gardeux *et al.*,

2017). Continuous Coloring of a Custom Gene Set - Categorical Gene Metadata:

_Chromosomes (either X or Y) was used under the Visualization tab. SVGs can be saved

directly from the website. The enrichment analysis performed to produce these graphs mirrors

Seurat's AddModuleScore function and is detailed in the Materials and Methods of Li et al.,

2022 (see header: Metabolic clustering using ASAP).

1064

On occasion, use of an alternative Assay ("log.counts") in the Seurat Object allowed for

visualization of low levels of gene expression in spermatocytes. This "log.counts" assay

contains a matrix of $\log_2(\text{counts} + 1)$, or log-transformed raw counts. This was done for plotting

or to perform analyses focused on promiscuous expression in spermatocytes (Figures 3F-G;

Figure 3 - figure supplement 2A-D,G,I; Figure 3 - source data 1). Figure 3 code describes how

this assay was added and shows how to utilize this information when needed.

1071

Cell type-specific transcription factors were taken from Li et al., 2022, Table S3 and are

available at https://doi.org/10.5061/dryad.m63xsj454 (TFs_list_500.txt). Each gene was

classified as being scored as expressed only in mature spermatocytes, both in other cell types

and mature spermatocytes, or not in germ line cell types as per assignments in the Fly Cell

Atlas Table S3 (Li et al., 2022). Figure 3 - source data 1 was generated by calculating average

log-transformed expression of each transcription factor per Leiden 0.4 cluster. Subtracting the

value in cluster 3 from that of cluster 2 yielded the upregulation of each gene in mature

spermatocytes relative to spermatogonia and early spermatocytes (Figure 3 - source data 1).

The DotPlot in Figure 3I, in contrast, utilized only the testis and seminal vesicle data to

summarize the phenomenon seen with the Fly Cell Atlas data using just the nuclei derived from

the dissected testis and seminal vesicle sample. A subset of clusters for the cell types of interest

at Leiden 0.4, including clusters 2 and 3 as mentioned above, were used to generate the plot.

1084

Tissue isolation and cell dissociation for scRNA-seq

The testis dissociation protocol was adapted from Witt et al., 2019. Fresh maceration buffer

(10mL Trypsin LE (Gibco) with 20 mg collagenase (Gibco)) was prepared on the day of the

dissection. Testes were hand dissected from 1-5 day old male flies in 1x phosphate-buffered

saline (PBS), separated from seminal vesicles and transferred immediately into tubes filled with

cold PBS, on ice. Testes were kept in PBS for a maximum of 30 minutes. Samples were

centrifuged at 135 rcf and the PBS removed and replaced with 400 µL maceration buffer.

Testes were incubated in maceration buffer for 30 minutes with gentle vortexing every 10

minutes at room temperature. Following incubation, samples were pipetted up and down for 15

minutes until all visible chunks were gone and the sample was in approximately a single-cell

suspension. Sample was filtered through a 35 µm filter into a polystyrene tube, then transferred

into a microcentrifuge tube. After the sample was centrifuged at 135 rcf for 7 minutes, the

supernatant was removed and the pellet resuspended in 1mL calcium- and magnesium-free

Hanks' Balanced Salt Solution (HBSS). The sample was spun a final time at 137 rcf for 7

minutes. All but 50 µL of the HBSS supernatant was removed, and the cell pellet was

resuspended in the remaining 50 µL. Cell viability and density was then assayed on a

hemocytometer using DIC imaging and Trypan Blue stain.

1102

1103   *Library preparation and sequencing for scRNA-seq*

1104   Cells were processed using the 10x Genomics Chromium Controller and Chromium Single Cell

1105   Library and Gel Bead Kit following standard manufacturer's protocol. Amplified cDNA libraries

1106   were quantified by bioanalyzer and size selected using AMPure beads. Samples were

1107   sequenced on a NovaSeq SP.

1108

Mapping and preprocessing of scRNA-seq data

Reads were mapped to the DM3 reference genome using the 10X CellRanger pipeline with

default parameters. The resulting feature matrix (default output, kept in

outs/filtered_feature_bc_matrix, and featuring barcodes.tsv.gz, features.tsv.gz, and

matrix.mtx.gz) was read into the R package Monocle3 using load_cellranger_data. The resulting

cell data set (cds) object was processed using 100 dimensions, and underwent dimensionality

reduction using the UMAP method. Germline cells were identified on the basis of super-cluster

(in Monocle3, "partition") identity, with 100 dimensions used to identify partitions. The germline

cells were then subsequently clustered using the Monocle3 "cluster_cells" command, with

resolution = 0.003.

1119

Trajectory analysis of FCA snRNA-seq data

The publicly available "Testis, 10X, Relaxed" loom file of snRNA-seq data from the Fly Cell Atlas

website (https://flycellatlas.org/; 10.1101/2021.07.04.451050) was read into Monocle3, and

preprocessed using 50 dimensions. This dimension number was determined empirically as it

resulted in connected clusters that represented the primary lineages (germline and cyst). UMAP

dimensionality reduction and clustering was performed with a resolution of 0.0002, again

determined empirically to represent biologically significant clusters that approximated the

original annotation. A trajectory graph was generated from data with "learn_graph". Pseudotime

was calculated with "order_cells", with the first (base) node selected as the root in Monocle3's

interactive mode. Pseudotime parameters were then subsequently visualized on the original

projection by adding a "pseudotime" slot to the FCA Seurat object. "learn_graph" and

"order_cells" were likewise run on the scRNA-seq dataset, with the single most base node again

selected as the root.

1133

A list of genes that change expression level dynamically along pseudotime was generated from

the full join of genes that vary along pseudotime in the single cell and nucleus datasets,

according to the graph_test function in Monocle3 (parameters: neighbor_graph =

"principal_graph", method = "Moran_I" cores = 4; selected genes with q_value = 0 and morans_I

= 0.25). Additionally, several genes encoding transcripts that were annotated as enriched in late

pseudotime in the original FCA paper were added to the analysis. Genes (representing rows on

the Figure 4G-H heatmaps) were ordered according to pseudotime point of peak expression

averaged between the two datasets. A z-score for each gene expression for each dataset,

smoothed across pseudotime using R function "smooth.spline" with 3 degrees of freedom was

calculated and the heatmap was generated using the R package ComplexHeatmap.

1144

Aligning trajectories for scRNA-seq and FCA snRNA-seq data by dynamic time warping

The trajectories for the scRNA-seq and snRNA-seq data were aligned on a common "warped"

time scale using a Dynamic Time Warping based procedure (adapted from Alpert et al., 2018;

Cacchiarelli et al., 2018). The smoothened gene expression (as shown in the heatmaps) for all

germline cells in each of the two monocle3 trajectories was used for alignment using the dtw

function in R package dtw. The Pearson's correlation based distance and "symmetric2" step

parameters were used in dtw. The aligned time scale returned by dtw was used as the warped

pseudotime. Transcript expression levels could then be plotted on the same axis.

1153

Code and Data Availability

Scripts written in the programming language R, necessary to reproduce the analyses in this

manuscript are contained in a folder hosted at:

https://doi.org/10.5061/dryad.m63xsj454. All additional data files, including the Seurat object for

the snRNA-seq data (FCAloomToSeurat2TFP_Annotations.rds), the Monocle3 object for the

scRNA-seq data (single_cell_monocle_obj.Robj), files for the warped pseudotime data, the full

transcription factor list, and the 3D projection of the cyst cell lineage, as well as the original

mapped gene expression matrix for the scRNA-seq data, are available at

https://doi.org/10.5061/dryad.m63xsj454. Both the raw and the filtered data from the FCA

snRNA-seq analysis are publicly available as indicated in the Results section of the text. The

raw and processed scRNAseq data can be found at the GEO accession GSE220615.

1165

Acknowledgements

We especially acknowledge and thank the Fly Cell Atlas project, the Chan-Zuckerberg Biohub,

the NIH sequencing facility and the laboratory of Steve Quake for the starting testis plus seminal

vesicle snRNA-seq dataset. We thank Stein Aerts and Bart Deplancke for supporting the efforts

of J.J. and W.S., respectively. We acknowledge the Whitehead Institute Genome Technology

Core for assistance in library preparation and sequencing of the scRNA-seq dataset, the

Bloomington Drosophila Stock Center and Kyoto DGGR for *Drosophila* strains, the

Developmental Studies Hybridoma Bank for various antibodies, and Aaron Allen for advice and

initial code to transfer the SCope loom file format into SEURAT. S.D. thanks Dan Beiting at

PennVet for his DIY_Transcriptomics course.

1176

1177   Competing Interests

1178   No competing interests declared

1179

Figure Legends

Figure 1 with one supplement

The snRNA-seq landscape of the testis

A) Illustration of adult *Drosophila* testis showing hub (green), germ cell lineage (blue), cyst cell lineage (yellow), terminal epithelium (pink) and seminal vesicle (dark gray). **B)** UMAP of FCA snRNAseq data from the testis plus seminal vesicle (relaxed version). Blue: germ cell lineage; Yellow: cyst cell lineage; Pink: terminal epithelial cells of testis (te); Dark gray: seminal vesicle (sv). Other cell types as listed in **G**. **(C-F)** UMAP plots of snRNA-seq data showing expression of: **C)** *vasa* (*vas*), **D)** traffic *jam* (*tj*), **E)** *string* (*stg*), **F)** *escargot* (*esg*). Red arrows: proliferating cells. Red arrowhead: hub. **G)** UMAP (as in **B**) with Leiden 6.0 clusters of germ and cyst cell lineages labeled (sg: Spermatogonia; mg: Mid-late proliferating spermatogonia; sp: Spermatocytes; s: Spermatids; es: Early elongation-stage spermatids; ms: Early-mid elongation stage spermatids; ls: Mid-late elongation-stage spermatids; hb: Germinal proliferation center hub; cs: Cyst stem cells, c1: Early cyst cells 1; c2: Early cyst cells 2; sa: Cyst cell with spermatocytes branch A; sb: Cyst cell with spermatocytes branch B; ca: Cyst cell branch a; cb: Cyst cell branch b; sac: Elongating spermatid-associated cyst cells; hcc: Head cyst cells; tcc: Tail cyst cells; te: Terminal epithelial cells of testis; sv: Seminal vesicle; ep: Male gonad associated epithelium; sr: Secretory cells of the male reproductive tract; mc: Muscle cells; hm: Hemocytes; nr: Neurons; pc: Pigment cells;, tr: Trachea; ft: Fat body). (see also Figure 1 – figure supplement 1).

Figure 2 with one supplement

Characteristics of the germline lineage

A) Germline portion of the UMAP generated by Seurat from clustering of the full testis plus

seminal vesicle dataset at Leiden 6.0 resolution. **B-C)** Apical tips of testes showing localized

expression of **B)** *aub* (magenta) and *fzo* (yellow) mRNA and **C)** *kmg* mRNA visualized by *in situ*

hybridization. Apical-most dotted line demarcates germ line stem cells (GSCs) around the hub

from spermatogonia. Lower dotted line demarcates spermatogonia and cells in premeiotic S

phase from young spermatocytes. Scale bars, 10 µm. **D-K)** Feature plots generated by Seurat

showing expression levels of *zpg, aub, kmg, rbp4, aly, CycB, fzo,* and *p-cup* in the germline

UMAP. Navy blue gradient bars: relative expression level for the indicated gene. **L)** Dot plot

generated by Seurat showing expression levels of selected germline markers by cluster as

nuclei progress from spermatogonia to spermatid. Color intensity: level of expression of the

indicated gene averaged over all the nuclei in a given cluster relative to the level for that gene in

other germline clusters. Size of dots: percent of nuclei in specified cluster in which expression

of the gene was detected. (see also Figure 2 - figure supplement 1).

1219

1220

1221

Figure 3 with two supplements

Features of the spermatocyte transcription program

(A,B) UMAPs of snRNA-seq data showing: **A)** number of genes detected as expressed and **B)**

Number of unique molecular identifiers (UMIs) detected per nucleus. **(C)** Plot of average

number of genes expressed (dot size) and UMIs detected per nucleus per germline-annotated

Leiden 6.0 cluster, ordered by estimated progression of germ line differentiation. **(D,E)** UMAPs

of snRNA-seq data showing average expression of: **D)** Y chromosome tor **E)** X chromosome

encoded transcripts relative to an expression-matched control set (gene sets with binned

expression matching transcript lists). (Note the dramatically different relative expression scales

in **D** vs **E**.) UMAPs taken directly from ASAP (Li *et al.*, 2022, Gardeux *et al.*, 2017). **(F,G)** UMAP

plots of snRNA-seq raw counts (log-transformed) showing expression of: **F)** *Hemolectin* (*Hml*) in

hemocytes and late spermatocytes and **G)** Myosin *heavy chain* (*Mhc*) in muscle and late

spermatocytes. Yellow: relative expression high. **H)** Testis hybridized *in situ* with biotinylated

antisense RNA probe to *Hml*, showing expression (blue) in spermatocytes. **I)** Dot Plot showing

expression of pairs of tissue specific markers across cell types in the testis plus seminal vesicle

snRNA-seq sample. Average expression of each gene in a given cell type (here, a cluster at

Leiden 0.4 resolution) denoted by color intensity. Percent of nuclei of the given cell type scored

as expressing each gene denoted by dot size. Colors along axes (see Figure **1B**) indicate the

relevant cell type for tissue-specific marker pairs. (see also Figure 3 - figure supplements 1,2

and Figure 3 – source data 1).

1242

1243

1244

Figure 4 with one supplement

Developmental transitions revealed by comparing sn and scRNA-seq

(A-D) UMAP plots of germline-annotated data from **A,B)** FCA snRNA-seq of adult testis plus

seminal vesicle and **C,D)** whole-cell scRNA-seq of adult testis. **A,C)** Color denotes germline

differentiation stage, with clustering in A as in Figure 4 - figure supplement 1. **B,D)** Color

denotes pseudotime, with the few nuclei lacking a calculable pseudotime value colored gray. **E)**

UMAPs of scRNA-seq data showing log10(Expression) levels of cell-stage diagnostic markers

zpg, *aub*, *fzo*, and *p-cup*. **F)** FISH of diagnostic genes *aub* and *fzo*. . Scale bar, 50µm. Yellow

arrowhead in **C,F,H)** marks the point at which *fzo* transcript drops, annotated as early round

spermatids based on **F. (G-H)** Heatmaps of row-normalized (z-score) gene expression over

pseudotime for: **G)** all germline-annotated single nuclei from panel A, **H)** all germline-annotated

single whole cells from panel C, with genes in same order as in G. X axes, pseudotime; Y axes,

genes. Vertical white line: nuclei (G) or cells (H) where level of *fzo* mRNA has dropped 50%

from peak to nadir (0 on Z score). *Fzo* transcript drop validated in the cell data, and marked with

yellow arrowhead. Top bars: cell identity for each column, colored as in panels A,C. Arrow

points to transition between clusters dominated by expression of genes up in spermatogonia to

clusters dominated by expression of genes upregulated in early spermatocytes (see Figure 4 -

figure supplement 1A,B). Gene markers as follows: black filled squares, transcripts validated in

Figure 2 - figure supplement 1. Arrowheads, *CycB* and *fzo*. Asterisks, *Mst77F* and *tomboy20*.

Circle, *l(2)41Ab*, validated in panel I. Black boxes: genes transcribed post-meiotically (see

Figure 5). **I)** FISH of *l(2)41Ab* intron (magenta) and *l(2)41Ab* exon (green). Arrowhead on whole

testis points to a round spermatid cyst, which lacks intronic signal. High magnification insets:

Side - Spermatocyte nucleus showing intronic and exonic probe signal. Bottom - elongating

spermatid cyst showing intronic probe signal only. Scalebars: whole testis, 50µm; spermatocyte,

5µm; spermatid, 20µm. **(J-K)** Comparison of gene expression over warped pseudotime for: **J)**

genes outlined by pink boxes in G,H;   **K)** genes outlined by green boxes in G,H. (see also

Figure 4 - figure supplement 1)

1272

1273

1274

Figure 5 with one supplement

The transcript landscape of haploid spermatids

A) Diagram of spermatid orientation in the testis at different stages of spermatid elongation.

Arrows: distal ends of spermatid cysts. (**B,C)** RNA FISH of representative transcripts (**B**: *f-cup*,

C: *loqs*) in whole testes showing different patterns of mRNA localization for post-meiotically

transcribed genes. DNA (magenta), target RNA (green). Asterisk: signal in spermatocytes as

surmised by relatively large nuclear profile revealed by absence of signal. Bar: 100µm. **D)**

smFISH for *wa-cup* (red) in whole testes. DNA (white). Bar: 100µm. (**E,F)** Left: smFISH for *soti*

E) or wa-cup **F**) in a single early elongating spermatid cyst (cyan dashed outline). RNA (red)

and DNA (white). Bar: 25µm. Right: Enlarged image of yellow dashed box showing spermatid

nuclei. Arrows: nuclear transcripts. Arrowheads: perinuclear granules. **G)** Dot plot for selected

spermatid transcribed genes showing expression levels in each germ cell cluster. **H**) Expression

over pseudotime for selected spermatid transcribed genes and fold change between late

pseudotime and previous expression maxima in early/mid pseudotime. (see also Figure 5 -

figure supplement 1).

1290

1291

1292

1293

Figure 6 with one supplement

Differentiation in the somatic cyst cell lineage

A) Cyst cell lineage portion of the UMAP from snRNA-seq data with Leiden 6.0 cluster

numbering. **B)** Expression of *tj* (blue), *eya* (orange), and *so* (red) projected on the UMAP

(heatmaps in Figure 6 - figure supplement 1C,D). **C)** 2D UMAP of cyst cell clusters newly

reprojected in their own gene expression space (Materials & Methods; note different axis

coordinates relative to panel A). Cluster colors correspond to panel A except for unidentifiable

clusters with low UMIs (gray). **D)** Schematic of cyst cell cluster progression from CySC, to

spermatogonia-associated (SgCC) to spermatocyte-associated (ScCC) to spermatid-associated

(SpCC) inferred from the 3D UMAP in panel C. Cluster numbers and colors as in A. Note two

splits, one earlier and one later in cyst lineage progression. **E)** Plot of average number of genes

(dot size) and average number of UMIs per cyst lineage-annotated cluster; with clusters ordered

by deduced progression of differentiation. Dotted and dashed lines under cluster numbers

represent the early split shown in D, while single (HCC) and double (TCC) solid lines represent

the later split. **F - T)** UMAPs compared to FISH (RNA), IF or live GFP emission (from tagged

protein) images. For each gene comparison, arrows, dashed arrows and/or arrowheads point

out the same cell type in the UMAP and its corresponding FISH and/ or GFP fluorescence

image. **F)** *piwi* mRNA expression projected on the UMAP. **(G,H)** *piwi* mRNA (FISH; green) and

DNA (magenta). **G**) whole testis, **H)** testis apical tip. **I)** Apical tip of testis expressing GFP

tagged Piwi protein. **J)** *eya* mRNA expression projected on the UMAP. **(K, L)** *eya* mRNA

(FISH; green) and DNA (magenta). **K)** Testis apical tip. **L)** Whole testis. Outline demarcates

cyst of spermatocytes showing *eya* transcript expression (see Figure 3 and its supplement). **M)**

Amph expression projected on the UMAP. **N)** Apical third of a testis expressing GFP-tagged

Amph protein**. O)** Schematic of head cyst cells (HCC, green solid outline) embedded (left) or

not embedded (right) in the terminal epithelium (TE, gray), with the tail cyst cell (TCC, green

dashed outline) extending away, containing either a pre-coiled (right) or coiled (left) spermatid

bundle (Sperm nuclei, magenta). Coracle (Cora, blue) marks the septate junction (SJ) between

TCC and HCC. **P)** *Nep4* mRNA expression projected on the UMAP. **Q)** Testis base showing

Nep4 mRNA (FISH, green), DNA (magenta), and Cora (septate junctions, cyan). Arrowhead

points to TCCs associated with coiled sperm. **R)** UMAP of *geko* mRNA expression. **S,T)** *geko*

mRNA (FISH, green), DNA (magenta), and Cora (septate junctions, cyan). **S)** Whole testis. **T)**

Testis base. Arrowhead points to TCCs associated with coiled sperm. **U)** Dotplot of gene

expression (Y axis) by cluster as cyst lineage progresses through differentiation (X axis, left to

right). Averaged expression in each cluster indicated by color scale. Percent of cells within a

cluster expressing the gene indicated by size of dot. Lines under cluster numbers as in E.

Asterisk denotes hub. Bars: 20μm. (see also Figure 6 - figure supplement 1)

1330

Figure 7 with two supplements

Characteristics of the Hub

(A-E) Expression of the indicated genes (*upd1*, *hh*, *CadN*, *Fas3*, *tup*) projected onto the testis

snRNA-seq UMAP with the 79 definitive hub nuclei outlined (small red box), with reprojection of

hub nuclei (larger red box). Color intensity corresponds to expression level, shown as

normalized average logFC.   Red dashed circles contain non-hub nuclei of cluster 90 (see text

and Figure S5). **F-F''')** Apical tip of adult testis carrying a Tup::GFP fusion transgene revealing

protein expression largely restricted to hub nuclei delimited by Fas3 (magenta). Nuclei (blue).

Scale bar: 10µm. **G-J)** Paired Venn diagrams, comparing up-regulated genes in the hub vs.

clusters containing either CySCs, spermatogonia, hemocytes or terminal epithelial cells.

Overlap in upregulated genes was greatest between Hub and CySCs. In each pairing, circle

size reflects the number of genes compared for each cluster. Genes are listed in Figure 7 -

source data 1; the specific genes for hub vs. CySCs in Figure 7 - source data 2. **K)** Dot plot of

expression of selected genes comparing hub to CySCs, the early Cyst lineage, Spermatogonia,

Hemocytes, and Terminal epithelium. (see also Figure 7 - figure supplements 1, 2).

1346

1347

1348

Figure 8 with one supplement

Characteristics of supporting epithelia

A) UMAP of non cyst cell lineage epithelial cells of the testis from the FCA snRNA-seq data with

Leiden 6.0 clusters. Note: identity of cluster 89 remains undefined.   **B - G)** UMAPs compared to

ISH images. For each comparison, arrows and/or arrowheads point out the same cell type in

the UMAP and its corresponding ISH image. **B)** *hh* mRNA expression projected on the UMAP.

C) FISH of *hh* mRNA (green) and DNA (magenta) showing the base of the testis including the

TE (arrowhead), testicular duct (TD), and SV (arrow). **D)** *MtnA* mRNA expression projected on

the UMAP. **E)** Colorimetric *in situ* hybridization of *MtnA* mRNA (blue) in an entire testis plus

seminal vesicle. Asterisk denotes hub. **F)** *Nep5* mRNA expression projected on the UMAP. **G)**

FISH of *Nep5* mRNA (green) and DNA (magenta) showing the SV. **H**) *Nep4* (blue) and *Nep5*

(red) mRNA expression projected on the UMAP with corresponding heatmap. Arrows and

arrowheads point to corresponding cell types in UMAPs and stained tissues.   Bars: 20μm. (see

also Figure 8 - figure supplement 1).

1363

Supplementary Figure Legends

Figure 1 – figure supplement 1
Expression of *vasa* in the snRNA-seq dataset
Violin plot of *vasa* expression in nuclei grouped and colored by annotation, sorted so that all germ line nuclei are on the left half of the plot. Numbers at the point for each cell type reflect the proportion of nuclei that expressed *vasa* at a level greater than 0.

Figure 2 – figure supplement 1
Comparison of snRNA-seq data clustered at Leiden 6.0 vs. lower resolution and new marker genes
A,B) UMAP clustered at **A)** Leiden 6.0 and **B)** Leiden 0.8. **C)** Dot Plot of selected marker gene expression per Leiden 0.8 cluster (compare to Leiden 6.0 in Figure 2L). Dot color intensity: Average expression level in all nuclei of cluster. Dot size: percent of cells in the cluster expressing the marker. **D)** Novel markers of early germ cells (top panels) and spermatocytes (middle panels), represented on UMAP of the germline-annotated nuclei. These four genes are marked with black squares in Figure 4G,H. Dot Plot (bottom panel) of novel marker gene expression per Leiden 0.8 cluster as in C. **E)** FISH of markers in **D** confirms expression patterns. *Pxt* and *Drep2* are expressed in early germ cells; *CG43317* and *Vha16-5* are expressed in spermatocytes. Scale bar, 10 microns.

Figure 3 – figure supplement 1
Magnitude of the mid-to-late spermatocytes transcription program
A) Average UMI and numbers of genes expressed per cluster (ordered by cluster number In Li, 2022) inFCA snRNA-seq data from testis, malpighian tubules, heart, and male reproductive tract. Dot size proportional to the number of genes detected as expressed (aveGenes). Note difference between panels in dot size scale bars. Across these tissues, Mid-to-late spermatocytes had the highest average UMI and number of genes expressed per cluster. The highest fat body signal was found to be due to contaminating spermatocytes, and is therefore not relevant here. In the FCA dataset from Li, 2022, Mid-to-late spermatocytes are represented by clusters 18, 29, 30, 31, 32; in our dataset, these correspond to clusters 51, 35, 33, 48, and some of 105 (Figure 2A and Figure 2 – figure supplement 1A). **B)** Germ line portion of UMAP from single cell sequencing (scRNAseq) of adult testes showing number of genes detected as expressed per cell. **C)** Violin plot of average number of UMIs per cell type for scRNA-seq data.

Figure 3 – figure supplement 2
Markers of other tissues also expressed in mid-to-late spermatocytes
(**A-D**) UMAPs of snRNA-seq raw counts (log-transformed) showing expression of: **A)** *Synaptotagmin 1* (*Syt1*); **B)** *grainy head* (*grh*); **C)** *eyes absent* (*eya*); **D)** *unpaired 1* (*upd1*). Yellow: relative expression high. **E)** Dot Plots showing expression of *Mhc*, *Hml*, *Syt1* and *grh* in spermatocytes detected in single cell RNA-seq (scRNA-seq) analysis of adult testes. **F)** Testis from a male mutant for the tMAC subunit *comr* hybridized *in situ* with biotinylated antisense RNA probe to *Hml*, showing expression in spermatocytes. **G)** UMAP of snRNA-seq raw counts (log-transformed) showing expression of *geko*. **H)** Testis showing *Visual system homeobox 1* (*Vsx1*) mRNA (FISH, green), Coracle (Cora) protein (red), and DNA (blue). White arrows:

1407　somatic cyst cells, white arrowheads: spermatocytes. **I**) UMAP of snRNA-seq raw counts (log-
1408　transformed) showing expression of *Vsx1*.
1409
1410　**Figure 4 – figure supplement 1**
1411　**Developmental transitions revealed by trajectory analysis**
1412　**A)** UMAP plot of germline-annotated data, as in Fig 4A, clustered using Monocle3. Colors and
1413　clusters same as in Fig 4. Unclustered nuclei labeled in grey. **B)** Dotplot showing transcriptional
1414　signatures of diagnostic marker genes in clustersdefined as in (**A**).
1415
Figure 5 – figure supplement 1
Reactivation of transcription in spermatids
(A-D) UMAP plots of *f-cup, loqs*, *wa-cup* and *soti*. Arrow: latest point in pseudotime. **E)** Scatter
plot showing that average Pearson correlation coefficient (measurement of how likely each
spermatid transcribed gene is to be expressed with every other spermatid transcribed gene) is
positively correlated with the average expression level (average expression of each spermatid
transcribed gene among spermatid clusters).
1423
Figure 6 – figure supplement 1
Cyst Lineage pseudotime and re-projection
A) UMAP of FCA testis plus seminal vesicle snRNA-seq data with cyst cell lineage colored by
pseudotime (scale: origin (purple) to end (yellow)). Nuclei in dark gray have been determined to
be a part of the cyst lineage, but did not have a calculable pseudotime value (see main text).
Nuclei in light gray are not a part of the cyst cell lineage and did not have a calculable
pseudotime value related to cyst cells. **B)** UMAP of cyst cell clusters re-projected in their own
gene expression space (3D representation available at
https://doi.org/10.5061/dryad.m63xsj454). Color denotes pseudotime (scale on right). Nuclei in
gray were designated as part of the cyst cell lineage but did not have a calculable pseudotime
value. **C)** *tj* (blue) and *eya* (orange) mRNA expression projected on the UMAP with
corresponding heatmap (refer to Figure 6B). **D)** *eya* (orange) and *so* (red) mRNA expression
projected on the UMAP with corresponding heatmap (refer to Figure 6B). **E - I)** UMAPs
compared to IF images. For each gene comparison, arrows, dashed arrows and/or arrowheads
point out the same cell type in the UMAP and its corresponding IF image. **E)** *CG3902* mRNA
expression projected on the UMAP. **F)** Apical tip of testis expressing GFP protein trap of
CG3902. **G)** *Akr1B* mRNA expression projected on the UMAP. **H-I)** GFP protein trap of Akr1B.
H) whole testis, and **I**) testis apical third view. Asterisk denotes hub. Bars: 20µm.
1442
Figure 7 – figure supplement 1
Hub justification using markers
Justification for excluding 41 nuclei originally assigned to cluster 90. **A)** Cluster 90 nuclei
(blue). Red box: 79 nuclei of the definitive hub. Solid and dashed red circles: 4 other spatially
separate domains of original cluster 90. Dashed circles: left wing (14 nuclei), right wing (21
nuclei). **B)** Expression of Rbp4, a gene highly expressed in the germline, and in both wings but
barely detected in the definitive hub. (**C-E**) Expression of three genes **C)** *zpg*, **D)** *p53*, **E)** *vas*,
all highly expressed in the germ line, detected in the right wing but barely detected in the
definitive hub. Scale bar: normalized logFC values.

Figure 7 – figure supplement 2
Further support for Hub identity using alternate clustering methodology
(A) Graph-based subclustering of the 120 nuclei originally assigned to cluster 90. (Top) Nuclei
colored according to cluster identity, based on highly expressed and highly variable genes in
those 120 nuclei. (Bottom) "True Hub" (ie, the 79 definitive hub nuclei) are colored in red. "True
Hub" annotation almost perfectly overlaps with cluster 0 in the graph-based subclustering. **(B)** A
dendrogram derided from hierarchical clustering of the 120 nuclei originally assigned to cluster
90. The 79 definitive hub nuclei largely cluster together (red text). Nuclei are named by their cell
barcode, with the suffix "_HUB" added to the 79 hub nuclei. The bolded lines indicate a single
clade composed entirely of definitive hub nuclei.

Figure 8 – figure supplement 1
RNA expression in non cyst cell testis epithelia
(A-C) UMAP of "mermaid" showing RNA counts (rather than normalized expression) for the
indicated marker genes: **A)** *hh*, **B)** *MtnA*, **C)** *Nep5*). Arrows and arrowheads point to the same
region on the corresponding UMAP in Figure 8B,D,F.
1467

List of source data files:

1469 A 'readMe' tab in each file describes the purpose and structure of that source.
1470
1471 Figure 3 - source data 1: Average expression of 496 transcription factors in Leiden 0.4 clusters from Li
1472 et al., 2022.
1473
1474 Figure 5 - source data 1: List of 162 spermatid transcribed genes.
1475
1476 Figure 7 - source data 1: Differentially-expressed genes in Hub, CySCs, Spermatogonia, Terminal
1477 epithelium, or Hemocytes.
1478
1479 Figure 7 - source data 2: Differentially-expressed gene comparison between Hub and CySC .
1480

References

Alpert, A., Moore, L.S., Dubovik, T., and Shen-Orr, S.S. (2018). Alignment of single-cell
trajectories to compare cellular expression dynamics. Nat. Methods *15*, 267–270.

Alphey, L., Jimenez, J., White-Cooper, H., Dawson, I., Nurse, P., and Glover, D.M. (1992).
twine, a cdc25 homolog that functions in the male and female germline of Drosophila. Cell *69*,
977–988.

Amoyel, M., Sanny, J., Burel, M., and Bach, E.A. (2013). Hedgehog is required for CySC self-
renewal but does not contribute to the GSC niche in the Drosophila testis. Development *140*,
56–65.

Anllo, L., and DiNardo, S. (2022). Visceral mesoderm signaling regulates assembly position and
function of the Drosophila testis niche. BioRxiv 2021.09.08.459436.

Anllo, L., Plasschaert, L.W., Sui, J., and DiNardo, S. (2019). Live imaging reveals hub cell
assembly and compaction dynamics during morphogenesis of the Drosophila testis niche. Dev.
Biol. *446*, 102–118.

Barckmann, B., Chen, X., Kaiser, S., Jayaramaiah-Raja, S., Rathke, C., Dottermusch-Heidel,
C., Fuller, M.T., and Renkawitz-Pohl, R. (2013). Three levels of regulation lead to protamine and
Mst77F expression in Drosophila. Dev. Biol. *377*, 33–45.

Barreau, C., Benson, E., Gudmannsdottir, E., Newton, F., and White-Cooper, H. (2008). Post-
meiotic transcription in Drosophila testes. Development *135*, 1897–1902.

Boukhatmi, H., Frendo, J.L., Enriquez, J., Crozatier, M., Dubois, L., and Vincent, A. (2012).
Tup/Islet1 integrates time and position to specify muscle identity in Drosophila. Development
139, 3572–3582.

Boyle, M., Wong, C., Rocha, M., and Jones, D.L. (2007). Decline in Self-Renewal Factors
Contributes to Aging of the Stem Cell Niche in the Drosophila Testis. Cell Stem Cell *1*, 470–478.

1505

Brower, D.L., Smith, R.J., and Wilcox, M. (1981). Differentiation within the gonads of Drosophila
revealed by immunofluorescence. J. Embryol. Exp. Morphol. *63*, 233–242.

Cacchiarelli, D., Qui, X., Mikkelsent, T.S., Rinn, J.L., and Trapnell, C. (2018). Aligning Single-
Cell Developmental and Reprogramming Trajectories Identifies Molecular Determinants of
Myogenic Reprogramming Outcome. Cell Syst. 258–268.

Cao, C., Ma, Q., Mo, S., Shu, G., Liu, Q., Ye, J., and Gui, Y. (2021). Single-Cell RNA
Sequencing Defines the Regulation of Spermatogenesis by Sertoli-Cell Androgen Signaling .
Front. Cell Dev. Biol.   *9*.

Cao, J., Packer, J., Ramani, V., Cusanovich, D., Chau, H., Riza, D., Xiaojie, Q., Choli, L., N.,
F.S., J., S.F., et al. (2017). Comprehensive single-cell transcriptional profiling of a multicellular
organism. Science *357*, 661–667.

Cao, J., Spielmann, M., Qiu, X., Huang, X., Ibrahim, D.M., Hill, A.J., Zhang, F., Mundlos, S.,
Christiansen, L., Steemers, F.J., et al. (2019). The single-cell transcriptional landscape of
mammalian organogenesis. Nature *566*, 496–502.

Chandley, A.C., and Bateman, A.J. (1962). Timing of Spermatogenesis in Drosophila
melanogaster using Tritiated Thymidine. Nature *193*, 299–300.

Chen, Y., Zheng, Y., Gao, Y., Lin, Z., Yang, S., Wang, T., Wang, Q., Xie, N., Hua, R., Liu, M., et
al. (2018). Single-cell RNA-seq uncovers dynamic processes and critical regulators in mouse
spermatogenesis. Cell Res. *28*, 879–896.

DeFalco, T., Camara, N., Le Bras, S., and Van Doren, M. (2008). Nonautonomous Sex
Determination Controls Sexually Dimorphic Development of the Drosophila Gonad. Dev. Cell
14, 275–286.

DiNardo, S., Okegbe, T., Wingert, L., Freilich, S., and Terry, N. (2011). lines and bowl affect the
specification of cyst stem cells and niche cells in the Drosophila testis. Development *138*, 1687–
1696.

Edgar, B.A., and O'Farrell, P.H. (1990). The three postblastoderm cell cycles of Drosophila
embryogenesis are regulated in G2 by string. Cell *62*, 469–480.

Fabrizio, J.J., Boyle, M., and DiNardo, S. (2003). A somatic role for eyes absent (eya) and sine
oculis (so) in drosophila spermatocyte development. Dev. Biol. *258,* 117–128.

Fairchild, M.J., Smendziuk, C.M., and Tanentzapf, G. (2015). A somatic permeability barrier
around the germline is essential for Drosophila spermatogenesis. Development *142*, 268–281.

Fincher, C.T., Wurtzel, O., de Hoog, T., Kravarik, K.M., and Reddien, P.W. (2018). Cell type
transcriptome atlas for the planarian Schmidtea mediterranea. Science *1736*, 1–19.

Fingerhut, J.M., and Yamashita, Y.M. (2020). mRNA localization mediates maturation of
cytoplasmic cilia in Drosophila spermatogenesis. J. Cell Biol. *219*, e202003084.

Fingerhut, J.M., Moran, J. V., and Yamashita, Y.M. (2019). Satellite DNA-containing gigantic
introns in a unique gene expression program during Drosophila spermatogenesis. PLoS Genet.
15, 1–23.

Fuller, M.T. (1993). Spermatogenesis, in The Development of Drosophila (Cold Spring Harbor,
NY: Cold Spring Harbor Laboratory Press).

Fuller, M.T. (1998). Genetic control of cell proliferation and differentiation in Drosophila
spermatogenesis. Semin. Cell Dev. Biol. *9*, 433–444.

Fuse, N., Hirose, S., and Hayashi, S. (1994). Diploidy of Drosophila imaginal cells is maintained
by a transcriptional repressor encoded by escargot. Genes Dev. *8*, 2270–2281.

Gan, Q., Chepelev, I., Wei, G., Tarayrah, L., Cui, K., Zhao, K., and Chen, X. (2010). Dynamic
regulation of alternative splicing and chromatin structure in Drosophila gonads revealed by
RNA-seq. Cell Res. *20*, 763–783.

Ghosh-Roy, A., Kulkarni, M., Kumar, V., Shirolikar, S., and Ray, K. (2004). Cytoplasmic Dynein–
Dynactin Complex Is Required for Spermatid Growth but Not Axoneme Assembly in Drosophila.

Mol. Biol. Cell *15*, 2470–2483.

Gönczy, P., Viswanathan, S., and DiNardo, S. (1992). Probing spermatogenesis in Drosophila
with P-element enhancer detectors. Development *114*, 89–98.

Gonzalez, J., Qi, H., Liu, N., and Lin, H. (2015). Piwi Is a Key Regulator of Both Somatic and
Germline Stem Cells in the Drosophila Testis. Cell Rep. *12*, 150–161.

Gould-Somero, M., and Holland, L. (1974). The timing of RNA synthesis for spermiogenesis in
organ cultures ofDrosophila melanogaster testes. Wilhelm Roux. Arch. Entwickl. Mech. Org.
174, 133–148.

Green, C.D., Ma, Q., Manske, G.L., Shami, A.N., Zheng, X., Marini, S., Moritz, L., Sultan, C.,
Gurczynski, S.J., Moore, B.B., et al. (2018). A Comprehensive Roadmap of Murine
Spermatogenesis Defined by Single-Cell RNA-Seq. Dev. Cell *46*, 651-667.e10.

Greenspan, L.J., de Cuevas, M., Le, K.H., Viveiros, J.M., and Matunis, E.L. (2022). Activation of
the EGFR/MAPK pathway drives transdifferentiation of quiescent niche cells to stem cells in the
Drosophila testis niche. Elife *11*, e70810.

Guo, J., Nie, X., Giebler, M., Mlcochova, H., Wang, Y., Grow, E.J., Kim, R., Tharmalingam, M.,
Matilionyte, G., Lindskog, C., et al. (2020). The Dynamic Transcriptional Cell Atlas of Testis
Development during Human Puberty. Cell Stem Cell *26*, 262-276.e4.

Haber, A.L., Biton, M., Rogel, N., Herbst, R.H., Shekhar, K., Smillie, C., Burgin, G., Delorey,
T.M., Howitt, M.R., Katz, Y., et al. (2017). A single-cell survey of the small intestinal epithelium.
Nature *551*, 333–339.

Hardy, R.W., Tokuyasu, K.T., Lindsley, D.L., and Garavito, M. (1979). The germinal proliferation
center in the testis of Drosophila melanogaster. J. Ultrastruct. Res. *69*, 180–190.

Herrera, S.C., Sainz de la Maza, D., Grmai, L., Margolis, S., Plessel, R., Burel, M., O'Connor,
M., Amoyel, M., and Bach, E.A. (2021). Proliferative stem cells maintain quiescence of their
niche by secreting the Activin inhibitor Follistatin. Dev. Cell *56*, 2284-2294.e6.

Hétié, P., de Cuevas, M., and Matunis, E. (2014). Conversion of Quiescent Niche Cells to

Somatic Stem Cells Causes Ectopic Niche Formation in the Drosophila Testis. Cell Rep. *7*,

715–721.

Hof-Michel, S., and Bökel, C. (2020). Transcriptome analysis of somatic cell populations in the

Drosophila testis links metabolism and stemness. BioRxiv 2020.02.24.962639.

Hwa, J.J., Hiller, M.A., Fuller, M.T., and Santel, A. (2002). Differential expression of the

Drosophila mitofusin genes fuzzy onions (fzo) and dmfn. Mech. Dev. *116*, 213–216.

Hwa, J.J., Zhu, A.J., Hiller, M.A., Kon, C.Y., Fuller, M.T., and Santel, A. (2004). Germ-line

specific variants of components of the mitochondrial outer membrane import machinery in

Drosophila. FEBS Lett. *572*, 141–146.

Jenkins, N.A., Kaumeyer, J.F., Young, E.M., and Raff, R.A. (1978). A test for masked message:

The template activity of messenger ribonucleoprotein particles isolated from sea urchin eggs.

Dev. Biol. *63*, 279–298.

Kaplan, Y., Gibbs-Bar, L., Kalifa, Y., Feinstein-Rotkopf, Y., and Arama, E. (2010). Gradients of a

Ubiquitin E3 Ligase Inhibitor and a Caspase Inhibitor Determine Differentiation or Death in

Spermatids. Dev. Cell *19*, 160–173.

Kapoor, T., Dubey, P., Shirolikar, S., and Ray, K. (2021). An actomyosin clamp assembled by

the Amphiphysin-Rho1-Dia/DAAM-Rok pathway reinforces somatic cell membrane folded

around spermatid heads. Cell Rep. *34*, 108918.

Kawase, E., Wong, M.D., Ding, B.C., and Xie, T. (2004). Gbb/Bmp signaling is essential for

maintaining germline stem cells and for repressing bam transcription in the Drosophila testis.

Development *131*, 1365–1375.

Kiger, A., Leanne, J.D., Cordula, S., Rogers, M., and Fuller, M. (2001). Stem Cell Self-Renewal

Specified by JAK-STAT Activation in Response to a Support Cell Cue. Science *294*, 2542–

2545.

Kudron, M.M., Victorsen, A., Gevirtzman, L., Hillier, L.W., Fisher, W.W., Vafeados, D., Kirkey,
M., Hammonds, A.S., Gersch, J., Ammouri, H., et al. (2018). The ModERN Resource: Genome-
Wide Binding Profiles for Hundreds of Drosophila and Caenorhabditis elegans Transcription
Factors. Genetics *208*, 937–949.

Kuhn, R., Kuhn, C., Börsch, D., Glätzer, K.H., Schäfer, U., and Schäfer, M. (1991). A cluster of
four genes selectively expressed in the male germ line of Drosophila melanogaster. Mech. Dev.
35, 143–151.

Law, N.C., Oatley, M.J., and Oatley, J.M. (2019). Developmental kinetics and transcriptome
dynamics of stem cell specification in the spermatogenic lineage. Nat. Commun. *10*, 2787.

Leatherman, J.L., and DiNardo, S. (2010). Germline self-renewal requires cyst stem cells and
stat regulates niche adhesion in Drosophila testes. Nat. Cell Biol. *12*, 806–811.

Le Bras, S., and Van Doren, M. (2006). Development of the male germline stem cell niche in
Drosophila. Dev. Biol. *294*, 92–103.

Li, H., Janssens, J., De Waegeneer, M., Sai, S.K., Davie, K., Gardeux, V., Saelens, W., Fabrice,
D., Brbić, M., Spanier, K., et al. (2022). Fly Cell Atlas: A single-nucleus transcriptomic atlas of
the adult fruit fly. Science *375*, eabk2432.

Li, M.A., Alls, J.D., Avancini, R.M., Koo, K., and Godt, D. (2003). The large Maf factor Traffic
Jam controls gonad morphogenesis in Drosophila. Nat. Cell Biol. *5*, 994–1000.

Liu, Y., Li, J.S.S., Rodiger, J., Comjean, A., Attrill, H., Antonazzo, G., Brown, N.H., Hu, Y., and
Perrimon, N. (2022). FlyPhoneDB: an integrated web-based resource for cell–cell
communication prediction in Drosophila. Genetics *220*, iyab235.

Lowe, N., Rees, J.S., Roote, J., Ryder, E., Armean, I.M., Johnson, G., Drummond, E., Spriggs,
H., Drummond, J., Magbanua, J.P., et al. (2014). Analysis of the expression patterns,
subcellular localisations and interaction partners of Drosophila proteins using a pigP protein trap
library. Development *141*, 3994–4005.

Lu, D., Sin, H.S., Lu, C., and Fuller, M.T. (2020). Developmental regulation of cell type-specific

transcription by novel promoter-proximal sequence elements. Genes Dev. *34*, 663–677.

Mahadevaraju, S., Fear, J.M., Akeju, M., Galletta, B.J., Pinheiro, M.M.L.S., Avelino, C.C., Cabral-de-Mello, D.C., Conlon, K., Dell'Orso, S., Demere, Z., et al. (2021). Dynamic sex chromosome expression in Drosophila male germ cells. Nat. Commun. *12*, 892.

McLaughlin, C.N., Brbić, M., Xie, Q., Li, T., Horns, F., Kolluru, S.S., Kebschull, J.M., Vacek, D., Xie, A., Li, J., et al. (2021). Single-cell transcriptomes of developing and adult olfactory receptor neurons in Drosophila. Elife *10*, e63856.

Michel, M., Kupinski, A.P., Raabe, I., and Bökel, C. (2012). Hh signalling is essential for somatic stem cell maintenance in the Drosophila testis niche. Development *139*, 2663–2669.

Morris, C.A., Benson, E., and White-Cooper, H. (2009). Determination of gene expression patterns using in situ hybridization to Drosophila testes. Nat. Protoc. *4*, 1807–1819.

Noguchi, T., Koizumi, M., and Hayashi, S. (2011). Sustained elongation of sperm tail promoted by local remodeling of giant mitochondria in Drosophila. Curr. Biol. *21*, 805–814.

Okegbe, T.C., and DiNardo, S. (2011). The endoderm specifies the mesodermal niche for the germline in Drosophila via Delta-Notch signaling. Development *138*, 1259–1267.

Olivieri, G., and Olivieri, A. (1965). Autoradiographic study of nucleic acid synthesis during spermatogenesis in Drosophila melanogaster. Mutat. Res. Mol. Mech. Mutagen. *2*, 366–380.

Papagiannouli, F., and Mechler, B.M. (2009). discs large regulates somatic cyst cell survival and expansion in Drosophila testis. Cell Res. *19*, 1139–1149.

Plass, M., Solana, J., Wolf, F.A., Ayoub, S., Misios, A., Glažar, P., Theis, F.J., Kocks, C., and Rajewsky, N. (2018). Cell type atlas and lineage tree of a whole complex animal by single-cell transcriptomics. *1723*, 1–17.

Qadir, M.M.F., Álvarez-Cubela, S., Klein, D., van Dijk, J., Muñiz-Anquela, R., Moreno-Hernández, Y.B., Lanzoni, G., Sadiq, S., Navarro-Rubio, B., García, M.T., et al. (2020). Single-

cell resolution analysis of the human pancreatic ductal progenitor cell niche. Proc. Natl. Acad.

Sci. *117,* 10876–10887.

Raja, J.S., and Renkawitz-Pohl, R. (2006). Replacement by Drosophila melanogaster

Protamines and Mst77F of Histones during Chromatin Condensation in Late Spermatids and

Role of Sesame in the Removal of These Proteins from the Male Pronucleus. Mol. Cell. Biol. *26,*

3682.

Renault, A.D. (2012). vasa is expressed in somatic cells of the embryonic gonad in a sex-

specific manner in Drosophila melanogaster. Biol. Open *1,* 1043–1048.

Rust, K., Byrnes, L.E., Yu, K.S., Park, J.S., Sneddon, J.B., Tward, A.D., and Nystul, T.G.

(2020). A single-cell atlas and lineage analysis of the adult Drosophila ovary. Nat. Commun. *11.*

Satija, R., Farrell, J.A., Gennert, D., Schier, A.F., and Regev, A. (2015). Spatial reconstruction

of single-cell gene expression data. Nat. Biotechnol. *33,* 495–502.

Schäfer, M., Kuhn, R., Bosse, F., and Schäfer, U. (1990). A conserved element in the leader

mediates post-meiotic translation as well as cytoplasmic polyadenylation of a Drosophila

spermatocyte mRNA. EMBO (Eur. Mol. Biol. Organ.) J. *9,* 4519–4525.

Schäfer, M., Nayernia, K., Engel, W., and Schäfer, U. (1995). Translational Control in

Spermatogenesis. Dev. Biol. *172,* 344–352.

Schaum, N., Karkanias, J., Neff, N.F., May, A.P., Quake, S.R., Wyss-Coray, T., Darmanis, S.,

Batson, J., Botvinnik, O., Chen, M.B., et al. (2018). Single-cell transcriptomics of 20 mouse

organs creates a Tabula Muris. Nature *562,* 367–372.

Sebé-Pedrós, A., Saudemont, B., Chomsky, E., Plessier, F., Mailhé, M.P., Renno, J., Loe-Mie,

Y., Lifshitz, A., Mukamel, Z., Schmutz, S., et al. (2018). Cnidarian Cell Type Diversity and

Regulation Revealed by Whole-Organism Single-Cell RNA-Seq. Cell *173,* 1520-1534.e20.

She, M., Tang, M., Jiang, T., and Zeng, Q. (2021). The Roles of the LIM Domain Proteins in

Drosophila Cardiac and Hematopoietic Morphogenesis   . Front. Cardiovasc. Med.   *8.*

Shi, Z., Lim, C., Tran, V., Cui, K., Zhao, K., and Chen, X. (2020). Single-cyst transcriptome

analysis of Drosophila male germline stem cell lineage. Development *147*, dev184259.

Shiraiwa, T., Nitasaka, E., and Yamazaki, T. (2000). Geko, a Novel Gene Involved in Olfaction

in Drosophila Melanogaster. J. Neurogenet. *14*, 145–164.

Shivdasani, A.A., and Ingham, P.W. (2003). Regulation of Stem Cell Maintenance and Transit

Amplifying Cell Proliferation by TGF-β Signaling in Drosophila Spermatogenesis. Curr. Biol. *13*,

2065–2072.

Siebert, S., Farrell, J. a., Cazet, J.F., Abeykoon, Y., Primack, A.S., Schnitzler, C.E., and Juliano,

C.E. (2019). Stem cell differentiation trajectories in Hydra resolved at single-cell resolution.

Science *365*, 1–8.

Sitnik, J.L., Francis, C., Hens, K., Huybrechts, R., Wolfner, M.F., and Callaerts, P. (2014).

Neprilysins: An Evolutionarily Conserved Family of Metalloproteases That Play Important Roles

in Reproduction in Drosophila. Genetics *196*, 781–797.

Soumillon, M., Necsulea, A., Weier, M., Brawand, D., Zhang, X., Gu, H., Barthès, P., Kokkinaki,

M., Nef, S., Gnirke, A., et al. (2013). Cellular Source and Mechanisms of High Transcriptome

Complexity in the Mammalian Testis. Cell Rep. *3*, 2179–2190.

Susic-Jung, L., Hornbruch-Freitag, C., Kuckwa, J., Rexer, K.-H., Lammel, U., and Renkawitz-

Pohl, R. (2012). Multinucleated smooth muscles and mononucleated as well as multinucleated

striated muscles develop during establishment of the male reproductive organs of Drosophila

melanogaster. Dev. Biol. *370*, 86–97.

Tanentzapf, G., Devenport, D., Godt, D., and Brown, N.H. (2007). Integrin-dependent anchoring

of a stem-cell niche. Nat. Cell Biol. *9*, 1413–1418.

Tazuke, S.I., Schulz, C., Gilboa, L., Fogarty, M., Mahowald, A.P., Guichet, A., Ephrussi, A.,

Wood, C.G., Lehmann, R., and Fuller, M.T. (2002). A germline-specific gap junction protein

required for survival of differentiating early germ cells. Development *129*, 2529–2539.

Thor, S., and Thomas, J.B. (1997). The Drosophila islet Gene Governs Axon Pathfinding and
Neurotransmitter Identity. Neuron *18*, 397–409.

Tokuyasu, K.T. (1975). Dynamics of spermiogenesis in Drosophila melanogaster. VI.
Significance of "onion" nebenkern formation. J. Ultrastruct. Res. *53*, 93–112.

Tokuyasu, K.T., Peacock, W.J., and Hardy, R.W. (1972). Dynamics of spermiogenesis in
Drosophila melanogaster. II. Coiling process. Z. Zellforsch. Mikrosk. Anat. *127*, 492–525.

Trapnell, C., Cacchiarelli, D., Grimsby, J., Pokharel, P., Li, S., Morse, M., Lennon, N.J., Livak,
K.J., Mikkelsen, T.S., and Rinn, J.L. (2014). The dynamics and regulators of cell fate decisions
are revealed by pseudotemporal ordering of single cells. Nat. Biotechnol. *32*, 381–386.

Tulina, N., and Matunis, E. (2001). Control of Stem Cell Self-Renewal in Drosophila
Spermatogenesis by JAK-STAT Signaling. Science *294*, 2546–2549.

Vedelek, V., Bodai, L., Grézal, G., Kovács, B., Boros, I.M., Laurinyecz, B., and Sinka, R. (2018).
Analysis of Drosophila melanogaster testis transcriptome. BMC Genomics *19*, 697.

Voog, J., Sandall, S.L., Hime, G.R., Resende, L.P.F., Loza-Coll, M., Aslanian, A., Yates, J.R.,
Hunter, T., Fuller, M.T., and Jones, D.L. (2014). Escargot Restricts Niche Cell to Stem Cell
Conversion in the Drosophila Testis. Cell Rep. *7*, 722–734.

Wasbrough, E.R., Dorus, S., Hester, S., Howard-Murkin, J., Lilley, K., Wilkin, E., Polpitiya, A.,
Petritis, K., and Karr, T.L. (2010). The Drosophila melanogaster sperm proteome-II (DmSP-II).
J. Proteomics *73*, 2171–2185.

Wendt, G., Zhao, L., Chen, R., Liu, C., O'Donoghue, A., Caffrey, C., Reese, M., and Collins, J.
(2020). A single-cell RNA-seq atlas of Schistosoma mansoni identifies a key regulator of blood
feeding. Science *369*, 1644–1649.

White-Cooper, H., Schafer, M.A., Alphey, L.S., and Fuller, M.T. (1998). Transcriptional and
post-transcriptional control mechanisms coordinate the onset of spermatid differentiation with
meiosis I in Drosophila. Development *125*, 125–134.

Wilk, R., Hu, J., and Krause, H.M. (2017). In Situ Hybridization: Fruit Fly Embryos and Tissues.
Curr. Protoc. Essent. Lab. Tech. *15*, 9.3.1-9.3.26.

Witt, E., Benjamin, S., Svetec, N., and Zhao, L. (2019). Testis single-cell RNA-seq reveals the
dynamics of de novo gene transcription and germline mutational bias in drosophila. Elife *8*, 1–
22.

Xu, S., Tyagi, S. and Schedl, P. (2014). Spermatid cyst polarization in Drosophila depends upon
apkc and the CPEB family translational regulator orb2. PLoS Genet. 10, e1004380

Zoller, R., and Schulz, C. (2012). The Drosophila cyst stem cell lineage. Spermatogenesis *2*,
145–157.

Zur Lage, P., Newton, F.G., and Jarman, A.P. (2019). Survey of the Ciliary Motility Machinery of
Drosophila Sperm and Ciliated Mechanosensory Neurons Reveals Unexpected Cell-Type
Specific Variations: A Model for Motile Ciliopathies. Front. Genet. *10*, 24.

1742

Figure 1

Figure 2

Figure 3

Figure 4.

Figure 5

A

Distal Ends of Elongating Spermatids

B *f-cup* DNA

C *loqs* DNA

D *wa-cup* DNA

E *soti* DNA

F *wa-cup* DNA

G

Features: Pp2C1, CG6701, Parp16, Glut3

Percent Expressed: 25, 50, 75, 100

Average Expression: 2, 1, 0

25 22 5 78 40 97 41 51 73 35 33 48 105 64 45 13 21 56 100 3 106 16 55 63 39 27 83 15 0 28 26 24 18 50 17 96 12 30 37 75 66 10 46

GSC/SG   Spermatocyte   Spermatid

H

Pp2C1 — Fold Change 4.03

CG6701 — 1.74

Parp16 — 8.65

Glut3 — 7.65

Expression vs Pseudotime

Figure 6

U Expression by cluster as cyst lineage progresses

Figure 7

Figure 8