Rhizospheric miRNAs affect the plant microbiota.

Harriet Middleton[1,2]#, Jessica Ann Dozois[2]#, Cécile Monard[1], Virginie Daburon[1], Emmanuel Clostres[1], Julien Tremblay[2], Jean-Philippe Combier[3], Étienne Yergeau[2]* and Abdelhak El Amrani[1]*

[1]ECOBIO, CNRS UMR 6553, Université de Rennes, Campus Beaulieu, Rennes, France

[2]Institut National de la Recherche Scientifique, Centre Armand-Frappier Santé Biotechnologie, Laval, Québec, Canada.

[3] LRSV, UMR CNRS/UT3/INP 5546, Auzeville-Tolosane, France.

#Contributed equally.

*Corresponding authors: abdelhak.elamrani@univ-rennes1.fr and Etienne.Yergeau@inrs.ca

Abstract

Recently, small RNAs have been shown to play important roles in cross-kingdom

communication, notably in plant-pathogen relationships. Plant miRNAs were even shown to

regulate gene expression in the gut microbiota. But what impact do they have on the plant

microbiota? Here we hypothesized that plant miRNAs can be found in the rhizosphere of

plants, where they are taken up by rhizosphere bacteria, influencing their gene expression,

thereby shaping the rhizosphere bacterial community. We found plant miRNAs in the

rhizosphere of *Arabidopsis thaliana* and *Brachypodium distachyon*. These plant miRNAs

were also found in rhizosphere bacteria, and fluorescent synthetic miRNAs were taken up by

cultivated bacteria. A mixture of five plant miRNAs modulated the expression of more than a

hundred genes in *Variovorax paradoxus*, whereas no effect was observed in *Bacillus*

mycoides. Similarly, when *V. paradoxus* was grown in the rhizosphere of *Arabidopsis* that

overexpressed a miRNA, it changed its gene expression profile. The rhizosphere bacterial

communities of *Arabidopsis* mutants that were impaired in their miRNA or small RNA

pathways differed from wildtype plants. Similarly, bacterial communities of *Arabidopsis*

overexpressing specific miRNAs diverged from control plants. Finally, the growth and the

abundance of specific ASVs of a simplified soil community were affected by exposure to a

mixture of synthetic plant miRNAs. Taken together, our results support a paradigm shift in

plant-bacteria interactions in the rhizosphere, adding miRNAs to the plant tools shaping

microbial assembly.

Keywords: plant miRNAs, rhizosphere, bacterial communities, *Variovorax*, transcriptomics.

Introduction

MicroRNAs (miRNAs) are small ~21nt non-coding RNAs that control target gene

expression, through sequence complementarity. Their roles in plants vary from regulating

developmental processes to responding to abiotic and biotic stresses. Plants also use

miRNAs to interact with pathogens: for example, cotton plants use them to inhibit the fungal

pathogen *Verticillium dahlia* [1] and *Arabidopsis thaliana* and tomato to inhibit another fungal

pathogen, *Botrytis cinerea* [2-4]. The more general role of plant miRNAs in shaping the

microbiota is, however, not known. We had hypothesized that miRNAs could be a novel, key

mediator in the rhizosphere [5], as it was shown to be in the mammalian gut. In mice and

human, both intestinal miRNAs and ingested plant-derived miRNAs regulate gut bacterial

gene expression and growth, shaping the gut microbiota [6, 7] [8].

Arabidopsis roots contain many miRNAs, of which over half are expressed in a

tissue-specific manner, with several being enriched at the root tip, in the early meristematic

zone [9]. This zone is also a recognized hotspot for plant-driven microbial selection [10]. The

current paradigm is that the plant selects the rhizospheric microbiota through

rhizodeposition, including exudates such as sugars, peptides, amino acids, nucleic acids,

nucleotides, fatty acids or secondary metabolites [11]. Here, we hypothesize that plant

miRNAs 1) can be found in the rhizosphere of plants, 2) are taken up by rhizosphere

bacteria, 3) influence the gene expression of rhizosphere bacteria, and 4) shape the

bacterial community. We report several independent experiments designed to test these

hypotheses.

Methods

The full method description is available as supplementary material.

Detection of plant miRNAs in the rhizosphere and roots.

For rhizosphere analyses, triplicate *Arabidopsis thaliana* (Col-0) and *Brachypodium*

distachyon (Bd21-3) were grown for a month, alongside unplanted soils. For root analyses,

we grew ten replicated *Aradidopsis thaliana* (Col-0) plants for 21 days, with unplanted

controls. RNA was extracted from the plant roots and rhizosphere and from the unplanted

soils and sent for small RNA sequencing. Sequences between 18 and 27 nucleotides were

mapped against *Arabidopsis* and *Brachypodium* genomes and were assigned to known

miRNAs. From these miRNAs we constructed an abundance table that was used to compare

the rhizosphere to the unplanted controls. Rhizospheric and root miRNAs were defined as

the miRNAs that had at least 10 reads for each of the rhizosphere or root samples and a

maximum of one read across all the bulk soil samples.

Internalization of plant miRNAs by bacteria.

Detection of plant miRNAs in rhizospheric bacteria.

We extracted bacterial cells from the rhizosphere soil of one-month old *A. thaliana* (Col-0)

using a Nycodenz gradient. RNA was extracted from the bacterial pellet [12], sequenced and

processed as described above, but with a lower threshold because of the lower number of

plant miRNA sequences retrieved. Only the miRNAs that were represented by at least 5

reads in each rhizosphere samples and absent in the bulk soil samples were kept. To further

confirm that miRNAs found in the rhizospheric bacteria originated from the plant and not the

bacteria, miRNA sequences were searched for on the + and - strands of 3,837 bacterial

genomes [13], of which 1,160 were isolated from plants.

Confocal microscopy.

Variovorax paradoxus EPS and *Bacillus mycoides* YL123 were grown overnight in liquid

media. The cultures were incubated for 4 hours with 3'-Cy5 fluorescent miRNAs (ath-

miR159a or a scrambled control – same nucleic acid content but in different order), or a

pCp-Cy5 control. Twenty minutes before visualization, the cultures were also treated with

MitoTracker Green FM (Invitrogen). We visualized the washed and concentrated culture

using a confocal microscope (Zeiss LSM780).

Flow cytometry.

We prepared the cultures as described in the confocal microscopy section. The cells were

fixed with 4% Paraformaldehyde (PFA) and then stained with a DNA marker (Hoechst

33342). The cells were diluted in PBS and processed by flow cytometry (BD LSRFortessa).

We used a variety of controls to ensure that our statistical analyses were conducted on

bacteria that were positive for our green (MitoTraker), red (Cy5 tagged miRNAs) and blue

(Hoechst) markers.

Effect of miRNAs on bacterial gene expression.

Bacterial transcriptomic response to miRNAs.

Variovorax paradoxus EPS and *Bacillus mycoides* YL123 were grown until the exponential

phase, when they were exposed to a mix of the 6 most abundant rhizospheric miRNAs:

miR159a, miR159b, miR159c, miR161.1, miR158a and miR165b, or a control mix of 6

scrambled miRNAs. The cultures were sampled after 20 min and 120 min of incubation,

pelleted, and the RNA of the cell pellet was extracted and sequenced. The transcripts were

mapped on their respective genomes, and differential expression analyses (plant miRNA vs.

scrambled miRNA) were done using DESeq2.

In silico analysis of miRNA targets.

To predict potential targets of the six miRNA used on the genome of *Variovorax paradoxus*,

we implemented a workflow [14] named *mirnatarget 1.0 : miRNA target finder*

(https://github.com/jtremblay/MiRNATarget), which was inspired from the plant miRNA target

finder, *psRNAtarget* [15].

In vitro miPEP treatment and transcriptomic experiment.

Approximately 50 *Arabidopsis thaliana* (Col-0) surface-sterilized seeds were grown

axenically in Petri dishes in a growth chamber. After twenty days of growth, the plants were

treated twice within 24 hours by inoculating miPEP159c, a scrambled miPEP, or water at the

plant crown. One hour after the second miPEP treatment, *Variovorax paradoxus* was

inoculated along the roots of the seedlings. Two hours after the bacterial inoculation, the

plants and rhizospheres were sampled, and RNA was extracted. From the potential targets

of miR159c found above, three were selected for RT-qPCR analyses in the rhizosphere –

alpha-2-macroglubulin, phosphatidate cytydylyltransferase, and LysR. We also quantified the

abundance in plant tissues of the primary transcript to mir159c, pri-miR159c.

Effect of miRNAs on the bacterial community.

Arabidopsis mutant experiment.

Five *A. thaliana* mutants were chosen. *RTL1* mutant over-expresses RTL1 protein which

results in a suppression of siRNA pathway without affecting miRNAs [16]. *RTL1myc* over-

expresses RTL1 protein flagged with Myc epitope, rendering RTL1 less active, so siRNA

pathway is less suppressed than with *RTL1* mutant. *Ago1-27* mutant has AGO protein

function partially impaired and is completely post-transcription gene silencing (PTGS)

deficient [17]. *Dcl1-2* mutant has total loss of function of DCL1 protein resulting in low levels

of miRNA and developmental problems [18]. *Hen1-4* mutant is miRNA defective but is also

affected in some siRNA – PTGS [19]. HEN1 methylates siRNA and miRNA to maintain their

levels and size, but also to protect them from uridylation and subsequent degradation [20].

The plants were grown for a month, after which the roots and attached rhizosphere were

sampled, its DNA extracted, amplified using 16S rRNA gene primers, and sequenced. The

same primers were used in real-time quantitative PCR to quantify bacterial abundance.

Amplicon sequencing data was processed with AmpliconTagger [21] and the R package

"phyloseq" v 1.32.0 [22].

miPEP experiment.

We treated *Arabidopsis thaliana* Col-0 with 500 μL of water (control condition) or a miPEP

solution (20 μM of miPEP159a, miPEP159b, or miPEP159c), applied at the base of the

crown, 3 times a week for a total of 10 applications. We then extracted the DNA from the

roots and the attached rhizosphere and sequenced and quantified the 16S RNA gene as

described above.

Simplified soil community experiment.

We created a simplified soil community by inoculating five different growth media with 2 g of

agricultural soil. The cultures were normalized to the same optical density, pooled, pelleted

and suspended in PBS. The cells were inoculated in a 96-wells plate containing a mixture of

17 amino acids as nitrogen source. Five wells were treated with a mixture of rhizospheric

miRNAs (ath-miR158a-3p, ath-miR158b, ath-miR159a, ath-miR827, and ath-miR5642b) and

five wells were treated with a mixture of scrambled miRNAs. These miRNAs were all found

in the rhizosphere (but for some below the stringent threshold used above) and were

predicted to target bacterial genes associated with nitrogen cycling. We measured bacterial

growth every hour for 52 hours, after which we sampled the bacteria, extracted the DNA,

and amplified and sequenced the 16S rRNA gene as described above.

Results

Plant miRNAs are present in the rhizosphere.

We sequenced small RNA extracted from *A. thaliana* rhizosphere soil and unplanted soils. One-hundred-eleven ath-miRNAs (mapped on *A. thaliana*'s genome) were detected, of which 14 were present in the rhizosphere with more than 10 reads per sample and absent in unplanted soil (Fig. 1a). The most abundant miRNAs in the rhizosphere were ath-miR158a-3p, ath-miR161.1, and various members of the miR159, miR166 and miR165 families (Fig. 1a). We then sequenced the rhizosphere miRNAs of a second model plant, *Brachypodium distachyon*. Ten bdi-miRNAs (mapped on *B. distachyon*'s genome) out of 81 miRNAs were present in the rhizosphere with more than 10 reads per sample and absent in unplanted soil (Fig. 1b). The most abundant miRNAs were bdi-miR159b-3p, bdi-miR156, bdi-miR166, bdi-miR396, and bdi-miR167. Amongst the rhizospheric miRNAs detected, four were common between *A. thaliana* and *B. distachyon*: miR159b-3p, miR167, miR166, and miR396.

We then grew more *Arabidopsis* plants and sequenced their root miRNAs to confirm that the rhizospheric miRNAs could be coming from the plant roots. Eleven miRNAs were represented by at least 10 reads in each root sample. There was a clear dominance of ath-miR165, ath-miR166, ath-miR159a, ath-miR160, and ath-miR159b.3p (Fig. 1c). Among these 11 miRNAs, 7 were common with the miRNAs found in *Arabidopsis* rhizosphere (ath-miR158a-3p, ath-miR159a, ath-miR159b-3p, ath-miR161.1, ath-miR162, ath-miR165, ath-miR166), which included most of the top five most abundant root and rhizosphere miRNAs.

A. thaliana miRNAs are internalized by soil bacteria.

Bacterial cells were isolated from the rhizosphere of one month-old *A. thaliana*, washed and their RNA content was extracted and sequenced to identify potentially internalized plant miRNAs. The miRNAs were mapped against *A. thaliana* TAIR10.1 genome, identifying a total of 34 ath-miRNAs. Five miRNAs – namely ath-miR158a-3p, ath-miR161.1, miR162, ath-miR159b-3p and ath-miR159a – were represented by at least 5

reads per rhizosphere samples and absent in bacteria extracted from the unplanted soil (Fig.

2a). Four out of these five miRNAs were among the five most abundant miRNAs found in the

rhizosphere of *A. thaliana* (Fig. 1a) and were found in a similar order of relative abundance.

To ensure that these miRNAs did not come from the bacteria themselves, their sequences

were searched for in 3,837 soil bacterial genomes, of which 1,160 were of bacteria isolated

from plants [13]. No matches were found, meaning that the miRNAs detected in the bacteria

could not be produced by the bacteria.

We then exposed two typical rhizosphere bacteria, *Variovorax paradoxus* EPS [23]

and *Bacillus mycoides* [24] to a Cy5-tagged synthetic ath-miR159a and visualized its

localization using confocal microscopy. Images show a clear localization of the miRNA inside

many bacterial cells (Fig. 2b-c). Flow cytometry confirmed that an average of 6.51%

Variovorax cells contained the Cy5 signal from plant miRNAs (average median fluorescence

intensity= 123.00), compared to 4.95% of *Bacillus* cells (average median fluorescence

intensity= 85.75). The scrambled miRNA, containing the same nucleotides as ath-miR159a

but in a different order, was also internalized (Supplementary Figure S1), suggesting a

general sequence-independent internalization mechanism for miRNAs. *Variovorax*, in

contrast to *Bacillus*, internalized more efficiently the tagged miRNAs than the tagged single

nucleotide. Indeed, a comparable amount of pCp-Cy5 was also internalized by *Bacillus* (on

average 5.04%, average median fluorescence intensity=75.8), but this was an order of

magnitude lower for *Variovorax* (on average 0.67%, average median fluorescence

intensity=81.5) (Supplementary Figure S1 and S2). This suggests that *Variovorax* actively

internalized the miRNAs, whereas it might have passively entered *Bacillus* either as intact

miRNAs or degradation products of the miRNA.

Plant miRNAs shifts rhizosphere bacterial gene expression.

We then incubated *Variovorax* and *Bacillus* with a synthetic mixture of the six most

abundant *A. thaliana* rhizosphere miRNAs: miR159a, miR159b, miR159c, miR161.1,

miR158a and miR165b, or a mixture of scrambled miRNAs at the same concentration.

Bacillus did not respond to the treatment – no gene was significantly differentially expressed

following incubation with the synthetic miRNAs. In contrast, *Variovorax* showed important

changes in response to the miRNA confrontation. After 20 min of incubation, the expression

of 79 genes was significantly lower in the plant miRNA-treated cultures and the expression

of 44 genes was significantly higher (adjusted *P*<0.05, Fig. 3a). After 120 min of incubation,

the expression of 24 genes was significantly lower in the plant miRNA-treated cultures and

the expression of 104 genes was significantly higher. Many genes were repressed after 20

min following the addition of the synthetic plant miRNAs to the bacterial culture, whereas

after 120 min, more genes presented an increased than decreased expression. Only one

gene was differentially expressed at both time points, a gene coding for a methionine

synthase (VARPA_RS01000), which was overexpressed in response to the plant miRNA

treatment.

Using the rules of a plant small RNA target finder, psRNAtarget, we compared the

differentially expressed genes from the previous experiment with the predicted target genes

for the six miRNAs used. The six miRNAs were predicted to target 237 sequences in the

Variovorax paradoxus EPS genome. Amongst these, 100 targets were positioned too far

from any coding sequence (CDS), so they were removed from the following analysis,

resulting in 137 potential targets. Amongst the 123 genes differentially expressed at 20 min,

only two were predicted as targets *in silico*: VARPA_RS05960 (targeted by miR165), coding

for a L-iditol 2-dehydrogenase, and VARPA_RS26385 (targeted by miR159a, b and c),

coding for a phosphatidate cytidylyltransferase. Only two genes were predicted as targets

amongst the 128 genes at 120 min: VARPA_RS00680 (targeted by miR165), coding for a

hypothetical protein, and VARPA_RS22555 (targeted by miR158a-3p and miR159c), coding

for a non-ribosomal peptide synthetase.

To confirm the effect of miRNAs on the bacterial transcriptome *in planta*, we

inoculated the miRNA-encoded peptide (miPEP) miPEP159c to *Arabidopsis* plants growing

in vitro and inoculated with *Variovorax*. miPEPs increase the expression of specific plant

miRNAs [25, 26]. We selected miR159c because it was among the most abundant miRNAs

in the rhizosphere, was in the mixture of miRNAs that modulated the gene expression of

Variovorax and was predicted to target several key genes. The relative expression of the

corresponding precursor miRNA (pri-miR159c) in the *Arabidopsis* plant tissue increased as

compared to the water control (t-test: t=3.31, P= 0.0145; Fig. 3c). We then quantified the

expression of three *Variovorax* genes determined to be potential targets of the miR159c

according to our bioinformatic and transcriptomic analyses. One hundred and twenty

minutes after the miPEP159c application, the relative expression of the LysR and the

phosphatidate cytidylyltransferase (CdsA) genes decreased by a factor 2.69 and 2.16,

respectively (t-test: t=4.27, *P*=0.00537, and t=3.43, P=0.00647, respectively, Fig. 3c),

whereas the expression of alpha-2 macroglobulin gene increased by a factor 1.30 (t-test:

t=2.92, P=0.0462, Fig. 3c) in comparison with the water control. The relative expressions of

the three genes and the pri-miR159c following the application of the scrambled miPEP

control (same amino acid composition as the miPEP, but in different order) were not

significantly different from the water control.

Plant miRNAs influence the rhizosphere bacterial community.

To investigate the role of plant small RNAs on the rhizospheric microbial diversity, we

grew *A. thaliana* mutants with disturbed miRNA and/or siRNA biosynthesis pathways and

analyzed their rhizospheric microbial communities by 16S rRNA gene amplicons

sequencing. The rhizosphere bacterial communities varied across the different genotypes

(Permanova: *P*<0.05; Fig 4a). In PCoA ordinations, *ago1-27* and *RTL1myc* mutants'

communities were more like unplanted soil communities than those of WT plants (not

shown). Bacterial diversity was higher in the rhizosphere of most mutant plants as compared

with the WT plants (Fig. 4b).

We then treated soil-grown *Arabidopsis* with miPEPs to further test for the effect of

the overexpression of specific miRNAs on the rhizospheric bacterial community. The

application of three miPEPs (miPEP159a, miPEP159b, and miPEP159c) on the crown of

Arabidopsis growing in soil changed the bacterial community in the roots/rhizoplane

(Permanova: R^2=0.160, P=0.031). All pairwise comparisons with the water control were

significant or nearly significant (pairwise Permanova: miPEP159a R^2=0.106, adjusted

P=0.083; miPEP159b R^2=0.164, adjusted P=0.025; miPEP159c R^2=0.154, adjusted

P=0.033). At the phylum level, the application of the miPEPs increased the relative

abundance of the *Proteobacteria* (F=8.91, P=0.00029) and decreased the relative

abundance of the *Planctomycetes* (F=40.78, P=3.64 x 10^{-10}) (Fig. 4c). This was due to

significant (adjusted P <0.05) differences between the water control and each of the miPEP

treatments in post-hoc Tukey HSD tests. The application of the miPEP on the crown of

Arabidopsis did not, however, affect the bacterial alpha diversity nor abundance.

Finally, we subjected *in vitro* a simplified microbial community that was enriched from

an agricultural soil to a mix of five synthetic plant miRNAs (ath-miR158a-3p, ath-miR158b,

ath-miR159a, ath-miR827, and ath-miR5642b), or a mix of their scrambled counterparts.

Exposure to the plant miRNAs significantly disturbed the growth of the microbial

communities during the log phase (t-test: P<0.05, Fig. 4d). The bacterial community

contained twenty ASVs at the endpoint of the incubation (52 hours) and we found shifts in

the bacterial community composition due to the plant miRNA exposure (Fig. 4e) and

significant shifts in four ASVs related to the *Enterobacter*, *Acinetobacter*, *Citrobacter* genera,

and to the *Enterobacteriaceae* family (Fig. 4f).

Discussion

Using multiple lines of independent evidence, we confirmed our four hypotheses. Plant miRNAs are present in the rhizosphere (1) and are taken up by rhizobacteria (2), which induce changes in their transcriptome (3), leading to shifts in the bacterial community (4). Plants and pathogenic fungi interact using small RNAs [1-4, 27-29] and gut bacteria are influenced by host miRNAs [6, 7], but this is the first report of this mechanism for plant-bacterial community interactions in the rhizosphere.

We detected for the first time plant miRNAs in the rhizosphere. The two model plants, *A. thaliana* and *B. distachyon* harbored a similar complement of plant miRNAs in their rhizospheres. Although it would require confirmation from more plant species, the presence of similar miRNAs in the rhizosphere of a dicotyledon and a monocotyledon suggests a conserved feature among land plants. All the major miRNAs that we found in the rhizosphere of *Arabidopsis* were also detected in the roots. This agrees well with previous reports of root miRNAs, where the two most abundant rhizospheric miRNAs ath-miR158a and ath-miR161.1, were highly enriched in the early meristematic zone [9]. Many root exudates, such as extracellular DNA, soluble compounds, and mucilage, are produced and secreted, by border cells, in this region of the root tip [30]. Even though there was a large overlap between the root and rhizosphere miRNAs in our two experiments, the relative abundance of the miRNAs was not the same, alluding to a potential selection mechanism for the miRNAs that make it to the rhizosphere. Alternatively, this pattern could also be explained by different half-life in the rhizosphere, preferential uptake by bacteria, the sequencing of entire roots, or by the slightly different conditions under which the two experiments were run.

Bacteria growing in the rhizosphere of *Arabidopsis* contained plant miRNA inside their cells and isolated rhizosphere bacteria took up a fluorescent synthetic plant miRNA. The sequence of the miRNA did not affect the uptake, as the scrambled miRNA was taken up just as efficiently as the plant miRNA. Both our Gram-positive and Gram-negative model bacteria showed that they could take up the miRNAs. The incorporation of eukaryote

miRNAs in bacteria is consistent with their ability to absorb environmental nucleic acids,

such as extracellular DNA, through natural competence. Alternatively, *in planta*, in the same

way that some bacteria secrete small RNAs in outer membrane vesicles [31], bacteria may

internalize external DNA *via* vesiduction [32], *i.e.* membrane fusion of a vesicle containing

DNA or RNA. The use of vesicles seems, however, not necessary for plant-microbe miRNA-

based interactions, as naked miRNAs were efficiently taken up by bacteria, and led to

transcriptomic and community shifts.

In plants, miRNA induce mRNA cleavage or translation inhibition, through near

perfect sequence complementarity [33]. Studies in the human gut also suggested that host

miRNA interacts with bacterial mRNA through sequence complementarity [6, 8], so we used

the rules for plant miRNA based on sequence homology, to search for targets in bacterial

genomes. Using this model to select our target genes for the *in vitro* miPEP experiment, we

found that two out of three targets were indeed inhibited following miPEP application and

increased expression of the precursor miR159c. The third gene, encoding for alpha-2-

macroglobulin, was, however, overexpressed in the presence of the miPEP. In bacteria, non-

coding small RNAs can sometimes induce expression of target mRNAs [6, 31].

Even though the miPEP experiment showed that prediction tools worked well when

focusing on a few genes, *Variovorax paradoxus* differentially expressed only 4 of the 137

predicted targets and differentially expressed another 247 non-predicted genes when

exposed to a mixture of six rhizospheric miRNAs. This suggests that either 1) the bacteria

adjusted their transcriptome in response to the shift in the expression of the miRNA-targeted

gene, 2) as it is often the case in plants [34], the miRNAs targeted bacterial transcription

factors, such as the ones from the LysR family that were differentially expressed at both time

points, 3) the miRNAs led to the production of secondary siRNA, as previously shown in

plants for two of our rhizospheric miRNAs, miR165 and miR161.1 [35], 4) target genes were

translationally repressed, which would be undetectable with transcriptomics, though this is

less common in plants than mRNA cleavage [33], 5) miRNAs protected targets from

repression, as it was shown for arbuscular mycorrhizal fungi [36], or 6) since many predicted

targets of the rhizospheric miRNAs were not in CDS, miRNAs could have affected DNA

methylation [37] or interacted with gene promoters [38]. Clearly, further investigation is

needed to clarify how eukaryotic miRNAs affect bacteria.

In contrast to *Variovorax*, plant miRNAs did not impact the transcriptome of *Bacillus*.

In our simplified soil community, only 4 out of the 20 ASVs were significantly impacted by

plant miRNAs, and, similarly, intestinal miRNAs only impacted the growth of specific

bacterial strains [6]. Plant exosomes containing miRNAs are also preferentially taken up by

some bacteria, affecting their gene expression and activity [8]. Our experiments were,

however, carried out using naked miRNAs, excluding this explanation. We first thought that

this could be related to differences in the cell wall that made miRNA entry impossible for

Gram-positive, but our microscopy work disproved that. Another possibility is that the

mechanism of interaction differs between the two groups of bacteria. In bacteria, chaperone

proteins such as RNA-binding Hfq, ProQ, or CrsA proteins protect small RNAs and stabilize

their interaction with mRNA, improving the formation of sRNA-mRNA duplexes that lead to

gene silencing [39]. This crucial role of chaperones in sRNA-mediated interactions was only

reported for a handful of Gram-positive bacteria [40, 41]. Alternatively, competence for DNA

uptake depends on environmental conditions, such as stress, nutrient availability, and cell

density [42]. The two bacteria tested might have different cues to initiate nucleic acid uptake,

and the growing conditions might not have been met during the transcriptomic experiment to

trigger this behavior in *Bacillus*. In any case, the differential transcriptomic response to

miRNAs of the two bacteria tested suggests a selective mechanism in the rhizosphere. Any

effect on a bacterium could have cascading effects on the rest of the community.

Arabidopsis miRNAs impacted the bacterial community in the root environment. We

conducted three complementary experiments to prove this point. First, we examined the

root-associated bacterial community of *A. thaliana* mutants affected in the biosynthesis of

miRNA and/or siRNA. Many of these mutants had disrupted bacterial communities

compared to wild-type plants. One of the most relevant mutants, the *dcl1-2* mutant, which is

specifically impaired in miRNA production, was severely affected in its community

composition at the phylum level and harbored a more diversified community. Microbial

communities in the roots and rhizosphere of *ago1-27* and *RTL1myc* mutants resembled

those of an unplanted soil more than those of wild-type plants. This suggests that mutations

in small RNA related pathways lead to a certain dysbiosis in the roots and rhizosphere

microbiota. Bacterial diversity was higher in the root environment of mutant plants which

could reflect a weaker selection from these plants because of their lack of miRNAs and/or

siRNAs. The mutations used are, however, pleiotropic, and plants were severely affected in

their phenotype, which could have also affected the bacterial community.

Second, the bacterial community associated with soil-grown *Arabidopsis* responded

significantly to miPEP application. As we showed in the *in vitro* experiment using

miPEP159c, miPEPs stimulate the production their corresponding miRNA in plant tissues.

This means that the up-regulation of a single miRNA could lead to changes in the bacterial

community. Other than the direct effect of the overexpressed miRNA on the bacteria, the

shifts observed in bacterial community could, however, also be explained by other factors.

For instance, plant miRNAs alter various physiological processes within the plant, such as

root development and plant immune response [34]. These changes induced by the

overexpression of some miRNAs would also lead to large shifts in the plant microbiota.

Third, to exclude most of the indirect plant-mediated effects of the mutant and the

miPEP experiments, we did an *in vitro* experiment with a simplified soil-derived bacterial

community of twenty ASVs exposed to a mixture of synthetic miRNAs. The plant miRNAs

affected the abundance of five ASVs and the growth of the community during the log phase.

This shows that plant miRNAs directly affect bacterial communities. At the individual level, a

bacterium could change in relative abundance because of 1) direct effect of the miRNA on

its growth or 2) changes in the relative abundance of other bacteria with which it interacts.

Some species can have a keystone role in interaction networks [43], and shift in these

species would influence the entire community. For instance, the presence of *Variovorax* in

the rhizosphere of *Arabidopsis* counteracted the root growth inhibition induced by many

other members of the community [44]. Plant miPEPs, through their effect on miRNAs, also

modulated the interactions between plants and key root symbionts, such as arbuscular mycorrhizal fungi [36] and rhizobia [45]. This effect would profoundly alter the microbial community, even if the miRNA had only affected a single keystone species.

We showed here for the first time that plant miRNAs are found in the rhizosphere of two model plants, that they are internalized in rhizosphere bacteria, that they affect the transcriptome of rhizosphere bacteria, and that they modulate soil microbial community composition and growth. The rhizosphere effect is thought to be mainly due to the rhizodeposition of various small organic molecules. Our study suggests a novel molecule, miRNAs, that plants could use to interact with their microbiota, challenging the current paradigm of rhizosphere microbial assembly. Many questions remain to be answered, but our study has shown beyond any doubt that plant miRNAs can shape the rhizosphere bacterial community.

Acknowledgments

We want to thank Hervé Vaucheret and Taline Elmayan (IJPB, INRAE, Versailles, France) for sharing their *Arabidopsis* mutant seeds with us and Paul M. Orwin for providing the *Variovorax paradoxus* EPS strain. We also thank Christophe Penno for the helpful discussions. We also thank Jessy Tremblay (INRS, Laval, Canada) for the expertise and support regarding the confocal microscopy and flow cytometry experiments.

Funding

This work was financially supported by a CNRS MITI (Mission pour les Initiatives Transverses et Interdisciplinaires 2020-2021), by a IEA (International Emerging Actions) grant (France), by the French National program EC2CO (Ecosphère Continentale et Côtière 2024-2025), a CRSNG discovery grant (RGPIN-2020-05723), a FRQNT Samuel-de-Champlain grant (2019-FQ-264131). We acknowledge Compute Canada for access to the Graham compute cluster.

Conflict of interest statement

The authors report no conflict of interest.

Data availability statement

We deposited the sequence data generated in this study under NCBI BioProject accessions

PRJNA836586 (rhizosphere miRNAs), PRJNA1107220 (root miRNAs), PRJNA1111839

(soil-extracted bacteria miRNAs), PRJNA1111831 (mutant and miPEP 16S rRNA gene

amplicons), PRJNA1111829 (simplified soil community 16S rRNA gene amplicons), and

PRJNA1111827 (*Bacillus* and *Variovorax* transcriptomics). The R code used to manipulate

the data and generate the figures is available on GitHub (https://github.com/le-labo-

yergeau/Middleton_miRNA) with the accompanying data being available on Zenodo

(https://zenodo.org/doi/10.5281/zenodo.11105307).

References

445   1.   Zhang T, Zhao Y-L, Zhao J-H, Wang S, Jin Y, Chen Z-Q et al. Cotton plants export
446   microRNAs to inhibit virulence gene expression in a fungal pathogen. Nat Plants.
447   2016;2:16153.

448   2.   Cai Q, Qiao L, Wang M, He B, Lin F-M, Palmquist J et al. Plants send small RNAs in
449   extracellular vesicles to fungal pathogen to silence virulence genes. Science.
450   2018;360:1126-1129.

451   3.   Qin S, Veloso J, Puccetti G, van Kan JAL. Molecular characterization of cross-
452   kingdom RNA interference in Botrytis cinerea by tomato small RNAs. Front Plant Sci.
453   2023;14:

454   4.   Wu F, Huang Y, Jiang W, Jin W. Genome-wide identification and validation of
455   tomato-encoded sRNA as the cross-species antifungal factors targeting the virulence
456   genes of Botrytis cinerea. Front Plant Sci. 2023;14:

457   5.   Middleton H, Yergeau É, Monard C, Combier J-P, El Amrani A. Rhizospheric Plant–
458   Microbe Interactions: miRNAs as a Key Mediator. Trends Plant Sci. 2021;26:132-
459   141.

460   6.   Liu S, da Cunha AP, Rezende RM, Cialic R, Wei Z, Bry L et al. The Host Shapes the
461   Gut Microbiota via Fecal MicroRNA. Cell Host Microbe. 2016;19:32-43.

462   7.   Liu S, Rezende RM, Moreira TG, Tankou SK, Cox LM, Wu M et al. Oral
463   Administration of miR-30d from Feces of MS Patients Suppresses MS-like Symptoms
464   in Mice by Expanding Akkermansia muciniphila. Cell Host Microbe. 2019;26:779-794.
465   e8.

466   8.   Teng Y, Ren Y, Sayed M, Hu X, Lei C, Kumar A et al. Plant-Derived Exosomal
467   MicroRNAs Shape the Gut Microbiota. Cell host & microbe. 2018;24:637-652.e8.

468   9.   Breakfield NW, Corcoran DL, Petricka JJ, Shen J, Sae-Seaw J, Rubio-Somoza I et
469   al. High-resolution experimental and computational profiling of tissue-specific known
470   and novel miRNAs in Arabidopsis. Genome Res. 2012;22:163-176.

471 10.   DeAngelis KM, Brodie EL, DeSantis TZ, Andersen GL, Lindow SE, Firestone MK.
472   Selective progressive response of soil microbial community to wild oat roots. ISME J.
473   2009;3:168–178.

474 11.   Tian T, Reverdy A, She Q, Sun B, Chai Y. The role of rhizodeposits in shaping
475   rhizomicrobiome. Environ Microbiol Rep. 2020;12:160-172.

476 12.   Griffiths RI, Whiteley AS, O'Donnell AG, Bailey MJ. Rapid method for coextraction of
477   DNA and RNA from natural environments for analysis of ribosomal DNA- and rRNA-
478   based microbial community composition. Appl Environ Microb. 2000;66:5488–5491.

479 13.   Levy A, Salas Gonzalez I, Mittelviefhaus M, Clingenpeel S, Herrera Paredes S, Miao
480   J et al. Genomic features of bacterial adaptation to plants. Nat Genetics. 2017;

481 14.   Penno C, Tremblay J, O'Connell Motherway M, Daburon V, El Amrani A. Analysis of
482   Small Non-coding RNAs as Signaling Intermediates of Environmentally Integrated
483   Responses to Abiotic Stress. Methods Mol Biol. 2023;2642:403-427.

484 15.   Dai X, Zhuang Z, Zhao PX. psRNATarget: a plant small RNA target analysis server
485   (2017 release). Nucl Acids Res. 2018;

486 16.   Shamandi N, Zytnicki M, Charbonnel C, Elvira-Matelot E, Bochnakian A, Comella P
487   et al. Plants Encode a General siRNA Suppressor That Is Induced and Suppressed
488   by Viruses. PLoS Biol. 2015;13:e1002326.

489 17.   Morel JB, Godon C, Mourrain P, Béclin C, Boutet S, Feuerbach F et al. Fertile
490   hypomorphic ARGONAUTE (ago1) mutants impaired in post-transcriptional gene
491   silencing and virus resistance. Plant Cell. 2002;14:629-639.

492 18.   Xie Z, Kasschau KD, Carrington JC. Negative feedback regulation of Dicer-Like1 in
493   Arabidopsis by microRNA-guided mRNA degradation. Curr Biol. 2003;13:784-9.

494 19.   Boutet S, Vazquez F, Liu J, Béclin C, Fagard M, Gratias A et al. Arabidopsis HEN1: a
495   genetic link between endogenous miRNA controlling development and siRNA
496   controlling transgene silencing and virus resistance. Curr Biol. 2003;13:843-848.

20. Li J, Yang Z, Yu B, Liu J, Chen X. Methylation protects miRNAs and siRNAs from a
3'-end uridylation activity in Arabidopsis. Curr Biol. 2005;15:1501-7.
21. Tremblay J, Yergeau É. Systematic processing of rRNA gene amplicon sequencing
data. GigaScience. 2019;8:giz146.
22. McMurdie PJ, Holmes S. phyloseq: an R package for reproducible interactive
analysis and graphics of microbiome census data. PLoS One. 2013;8:e61217.
23. Han JI, Spain JC, Leadbetter JR, Ovchinnikova G, Goodwin LA, Han CS et al.
Genome of the Root-Associated Plant Growth-Promoting Bacterium Variovorax
paradoxus Strain EPS. Genome Announc. 2013;1:
24. Agoussar A, Azarbad H, Tremblay J, Yergeau É. The resistance of the wheat
microbial community to water stress is more influenced by plant compartment than
reduced water availability. FEMS Microbiol Ecol. 2021;97:fiab149.
25. Lauressergues D, Couzigou J-M, San Clemente H, Martinez Y, Dunand C, Bécard G
et al. Primary transcripts of microRNAs encode regulatory peptides. Nature.
2015;520:90.
26. Couzigou J-M, Lauressergues D, Bécard G, Combier J-P. miRNA-encoded peptides
(miPEPs): A new tool to analyze the roles of miRNAs in plant biology. RNA Biol.
2015;12:1178-1180.
27. Wang B, Sun Y, Song N, Zhao M, Liu R, Feng H et al. Puccinia striiformis f. sp. tritici
microRNA-like RNA 1 (Pst-milR1), an important pathogenicity factor of Pst, impairs
wheat resistance to Pst by suppressing the wheat pathogenesis-related 2 gene. New
Phytol. 2017;215:338-350.
28. Wang M, Weiberg A, Lin FM, Thomma BP, Huang HD, Jin H. Bidirectional cross-
kingdom RNAi and fungal uptake of external RNAs confer plant protection. Nat
Plants. 2016;2:16151.
29. Wang M, Weiberg A, Dellota E, Jr., Yamane D, Jin H. Botrytis small RNA Bc-siR37
suppresses plant defense genes by cross-kingdom RNAi. RNA Biol. 2017;14:421-
428.
30. Sasse J, Martinoia E, Northen T. Feed Your Friends: Do Plant Exudates Shape the
Root Microbiome? Trends Plant Sci. 2018;23:25-41.
31. Choi J-W, Um J-H, Cho J-H, Lee H-J. Tiny RNAs and their voyage via extracellular
vesicles: Secretion of bacterial small RNA and eukaryotic microRNA. Exp Biol Med.
2017;242:1475-1481.
32. Soler N, Forterre P. Vesiduction: the fourth way of HGT. Environ Microbiol.
2020;22:2457-2460.
33. Yu Y, Jia T, Chen X. The 'how' and 'where' of plant microRNAs. New Phytol.
2017;216:1002-1017.
34. Tang J, Chu C. MicroRNAs in crop improvement: fine-tuners for complex traits. Nat
Plants. 2017;3:17077.
35. Manavella PA, Koenig D, Weigel D. Plant secondary siRNA production determined
by microRNA-duplex structure. Proc Natl Acad Sci. 2012;109:2461-2466.
36. Couzigou JM, Lauressergues D, André O, Gutjahr C, Guillotin B, Bécard G et al.
Positive Gene Regulation by a Natural Protective miRNA Enables Arbuscular
Mycorrhizal Symbiosis. Cell Host & Microbe. 2017;21:106-112.
37. Wu L, Zhou H, Zhang Q, Zhang J, Ni F, Liu C et al. DNA Methylation Mediated by a
MicroRNA Pathway. Mol Cell. 2010;38:465-475.
38. Toscano-Garibay JD, Aquino-Jarquin G. Transcriptional regulation mechanism
mediated by miRNA–DNA•DNA triplex structure stabilized by Argonaute. Biochimica
et Biophysica Acta. 2014;1839:1079-1083.
39. Boutet E, Djerroud S, Perreault J. Small RNAs beyond Model Organisms: Have We
Only Scratched the Surface? Int J Mol Sci. 2022;23:4448.
40. Nielsen JS, Lei LK, Ebersbach T, Olsen AS, Klitgaard JK, Valentin-Hansen P et al.
Defining a role for Hfq in Gram-positive bacteria: evidence for Hfq-dependent
antisense regulation in Listeria monocytogenes. Nucl Acids Res. 2010;38:907-919.

41. Müller P, Gimpel M, Wildenhain T, Brantl S. A new role for CsrA: promotion of
complex formation between an sRNA and its mRNA target in Bacillus subtilis. RNA
Biol. 2019;16:972-987.
42. Blokesch M. Natural competence for transformation. Curr Biol. 2016;26:R1126-
R1130.
43. Carlström CI, Field CM, Bortfeld-Miller M, Müller B, Sunagawa S, Vorholt JA.
Synthetic microbiota reveal priority effects and keystone strains in the Arabidopsis
phyllosphere. Nat Ecol Evol. 2019;3:1445-1454.
44. Finkel OM, Salas-González I, Castrillo G, Conway JM, Law TF, Teixeira PJPL et al.
A single bacterial genus maintains root growth in a complex microbiome. Nature.
2020;587:103-108.
45. Couzigou J-M, André O, Guillotin B, Alexandre M, Combier J-P. Use of microRNA-
encoded peptide miPEP172c to stimulate nodulation in soybean. New Phytol.
2016;211:379-381.
565

Figure legends.

Figure 1: Plant miRNAs are present in the rhizosphere and absent in unplanted soil.

Relative abundance of the plant miRNAs found in the rhizosphere of *Arabidopsis thaliana*

(A), *Brachypodium distachyon* (B) and in the roots of *Arabidopsis thaliana* (C), while

completely absent from unplanted soil.

Figure 2: Plant miRNAs are internalized by rhizosphere bacteria. Relative abundance of

Arabidopsis thaliana miRNAs found in cells pelleted from the rhizosphere and absent in cells

pelleted from unplanted soil (A) and confocal microscopy images of *Variovorax paradoxus*

(B) and *Bacillus mycoides* (C) after a 4-hour exposure to ath-miR159a tagged with Cy5.

Arrows indicate cells containing the fluorescent molecule.

Figure 3: Plant miRNAs affect the transcriptome of a rhizosphere bacterium. Gene

expression of *Variovorax praradoxus* after 20 min (A) and 120 min (B) exposure to a mixture

of five synthetic miRNAs. (C): Arabidopsis thaliana expression of pri-miR159c and

rhizospheric *Variovorax paradoxus* expression of CdsA, Alpha-2-macroglobulin and LysR

after exposure of the plant to miPEP159c.

Figure 4: Plant miRNAs affect the bacterial community. Phylum-level bacterial

community composition (A) and Shannon diversity (B) for different *Arabidopsis* mutants with

impaired small RNAs pathways. (C): Phylum-level bacterial community composition for

Arabidopsis plants inoculated with miPEP159a, miPEP159b or miPEP159c. Growth curve

(D), community composition at genus level (E), and relative abundance of significantly

affected ASVs (F) for a simplified soil microbial community exposed to a mixture of synthetic

plant or scrambled miRNAs

589 **Figure 1.**

591
592

593 **Figure 2.**

A

B

C

594
595

Figure 3.

Figure 4.

603 **Figure S1.**

604
605

606   **Figure S2.**

607
608

Supplementary methods
Detection of plant miRNAs in the roots and the rhizosphere

Plant growth conditions.

For the detection of miRNA in the rhizosphere, *Arabidopsis thaliana* (Col-0) and *Brachypodium distachyon* (Bd21-3) were planted in triplicates and grown for a month in a growth cabinet, alongside unplanted soils. For the detection of miRNA in roots, *Arabidopsis thaliana* (Col-0) was planted in ten different experimental blocks and grown for 21 days in a growth cabinet, alongside unplanted soils. The program of the growth cabinets was: 12 hours of daylight, 3 hours of twilight, 3 hours of dawn light, and 6 hours of darkness at a constant humidity of 70%. The temperature varied in accordance with the light cycle between 20-25 °C.

Sampling, RNA extraction and sequencing

The rhizospheres and control soils were sampled and flash-frozen in liquid nitrogen. RNA was then extracted from 2 g of soil using MOBIO RNA Power soil kit, eluted in 100 µL, following the manufacturer's guidelines. The roots were gently and quickly washed with sterile PBS, placed in a sterile tube and flash frozen with liquid nitrogen. The roots were lyophilized for three hours, crushed with a micro pestle, and RNA was extracted using the Qiagen RNeasy Plant Mini kit, following the manufacturer's guidelines. The RNA was treated with DNase (Turbo DNA-free kit, Ambion), purified and concentrated (RNA Clean & Concentrator 25µl, ZymoResearch). We evaluated RNA quality with Nanodrop and 4% agarose gels, before sending it for sequencing on an Illumina HiSeq 2500 in single read mode (1x50bp) at the Centre d'expertise et de services Génome Québec (Montreal, Canada).

Bioinformatic analyses

Reads were trimmed using Trimmomatic [1] (v0.39) to remove the sequencing adapters with the following Illumina clip settings :5:10:4 and to remove the 5' and 3' flanking sequences containing bases having quality scores below 33 and 30, respectively. Common contaminants, such as Illumina adapters and PhiX spike-in sequences, were removed using

BBDuk (BBMap v38.11, https://github.com/BioInfoTools/BBMap), with key parameters: k=20

and minkmerhits=1. Remaining sequences were then filtered by size such that reads having

a length between 18 and 27 nucleotides were selected. Selected reads were mapped

against either *A. thaliana* (TAIR10/GCA_000001735.1) or *B. distachyon*

(bd21/GCF_000005505.1) reference genomes, which are the same as those used in

miRbase [2], using BWA-aln [3] with the following parameters: mismatch=1 and seed=5.

Using BEDTools v.2.29.2 [4] with -F 0.95 argument, a number of reads aligned inside

miRNA coordinates were determined, meaning that at least 95% of each read had to belong

inside the miRNA coordinates to be considered a valid hit. These assigned miRNAs which

were mapped against the reference database were kept for abundance estimation across

samples.

Statistical analyses

Rhizospheric and root miRNAs were defined as the miRNAs that had at least 10 reads for

each the rhizosphere or root samples and a maximum of one read across all the bulk soil

samples.

Internalization of plant miRNAs by bacteria

Detection of plant miRNAs in rhizospheric bacteria

Plant growth and bacteria isolation. *A. thaliana* Col-0 was grown for a month, then the

rhizosphere was sampled. Bacteria were isolated from 5 g of rhizospheric soil, using a

density gradient. Samples were added to 30 mL of 0.2% sodium hexametaphosphate and

rotated on a spinning wheel for 2 h. Samples were then centrifuged at 18 *g* for 1 min at

10°C, the supernatants were pooled two-by-two and centrifuged at 2824 x *g* for 20 min at

10°C. The microbe-containing pellet was resuspended in 10 mL of 0.8% NaCl and samples

were again pooled by two. The resuspended solution was then transferred onto 10 mL of

sterile Nycodenz and was centrifuged at 3220 *g* for 40 min. The bacterial layer was collected

and a volume of 0.8% NaCl sufficient for a total of 35 mL was added. This solution was

centrifuged at 3220 *g* for 15 min at 10°C. Finally, the pellet was resuspended in 1 mL of

NaCl 0.8% to wash the residual Nycodenz away and centrifuged at 1700 *g* for 5 min at 10°C.

The final pellet was resuspended in 100 μL TE buffer and kept at -20°C. Eight bacterial

samples were obtained from the rhizosphere and six from unplanted soils.

<u>RNA extraction and sequencing.</u> RNA was extracted from this bacterial solution as follows:

the 100 μL were transferred into a MP Biomedicals™ Lysing Matrix E tube with 250 μL of

warmed 10% CTAB (Cetyl Trimethyl Ammonium Bromide) - 0.7 M NaCl, followed by 250 μL

of 240 mM K_2HPO_4/KH_2PO_4 (pH 8.0) and 500 μL of phenol-chloroform-isoamyl alcohol

(25:24:1) (pH 8.0). Samples were then bead-beaten using TissueLyser II (QIAGEN) at 30 Hz

for 3 min. The tubes were centrifuged at 16000 *g* for 10 min at 4°C and the supernatant was

transferred into a new tube. One volume of chloroform-isoamyl alcohol (24:1) was added to

the supernatant and centrifuged again at 16,000 *g* for 10 min at 4°C. The supernatant was

collected and 2 volumes of 30% PEG 6000-1.6 M NaCl were added, and the tube was

inverted to mix and was left at 4°C for at least 2 h. The tubes were then centrifuged at

18,000 *g* for 30 min at 4°C to precipitate RNA. The RNA pellet was then washed with 500 μL

of 70% ethanol and centrifuged at 18,000 *g* for 10 min at 4°C. The pellet was then air dried

and resuspended in 50 μL of RNAse-free water. RNA quality and quantity was verified using

a 1% agarose gel and Nanodrop, and then sent for sequencing on an Illumina NovaSeq

6000 using paired-end 100 bp with size-selected library.

<u>Bioinformatic analyses.</u> Sequencing results were processed as stated above with the

rhizospheric miRNAs. To confirm that miRNAs found in the rhizospheric bacteria originated

from the plant and not the bacteria, miRNA sequences were searched for on the + and -

strands of 3,837 bacterial genomes [5], of which 1,160 bacteria were isolated from plants.

These miRNA sequences were also searched for in the genome of *A. thaliana* TAIR10.1 in

which they were found at their expected position.

<u>Statistical analyses.</u> We used the same strategy as above to define the internalized

rhizospheric miRNAs, but with a lower threshold because there were less plant miRNA reads

produced for this experiment. We defined the internalized miRNAs as the miRNAs that were

represented by at least 5 reads in each rhizosphere microbial pellets and absent in the bulk

soil microbial pellets.

Confocal microscopy.

Variovorax paradoxus EPS and *Bacillus mycoides* YL123 were cultured on solid media

(Bacto Yeast Extract Agar for *V. paradoxus* EPS and Tryptic Soy Agar for *B. mycoides*

YL123) for 48 hours and then a few colonies were inoculated in 5 mL of the equivalent liquid

media and grown overnight (250 rpm 28 ºC). The cultures were then standardized to an

OD600 of 0.200, treated with one of the 3'-Cy5-labelled oligos (plant ath-miR159a-Cy5,

scrambled ath-miR159a-Cy5 or a modified version of cytidine pCp-Cy5, final concentration 2

μM) (Supplementary Table S1) and incubated for a total of 3 hours and 40 minutes (the

cultures were within the log phase) after which they were treated with MitoTracker Green FM

(Invitrogen, final concentration 2 μM). The cultures were immediately reincubated for 20

minutes, pelleted, washed with sterile PBS, pelleted again, and concentrated 10 times in

sterile PBS. The cells (4.5 μl of washed and concentrated culture) were then placed on a

glass bottom dish, covered with a thin slice of 2% agarose, and visualized using a confocal

microscope (Zeiss LSM780). The experiment was repeated three independent times.

Flow cytometry.

We prepared the bacterial cultures as for the confocal microscopy. After that, we took 5.5 μl

of washed and concentrated culture and incubated it for 1 h and 30 min (room temperature).

Then, we treated the bacteria with 4% Paraformaldehyde (PFA) for 40 min to fix them. We

then stained the cells with a DNA marker (Hoechst 33342) for 35 min. After which we diluted

the cells in sterile PBS to reach a final volume of 350 μl and processed the samples with a

flow cytometer (BD LSRFortessa). For each sample, 50 000 events were measured. We

used a variety of controls, such as cells only stained with Hoechst, cells only stained with

MitoTracker GreenFM, cells stained with Hoechst + Mitotracker GreenFM and cells without

dye to set appropriate gates and to ensure that our analyses were conducted on bacteria

that were positive for our green (MitoTracker), red (Cy5 tagged miRNAs) and blue (Hoechst)

markers. For each experiment, duplicates for each strain were prepared. The experiment

was repeated five independent times for *V. paradoxus* EPS and four times for *B. mycoides*

YL123. For our analyses, we used the mean of the duplicates and tested the differences

between the strains and treatments with two Kruskal-Wallis tests. As a post-hoc test, we

used Dunn's test for pairwise multiple comparisons (significance=p adjusted < 0.05).

Effect of miRNAs on bacterial gene expression

In silico analysis of miRNA targets

To predict potential plant miRNA targets in bacterial genomes, we implemented a workflow

named *mirnatarget 1.0 : miRNA target finder* (https://github.com/jtremblay/MiRNATarget),

which was inspired from the plant miRNA target finder, *psRNAtarget* [6]. This tool is based

on specific pairing patterns between plant miRNA and targets [7] and implements

SSEARCH36 (from the fasta36 v36.3.8 package) which uses the Smith-Waterman local

alignment algorithm, resulting in optimal alignments [8]. Multiple targets for each miRNA can

be identified. Based on the *psRNAtarget* rules for miRNA-target recognition, an *e*-value was

calculated for each alignment and only those with an *e*-value ≤ 5 were retained. The

positions of the targeted regions were noted with respect to the neighbouring coding

sequences (CDS): the target sequence could be inside a CDS, at the 5' or 3' flanking

regions (FLR) or in other short/very short inter-CDS regions, overlapping CDSs or

elsewhere, further away from any CDS. Targets in the latter category were positioned at a

distance superior to 100 nt from the 5' end or at a distance superior to 350 nt from the 3' end

of the CDS. These distances were chosen as they correspond to the longest UTRs that are

implicated in gene regulation. Targets in this category were discarded from the analysis,

because it was considered that a miRNA in that area would not affect mRNA translation. The

annotation of targeted regions and extraction of the closest CDSs were carried out with our

tool "align2cdsRegions" (https://github.com/ntzv-git/align2cdsRegions). Sequences of

targeted CDSs were retrieved using the *getfasta* tool from BEDTools [4] (v2.30.0) and GO

(Gene Ontology) terms were attributed using InterProScan (v5.59-91.0) [9]. For each domain

of GO (*i.e.* Biological Processes, Cellular Component, Molecular Function) and for each

targeted CDS, a GO ancestor and their description were associated, and a level of

hierarchical ontology was attributed [10].

Bacterial transcriptomic response to miRNAs

<u>Bacterial growth and miRNA treatment.</u> *Variovorax paradoxus* EPS was grown in 5% yeast

extract (YE), at 30°C with 250 rpm shaking. *Bacillus mycoides* YL123 was grown in tryptic

soy broth (TSB), at 25°C with 250 rpm shaking. Bacterial stocks were plated on YE or TSB

and single colonies were selected for liquid culture overnight (*n* = 3), in 20 mL YE or TSB. In

the morning, cultures were normalised as such: the optical density (OD) of overnight cultures

was measured at 600 nm, which were then centrifuged at 5000 rpm for 5 min to collect the

bacterial pellet. The supernatant was removed, and a calculated volume of medium was

added to the pellet to reach OD = 1 for *B. mycoides* (*i.e.* 7.36 x 10^8CFU/mL) and OD = 0.3

for *V. paradoxus* (*i.e.* 9.99 x 10^8CFU/mL). Bacterial cultures were left to grow until the

exponential phase was reached, which was determined previously (4-5 h for both strains).

OD was then measured to check that the maximum number of cells/mL for RNA extraction

was not exceeded (10^9 cells/mL). At this point, 1 µg of the miRNA mix, or scrambled mix was

applied to the cultures and gently mixed (15 µL of each miRNA solution diluted at 10 µM).

Cultures were normalised to reach the same number of bacterial cells per mL. The volume of

the cultures was varied to mimic natural differences in rhizosphere volume. Total

concentration of miRNAs ranged between 65 nM-104 nM for *B. mycoides* and 45 nM-85 nM

for *V. paradoxus*. The same cultures were sampled after 20 min and 120 min of incubation,

to evaluate the evolution of gene expression in a same biological sample in response to a

short and long confrontation with miRNAs. The miRNA mix was composed of 6 rhizospheric

miRNAs : miR159a, miR159b, miR159c, miR161.1, miR158a and miR165b (Supplementary

Table S1). The scrambled mix was composed of 6 RNA sequences with the same nucleic

acids as the respective miRNAs but in a random order (Supplementary Table S1). To

resemble mature plant miRNAs found in the rhizosphere, these synthetic miRNAs were

single-stranded with a 3' methyl group. They were synthesised by Integrated DNA
Technologies (IDT).

RNA extraction. After incubation, 1 mL of every culture was sampled and centrifuged at 5000
rpm for 5 min. The bacterial pellet was resuspended in 100 µL of a lysozyme solution: 1
mg/mL for *V. paradoxus* (Gram -) and 3 mg/mL for *B. mycoides* (Gram +). After vortexing the
resuspension, the samples were left to incubate for 15 min at room temperature. The
following steps used solutions from the RNeasy Plant Mini Kit (QIAGEN). The guanidine-
thiocyanate-based buffer RLT was added to the samples for further lysis (600 µL) and
vortexed for 10 s. For *B. mycoides* samples, an additional step was necessary: the 700 µL
were transferred to a sterile 2 mL Safe-Lock tube containing 100 mg glass beads (0.1 mm
diameter) and 100 mg zirconia/silica beads (0.5 mm diameter). Beads were resuspended by
pipetting then cells were disrupted using Fast Prep for 50 s (5 m/s), followed by 5 min on ice
and another 50 s of Fast Prep. Tubes were centrifuged at 10 000 *g* for 1 min and
supernatant was transferred to a new tube. The next steps were common to both strains. An
equal volume of 70% ethanol was added to the lysed samples and mixed by pipetting. The
lysate was transferred to a RNeasy spin column (pink column) in a 2 mL collection tube and
centrifuged at 10 000 rpm for 30 s, then the flow-through was discarded. Columns were first
washed with 700 µL of the RW1 wash buffer, followed by 500 µL RPE buffer and again with
500 µL RPE buffer. Collection tube was changed, and the column was centrifuged for 1 min
at maximum speed to dry the membrane off. To elute RNA, columns were placed in new 1.5
mL tubes and 30 µL of RNAse-free water was added before centrifugation. The 30 µL were
then reloaded and centrifuged one last time.

RNA-seq & sequence analysis. Total RNA was sent for sequencing at the Centre d'expertise
et de services Génome Québec (Montreal, Canada), where ribosomal RNAs were depleted
and messenger RNAs were sequenced. RNA-seq raw reads were processed with
ShotgunMG [11]. Reads were trimmed and filtered for quality (Trimmomatic v.39) and
mapped (BWA v0.7.17) against their respective reference genome. The genome of *B.*
mycoides YL123 originated from our lab, was sequenced, processed and submitted to the

NCBI through the Prokaryotic Genome Annotation Pipeline (PGAP) under the assembly

accession GCA_024297165.1 on GenBank. The *V. paradoxus* EPS genome was published

with accession number NC_014931 [12]. BedTools v2.23.0 was used to generate read

count files from each .bam file that were then merged to generate a raw count matrix. The

latter file was used as input data for a differential gene expression analysis using DESEq2

(Bioconductor, version 1.36.0). For each strain and each time point, gene expression of

samples treated with synthetic miRNAs was compared to those treated with scrambled

synthetic miRNAs.

In vitro miPEP transcriptomic experiment

<u>*Arabidopsis* culture *in vitro.*</u> Approximately 50 *Arabidopsis thaliana* col-0 seeds were surface-

sterilised by soaking them in 70% ethanol for 5 minutes, followed by 20 minutes in 0.5%

sodium hypochlorite. Seeds were then rinsed six times with sterile distilled water and

resuspended in 1 mL of water, which was poured onto a Petri dish, with some culture

medium, to let the water evaporate under sterile conditions. The Petri dish was sealed with

parafilm and placed at 4°C for 3 days, for the stratification process, allowing a synchronous

germination. The culture medium was composed of 1.2 g of Hoagland's No. 2 basal salt

mixture (Sigma-Aldrich, reference H2395), 8 g of agar 'suitable for plant cell culture' (Sigma-

Aldrich, ref. A8678) and 500 mL of MilliQ water. The pH of the medium was adjusted with

HCl to 5.8-5.9, before adding the final 250 mL of water, resulting in 750 mL of medium. The

solution was autoclaved and poured into square petri dishes for plant culture. When

hardened, around 2 cm of solid medium was cut out, creating a ridge where the surface-

sterilized seeds were positioned (10-15 seeds/dish). Petri dishes with these positioned

seeds were sealed with microtape and placed vertically in a growth chamber, with 16 h of

daylight at 21°C, 8 h of darkness at 19°C, with 25% relative humidity.

<u>miPEP treatment.</u> Twenty days after the seeds were placed in the growth chamber, the

seedlings had developed at least 6 leaves and were ready to be treated. The miPEP

treatment consisted in applying 100 µL of a 10 µM solution to each plant, starting at the tip of

the root to the base (bottom-to-top). A second treatment was applied 24 h later, before the

microbial inoculation. There were three conditions: miPEP159c treatment

(MQNLRVHVFLIESARC), a scrambled version of miPEP159c and a water control, each

treatment was applied to 7 Petri dishes (biological replicates), containing 10-15 plants each

(pseudo-replicates). The use of a scrambled control as well as a solvent control (*i.e.* water)

was to ensure that any "miPEP-effect" was indeed due to its action on the plant and not *via*

some nutritive or anti-microbial effect of the peptide.

<u>Bacterial growth and inoculation.</u> *Variovorax paradoxus* EPS was first inoculated, from a -

20°C stock, on a yeast extract Petri dish (5 g/L) to check that colonies are homogenous. A

single colony was transferred into a liquid culture of yeast extract to grow overnight at 30°C

and 110 rpm. The overnight culture's optical density was measured at 600 nm to estimate

the number of bacterial cells added to each plant: at OD = 1, the estimated cellular

concentration is 3.33 x 10^9 CFU/mL. 100 µL (with OD = 0.418 *i.e.* ~1.39 x 10^8 cells) of the

overnight culture was directly applied to each plant, from the tip to the base of the root, one

hour after the second miPEP treatment.

<u>Sampling.</u> Two hours after the bacterial inoculation, the plants and rhizospheres were

sampled. The rhizosphere samples were isolated using a sterile spatula by extracting small

squares of medium around the tip of the roots. Whole plants were gently extracted from the

medium, by pulling on the aerial system. Samples were flash-frozen in liquid nitrogen and

ground shortly after, using a sterile mortar and pestle. The resulting fine powder was stored

at -80°C.

<u>RNA extraction.</u> The NucleoSpin ® RNA kit (MACHEREY-NAGEL, Düren, Germany) was

used to extract RNA from ~100 mg of previously ground powder, with some changes to the

manufacturer's protocol. During the cell lysis, 400 µL of RA1 solution were used instead of

350µL, followed by 400 µL of ethanol to adjust the RNA binding conditions. The following

steps remained the same, up until the elution, where only 30µL of water was used and left to

incubate on the column before centrifugation, resulting in better yields. DNase reaction was

performed on-column during the protocol.

Target prediction. Potential targets of miRNA159c in the genome of *Variovorax paradoxus*

EPS were identified using the online *psRNAtarget* tool. On the server, the ath-miRNA159c

sequence (UUUGGAUUGAAGGGAGCUCCU), obtained from miRBase, and the accession

number of the EPS genome (CP002417.1) were uploaded for analysis, with the default

parameters of the Schema V2 of the 2017 release. Three of the targeted genes were

selected for *in vivo* quantification.

RT-qPCR. Retrotranscription was performed using 50 ng of total RNA, more or less diluted

to start the reaction with 12 µL of RNA, to which was added 1 µL dNTPs (10 mM) and 1 µL

of random primers (200 ng), followed by a 5-minute incubation at 65°C. Samples were then

placed on ice for 1min. Sequentially, 4 µL of 5X First-Strand Buffer, 1 µL of DTT (0,1 M) and

1 µL of SuperScript™ Reverse Transcriptase (200 U/µL, Invitrogen) were mixed in by

pipetting and the samples incubated for 5 min at 25°C, followed by 1 h at 60°C and 15 min at

70°C. Samples were then placed on ice and 100 µL of RNAse-free water was added to the

25 µL of newly synthesised cDNA. Quantitative PCR was performed using a reaction mix

composed of 5 µL of iTaq Universal SYBR Green Supermix (Bio-Rad Laboratories, ref.

1725124), 0.6 µL of primer mix (10 µM, forward & reverse) and 3.4 µL of RNAse-free water,

to which 1 µL of cDNA was added. All measurements were performed using three technical

replicates, in 384-well plates, on a Roche LightCycler ® 340 thermocycler. The qPCR was

programmed as follows : a first step of polymerase activation and DNA denaturation at 95°C

for 5 min; then the cycle commenced with further denaturation at 95°C for 10 sec, an

annealing step at Tm for 20 sec and an extension step at 72°C for 30 sec; after 40 cycles, a

final melting curve was produced: from 65°C to 97°C, with 5 acquisitions per °C. The melting

temperature (Tm) for each primer set was adjusted to optimise the amplification specificity

and their efficiency was validated beforehand and can be found with primer sequences in

Supplementary Table S2. Using a dilution range of a cDNA mixture, each primer set's

efficiency was calculated using the slope of the x=log(dilution); y=Ct, in the following

equation: $(10^{-1/(slope)}-1)*100$, which should be in the 90-110% range. A melting curve

was produced at the end of each qPCR program to ensure that a single gene was amplified.

In some cases, the amplicon was verified by gel electrophoresis.

<u>Data analysis.</u> After RNA extraction and RT, two samples were discarded from further

analysis due to experimental mishaps, resulting in 6 replicates in the "miPEP" and "water"

conditions and 7 replicates in the "scrambled" condition. Quantification of pri-miR159c only

succeeded in 6 out of 7 "scrambled" replicates and 5 out of 6 "water" replicates. Bacterial

genes, *i.e.* Lys R, phosphatidate and alpha-macroglobulin, were quantified relative to RecA

and GyrA reference genes, whereas the plant pri-miRNA159c was quantified relative to a

plant reference gene, cyclophilin. The number of PCR cycles necessary for the fluorescence

to emerge from the background noise is referred to as the "*Ct*". Noise was automatically

determined by the LightCycler 480 software and *Ct*s were calculated with the Abs Quant/2nd

Derivative Max - high confidence mode. A coefficient of variation (CV) was calculated for the

three technical replicates and a threshold of 2% was applied : extreme technical replicates

were excluded, if this limit was exceeded. The remaining technical replicates were averaged

to create the *Ct* value for each sample. The analysis of pri-miR159c expression was done

using the Pfaffl method, which takes into account the efficiency of each primer set [13]. This

method calculates a relative expression ratio based on the difference of *Ct*s of a gene of

interest in a sample *versus* a calibrator, and in comparison, with a reference gene. The

analysis of bacterial genes relied on two reference genes, which called for a slightly modified

version of the Pfaffl method. To calculate a relative expression ratio, we had to normalise our

qPCR data using the geometric mean of both reference genes [14]. In both cases, we

averaged the samples treated with water to create a calibrator.

<u>Statistical analyses.</u> Statistical analyses were performed on the resulting relative expression

ratios, comparing group-to-group, using a Welch Two Sample t-test or a Wilcoxon rank sum

exact test, depending on if the data were normal.

Effect of miRNAs on the bacterial community

Arabidopsis mutant experiment

Plant growth and mutant description. Five *A. thaliana* mutants were grown in a mix of soil

and sand (2:1) for a month, in individual pots. Mutants were provided by Hervé Vaucheret

and Taline Elmayan (IJPB, INRAE, Versailles, France). *RTL1* mutant over-expresses RTL1

protein which results in a suppression of siRNA pathway without affecting miRNAs [15].

RTL1myc over-expresses RTL1 protein flagged with Myc epitope, rendering RTL1 less

active, so siRNA pathway is less suppressed than with *RTL1* mutant. *Ago1-27* mutant has

AGO protein function partially impaired and is completely post-transcription gene silencing

(PTGS) deficient [16]. *Dcl1-2* mutant has total loss of function of DCL1 protein resulting in

low levels of miRNA and developmental problems [17]. *Hen1-4* mutant is miRNA defective

but is also affected in some siRNA – PTGS [18]. HEN1 methylates siRNA and miRNA to

maintain their levels and size, but also to protect them from uridylation and subsequent

degradation [19].

DNA extraction, sequencing and qPCR. We grew *A. thaliana* mutants and wild-type plants in

quadruplicate for a month and sampled the root, rhizoplane and closely adhering

rhizosphere. Two of the wild-type plants were lost during the experiment and subsequent

analyses. DNA was extracted using NucleoSpin Plant II kit (Macherey-Nagel). DNA was sent

for 16S rRNA gene amplicon (primers 341F: 5'- CCTACGGGNGGCWGCAG- 3' and 534R:

5'- ATTACCGCGGCTGCTGGCA – 3') sequencing on an Illumina MiSeq in paired-end mode

(2x250 bp) at the Centre d'expertise et de services Génome Québec (Montreal, Canada). In

parallel, 16S rRNA gene was quantified by qPCR, using the same primers as for

sequencing. Within each well was 0.1 µL of each primer (10 µM), 4 µL LightCycler® 480

SYBR Green I Master (final volume 6 µL PCR mix) and 2 µL of DNA (25 ng.µL^{-1}). We used

the following 40-cycle program: pre-incubation (95°C- 4 min) / amplification (95°C – 30 s ;

49°C – 1 min ; 72°C – 1 min) / 72°C-10 min/ melting curve (95°C – 5 s ; 49°C – 1 min ; 97°C

– continuous 5 measures/ °C) / cooling (40°C – 30 s). At the last step of amplification, a

single measurement was performed and then continuously during the melting curve step. A

minimum of 3 technical replicates was quantified. The number of copies of 16S rRNA gene

was determined in comparison with a standard curve using serial dilutions of plasmids with

cloned fragments ($R^2 = 0.994$).

<u>Amplicon sequences processing.</u> Amplicon sequencing data was processed with

AmpliconTagger [20]. Briefly, contaminants and unpaired reads were removed, and

remaining sequences were trimmed to remove adaptors and primers. Reads were then

filtered for quality control such that reads having at least one N or having average phred

score quality less than 20 were left out. A total of 3,776,292 reads passed the quality control

and 1,230,944 sequences were successfully processed to generate ASVs (DADA2) [21].

ASVs were filtered for chimaeras using DADA2's internal removeBimeraDeNovo (method =

'consensus') workflow followed by VSEARCH's [22] UCHIME *de novo*. Each remaining ASV

was assigned a taxonomic lineage by using the RDP classifier with the SILVA [23] R138.

ASVs assigned to bacterial/archaeal and rendered into ASV abundance tables.

<u>Microbial community analyses.</u> Analyses of ASVs composing the microbial communities

were performed using the R package "phyloseq" v 1.32.0 [24]. After checking the rarefaction

curves, all samples seemed sufficiently sequenced. For Shannon diversity index calculation,

all reads were rarefied to the lowest number of reads found in a sample. Significance of

differences in alpha-diversity between mutants was tested using a linear model, whilst

checking the normal distribution of residuals and using wild-type (WT) samples as a

reference. If residuals did not follow a normal distribution, a general linear model (GLM) with

gamma distribution and "log" or "link" scale were used. All tests performed were, by default,

two-sided.

<u>Statistical analyses.</u> The effect of miRNA/siRNA mutation in *A. thaliana* on the structure of

microbial communities was visualized using Principal Coordinate Analysis (PCoA)

ordinations of Bray-Curtis dissimilarities. Permutational Multivariate Analysis of Variance

(Permanova) was used to determine the statistical impact of miRNA/siRNA mutations on the

microbiota structure, alongside checking group dispersion and pairwise tests.

miPEP experiment

<u>Plant growth and miPEP treatment.</u> *Arabidopsis thaliana* Col-0 were sowed on a mix of

potting soil and sifted sand (<2.2 mm) (ratio 2:1), in a greenhouse. After germination,

seedlings were transferred into individual pots (3 cm diameter, 5 cm depth) that were moved

to a growth chamber under 16 h of daylight, 8 h of darkness, at 20°C. When the plants

reached the 6-leaf stage, the miPEP treatments started. Plants (16 replicates/condition)

were treated with 500 µL of water (control condition) or a miPEP solution (20 µM of

miPEP159a: MTWPLLSLSFLLSKYV, miPEP159b: MGLRKVLEMNTIFDSLFLSH or

miPEP159c: MQNLRVHVFLIESARC), applied at the base of the crown, 3 times a week for a

total of 10 applications. After each application, the trays in the chamber were moved around

to diminish any border effect.

<u>Sampling & DNA-RNA extraction.</u> Once the treatments were over, each individual pot was

spilled on a sieve and the aerial part was separated from the root system. The roots of two

plants were pooled together and briefly rinsed in 10 mL of PBS (1X). The roots and the

attached soil were then separated from the rinsing solution, using a funnel and a sterile

compress, and were subsequently dried using absorbent paper. The dried roots and

rhizosphere were flash frozen in liquid nitrogen and stored at -80°C. DNA was extracted

using the NucleoSpin® Plant II mini kit (MACHEREY-NAGEL, Düren, Germany) whereas

RNA was isolated with a homemade protocol [25]. The integrity and quantity of the extracted

DNA and RNA was estimated using Nanodrop and by running a 1% agarose gel. Amplicon

sequencing, processing, qPCR quantification, and analyses were performed as described

above.

Simplified soil community experiment

<u>Soil microbes enrichment.</u>

The soil microbes were enriched from five different media: 1) Tryptic Soy Broth, 2) Potato

Dextrose Broth, 3) Minimal medium + Carbon solution + NH4NO3, 4) Minimal medium +

Carbon solution + urea and 5) Minimal medium + Carbon solution + amino acids. The

minimal media were supplemented with a solution of artificial carbon-rich root exudates: 20

mM glucose 20 mM fructose, 12 mM sucrose, 20 mM lactic acid, 12 mM citric acid, 16 mM

succinic acid. For these media, the final concentration was 0.5 g/L of C and 0.1 g/L of N. To

culture the soil microbes, 2 g of sieved agricultural soil (sampled at the Armand-Frappier

experimental field: 45.5416N, -73.7173E) was incubated in 20 mL of each media, for 28 h

(200 rpm, 25 ⁰C). The cultures were, filtered 30 µm, normalized to the same optical density,

pelleted (4 ⁰C, 15 min, 4700 g), suspended in PBS, pooled and aliquoted into sterile

cryotubes. A cryoprotective solution 2x (0.6% (w/v) Tryptic Soy Broth, 10% (v/v) DMSO and

2% (w/v) trehalose) was added to the cultures (1:1 (v/v)) and the cryotubes were gently

mixed (5x inverting), left to equilibrate for 20 min and placed at -80 ⁰C [26]. Cells were

revived, inoculated in 5 mL of TSB overnight, washed, normalized to OD600 0.200 and

inoculated in a 96-wells plate containing a mixture of 17 amino acids as nitrogen source and

artificial root exudates as a carbon source.

<u>Microbial growth and miRNA exposure.</u>

To investigate the potential modification of microbial activity by plant miRNAs, we revived,

cultured overnight (in 5 mL of TSB, 28 ⁰C, 200 rpm), washed (2x) and normalized the soil

microbes (OD600 0.200). We then cultured them in a medium containing an equimolar

mixture of 17 L-amino acids (15 mM), artificial root exudates (15 mM) and a miRNA

treatment that had a final concentration of 10 µM (five biological replicates were prepared).

The miRNA treatment consisted of an equimolar mix of five miRNAs (ath-miR158a-3p, ath-

miR158b, ath-miR159a, ath-miR827, and ath-miR5642b) of either the plant mimics or

scrambled controls (Supplementary Table S1). Microbial activity was quantified by

introducing a tetrazolium dye, which undergoes a color change to purple in the presence of

dehydrogenases, subsequently intensifying the optical density at 600 nm. Optical density

measurements were recorded hourly using a plate reader. The experiment was concluded

after 52 hours.

<u>16S rRNA gene amplicon sequencing.</u>

To determine whether alterations in microbial activity were indicative of changes within the

bacterial community, we proceeded with a microvolume physical DNA extraction [27] from

the cultures and prepared the libraries to sequence the V4-V5 region of the 16S rRNA gene

(primers 515F-Y: GTGYCAGCMGCCGCGGTAA and 926R: CCGYCAATTYMTTTRAGTTT)

[28] using the Illumina MiSeq platform at the Centre d'expertise et de services de Génome

Québec, Montreal, Canada. The sequences were processed with the AmpliconTagger

pipeline [20], and analysed as described above.

References

1031

1.   Bolger AM, Lohse M, Usadel B. Trimmomatic: a flexible trimmer for Illumina
sequence data. Bioinformatics. 2014;30:2114-2120.
2.   Kozomara A, Birgaoanu M, Griffiths-Jones S. miRBase: from microRNA sequences
to function. Nucl Acids Res. 2019;47:D155-D162.
3.   Li H, Durbin R. Fast and accurate short read alignment with Burrows-Wheeler
transform. Bioinformatics. 2009;25:1754-60.
4.   Quinlan AR, Hall IM. BEDTools: a flexible suite of utilities for comparing genomic
features. Bioinformatics. 2010;26:841-842.
5.   Levy A, Salas Gonzalez I, Mittelviefhaus M, Clingenpeel S, Herrera Paredes S, Miao
J et al. Genomic features of bacterial adaptation to plants. Nat Genetics. 2017;
6.   Dai X, Zhuang Z, Zhao PX. psRNATarget: a plant small RNA target analysis server
(2017 release). Nucl Acids Res. 2018;
7.   Axtell MJ. Classification and Comparison of Small RNAs from Plants. Ann Rev Plant
Biol. 2013;64:137-159.
8.   Pearson WR. Finding Protein and Nucleotide Similarities with FASTA. Curr Protoc
Bioinformatics. 2016;53:3 9 1-3 9 25.
9.   Jones P, Binns D, Chang H-Y, Fraser M, Li W, McAnulla C et al. InterProScan 5:
genome-scale protein function classification. Bioinformatics. 2014;30:1236-1240.
10.   Manjang K, Tripathi S, Yli-Harja O, Dehmer M, Emmert-Streib F. Graph-based
exploitation of gene ontology using GOxploreR for scrutinizing biological significance.
Sci Rep. 2020;10:16672.
11.   Tremblay J, Schreiber L, Greer C. High resolution shotgun metagenomics the more
data the better? Brief Bioinfo. 2022;23:bbac443.
12.   Han J-I, Choi H-K, Lee S-W, Orwin PM, Kim J, LaRoe SL et al. Complete genome
sequence of the metabolically versatile plant growth-promoting endophyte *Variovorax*
paradoxus S110. J Bacteriol. 2011;193:1183-1190.
13.   Pfaffl MW. A new mathematical model for relative quantification in real-time RT-PCR.
Nucl Acids Res. 2001;29:e45.
14.   Vandesompele J, De Preter K, Pattyn F, Poppe B, Van Roy N, De Paepe A et al.
Accurate normalization of real-time quantitative RT-PCR data by geometric
averaging of multiple internal control genes. Genome Biol. 2002;3:RESEARCH0034.
15.   Shamandi N, Zytnicki M, Charbonnel C, Elvira-Matelot E, Bochnakian A, Comella P
et al. Plants Encode a General siRNA Suppressor That Is Induced and Suppressed
by Viruses. PLoS Biol. 2015;13:e1002326.
16.   Morel JB, Godon C, Mourrain P, Béclin C, Boutet S, Feuerbach F et al. Fertile
hypomorphic ARGONAUTE (ago1) mutants impaired in post-transcriptional gene
silencing and virus resistance. Plant Cell. 2002;14:629-639.
17.   Xie Z, Kasschau KD, Carrington JC. Negative feedback regulation of Dicer-Like1 in
Arabidopsis by microRNA-guided mRNA degradation. Curr Biol. 2003;13:784-9.
18.   Boutet S, Vazquez F, Liu J, Béclin C, Fagard M, Gratias A et al. Arabidopsis HEN1: a
genetic link between endogenous miRNA controlling development and siRNA
controlling transgene silencing and virus resistance. Curr Biol. 2003;13:843-848.
19.   Li J, Yang Z, Yu B, Liu J, Chen X. Methylation protects miRNAs and siRNAs from a
3'-end uridylation activity in Arabidopsis. Curr Biol. 2005;15:1501-7.
20.   Tremblay J, Yergeau É. Systematic processing of rRNA gene amplicon sequencing
data. GigaScience. 2019;8:giz146.
21.   Callahan BJ, McMurdie PJ, Rosen MJ, Han AW, Johnson AJA, Holmes SP. DADA2:
high-resolution sample inference from Illumina amplicon data. Nat Meth.
2016;13:581.
22.   Rognes T, Flouri T, Nichols B, Quince C, Mahé F. VSEARCH: a versatile open
source tool for metagenomics. PeerJ. 2016;4:e2584.

23.   Quast C, Pruesse E, Yilmaz P, Gerken J, Schweer T, Yarza P et al. The SILVA
ribosomal RNA gene database project: improved data processing and web-based
tools. Nucl Acids Res. 2012;41:D590-D596.
24.   McMurdie PJ, Holmes S. phyloseq: an R package for reproducible interactive
analysis and graphics of microbiome census data. PLoS One. 2013;8:e61217.
25.   Griffiths RI, Whiteley AS, O'Donnell AG, Bailey MJ. Rapid method for coextraction of
DNA and RNA from natural environments for analysis of ribosomal DNA- and rRNA-
based microbial community composition. Appl Environ Microb. 2000;66:5488–5491.
26.   Kerckhof FM, Courtens EN, Geirnaert A, Hoefman S, Ho A, Vilchez-Vargas R et al.
Optimized cryopreservation of mixed microbial communities for conserved
functionality and diversity. PLoS ONE. 2014;9:e99517.
27.   Bramucci AR, Focardi A, Rinke C, Hugenholtz P, Tyson GW, Seymour JR et al.
Microvolume DNA extraction methods for microscale amplicon and metagenomic
studies. ISME Commun. 2021;1:79.
28.   Parada AE, Needham DM, Fuhrman JA. Every base matters: assessing small
subunit rRNA primers for marine microbiomes with mock communities, time series
and global field samples. Environ Microbiol. 2016;18:1403-1414.
1100
1101

1102 **Supplementary Table S1**: Sequences of the single-stranded synthetic miRNAs.

1103

miRNA	Sequences
ath-miR158a-3p	UCCCAAAUGUAGACAAAGCA
ath-miR158b	CCCCAAAUGUAGACAAAGCA
ath-miR159a	UUUGGAUUGAAGGGAGCUCUA
ath-miR159b-3p	UUUGGAUUGAAGGGAGCUCUU
ath-miR159c	UUUGGAUUGAAGGGAGCUCCU
ath-miR161.1	UGAAAGUGACUACAUCGGGGU
ath-miR165b	UCGGACCAGGCUUCAUCCCCC
ath-miR827	UUAGAUGACCAUCAACAAACU
ath-miR5642b	UCUCGCGCUUGUACGGCUUU
sc-ath-miR158a-3p	CAACAAGAUGACAAUGUCAC UGAAAACAUACAUCAGACCG*
sc-ath-miR158b	GACAAGCUCCACAUACAGAA
sc-ath-miR159a	AGGGAAGUAUUGUCGUGACUU AGAGAGACCUUGGGUUGAUUU**
sc-ath-miR159b-3p	UGGGGACUGUAUCUUAGUUAG
sc-ath-miR159c	CAGGCCUUGGUGUUAUAUGAG
sc-ath-miR161.1	CUGCAGCUAGUUGGGUAAAGA
sc-ath-miR165b	CUCCACCGAGGUCCUCCACGU
sc-ath-miR827	GAUCCAAAAGCUAAUUUCAC
sc-ath-miR5642b	ACGUCGUAAAACUUCACGUG

1119

1120 * and **: Scramble miRNAs used in the simplified soil community experiment
1121 ** Scramble miRNA ath-miR159a used during confocal microscopy imaging and flow cytometry.
1122 Note: all sequences were methylated at their 3' end.

1123

1124

1125

1126 **Supplementary Table S1**: Primers used for qPCR in *Arabidopsis thaliana* and *Variovorax*

1127 *paradoxus* EPS for the *in vitro* miPEP transcriptomic experiment. Most primers were

Amplified gene	Amplicon size	Forward primer 5'-3'	Reverse primer 5'-3'	Tm	Reference
pri-miR159c	178bp	AGCTCCTTTTCTTCTTCTCTTAAT	CGTCTTCTCGTAAATAAACAACATT	53°C	Allen, *et al*., 2010
cyclophilin	62bp	TGGACCAGGTGTACTTTCAATGG	CCACTGTCTGCAATTACGACTTTG	53°C	Allen, *et al*., 2010
LysR	96bp	GTAGGTGAAGCGGGTGAGG	TCAAGGTGAAAGCCCACGTC	53°C	from EPS genome
phosphatidate	90bp	TTCATCAGCCTGGCCTACAC	CGGGGCAGTACTTCCTGATC	54°C	from EPS genome
alpha-macroglobulin	82bp	CAGGCTGGTGATCTGAAGGG	AATGAGGGAACGAAACGGGTC	53°C	from EPS genome
RecA	99bp	CCCTTCTTGATGGTGCCGAT	TGATGTTCGGTTCGCCTGAA	57°C	from EPS genome
GyrA	93bp	CCGAACTCCTGCTTGATGGT	CAGAGATCGACGACCTGCTC	57°C	from EPS genome

1128 designed based on the EPS genome, except for plant gene primers.

1129

1130

1131 Allen R. S., Li J., Alonso-Peral M. M., White R. G., Gubler F., Millar A. A. (2010). MicroR159

1132 regulation of most conserved targets in *Arabidopsis* has negligible phenotypic effects.

1133 Silence 1:18. doi: 10.1186/1758-907X-1-18

1134