Physiological traits and their relationships vary along an aridity gradient

within and among Fijian bee species

Carmen RB da Silva[1]*, Julian E Beaman[2], Marika Tuiwawa[3], Mark I Stevens[4,5],

Rosalyn Gloag[6], Vanessa Kellermann[1], Lesley A Alton[1]

[1]*School of Biological Sciences, Monash University, Melbourne, Victoria, Australia*

[2] *College of Science and Engineering, Flinders University, Bedford Park, South*

Australia, Australia

[3]*The University of the South Pacific Herbarium, Viti Levu, Fiji*

[4]*Securing Antarctica's Environmental Future, Biological Sciences, South Australian*

Museum, Adelaide, South Australia, Australia

[5]*School of Biological Sciences, University of Adelaide, South Australia, Australia*

[6]*School of Life and Environmental Sciences, University of Sydney, Sydney, New*

South Wales, Australia

Corresponding author: carmen.dasilva@monash.edu

25 **Key words:** metabolic cold adaptation, discontinuous gas exchange, hygric

26 hypothesis, bees, thermal performance, co-variation, metabolic rate, ventilation rate

Abstract

Temperature and water availability are independently hypothesised to be important
abiotic drivers of the evolution of metabolic rates and gas exchange patterns,
respectively. Specifically, the metabolic cold adaptation hypothesis (MCA) predicts
that cold environments select for faster metabolic rates to counter the thermodynamics
of biochemical reactions while the hygric hypothesis predicts that dry environments
select for discontinuous gas exchange to reduce water loss. Although these two
hypotheses consider different physiological traits and how they vary along different
abiotic gradients, metabolic rate drives gas exchange patterns in insects meaning
these two traits are inherently linked. Despite this link, the MCA and hygric hypotheses
are rarely considered together and the extent to which metabolic rates and respiratory
patterns vary and co-vary with temperature and aridity along climatic gradients
remains unclear. We tested the MCA and hygric hypotheses within a species of
endemic Fijian bee, *Homalictus fijiensis*, across an altitudinal gradient of 1100 m, and
among four Fijian bee species, including *H. fijiensis*, that inhabit different altitudinal
bands. In Fiji, environmental temperature is ~5°C lower in the central highlands than
in the coastal lowlands with the highlands receiving ~100 mm of additional precipitation
than the lowlands each month. We found an MCA-like pattern within *H. fijiensis* and
among Fijian bee species, where metabolic rate decreased with increasing
temperature, but precipitation also explained variation in metabolic rate. We also found
support for the hygric hypothesis within *H. fijiensis* and across species. Lastly, we
found that the relationship between metabolic rate and ventilation rate changed
depending on precipitation of the driest month within *H. fijiensis* and among bee
species. Bees that inhabited wet regions had low ventilation rates of greater volume
but bees that inhabited drier locations had rapid ventilations of smaller volume
potentially as a mechanism to exchange gas without desiccating. As the correlation
between metabolic rate and ventilation rate changes across the precipitation gradient,
it is possible that these two traits can evolve independently of each other in response
to abiotic variables despite being positively correlated.

Introduction

Variation in climate shapes species' physiological traits and geographic distributions [1–3]. For terrestrial ectotherms in particular, environmental temperature and aridity are hypothesised to be important abiotic drivers of the evolution of key physiological traits such as metabolic rate and respiratory patterns [4,5]. These two traits underpin the adaptive hypotheses known as the metabolic cold adaptation (MCA) and hygric hypotheses, which aim to explain how and why metabolic rate and respiratory patterns, respectively, vary among populations and species that occupy different climatic niches [5–10]. However, studies that seek to understand the environmental drivers of variation in these two traits rarely consider the MCA and hygric hypotheses together (but see [9,11]), despite the knowledge that increases in metabolic rate result in higher rates of ventilation [12–14]. Given the inherent link between metabolic rate and respiratory pattern, it is plausible that these two traits have co-evolved meaning that selection on metabolic rate has potentially influenced respiratory patterns, and vice versa. However, even though these traits are linked, selection could be acting on each trait independently, causing a change in the relationship between metabolic rate and ventilation rate across an environmental gradient. Thus, to better understand the evolution of one trait requires consideration of the other to determine how environments have shaped trait variation and covariation.

The metabolic cold adaptation hypothesis predicts that organisms from cold environments (such as high latitudes and altitudes) will have higher metabolic rates than those from warm environments, when measured at the same temperature. This pattern is expected to arise as a mechanism to maintain physiological function in cold environments because acute exposure to colder temperatures causes biochemical reaction rates to slow [5,6,15,16]. However, in terrestrial organisms, the pattern described by the MCA hypothesis may also emerge as a consequence of organisms depressing their metabolic rate in response to high temperatures to prevent desiccation [17]. This is because high air temperatures result in high vapour pressure deficits (VPDs) that can lead to higher rates of water loss from wet respiratory surfaces [10]. Organisms living in warmer environments that are also more desiccating could therefore slow their metabolic rates, reduce their ventilation rate, and breathe discontinuously (that is, alternate between periods of ventilation and breath-holding).

The hygric hypothesis makes the explicit prediction that discontinuous gas exchange
in resting tracheate arthropods, such as insects, is an adaptation to limit respiratory
water loss. As such, drier environments are expected to favour arthropods that breathe
discontinuously, and by extension, those that hold their breath for longer (i.e., have a
lower ventilation rate) [4,9,18].

In insects, both the MCA and hygric hypotheses tend to be supported by broad-scale
comparative analyses [6,18], but not always [19]). In contrast, experimental evolution
studies have been unable to replicate the macrophysiological pattern described by the
MCA hypothesis by manipulating temperature in controlled laboratory environments
[7,20]. Such research indicates that temperature alone may not be responsible for
driving the evolution of metabolic rate. The hygric hypothesis similarly suffers from
equivocal support [21], and alternative adaptive hypotheses propose that
discontinuous gas exchange has evolved as a mechanism to improve gas exchange
in underground environments, or to reduce oxidative tissue damage [8,22–24]. It has
also been suggested that discontinuous gas exchange is not an adaptation to
environmental conditions, but instead arises as a consequence of neural control of
gas exchange during rest [25,26]. Given that there is mixed support for both the MCA
and hygric hypotheses, and that selection rarely acts on single traits independently
[27–29], we propose that these hypotheses should be considered simultaneously to
better understand how metabolic rate and respiratory patterns vary and co-vary along
climatic gradients.

Here we examine how metabolic rate and respiratory patterns vary and co-vary within
and among bee species that breathe discontinuously [30] and exist along an altitudinal
gradient on the tropical mountainous island of Viti Levu, Fiji. Altitudinal gradients are
powerful for testing hypotheses concerning climatic drivers of trait variation because
the spatial rate of change in climate is greater along altitudinal gradients than
latitudinal gradients, while habitat composition and photoperiod vary less [31]. Viti
Levu is divided by a mountain range that runs north to south and the highest peak is
1328 m above sea level (a.s.l). The highlands in the centre of the island are
characterised by cloud forests that are ~ 5°C cooler and receive ~100 mm less
precipitation per month than the coastal lowlands (Fig. 1). Due to the mountain range
that divides Viti Levu, there is also a rain shadow on the western side of the island that

receives significantly less rainfall than the eastern side (Fig. 1c). The bee species that inhabit Viti Levu occupy different altitudinal ranges: the endemic *Homalictus fijiensis*, Perkins & Cheesman (1928), is found along the entire altitudinal gradient while 22 other endemic *Homalictus* species are restricted to the highlands, and the invasive *Braunsapis puangensis*, Cockerell (1928), is generally restricted to the lowlands [32–34].

Figure 1. Abiotic variables across Viti Levu, Fiji: a) altitude (m asl), b) temperature of the coolest and driest month (July) (°C), c) precipitation of the driest month (July) (mm) and, d) vapour pressure deficit (July) (kPa). e) Collection altitude of each species (Hf: *Homalictus fijiensis*, Bp: *Braunsapis puangensis*, Hg: *Homalictus groomi*, Ht: *Homalictus tuiwawae*) where circle size indicates magnitude of sample size. f) Relationship between altitude and environmental temperature, g) relationship between altitude and precipitation, h) relationship between altitude and vapour pressure deficit (VPD). Filled circles indicate collection sites on the wet side of Viti Levu and open circles indicate sample sites on the dry side of Viti Levu.

To examine the climatic drivers of variation and co-variation in metabolic rate and respiratory patterns of bees, we collected individuals of *H. fijiensis*, *H. groomi*, Dorey et al., (2019), *H. tuiwawae*, Dorey et al., (2019), and *B. puangensis* from multiple sites on either side of the rain shadow along the altitudinal gradient of Viti Levu (Fig. S1e). We measured the metabolic rates of 203 individual bees at 25°C as rates of carbon dioxide production using flow-through respirometry. Of those bees that displayed discontinuous gas exchange, which was all but four individuals, we calculated their ventilation rate. We examined the relationships between the metabolic rate and

ventilation rate of bees and the climatic variables at their collection site using a strong inference framework whereby a set of models were selected *a priori* based on the explicit predictions of the MCA and hygric hypotheses (see Table 1 & Supplementary Box 1 for list of models tested). We accepted support for the MCA hypothesis if bees from cooler climates had higher metabolic rates and the only climate variable in the best model was temperature (Figure 2A). We accepted support for the hygric hypothesis if bees from drier climates had lower ventilation rates and the only climate variable in the best model was either vapour pressure deficit (VPD) or precipitation (Figure 2B), both of which are assumed to be indicative of aridity. In addition to testing the predictions of the MCA and hygric hypothesis, we also tested the hypothesis that ventilation rate and metabolic rate will be positively correlated, but that the strength of the relationship between these two traits will vary depending on the aridity of the environment (Figure 2C). All else being equal, insects with higher metabolic rates breathe more frequently [12–14], but we expect that in drier environments there will be stronger selection for insects with high metabolic rates to maintain lower ventilation rates to minimise water loss, which will result in a shallower relationship between ventilation rate and metabolic rate. In wetter environments where the risk of desiccation is lower, we expect that selection on low ventilation rate will be relaxed resulting in a steeper relationship between ventilation rate and metabolic rate.

Figure 2. A) Conceptual illustration of the relationship between metabolic rate and temperature along an altitudinal gradient if the metabolic cold adaptation (MCA) hypothesis is supported. B) Relationship between ventilation rate and aridity (precipitation or inverse vapour pressure deficit (VPD)) if the hygric hypothesis is supported. C) Change in relationship between metabolic rate and ventilation rate depending on ardity (precipitation or VPD). Where there is reduced selective pressure to avoid desiccation, we expect that selective pressure on metabolic rate and ventilation rate will be relaxed, resulting in a steep relationship between metabolic rate and ventilation rate in areas of high precipitation or low VPD. *Homalictus fijiensis* photograph by James B. Dorey.

Methods

Animal collection

Bees were collected over 10 days in September 2019 on Viti Levu, Fiji. Viti Levu is divided by a mountain range that runs north to south, which creates a rain shadow on the western side of the island that receives significantly less rainfall than the eastern side (Fig. S1C). We collected bees from 16 sites along an altitudinal gradient of 6–1072 m above sea level (a.s.l.), with 10 of these sites being on the dry western side of the island and the other 6 sites being on the wet eastern side. Sites were sampled in a haphazard order over the 10 days and bees were collected by sweep-netting vegetation along road verges or in native vegetation. Individuals of *Braunsapis puangensis* were also collected directly from stem nests. The males and females of each species collected are known to be active year-round and are multivoltine (as per observation of M.P. Schwarz on multiple previous research trips to Fiji). Immediately upon capture, bees were placed in individual vials and stored in an insulated container. Inside the vials, bees were provided with a small piece of paper towel dipped in a 20% sucrose solution. On the day of collection, bees were transported to the laboratory to be measured that same evening.

Climatic variables

For each collection site, we downloaded historical climate data (1970-2000) at a 30-arc-second resolution from WorldClim2 [35] using the R package *raster* [36]. Since the MCA and hygric hypotheses make explicit predictions about how cool and dry climates shape metabolic rates and ventilation rates, respectively, we characterised the climates at each collection site using the variables of mean temperature (°C) of the coolest month (T_E) (July), mean precipitation (mm) of the driest month (July), and mean vapour pressure (VP, kPa) of the driest month (July) (Figure 1). VP was used to calculate the vapour pressure deficit (VPD, kPa) (drying power of the air) for bees during the driest month assuming that their respiratory surfaces are saturated with

water vapour (saturated water vapour pressure, SVP, kPa) and that their body
temperature matches that of the environment (T_E, mean temperature of the driest and
coolest month, °C) using the following equations:

$$SVP = 0.611exp\left(\frac{17.27 \times Te}{Te + 237.3}\right), [37] \tag{1}$$

$$VPD = SVP - VP \tag{2}$$

The range of environmental variables that each species was collected from is shown
in Supplementary Table 1.

Trait measurements

We measured the metabolic rate and ventilation rate of individual bees by measuring
their rates of carbon dioxide production (VCO_2, µl h^{-1}) at 25°C using a seven-channel
flow-through respirometry system. The respirometry system was supplied with room
air that was pushed through columns of soda lime and Drierite® to remove carbon
dioxide and water vapour, respectively. Flow rates through each of the seven channels
was regulated nominally to 100 mL min^{-1} by a mass flow controller (Aalborg, Model
GFC17, Orangeburg, NY, USA). The volumetric flow rates produced by the flow
controllers was measured using a Gilian Gilibrator-2 NIOSH Primary Standard Air Flow
Calibrator with a low-flow cell (Sensidyne, LP, St Petersburg, FL, USA) and corrected
to standard temperature pressure, STP (101.3 kPa and 0°C). To prevent bees from
desiccating during measurements, the air was re-humidified after the flow controllers
by passing the air through a humidifying chamber (a syringe of wetted cotton). The
humidified air then flowed through a respirometry chamber (2.5 ml plastic syringe)
containing an individual bee. Respirometry chambers were situated inside a
temperature-controlled cabinet that maintained air temperature to 25 ± 1°C and kept
bees in the dark. The excurrent air from the respirometry chamber then flowed through
one of seven infrared CO_2/H_2O gas analysers (LI-COR, Model LI-840A, Lincoln,
Nebraska, USA) that were calibrated with precision span gases (5.0 and 30.4 ppm

CO$_2$, Alphagaz – Air Liquide, Victoria, Australia) and measured CO$_2$ concentrations at a sampling rate of 1 Hz.

The fractional CO$_2$ concentration of the excurrent air (F$_{ECO2}$) from each respirometry chamber was recorded for 50 min. The fractional CO$_2$ concentration of the incurrent air when it was stable was measured for 2 min before and after each 50-min measurement period with an empty chamber, and a linear model was fitted to these data to estimate the CO$_2$ concentration of the incurrent air during the 50-min measurement period (F$_{iCO2}$). The rate of carbon dioxide production (VCO$_2$, µl h^{-1}) was then calculated using equation 3 (Lighton 2018) where FR is the flow rate (µl h^{-1}) corrected to STP (101.3 kPa and 0°C) accounting for water vapour dilution:

$$V_{CO2} = FR(F_{ECO2} - F_{iCO_2}) \qquad (3)$$

The first 15 min of the 50-min measurement period was considered a settling period and was discarded from analysis. The lowest averaged VCO$_2$ over 20 min during the final 35 min of the measurement period was taken as the measure of metabolic rate for each individual. Bees that breathed discontinuously produced traces of carbon dioxide production that showed periods of minimal gas exchange interspersed by peaks in carbon dioxide production (Fig. S2). The number of peaks of carbon dioxide production observed during the 20-min period was used to calculate the ventilation rate (h^{-1}) of the bee (Supplementary Figure 1). In all, 96% of bees exhibited discontinuous gas exchange patterns. Those that had ventilations rates greater than 120 h^{-1} (n = 8) were removed from the analysis as it is likely they were stressed [24].

Immediately following metabolic rate measurements, bees were preserved in 100% ethanol to be sexed and weighed at a later date. Bees were sexed by visual assessment by examining the number of antennal segments (males have 13 and females have 12). Prior to weighing, the right hind leg of each individual was detached for species identification (see below). Dry mass was measured two months after metabolic rate measurements with specimens dried at 60°C for 48 h prior to being weighed (XP2U Ultra Micro Balance, Mettler Toledo, Victoria, Australia).

Bees were measured during the evening on the day that bees were collected from the
field. Due to the logistical challenges of bee collection on Viti Levu, bees could not be
collected from all sites each day. Consequently, on some nights, bees from multiple
sites could be measured, but on other nights only bees from one site could be
measured. Since the species and sex of the bees measured each night were mostly
unknown (see below), measurements were conducted blind to the species and sex of
bees in most cases (*B. puangensis* is easily differentiated from *Homalictus* species).
After species were identified (see below), only those species with sample sizes greater
than 10 were retained for analysis. In total, we measured 212 individuals of four
species: *H. fijiensis* (*n* = 124), *H. tuiwawae* (*n* = 21), *H. groomi* (*n* = 16), and *B.*
puangensis (*n* = 42).
295

Species identification

Homalictus species living at high altitudes are cryptic and, in many cases, can only be
identified via male genitalia morphology, or via genetic comparison. To identify our
Homalictus species collected from altitudes above 300 m a.s.l., we sent tissue
samples (the right hind leg) to the Canadian Centre for DNA Barcoding (CCDB) at the
University of Guelph, Ontario, Canada, to have the mtDNA COI gene fragment
sequenced. A total of 84 samples were DNA barcoded using single-molecule real-time
sequencing (SMRT49) using the PacBio Sequel platform (Pacific Biosciences, Menlo
Park, CA, USA). Sequences were aligned to existing COI sequences taken from
species identified by [33,38]. The invasive bee species, *B. puangensis*, which is
generally restricted to the lowlands, is similar in size to *H. fijiensis*, but is
distinguishable by its black cuticle and white facial marking [30,32].

Statistical analyses

We examined variation and co-variation in metabolic rate and ventilation rate within *H.*
fijiensis and among species across climatic space using linear mixed models using the
nlme package [39] in R Version 4.1.0 [40]. Using a strong inference approach, we
tested suites of six models to test specific hypotheses that would support or reject the

MCA and hygric hypotheses, and to test whether the relationship between ventilation
rate and metabolic rate changes across the aridity gradient (Table 1; see rationale
behind hypothesis tests in Supplementary Box 1). The sixth model in each set was a
null model that included altitude and no other environmental variables (Table 1). All
models included $\log_{10}$ transformed dry body mass, sex and altitude as predictor
variables, as well as the hypothesis of interest variable(s). Metabolic rate was also
$\log_{10}$ transformed to ensure linear mixed effect model assumptions were met. We did
not include VPD and temperature in the same models as VPD is strongly correlated
with temperature. Models testing the hygric hypothesis included $\log_{10}$ transformed
metabolic rate as a predictor variable to account for the effect of metabolic rate on
ventilation rate as they are positively correlated. Multispecies models testing the MCA,
hygric hypothesis and the way in which metabolic rate and ventilation rate covary
included species as a factor within the models. Measurement block and channel were
included as random factors within each model.

We compared the Akaike information criterion (AIC) of each model to determine which
hypothesis best explained the variation in metabolic rate, ventilation rate and how traits
covary across abiotic variables as per [18,41] using the MuMIn package [42]. We then
evaluated the direction of the relationships between the traits of interest and abiotic
variables within the best fitting models to make inferences on whether metabolic rate
and ventilation rate are best explained by the MCA and hygric hypotheses,
respectively, and if the relationship between metabolic rate and ventilation rate
remains consistent across abiotic variables or if it becomes uncoupled. All models
included $\log_{10}$ transformed body mass, sex and altitude as predictor variables, and an
interaction between $\log_{10}$ transformed body mass and species were included within
the multispecies models. The multispecies models testing the hygric hypothesis also
included an interaction between $\log_{10}$ transformed metabolic rate and species. Only
species that had sample sizes over 10 individuals were included in the multispecies
analysis (four species). The packages ggplot2 [43] and cowplot [44] were used to
produce data figures throughout the results.

346
347
348
349

Table 1. Hypotheses that were compared to test the metabolic cold adaptation hypothesis, the hygric hypothesis, and examine if the way in which metabolic rate and ventilation rate covaries changes with aridity. **-** indicates a negative relationship for the hypothesis to be supported. **+** indicates a positive relationship for hypothesis to be supported. o indicates we expect no relationship between the response variable and the predictor variable.

Hypothesis predictor variable	Metabolic rate (MR) (testing MCA hypothesis)	Ventilation rate (VR) (testing hygric hypothesis)	VR ~ MR * predictor variable * species (Trait covariation depends on aridity)
T_{MIN} (mean temperature of coldest month, July)	-	o	o
P_{DRY} (precipitation of the driest month, July)	o	+	VR ~ MR relationship is steepest in low P_{DRY} environments
VPD (vapour Pressure deficit of the driest month, July)	o	-	VR ~ MR relationship steepest low VPD locations.
T_{MIN} + P_{DRY}	If T_{MIN} has a negative effect on MR, an MCA like pattern arises.	If P_{DRY} has a positive effect on VR, a hygric like pattern arises.	NA
T_{MIN} * P_{DRY}	If MR is highest at the lowest T_{MIN} * P_{DRY} combination an MCA like pattern will arise.	If VR is highest at the highest P_{DRY} * T_{MIN} combination, a hygric like pattern will arise.	VR ~ MR relationship steepest in cool wet locations.
Altitude (null hypothesis)	o	o	o

Results

Testing the MCA within and among species

The hypothesis that best explained variation in metabolic rate within *H. fijiensis* and across species included average temperature of the coldest month (*H. fijiensis*: estimate = -0.049 $\pm$ SE = 0.024, df = 89, t = -2.04, P = 0.045) (among species: (estimate = -0.065 $\pm$ SE = 0.026, df = 160, t = -2.54, P = 0.012), and average precipitation of the driest month as the test variables (*H. fijiensis*: estimate = -0.002 $\pm$ SE = 0.0004, df = 89, t = -3.71, P < 0.005) (among species: estimate = -0.002 $\pm$ SE = 0.0004, df = 160, t = -4.53, P < 0.005) (Table 2 and see full model summary in Supplementary Table 2 & 3). Metabolic rate was negatively correlated with both

temperature and precipitation, therefore an MCA like pattern is observed (Figure 3A &

3C), but the MCA hypothesis is not supported in its purest form. Within the *H. fijiensis*

model comparison, two other models had ΔAIC values < 2, indicating that these

models had similar model fit to the best fitting model (Table 2). However, two of the

top three models included average temperature of the coldest month and all three

included precipitation of the driest month, therefore, it is likely that both temperature

and precipitation explain variation in metabolic rate in *H. fijiensis*, as explained by the

additive top model.

Table 2. Testing the MCA within *H. fijiensis* (A) and among species (B). Model
comparison table. Best fitting model with lowest AIC value is in bold. T_e =
environmental temperature (°C), P_{dry}= mean precipitation of the driest month (mm),
VPD = vapour pressure deficit (kPa). All models also included $\log_{10}$ transformed body
mass, sex and altitude as predictor variables. wi indicates AIC model weight.

A) *Homalictus fijiensis* metabolic rate model comparison					
Model environmental factors	df	AIC	AICc	ΔAICc	wi
T_e + P_{dry}	**9**	**-128.3279**	**-126.7**	**0**	**0.53**
P_{dry}	10	-126.8611	-125.1	1.62	0.24
T_e x P_{dry}	8	-126.3840	-124.9	1.83	0.21
Altitude	7	-119.1432	-118.2	8.57	0.007
VPD	8	-117.9569	-116.7	10.04	0.004
T_e	8	-117.2654	-116.0	10.74	0.002
B) Multispecies metabolic rate model comparison					
Model environmental factors	df	AIC	AICc	ΔAICc	wi
T_e + P_{dry}	**12**	**-167.6970**	**-166.1**	**0**	**0.69**
T_e x P_{dry}	13	-165.7632	-163.8	2.22	0.23
P_{dry}	11	-163.1104	-161.7	4.33	0.08
Altitude	10	-153.1987	-152.1	14.00	0.001
VPD	11	-151.5211	-150.1	15.92	0.00
T_e	11	-151.2203	-149.8	16.22	0.00

383
384 **Figure 3.** A) *Homalictus fijiensis* residual metabolic rate (mL min⁻¹) across residual
385 average temperature of the coldest month, and B) residual precipitation of the driest
386 month (mm). C) Residual metabolic rate of multiple Fijian bee species across residual
387 temperature of the coldest month, and D) residual precipitation of the driest month.
388
389
390
Testing the hygric hypothesis within and across species

The model that best explained variation in ventilation rate within *H. fijiensis* and across

species included precipitation of the driest month (*H. fijiensis*: estimate = 0.0005 ± SE

= 0.0001, df = 89, t = 3.3, P < 0.005) (across species: estimate = 0.0004 ± SE =

0.00017, df = 160, t = 2.55, P = 0.012) (Table 3; see Supplementary Table 4 & 5 for

full model coefficient summaries). Where ventilation rate increased with precipitation,

supporting the hygric hypothesis (Figure 4).

Table 3. Testing the hygric hypothesis within *Homalictus fijiensis* (A) and across Fijian bee species (B). Best fitting model with lowest AIC value is in bold. T_e = environmental temperature (°C), P_{dry}= mean precipitation of the driest month (mm), VPD = vapour pressure deficit (kPa). All models also included $\log_{10}$ transformed body mass, sex, altitude and $\log_{10}$ transformed metabolic rate as predictor variables.

A) *Homalictus fijiensis* ventilation rate model comparison					
Model environmental factors	df	AIC	AICc	ΔAICc	wi
P_{dry}	**9**	**-345.05**	**-351.8**	**0.000**	**0.68**
T_e + P_{dry}	9	-353.43	-349.6	2.22	0.22
T_e x P_{dry}	9	-344.03	-347.3	4.55	0.07
Altitude	10	-351.57	-343.9	7.96	0.013
T_e	11	-349.66	-343.5	8.37	0.010
VPD	8	-345.14	-342.5	9.40	0.006
B) Multispecies ventilation rate model comparison					
Model environmental factors	df	AIC	AICc	ΔAICc	wi
P_{dry}	**12**	**-493.97**	**-492.3**	**0.00**	**0.56**
T_e + P_{dry}	13	-491.99	-490.1	2.26	0.18
T_e	12	-490.23	-488.6	3.47	0.08
Altitude	14	-489.99	-488.0	4.35	0.06
T_e x P_{dry}	12	-489.36	-487.8	4.57	0.057
VPD	11	-489.26	-487.6	4.71	0.053

Figure 4. A) *Homalictus fijiensis* ventilation rate (ventilations min^{-1}) model residuals
across precipitation of the driest month model residuals (mm) (for data visualisation
two outlier points were removed with residual precipitation values less than -10,
however, these individuals can be observed in panel B). B) Multispecies metabolic
rate model residuals across precipitation of the driest month (mm) model residuals (*H.*
fijiensis residual ventilation rate differs slightly in panels A and B due to different
residuals being generated from different model structures).

Does the relationship between metabolic rate and ventilation rate co-vary with abiotic

variables?

The model that best explained the the way in which ventilation rate covaries with

metabolic rate across climatic variables within *H. fijiensis* and across bee species

included an interaction between log$_{10}$ transformed metabolic rate and precipitation of

the driest month as the test variables (*H. fijiensis*: estimate = -0.0004 $\pm$ SE = 0.0005,

df = 88, t = -0.81, P = 0.41) (among species: estimate = -0.0009 $\pm$ SE = 0.0005, df =

159, t = -1.68, P = 0.096) (see Supplementary Tables 6 & 7 for full model coefficient

summaries). Despite the non-significant interaction term between metabolic rate and

precipitation of the driest month in both the within *H. fijiensis* and across bee species

models, these models had notably lower AICc values than all other test hypotheses

(Table 4). The negative model estimate values stated above indicate that bees that

inhabit drier environments have higher ventilation rates per mL of CO_2 exchanged per

minute than bees that inhabit wetter environments within species and among species

(Figure 5).

Table 4. A) *Homalictus fijiensis* trait covariance model comparison table. B)
Multispecies trait covariation model comparison. Best fitting model with lowest AICc
(sample size corrected AIC) value is in bold. T_e = environmental temperature (°C),
P_{dry}= mean precipitation of the driest month (mm), VPD = vapour pressure deficit
(kPa).

A) *Homalictus fijiensis* trait covariance model comparison					
Testing how trait covariance changes with climatic variables	df	AIC	AICc	ΔAICc	wi
MR x P_{dry}	**10**	**-352.12**	**-350.2**	**0**	**0.93**
Altitude	8	-345.14	-343.9	-6.29	0.04
MR x T_e	10	-343.21	-341.3	-8.91	0.01
MR x T_e x P_{dry}	14	-344.36	-340.5	-9.67	0.007
MR x VPD	10	-342.34	-340.4	-9.78	0.007
B) Multispecies trait covariance model comparison					
Testing how trait covariance changes with climatic variables	df	AIC	AICc	ΔAICc	wi
MR x P_{dry}	**13**	**-507.33**	**-505.4**	**0.00**	**0.94**
MR x T_e x P_{dry}	17	-501.45	-498.1	7.26	0.025
Altitude	12	-498.41	-496.8	8.63	0.013
MR x T_e	13	-498.34	-496.4	8.99	0.011
MR x VPD	13	-497.19	-495.3	10.14	0.006

Figure 5. A) *Homalictus fijiensis* residual ventilation rate (ventilations min[-1]) by residual
metabolic rate (mL min[-1]). B) Multispecies relationship between ventilation rate and
metabolic across precipitation of the driest month. To visually show how the
relationship between metabolic rate and ventilation rate changes depending on
precipitation of the driest month we grouped *H. fijiensis* into precipitation groups, 'dry
= 57 - 124 mm' or 'wet = 125 - 190 mm'. This is for visual representation only, and the
parameter estimates from the model should be used to gain a true understanding of
how the relationship between metabolic rate and ventilation rate changes across the
precipitation of the driest month gradient.
452
453
454
455
456 **Discussion**
457
Adaptive hypotheses that explain variation in physiological traits along climatic
gradients make explicit predictions based on the expected direct effects of abiotic
variables on physiological function. The metabolic cold adaptation (MCA) and hygric

hypotheses, for example, predict that metabolic rates and gas exchange patterns
evolve in response to the effects of temperature and water availability on these two
traits, respectively [6,10]. Whether these hypotheses are supported within and among
species and across environments has been the centre of debate over the last few
decades [6,7,10,19,45]. However, despite metabolic rate and gas exchange patterns
being inherently linked, how they covary across climatic conditions is rarely
considered. Understanding how climate shapes trait relationships is likely to be critical
for improving our understanding of the mechanisms that underpin species distributions
and survival with changing environments [46,47]. We discuss how metabolic rate and
ventilation rate change across climatic gradients independently and then
simultaneously to explore how these traits might evolve together.
472
We found that the MCA hypothesis was not supported in its purest form in Fijian bees.
Although we found a negative relationship between metabolic rate and temperature,
as predicted by the MCA hypothesis, the best model that explained variation in
metabolic rate also included precipitation. We expected that if precipitation did explain
variation in metabolic rate, that metabolic rate would be lowest in dry environments
and highest in wet environments as a mechanism to avoid desiccation (Supplementary
Box 1). However, we found that metabolic rate was negatively correlated with
precipitation (Figure 3B & 3D). This unexpected pattern might be explained by
variation in environmental productivity. Productivity is usually greatest in regions of
high precipitation, and biodiversity is often positively correlated with productivity [48].
In Fiji, precipitation increases with altitude and so does native bee species diversity,
where only one endemic species, *H. fijiensis,* is common in the lowlands and ~ 22
other endemic *Homalictus* species are found in the highlands (> 800 m a.s.l.) [33,38].
Land surface area decreases with altitude in Fiji, leading to an environment that is
small, but productive, and contains high endemic bee diversity. Therefore, potentially
high competition for resources between species has led to an environment where a
lack of energetic resources could be reducing metabolic rates in bees that inhabit high
precipitation high altitude regions. Low metabolic rates might be favoured in
environments with high competition or fewer resources following the 'compensation'
hypothesis which suggests that individuals with low resting metabolic rates will have
higher fitness because they have lower self maintenance costs and can devote more
energy towards growth and reproduction [49]. However, this hypothesis contradicts

findings by [50] which shows that individuals with higher metabolic rates have greater
competitive ability.
497
Variation in ventilation rate within *H. fijiensis* and across Fijian bee species was also
best explained by precipitation of the driest month (Table 3 & Figure 4). Ventilation
rate increased with precipitation, supporting the idea that ventilation rate will be
suppressed in dry environments to avoid desiccating (hygric hypothesis), but selective
pressure on ventilation rate will be relaxed in wet environments. Generally, the hygric
hypothesis is tested by examining the type of breathing pattern that insects employ
across different environments (e.g. continuous, cyclic or discontinuous), but here we
show that species that exhibit discontinuous gas exchange can further regulate the
frequency at which they exchange gas, likely in response to aridity. However, it is
interesting to note that a model that included vapour pressure deficit (drying power of
the air), the other aridity metric, offered the least amount of support in explaining
variation in metabolic rate and ventilation rate within *H. fijiensis* and across bee
species. Because of the rain shadow in Fiji, vapour pressure deficit and precipitation
are only weakly correlated, as can be observed in Figure 1. Thus, bees that inhabit
regions that experience more rainfall might have greater access to water in the
environment, therefore reducing desiccation stress, even if the drying power of the air
remains high.
515
When we examined how metabolic rate and ventilation rate covary across climatic
space, we found that within *H. fijiensis* and across species, those that inhabit the driest
environments have higher ventilation rates per unit CO_2 expelled (steeper ventilation
rate ~ metabolic rate slopes) (Figure 5). Therefore, bees that inhabit drier regions have
high ventilation rates but only exchange a small volume of CO_2 per ventilation, and
that bees in wet environments have low ventilation rates, but each ventilation
exchanges a greater volume of gas. Therefore, we suspect that bees in dry
environments are using shallow ventilations as a mechanism to breathe while limiting
the amount of water that can leave their system and reduce the volume of air that can
diffuse through their tracheole system and desiccate them. Previous studies have
found that some insects fill their tracheoles with water to reduce the surface area of
their respirometry surfaces, potentially as a mechanism to avoid desiccating [51].
Because we re-humidified our respirometry system to avoid desiccating the bees, we

were unable to measure rates of water loss while measuring metabolic rate, which
would have been useful in determining if bees with high frequency small volume
ventilation rates lose less water than bees with slower frequency high volume
ventilations. Future studies should seek to examine how rates of water loss correlate
with metabolic rate-ventilation rate patterns in the future.

As we found that the correlation between metabolic rate and ventilation rate changed
across the precipitation gradient, it suggests that while ventilation rate and metabolic
rate are linked (they are positively correlated across all species) we could be observing
a physiological trait trade-off. While we cannot determine whether the change in
correlation between traits across the precipitation gradient is due to correlational
selection or genetic correlation, the fact that species from different environments have
different trait relationships suggests that there is scope for metabolic rate and
ventilation rate to evolve independently of each other in response to variation in
climate. However, while the two traits might be evolving independently of each other,
they remain correlated, and the extent to which each trait can evolve independently in
response to variation in climate remains an exciting research question for the future.

We believe it is critical to examine organisms as a mosaic of traits to improve our
understanding of organismal responses to changing environments [47,52,53]. Yet
traits are still commonly measured and considered in isolation with very few examples
of how traits coevolve across environmental space, especially in wild populations [but
see 47,54]. Our study shows that the correlation between metabolic rate and
ventilation rate changes across a precipitation gradient, and that these traits could be
evolving independently of each other in response to abiotic factors. Further work is
needed to determine whether the observed phenotypic correlation is underpinned by
the same or different alleles and to examine evolutionary responses of linked traits
that are important for species survival with climate change. Variation in the strength of
trait relationships could have implications for the way in which species evolve with
further anthropogenic climate change [55]. As organisms are composed of multiple
interacting traits and adaptation is a multidimensional problem, we believe it is critical
for research to explore how climate tolerance traits are linked and evolve across
landscapes together as we move further into the Anthropocene.

Acknowledgements

We would like to thank Craig White and Michael Schwarz for helpful discussions and providing comments on an early version of the manuscript, as well as James Dorey for providing bee photographs. We would also like to thank the Navai village for their support. This research was funded by an Endeavour Leadership Postdoctoral Scholarship and a Company of Biologist Travelling Fellowship to CdS. Funding from Monash University (Advancing Women in Science Grant) was awarded to VK, funding from the Australian Research Council was awarded to LAA (DP180103925) and VK (DP200101272). This research was also partially funded by the Australian Department of Foreign Affairs and Trade (via the New Colombo Plan program).

References

1. Diamond SE, Martin RA. 2021 Physiological adaptation to cities as a proxy to forecast global-scale responses to climate change. *J. Exp. Biol.* **224**, jeb229336.

2. Hoffmann AA, Sgro CM. 2011 Climate change and evolutionary adaptation. *Nature* **470**, 479–485.

3. Kellermann V, Overgaard J, Hoffmann AA, Fløjgaard C, Svenning J-C, Loeschcke V. 2012 Upper thermal limits of Drosophila are linked to species distributions and strongly constrained phylogenetically. *Proc. Natl. Acad. Sci.* **109**, 16228–16233.

4. Chown SL, Sørensen JG, Terblanche JS. 2011 Water loss in insects: an environmental change perspective. *J. Insect Physiol.* **57**, 1070–1084.

5. White CR, Alton LA, Frappell PB. 2012 Metabolic cold adaptation in fishes occurs at the level of whole animal, mitochondria and enzyme. *Proc. R. Soc. B Biol. Sci.* **279**, 1740–1747.

6. Addo-Bediako A, Chown SL, Gaston KJ. 2002 Metabolic cold adaptation in insects: a large-scale perspective. *Funct. Ecol.* **16**, 332–338.

7. Alton LA, Condon C, White CR, Angilletta Jr MJ. 2017 Colder environments did not select for a faster metabolism during experimental evolution of Drosophila melanogaster. *Evolution* **71**, 145–152.

8. Chown SL, Gibbs AG, Hetz SK, Klok CJ, Lighton JR, Marais E. 2006 Discontinuous gas exchange in insects: a clarification of hypotheses and approaches. *Physiol. Biochem. Zool.* **79**, 333–343.

9. Terblanche JS, Clusella-Trullas S, Chown SL. 2010 Phenotypic plasticity of gas exchange pattern and water loss in Scarabaeus spretus (Coleoptera: Scarabaeidae): deconstructing

the basis for metabolic rate variation. *J. Exp. Biol.* **213**, 2940–2949.

10. White CR, Blackburn TM, Terblanche JS, Marais E, Gibernau M, Chown SL. 2007
Evolutionary responses of discontinuous gas exchange in insects. *Proc. Natl. Acad. Sci.*
104, 8357–8361.

11. Davis AL, Chown SL, McGeoch MA, Scholtz CH. 2000 A comparative analysis of
metabolic rate in six Scarabaeus species (Coleoptera: Scarabaeidae) from southern
Africa: further caveats when inferring adaptation. *J. Insect Physiol.* **46**, 553–562.

12. Contreras HL, Bradley TJ. 2009 Metabolic rate controls respiratory pattern in insects. *J.*
Exp. Biol. **212**, 424–428.

13. Contreras HL, Bradley TJ. 2010 Transitions in insect respiratory patterns are controlled by
changes in metabolic rate. *J. Insect Physiol.* **56**, 522–528.

14. Schimpf NG, Matthews PG, White CR. 2012 Cockroaches that exchange respiratory
gases discontinuously survive food and water restriction. *Evol. Int. J. Org. Evol.* **66**, 597–
604.

15. Gaston KJ *et al.* 2009 Macrophysiology: a conceptual reunification. *Am. Nat.* **174**, 595–
612.

16. Krogh A. 1916 *The respiratory exchange of animals and man*. Longmans, Green.

17. Addo-Bediako A, Chown SL, Gaston KJ. 2001 Revisiting water loss in insects: a large
scale view. *J. Insect Physiol.* **47**, 1377–1388.

18. White CR, Blackburn TM, Terblanche JS, Marais E, Gibernau M, Chown SL. 2007
Evolutionary responses of discontinuous gas exchange in insects. *Proc. Natl. Acad. Sci.*
104, 8357–8361.

19. Messamah B, Kellermann V, Malte H, Loeschcke V, Overgaard J. 2017 Metabolic cold
adaptation contributes little to the interspecific variation in metabolic rates of 65 species
of Drosophilidae. *J. Insect Physiol.* **98**, 309–316.

20. Mallard F, Nolte V, Tobler R, Kapun M, Schlötterer C. 2018 A simple genetic basis of
adaptation to a novel thermal environment results in complex metabolic rewiring in
Drosophila. *Genome Biol.* **19**, 1–15.

21. Chown SL. 2002 Respiratory water loss in insects. *Comp. Biochem. Physiol. A. Mol.*
Integr. Physiol. **133**, 791–804.

22. Buck J, Keister M, Specht H. 1953 Discontinuous respiration in diapausing Agapema
pupae. In *Anatomical record*, pp. 541–541. Wiley-Liss Div John Wiley & Sons Inc, 605
Third Ave, New York, NY 10158-0012.

23. Hetz SK, Bradley TJ. 2005 Insects breathe discontinuously to avoid oxygen toxicity.
Nature **433**, 516–519.

24. Lighton JR. 1998 Notes from underground: towards ultimate hypotheses of cyclic,
discontinuous gas-exchange in tracheate arthropods. *Am. Zool.* **38**, 483–491.

25. Matthews PG, White CR. 2011 Discontinuous gas exchange in insects: is it all in their
636   heads? *Am. Nat.* **177**, 130–134.

637 26. Rowe TT, Gutbrod MS, Matthews PG. 2022 Discontinuous gas exchange in Madagascan
638   hissing cockroaches is not a consequence of hysteresis around a fixed PCO2 threshold.
639   *J. Exp. Biol.*

640 27. Blows MW. 2007 A tale of two matrices: multivariate approaches in evolutionary biology.
641   *J. Evol. Biol.* **20**, 1–8.

642 28. Calsbeek R, Irschick DJ. 2007 The quick and the dead: correlational selection on
643   morphology, performance, and habitat use in island lizards. *Evol. Int. J. Org. Evol.* **61**,
644   2493–2503.

645 29. Lande R, Arnold SJ. 1983 The measurement of selection on correlated characters.
646   *Evolution* , 1210–1226.

647 30. da Silva CR *et al.* 2021 Climate change and invasive species: a physiological performance
648   comparison of invasive and endemic bees in Fiji. *J. Exp. Biol.* **224**, jeb230326.

649 31. De Frenne P *et al.* 2013 Latitudinal gradients as natural laboratories to infer species'
650   responses to temperature. *J. Ecol.* **101**, 784–795.

651 32. da Silva CR, Groom SV, Stevens MI, Schwarz MP. 2016 Current status of the introduced
652   allodapine bee *Braunsapis puangensis* (Hymenoptera: Apidae) in Fiji. *Austral Entomol.*
653   **55**, 43–48.

654 33. Dorey JB *et al.* 2020 Radiation of tropical island bees and the role of phylogenetic niche
655   conservatism as an important driver of biodiversity. *Proc. R. Soc. B* **287**, 20200045.

656 34. Groom SV, Stevens MI, Schwarz MP. 2013 Diversification of Fijian halictine bees: insights
657   into a recent island radiation. *Mol. Phylogenet. Evol.* **68**, 582–594.

658 35. Fick SE, Hijmans RJ. 2017 WorldClim 2: new 1-km spatial resolution climate surfaces for
659   global land areas. *Int. J. Climatol.* **37**, 4302–4315.

660 36. Hijmans RJ *et al.* 2015 Package 'raster'. *R Package* **734**.

661 37. Monteith J, Unsworth M. 2013 *Principles of environmental physics: plants, animals, and*
662   *the atmosphere*. Academic Press.

663 38. Dorey JB, Groom SV, Velasco-Castrillón A, Stevens MI, Lee MS, Schwarz MP. 2021
664   Holocene population expansion of a tropical bee coincides with early human colonization
665   of Fiji rather than climate change. *Mol. Ecol.* **30**, 4005–4022.

666 39. Pinheiro J, Bates D, DebRoy S, Sarkar D, Heisterkamp S, Van Willigen B, Maintainer R.
667   2017 Package 'nlme'. *Linear Nonlinear Mix. Eff. Models Version* **3**.

668 40. R Development Core Team. 2019 *R: a language and environment for statistical*
669   *computing*. Vienna, Austria.

670 41. Burnham KP, Anderson DR. 2002 A practical information-theoretic approach. *Model Sel.*
671   *Multimodel Inference* **2**.

42. Barton K, Barton MK. 2015 Package 'mumin'. *Version* **1**, 439.

43. Wickham H. 2011 ggplot2. *Wiley Interdiscip. Rev. Comput. Stat.* **3**, 180–185.

44. Wilke CO, Wickham H, Wilke MCO. 2019 Package 'cowplot'. *Streamlined Plot Theme Plot*
Annot. 'ggplot2

45. Clarke A. 2003 Costs and consequences of evolutionary temperature adaptation. *Trends*
Ecol. Evol. **18**, 573–581.

46. Hellmann JJ, Pineda-Krch M. 2007 Constraints and reinforcement on adaptation under
climate change: selection of genetically correlated traits. *Biol. Conserv.* **137**, 599–609.

47. Svensson EI *et al.* 2021 Correlational selection in the age of genomics. *Nat. Ecol. Evol.* **5**,
562–573.

48. Fei S, Jo I, Guo Q, Wardle DA, Fang J, Chen A, Oswalt CM, Brockerhoff EG. 2018 Impacts
of climate on the biodiversity-productivity relationship in natural forests. *Nat. Commun.* **9**,
1–7.

49. Burton T, Killen SS, Armstrong JD, Metcalfe NB. 2011 What causes intraspecific variation
in resting metabolic rate and what are its ecological consequences? *Proc. R. Soc. B Biol.*
Sci. **278**, 3465–3473.

50. Pettersen AK, Hall MD, White CR, Marshall DJ. 2020 Metabolic rate, context-dependent
selection, and the competition-colonization trade-off. *Evol. Lett.* **4**, 333–344.

51. Wigglesworth VB. 1931 The extent of air in the tracheoles of some terrestrial insects. *Proc.*
R. Soc. Lond. Ser. B Contain. Pap. Biol. Character **109**, 354–359.

52. Endler JA. 1995 Multiple-trait coevolution and environmental gradients in guppies. *Trends*
Ecol. Evol. **10**, 22–29.

53. Roff DA, Fairbairn DJ. 2012 A test of the hypothesis that correlational selection generates
genetic correlations. *Evol. Int. J. Org. Evol.* **66**, 2953–2960.

54. Sgro CM, Blows MW. 2004 The genetic covariance among clinal environments after
adaptation to an environmental gradient in Drosophila serrata. *Genetics* **167**, 1281–1291.

55. Chantepie S, Chevin L. 2020 How does the strength of selection influence genetic
correlations? *Evol. Lett.* **4**, 468–478.

56. Angilletta Jr MJ, Huey RB, Frazier MR. 2010 Thermodynamic effects on organismal
performance: is hotter better? *Physiol. Biochem. Zool.* **83**, 197–206.