1 # Mechanisms of circadian clock timing and entrainment revealed by
2 # the structure of Cryptochrome bound to Timeless

4 Changfan Lin[†], Shi Feng[†], Cristina C. DeOliveira and Brian R. Crane
5 [†]Equal Contribution
6
7 Department of Chemistry and Chemical Biology, Cornell University, Ithaca NY 14853

8
9

Circadian rhythms influence many behaviors and diseases. They arise from
oscillations in gene expression caused by repressor proteins that directly inhibit
transcription of their own genes. The circadian clock in flies offers a valuable model
for studying these processes, wherein Timeless (TIM) plays a critical role in
mediating nuclear entry of the transcriptional repressor Period (PER) and the
photoreceptor Cryptochrome (CRY) entrains the clock by triggering TIM
degradation in light. The cryo-EM structure of the CRY:TIM complex reveals how a
light-sensing cryptochrome recognizes its target. CRY engages a continuous core
of N-terminal TIM armadillo (ARM) repeats, resembling how photolyases recognize
damaged DNA, and binds a C-terminal TIM helix reminiscent of the interactions
between light-insensitive CRYs and their partners in mammals. The structure
highlights how the CRY flavin cofactor undergoes substantial conformational
changes that couple to large-scale rearrangements at the molecular interface, and
how a phosphorylated segment in TIM impacts clock period by regulating the
binding of importin-α and the nuclear import of TIM and PER. Moreover, the
structure reveals that the TIM N-terminus inserts into the restructured CRY pocket
to replace the autoinhibitory C-terminal tail released by light, thereby explaining
how the LS-TIM polymorphism adapts flies to different climates.

Circadian rhythms rely on cell-autonomous molecular clocks composed of transcriptional-

translational feedback loops[1,2]. In the paradigm clock of *Drosophila melanogaster* (**Figure 1a**),

the transcriptional repressor proteins Period (PER) and Timeless (TIM) associate and accumulate

in the cytosol before timed transport to the nucleus where they inhibit the transcription factors

Clock (CLK) and Cycle (CYC)[2,3] and thereby downregulate the expression of *per*, *tim* and other

clock-controlled genes[2,3] (**Figure 1a**). A cascade of TIM phosphorylation by the SHAGGY (SGG)

and CK2 kinases gates nuclear entry of PER:TIM to properly time the circadian oscillator[4]. TIM

co-transports PER by binding directly to the nuclear import factor importin-α1[5].

The fly clock is entrained by the flavoprotein Cryptochrome (CRY, **Figure 1a**), which binds to TIM

in the light (**Figure 1b**) and directs it for proteasomal degradation[2,3]. CRYs share a flavin-binding

Photolyase Homology Region (PHR) with the DNA repair enzymes photolyase (PL) but also

contain a CRY C-terminal extension (CCE) that is specific for the function of a given CRY[6-8]. Type

I CRYs from plants and animals sense light to reset the circadian clock; Type II CRYs from

mammalians repress transcription in a light-independent manner by binding the PER proteins and

directly inhibiting CLK[6,7]. In Drosophila, photoreduction of the CRY FAD to the anionic

semiquinone (ASQ) by a tryptophan residue tetrad (W342, W397, W420, W394) triggers release

of a C-terminal tail (CTT) helix (**Figure 1b**)[9-12,13-15]. Photoreduced CRY resets TIM levels by

binding TIM and recruiting the E3 ubiquitin-ligase Jetlag (JET, **Figure 1a**)[16,17]. Recognition of TIM

by CRY is hence the molecular event that sensitizes the clock to light. Although light-sensing

CRYs have also been developed as powerful tools for optogenetic applications[18-20]; there is little

direct structural information on how light-activated CRYs recognize their targets.

To overcome the challenge of isolating sufficient amounts of TIM (1398 residues) for structural

characterization by cryo-electron microscopy (cryo-EM), we developed an insect (Drosophila S2)

cell-based expression and nanobody-based affinity purification workflow for isolation of the

CRY:TIM complex (264 kDa). To produce CRY:TIM, a CRY variant that lacks the 22-residue

CCE[21] and therefore binds TIM constitutively[22,23] was co-transfected and expressed in S2 cells

with S-TIM, the more stable of two TIM isoforms (**Supplementary Methods**). Nanobody-based

affinity resin directed at a 13-residue ALFA tag allowed enrichment and elution of the complex.

Complex homogeneity was further improved, and aggregation avoided by crosslinking with the

lysine-specific reagent disuccinimidyl sulfoxide (DSSO) followed by SEC separation (**Extended**

Data Fig. 1). Eluted samples were applied directly to glow-discharged Quantifoil® grids, vitrified,

and imaged on a 200 keV Talos Arctica microscope equipped with a Gatan Quantum LS energy

filter and K3 direct electron camera. Two out of three 3D classes were refined to yield global 3.48

Å (~98 k particles) and 3.3 Å (~160 k particles) maps of the complex (**Extended Data Fig. 1 and**

Supplementary Table 1). We built an atomic model based on better resolved TIM density in the

3.3 Å map (**Supplementary Movie 1**). Local resolution at the complex interface and within the

body exceeds 2.5 Å.

The core of TIM revealed in the CRY:TIM complex (residues 1-248; 575-783; and 992-1128,

Figure 1b, Extended Data Fig. 2, and Supplementary Movie 1) forms a continuous right-

handed supercoil of armadillo (ARM) 3-helix repeats[24] in which two ARM domains discontinuous

in sequence (ARM$_1$ and ARM$_2$, **Fig. 1c**) associate by swapping their terminal helices (**Extended**

Fig. 2). TIM resembles mammalian TIM (mTIM, see below)[25] but contains an N-terminal extension

that binds to CRY (**Figure 1d**). The CRY PHR comprises an N-terminal α/β domain and a C-

terminal helical domain that harbors the FAD isoalloxazine ring in a central 4-helix bundle of α13-

α16 (**Figure 1b,c, Extended Data Fig. 3, and Supplementary Movie 1**)[26-28]. Four helices (α8-

α11) N-terminal to the helix bundle form the top of the flavin binding pocket and interact with the

FAD phosphate groups and adenosine moiety (**Extended Data Fig. 3**). In this cofactor-capping

region the phosphate binding loop (residues 249-263) following α8 and the protrusion motif

(residues 288-306) following α10, distinguish CRYs from PLs (**Figure 1c**)[26]. The phosphate-

binding loop, the protrusion motif, and the Ser-rich C-terminal lid (residues 420-446) together

comprise the C-terminal coupled motif (CCM, **Figure 1c and Extended Data Fig. 3**) that changes

proteolytic sensitivity when CRY becomes light activated [10].

TIM forms a discontinuous interface with CRY both at the N-terminus and C-terminus (**Figures**

1b-e). The primary CRY:TIM interface buries ~1862 Å^2 (49 residues) and 1807 Å^2 (52 residues)

of solvent accessible surface area on TIM and CRY, respectively. TIM primarily binds to CRY

through the first three helices of the N-terminal ARM domain (**Figure 1d,e**). ARM$_1$-α1 binds across

CRY α17 and inserts directly into the flavin binding pocket replacing the CRY CTT (**Figures 1d,**

2a, and Extended Data Fig. 4a). The contact is dominated by TIM ARM$_1$-α1-3, ARM$_1$-α7 and the

ARM$_1$-α9-10 loop which mesh against elements of the CRY CCM (**Figure 1e and Extended Data**

Figs. 2 and 4). At the center of the interface, Met1 and Asp2 extend out from α1 with Met1

packing against the Trp4 indole and embedding among Tyr250 and Leu251 of the CRY phosphate

binding loop, Gly310 on α11, and Tyr158-Gln159 on the α5-α6 loop that extends down from the

α/β domain (**Extended Data Figs. 3 and 5**). TIM Leu4 and Leu5 insert deep into the pocket

adjacent to CRY Trp314 and the FAD adenine moiety. ARM_1-α2 contacts the C-terminal lid and

ARM_1-α7 contacts the protrusion motif, wherein the α10 helix lengthens to project the tip region

toward the Met-rich ARM_1-α9-α10 loop at the interface periphery (**Extended Data Figs. 4c,d**).

The recognition of TIM ARM_1-a1 by CRY (**Figure 1d**) explains an evolutionary adaptation in light-

sensing by flies. A *Drosophila melanogaster tim* allele (*ls-tim*) causes differential expression of

two isoforms; S-TIM, which begins at the N-terminal Met residue found in the structure, and L-

TIM, which extends by an additional 23 N-terminal residues owing to introduction of an alternative

translational start site (**Figure 1b**)[29,30]. The frequency of *ls-tim* correlates with geographical

latitude and may confer a beneficial preference for more robust rhythms and a diapause state of

dormancy in climates of more variable light and cooler temperatures. L-TIM reduces interaction

with CRY by yeast two-hybrid, has greater stability in fly heads, and diminishes clock responses

to light [17,31]. The Met1 amino-terminus projects into a solvent channel at the interface of the two

proteins that could therefore accommodate the L-TIM extension (**Figure 1d and Extended Data**

Fig. 6a). However, the recognition of S-TIM by CRY through the N-terminal Met residue will be

altered by the L-TIM extension (**Figure 1d)**, thereby explaining the capacity of *ls-tim* to alter fly

light sensitivity.

In addition to the primary interaction between TIM ARM_1 and CRY, strong electron density for a

TIM helical segment is evident among the C-terminal lid, the C-terminal end of the last CRY PHR

helix (α20) and the α5-α6 loop that projects down from the α/β domain (**Figure 1e and Extended**

Data Fig. 3). This TIM helix buries 496 Å^2 of surface area and binds at the position occupied by

the residues that link α20 to the CTT in the dark-state structure, thereby supporting results from

interaction studies[23] and providing further evidence that the CTT undocks from the flavin pocket

upon light activation[10,12,14]. There are no obvious connections of this density to the ordered TIM

ARM core and thus its sequence assignment is provisional; however, after testing many helical

models and sequence registers, TIM 1163-1174 provided the most reasonable fit to the density

that also adhered to the stereochemical constraints of the site (**Extended Data Fig. 7a**). To further

validate this assignment, we performed the SWFTI assay (**Extended Data Fig. 8**) with TIM

truncations and the CRY variant H377L that displays tight binding to TIM[13]. CRY bound TIM 1-

1132 and 571-1398 similarly, but with reduced affinity compared to full-length S-TIM. TIM 1133-

1398, which contains the proposed peripheral helix, alone bound to CRY-H377L (**Extended Data**

Fig. 8). A known mutation of human TIM that removes the analogous C-terminal region (R1081X)

causes familial advanced sleep phase by reducing interactions between mammalian TIM and

mCRY2[32]. Thus, although the roles of mCRY and CRY are not the same in animals and insects[2],

they may produce similar interactions with the otherwise disordered C-termini of their respective

TIM proteins. These interactions may include competition with PER because TIM 1163-1174

associates with CRY in a region that coincides with where mCRY binds the mammalian Period

proteins (**Extended Data Fig. 9a**)[33-35]

TIM and mTIM have a similar topology in their ARM repeats (**Extended Data Fig. 9b**) and also

share a C-terminal poly-(ADP-ribose) polymerase – 1 (PARP-1) binding (PAB) domain[36], which

includes a conserved stretch of the Timeless C region (residues 1051-1096[32]) that forms a helix

and loop structure (**Extended Data Fig. 9c**). However, the TIM PAB is unlikely to bind to PARP-

1 because it does not compose the required interface displayed by mTIM, despite high structural

similarity of the Drosophila and mammalian PARP-1 proteins. In fact, mTIM is more closely related

to the Drosophila Timeout protein (TIM-2) than to TIM itself[25,32]. Although mTIM and TIM-2 play

some role in circadian rhythms neither are central constituents of the core mammalian oscillator[2].

In contrast, mTIM and its homologs (TOF1 in yeast, SWI1 in and TIM-1 in *C. elegans*) stabilize

replication forks during DNA replication, pair sister chromatids and regulate S-phase

checkpoints[37]. Tof-1 participates in the large replisome complex in which it binds directly to dsDNA

(**Extended Data Fig. 9d**)[38]. Thus, details of the respective structures indicate that TIM is likely

functionally distinct from mTIM, Tof-1, and TIM-2, in keeping with its role in the stabilization and

co-transport of PER.

Pronounced changes in the CRY cofactor binding pocket (**Figure 2a**) involve repositioning of the

adenosine and ribose moieties, a large switch in the conformation of the diphosphate groups,

refolding of the C-terminal lid to interact with both the TIM N-terminus and ARM_1-α2 and collapse

of the phosphate binding loop into the flavin pocket to allow Asn253, Gln254 and Gln311 to

hydrogen bond with the FAD phosphates (**Extended Data Fig. 5a**). Tyr250 and Leu251 interact

with the FAD pocket and make space for the top of TIM ARM_1-α3 to insert Phe58 behind the new

conformation of the phosphate-binding loop (**Figure 2b and Extended Data Fig. 5a**). In addition,

α8 tilts 20° towards the FAD diphosphates as Arg237 replaces the FAD Mg^{2+} counter ion, for

which no electron density is observed (**Figure 2c and Extended Data Fig. 5b**). The FAD adenine

ring slides 1.5 Å relative to the unbound structure, an adjustment compensated by a shift of

Phe318 and harboring helix α11 (**Extended Data Fig. 5b**). The 262-to-264 loop also changes

structure to accommodate the new diphosphate conformation and Gln311 swivels away from TIM

Leu4 (**Extended Data Fig. 5a,c**). A CTT-like sequence on TIM (617-626) that in isolation binds

light-state CRY [10] locates in the ARM core remote from the interface (**Figure 2a**).

The CRY:TIM complex was purified under light, although the cofactor redox state could not be

verified on the cryo-EM grids. Nevertheless, TIM likely stabilizes features of the light-state

structure. For example, the site of the light-insensitive *cryb* mutation[39], an Asp410-to-Arg381 salt

bridge proposed from MD simulations to restructure during light activation[40], appears to break in

response to the new conformation of the ribose backbone (**Figure 2d**), in keeping with light driving

undocking of the CTT and subsequent rearrangement of the CCM. Two conserved histidine

residues within the flavin pocket, His377 and His378, participate in the PL catalytic mechanism

and are implicated in CTT release and TIM binding[13,15,41]. In the unbound state His378 ND1

hydrogen bonds with O2' of the FAD ribose, but in the TIM complex, His378 tilts away from the

ribose hydroxyl to hydrogen bond to the TIM Leu5 carbonyl with NE2 (**Figures 2b,c and**

Extended Data Fig. 5b). Substitutions at His378 still allow CTT release, but on/off switching is

less robust[13,41]. Protonation of His378 NE2 is likely required for it to interact with the TIM backbone

carbonyl and His378 protonation may be responsive to the flavin redox state. The H377L

substitution substantially increases the binding affinity to TIM in both light and dark[13]. Surprisingly,

His377 does not contact TIM directly but rather anchors the restructured phosphate binding loop

by contacting several residues and hydrogen bonding to the Asn253 carbonyl (**Figure 2c**).

The interaction of CRY with TIM corresponds closely to how 6-4 photolyase recognizes the 6-4

crosslinked pyrimidine lesion in damaged DNA (**Figure 3a**). Indeed, TIM Leu4 and Leu5

superimpose with the crosslinked base and the TIM ARM$_1$-a1 polypeptide closely follows the path

of 5'-to-3' strand-separated DNA (**Figure 3a**)[42]. Key residues in the pocket also share identity and

similar positions, with the notable exception of PL Tyr306, which as a Phe in CRY contacts the

Leu5 side-chain. TIM and FAD also combine in CRY to mimic interactions of the FBXL3 E3 ligase

with apo-mCRYs for their ubiquitin mediated degradation (**Figure 3b**)[43]. Despite these similarities

at the primary cofactor pocket, the phosphate binding loop, protrusion motif and C-terminal lid

differ in both sequence and conformation among PLs, mCRYs and CRY. CRY also contains a

secondary pocket between its two domains that in PLs binds antenna cofactors and in mCRYs

mediates contacts with other clock components[6,44]. No cofactors or other constituents are found

within the secondary pocket of the CRY:TIM structure; however, it composes a deep cleft typical

of a recognition motif (**Extended Data Fig. 6b**).

The TIM structure provides insight into the gating of TIM:PER nuclear entry by the interaction
between TIM and importin-α1 (IMPα1). TIM is homologous to the β-catenins, their related family
members (HMP-2, p120, plakoglobin, plakophilin, and SYS-1) as well as the importin-α/β
karyopherins involved in nuclear transport[5,24,45]. These ARM proteins conserve a positively
charged groove on the concave side of their super-helical crescents where they bind extended
peptide chains, such as nuclear localization signals (**Figure 4a,b and Extended Data Fig. 10**)[46,47].
Two mutations known to block nuclear entry by reducing interactions between TIM and importin-
α1 (Thr90Ala and Pro92Leu; S-TIM numbering)[5] localize within the TIM groove (**Figure 4b**) but
do not interact with the TIM nuclear localization sequence, which rather extends in a helix along
the convex side of the protein (**Figure 4c**). An extended polypeptide with α-helical character
binds into the TIM groove adjacent to catenin-conserved Arg (84, 165) and Asn (88, 166) residues
(**Extended Data Fig. 10**), and Thr90/Pro92 (**Figure 4b**). Crosslinking mass spectrometry
revealed that TIM Lys313 and Lys328 in the disordered TIM domain (249-545) react with Lys
residues in the groove and toward the bottom of the crescent (**Figure 4c and Extended Data Fig.**
10), whereas Lys residues in other disordered regions (400-544; 784-991; 1129-1398) show little
crosslinking to TIM or CRY. After extensive modeling, the sequence Ser283-Thr303 was
provisionally assigned as the most reasonable fit to the polypeptide density (**Extended Data Fig.**
7b). This assignment places Asp287 and Asn291 adjacent to the conserved Asn/Arg residues
and occludes the Thr90 and Pro92 residues involved in importin-α1 binding (**Figure 4b**).
207
The Ser283-Thr303 polypeptide contains Ser286 and Ser290, which belong to the TIM-ST stretch
of SGG and CK2 phosphorylation sites that regulate nuclear import (TIM-ST: SGG: S268, T278;
CK2: T282, S286, S290; **Figures 1b and 4b**)[4]. Phosphorylation of these residues in pacemaker
neurons (s-LVs) increases nuclear entry rates[4]. The structure suggests that phosphorylation of
the TIM-ST displaces this segment from the groove to allow engagement of importin-α1 (**Figure**
4d). Thus, key delays that tune the clock period likely depend on the interactions of a TIM
autoinhibitory segment that alter upon its timed phosphorylation by circadian kinases.
215
The CRY:TIM structure reveals how a light-sensing cryptochrome engages its target. It also
provides insight into how evolutionary relationships among the component proteins comprise
regulatory circuits. Swapping the auxiliary CTT of CRY for the variable TIM N-terminus not only
sets the circadian oscillator but also tunes its light sensitivity. In this interaction, the ability of TIM
to serve as a peptide mimetic for a PL DNA substrate underscores how conserved structural
features of PLs are also utilized by CRYs. In gating its partnership with CRY, TIM shares an

autoinhibitory mechanism that resembles that which regulates importin-α[47-49]. The striking changes in FAD geometry that couple to CRY conformational rearrangement and TIM binding were unexpected. These new insights raise questions as to whether other flavoprotein light-sensors share similar mechanisms to convey light or redox signals from their cofactors.

References

Takahashi, J. S. Transcriptional architecture of the mammalian circadian clock. *Nature Reviews Genetics* **18**, 164-179 (2017).

Crane, B. R. & Young, M. W. interactive features of proteins composing eukaryotic circadian clocks. *Ann. Rev. Biochem.* **83** (2014).

Foley, L. E. & Emery, P. Drosophila Cryptochrome: Variations in blue. *Journal of Biological Rhythms* **35**, 16-27 (2020).

Top, D., Harms, E., Syed, S., Adams, E. L. & Saez, L. GSK-3 and CK2 kinases converge on Timeless to regulate the master clock. *Cell Reports* **16**, 357-367 (2016).

Jang, A. R., Moravcevic, K., Saez, L., Young, M. W. & Sehgal, A. Drosophila TIM binds Importin α1, and acts as an adapter to transport PER to the nucleus. *Plos Genetics* **11** (2015).

Chaves, I. *et al.* The cryptochromes: blue light photoreceptors in plants and animals. *Annual review of plant biology* **62**, 335-364 (2011).

Ozturk, N. Phylogenetic and functional classification of the photolyase/cryptochrome family. *Photochemistry and Photobiology* **93**, 104-111 (2017).

Parico, G. C. G. *et al.* The human CRY1 tail controls circadian timing by regulating its association with CLOCK:BMAL1. *Proc. Natl. Acad. Sci. US.* **117**, 27971-27979 (2020).

Berndt, A. *et al.* A novel photoreaction mechanism for the circadian blue light photoreceptor Drosophila cryptochrome. *J Biol Chem* **282**, 13011-13021 (2007).

Vaidya, A. T. *et al.* Flavin reduction activates Drosophila cryptochrome. *Proc Natl Acad Sci USA* **110**, 20455-20460 (2013).

Ozturk, N., Selby, C.P., Annayev, Y., Zhong, D., Sancar, A. Reaction Mechanism of Drosopila cryptochrome. *Proc Natl Acad Sci USA* **108**, 516-521 (2011).

Lin, C. F., Top, D., Manahan, C. C., Young, M. W. & Crane, B. R. Circadian clock activity of cryptochrome relies on tryptophan-mediated photoreduction. *Proc. Natl. Acad. Sci. USA* **115**, 3822-3827 (2018).

Lin, C., Schneps, C. M., Chandrasekaran, S., Ganguly, A. & Crane, B. R. Mechanistic insight into light-dependent recognition of Timeless by Drosophila Cryptochrome. *Structure* doi: 10.1016/j.str.2022.03.010 (2022).

14 Chandrasekaran, S. *et al.* Tuning flavin environment to detect and control light-induced
conformational switching in Drosophila cryptochrome. *Communications biology* **4**, 1-12
(2021).

15 Ganguly, A. *et al.* Changes in active site histidine hydrogen bonding trigger
cryptochrome activation. *Proc. Natl. Acad. Sci. U. S. A.* **113**, 10073-10078 (2016).

16 Koh, K., Zheng, X. Z. & Sehgal, A. JETLAG resets the Drosophila circadian clock by
promoting light-induced degradation of TIMELESS. *Science* **312**, 1809-1812 (2006).

17 Peschel, N., Chen, K. F., Szabo, G. & Stanewsky, R. Light-dependent interactions
between the Drosophila circadian clock factors cryptochrome, jetlag, and timeless.
Current Biology **19**, 241-247 (2009).

18 Zhou, X. X. *et al.* A single-chain photoswitchable CRISPR-Cas9 architecture for light-
inducible gene editing and transcription. *ACS Chem Biol* **13**, 443-448 (2018).

19 Yu, D. *et al.* Optogenetic activation of intracellular antibodies for direct modulation of
endogenous proteins. *Nat Methods* **16**, 1095-1100 (2019).

20 Gil, A. A. *et al.* Optogenetic control of protein binding using light-switchable nanobodies.
Nat Commun **11**, 4044 (2020).

21 Busza, A., Emery-Le, M., Rosbash, M. & Emery, P. Roles of the two Drosophila
Cryptochrome structural domains in circadian photoreception. *Science* **304**, 1503-1506
(2004).

22 Dissel, S. *et al.* A constitutively active cryptochrome in Drosophila melanogaster. *Nat.
Neurosci.* **7**, 834-840 (2004).

23 Hemsley, M. J. *et al.* Linear motifs in the C-terminus of D-melanogaster cryptochrome.
Biochemical and Biophysical Research Communications **355**, 531-537 (2007).

24 Vodovar, N., Clayton, J. D., Costa, R., Odell, M. & Kyriacou, C. P. The Drosophila clock
protein Timeless is a member of the Arm/HEAT family. *Current Biology* **12**, R610-R611
(2002).

25 Holzer, S. *et al.* Crystal structure of the N-terminal domain of human Timeless and its
interaction with Tipin. *Nucleic Acids Research* **45**, 5555-5563 (2017).

26 Zoltowski, B. D. *et al.* Structure of full-length Drosophila cryptochrome. *Nature* **480**, 396-
399 (2011).

27 Levy, C. *et al.* Updated structure of Drosophila cryptochrome. *Nature* **495**, E3-4 (2013).

28 Czarna, A. *et al.* Structures of Drosophila cryptochrome and mouse cryptochrome1
provide insight into circadian function. *Cell* **153**, 1394-1405 (2013).

29 Tauber, E. *et al.* Natural selection favors a newly derived timeless allele in Drosophila
melanogaster. *Science* **316**, 1895-1898 (2007).

30   Deppisch, P. *et al.* Adaptation of Drosophila melanogaster to long photoperiods of high-
latitude summers is facilitated by the ls-timeless allele. *J. Biol. Rhythms* **37**, 185-201
(2022).

31   Sandrelli, F. *et al.* A molecular basis for natural selection at the timeless locus in
Drosophila melanogaster. *Science* **316**, 1898-1900 (2007).

32   Kurien, P. *et al.* TIMELESS mutation alters phase responsiveness and causes advanced
sleep phase. *Proc. Natl. Acad. Sci. USA* **116**, 12045-12053 (2019).

33   Schmalen, I. *et al.* Interaction of circadian clock proteins CRY1 and PER2 Is modulated
by zinc binding and disulfide bond formation. *Cell* **157**, 1203-1215 (2014).

34   Fribourgh, J. L. *et al.* Dynamics at the serine loop underlie differential affinity of
cryptochromes for CLOCK:BMAL1 to control circadian timing. *Elife* **9** (2020).

35   Nangle, S. N. *et al.* Molecular assembly of the period-cryptochrome circadian
transcriptional repressor complex. *Elife* **3** (2014).

36   Xie, S. *et al.* Timeless interacts with PARP-1 to promote homologous recombination
repair. *Molecular Cell* **60**, 163-176 (2015).

37   Engelen, E. *et al.* Mammalian Timeless Is Involved in period determination and DNA
damage-dependent phase advancing of the circadian clock. *PLoS One* **8** (2013).

38   Baretic, D. *et al.* Cryo-EM Structure of the fork protection complex bound to CMG at a
replication fork. *Molecular Cell* **78**, 926-+ (2020).

39   Stanewsky, R. *et al.* The *cry(b)* mutation identifies cryptochrome as a circadian
photoreceptor in Drosophila. *Cell* **95**, 681-692 (1998).

40   Wang, Y. J., Veglia, G., Zhong, D. P. & Gao, J. L. Activation mechanism of Drosophila
cryptochrome through an allosteric switch. *Science Advances* **7** (2021).

41   Berntsson, O. *et al.* Photoactivation of Drosophila melanogaster cryptochrome through
sequential conformational transitions. *Science Advances* **5** (2019).

42   Glas, A. F., Schneider, S., Maul, M. J., Hennecke, U. & Carell, T. Crystal structure of the
T(6-4)C lesion in complex with a (6-4) DNA photolyase and repair of UV-induced (6-4)
and dewar photolesions. *Chemistry-a European Journal* **15**, 10387-10396 (2009).

43   Nangle, S., Xing, W. M. & Zheng, N. Crystal structure of mammalian cryptochrome in
complex with a small molecule competitor of its ubiquitin ligase. *Cell Research* **23**, 1417-
1419 (2013).

44   Rosensweig, C. *et al.* An evolutionary hotspot defines functional differences between
cryptochromes. *Nature Communications* **9** (2018).

45   Tewari, R., Bailes, E., Bunting, K. A. & Coates, J. C. Armadillo-repeat protein functions:
questions for little creatures. *Trends in Cell Biology* **20**, 470-481 (2010).

46   Choi, H. J. *et al.* A conserved phosphorylation switch controls the interaction between
329   cadherin and β-catenin In Vitro and In Vivo. *Developmental Cell* **33**, 82-93 (2015).

330   47   Goldfarb, D. S., Corbett, A. H., Mason, D. A., Harreman, M. T. & Adam, S. A. Importin-α:
331   a multipurpose nuclear-transport receptor. *Trends in Cell Biology* **14**, 505-514 (2004).

332   48   Catimel, B. *et al.* Biophysical characterization of interactions involving importin-alpha
333   during nuclear import. *Journal of Biological Chemistry* **276**, 34189-34198 (2001).

334   49   Harreman, M. T., Hodel, M. R., Fanara, P., Hodel, A. E. & Corbett, A. H. The auto-
335   inhibitory function of importin-α is essential in Vivo. *Journal of Biological Chemistry* **278**,
336   5854-5863 (2003).

337   50   Holm, L. & Laakso, L. M. Dali server update. *Nucleic Acids Research* **44**, W351-W355
338   (2016).

339
340

Figure 1. Cryo-EM structure of the CRY:TIM complex. (**a**) Drosophila Cryptochrome (CRY)

entrains the circadian clock by initiating degradation of Timeless (TIM) in light. Jetlag (JET),

Period (PER), Doubletime (DB), Cycle (CYC), Clock (CLK) and Importin-α1 (IMPα1) compose the

core circadian oscillator. (**b**) CRY:TIM docked into the electron density map. Dark-state CRY

shown at the bottom (grey, PDB ID: 4GU5). (**c**) Domain arrangements and key regions of CRY

and TIM. Features of TIM: CRY-binding domains (CRY-BD1, 1-234, CRY-BD2, 1133-1398),

serine/threonine-rich region (ST, 270-290) with key phosphorylation sites starred (S274, T278,

T282, S286 and S290), ARM1 (1-248), ARM2 (578-866), nuclear localization sequence (NLS,

545-578), Parp-1 binding domain (PAB, 986-1142; TIM-C, 1051-1096), 23 residue L-TIM

extension (L, (23aa)), disordered regions are dashed. Features of CRY: Phosphate binding loop

(PL, 249-263), Protrusion motif (PM, 288-306), C-terminal lid (CL, 420-446), C-terminal tail (CTT,

519-542, absent in CRYΔ). (**d**) N-terminus of TIM inserts into the primary flavin binding pocket.

2.4 Å resolution local electron density is shown with notable hydrogen bonds (yellow dashes) and

hydrophobic contacts (green circle). (**e**) Interface between CRY and TIM (blue shaded area), TIM

groove peptide (in navy blue) and peripheral helix (fitted with 1163-1174 putatively, blue). Inset

shows the interactions between TIM 1163-1174 and CRY superimposed with electron density.

Figure 2. Conformational changes of FAD to the CRY C-terminal coupled motif upon

binding TIM. (**a**) CRY:TIM (CRY, yellow; TIM, blue) in superposition with dark-state CRY. The

N-terminus of TIM replaces the CRY CTT (red), as depicted in the schematic below. Structural

elements designated as in **Figure 1**. (**b**) The phosphate binding loop (dark pink) collapses into

the flavin pocket and N253 and E254 hydrogen bond (yellow dashes) with the FAD diphosphate

conformation (red, orange) that differs substantially from that of the dark state (white bonds). (**c**)

R237 replaces the Mg^{2+} counter ion and H377 anchors the new position of the loop. The D410-

R381 salt bridge (top) appears to break in response to the new conformation of the ribose

backbone (bottom). Reduced side-chain density of R381 suggests increased flexibility in the TIM

complex.

Figure 3. Conservation of recognition modes by CRY family members. (**a**) The interaction of

CRY with TIM corresponds closely to how 6-4 photolyase recognizes the 6-4 pyrimidine lesion in

damaged DNA (PDB ID:3CVU). TIM Leu4 and Leu5 superimpose on the crosslinked bases and

the TIM ARM1- α1 polypeptide closely follows the path of 5'-to-3' strand-separated DNA. (**b**) The

TIM and FAD contacts of CRY mimic the interaction of apo-mCRY with the C-terminus of the

FBXL3 E3 ligase (PDB ID: 4I6J).

Figure 4. A structural basis for regulation of TIM nuclear entry. (**a**) Electron density (dark

blue) for an extended polypeptide resides within a groove wherein members of the β-catenin and

α-importin proteins bind their target sequences. (**b**) The TIM groove peptide density shown with

the sequence assignment S283-T303. The peptide interacts with β-catenin-conserved Asn and

Arg residues and buries residues important for importin-α1 binding (T90 and P92). (**c**) DSSO-

mediated lysine crosslinking patterns of disordered residues K313 and K238 shown on the TIM

structure. K313 and K328 produce a pattern of crosslinking that localizes them to a position

consistent with placement of the S283-T303 groove peptide. The TIM NLS lies in a helix adjacent

to the groove but does not occlude T90 and P92. (**d**) Schematic of proposed nuclear entry

mechanism of TIM. Phosphorylation of the TIM-ST region by SGG and CK2 releases the ST

region from TIM groove to expose TIM T90 and P92. Importin-α1 recognizes TIM by binding

adjacent to Thr90 and Pro92, while also binding the TIM NLS within a structurally analogous ARM

groove of its own.

Methods

Cloning

All DNA constructs (shown above) were built by a modified Gibson assembly method and Q5

polymerase-based mutagenesis with home-made reaction mixture, as reported previously[51].

Transient co-transfection of S2 cells

S2 cells were maintained at 27 °C in 90% Schneider's Drosophila Medium (Cat# 21720024,

ThermoFisher) suppled with 10% heat-inactivated Fetal Bovine Serum (FBS, Cat# F4135, sigma),

as reported previously[51]. pAC5.1 vectors of CLIP-CRYΔ and TIM-SNAP-HA-ALFA were delivered

into S2 cells using the TransIT-Insect Transfection Reagent (Cat# MIR6100, Mirus) following a

modified version of the manufacturer's instruction. Briefly, about twenty 100 mm culture dishes

(Cat# 353003, Corning) with 1000×10^4 cells in 10 ml fresh growth medium were prepared on day

1. About 20 hours later, 10 µg DNA in total (CRYΔ: TIM mass ratio 1:1) were transfected by the

TransIT-Insect reagent in each 100 mm dish. 72 hours after transfection, cells were harvested by

centrifugation with 500 × g for 5 minutes at room temperature, washed once with chilled PBS and

then stored at -80 °C until lysis.

Nanobody-based affinity purification of the CRY:TIM complex

Two cycles of freeze-thaw were used to lyse the S2 cells as follows: S2 cell pellets from 200 ml

of culture were resuspended with 16 ml lysis buffer (50 mM Tris, pH 8, NaCl 150 mM, 20% glycerol,

1x protease inhibitor (Cat# A32965, ThermoFisher)) and flash frozen in liquid nitrogen. The

sample was then quickly thawed as 800 ul aliquots in a room temperature water bath. The lysate

aliquot was resupended (without vortexing) and the freeze-thaw cycle repeated. 100 U/ml

benzonase (sc-202391, Santa Cruz Biotechnology) was added to the sample and incubated at

4 °C for 45 min to remove DNA/RNA contamination. The cell lysate was spun at 20000 x g for 20

minutes at 4 °C and further filtered with a 0.45 micron PVDF syringe filter (Cat# SLHV033NB,

Millipore Sigma). 160 µL settled ALFA resin (pre-washed with lysis buffer A) was incubated with

the cell lysate at 4 °C for 1 h on an end-to-end rotator (Cat# 88881001, ThermoFisher scientific).

The slurry was transferred to a 15 ml falcon tube, centrifuged at 1000 x g for 1 min and wash the

resin with 7 mL wash buffer (HEPES 50 mM, pH 7.9, NaCl 150 mM, 10% glycerol). This process

was repeated 3 times. Then the resin was resuspended in 1 mL wash buffer and split into two

and applied to separate spin columns (Cat# 6572, BioVision), with 80 µL resin applied to each

column. The spin columns were washed 4 times with 500 ul wash buffer. 500 µL elution buffer

(wash buffer with 200 uM ALFA peptide) was added to 80 µL resin in each spin column and

incubated in the cold room for 20 min on an end-to-end rotator. The columns were spun at 1000

x g for 1 min to collect the elution. The elution was repeated 3 times and all of the elutions were

then combined (3 mL in total).

DSSO crosslinking coupled with SEC

The elution was concentrated to about 70 to 80 μL (while maintaining the final protein

concentration at less than 2 mg/mL to ensure efficient crosslinking) with a 30 kDa vivaspin

concentrator (Cat# 28932235, Cytiva). 50 mM stock solution of DSSO was prepared in dimethyl

sulfoxide (DMSO) following the manufacture's instructions. DSSO was added to the elution

sample to a final concentration of 1 mM and incubated at room temperature for 45 minutes,

followed by quenching for 5 min at room temperature with addition of 1 M Tris buffer (pH 8) to

produce a final concentration of 20 mM Tris. The crosslinked complex was concentrated by 50

kDa vivaspin (Cat# 28932236, Cytiva) and purified by a Superose SEC column 6 10/300 (Cat#

29091596, Cytiva) or 3.2/300 (Cat# 29091598, Cytiva). The protein complex was eluted in SEC

buffer (20 mM HEPES, pH 7.9, 150 mM NaCl) and concentrated to ~40 μL by a 50 kDa cutoff

vivaspin concentrator.

Cross-Linking Mass Spectrometry (XL-MS)

SEC-purified crosslinked product was mixed with 4x Laemmli Sample Buffer (Cat# 1610747,

BioRad) and 2.5% 2-mercaptoethanol (BME, Cat# 1610710, BioRad) and then boiled for 6 min.

The sample was then loaded on an SDS-PAGE gel and stained with QC Colloidal Coomassie

Stain (Cat #1610803, BioRad). The band of the complex (~ 20 ug protein sample) was cut out

and prepared for MS-MS analysis of crosslinked peptides as previously reported[52] and described

below.

In-gel digestion of SDS gel bands

The protein band (~ 20 μg protein) was sliced into ~1 mm cubes. The excised gel pieces were

washed consecutively with 400 μL deionized water followed by 50 mM ammonium bicarbonate,

50% acetonitrile and finally 100% acetonitrile. The dehydrated gel pieces were reduced with 200

μL of 10 mM DTT in 100 mM ammonium bicarbonate for 45 min at 56 °C, followed by alkylation

with 200 μL of 55 mM iodoacetamide in 100 mM ammonium bicarbonate at room temperature in

the dark for 45 min. Wash steps were repeated as described above. The gel was then dried and

rehydrated with 100 μL LysC at 10 ng/μL in 50 mM ammonium bicarbonate containing 10%

acetonitrile (ACN) and incubated at 37 °C for 3 h. After the 3 h incubation, 30 μL Trypsin at 43

ug/uL in 50 mM ammonium bicarbonate was added and the sample was continued to incubate at

37 °C for 18 h. The digested peptides were extracted from the gel twice with 200 μL of 50%

acetonitrile containing 5% formic acid and once with 200 μL of 75% acetonitrile containing 5%

formic acid. Extractions from each sample were pooled together and then divided equally in half.

One half was then filtered with 0.22 μm spin filter (Costar Spin-X from Corning), dried to dryness

in the speed vacuum and analyzed as LysC/Trypsin digests. The second half was dried to

dryness in the speed vacuum then reconstituted with 30 µL GluC at 33 ng/µL in 50mM ammonium

bicarbonate (pH 7.9), incubated at 37 °C for 18 h. After incubation, formic acid was added to stop

enzyme reaction. The digests were filtered with 0.22 µm spin filter, dried to dryness in the speed

vacuum and analyzed as LysC/Trypsin/GluC digests. Each sample was reconstituted in 0.5%

formic acid prior to LC- MS/MS analysis.

Crosslinks identification by nanoLC-MS/MS

The digested product was characterized by NanoLC-MS/MS analysis at the Cornell Biotechnology

Resource Center. The analysis was carried out using an Orbitrap Fusion™ Tribrid™ (Thermo-

Fisher Scientific, San Jose, CA) mass spectrometer equipped with a nanospray Flex Ion Source,

and coupled with a Dionex UltiMate 3000 RSLCnano system (Thermo, Sunnyvale, CA). Each

sample was loaded onto a nano Viper PepMap C18 trapping column (5 µm, 100 µm × 20 mm,

100 Å, Thermo Fisher Scientific) at 20 μL/min flow rate for rapid sample loading. After 3 minutes,

the valve switched to allow peptides to be separated on an Acclaim PepMap C18 nano column

(2 µm, 75 µm x 25 cm, Thermo Fisher Scientific) at 35 °C in a 90 min gradient of 5% to 40% buffer

B (98% ACN with 0.1% formic acid) at 300 nL/min. The Orbitrap Fusion was operated in positive

ion mode with nano spray voltage set at 1.7 kV and source temperature at 275 °C. External

calibrations for Fourier transform, ion-trap and quadrupole mass analyzers were performed prior

to the analysis. Samples were analyzed using the CID-MS^2-MS^3 workflow, in which MS scan

range was set to 375–1500 m/z and the resolution was set to 60,000. Precursor ions with charge

states 3-8 were selected for CID MS^2 acquisitions in Orbitrap analyzer with a resolution of 30,000

and an AGC target of 5 ×10^4. The precursor isolation width was 1.6 m/z and the maximum

injection time was 100 ms. The CID MS^2 normalized collision energy was set to 25%. Targeted

mass-difference triggered HCD-MS^3 spectra were acquired in the ion trap with AGC target of 1

×10^4. When a unique mass difference (Δ=31.9721 Da) was observed in CID-MS^2 spectra for ions

with a single charge and multiple charges, the HCD collision energy at 38% and 28% was used

respectively, for the subsequent MS^3 scans. All data were acquired under Xcalibur 4.3 operation

software and Orbitrap Fusion Tune application v3.4 (Thermo-Fisher Scientific).

MS Data analysis

All MS, MS^2 and MS^3 raw spectra from each sample were searched using Proteome Discoverer

2.4 (Thermo-Fisher Scientific, San Jose, CA) with XlinkX v2.0 algorithm for identification of

crosslinked peptides. The search parameters were as follow: four missed cleavage for either

double digestion or triple digestion with fixed carbamidomethyl modification of cysteine, variable
modifications of methionine oxidation, asparagine and glutamine deamidation, protein N-terminal
acetylation. The peptide mass tolerance was 10 ppm, and MS^2 and MS^3 fragment mass tolerance
was 20 ppm and 0.6 Da, respectively. The *Drosophila melanogaster* database with added
targeted protein sequences was used for PD 2.4 database search with 1% FDR for report of
crosslink results. Identified crosslinked peptides were filtered for Max. XlinkX Score > 40
containing at least two identified MS^3 spectra for each pair of crosslinked peptides. Search Results
were exported by the software as a spreadsheet. The crosslinked peptides were analyzed by
xiNET for generating the viewer of inter or inner crosslinks between two target proteins.
506

SWFTI assay
S2 cells were co-transfected as described above with pAC 5.1 vectors of CLIP-CRY-377L and
TIM-SNAP-HA-ALFA variants at a mass ratio of 3:1. The pulldown assay was carried out as
follows, similar to that previously described[51]. 5 mL of S2 cell pellets were lysed with 400 μL lysis
buffer containing non-ionic detergent (50 mM Tris buffer, pH 8, 150 mM NaCl, 1% IGEPAL CA-
630 (Cat# I8896, Sigma Aldrich), 10% glycerol and 1x protease inhibitor (Cat# A32965,
ThermoFisher)) was added to cell pellets harvested from 5 mL cells. EDTA was omitted to avoid
disturbing SNAP/CLIP dye labeling. All buffer pHs were adjusted at the working temperature. After
cells were incubated with lysis buffer on ice for 30 mins (flicking the tube every 10 min to avoid
vortexing), cell debris was removed by spinning at 15,000 × g for 15 minutes at 4°C.
To label the lysate sample with SNAP/CLIP dyes, 1 part of lysate was diluted with 2 parts of
labeling buffer (50 mM Tris buffer, pH 8, 150 mM NaCl, 10% glycerol) to decrease the
concentration of non-ionic detergent to less than 0.5%, as recommended by the manufacturer
(New England Biolabs). Then 27 μL of diluted lysate was mixed with 1 μL of 0.1 mM SNAP dye
(Cat# S9102S, SNAP-Cell 647-SiR), 1 ul 0.1 mM CLIP dye (Cat# S9217S, CLIP-Cell 505) and 1
ul 30 mM DTT and incubated in the dark at room temperature for 1 hour. 10 uL 4x Laemmli sample
buffer (Cat# 1610747, BioRad) and 1.5 μL 2-mercaptoethanol (BME) was added to the 30 μL mix,
boiled at 95 °C for 5 minutes. After labeling with the SNAP/CLIP substrate, the sample was
protected from bright light to avoid photobleaching. 20 μL of the sample was loaded on a gradient
stain-free gel (Cat# 4568095, BioRad) to perform SDS-PAGE. The result was imaged with
Chemidoc (BioRad) and quantified by image lab software (BioRad).
528

To label CRY and TIM on the affinity resin, ~350 μL cell lysate was mixed with 10 μL pre-washed
magnetic HA resin slurry (Cat# 88836, ThermoFisher) and incubated on an end-to-end rotator

overnight at 4 °C. The dark sample was covered by foil and kept in the dark. Next day, the resin

was washed with room temperature 500 µL TBS-T (20 mM Tris buffer, pH 7.6, 150 mM NaCl,

0.05% Tween-20) four times. Then HA resin was resuspended in 27 µL labeling buffer, 1 µL 0.1

mM SNAP dye, 1 µL 0.1 mM CLIP dye and 1 µL 30 mM DTT. The tube was incubated in the dark

on an end-to-end rotator at room temperature for one hour. Subsequently, 10 µL 4x Laemelli

sample buffer and 1.5 µL BME were added to the 30 µL resin mix. The tube was boiled at 95 °C

for 10 minutes to denature and elute CRY and TIM from HA resin. SDS-PAGE and imaging were

performed the same as above. Effective KD values were calculated as described elsewhere[51].

Negative Stain Electron Microscopy

Initial CRY:TIM samples were screened by negative-stain. Formvar/carbon 200 mesh copper

grids (Electron Microscopy Sciences) was plasma cleaned by the GloQube glow discharge

system (Electron Microscopy Sciences). Ten µL of 0.05 mg/ml CRY:TIM solution was applied

onto the grid, incubated for 30 seconds, and the excess solution was blotted with filter paper. 10

µl of 2% uranyl acetate was then applied to the grid twice for 30-second incubation followed by

blotting. The grid was air-dried for 5 minutes and visualized by Philips Morgagni 268 Transmission

Electron Microscope at 100 keV.

Cryo-EM grid preparation and data collection of CRY:TIM complex

Four µL of 0.5 mg/ml CRY:TIM solution was applied to glow-discharged grids (300 mesh

Quantifoil Au, R1.2/1.3) that were purchased from Electron Microscopy Sciences. After 30

seconds incubation, the excess solution was blotted for 4 sec with filter paper by a Vitrobot Mark

IV (Thermo Fischer Scientific). Grids were subsequently vitrified in liquid ethane and stored in

liquid nitrogen. Cryo-EM images of CRY:TIM complex were collected on a Talos Arctica (Thermo

Fischer Scientific) operated at 200 keV at a nominal magnification of 79,000x with Gatan GIF

Quantum LS Imaging energy filter, using a Gatan K3 direct electron camera in counting mode,

corresponding to a pixel size of 0.516 Å. A total of 1636 images stacks were obtained with a

defocus rage of -0.8 to -2.0 µm by EPU (Thermo Fischer Scientific). Each stack movie was

recorded for a total dose of 53 e-/Å2.

Cryo-EM data processing of the CRY:TIM complex

Dose fractionated image stacks were subjected to motioncor2[53] wrapped in Relion 3.1.3 [54]

followed by patch CTF estimation in CryoSparc v3.2.0[55]. The micrographs whose CTF estimation

is greater than 4 Å were chose for later processing, which yielded 1473 micrographs. Blob picker

in CryoSparc was used on the three hundred randomly selected micrographs, which was followed

by 2D classification and generation of ~23k particles for the first-round Topaz train[56]. Then the

trained model was applied to all 1473 micrographs and ~48k particles were extracted and selected

by 2D classification for the second-round of Topaz train particle identification. After two rounds of

Topaz train and extraction, in totality ~2M particles were picked. Following 2D classification non-

relevant particles were removed and only the best classes were kept (~350k particles). The

particle stack was subjected to Bayesian polishing in RELION 3.1.3 and imported back to

CryoSparc for the further Ab Initio reconstruction and Heterogenous refinement with 3 classes.

The second and third classes were selected, and multiple rounds of Homogenous and Non-

uniform Refinement were performed to yield a global 3.48 Å (~98k particles) and a global 3.3 Å

(~160k particles) map using "gold standard" FSC at 0.143 cut-off, respectively. The 3.3 Å map is

subjected to the model building. C1 symmetry was used throughout data processing. The

workflow is provided in Extended Figure 1.

Model building and refinement of the CRY:TIM complex

The initial CRY model is adapted from the PDB entry 4GU5, and the initial TIM model was fetched

from Alphafold Protein Structure Database entry P49021[57]. Both models were manually docked

into the map by UCSF ChimeraX[58]. The model was subjected to ISOLDE[59] to fit the loops which

undergo conformational changes into the map density. Intrinsically disordered loops of TIM which

were not visualized in the map were not included in the model. The CRY:TIM model was subjected

to multiple rounds of automatic refinement using Phenix real space refinement, and manually

building using Coot[60-62]. Validation was performed by MolProbity[63]. The final refinement statistics

is provided in Table S1.

The unassigned polypeptide in the TIM grove was built in both directions and the sequence

register slid along the backbone for a 150 residue stretch in the 250-400 range and each register

real-spaced refined against the electron density and Q-score [64] calculated. Given the helical

nature of the density at one end of the groove, directionality of the sequence was best fit with the

polypeptide having its N-terminal end closest to CRY and the C-terminal end running toward the

Lys313/328 crosslinking partners at the base of the groove. Side chain density in the helical region

showed a variable size pattern that was inconsistent with the repetitive patterns of much of the

disordered region beyond residue 350. C-terminal to the groove peptide Lys313 and Lys328 are

unresolved, but should be closely positioned to their crosslinking partners. To assign sequence

to the disconnected TIM helix binding the PHR C-terminus secondary structure predictions on the

regions of TIM not resolved in the ordered core were performed by RAPTORX[65]. All predicted

helical sequences were fit into the density and refined in real-space with a range of sequence

registers. As an additional check Alphafold2.0[57] was used to predict interactions between CRY

and the disordered region of TIM, with the highest scoring predictions also tested for agreement

with the electron density map. Although Alphafold did predict interactions of TIM in the region of

the peripheral helix, none of the predictions aligned well with the observed density. The

disconnected helix displays strong side-chain density, which sets its directionality and ruled out

many sequence placements. Given the resolution limitations of the cryo-EM maps, we consider

the sequence assignments of both of these helical regions tentative.

Methods References

51   Lin, C., Schneps, C. M., Chandrasekaran, S., Ganguly, A. & Crane, B. R. Mechanistic
insight into light-dependent recognition of Timeless by Drosophila Cryptochrome.
Structure doi:10.1016/j.str.2022.03.010 (2022).

52   Muok, A. R. *et al.* Engineered chemotaxis core signaling units indicate a constrained
kinase-off state. *Science Signaling* **13** (2020).

53   Zheng, S. Q. *et al.* MotionCor2: anisotropic correction of beam-induced motion for
improved cryo-electron microscopy. *Nat Methods* **14**, 331-332 (2017).

54   Scheres, S. H. RELION: implementation of a Bayesian approach to cryo-EM structure
determination. *J Struct Biol* **180**, 519-530 (2012).

55   Punjani, A., Rubinstein, J. L., Fleet, D. J. & Brubaker, M. A. cryoSPARC: algorithms for
rapid unsupervised cryo-EM structure determination. *Nat Methods* **14**, 290-296 (2017).

56   Bepler, T. *et al.* Positive-unlabeled convolutional neural networks for particle picking in
cryo-electron micrographs. *Nat Methods* **16**, 1153-1160 (2019).

57   Jumper, J. *et al.* Highly accurate protein structure prediction with AlphaFold. *Nature* **596**,
583-589 (2021).

58   Pettersen, E. F. *et al.* UCSF ChimeraX: Structure visualization for researchers,
educators, and developers. *Protein Sci* **30**, 70-82 (2021).

59   Croll, T. I. ISOLDE: a physically realistic environment for model building into low-
resolution electron-density maps. *Acta Crystallogr D Struct Biol* **74**, 519-530 (2018).

60   Morin, A. *et al.* Collaboration gets the most out of software. *Elife* **2**, e01456 (2013).

61   Emsley, P., Lohkamp, B., Scott, W. G. & Cowtan, K. Features and development of Coot.
Acta Crystallogr D Biol Crystallogr **66**, 486-501 (2010).

62   Liebschner, D. *et al.* Macromolecular structure determination using X-rays, neutrons and
electrons: recent developments in Phenix. *Acta Crystallogr D Struct Biol* **75**, 861-877
(2019).

63   Chen, V. B. *et al.* MolProbity: all-atom structure validation for macromolecular
crystallography. *Acta Crystallogr D Biol Crystallogr* **66**, 12-21 (2010).

64   Pintilie, G. *et al.* Measurement of atom resolvability in cryo-EM maps with Q-scores. *Nat
Methods* **17**, 328-334 (2020).

65   Kallberg, M. *et al.* Template-based protein structure modeling using the RaptorX web
server. *Nature Protocols* **7**, 1511-1522 (2012).

642
Data availability The cryo-EM density maps and the atomic model for TIM:CRY have been
deposited in the EM Data Bank (EMDB: EMD-27335) and Protein Data Bank (PDB: 8DD7),
respectively.   The mass spectrometry proteomics data have been deposited to the
ProteomeXchange Consortium via the PRIDE partner repository with the dataset identifier
PXD034054.

Acknowledgements We thank the Proteomics and Metabolomics Facility of Cornell University
for providing the mass spectrometry data; Katherine Spoth and Mariena Silvestray-Ramos for
help with EM instrumentation; Yitong Li for suggesting use of the ALFA Tag and experimental
assistance, Greg Merz, Thuy-Tien Nguyen, and Cody Aplin for general aid; and Richard Cerione
and Michael Young for helpful feedback. Figures 1A, inlet in 2B, 4D, and S1 were created with
BioRender.com. This work was supported by NIH grants R35GM122535 to BRC and 1S10
OD017992-01 for the Orbitrap Fusion mass spectrometer. This work made use of the Cornell
Center for Materials Research Shared Facilities which are supported through the NSF (DMR-
1719875).

Author contributions C.L., S.F., C.C.D, and B.R.C. designed research; C.L., S.F., C.C.D, and
B.R.C. conducted research; C.L., S.F., C.C.D, and B.R.C. analyzed data; C.L., S.F., C.C.D, and
B.R.C. wrote the manuscript.

Competing interests The authors declare that they have no competing interests.

Extended Data Figure 1. CRY:TIM purification and cryo-EM workflow. Purification of
CRY:TIM complexes for cryo-EM. (a) SDS-PAGE gel showing complex after elution from the
ALFA resin, followed by crosslinking, followed by SEC. (b) SEC trace of purified complex, peak I:
aggregates, peak II: CRY-TIM complex, and peak III: excessive ALFA peptide. (c) Processing
workflow for CRY:TIM collected on a Talos Arctica (Thermo Fischer Scientific) operated at 200
keV using a Gatan K3 direct electron camera. The red box indicates the final reconstruction in
this study. Data processing was carried out in CryoSparc and RELION. Motion Correlation was
performed by RELION (MotionCor2); with Contrast Transfer Function (CTF) applied in CryoSparc.
Topaz was used for particle classification. Overall resolution determination in CryoSparc by Gold
Standard Fourier Shell Correlation (GSFSC) is shown at the mid-bottom left. Corrected values
give 3.30 Å overall resolution. Local resolution exceeds 2.5 Å, as shown on the bottom right.
Particle orientational sampling is reasonably uniform as shown at the bottom left. Number of
particles (images) holding a given azimuthal and elevation angle are indicated in the 2-
dimensional histogram.

Extended Data Figure 2. TIM domain construction and secondary structure assignments.
Domain construction and secondary structure assignments of TIM.
680
681   **Extended Data Figure 3. CRY domain construction and secondary structure assignments.**
682   Domain construction and secondary structure assignments of CRY.
683
Extended Data Figure 4. Key interactions at the CRY:TIM interface. (a) Interactions between
the bottom half of TIM ARM$_1$-α1 with CRY. ARM1- α1 binds across CRY α14 and α17. The
guanidinium of CRY R446 (on α17) forms π-cation interaction with TIM Y11. TIM Y11 participates
in hydrophobic contacts with CRY P257, L443 and L447. On TIM α1, TIM P8 stacks against CRY
α12 and TIM S15 hydrogen bonds to the backbone carbonyl of CRY A442. (b) In the β-hairpin
connecting ARM$_1$- α1 to ARM$_1$- α2 E21 salt-bridges to R474 and C19 interacts with K445. (c)
Interactions of the CRY protrusion motif. TIM R109, F113, and H110 interact with the protrusion
motif of CRY. TIM ARM$_2$-α3 H110 stacks with CRY H307. Four Met residues (M179, M181 M182,
M185) in the ARM$_3$-α2-α3 loop provide side-chain interactions with the tip of the CRY protrusion
motif. The CRY α10 helix lengthens to residue A295 compared to the dark state and the following
300-306 residues interact directly with ARM$_2$-α3. (d) Interactions of the CRY C-terminal lid and
α20 helix. The restructured C-terminal lid interacts with residues on TIM ARM$_1$- α2. For examples,
L34 make a hydrophobic contact with I436 and E37 hydrogen bonds to the backbone of S434. At

the end of ARM$_1$-α2, Asn30 contacts the N-terminal end of CRY α20. A DSSO crosslink observed

between Lys510 on CRY a20 and Lys41 on ARM1-α3 is fully consistent with the observed

interface.

Extended Data Figure 5. Changes in flavin binding pocket for CRY dark and bound states.

Comparison of the flavin pocket in CRY:TIM compared to in dark-state CRY (PDB 4GU5). (a) In

CRY:TIM, Y250 and L251 interact with the FAD pocket and make space for the top of TIM ARM$_1$-

α3 to insert F58 behind the new conformation of the phosphate-binding loop. (b) The FAD adenine

ring of CRY:TIM slides 1.5 Å relative to the unbound structure. CRY F318 shifts away from its

original position and harbors helix α11 to compensate for the FAD adenine ring adjustment,

following the protrusion motif. (c) G311 swivels away from the inserted TIM L4. New diphosphate

conformations cause rearrangement of the CRY 262-to-264 loop.

Extended Data Figure 6. Exit channel at the CRY:TIM interface and CRY secondary pocket.

(a) Putative exit channel for the additional 23 N-terminal residues of L-TIM. The first residue of

TIM (M1) is colored as red. A solvent accessible channel at the interface between the two proteins

is marked as the green arrow. (b) Electron density for the secondary binding pocket of CRY where

antenna cofactors bind in photolyases and mCRY interact with other Clock proteins (**red** box)

CRY (**yellow**), TIM (**periwinkle blue**), TIM 1163-1174 (**blue**), TIM groove (**navy blue**). Although

the pocket is well formed, there is no obvious density for additional moieties.

Extended Data Figure 7. TIM peripheral helix and groove peptide fitting to the electron

density map and associated Q-scores. (a) Real-space electron density fitting metrics for the

top scoring sequence assignments for the TIM peripheral helix. Real space density agreements

were calculated in Q-scores and are shown above for sidechains and below for mainchains. (b)

Real-space electron density fitting metrics for the top scoring sequence assignments for the TIM

groove-binding polypeptide. Real space density agreements were calculated in Q-scores and are

shown above for sidechains and below for mainchains. The TIM groove binding polypetide was

tested in both directions.

Extended Data Figure 8. Relative binding affinities between fragments of TIM and CRY

H377L by the SWFTI assay. (a) The CRY H377L variant enhances binding affinity for TIM by

stabilizing the TIM-binding conformation of CRY and thereby increases TIM expression levels and

overall facilitates detection of weaker binding variants. sample size = 3. One-way anova was used
to analyze p value. * represents p<0.05, ** p<0.01 (b) Pull down results of CRY-H377L with TIM-
1133-1398. CRY-H377L interacts with the C-terminal regions of TIM. (c) Quantification of the
binding strength between CRY-H377L and TIM (1133-1398) compared to the negative control
(CRY-H377L only with the same amount of resin). The binding strength is defined as the amount
of CRY on the resin divided by the amount of CRY in the lysate sample. The negative control is
normalized to 1. Sample size = 3. Two-tailed t-test yields a p value < 0.0001 (****). Binding
affinities could not be derived from this experiment owing to the challenge of producing excessive
CRY relative to TIM 1133-1398.

Extended Data Figure 9. Comparisons of structural features in the CRY:TIM complex to
those of CRY and TIM homologues. (a) Structural comparison of CRY:TIM with mammalian
CRY bound with mammalian PER. TIM 1163-1174 (**blue**) follows the backbone trace of mPER
(CRYΔ (**yellow**), TIM BD2 1163-1174 (**blue**)). 4CT0: Crystal Structure of Mouse Cryptochrome1
in Complex with Period2 (**magenta**). 4U8H: Crystal Structure of Mammalian Period-
Cryptochrome Complex (**green**). 6OF7: Crystal structure of the CRY1-PER2 complex (**orange**
red). (b) Structural comparison of CRY:TIM (CRY (**yellow**), TIM (**periwinkle blue**), TIM 1163-
1174 (**blue**), TIM groove peptide (**navy blue**)) with the crystal structure of the N-terminal domain
of human Timeless (PDB ID: 5MQI, **salmon**). Note that mammalian TIM does not contain the N-
terminal residues that bind into the CRY flavin pocket. (c) Structural superposition of TIM PAB
(PARP-1 binding) domain with the structure of human TIM PAB bound to the PARP-1 catalytic
domain. Red box highlights the TIM α-helix occupying the same space required for PARP-1
binding. TIM (this study, **periwinkle blue**), 4XHU (PAB:**sky blue**, PARP-1:**charcoal**). (d)
Superposition of TIM and Tof1 within the replisome.

Extended Data Figure 10. Interactions between TIM and its homologs with extended
polypeptides in their respective ARM grooves in relation to crosslinking map of CRY:TIM.
(a) Sequence alignment of top TIM structural homologs as calculated by the DALI server[50]. Boxes
outline the conserved R-XXX-N and RN residues within the ARM motifs that interact with the
extended polypeptide chains of binding partners. Protein Data Bank (PDB) codes are shown on
the left. (b) The extended peptide bound in the TIM groove resembles interactions found in other
ARM-repeat proteins such as Importin-α1, β-catenin, and yeast TIM proteins. ARM-repeat
proteins were structurally superimposed by their ARM-repeat cores to show the alignment of the
extended peptides that bind in the crescent groove common to the family. For clarity only the

groove peptides are displaced. Extended polypeptides found in yeast TOF1 (6SKL) and β-catenin

(3IFQ) and TIM travel on the same lateral axis anti-parallel to the other importin- α, β-catenin, and

TIM protein homologues. The PDB labels indicate the N-terminus of the loop. 6SKL: Cryo-EM

structure of the CMG Fork Protection Complex at a replication fork - Conformation 1 (lavender),

3IFQ: Interaction of plakoglobin and β-catenin with desmosomal cadherins (**gold**), 1JDH: Crystal

structure of β-catenin and htcf-4 (**purple**), 3L3Q: Mouse importin α-pepTM NLS peptide complex

(gray), 3KND: TPX2:importin-αlpha complex (**magenta**), 1IQ1: Crystal structure of the importin-

α(44-54)-importin-α(70-529) complex (**salmon**), 3UKX: Mouse importin-α:Bimax2 peptide

complex (**sky blue**). (c) Map of DSSO lysine residue crosslinking defined by mass spectroscopy

for the CRY:TIM complex. For the LysC+trypsin digested sample, although +2-charged peptides

were excluded in the analysis, 59.79% sequence coverage for CRY and 49.81% for TIM were

obtained. For the LysC+trypsin+GluC digested sample, 25.07% sequence coverage was obtained

for CRY and 25.53% sequence coverage was obtained for TIM. Summary of the cross-links are

given in the attached excel files and the raw data has been submitted to the PRIDE database.

Figure 1. Cryo-EM structure of the CRY:TIM complex. (**a**) Drosophila Cryptochrome (CRY) entrains the circadian clock by initiating degradation of Timeless (TIM) in light. Jetlag (JET), Period (PER), Doubletime (DB), Cycle (CYC), Clock (CLK) and Importin-α1 (IMPα1) compose the core circadian oscillator. (**b**) CRY:TIM docked into the electron density map. Dark-state CRY shown at the bottom (grey, PDB ID: 4GU5). (**c**) Domain arrangements and key regions of CRY and TIM. Features of TIM: CRY-binding domains (CRY-BD1, 1-234, CRY-BD2, 1133-1398), serine/threonine-rich region (ST, 270-290) with key phosphorylation sites starred (S274, T278, T282, S286 and S290), ARM1 (1-248), ARM2 (578-866), nuclear localization sequence (NLS, 545-578), Parp-1 binding domain (PAB, 986-1142; TIM-C, 1051-1096), 23 residue L-TIM extension (L, (23aa)), disordered regions are dashed. Features of CRY: Phosphate binding loop (PL, 249-263), Protrusion motif (PM, 288-306), C-terminal lid (CL, 420-446), C-terminal tail (CTT, 519-542, absent in CRYΔ). (**d**) N-terminus of TIM inserts into the primary flavin binding pocket. 2.4 Å resolution local electron density is shown with notable hydrogen bonds (yellow dashes) and hydrophobic contacts (green circle). (**e**) Interface between CRY and TIM (blue shaded area), TIM groove peptide (in navy blue) and peripheral helix (fitted with 1163-1174 putatively, blue). Inset shows the interactions between TIM 1163-1174 and CRY superimposed with electron density.

Figure 2. Conformational changes of FAD to the CRY C-terminal coupled motif upon binding TIM. (**a**) CRY:TIM (CRY, yellow; TIM, blue) in superposition with dark-state CRY. The N-terminus of TIM replaces the CRY CTT (red), as depicted in the schematic below. Structural elements designated as in **Figure 1**. (**b**) The phosphate binding loop (dark pink) collapses into the flavin pocket and N253 and E254 hydrogen bond (yellow dashes) with the FAD diphosphate conformation (red, orange) that differs substantially from that of the dark state (white bonds). (**c**) R237 replaces the Mg^{2+} counter ion and H377 anchors the new position of the loop. The D410-R381 salt bridge (top) appears to break in response to the new conformation of the ribose backbone (bottom). Reduced side-chain density of R381 suggests increased flexibility in the TIM complex.

a

Damaged DNA

TIM N-term

CRY

6-4 Photolyase

TIM N-term

6-4 cross-linked pyrimidine

L4

L5

FAD

b

FBXL-3 C-term

TIM ARM$_1$-α3

TIM ARM$_1$-α1

CRY FAD

Figure 3. Conservation of recognition modes by CRY family members. (**a**) The interaction of CRY with TIM corresponds closely to how 6-4 photolyase recognizes the 6-4 pyrimidine lesion in damaged DNA (PDB ID:3CVU). TIM Leu4 and Leu5 superimpose on the crosslinked bases and the TIM ARM1- α1 polypeptide closely follows the path of 5'-to-3' strand-separated DNA. (**b**) The TIM and FAD contacts of CRY mimic the interaction of apo-mCRY with the C-terminus of the FBXL3 E3 ligase (PDB ID: 4I6J).

Figure 4. A structural basis for regulation of TIM nuclear entry. (**a**) Electron density (dark blue) for an extended polypeptide resides within a groove wherein members of the β-catenin and α-importin proteins bind their target sequences. (**b**) The TIM groove peptide density shown with the sequence assignment S283-T303. The peptide interacts with β-catenin-conserved Asn and Arg residues and buries residues important for importin-α1 binding (T90 and P92). (**c**) DSSO-mediated lysine crosslinking patterns of disordered residues K313 and K238 shown on the TIM structure. K313 and K328 produce a pattern of crosslinking that localizes them to a position consistent with placement of the S283-T303 groove peptide. The TIM NLS lies in a helix adjacent to the groove but does not occlude T90 and P92. (**d**) Schematic of proposed nuclear entry mechanism of TIM. Phosphorylation of the TIM-ST region by SGG and CK2 releases the ST region from TIM groove to expose TIM T90 and P92. Importin-α1 recognizes TIM by binding adjacent to Thr90 and Pro92, while also binding the TIM NLS within a structurally analogous ARM groove of its own.

Extended Data Fig. 1

Extended Data Figure 1. CRY:TIM purification and cryo-EM workflow. Purification of CRY:TIM complexes for cryo-EM. (a) SDS-PAGE gel showing complex after elution from the ALFA resin, followed by crosslinking, followed by SEC. (b) SEC trace of purified complex, peak I: aggregates, peak II: CRY-TIM complex, and peak III: excessive ALFA peptide. (c) Processing workflow for CRY:TIM collected on a Talos Arctica (Thermo Fischer Scientific) operated at 200 keV using a Gatan K3 direct electron camera. The red box indicates the final reconstruction in this study. Data processing was carried out in CryoSparc and RELION. Motion Correlation was performed by RELION (MotionCor2); with Contrast Transfer Function (CTF) applied in CryoSparc. Topaz was used for particle classification. Overall resolution determination in CryoSparc by Gold Standard Fourier Shell Correlation (GSFSC) is shown at the mid-bottom left. Corrected values give 3.30 Å overall resolution. Local resolution exceeds 2.5 Å, as shown on the bottom right. Particle orientational sampling is reasonably uniform as shown at the bottom left. Number of particles (images) holding a given azimuthal and elevation angle are indicated in the 2-dimensional histogram.

TIM ARM₁:
$\alpha 1$: 1 - 17
$\alpha 2$: 31 - 44
$\alpha 3$: 51 - 59
$\alpha 4$: 61 - 72
$\alpha 5$: 76 - 89
$\alpha 6$: 92 - 102
$\alpha 7$: 106 -126
$\alpha 8$: 129 - 145
$\alpha 9$: 151 - 177
$\alpha 10$: 185 - 201
$\alpha 11$: 204 - 213
$\alpha 12$: 217 - 231
$\alpha 13$: 237 - 247

TIM ARM₁ (1-248)

TIM ARM₂ (545-866)

TIM ARM₂:
NLS: 546 - 570
$\alpha 1$: 578 - 595
$\alpha 2$: 597 - 608
$\alpha 3$: 618 - 634
$\alpha 4$: 648 - 668
$\alpha 5$: 677 - 702
$\alpha 6$: 709 - 722
$\alpha 7$: 726 - 736
$\alpha 8$: 745 - 765
$\alpha 9$: 787 - 797
$\alpha 10$: 799 - 802
$\alpha 11$: 807 - 818
$\alpha 12$: 830 - 842
$\alpha 13$: 848 - 862

TIM PAB-C:
$\alpha 1$: 991 - 1002
$\alpha 2$: 1006 - 1017
$\alpha 3$: 1020 - 1027
$\alpha 4$: 1040 - 1046
$\alpha 5$: 1059 - 1066
$\alpha 6$: 1068 - 1076
$\alpha 7$: 1103 -1107

TIM PAB-C (986-1142)

Extended Data Fig. 2

Extended Data Figure 2. TIM domain construction and secondary structure assignments. Domain construction and secondary structure assignment of TIM.

CRY N-terminal
α/β domain:
β1: 6 - 11
α1: 21 - 32
β2: 35 - 42
α2: 54 - 75
β3: 82 - 85
α3: 88 - 98
β4: 101 - 106
α4: 112 - 128
β5: 131 - 135

Linker region (not highlighted)
α5: 143 - 149
α6: 159 - 168
α7: 190 - 195

α/β domain (8-137)

Protrusion motif (PM)

TIM BD2 (1163-1174)

to C-terminal tail (CTT)

CRY α-helical domain (FAD binding):
α8: 227 - 247
α9: 267 - 272
α10: 277 - 288
(277-293 in CRY:TIM complex)
α11: 310 - 321
α12: 347 - 355
α13: 361 - 372
α14: 378 - 388
α15: 397 - 406
α16: 413 - 424
α17: 440 - 447
α18: 452 - 457
α19: 477 - 482
α20: 498 - 515

Phosphate binding loop (PL)

C-terminal lid (CL)

CRY α-helical domain (225-517)

View direction

Extended Data Fig. 3

Extended Data Figure 3. CRY domain construction and secondary structure assignments. Domain construction and secondary structure assignment of CRY.

Extended Data Fig. 4

Extended Data Figure 4. Key interactions at the CRY:TIM interface. (a) Interactions between the bottom half of TIM ARM$_1$-α1 with CRY. ARM1- α1 binds across CRY α14 and α17. The guanidinium of CRY R446 (on α17) forms π-cation interaction with TIM Y11. TIM Y11 participates in hydrophobic contacts with CRY P257, L443 and L447. On TIM α1, TIM P8 stacks against CRY α12 and TIM S15 hydrogen bonds to the backbone carbonyl of CRY A442. (b) In the β-hairpin connecting ARM$_1$- α1 to ARM$_1$- α2 E21 salt-bridges to R474 and C19 interacts with K445. (c) Interactions of the CRY protrusion motif. TIM R109, F113, and H110 interact with the protrusion motif of CRY. TIM ARM$_2$-α3 H110 stacks with CRY H307. Four Met residues (M179, M181 M182, M185) in the ARM$_3$-α2-α3 loop provide side-chain interactions with the tip of the CRY protrusion motif. The CRY α10 helix lengthens to residue A295 compared to the dark state and the following 300-306 residues interact directly with ARM$_2$-α3. (d) Interactions of the CRY C-terminal lid and α20 helix. The restructured C-terminal lid interacts with residues on TIM ARM$_1$- α2. For examples, L34 make a hydrophobic contact with I436 and E37 hydrogen bonds to the backbone of S434. At the end of ARM$_1$- α2, Asn30 contacts the N-terminal end of CRY α20. A DSSO crosslink observed between Lys510 on CRY a20 and Lys41 on ARM1-α3 is fully consistent with the observed interface.

Extended Data Fig. 5

Extended Data Figure 5. Changes in flavin binding pocket for CRY dark and bound states. Comparison of the flavin pocket in CRY:TIM compared to in dark-state CRY (PDB 4GU5). (a) In CRY:TIM, Y250 and L251 interact with the FAD pocket and make space for the top of TIM ARM_1-α3 to insert F58 behind the new conformation of the phosphate-binding loop. (b) The FAD adenine ring of CRY:TIM slides 1.5 Å relative to the unbound structure. CRY F318 shifts away from its original position and harbors helix α11 to compensate for the FAD adenine ring adjustment, following the protrusion motif. (c) G311 swivels away from the inserted TIM L4. New diphosphate conformations cause rearrangement of the CRY 262-to-264 loop.

a

b

Extended Data Fig. 6

Extended Data Figure 6. Exit channel at the CRY:TIM interface and CRY secondary pocket. (a) Putative exit channel for the additional 23 N-terminal residues of L-TIM. The first residue of TIM (M1) is colored as red. A solvent accessible channel at the interface between the two proteins is marked as the green arrow. (b) Electron density for the secondary binding pocket of CRY where antenna cofactors bind in photolyases and mCRY interact with other Clock proteins (**red** box) CRY (**yellow**), TIM (**periwinkle blue**), TIM 1163-1174 (**blue**), TIM groove (**navy blue**). Although the pocket is well formed, there is no obvious density for additional moieties.

Extended Data Fig. 7

Extended Data Figure 7. TIM peripheral helix and groove peptide fitting to the electron density map and associated Q-scores. (a) Real-space electron density fitting metrics for the top scoring sequence assignments for the TIM peripheral helix. Real space density agreements were calculated in Q-scores and are shown above for sidechains and below for mainchains. (b) Real-space electron density fitting metrics for the top scoring sequence assignments for the TIM groove-binding polypeptide. Real space density agreements were calculated in Q-scores and are shown above for sidechains and below for mainchains. The TIM groove binding polypetide was tested in both directions.

a

b

c

Lysate

Resin

TIM (bait)   CRY

Extended Data Fig. 8

Extended Data Figure 8. Relative binding affinities between fragments of TIM and CRY H377L by the SWFTI assay. (a) The CRY H377L variant enhances binding affinity for TIM by stabilizing the TIM-binding conformation of CRY and thereby increases TIM expression levels and overall facilitates detection of weaker binding variants. sample size = 3. One-way anova was used to analyze p value. * represents $p<0.05$, ** $p<0.01$ (b) Pull down results of CRY-H377L with TIM-1133-1398. CRY-H377L interacts with the C-terminal regions of TIM. (c) Quantification of the binding strength between CRY-H377L and TIM (1133-1398) compared to the negative control (CRY-H377L only with the same amount of resin). The binding strength is defined as the amount of CRY on the resin divided by the amount of CRY in the lysate sample. The negative control is normalized to 1. Sample size = 3. Two-tailed t-test yields a p value < 0.0001 (****). Binding affinities could not be derived from this experiment owing to the challenge of producing excessive CRY relative to TIM 1133-1398.

Extended Data Fig. 9

Extended Data Figure 9. Comparisons of structural features in the CRY:TIM complex to those of CRY and TIM homologues. (a) Structural comparison of CRY:TIM with mammalian CRY bound with mammalian PER. TIM 1163-1174 (**blue**) follows the backbone trace of mPER (CRYΔ (**yellow**), TIM BD2 1163-1174 (**blue**)). 4CT0: Crystal Structure of Mouse Cryptochrome1 in Complex with Period2 (**magenta**). 4U8H: Crystal Structure of Mammalian Period-Cryptochrome Complex (**green**). 6OF7: Crystal structure of the CRY1-PER2 complex (**orange red**). (b) Structural comparison of CRY:TIM (CRY (**yellow**), TIM (**periwinkle blue**), TIM 1163-1174 (**blue**), TIM groove peptide (**navy blue**)) with the crystal structure of the N-terminal domain of human Timeless (PDB ID: 5MQI, **salmon**). Note that mammalian TIM does not contain the N-terminal residues that bind into the CRY flavin pocket. (c) Structural superposition of TIM PAB (PARP-1 binding) domain with the structure of human TIM PAB bound to the PARP-1 catalytic domain. Red box highlights the TIM α-helix occupying the same space required for PARP-1 binding. TIM (this study, **periwinkle blue**), 4XHU (PAB:**sky blue**, PARP-1:**charcoal**). (d) Superposition of TIM and Tof1 within the replisome.

a

b

c

CRY
TIM
LysC+trypsin
LysC+trypsin+GluC

Extended Data Fig. 10

Extended Data Figure 10. Interactions between TIM and its homologs with extended polypeptides in their respective ARM grooves in relation to crosslinking map of CRY:TIM. (a) Sequence alignment of top TIM structural homologs as calculated by the DALI server[50]. Boxes outline the conserved R-XXX-N and RN residues within the ARM motifs that interact with the extended polypeptide chains of binding partners. Protein Data Bank (PDB) codes are shown on the left. (b) The extended peptide bound in the TIM groove resembles interactions found in other ARM-repeat proteins such as Importin-α1, β-catenin, and yeast TIM proteins. ARM-repeat proteins were structurally superimposed by their ARM-repeat cores to show the alignment of the extended peptides that bind in the crescent groove common to the family. For clarity only the groove peptides are displaced. Extended polypeptides found in yeast TOF1 (6SKL) and β-catenin (3IFQ) and TIM travel on the same lateral axis anti-parallel to the other importin-α, β-catenin, and TIM protein homologues. The PDB labels indicate the N-terminus of the loop. 6SKL: Cryo-EM structure of the CMG Fork Protection Complex at a replication fork - Conformation 1 (lavender), 3IFQ: Interaction of plakoglobin and β-catenin with desmosomal cadherins (**gold**), 1JDH: Crystal structure of β-catenin and htcf-4 (**purple**), 3L3Q: Mouse importin α-pepTM NLS peptide complex (gray), 3KND: TPX2:importin-αlpha complex (**magenta**), 1IQ1: Crystal structure of the importin-α(44-54)-importin-α(70-529) complex (**salmon**), 3UKX: Mouse importin-α:Bimax2 peptide complex (**sky blue**). (c) Map of DSSO lysine residue crosslinking defined by mass spectroscopy for the CRY:TIM complex. For the LysC+trypsin digested sample, although +2-charged peptides were excluded in the analysis, 59.79% sequence coverage for CRY and 49.81% for TIM were obtained. For the LysC+trypsin+GluC digested sample, 25.07% sequence coverage was obtained for CRY and 25.53% sequence coverage was obtained for TIM. Summary of the cross-links are given in the attached excel files and the raw data has been submitted to the PRIDE database.