1

Single-cell transcriptome analysis reveals heterogeneity and a dynamic

regenerative response of quiescent radial glia in adult zebrafish brain

Manana Kutsia, Yuki Takeuchi, Nishtha Ranawat, and Ichiro Masai*

Developmental Neurobiology Unit, Okinawa Institute of Science and Technology
Graduate University, Tancha 1919-1, Onna, Okinawa 904-0495, Japan

* Correspondence should be addressed: masai@oist.jp

Running Title: Heterogeneity and regenerative response of quiescent radial glia in zebrafish

Key Words: radial glia, adult neural stem cell, neural regeneration, microglia, zebrafish

Word count of the main text (Introduction/Results/Discussion): 9,125

Abstract

In zebrafish telencephalon, radial glial cells (RGs) show a remarkable ability to regenerate damaged neural tissue by re-initiating cell proliferation to produce neural precursors to rebuild the lost neural circuit. However, it is not fully understood how RGs respond to brain damage to initiate this regenerative response. Here we applied single-cell transcriptomics to RGs in adult zebrafish telencephalon and identified five RG subtypes, which are classified into four quiescent RGs (qRGs) and one proliferating RG (pRG). The four qRGs differentially express distinct subsets of qRG markers, suggesting heterogeneity of qRG in zebrafish adult brain. Interestingly, one qRG subtype shows high expression of ribosomal proteins, and its fraction increases in response to brain damage. Consistently, the mTOR pathway is activated in RGs near the injury site. It was reported that inflammatory responses of brain-resident immune cells, microglia, are required for inducing regenerative responses of RGs in zebrafish. Genetical elimination of microglia not only suppressed the damage-induced regenerative response of RGs, but also decreased the fraction of the ribosomal expression-enriched qRGs. Our pseudo-time analysis suggests that putative dormant RGs produce ribosomal expression-enriched qRGs through activation of ribosomal genesis, as well as suppression of cholesterol biogenesis, and pRGs through activation of the JAK/STAT pathway. Our findings reveal heterogeneity of qRGs in adult zebrafish brain and their dynamic regenerative response to brain damage.

(219 words)

Introduction

In mammals, most neurons in the brain are generated during embryonic development; however, production of new neurons is sustained throughout adulthood in only two regions of the telencephalon, the ventricular/subventricular zone (V/SVZ) of the lateral ventricle and the subgranular zone (SGZ) of the dentate gyrus of the hippocampus (Kriegstein and Alvarez-Buylla, 2009; Zhao et al., 2008). In the SGZ, radial glial stem cells (Type 1 cells) generate amplifying progenitor cells (Type 2 cells) and then neuroblasts (Type 3 cells), which eventually generate granule interneurons. Dentate gyrus granule neurons extend axons called mossy fibers to target pyramidal neurons in the CA3 and CA2 hippocampal regions (Cope and Gould, 2019). In the V/SVZ, neural stem cells (B1 cells) generate transit-amplifying cells (C cells). C cells subsequently generate neuroblasts (A cells), which migrate toward the olfactory bulb, where they differentiate into local interneurons (Otsuki and Brand, 2020).

In contrast to the limited neurogenic capability of adult mammals, teleosts such as zebrafish show remarkably higher proliferative and regenerative potential in the nervous system (Grandel and Brand, 2013; Lindsey and Tropepe, 2006). In zebrafish, composition of neurogenic niches and the cellular nature of neurogenic progenitors have been intensively studied. Classic analysis of zebrafish adult brain with proliferative markers identified 16 different proliferative zones, which produce newborn neurons (Adolf et al., 2006; Grandel et al., 2006). In zebrafish, two distinct neurogenic proliferation zones are formed at the ventricular surface of the dorsal and ventral telencephalon, respectively (Barbosa and Ninkovic, 2016; Jurisch-Yaksi et al., 2020). In the dorsal telencephalon, called the pallium, radial glial cells (RGs) are located at the ventricular surface and extend long radial processes, whose endfeet reach the pial surface close to a blood vessel at the lateral margin of the dorsal telencephalon (Ganz et al., 2010; Marz et al., 2010). These pallial ventricular RGs highly express glial markers such as GFAP, Vimentin and S100β, and generate newborn neurons. In the lateral domain of the pallium, the same process operates, but RGs are generated during juvenile and adult stages from neuroepithelial progenitor cells located at the interface between the pallium and subpallium (Dirian et al., 2014), suggesting that the source of RGs is different from that of RGs in the medial and dorsal pallium. On the other hand, in the ventral telencephalon, called the subpallium, neuroepithelial-like progenitor cells

are located in the ventricular region (Ganz et al., 2010; Marz et al., 2010). They rapidly

proliferate to generate neuroblasts migrating toward the olfactory bulb, so this

neurogenic niche may share characteristics with glial marker-negative proliferating

neuroblasts in mouse V/SVZ. Thus, there is heterogeneity in progenitor cell types in the

adult zebrafish telencephalon.

In general, severe trauma brain injury (TBI) causes direct damage to neural

tissue in the form of primary lesions (Xiong et al., 2013). Following these primary

injuries, lasting damage is sustained through a series of complex events called

secondary injuries (Jassam et al., 2017). In mammals, brain damage induces acute

inflammation, microglial activation, and reactive astrogliosis, leading to scar formation,

which limits spread of damage into unaffected CNS area. However, significant

regeneration does not occur. In contrast, stab injuries to the zebrafish telencephalon

cause an acute inflammatory response and reactive gliosis; however, neither persistent

gliosis, fibrotic scar formation nor chronic inflammation occurs (Baumgart et al., 2012;

Kishimoto et al., 2012; Kroehne et al., 2011). Rather, RGs upregulate cell proliferation

at 2-3 days post-lesion (dpl), and subsequently promote neurogenesis, which leads to

rebuild neural circuits of damaged tissue.

Quantitative clonal analysis of RGs in zebrafish dorsal pallium revealed that

RGs comprise two groups of neural stem cells (NSCs): deeply quiescent and self-

renewing neural stem cells called reservoir NSCs and neural stem cells called

operational NSCs, which promote neurogenesis. Operational NSCs undergo various cell

division modes: symmetric gliogenic division, asymmetric neurogenic division,

symmetric neurogenic division, and direct conversion to neurons without cell division,

in normal homeostasis (Rothenaigner et al., 2011; Than-Trong et al., 2020).

Furthermore, *in vivo* live imaging of RGs in zebrafish dorsal pallium confirmed that

neurons are generated by both direct conversion from NSCs to post-mitotic neurons and

asymmetric neurogenic division to generate newborn neurons or via intermediate

progenitors (Barbosa et al., 2015). Interestingly, in injured telencephalon, RGs generate

neural progenitors migrating toward injured sites by expanding a new type of symmetric

cell division, which depletes the pool of NSCs, but generates two neural progenitor cells

that are never observed under normal homeostatic conditions. Thus, compared with

NSCs in mouse V/SVZ and SGZ, zebrafish dorsal pallium NSCs show unique modes of

cell division in intact and injured conditions (Barbosa and Ninkovic, 2016).

In zebrafish, the process controlling NSC activation from quiescent RGs (qRGs)

to proliferating RGs (pRGs) and non-glial proliferating neuroblasts is regulated by the

Notch signaling pathway. In the pallium of zebrafish telencephalon, Notch 3 is

prominently expressed in qRGs and pRGs, whereas Notch 1a/1b are expressed in pRGs

and non-glial progenitor cells (Alunni et al., 2013; de Oliveira-Carlos et al., 2013).

Indeed, Notch 3 inhibits qRGs from entering symmetric gliogenic cell division, which

limits amplifying neurogenic division, whereas Notch1a stimulates qRGs to

differentiate into pRGs (Alunni et al., 2013). RNA-seq analysis confirmed that Notch 3

and its downstream transcription factor Hey1 promote stemness and quiescence of

qRGs in the pallium of the telencephalon (Than-Trong et al., 2018). Recently, single-

cell transcriptomic analysis has been applied to NSCs in mouse brain (Basak et al.,

2018; Dulken et al., 2017; Llorens-Bobadilla et al., 2015) and RGs in zebrafish

telencephalon (Cosacak et al., 2019; Lange et al., 2020). scRNA-seq analysis of RGs

using zebrafish transgenic lines *Tg[her4.1:mCherry; elavl3:GFP]*, which differentially

labels RGs, newborn neurons, and mature neurons, revealed the diversity of RG

progeny, consisting of qRGs, pRGs, newborn neurons, and mature neurons (Lange et

al., 2020). This study also identified 47 genes that are differentially expressed in qRGs

and pRGs. Another scRNA-seq analysis of RGs using zebrafish *Tg[her4.1:GFP]*,

identified nine RG clusters. However, the diversity of these clusters is likely to

represent heterogeneity of RGs corresponding to different ventricular domains of the

telencephalon (Cosacak et al., 2019). Thus, it is still unknown how many cellular states

can be specified in the NSC activation process from qRGs to pRGs in normal intact and

injured zebrafish telencephalon.

Inflammation occurs rapidly following brain injury and involves activation of

different types of immune cells. Although the immune system is beneficial for clearance

of neurotoxic debris induced by brain damage, several mouse studies suggest that

inflammation is largely unfavorable for neural precursor cell proliferation, which is

required for tissue regeneration. Microglial activation in the brain reduces hippocampal

neurogenesis, and treatment of anti-inflammatory drugs restores it (Ekdahl et al., 2003;

Monje et al., 2003). Such detrimental immune responses are involved in brain damage

pathogenesis and become an obstacle to recovery in mammals. However, surprisingly,
in zebrafish, the immune response after traumatic brain injury is required to initiate the
regenerative response of RGs in the telencephalon (Kyritsis et al., 2012), suggesting an
interaction between RGs and immune cells, including microglia. However, it remains to
be seen how immune cells initiate regenerative responses of RGs.

In this study, to understand how RGs respond to brain damage to initiate a
regenerative response, we applied single-cell transcriptomics to RGs in zebrafish adult
telencephalon using the zebrafish transgenic line *Tg[GFAP:dTomato]*. We identified
five RG subtypes, which are classified into four quiescent RGs (qRG) and one
proliferating RG (pRG). The four qRGs express distinct subsets of qRG markers,
suggesting heterogeneity of qRGs in adult zebrafish brain. Interestingly, one qRG
subtype shows high expression of ribosomal proteins, and its fraction increases in
response to brain damage. Consistently, the mTOR pathway is activated in RGs near the
injury site. Next, to investigate the role of microglia in regenerative responses of RGs,
we eliminated microglia. We confirmed that microglial elimination suppresses damage-
induced regenerative responses of RGs. Interestingly, our scRNA-seq analysis revealed
that the fraction of ribosomal expression-enriched qRGs is markedly decreased in
microglia-depleted injury brain, indicating that the transition of ribosomal expression-
enriched qRGs depends on microglia function. Our pseudo-time analysis suggests that
putative dormant RGs produce the ribosomal expression-enriched qRGs through
activation of ribosomal genesis as well as suppression of cholesterol biogenesis, and
pRGs through activation of the JAK/STAT pathway. Our findings reveal heterogeneity
of qRGs in adult zebrafish brain and their dynamic regenerative responses to brain
damage.

Results

TBI induces a proliferative response of RGs in the telencephalon of adult zebrafish

In zebrafish telencephalon, RGs function as neural stem cells to regenerate damaged

neural tissue by re-entering cell cycle and producing neural progenitor cells (Barbosa

and Ninkovic, 2016; Kishimoto et al., 2012; Kroehne et al., 2011; Than-Trong and

Bally-Cuif, 2015). To understand this mechanism, stab-mediated injury was employed

in the dorsal region of a hemisphere of the telencephalon of adult zebrafish, 3 – 7

months post-fertilization (mpf) (Fig. 1A). After BrdU treatment for 12 h, injured fish

were sacrificed at 1, 3, 7 and 14 dpl, and the telencephalon was dissected and fixed with

paraformaldehyde (PFA) (Fig. 1B). Using a vibratome, 100-μm slices covering the

telencephalon along the anterior and posterior axes were prepared and labeled with anti-

GFAP and ani-BrdU antibodies. GFAP is a glial maker and is expressed in all RGs of

the zebrafish telencephalon (Lam et al., 2009). RG cell bodies are located in the

ventricular zone of the telencephalon and extend processes to the basal area of the brain

(Ganz et al., 2010; Marz et al., 2010) (Fig. 1C). Double labeling of the telencephalon

with anti-BrdU and anti-GFAP antibodies visualized proliferating RGs, which re-

entered cell-cycle progression (Fig. 1D). We examined BrdU$^+$ RGs in telencephalic

slices at 1, 3, 7, and 14 dpl (Fig. 1E) and counted the total number of BrdU$^+$ RGs (Fig.

1F). The number of BrdU$^+$ RGs was significantly higher at 3 dpl than that of non-

injured controls. However, there was no significant difference at other time points

between injured samples and non-injured controls. We also confirmed that the number

of BrdU$^+$ RGs was significantly higher in lesioned hemispheres than in control intact

hemispheres (Fig. 1G). These data suggest that cell proliferation is activated in RGs at 3

dpl, which is consistent with previous reports (Kishimoto et al., 2012; Kroehne et al.,

2011). The telencephalon is divided into two major areas, the dorsal telencephalic area

(D) and the ventral telencephalic area (V) (Ganz et al., 2010; Than-Trong and Bally-

Cuif, 2015) (Fig. 1C). Accordingly, the ventricular zone of D is divided into three areas,

the medial (Dm), dorsal (Dd), and lateral zones of D (Dl). Similarly, the ventricular

zone of V is divided into two areas, the dorsal nucleus (Vd) and the ventral nucleus of V

(Vv). We examined which ventricular zone area most actively responds to TBI. RGs in

the Dd region responded significantly to TBI and incorporated BrdU, whereas RGs in

other areas did not (Fig. 1H). This is reasonable, because the Dd region is closest to the

site of the administered stab injury. Taken together, TBI induces a regenerative
response of RGs in adult zebrafish telencephalon.

scRNA-seq analysis on non-injured controls and 3-dpl RGs revealed five distinct
populations of RGs
To identify the mechanism that regulates the regenerative response of RGs, we applied
scRNA-seq analysis to RGs in 3-dpl and non-injured control telencephalons (Fig. 2A).
Stab injuries were inflicted in telencephalons of 5-mpf adult zebrafish carrying a
transgene *Tg[gfap:dTomato]*, which expresses dimeric Tomato (dTomato) in RGs under
the control of the *gfap* promoter (Satou et al., 2012; Than-Trong et al., 2020). At 3 dpl,
telencephalons were homogenized with papain to prepare cell suspensions, which were
subsequently subjected to flow cytometry-assisted cell sorting (Fig. 2-figure supplement
1AC). We used a 10x genomic system to prepare scRNA-seq libraries. After read
sequencing was done, reads were mapped to the zebrafish genomic database GRCz11.
Next, we evaluated nUMI, nGene, and mitochondrial genes in both datasets separately
to ensure that all data were comparable (Fig. 2-figure supplement 1D). After removing
cells that exhibited unique gene counts below the minimum threshold (200 genes) or
above the maximum threshold (2500 genes), or mitochondrial genes above the
maximum threshold (5% counts), we obtained 567 and 747 cells from non-injured
control and 3-dpl samples, respectively. We conducted unbiased clustering of the
datasets from non-injured control and 3-dpl samples separately, using the Seurat
package (Butler et al., 2018) . In the non-injured control sample, cells were classified
into 6 clusters (Fig. 2-figure supplement 2A). Based on a previous report on zebrafish
genes specifically expressed in qRGs, pRGs, neuroblasts, newborn neurons (NBNs),
ependymal cells, and oligodendrocyte precursor cells (OPCs) (Lange et al., 2020), 6
clusters corresponded to 3 types of qRGs, pRG/Neuroblast/NBN, ependymal cells/OPC,
and probably pineal gland (Fig. 2-figure supplement 2B, Supplementary table 1). On the
other hand, in the 3-dpl sample, cells were classified into 7 clusters (Fig. 2-figure
supplement 3A), which correspond to 4 types of qRGs, qRG/OPC,
pRG/Neuroblast/NBN, ependymal cells (Fig. 2-figure supplement 3B, Supplementary
table 2). Thus, each datasets had sufficient quality to apply further clustering analysis of
merged datasets.

Next, we utilized the Harmony merging algorithm and applied clustering

analysis to the two merged datasets, non-injured controls and 3-dpl samples (Korsunsky

et al., 2019). gfap:dTomato$^+$ cells were classified into 9 clusters, corresponding to 4

types of qRGs, pRG/Neuroblast/MBN, MN/NBN, OPC, ependymal cells, and pineal

gland (Fig. 2B). Heat map analysis of TOP10 genes differentially expressed in each

cluster confirmed clear separation of each cluster (Fig.2-figure supplement 4). We also

confirmed how each of the 9 clusters of the merged datasets corresponded to 6 clusters

of non-injured control cells (Fig.2-figure supplement 2C) and 7 clusters of 3-dpl cells

(Fig.2-figure supplement 3C).

Four qRGs differentially express a distinct subset of qRG-specific genes

4 qRG clusters express qRG markers, and two of them also express neuroepithelial

markers, *krt8* and *zic1*, respectively (Fig. 2C). So, we named these two clusters

qRG/NeuroEp1 and qRG/NeuroEp2. In other two qRG clusters, one expressed *lgals2a*

and the other expressed *s100β* in TOP10 genes (Fig.2-figure supplement 4), which are

named qRG1 and qRG2, respectively (Fig. 2C). To further characterize the 4 qRG

clusters, we focused on 47 genes that were reported to be specifically expressed in

ccnd1-negative qRG (Lange et al., 2020). Among these 47 genes, only one gene, *drg3a*,

was not included for analysis. Interestingly, the other 46 genes were differentially

expressed in each of the 4 qRG clusters (Fig. 2D). These data suggest that

qRG/NeuroEp1, qRG/NeuroEp2, qRG1 and qRG2 correspond to different states of

qRGs.

Interestingly, these 46 qRG markers are separated into 4 gene modules, each of

which were highly expressed in qRG/NeuroEp1, qRG/NeuroEp2, qRG1 and qRG2 (Fig.

2-figure supplement 5A). Gene module 1 contains 6 qRG genes: *hipk2*, *mfge8a*, *luzp2*,

ptn, *si:ch211-251b21.1*, and *s1pr1*, that are prominently expressed in qRG/NeuroEp1

(Fig. 2-figure supplement 5BC). Gene module 2 contains 14 qRG genes: *gmnn*, *grm2b*,

slc6a11b, cxcl12a, slc4a4a, slc1a2b, ppap2d, atp1b4, cx43, atp1a1b, hepacama, psap,

fgfbp3, and *aldocb*, that are prominently expressed in qRG/NeuroEp2 (Fig. 2-figure

supplement 5D). Gene module 3 contains 10 qRG genes: *fabp7a, mt-co3, mt-co2,*

lgals2a, selenop, mt-nd2, mt-atp6, mt-atp8, zgc:153704, and *mgll*, that are prominently

expressed in qRG1 (Fig. 2-figure supplement 5E). Gene module 4 contains 16 qRG

genes: *b2ml, tegt, itm2ba, ptgdsb.2, ptgdsb.1, mt2, anxa11b, glula, sdcbp2, mt-nd1, mt-*

cyb, ckbb, acbd7, s100b, gapdhs, and *efhd1*, that are prominently expressed in qRG2

(Fig. 2-figure supplement 5F).

Previous scRNA-seq analysis of zebrafish telencephalon revealed that

heterogeneity of qRG clusters mostly depends on telencephalic ventricular domains,

where each qRG subtype is positioned (Cosacak et al., 2019). Accordingly, we

examined whether these 4 qRG clusters are linked to different telencephalic ventricular

domains. We focused on 8 telencephalic ventricular regional markers used in Cosacak

et al. (2019): *foxp4* (Vd), *Pou3f1* (Dd, Dm, Dl), *dmrta2* (Dd, Dm), *gsx2* (Dm), *nr2f1b*

(Dd), *foxj1a* (Vd), *zic1* (Vv), and *iqgap2* (pre-neuroblasts/neuroblasts), and examined

their expression in 4 qRG clusters (Fig. 2-figure supplement 6A). *zic1* was specifically

expressed in qRG/NeuroEp2 and qRG2 (Fig. 2-figure supplement 6BC). *gsx2* was

prominently expressed in qRG/NeuroEp2 and qRG2, but weakly expressed in a

relatively small fraction of qRG/NeuroEp1 (Fig. 2-figure supplement 6BC). *foxj1a* was

specifically expressed in qRG/NeuroEp1 (Fig. 2-figure supplement 6BC). *nr2f1b* was

expressed in a small fraction of qRG1, whereas *pou3f1, dmrta2*, and *foxp4* were

expressed in a small fraction of qRG/NeuroEp1 and pRG/Neuroblast/NBN (Fig. 2-

figure supplement 6BC). These data suggest that qRG/NeuroEp1, qRG1 and

pRG/Neuroblast/NBN are expressed in the dorsal ventricular zone (Dm, Dd, and Dl) as

well as Vd, whereas qRG/NeuroEp2 and qRG2 are expressed in Dm and Vv. These data

suggest that qRG/NeuroEp1, qRG1 and pRG/Neuroblast/NBN link each other in terms

of ventricular domains, especially Vd, Dm, Dd, and Dl. This is consistent with our

observation that RG proliferation is significantly increased in the Dd area after TBI (Fig.

1H).

In zebrafish telencephalon, Notch3 and its downstream target, Hey1, are

essential for maintenance of quiescence and stemness of RGs (Than-Trong et al., 2018).

gfap, *her4.1, her6, her9*, *sox2,* and *nestin* are expressed in RGs (Than-Trong and Bally-

Cuif, 2015; Than-Trong et al., 2020). *vimentin* is strongly expressed in RGs of the

dorsal ventricular domain and Vd, but weakly expressed in Vv (Ganz et al., 2010). The

JNK-Jun pathway mediates stress responses after neuronal injury (Coffey, 2014).

Therefore, we examined expression of these genes in 4 qRG and pRG clusters.

Interestingly, these genes are classified into 4 modules (Fig. 2-figure supplement 7A).

notch3, hey1, sox2 and *her6* are expressed in qRG/NeuroEp1, qRG/NeuroEp2, and

qRG2; however, their expression is low in qRG1 and pRG (Fig. 2-figure supplement

7BC), suggesting that stemness and quiescence start to be lost in qRG1. *vimentin, nestin,*

her4.1 are expressed very little in qRG/NeuroEp2 and qRG2, whereas they are highly

expressed in qRG/NeuroEp1, qRG1, pRG (Fig. 2-figure supplement 7BC), suggesting

that regenerative responses start in qRG/NeuroEp1, qRG1 and pRG. Furthermore, a low

level of *vimentin* expression is consistent with our regional marker analysis, indicating

that qRG/NeuroEp2 and qRG2 have a more ventral subpallium character (Fig. 2-figure

supplement 6). Interestingly, *jun* is expressed in qRG/NeuroEp1, qRG/NeuroEp2, pRG,

but little expressed in qRG1 and qRG2 (Fig. 2-figure supplement 7BC), suggesting that

qRG1 is not exposed to a stress response.

308   Next, we examined pRG markers in 4 qRG clusters. Here we focused on 47

genes that were reported to be specifically expressed in ccnd1-positive pRGs (Lange et

al., 2020). Among these 47 pRG genes, expression of eight genes (*si:ch211-193l2.3,*

si:ch211-193l2.4, si:ch211-193l2.5, si:ch211-193l2.6, si:ch211156b7.4, prrc2b, tcp1,

and *h3f3a*) was not detected for pRG analysis. Thus, we investigated expression of the

remaining 39 pRG-specific genes in 4 qRG clusters and pRG (Fig. 2D). These 39 genes

are classified into three groups, Gene modules 1-3, depending on their expression

profile in qRG clusters (Fig. 2-figure supplement 8A). Gene module 1 contains 27

genes that are highly expressed in pRG, but little expressed in all qRG clusters (Fig. 2-

figure supplement 8BC). Gene module 2 contains 5 genes (*mibp2, rps23, rpl22l2,*

nap1l1, and *rsl1d1*) that are highly expressed in both pRG and qRG1 (Fig. 2-figure

supplement 8D). Nap1l1 is a nucleosome assembly protein 1-like 1 and promotes cell

proliferation of neuronal progenitors and neurogenesis (Qiao et al., 2018). Rsl1d1 is a

ribosomal L1 domain-containing 1 that directly interacts with Mdm2 to promote tp53

degradation (Ding et al., 2021). In addition, increased expression of ribosomal proteins

such as *rps23* and *rpl22l2* indicates energy metabolism in qRG1. Thus, these genes may

promote cell proliferation and survival in qRG1. Gene module 3 contains 7 genes (*actl6,*

ccnd2a, sox4a, arrdc3b, kdm6ba, ascl1b, and *boc*) that are highly expressed in pRG,

qRG/NeuroEp1, and qRG/NeuroEp2 (Fig. 2-figure supplement 8E). In zebrafish,

Kdm6ba is a histone demethylase JMJB3 and is essential for tissue regeneration

(Stewart et al., 2009). ascl1 is a master regulator of neuronal regeneration

(Ramachandran et al., 2011). So, elevated expression of *kdm6ba* and *ascl1b* may
indicate that a regenerative response starts to be activated in qRG/NeuroEp1.

Each RG fraction is dynamically changed in response to TBI
We compared the fraction of each cluster in non-injured controls and 3-dpl conditions
(Fig. 2E). First, the fraction of the pRG/Neuroblast/NBN cluster was increased in 3-dpl
conditions (7.7%), compared with non-injured controls (4.4%). Accordingly, the
fraction of MN/NBN was also increased in in 3-dpl conditions (1.2%), compared with
non-injured controls (0.7%). This is consistent with our observation that a higher
number of RGs enter cell proliferation to regenerate neurons in response to TBI at 3 dpl
(Fig. 1E). Another drastic increase was detected in the fraction of qRG1 in 3-dpl
conditions (27.0%), compared with that of non-injured controls (12.0%) (Fig. 2E). On
the other hand, the fraction of qRG/Ep2 was decreased in 3-dpl conditions (12.0%),
compared with non-injured controls (25.0%). The fraction of qRG2 was also decreased
in 3-dpl conditions (3.6%), compared with non-injured controls (8.8%). Since qRG1
seems to be spatially separated from qRG2 and qRG/NeuroEp2 (Fig. 2-figure
supplement 6), the reduction of the qRG2 and qRG/NeuroEp2 fractions may result from
the increase of the qRG1 and pRG/Neuroblast/NBN fractions. The qRG/NeuroEP1
fraction was not changed between 3-dpl conditions (40.0%) and non-injured controls
(38.0%). In summary, in response to TBI, qRG2 and qRG/Ep2 were decreased, whereas
qRG1, pRG/Neuroblast/NBN, MN/NBN were increased.

qRG1 is a ribosomal protein-enriched RG that increased in response to TBI
Since about 70% of the TOP100 differentially expressed genes of qRG1 are ribosomal
proteins (Supplemental Table 3), qRG1 shows a high expression of ribosomal proteins,
which is shared with pRG, but not with other qRG populations (Fig. 2-figure
supplement 8D). In addition, qRG1 does not express Notch3, but expresses *her4.1*, and
vimentin (Fig.2-figure supplement 7). Since Notch3 is required to maintain quiescence
and stemness of NSCs (Alunni et al., 2013; Than-Trong et al., 2020), these data suggest
that qRG1 is in transition from a dormant qRG state. Furthermore, previous scRNA-seq
analyses using adult mouse brain revealed that dormant NSCs enter a primed-quiescent
state before activation, which involves downregulation of Notch signaling and

concomitant upregulation of ribosomal protein expression, leading to elevated protein
synthesis (Basak et al., 2018; Dulken et al., 2017; Llorens-Bobadilla et al., 2015). Thus,
these primed quiescent NSCs in mice have a similar transcriptomic profile to qRG1 in
zebrafish. Since the qRG1 fraction increases in response to TBI, it is expected that
protein translation by upregulated ribosomal genesis is elevated in TBI.

To confirm this possibility, we examined whether the mammalian target of
rapamycin (mTOR) signaling is activated in telencephalon in response to TBI. mTOR
signaling is important for stem cell self-renewal and differentiation in mammalian brain
development (Meng et al., 2018; Saxton and Sabatini, 2017). During activation of
mTOR signaling, mammalian TOR complex 1 (mTORC1) activates ribosomal protein
S6 kinase (S6K) (Magnuson et al., 2012), which subsequently induces phosphorylation
of ribosomal protein S6 (rpS6), a component of the S40 ribosome (Ruvinsky and
Meyuhas, 2006). Furthermore, rpS6 phosphorylation by S6K is required for
regeneration of zebrafish fin (Hirose et al., 2014), heart (Miklas et al., 2022), and retinal
pigment epithelium (Lu et al., 2022). Thus, we conducted labeling of zebrafish
telencephalon at 3 dpl with anti-phospho-rpS6 antibody. Phosphorylation of rpS6 was
increased in the dorsal ventricular zone (Dm, Dd, and Dl) of lesioned hemispheres,
compared with non-injured control hemispheres (Fig. 3A). The fraction of p-rpS6$^+$ area
in GFAP$^+$ area is significantly higher in lesioned hemispheres than in intact non-injured
hemispheres (Fig. 3B), suggesting that the mTOR-activated population of RGs is
increased by TBI. These data suggest that mTOR signaling is activated in response to
TBI in zebrafish telencephalon.

Next, we examined whether qRG1 cells increase in response to TBI in zebrafish
telencephalon *in vivo*. One of the qRG markers, *lgals2a*, is strongly expressed in qRG1
and moderately expressed in qRG2, but very low in qRG/Ep1 and qRG/Ep2 (Fig. 2D
and Fig.2-figure supplement 5AE). *lgals2a* expression was upregulated in the dorsal
ventricular zone (Dd) of lesioned hemispheres, compared with non-injured control
hemispheres, in which *lgals2a* expression is very low (Fig. 3CD). It was reported that
gstm.1 is expressed in primed quiescent neural stem cells in mice (Basak et al., 2018).
scRNA-seq analysis revealed that *gstm.1* is one of the differentially expressed genes of
qRG1 (Supplementary Table 3). *gstm.1* expression was highly upregulated in RGs of
both lesioned hemispheres and non-injured hemispheres in TBI-induced brain,

compared with non-injured control brains, in which *gstm.1* expression is very low (Fig.

3EF). These data suggest that ribosomal protein enriched qRG1 is increased in the

ventricular zone of the dorsal telencephalon in response to TBI.

Pseudo-time analysis reveals that qRG1 and pRG are generated from

qRG/NeuroEp1

qRG1 in zebrafish and primed quiescent NSCs in mice share upregulation of ribosomal

proteins and downregulation of Notch signaling (Dulken et al., 2017; Llorens-Bobadilla

et al., 2015), suggesting that qRG1 may be in a transient state from dormant qRG

toward pRG. To examine this possibility, we conducted pseudo-time analysis using

Monocle3 (Trapnell et al., 2014). We determined qRG/NeuroEp1 as the starting point

for pseudo-time analysis, because qRG/NeuroEp1 expresses dormant markers of qRG

such as *notch3* (Alunni et al., 2013; Than-Trong et al., 2018) (Fig. 2-figure supplement

7AC) and *mfge8a* (Zhou et al., 2018) (Fig. 2-figure supplement 5AC). In addition, two

qRG markers highly expressed in qRG/NeuroEp1, *si:ch211-251b21.1* and *s1pr1* (Fig.

2-figure supplement 5AC), are downregulated in *Notch3*$^{-/-}$ mutants (Than-Trong et al.,

2018), suggesting that Notch3 promotes expression of these qRG markers in

qRG/NeuroEp1. Our pseudo-time transcriptome analysis revealed that qRG/NeuroEp1

was split into three branches in non-injured controls, which lead to

pRG/Neuroblast/NBN, qRG1, and qRG/NeuroEp2, respectively (Fig. 4A and Fig. 4-

figure supplement 1A). On the other hand, in 3-dpl conditions, qRG/NeuroEp1 was split

into two branches, which lead to pRG/Neuroblast/NBN and qRG1, respectively, and

qRG/NeuroEp2 was not mapped on either pseudo-time branches (Fig. 4A and Fig. 4-

figure supplement 1A). Gene expression dynamics on UMAP (Fig. 4-figure supplement

1B) and a Dot-plotting graph (Fig. 4B) confirmed different gene expression dynamics

between a branch moving toward qRG1 (Branch1) and a branch moving toward

pRG/Neuroblast/NBN (Branch2). These data suggest that qRG1 is not an intermediate

state from qRG/NeuroEp1 into pRG/Neuroblast/NBN, but that qRG1 and pRG are

produced via two separate lineages from qRG/NeuroEp1. However, Monocle3 UMAP

indicates that a small population of pRG/Neuroblast/NBN cells are associated with a

Branch1 trajectory path and spatially segregated with a dense population of

pRG/Neuroblast/NBN cells mapped on a Branch2 trajectory path, suggesting the

possibility that qRG1 cells change into pRGs/Neuroblasts/NBNs (Fig. 4-figure
supplement 2A). To clarify this inconsistency, we extracted both Branch1- and
Branch2-associated pRG/Neuroblast/NBN cells and examined their gene expression
profiles. We found that differentially expressed genes differ completely between
Branch1- and Branch2-associated pRG/Neuroblast/NBN cells. Branch1-associated
pRG/Neuroblast/NBN cells showed a very similar gene expression profile to qRG1,
indicating that Branch1-associated pRG/Neuroblast/NBN cells are qRG1 (Fig. 4-figure
supplement 2BC). These data suggest that qRG/NeuroEp1 is a source of both qRG1 and
pRG/Neuroblast/NBN, and that these two RG types are differentially generated from
qRG/NeuroEp1.

Interestingly, the trajectory path from qRG/NeuroEp1 to qRG/NeuroEp2
disappeared in TBI conditions (Fig. 4A). The qRG1 population is increased by TBI,
whereas qRG/NeuorEp2 is decreased by it (Fig. 2E). In adult zebrafish pallium, 17% of
RGs produce neurons by direct conversion; however, the percentage of RGs undergoing
direct conversion was markedly deceased to 3 % after TBI (Barbosa et al., 2015).
Recently it was reported that high aryl hydrocarbon receptor (AhR) signaling promotes
the direct conversion of a specific subset of RGs into post-mitotic neurons, while low
AhR signaling promotes proliferation of RGs (Di Giaimo et al., 2018). After brain
damage, AhR signaling is promptly suppressed in 1–5 dpl and recovered to a normal
level at 7 dpl in zebrafish, indicating that brain damage transiently inhibits AhR
signaling, leading to decrease of the rate of direct conversion. We found that a major
mediator of AhR signaling in zebrafish, Ahr2, and its transcriptional target, cytochrome
P450 1B1 oxidase (Cyp1b1), are highly expressed in qRG/NeuroEp2 but not in other
qRGs (Fig. 4-figure supplement 3). qRG/NeuroEp2 cells are located in Dm and Vv
(Fig.2-figure supplement 7). These data raise the possibility that Dm-associated
qRG/NeuroEp2 is a major source of direct conversion, which may explain the
disappearance of the trajectory path from qRG/NeuroEp1 to qRG/NeuroEp2 after TBI
as well as decrease of qRG/NeuorEp2 fraction after TBI. Further *in vivo* analysis will be
necessary to clarify this point.

TBI dynamically modifies regulatory gene expression in the transition from
qRG/NeuroEp1 to qRG1 and pRG

To determine how TBI promotes the transition from qRG/NeuroEp1 to qRG1 and pRG,

we determined upregulated and downregulated genes in qRG/NeuroEp1 in the injured

condition, compared with the non-injured control condition (Fig. 5A). First, we selected

genes whose mRNA expression was significantly upregulated in 3-dpl conditions,

compared with non-injured control conditions, and applied gene ontogeny (GO)

analysis (Fig. 5B). These upregulated genes are classified into several regulatory

pathways (Fig. 5-figure supplement 1A), for example, the JAK-STAT pathway (Fig. 5 -

figure supplement 1BC), innate immune response (Fig. 5-figure supplement 1D), and

ribosomal metabolism including translation, ribosomal biogenesis, ribosome assembly

and cell-cycle regulation (Fig. 5-figure supplement 1E).

Next, among those, the TOP30 upregulated genes were selected. We examined

mRNA expression levels of these upregulated genes in each of 4 qRG clusters and the

pRG cluster between injured and non-injured conditions (Fig. 5C). Interesting, the

TOP30 upregulated genes are classified into three groups. The first group has 17 genes,

which are highly expressed in both qRG1 and pRG clusters in the injured condition

(Fig. 5C). Most of these genes are ribosomal genes, suggesting that ribosomal

biogenesis is enhanced in the transition lineage from qRG/NeuroEp1 to both qRG1 and

pRG. In addition, protein folding regulator *ppiaa* and *vimentin* are found in this first

group. The second group has 2 genes, *ift45* and *b2m*, whose mRNA expression is higher

in qRG1 than in pRG (Fig. 5C), suggesting that these two genes may promote the

transition to qRG1. The third group has 11 genes that are highly expressed in pRG, but

not in qRG1 (Fig. 5B). The third group contains regulators for the JAK-STAT pathway

(*socs3b* and *stat1*), protein folding (*hsp70l*), and innate immune response (*cxcl18b*),

suggesting that these regulatory pathways are involved in the transition to pRG. We also

confirmed the mRNA expression profile of these three groups of genes along the

pseudo-time lineage toward qRG1 (branch1) and pRG (branch2) (Fig. 5-figure

supplement 2).

Next, we focused on downregulated genes and selected TOP12 downregulated

genes, whose mRNA expression was significantly downregulated in the injured

condition. We performed GO enrichment analysis on these downregulated genes (Fig.

5-figure supplement 3A). Interestingly, 5 genes (*msmo1, sc5d, hmgcs1, cyp51*, and

insig1) are enzymes involved in cholesterol synthesis (Dave and Peeples, 2021) (Fig. 5-

figure supplement 3BC), and one gene (*lipg*) is involved in lipid metabolism (Yu et al.,

2018) (Fig. 5-figure supplement 3D), suggesting that cholesterol synthesis is suppressed

in qRG/NeuroEp1 in the injured condition. In addition, neurogenic inhibitor *id2b* is also

suppressed in the injured condition (Fig. 5-figure supplement 3E). We also examined

mRNA expression levels of these downregulated genes in each of 4 qRG clusters and

pRG between injured and non-injured conditions (Fig. 5C). These 12 downregulated

genes form one group, whose mRNA expression is very low in both qRG1 and pRG

(Fig. 5C). We also confirmed mRNA expression profiles of these 12 genes along the

pseudo-time lineage toward qRG1 (branch1) and pRG (branch2) (Fig. 5-figure

supplement 4).

In summary, upregulation of ribosomal metabolism regulators, *ppiaa* and

vimentin and downregulation of cholesterol biogenesis genes and *id2b* are likely

associated with the transition from qRG/NeruoEp1 to both qRG1 and pRG (Fig. 5D).

Upregulation of *ift45* and *b2m* may be associated with transition to qRG1, whereas

upregulation of JAK-STAT pathway regulators, *cxcl18b*, and *hsp70l* may be associated

with the transition to pRG (Fig. 5D).

Microglia are activated at 1 dpl in response to TBI

In zebrafish, activation of an inflammatory response is required for a regenerative

response of RGs after TBI (Kyritsis et al., 2012). First, we examined whether microglia

are activated by TBI in zebrafish telencephalon. Microglia show ramified morphologies

in a resting state, whereas they become rod-shape or adopt an amoeboid-like

morphology in a reactive state. Therefore, we examined microglial morphology by

measuring the cell sphericity index, which is defined as the ratio of the surface area of a

sphere of the same volume as the given object to the surface area of the object

(Fernandez-Arjona et al., 2017). We scanned microglia in adult telencephalon using the

transgenic line *Tg[mpeg1.1:EGFP]*, which visualizes microglia (Ranawat and Masai,

2021), and calculated their sphericity indices. Indeed, microglia show a range of

sphericity from 0.2 to 0.7 along the axis of cell shape complexity from high to low (Fig.

6-figure supplement 1A). Next, we examined the spatiotemporal profile of the

sphericity of microglia in the telencephalon where TBI was induced in one hemisphere.

First, we examined the sphericity of microglia located in the Dd area adjacent to the

injured site and compared their sphericity distribution with that of microglia located in

the equivalent Dd area of the opposite non-injured hemisphere (Fig. 6-figure

supplement 1B). The sphericity index for microglia in lesioned areas was shifted to a

higher range than that of microglia in control areas at 1 dpl, but not at 3 dpl (Fig. 6-

figure supplement 1C), suggesting that microglia are activated at 1 dpl. To examine this

more precisely, we focused on microglia with high sphericity, which were defined as

more than 0.6 (Fig. 6-figure supplement 1D). The fraction of microglia located in the

lesioned hemispheres in both lesioned and non-injured control hemispheres was

significantly higher for high microglia sphericity at 1 dpl than in non-injured control

conditions, but not at 3 dpl (Fig. 6-figure supplement 1E). Biased distribution of total

microglia in the lesioned hemispheres was not observed at both 1 dpl and 3 dpl (Fig. 6-

figure supplement 1E). These observations suggest that microglia are activated in

response to TBI at 1 dpl.

Genetic elimination of microglia with the NTR-MTZ system

Next, to determine the role of microglia in TBI-mediated neuronal regeneration, we

eliminated microglia in adult zebrafish telencephalon. We used *Tg[mpeg1.1:NTR-*

eYFP], which expresses nitroreductase (NTR) under the control of the *mpeg1.1*

promoter. NTR-mediated conversion of metronidazole (MTZ) results in cytotoxic

reagent formation, which causes death of NTR-expressing cells (Petrie et al., 2014). We

applied 2.5 mM MTZ within 2 or 3 days to *Tg[mpeg1.1:NTR-eYFP]* transgenic

zebrafish at 4 – 7 mpf and examined whether microglia are eliminated (Fig. 6-figure

supplement 2A). The number of microglia was decreased in telencephalon of

Tg[mpeg1.1:NTR-eYFP] transgenic zebrafish treated with MTZ for 2 days compared

with those with no MTZ treatment, although the difference was not significant

(p=0.0752) (Fig. 6-figure supplement 2B). On the other hand, the number of microglia

was significantly decreased in telencephalon of *Tg[mpeg1.1:NTR-eYFP]* transgenic

zebrafish treated with MTZ for 3 days, compared with those with no MTZ treatment

(Fig. 6-figure supplement 2B). We also examined microglial sphericity in telencephalon

of MTZ-treated *Tg[mpeg1.1:NTR-eYFP]* transgenic zebrafish. Microglial sphericity

was significantly increased in *Tg[mpeg1.1:NTR-eYFP]* transgenic zebrafish treated with

MTZ for 2 days compared with those with no MTZ treatment (Fig. 6-figure supplement

2C), but not in *Tg[mpeg1.1:NTR-eYFP]* transgenic zebrafish treated with MTZ for 3 days (p=0.0592) (Fig. 6-figure supplement 2C). Although the number of microglia was not significantly decreased 2 days after MTZ treatment, microglia are likely to become unhealthy, suggesting that elimination of microglia starts 2 days after MTZ treatment. Since treatment with MTZ at 2.5 mM for 3 days or at 5 mM for 2 days affected telencephalon morphology (data not shown), we decided to apply MTZ at 2.5 mM within 2 days for all MTZ-related experiments.

Microglia elimination suppresses the TBI-induced proliferative response of RGs

To examine the role of microglia in the TBI-induced proliferative response of RGs, we applied MTZ at 2.5 mM for 2 days to *Tg[mpeg1.1:NTR-eYFP]* transgenic zebrafish at 4 – 7 mpf, and then introduced TBI. We applied BrdU for 12 h prior to dissection of their telencephalons, which were fixed with PFA at 1, 3, 7, and 14 dpl (Fig. 6A). Fixed telencephalons were sliced at 100 μm with a vibratome. Slices were labeled with anti-GFAP and anti-BrdU antibodies (Fig. 6B). First, we evaluated TBI-induced RG proliferation in MTZ-treated transgenic telencephalons. Previously, our observations indicated that the number of BrdU$^+$ RGs increased significantly at 3 dpl telencephalons compared with non-injured controls (Fig. 1E and 1F). However, in MTZ-treated injured telencephalons of *Tg[mpeg1.1:NTR-eYFP]* transgenic fish, the number of BrdU$^+$ RGs increased slightly at 3 dpl, but not significantly compared to non-injured controls of MTZ-untreated, non-transgenic telencephalons (Fig. 6C). Next, we compared the number of BrdU$^+$ RGs between lesioned and non-injured hemispheres at 3 dpl. Previously, our observations indicated that the number of BrdU$^+$ RGs is significantly higher in lesioned hemispheres than in non-injured hemispheres in telencephalons (Fig. 1G). On the other hand, the number of BrdU$^+$ RGs in lesioned hemispheres of MTZ-treated *Tg[mpeg1.1:NTR-eYFP]* transgenic telencephalons was decreased compared with that of lesioned hemispheres in MTZ-untreated, non-transgenic telencephalons or in MTZ-treated, non-transgenic telencephalons (Fig. 6D); however, the number was still significantly higher than in a non-injured hemispheres of MTZ-treated *Tg[mpeg1.1:NTR-eYFP]* transgenic telencephalons (Fig. 6D). Previously, our observations indicated that RGs in the Dd area more efficiently responded to TBI and incorporated BrdU (Fig. 1H). A similar significant increase in the number of BrdU+

RGs was observed in the Dd area of MTZ-treated *Tg[mpeg1.1:NTR-eYFP]* transgenic

telencephalons compared with non-injured controls of MTZ-untreated, non-transgenic

telencephalons (Fig. 6E). However, the number of BrdU+ RGs in the Dd area was

significantly   decreased   in   MTZ-treated   *Tg[mpeg1.1:NTR-eYFP]*   transgenic

telencephalons, compared with that of MTZ-treated, non-transgenic telencephalons

(Fig. 6-figure supplement 3). These observations suggest that microglial elimination

affects the proliferative response of RGs in response to TBI, although a local response

was still retained to some degree.

The fractions of qRG1 and pRG decrease in response to TBI in microglia-depleted

telencephalons

We applied scRNA-seq analysis to zebrafish MTZ-treated *Tg[mpeg1.1:NTR-eYFP;*

gfap:dTomato] transgenic telencephalons with TBI treatment. Similar to scRNA-seq

analysis of TBI-induced and non-injured control telencephalons (Fig. 2A), stab injuries

were introduced in telencephalons of 4 mpf adult zebrafish. At 3 dpl, telencephalons

were homogenized to prepare cell suspensions, which were subjected to flow-

cytometry-assisted cell sorting to collect the *gfap:dTomato$^+$* RG fraction (Fig. 2-figure

supplement 1BC). We used a 10x genomics system to prepare scRNA-seq libraries.

After read sequencing and mapping to the zebrafish genomic database GRCz11, we

evaluated nUMI, nGene, and mitochondrial genes for two datasets, non-injured and 3-

dpl MTZ-treated *Tg[mpeg1.1:NTR-eYFP]* brains, to ensure that all our data were

comparable (Fig. 2-figure supplement 1E). We analyzed 992 cells from 3-dpl MTZ-

treated *Tg[mpeg1.1:NTR-eYFP]* brains following the thresholding for unique gene

counts and mitochondrial gene expression and obtained scRNA-seq data on RGs of 3-

dpl MTZ-treated *Tg[mpeg1.1:NTR-eYFP]* brains, which we called "3dpl+MTZ brains"

or "3dpl microglia-depleted brain" hereafter (supplementary table 4). We conducted

unbiased clustering of the dataset from 3-dpl+MTZ sample, using the Seurat package

(Butler et al., 2018). In 3-dpl+MTZ sample, cells were classified into 11 clusters (Fig.

7-figure supplement 1A), which correspond to 4 types of qRGs, pRG/Neuroblast/NBN,

3 types of MN/NBN, ependymal cells, pineal gland, microglia (MG)/macrophage (Fig.

7-figure supplement 1B, and Supplementary Table 4), confirming sufficient quality to

apply further clustering analysis of merged datasets. Then, we used the Harmony

617 merging algorithm to integrate two datasets: scRNA-seq data of RGs between 3-dpl

618 and 3dpl+MTZ brains, the former of which is shown in Fig. 2. After clustering analysis

619 of merged scRNA-seq data, we identified 11 clusters, including 6 qRG populations,

620 pRGs/Neuroblasts/NBNs, MNs/NBNs, ependymal cells, pineal gland cells,

621 microglia/macrophages (Fig. 7A, and Supplementary Table 5). We determined the

622 TOP10 differentially expressed genes for each cluster (Fig. 7-figure supplement 2) and

623 found that 6 qRG clusters contain qRG/NeuroEp1, qRG/NeuroEp2, qRG1 and RG2,

624 which are identified by RNA-seq analysis on merged data of 3-dpl and non-injured

625 controls (Fig. 2), and two new RG clusters, namely, qRG3 and qRG4 (Fig. 7-figure

626 supplement 2). Interestingly, in their TOP10 genes, qRG3 and qRG4 express *cxcl18b*

627 and *id2b*, respectively, which are upregulated and downregulated in qRG/NeuroEp1

628 under 3-dpl conditions (Fig. 5C, Fig. 5-figure supplement 1A, and Fig. 5-figure

629 supplement 3A). To clarify the origin of qRG3 and qRG4, we mapped 3-dpl

630 populations of qRG3 and qRG4 to UMAP of 3-dpl scRNA-seq data (Fig. 7-figure

631 supplement 3AB). We found that qRG3 corresponds to a part of qRG/NeuroEP1 in 3-

632 dpl UMAP, whereas qRG4 corresponds to a part of qRG/NeuroEP1 and OPC in 3-dpl

633 UMAP (Fig. 7-figure supplement 3C). Thus, four qRG populations, qRG/NeuroEP1,

634 qRG/NeuroEP2, qRG1 and qRG2, remain to be clustered in merged data of 3-dpl and

635 3dpl+MTZ brain, although small fractions of qRG/NeuroEP1 form two new clusters,

636 qRG3 and qRG4. We also confirmed how each of the 11 clusters of the merged datasets

637 corresponded to 11 clusters of 3 dpl+MTZ cells (Fig. 7-figure supplement 1D).

638   Next, we examined 46 qRG markers in 6 qRG clusters in merged data of 3-dpl

639 and 3dpl+MTZ brain (Fig. 7B). 46 qRG markers were split into four gene modules,

640 "Microglia-depleted (MD)-gene modules1–4". MD-gene module 1 contains 20 genes

641 that are highly expressed in qRG2. MD-gene module 1 contains 10 qRG markers of

642 "Gene module 4" and 8 qRG markers of "Gene module 3" in non-injured controls and

643 the 3-dpl merged dataset (Fig. 7B). "Gene module 3" genes are highly expressed in

644 qRG1; however, "Gene module 3"-derived genes except *lgals2a* of MD-gene module 1

645 seem to be downregulated in qRG1. DM-gene module 2 contains 13 genes that are

646 highly expressed in qRG/NeuroEp1, qRG3, and qRG4. MD-gene module 2 contains all

647 6 "Gene module 1" genes in non-injured controls and the 3-dpl merged dataset (Fig. 7B).

648 Since qRG3 and qRG4 are thought to be derived from qRG/NeuroEp1, it is reasonable

that DM-gene module 2 genes are highly expressed in qRG3 and qRG4. DM-gene

module 3 contains 12 genes that are highly expressed in qRG/NeuroEp1,

qRG/NeruoEp2, qRG3 and qRG4. All 12 genes of DM-gene module 3 are "Gene

module 2" genes in non-injured controls and the 3-dpl merged dataset (Fig. 7B),

suggesting that DM-gene module 3 is equivalent to "Gene module 2". DM-gene module

4 contains only one gene, *itm2ba*, which is highly expressed in all 6 qRG populations.

itm2ba belongs to "Gene module 4" of non-injured controls and the 3-dpl merged

dataset (Fig. 7). Thus, "Gene module 3" genes, which are highly expressed in qRG1, are

split into MD-gene modules 1–2, whereas "Gene module 1" genes, which are highly

expressed in qRG/NeuroEp1, are merged with "Gene module 4" genes to form DM-

gene module 2. Thus, microglia elimination rearranges the state of qRG/NeuroEp1 and

qRG1.

Next, we examined pRG markers. pRG markers were split into 4 gene modules,

DM-modules 1–4 (Fig. 7-figure supplement 4). DM-gene module 1 contains 11 genes

that are highly expressed in pRG/Neuroblast/NMN and MN/NBN, but show low

expression in all 6 qRG clusters. DM-gene module 2 contains 6 genes that are highly

expressed in pRG/Neuroblast/NMN and MN/NBN, as well as qRG1 and qRG2. DM-

gene module 3 contains 16 genes that are highly expressed only in

pRG/Neuroblast/NMN. Most genes in DM-gene modules 1–3 overlapped with pRG

markers of "Gene module 1" in non-injured controls and the 3-dpl merged dataset (Fig.

7-figure supplement 4). DM-gene module 4 contains 6 genes that are highly expressed

in pRG/Neuroblast/NMN as well as qRG/NeuroEp1, qRG/NeuroEp2, qRG3 and qRG4.

Five of these 6 genes overlap with "Gene module 3" genes in merged data of non-

injured and 3-dpl brains (Fig. 7-figure supplement 4). Thus, "Gene module 2" genes,

which are highly expressed in qRG1, are split into DM-gene modules 1–3.

Next, we compared the fraction of each cluster between 3-dpl and 3dpl+MTZ

conditions (Fig. 7C). Interestingly, the fraction of qRG1 was markedly decreased in

3dpl+MTZ conditions (10.0%), compared with 3-dpl (28.0%). Thus, qRG1 increases in

response to TBI in a microglia-dependent manner. In addition, the fraction of

qRG/NeuroEP2 and MN/NBN clusters were increased in 3dpl+MTZ conditions,

compared with 3-dpl.

To identify what kind of signaling pathway is a potential target of microglia

during the TBI-mediated transition from qRG/NeuroEp1 into either qRG1 or pRG, we

determined upregulated and downregulated genes in qRG/NeuroEp1 of 3dpl+MTZ

brain relative to 3-dpl brain (Fig. 8A). GO analysis revealed that genes related to

ribosomal metabolism (*rps* and *rpl* genes), chaperone cofactor-mediated protein folding

(*hsp70l, hsp70.1, hsp70.4, hsp90aa1.2*), response to cytokine and JAK-STAT pathway

(*cxcl12a, stat2, socs3a*), Jun (*jun, junba, junbb, jund*), Hes (*her4.3, her4.4, her6*) are

downregulated in 3-dpl+MTZ brain, compared with 3-dpl brain (Fig. 8B, Fig. 8-figure

supplement 1A). Downregulation of mRNA expression in qRG/NeuroEp1 was

confirmed on UMAP of 3-dpl and 3dpl+MTZ for the JAK-STAT pathway (Fig. 8-

figure supplement 1BC), for chaperone cofactor-mediated protein folding (Fig. 8-figure

supplement 1D), for ribosomal metabolism (Fig. 8-figure supplement 1E), for Hes (Fig.

8-figure supplement 1F), and for Jun (Fig. 8-figure supplement 1G). Furthermore, GO

analysis revealed that four genes related to wound response and sensory epithelial

regeneration, *hbegfa, mdka, hmgb1a*, and *lgals2a* are upregulated in qRG/NeuroEp1

(Fig. 8C, Fig. 8-figure supplement 2A). Upregulation of mRNA expression in

qRG/NeuroEp1 was confirmed in UMAP of 3-dpl and 3 dpl+MTZ (Fig. 8-figure

supplement 2BC) and dot-plot histogram (Fig. 8B).

We also confirm downregulation of these genes in dot-plot histogram (Fig. 8D).

These data suggest that microglia normally promote mRNA upregulation of genes

related to ribosomal genesis, *vimentin*, the JAK-STAT pathway, and *hsp70l* in response

to TBI (Fig. 8E). Upregulation of *cxcl18b, ppiaa, ift45,* and *b2m* mRNA and

downregulation of genes related to cholesterol biogenesis are induced by TBI,

independent of or upstream of microglia-mediated expansion of qRG1 and pRG. This

may be consistent with the observation that a new EP1-derived cluster, qRG3, expresses

cxcl18b (Fig. 7-figure supplement 2). In addition, a new EP1-derived cluster, qRG4,

expresses *id2b* (Fig. 7-figure supplement 2), it is likely that microglia elimination

affects *id2b* downregulation in qRG/NeuroEp1 or its derived qRG4 in response to TBI

(Fig. 8E). It is possible that microglia act on these clusters to change their states into

qRG1 and pRG during the TBI-mediated regenerative response of qRG.

Discussion

In this study, we analyzed scRNA of RGs in adult zebrafish telencephalon in response to TBI. In the merged data set of non-injured control and 3-dpl conditions, we identified 4 qRG clusters, qRG/NeuroEp1, qRG/NeuroEp2, qRG1 and qRG2. Previous scRNA analysis of RGs in zebrafish adult telencephalon revealed 47 qRG-specific genes (Lange et al., 2020). Interestingly, our 4 identified qRG clusters express different combinations of these qRG-specific genes, indicating heterogeneity of qRGs in zebrafish. Our regional marker analysis revealed that these qRG clusters are segregated into two groups, which are localized in dorsal and ventral telencephalic domains. *vimentin* and *zic1* are highly expressed in qRG of dorsal (pallium) and ventral (subpallium) domains of telencephalon, respectively. qRG1 and qRG/NeuroEp1 are likely to be pallial qRGs, because of *zic1*-negative and *vimentin*-positive clusters. On the other hand, qRG2 and qRG/NeuroEp2, seem to be subpallial qRGs, because of *zic1*-positive and *vimentin*-negative clusters. Thus, it is likely that qRG1 and qRG/NeuroEp1 are spatially segregated to qRG2 and qRG/NeuroEp2.

Among 4 qRG clusters, qRG/NeuroEp1 seems to be a dormant qRG, because qRG/NeuroEp1 prominently expresses *Notch3* and *mfge8*, which are required for stemness and quiescence, respectively (Than-Trong et al., 2018; Zhou et al., 2018). qRG1 is a ribosomal gene-enriched cluster and markedly increased in cell number in response to TBI. Consistently, qRGs expressing a mTOR marker, p-rpS6, are significantly increased in dorsal pallium of injured hemispheres, compared with non-injured control hemispheres. Pseudo-time analysis revealed that qRG/NeuroEp1 produces qRG1 and pRG via independent lineages. This is in contrast to primed quiescent NSCs in the V/SVZ of adult mouse brain, which shows a similarly ribosomal gene-enriched population and represents the intermediate state from dormant NSCs to active proliferative NSCs (Dulken et al., 2017; Llorens-Bobadilla et al., 2015).

What is the physiological role of qRG1 in the regenerative response of RGs after TBI? One of differentially expressed genes in qRG1 is *lgals2*, which encodes Galectin-2. Galectins are β-galactose-binding animal lectins (Nio-Kobayashi, 2017). Galectin-1 is expressed in V/SVZ astrocytes of adult mouse brain and promotes proliferation of NSCs in a carbohydrate-binding manner (Sakaguchi et al., 2006). Galectin-1 also promotes proliferation of neural progenitor cells in the SGZ of the dentate gyrus of

adult mouse brain (Kajitani et al., 2009). Galectin-1 and 3 are required for astrocyte

reactivity after adult mouse brain injury (Sirko et al., 2015). In animal models of

experimental autoimmune encephalomyelitis (EAE), Galectin-1 is secreted from

astrocytes, binds to M1 microglia, and negatively regulates M1 microglia-mediated

neurotoxicity, leading to promotion of M2 microglia-mediated neuroprotection

(Starossom et al., 2012). Thus, Galectins regulate astrocyte or NSC behaviors through

modulation of the immune system. In zebrafish retina, Galectin-2 expression is rapidly

induced in retinal NSCs called Müller glia after light-induced damage, and its

knockdown affected the rod regenerative response (Craig et al., 2010), suggesting that

upregulation of *lgals2* expression is the first regenerative response of retinal NSCs in

zebrafish. Indeed, our data showed that *lgals2*-expressing cells are induced only in the

ventricular region near the injury site of the brain, and that the number of *lgals2*-

expressing cells correlates with the size of injury (Fig. 3C). A possible scenario is that

damaged qRG/NeuroEp1 cells exit the dormant state and become qRG1, which

subsequently expresses Galectin-2. Galectin-2 secreted from qRG1 may act on

microglia to migrate into the damaged area for clearance of dead and damaged cells.

Galectin-2 may promote surrounding qRG/NeuroEp1 cells to differentiate into pRGs in

concert with microglia. If this is the case, qRG1 may function as a sensor to detect

damage levels and promote the transition from dormant qRGs to pRGs. It will be

interesting to examine whether the TBI-induced regenerative response of qRGs is

affected in adult zebrafish *lgals2* knockdown telencephalon.

764   Another important question is what kinds of regulatory factors are involved in

production of qRG1 and pRG by qRG/NeuroEp1. We examined up- and down-

regulated genes in qRG/NeuroEp1 at 3 dpl versus non-injured controls. GO analysis

revealed that four signaling pathways, ribosomal metabolism, JAK/STAT pathway,

cytokine signaling including *cxcl18b*, and chaperonin-mediated protein folding,

including *hsp70l* and *ppiaa*, are upregulated in 3-dpl qRG/NeuroEp1. In addition,

vimentin, *ift45* and *b2m* are upregulated. Interestingly, these upregulated genes are

classified into three groups. The first group (ribosomal genes, *vimentin, ppiaa*) is highly

expressed in both pRGs and qRG1 after injury. The second group (*ift45, b2m*) is highly

expressed in qRG1, but not pRG, and the third group (JAK-STAT, *cxcl18l, hsp70l*) is

highly expressed in pRG, but not qRG1 (Fig. 5C). It is possible that these three group

genes differentially generate both pRG and qRG1, qRG1 or pRG, respectively (Fig. 5D).

Furthermore, three signaling pathways: cholesterol biogenesis, a lipid metabolism factor

lipg, and a neurogenic inhibitor, *id2b*, are downregulated in 3-dpl qRG/NeuroEp1.

Almost all these downregulated genes show low expression in both qRG1 and pRG at 3

dpl, suggesting that their downregulation promotes the transition from qRG/NeuroEp1

to both qRG1 and pRG (Fig. 5D). At present, it is unknown how ribosomal metabolic

genes are upregulated in qRG/NeuroEp1 by TBI. It was reported that *mfge8* is highly

expressed in quiescent NSCs in the SGZ of adult mouse dentate gyrus, and that Mfge8

promotes NSC quiescence by suppressing the mTOR pathway in an autocrine manner

(Zhou et al., 2018). Since mfge8 is highly expressed in qRG/NeuroEp1, but

downregulated along the lineage toward qRG1 and pRG (Fig. 4B), downregulation of

mfge8 may activate the AKT/mTOR pathway to increase ribosomal genesis in both

qRG1 and pRG.

Another interesting finding is the downregulation of cholesterol synthesis genes.

ER membrane-localized cholesterol sensor protein (*insig1*) and four components of the

cholesterol synthesis pathway (*hmgcs1, cyp51, msmo1, sc5d*) are downregulated in

qRG/NeuroEp1 after injury. Furthermore, endothelial lipase (*lipg*), which promotes

lipid biosynthesis through cleavage of high-density-lipoprotein (HDL) (Yu et al., 2018),

is downregulated in qRG/NeuroEp1 after injury. Brain cholesterol metabolism is tightly

regulated through cooperation between neurons and different glial-cell types (Dai et al.,

2021; Dave and Peeples, 2021). Since peripheral cholesterol cannot enter the brain

across the blood-brain-barrier, cholesterol in adult brain is synthesized *de novo* in glial

cells, primarily astrocytes. Cholesterol is loaded onto apolipoprotein E (ApoE) in

astrocytes and secreted through ATP binding cassette (ABC) transporters. Neurons and

oligodendrocytes take up lipidated ApoE particles via receptor-mediated endocytosis, so

they depend on astrocytes for cholesterol. However, in chronic demyelinated lesions,

top-downregulated genes in astrocytes are involved in cholesterol synthesis (Itoh et al.,

2018), suggesting that reduction of cholesterol biogenesis is an initial response of

astrocytes after brain damage. At present, its physiological significance is unknown;

however, it was recently reported that the stem cell marker Prominin1 promotes axon

regeneration by downregulating cholesterol synthesis via Smad signaling (Lee et al.,

2020). Thus, cholesterol biogenesis was implicated in axonal regeneration. It will be

interesting to investigate how cholesterol regulates regenerative responses of RGs after
brain injury.

Previous studies revealed that acute inflammation is required for the
regenerative response of RGs to TBI in zebrafish (Kyritsis et al., 2012). We confirmed
that microglial depletion compromises the regenerative response of RGs to TBI. Next,
scRNA-seq analysis was applied to 3-dpl zebrafish telencephalon under microglia-
depleted conditions. In the merged dataset of GFAP:dTomato$^+$ cells in 3-dpl zebrafish
telencephalon with and without microglia, 11 clusters were identified. Among them,
qRGs are classified into 6 clusters. qRG/NeuroEp1, qRG/NeuroEp2, qRG1, qRG2,
qRG3 and qRG4. qRG3 and qRG4 are a part of qRG/NeuroEp1 in 3-dpl conditions,
suggesting that these clusters are derived from qRG/NeuroEp1. Interestingly, the
fraction of qRG1 is markedly decreased in 3-dpl microglia-depleted conditions,
compared with 3-dpl microglia-intact conditions, suggesting that expansion of qRG1 by
brain injury depends on microglia. To determine what kinds of signaling networks are
targeted by microglia during brain damage, upregulated and downregulated genes in
qRG/NeuroEp1 were compared between 3-dpl microglia-depleted conditions and
microglia-intact conditions. GO analysis revealed that downregulated genes pertain to
ribosomal metabolism, chaperone-mediated protein folding, cytokine and JAK-STAT
signaling, jun, and her4/6. These data suggest that upregulation of ribosomal
metabolism, chaperone-mediated protein folding, cytokine and JAK-STAT signaling in
qRG/NeuroEp1 after TBI depend on microglia. Only four genes in qRG/NeuroEp1 are
upregulated: *lgals2a, mdka, hmgb1a,* and *hbegfa*. Although the qRG1 fraction is
markedly decreased, the qRG1-specific marker *lgals2a* is highly expressed in
qRG/NeuroEp1. So, TBI may induce *lgals2a* transcription even in microglia-depleted
conditions; however, *lgals2a*-expressing cells fail to differentiate into qRG1, because
ribosomal genes and vimentin fail to be upregulated in qRG/NeuroEp1 after TBI.
cxcl18b expression in qRG/NeuroEp1 was not downregulated in 3-dpl microglia-
depleted conditions and *cxcl18b* is highly expressed in qRG3, which is likely to derive
from qRG/NeuroEp1. These observations suggest that *cxcl18b* upregulation of 3-dpl
qRG/NeuroEp1 occurs prior to microglial action. Similarly, expression of cholesterol
biogenesis genes was not upregulated in qRG/NeuroEp1 under microglia-depleted
conditions, suggesting that downregulation of cholesterol biogenesis gene expression

occurs prior to microglial action. Taken together, these data illustrate that modulation of

gene expression for cholesterol biogenesis, *ppiaa, cxcl18b, id2b, ift45*, and *b2m* occurs

prior to the interaction between microglia and qRG/NeuroEp1, whereas expression of

genes for ribosomal metabolism, vimentin, the JAK/STAT pathway, *hsp70l*, *junba/bb*,

and *her4/6* are modulated by the interaction between microglia and qRG/NeuroEp1 (Fig.

8E). In the future, further studies are necessary to identify the mechanism underlying

the interaction between microglia and dormant RGs during TBI.

Materials and Methods

Fish Strains

Zebrafish (*Danio rerio*) were maintained using standard procedures (Westerfield, 2000).

Adult zebrafish at 3-7 months old were used for all experiments. Female and male fish

were randomly selected for analysis. *okinawa wild-type* (*oki*) was used for the wild-type

strain. *Tg[GFAP:GFP]*[mi2001] (Bernardos and Raymond, 2006) and

Tg[GFAP:dTomato][nns17Tg] (Satou et al., 2012) were used to visualize qRG.

Tg[mpeg1.1:EGFP][oki053] (Ranawat and Masai, 2021) was used to visualize microglia.

Tg[mpeg1:NTR-EYFP][w202] (Petrie et al., 2014) was used for depletion of microglia.

Ethics statement

All zebrafish experiments were performed in accordance with the Animal Care and Use

Program of Okinawa Institute of Science and Technology Graduate University (OIST),

Japan, which is based on the Guide for the Care and Use of Laboratory Animals by the

National Research Council of the National Academies. The OIST animal care facility

has been accredited by the Association for Assessment and Accreditation of Laboratory

Animal Care (AAALAC International). All experimental protocols were approved by

the OIST Institutional Animal Care and Use Committee.

Stab lesion assay

Following anesthesia of adult zebrafish with Tricaine/MS222 (200μg/ml), a needle with

a length of 2 mm and an outer diameter of 300 μm was vertically inserted through the

skull into the medial region of the right hemisphere of the telencephalon. After

lesioning, fish were maintained in separate tanks until the times for analysis and were

sacrificed with ice water following Tricaine-mediated anesthesia. Heads were separated

from the bodies at the level of the gills, and fixed with 4% PFA for 2 h at room

temperature or 6 h at 4°C. Afterward, brains were dissected from skulls and fixed again

with 4% PFA overnight at 4°C. After injuring the fish, when mild hemorrhage was

noticed, they were permitted to recover in fresh water, which was daily changed.

Uninjured intact zebrafish were used for controls.

Histology and antibody labeling

Using a vibratome (Dosaka, EM Co. Ltd., Linear Slicer PRO7), 100-μm frontal sections

were prepared along the antero-posterior axis of the adult telencephalon. These

vibratome sections were used for immunolabeling according to a published protocol

(Schmidt et al., 2014). The following primary antibodies were used at an appropriate

dilution rate: anti-BrdU (Abcam, ab6326, 1:400), anti-HuC/D (ThermoFisher, A-

21271, 1:500), rabbit anti-GFP (ThermoFisher, A11122, 1:200), anti-GFAP (zrf1)

(ZIRC, 1:200), anti-phosphorylated rpS6 (Cell signaling, 2211,1:500). Anti-mouse,

anti-rabbit, and anti-rat antibodies conjugated with several Alexafluor dyes (Alexa 488,

Alexa 546, and Alexa 633) were used as secondary antibodies (ThermoFisher, 1:500).

Immunolabeled sections were mounted on glass slides in PBS buffer with 70% glycerol

for confocal LSM scanning. Vibratome brain slices were scanned with A1R confocal

LSM (Nikon). 3D images covering a large area were captured using a 20x/0.75

objective lens with tiling and stitching functions. Sample preparation and

immunolabeling of cryo-sections were carried out according to standard protocols

(Masai et al., 2003). Cryo-sectioned samples were scanned with confocal LSM, A1R

(Nikon).

In situ **hybridization**

cDNA fragments of *gstm.1* and *lgals2a* were amplified by RT-PCR using mixed cDNA

synthesized from the heads of 1-4 dpf embryos as a template and the following primers:

gstm.1 forward, 5'-ATGGCAATGAAATTGGCTTATTG-3'

gstm.1 reverse, 5'-TCACTCCTTCTTGTTTCCCCA-3'

lgals2a forward, 5'-ATGGCCGGTGTGCTTATACAGA-3'

lgals2a reverse, 5'-CTATTTAATTTCAACCCCTTGGAT-3'

Amplified fragments were subcloned into the PCRII-TOPO vector (Invitrogen).

Antisense and sense RNA probes were synthesized from the linearized plasmid template

by *in vitro* transcription using a DIG RNA Labeling Kit (Roche). *In situ* hybridization

was performed on cryo-sections of telencephalon as previously described (Masai et al.,

2003) with a minor modification. Hybridization was performed overnight at 60°C using

lgals2a, *gstm.1* antisense RNA probes at 0.52 ng/μL and 0.31 ng/μL, respectively.

BrdU incorporation, detection, and analysis

Adult fish were incubated in 10 mM BrdU-containing water for 12 h and sacrificed to dissect telencephalons and prepare vibratome- or cryo-sections for further analysis. To increase efficiency of anti-BrdU antibody labeling, vibratome- and cryo-sections were pretreated with 2N HCl for 1 h at room temperature to expose BrdU antigen from nuclear packed chromatin. BrdU-incorporated brain slices were labeled with anti-BrdU and anti-GFAP (zrf1) antibodies and scanned with confocal LSM A1R (Nikon) with large-scan combined tiling and stitching functions. GFAP and BrdU double-positive nuclei in the ventricular zone of the dorsal pallium were visualized by surface rendering using Imaris software (Bitplane) and used for statistical analysis using GraphPad Prism (ver.9).

Sphericity analysis of microglia

Vibratome brain slices of zebrafish transgenic line *Tg[mpeg1.1:EGFP]* were scanned with confocal LSM A1R (Nikon). After confocal scanning, target telencephalic areas were analyzed using Imaris software (Bitplane) to prepare surface rendering objects for each of *mpeg1.1:EGFP+* microglia. For sphericity analysis, the sphericity index was defined as the ratio of the surface area of a sphere of the same volume as a given object to the surface area of the object (Fernandez-Arjona et al., 2017). The sphericity index was calculated using surface area and volume data for each object. Statistical analysis was performed using GraphPad Prism (ver.9).

Surface-Rendering Process

To identify proliferating cells localized in the ventricular zone, the niche of RG, GFAP$^+$ surface was created, followed by creating BrdU$^+$ surfaces. Depending on the threshold used for creating GFAP and BrdU surfaces, a colocalization channel was created. The colocalized surface, which is identical to the BrdU$^+$ surface, but is not located in the parenchyma, was counted as proliferating cells in the ventricular zone. If necessary, threshold adjustment was carried out to decrease the noise level during surface-creation. To evaluate the sphericity of microglia, due to their complicated shapes and their close

association with the lesioned site, a manual threshold was specified across all sections

and samples for surface rendering.

Microglia depletion by NTR-MTZ system

To eliminate microglia from adult zebrafish brains, we used a transgenic line

Tg[mpeg1:NTR-eYFP] that expresses NTR under the control of the *mpeg1.1* promoter

(Petrie et al., 2014). Fish were maintained in water containing 2.5 mM MTZ from 2

days prior to injury until harvesting of brains. To evaluate levels of microglia

elimination, the transgenic line *Tg[mpeg1.1:NTR-eYFP]* was combined with another

transgenic line *Tg[mpeg1.1:EGFP]* (Ranawat and Masai, 2021), which labels microglia

or microglial precursor cells more efficiently. This double transgenic line was used for

sphericity analysis (Fig. 6-figure supplement 1) and for counting microglial numbers

(Fig. 6-figure supplement 2).

Fraction of p-rpS6-positive area in the GFAP-positive Dd ventricular zone

Cryosections of zebrafish adult telencephalons carrying the transgene *Tg[GFAP:EGFP]*

were labeled with anti-p-rpS6 antibody and scanned with confocal LSM. In the

scanning images, the ROI corresponding to the Dd ventricular domain of injured

hemispheres or non-injured control hemispheres was selected. Using Image-J (ver.1.44,

NIH), GFAP:EGFP$^+$ areas were outlined by changing the threshold adjustment to

remove background noise, and then the fraction of the GFAP:EGFP$^+$ area in the ROI

was determined. Similarly, the fraction of the p-rpS6$^+$ area in the same ROI was

determined. Using the area of the ROI, we calculated the area of p-rpS6$^+$ and

GFAP:EGFP$^+$ regions. The fraction of p-rpS6$^+$ in the GFAP:EGFP$^+$ area in the ROI of

each brain section was calculated. Statistical significance was determined using paired

t-tests with GraphPad Prism (ver. 9).

Fraction of *lgals2a-* and *gstm.1-positive* cells in the total hemisphere area

Cryosections of zebrafish adult telencephalon were labeled with *lgasl2a* or *gstm.1* RNA

probe. RNA signals were outlined using Image-J (ver.9) by changing the threshold

adjustment to remove background noise. The fraction of the mRNA-expressing area in

the total area of the injured or non-injured hemisphere was calculated. Statistical

significance was determined using one-way ANOVA with Tukey's multiple comparison

test, using GraphPad Prism (ver. 9).

976   Fluorescence-Activated Cell Sorting of RGs

Telencephalons were dissected in ice-cold Hank's Balanced Salt Solution (HBSS)

(0.137 M NaCl, 5.4 mM KCl, 0.25 mM Na_2HPO_4, 0.1% glucose, 0.44 mM KH_2PO_4, 1.3

mM $CaCl_2$, 1.0 mM $MgSO_4$, 4.2 mM $NaHCO_3$) from non-lesioned or 3-dpl

Tg[gfap:dTomato] fish. For the microglia-ablated system, MTZ-treated

Tg[gfap:dTomato; mpeg1.1:NTR-eYFP] fish were used. Seven telencephalons per

experimental condition were pooled and processed. Isolated telencephalons were

chemically dissociated for 30 - 40 min at room temperature in HBSS containing papain

(20 U/mL) and DNase (2000U/mL) with gentle trituration every 10 min. Cell

suspensions were filtered with a 40-μm cell strainer (Falcon). Then 10% FBS in L-15

medium was added to stop the enzymatic reaction. Dissociated cells were washed with

10% FBS/PBS solution, and the final pellet was resuspended in 10% FBS/PBS for cell

sorting. Before cell sorting, cells were again passed through a 40-μm cell strainer, and

Sytox blue dead cell stain (Invitrogen) was added to label dead cells. FACS sorting was

performed using a BD FACSAria II, and 9,000 – 20,000 $dTomato^+$ cells were collected

in a 10% FBS/PBS precoated tube. Collected cells were concentrated by centrifugation

at 300 x *g* for 5 min and resuspended in 50-60 μL of 10%FBS/PBS.

994   scRNA-seq library preparation

995   Collected $dTomato^+$ cells were loaded onto a 10X Chromium System using 10X v3

996   chemistry. Single-cell RNAseq libraries were prepared with 12 cycles of cDNA

997   amplification and 15 cycles of indexing PCR, and were sequenced with NovaSeq6000

998   (Illumia).

1000   Computational analysis

Sequencing data were analyzed using 10x Genomics Cell Ranger 3.1.0 (Zheng et al.,

2017) to perform alignment, filtering, barcode counting, and UMI counting with default

settings and the zebrafish reference genome (GRCz11). Filtered output matrices were

used for clustering and downstream analysis in the Seurat v4.0 R package (Hao et al.,
2021), where cells with a unique feature count between 200 and 2500 and a fraction of
mitochondrial transcripts less than 5% and genes expressed in at least 3 cells were used
for analysis. Each dataset was normalized using the SCTransform function with
regressing out of mitochondrial gene expression as a possible source of variation.
Control and 3-dpl datasets were merged using the Harmony algorithm (Korsunsky et al.,
2019), followed by principal component analysis. Top 20 principal components were
used for cluster generation with resolution = 0.5. UMAP was used to visualize resulting
clusters in low-dimensional space. To identify each cluster's differentially expressed
genes, the Findmarkers function with the Wilcoxon rank sum test was used.
Differentially expressing genes (DEGs) between controls and 3-dpl datasets in a cluster
were determined using the Findmarker function with DESeq2 test. To identify
upregulated and downregulated biological processes between control and 3-dpl datasets,
gene ontology (GO) enrichment analysis was performed with the list of significantly
upregulated and downregulated genes (adjusted p value < 0.05) using the clusterProfiler
R package (Yu et al., 2012). Trajectory inference was performed using the Monocle3 R
package (Qiu et al., 2017). OPC, Ependymal Cells, Pineal Gland, and MN/NBN clusters
were dropped from control and 3-dpl merged objects were analyzed with Seurat. Then
the object was split into control and 3-dpl. After individual Seurat objects were
converted to monocle3 objects with the as.cell_data_set function, preprocessing using
num_dim=10, dimensionality reduction, and UMAP projection was performed. To learn
the trajectory graph and to order the cells in pseudotime, learn_graph and order_cells
functions were used with default parameters.
1027

Statistical Analysis

Statistical analysis was conducted using GraphPad Prism 9.4.0. For analyzing
proliferative responses in 3-dpl injury conditions (Fig. 1) and 3-dpl+MTZ conditions
(Fig. 6), the same non-injured control sample was used for the comparison. In the case
of analyzing hemisphere-dependent responses, only sections with clear evidence of
lesion sites were included in calculations from each biological replicate. The number of
samples used per experiment is represented in dot-pot histogram and Source Data files.
Tests used to calculate statistical significance are mentioned in figure legends. All data

are presented as means ± SDs. A p-value of 0.05 was considered statistically significant.

No data point was removed based on variability. In histograms, p*< 0.05, p**< 0.01,

p***< 0.001, p****< 0.0001, or ns (not significant) is indicated.

1039

Acknowledgements

We thank Dr. Yuko Nishiwaki, Dr. Tetsuya Harakuni, and Mr. Yutaka Kojima for

supporting experiments. We thank Dr. Randall T. Moon and Dr. Jeanot Muster for

providing the *Tg[mpeg1.1:NTR-eYFP]* line. We thank the OIST Research Support

Division: the Sequencing Section (SQC) for assistance in sequencing, the Instrumental

Analysis Section (IAS) for assistance and support in FACS sorting, and the Scientific

Imaging Section (SIG) for assistance and support in imaging experiments.

1047

1048   **Competing interests**

1049   The authors declare no competing or financial interests.

1050

1051   **Funding**

1052   This work was supported by a grant from the Okinawa Institute of Science and

1053   Technology Graduate University to IM.

1054

1055

References

Adolf, B., Chapouton, P., Lam, C. S., Topp, S., Tannhauser, B., Strahle, U., Gotz, M.
and Bally-Cuif, L. (2006). Conserved and acquired features of adult
neurogenesis in the zebrafish telencephalon. *Dev Biol* **295**, 278-293.
Alunni, A., Krecsmarik, M., Bosco, A., Galant, S., Pan, L., Moens, C. B. and Bally-Cuif,
L. (2013). Notch3 signaling gates cell cycle entry and limits neural stem cell
amplification in the adult pallium. *Development* **140**, 3335-3347.
Barbosa, J. S. and Ninkovic, J. (2016). Adult neural stem cell behavior underlying
constitutive and restorative neurogenesis in zebrafish. *Neurogenesis (Austin)* **3**,
e1148101.
Barbosa, J. S., Sanchez-Gonzalez, R., Di Giaimo, R., Baumgart, E. V., Theis, F. J., Gotz,
M. and Ninkovic, J. (2015). Neurodevelopment. Live imaging of adult neural
stem cell behavior in the intact and injured zebrafish brain. *Science* **348**, 789-
793.
Basak, O., Krieger, T. G., Muraro, M. J., Wiebrands, K., Stange, D. E., Frias-Aldeguer,
J., Rivron, N. C., van de Wetering, M., van Es, J. H., van Oudenaarden, A., et al.
(2018). Troy+ brain stem cells cycle through quiescence and regulate their
number by sensing niche occupancy. *Proc Natl Acad Sci U S A* **115**, E610-E619.
Baumgart, E. V., Barbosa, J. S., Bally-Cuif, L., Gotz, M. and Ninkovic, J. (2012). Stab
wound injury of the zebrafish telencephalon: a model for comparative analysis
of reactive gliosis. *Glia* **60**, 343-357.
Bernardos, R. L. and Raymond, P. A. (2006). GFAP transgenic zebrafish. *Gene Expr*
Patterns **6**, 1007-1013.
Butler, A., Hoffman, P., Smibert, P., Papalexi, E. and Satija, R. (2018). Integrating
single-cell transcriptomic data across different conditions, technologies, and
species. *Nat Biotechnol* **36**, 411-420.
Coffey, E. T. (2014). Nuclear and cytosolic JNK signalling in neurons. *Nat Rev Neurosci*
15, 285-299.
Cope, E. C. and Gould, E. (2019). Adult Neurogenesis, Glia, and the Extracellular
Matrix. *Cell Stem Cell* **24**, 690-705.
Cosacak, M. I., Bhattarai, P., Reinhardt, S., Petzold, A., Dahl, A., Zhang, Y. and Kizil, C.
(2019). Single-Cell Transcriptomics Analyses of Neural Stem Cell Heterogeneity
and Contextual Plasticity in a Zebrafish Brain Model of Amyloid Toxicity. *Cell*
Rep **27**, 1307-1318 e1303.
Craig, S. E., Thummel, R., Ahmed, H., Vasta, G. R., Hyde, D. R. and Hitchcock, P. F.
(2010). The zebrafish galectin Drgal1-l2 is expressed by proliferating Muller glia
and photoreceptor progenitors and regulates the regeneration of rod
photoreceptors. *Invest Ophthalmol Vis Sci* **51**, 3244-3252.
Dai, L., Zou, L., Meng, L., Qiang, G., Yan, M. and Zhang, Z. (2021). Cholesterol
Metabolism in Neurodegenerative Diseases: Molecular Mechanisms and
Therapeutic Targets. *Mol Neurobiol* **58**, 2183-2201.
Dave, A. M. and Peeples, E. S. (2021). Cholesterol metabolism and brain injury in
neonatal encephalopathy. *Pediatr Res* **90**, 37-44.

de Oliveira-Carlos, V., Ganz, J., Hans, S., Kaslin, J. and Brand, M. (2013). Notch
receptor expression in neurogenic regions of the adult zebrafish brain. *PLoS*
One **8**, e73384.
Di Giaimo, R., Durovic, T., Barquin, P., Kociaj, A., Lepko, T., Aschenbroich, S., Breunig,
C. T., Irmler, M., Cernilogar, F. M., Schotta, G., et al. (2018). The Aryl
Hydrocarbon Receptor Pathway Defines the Time Frame for Restorative
Neurogenesis. *Cell Rep* **25**, 3241-3251 e3245.
Ding, L., Zhang, Z., Zhao, C., Chen, L., Chen, Z., Zhang, J., Liu, Y., Nie, Y., He, Y., Liao,
K., et al. (2021). Ribosomal L1 domain-containing protein 1 coordinates with
HDM2 to negatively regulate p53 in human colorectal Cancer cells. *J Exp Clin*
Cancer Res **40**, 245.
Dirian, L., Galant, S., Coolen, M., Chen, W., Bedu, S., Houart, C., Bally-Cuif, L. and
Foucher, I. (2014). Spatial regionalization and heterochrony in the formation of
adult pallial neural stem cells. *Dev Cell* **30**, 123-136.
Dulken, B. W., Leeman, D. S., Boutet, S. C., Hebestreit, K. and Brunet, A. (2017).
Single-Cell Transcriptomic Analysis Defines Heterogeneity and Transcriptional
Dynamics in the Adult Neural Stem Cell Lineage. *Cell Rep* **18**, 777-790.
Ekdahl, C. T., Claasen, J. H., Bonde, S., Kokaia, Z. and Lindvall, O. (2003). Inflammation
is detrimental for neurogenesis in adult brain. *Proc Natl Acad Sci U S A* **100**,
13632-13637.
Fernandez-Arjona, M. D. M., Grondona, J. M., Granados-Duran, P., Fernandez-
Llebrez, P. and Lopez-Avalos, M. D. (2017). Microglia Morphological
Categorization in a Rat Model of Neuroinflammation by Hierarchical Cluster
and Principal Components Analysis. *Front Cell Neurosci* **11**, 235.
Ganz, J., Kaslin, J., Hochmann, S., Freudenreich, D. and Brand, M. (2010).
Heterogeneity and Fgf dependence of adult neural progenitors in the zebrafish
telencephalon. *Glia* **58**, 1345-1363.
Grandel, H. and Brand, M. (2013). Comparative aspects of adult neural stem cell
activity in vertebrates. *Dev Genes Evol* **223**, 131-147.
Grandel, H., Kaslin, J., Ganz, J., Wenzel, I. and Brand, M. (2006). Neural stem cells and
neurogenesis in the adult zebrafish brain: Origin, proliferation dynamics,
migration and cell fate. *Dev Biol* **295**, 263-277.
Hao, Y., Hao, S., Andersen-Nissen, E., Mauck, W. M., 3rd, Zheng, S., Butler, A., Lee, M.
J., Wilk, A. J., Darby, C., Zager, M., et al. (2021). Integrated analysis of
multimodal single-cell data. *Cell* **184**, 3573-3587 e3529.
Hirose, K., Shiomi, T., Hozumi, S. and Kikuchi, Y. (2014). Mechanistic target of
rapamycin complex 1 signaling regulates cell proliferation, cell survival, and
differentiation in regenerating zebrafish fins. *BMC Dev Biol* **14**, 42.
Itoh, N., Itoh, Y., Tassoni, A., Ren, E., Kaito, M., Ohno, A., Ao, Y., Farkhondeh, V.,
Johnsonbaugh, H., Burda, J., et al. (2018). Cell-specific and region-specific
transcriptomics in the multiple sclerosis model: Focus on astrocytes. *Proc Natl*
Acad Sci U S A **115**, E302-E309.

Jassam, Y. N., Izzy, S., Whalen, M., McGavern, D. B. and El Khoury, J. (2017).
Neuroimmunology of Traumatic Brain Injury: Time for a Paradigm Shift. *Neuron*
95, 1246-1265.
Jurisch-Yaksi, N., Yaksi, E. and Kizil, C. (2020). Radial glia in the zebrafish brain:
Functional, structural, and physiological comparison with the mammalian glia.
Glia **68**, 2451-2470.
Kajitani, K., Nomaru, H., Ifuku, M., Yutsudo, N., Dan, Y., Miura, T., Tsuchimoto, D.,
Sakumi, K., Kadoya, T., Horie, H., et al. (2009). Galectin-1 promotes basal and
kainate-induced proliferation of neural progenitors in the dentate gyrus of
adult mouse hippocampus. *Cell Death Differ* **16**, 417-427.
Kishimoto, N., Shimizu, K. and Sawamoto, K. (2012). Neuronal regeneration in a
zebrafish model of adult brain injury. *Dis Model Mech* **5**, 200-209.
Korsunsky, I., Millard, N., Fan, J., Slowikowski, K., Zhang, F., Wei, K., Baglaenko, Y.,
Brenner, M., Loh, P. R. and Raychaudhuri, S. (2019). Fast, sensitive and
accurate integration of single-cell data with Harmony. *Nat Methods* **16**, 1289-
1296.
Kriegstein, A. and Alvarez-Buylla, A. (2009). The glial nature of embryonic and adult
neural stem cells. *Annu Rev Neurosci* **32**, 149-184.
Kroehne, V., Freudenreich, D., Hans, S., Kaslin, J. and Brand, M. (2011). Regeneration
of the adult zebrafish brain from neurogenic radial glia-type progenitors.
Development **138**, 4831-4841.
Kyritsis, N., Kizil, C., Zocher, S., Kroehne, V., Kaslin, J., Freudenreich, D., Iltzsche, A.
and Brand, M. (2012). Acute inflammation initiates the regenerative response
in the adult zebrafish brain. *Science* **338**, 1353-1356.
Lam, C. S., Marz, M. and Strahle, U. (2009). gfap and nestin reporter lines reveal
characteristics of neural progenitors in the adult zebrafish brain. *Dev Dyn* **238**,
475-486.
Lange, C., Rost, F., Machate, A., Reinhardt, S., Lesche, M., Weber, A., Kuscha, V.,
Dahl, A., Rulands, S. and Brand, M. (2020). Single cell sequencing of radial glia
progeny reveals the diversity of newborn neurons in the adult zebrafish brain.
Development **147**.
Lee, J., Shin, J. E., Lee, B., Kim, H., Jeon, Y., Ahn, S. H., Chi, S. W. and Cho, Y. (2020).
The stem cell marker Prom1 promotes axon regeneration by down-regulating
cholesterol synthesis via Smad signaling. *Proc Natl Acad Sci U S A* **117**, 15955-
15966.
Lindsey, B. W. and Tropepe, V. (2006). A comparative framework for understanding
the biological principles of adult neurogenesis. *Prog Neurobiol* **80**, 281-307.
Llorens-Bobadilla, E., Zhao, S., Baser, A., Saiz-Castro, G., Zwadlo, K. and Martin-
Villalba, A. (2015). Single-Cell Transcriptomics Reveals a Population of Dormant
Neural Stem Cells that Become Activated upon Brain Injury. *Cell Stem Cell* **17**,
329-340.
Lu, F., Leach, L. L. and Gross, J. M. (2022). mTOR activity is essential for retinal
pigment epithelium regeneration in zebrafish. *PLoS Genet* **18**, e1009628.

Magnuson, B., Ekim, B. and Fingar, D. C. (2012). Regulation and function of ribosomal
protein S6 kinase (S6K) within mTOR signalling networks. *Biochem J* **441**, 1-21.
Marz, M., Chapouton, P., Diotel, N., Vaillant, C., Hesl, B., Takamiya, M., Lam, C. S.,
Kah, O., Bally-Cuif, L. and Strahle, U. (2010). Heterogeneity in progenitor cell
subtypes in the ventricular zone of the zebrafish adult telencephalon. *Glia* **58**,
870-888.
Masai, I., Lele, Z., Yamaguchi, M., Komori, A., Nakata, A., Nishiwaki, Y., Wada, H.,
Tanaka, H., Nojima, Y., Hammerschmidt, M., et al. (2003). N-cadherin
mediates retinal lamination, maintenance of forebrain compartments and
patterning of retinal neurites. *Development* **130**, 2479-2494.
Meng, D., Frank, A. R. and Jewell, J. L. (2018). mTOR signaling in stem and progenitor
cells. *Development* **145**.
Miklas, J. W., Levy, S., Hofsteen, P., Mex, D. I., Clark, E., Muster, J., Robitaille, A. M.,
Sivaram, G., Abell, L., Goodson, J. M., et al. (2022). Amino acid primed mTOR
activity is essential for heart regeneration. *iScience* **25**, 103574.
Monje, M. L., Toda, H. and Palmer, T. D. (2003). Inflammatory blockade restores adult
hippocampal neurogenesis. *Science* **302**, 1760-1765.
Nio-Kobayashi, J. (2017). Tissue- and cell-specific localization of galectins, beta-
galactose-binding animal lectins, and their potential functions in health and
disease. *Anat Sci Int* **92**, 25-36.
Otsuki, L. and Brand, A. H. (2020). Quiescent Neural Stem Cells for Brain Repair and
Regeneration: Lessons from Model Systems. *Trends Neurosci* **43**, 213-226.
Petrie, T. A., Strand, N. S., Yang, C. T., Rabinowitz, J. S. and Moon, R. T. (2014).
Macrophages modulate adult zebrafish tail fin regeneration. *Development* **141**,
2581-2591.
Qiao, H., Li, Y., Feng, C., Duo, S., Ji, F. and Jiao, J. (2018). Nap1l1 Controls Embryonic
Neural Progenitor Cell Proliferation and Differentiation in the Developing Brain.
Cell Rep **22**, 2279-2293.
Qiu, X., Mao, Q., Tang, Y., Wang, L., Chawla, R., Pliner, H. A. and Trapnell, C. (2017).
Reversed graph embedding resolves complex single-cell trajectories. *Nat*
Methods **14**, 979-982.
Ramachandran, R., Zhao, X. F. and Goldman, D. (2011). Ascl1a/Dkk/beta-catenin
signaling pathway is necessary and glycogen synthase kinase-3beta inhibition is
sufficient for zebrafish retina regeneration. *Proc Natl Acad Sci U S A* **108**,
15858-15863.
Ranawat, N. and Masai, I. (2021). Mechanisms underlying microglial colonization of
developing neural retina in zebrafish. *Elife* **10**.
Rothenaigner, I., Krecsmarik, M., Hayes, J. A., Bahn, B., Lepier, A., Fortin, G., Gotz,
M., Jagasia, R. and Bally-Cuif, L. (2011). Clonal analysis by distinct viral vectors
identifies bona fide neural stem cells in the adult zebrafish telencephalon and
characterizes their division properties and fate. *Development* **138**, 1459-1469.
Ruvinsky, I. and Meyuhas, O. (2006). Ribosomal protein S6 phosphorylation: from
protein synthesis to cell size. *Trends Biochem Sci* **31**, 342-348.

Sakaguchi, M., Shingo, T., Shimazaki, T., Okano, H. J., Shiwa, M., Ishibashi, S., Oguro,
H., Ninomiya, M., Kadoya, T., Horie, H., et al. (2006). A carbohydrate-binding
protein, Galectin-1, promotes proliferation of adult neural stem cells. *Proc Natl*
Acad Sci U S A **103**, 7112-7117.
Satou, C., Kimura, Y. and Higashijima, S. (2012). Generation of multiple classes of V0
neurons in zebrafish spinal cord: progenitor heterogeneity and temporal
control of neuronal diversity. *J Neurosci* **32**, 1771-1783.
Saxton, R. A. and Sabatini, D. M. (2017). mTOR Signaling in Growth, Metabolism, and
Disease. *Cell* **168**, 960-976.
Schmidt, R., Beil, T., Strahle, U. and Rastegar, S. (2014). Stab Wound Injury of the
Zebrafish Adult Telencephalon: A Method to Investigate Vertebrate Brain
Neurogenesis and Regeneration. *Jove-J Vis Exp*.
Sirko, S., Irmler, M., Gascon, S., Bek, S., Schneider, S., Dimou, L., Obermann, J., De
Souza Paiva, D., Poirier, F., Beckers, J., et al. (2015). Astrocyte reactivity after
brain injury-: The role of galectins 1 and 3. *Glia* **63**, 2340-2361.
Starossom, S. C., Mascanfroni, I. D., Imitola, J., Cao, L., Raddassi, K., Hernandez, S. F.,
Bassil, R., Croci, D. O., Cerliani, J. P., Delacour, D., et al. (2012). Galectin-1
deactivates classically activated microglia and protects from inflammation-
induced neurodegeneration. *Immunity* **37**, 249-263.
Stewart, S., Tsun, Z. Y. and Izpisua Belmonte, J. C. (2009). A histone demethylase is
necessary for regeneration in zebrafish. *Proc Natl Acad Sci U S A* **106**, 19889-
19894.
Than-Trong, E. and Bally-Cuif, L. (2015). Radial glia and neural progenitors in the adult
zebrafish central nervous system. *Glia* **63**, 1406-1428.
Than-Trong, E., Kiani, B., Dray, N., Ortica, S., Simons, B., Rulands, S., Alunni, A. and
Bally-Cuif, L. (2020). Lineage hierarchies and stochasticity ensure the long-term
maintenance of adult neural stem cells. *Sci Adv* **6**, eaaz5424.
Than-Trong, E., Ortica-Gatti, S., Mella, S., Nepal, C., Alunni, A. and Bally-Cuif, L.
(2018). Neural stem cell quiescence and stemness are molecularly distinct
outputs of the Notch3 signalling cascade in the vertebrate adult brain.
Development **145**.
Trapnell, C., Cacchiarelli, D., Grimsby, J., Pokharel, P., Li, S., Morse, M., Lennon, N. J.,
Livak, K. J., Mikkelsen, T. S. and Rinn, J. L. (2014). The dynamics and regulators
of cell fate decisions are revealed by pseudotemporal ordering of single cells.
Nat Biotechnol **32**, 381-386.
Westerfield, M. (2000). *The zebrafish book. A guide for the laboratory use of zebrafish*
(Danio rerio). (4th ed. edn). Eugene.: Univ. of Oregon Press.
Xiong, Y., Mahmood, A. and Chopp, M. (2013). Animal models of traumatic brain
injury. *Nat Rev Neurosci* **14**, 128-142.
Yu, G. C., Wang, L. G., Han, Y. Y. and He, Q. Y. (2012). clusterProfiler: an R Package for
Comparing Biological Themes Among Gene Clusters. *Omics* **16**, 284-287.
Yu, J. E., Han, S. Y., Wolfson, B. and Zhou, Q. (2018). The role of endothelial lipase in
lipid metabolism, inflammation, and cancer. *Histol Histopathol* **33**, 1-10.

Zhao, C., Deng, W. and Gage, F. H. (2008). Mechanisms and functional implications of
adult neurogenesis. *Cell* **132**, 645-660.
Zheng, G. X., Terry, J. M., Belgrader, P., Ryvkin, P., Bent, Z. W., Wilson, R., Ziraldo, S.
B., Wheeler, T. D., McDermott, G. P., Zhu, J., et al. (2017). Massively parallel
digital transcriptional profiling of single cells. *Nat Commun* **8**, 14049.
Zhou, Y., Bond, A. M., Shade, J. E., Zhu, Y., Davis, C. O., Wang, X., Su, Y., Yoon, K. J.,
Phan, A. T., Chen, W. J., et al. (2018). Autocrine Mfge8 Signaling Prevents
Developmental Exhaustion of the Adult Neural Stem Cell Pool. *Cell Stem Cell* **23**,
444-452 e444.

1279

1280

Figure Legends

Figure 1. Stab injuries activate regenerative responses of RGs in adult zebrafish telencephalon.

(A) Injury application method. Using a needle, stab injury is introduced in the right hemisphere of the dorsal telencephalon (pallium) in adult zebrafish. Image was created using Biorender (https://biorender.com).

(B) Experimental design to investigate temporal profiles of regenerative responses of RGs to TBI. After TBI, BrdU is applied for 12 h prior to analysis at 1, 3, 7, and 14 dpl. Image was created using Biorender (https://biorender.com).

(C) Schematic drawing of zebrafish adult telencephalon. In zebrafish, two distinct neurogenic proliferation zones were formed in ventricular surfaces of the dorsal and ventral telencephalon, respectively. In the dorsal domain, called the pallium, an RG cell body (green circle) is located at the ventricular surface and extends a long radial process, the endfoot of which reaches the pial surface close to a blood vessel (orange circle) at the lateral margin of the dorsal telencephalon. In the ventral domain, called the subpallium, neuroepithelial-like progenitors (red circles) are located at the ventricular zone.

(D) Double labeling of the telencephalon with anti-GFAP (green) and anti-BrdU (magenta) antibodies. Top and bottom panels indicate original fluorescent images and surface-rendered object images, respectively. Only surface-rendered objects located in the ventricular zone (VZ) were used for analysis. Scales: 20μm.

(E) Double labeling of the telencephalon with anti-GFAP (white) and anti-BrdU (magenta) antibodies at 1, 3, 7, and 14 dpl. The left-most panel indicates non-injured telencephalon. Scales: 70μm.

(F) The number of BrdU+ cells in the ventricular zone of both hemispheres. The number is significantly higher at 3 dpl than in non-injured control. There is no significant difference between non-injured controls and other time points. mean±SD. $p* < 0.05$; one way ANOVA Dunnett's multiple comparison.

(G) Comparison of the number of BrdU$^+$ cells between lesioned and intact hemispheres at 3 dpl. The number was significantly higher in lesioned hemispheres. mean±SD. $p** < 0.01$; paired t-test.

(H) The number of BrdU$^+$ cells in different ventricular domains of the telencephalon: Dm, Dd, Dl, Vv and Vd domains. The number was significantly higher in the Dd domain at 3 dpl than in non-injured control brains. There was no significant difference in other domains between 3 dpl and non-injured controls, although the number was higher in Dm, Dl and Vv at 3 dpl. mean±SD. p*<0.05; unpaired students' t-test.

Source Data 1. Data for Figure 1FGH.

Figure 2. scRNA-seq analysis identifies 4 distinct qRG clusters in 3-dpl and non-injured control telencephalons.

(A) Experimental procedures of scRNA-seq analysis of *GFAP:dTomato*$^+$ cells in adult telencephalons in 3-dpl and non-injured control conditions. Image was created using Biorender (https://biorender.com).

(B) UMAP of scRNA-seq data. Nine clusters are identified: 4 qRG, pRG/Neuroblast/NBN, OPC, Ependymal cells, Pineal gland, MN/NBN.

(C) Feature plot of markers defining each of the 9 clusters.

(D) Dot plot analysis of 46 qRG (left) and 39 pRG (right) markers in 4 qRG (qRG/NeuroEp1, qRG/NeuroEp2, qRG1, and qRG2) and pRG. qRG and pRG markers are classified into 4 and 3 Gene modules, respectively.

(E) Fraction of each cluster in 3-dpl and non-injured control conditions. qRG1 and pRG fractions were increased at 3 dpl, whereas qRG/NeuroEp2 and qRG2 fractions were decreased at 3 dpl.

Figure 2-figure supplement 1. FACS cell sorting and scRNA-seq data quality check.

Figure2-figure supplement 2. Cell clusters in the non-injured control dataset.

Figure2-figure supplement 3. Cell clusters in the 3-dpl dataset.

Figure2-figure supplement 4. Heatmap of the top 10 markers for clusters in the non-injured control and 3-dpl merged datasets.

Figure2-figure supplement 5. Gene module analysis of qRG markers.

Figure2-figure supplement 6. Regional marker analysis of qRGs.

1343 **Figure2-figure supplement 7.** Expression analysis of genes responsible for quiescence

1344 and stemness of qRGs.

1345 **Figure2-figure supplement 8.** Gene module analysis of pRG markers.

1346

Figure 3. TBI promotes the mTOR signaling pathway and transcription of *lgals2a*

and *gstm.1* in the ventricular zone of the pallium.

(A) Labeling of 3-dpl *Tg[GFAP:EGFP]* (green) transgenic telencephalon with anti-p-

rpS6 antibody (magenta). Right panels show higher magnification images.

Arrowheads indicate p-rpS6$^+$ RG, which are increased in the injured hemisphere

(right). Scale: 70μm.

(B) Fraction of p-rpS6$^+$ area in the GFAP:EGFP$^+$ Dd area in non-injured control and

lesioned hemispheres. The fraction of p-rpS6$^+$ area is significantly higher in

lesioned hemispheres than in non-injured control hemispheres. p*<0.05; paired t-

test.

(C) *In situ* hybridization of adult zebrafish telencephalon with the *lgals2a* RNA probe

in non-injured control and 3-dpl conditions. The right panel shows a higher-

magnification image of a 3-dpl sample. An asterisk indicates the stab-induced

injury site. Black filled and open arrowheads indicate mRNA expression in Dd and

Dm ventricular domains, respectively. Scale: 100 μm.

(D) The fraction of *lgals2a*$^+$ cells in the hemisphere area. This fraction is markedly

higher in 3-dpl lesioned hemispheres. mean±SD. p*<0.05; One way ANOVA

Turkey's multiple comparison test.

(E) *In situ* hybridization of adult zebrafish telencephalon with the *gstm.1* RNA probe in

non-injured control and 3-dpl conditions. The right panel shows a higher-

magnification image of a 3-dpl sample. An asterisk indicates the stab-induced

injury site. Black filled and open arrowheads indicate mRNA expression in Dd and

Dm/Vd/Vv ventricular domains, respectively. Scale: 100 μm.

(F) The fraction of *gstm.1*+ cells in the hemisphere area. mean±SD. The fraction is

higher in 3-dpl control and lesioned hemispheres. p*<0.05; One way ANOVA

Turkey's multiple comparison test.

1373

1374 **Source Data 1.** Data for Figure 3B.

1375 **Source Data 2.** Data for Figure 3DF.

1376

Figure 4. Pseudo-time analysis indicates that qRG/NeuroEp1 generates qRG1 and

pRG in separate lineages.

(A) Pseudo-time trajectory generated with monocle 3 for non-injured control (left) and

3-dpl conditions (right). qRG/NeuroEp1 is selected as a root for pseudo-time

trajectory. Three branches are specified in non-injured control conditions, whereas

two branches are specified in 3-dpl conditions. Trajectories toward qRG1 and pRG

are observed as Branch1 and Branch2, respectively, in both conditions.

(B) Gene expression profile along pseudo-time trajectory of branch1(qRG1) and

branch2 (pRG) in non-injured control and 3-dpl conditions. Thirteen genes are

selected: *ascl1a*, *ccnd1*, *cx43*, *fabp7*, *her4.1*, *her9*, *junbb*, *lgals2a*, *mcm7*, *mfge8*,

notch3, *pcna*, and *sox2*.

1388

Figure 4-figure supplement 1. Pseudo-time analysis using Monocle3 for non-injured

control and 3-dpl conditions.

Figure 4-figure supplement 2. Gene expression analysis of branch1- and branch2-

associated pRG/Neuroblasts/NBN cells.

Figure 4-figure supplement 3. AhR signaling pathway is specifically activated in

qRG/NeuorEp2.

1395

Figure 5. DEG analysis of qRG/NeuroEp1 in response to TBI.

(A) Scatter plot visualization of differentially expressed genes (DEGs) in

qRG/NeuroEp1 in response to TBI. Upregulated (red) and downregulated (blue)

genes are mapped.

(B) GO analysis of DEGs of qRG/NeuroEp1 in response to TBI.

(C) Dot-plot visualization of up- and downregulated genes of qRG/NeuroEp1 in

response to TBI across all RG clusters in non-injured control and 3-dpl conditions.

TOP30 upregulated genes (red) and TOP12 downregulated genes (blue) are shown.

Upregulated genes are classified into three groups: (1) genes whose expression is

high in both qRG1 and pRG at 3 dpl, (2) genes whose expression is high in qRG1

at 3 dpl, and (3) genes whose expression is high in pRG at 3 dpl. All downregulated
genes show low expression in both qRG1 and pRG at 3 dpl. Genes nominated by
GO analysis are shown in the right-most column.
(D) Venn diagram of up- and downregulated genes in qRG/NeuroEp1 linked by the
transition to qRG1 and pRG/Neuroblast/NBN.
1411
Figure 5-figure supplement 1. Signaling network of upregulated genes in
qRG/NeuroEp1 in response to TBI.
Figure 5-figure supplement 2. Gene expression profiles of the TOP30 upregulated
genes in qRG/neuroEp1 along a pseudo-time trajectory.
Figure 5-figure supplement 3. Signaling network of downregulated genes in
qRG/NeuroEp1 in response to TBI.
Figure 5-figure supplement 4. Gene expression profiles of the TOP12 downregulated
genes in qRG/neuroEp1 along a pseudo-time trajectory.
1420
Figure 6. Microglial depletion compromises regenerative responses of RGs after
TBI.
(A) Experimental design to investigate the temporal profile of regenerative responses of
RGs to TBI. Stab injury was introduced through the skull into the right hemisphere
of the telencephalon. MTZ was applied to fish for 2 days prior to TBI. After TBI,
BrdU was applied for 12 h prior to analysis at 1, 3, 7, and 14 dpl. Image was created
using Biorender (https://biorender.com).
(B) Double labeling of microglia-depleted telencephalon with anti-GFAP (white) and
anti-BrdU (magenta) antibodies at 1, 3, 7, and 14 dpl. Scales: 70μm.
(C) The number of BrdU+ cells in the ventricular zones of both hemispheres. The non-
injured control (NTR–; MTZ–) is the same sample used for analysis in Fig. 1F.
There was no significant difference between non-injured control and 3 dpl brains.
mean±SD. One-way ANOVA with Dunnett's multiple comparisons test.
(D) Comparison of the number of $BrdU^+$ cells between lesioned and intact hemispheres
at 3 dpl in three different conditions: (1) NTR–; MTZ–, (2) NTR+; MTZ+, and (3)
NTR–; MTZ+. Microglia-intact control (NTR–; MTZ–) is the same sample used for

analysis in Fig. 1G. As in microglia-intact conditions (NTR–; MTZ– and NTR–;

MTZ+), the number was significantly higher in lesioned hemispheres than in non-

injured control hemispheres for NTR+; MTZ+. However, the BrdU+ cell number

was decreased in microglia-depleted conditions (NTR+; MTZ+), compared with

microglia-intact conditions (NTR–; MTZ– and NTR–; MTZ+).   mean±SD.

p*<0.05, p**<0.01; paired t-test.

(E) The number of BrdU$^+$ cells in different ventricular domains of telencephalon: Dm,

Dd, Dl, Vv and Vd domains. Non-injured control (NTR–; MTZ–) was the same

sample used for analysis in Fig. 1H. The number was significantly higher in the Dd

domain in 3-dpl microglia-depleted brain (NTR+; MTZ+) than in non-injured

control microglia-intact brain (NTR–; MTZ–). P**<0.01; unpaired students' t-test.

1448

1449 **Source data 1**. Data for Figure 6CDE

1450 **Source data 2.** Data for Figure 6-figure supplement 1B

1451 **Source data 3.** Data for Figure 6-figure supplement 1E

1452 **Source data 4.** Data for Figure 6-figure supplement 2BC

1453 **Source data 5.** Data for Figure 6-figure supplement 3B

1454

1455 **Figure 6-figure supplement 1.** Microglia are activated in lesioned hemispheres at 1dpl

1456 **Figure 6-figure supplement 2.** NTR-MTZ treatment affects microglial integrity and

1457 reduces their number.

1458 **Figure 6-figure supplement 3**. Microglial depletion reduces the proliferative response

1459 of RGs.

1460

Figure 7. scRNA-seq analysis reveals that qRG1 is decreased in 3-dpl microglia-

depleted telencephalon.

(A) UMAP of scRNA-seq data in the 3-dpl and 3 dpl+MTZ merged dataset. Twelve

clusters are identified: 6 qRG, pRG/Neuroblast/NBN, Ependymal cells, Pineal

gland, MN/NBN, Microglia (MG)/Macrophage.

(B) Dot-plot analysis of qRG marker expression in 6 qRG clusters,

pRG/Neuroblast/NBN, and MN/NBN of the 3-dpl and 3 dpl+MTZ merged dataset.

qRG marker genes are classified into 4 new gene modules, called DM-gene modules

1-4. qRG marker genes of "Gene module 3" in the non-injured control and 3-dpl

merged dataset are rearranged into two MD-gene modules 1 and 2. Furthermore,

qRG marker genes of "Gene module 1" in the non-injured control and 3-dpl merged

dataset are also merged with "Gene module 4" to form MD-gene module 2.

(C) Fractions of each cluster in the 3-dpl and 3 dpl+MTZ merged dataset. qRG1 is

decreased in the 3-dpl+MTZ condition, whereas qRG/NeuroEp2 and MN/NBN are

increased.

1476

Figure 7-figure supplement 1. Cell clusters in the 3 dpl+MTZ dataset.

Figure 7-figure supplement 2. Heatmap of the top 10 markers for clusters identified in

the 3-dpl and 3 dpl+MTZ merged dataset.

Figure 7-figure supplement 3. Clarifying origins of new qRG3 and qRG4 clusters in

the 3-dpl and 3dpl+MTZ merged dataset.

Figure 7-figure supplement 4. Dot-plot analysis of pRG marker expression in clusters

of the 3-dpl and 3 dpl+MTZ merged dataset.

1484

Figure 8. DEG analysis of qRG/NeuroEp1 in response to TBI.

(A) Scatter plot visualization of differentially expressed genes (DEGs) in

qRG/NeuroEp1 at 3 dpl under microglia-depleted conditions versus microglia-intact

conditions. Upregulated (red) and downregulated (blue) genes are mapped.

(B) GO analysis of downregulated genes in qRG/NeurEp1 in response to microglial

depletion.

(C) GO analysis of upregulated genes in qRG/NeurEp1 in response to microglial

depletion.

(D) Dot-plot visualization of DEGs of qRG/NeuroEp1 across all RG clusters at 3 dpl

under microglia-depleted and -intact conditions. Gene expression columns of

qRG/NeuroEp1 under the 3-dpl and 3 dpl+MTZ conditions are indicated by blue

and red squares, respectively. The TOP30 upregulated genes and TOP30

downregulated genes are shown. Genes nominated by GO analysis are shown in the

right-most column: 4 upregulated genes: *lgasl2a*, *mdka*, *hmgb1a*, and *hbegfa*, and

downregulated genes: ribosomal genes, *vimentin*, *hsp70.1*, *hsp 70.2*, *hsp70.3*, *hsp*

70l, *junba*, and *junbb*.

(E) Venn diagram of up- and downregulated genes in qRG/NeuroEp1 linked to the

transition to qRG1 and pRG/Neuroblast/NBN. Potential targets of microglia are

indicated by yellow squares.

1504

1505 **Figure 8-figure supplement 1.** qRG/NeuroEp1 downregulated genes in response to

1506 microglial depletion.

1507 **Figure 8-figure supplement 2.** qRG/NeuroEp1 upregulated genes in response to

1508 microglial depletion.

1509

1510

1511 **Legends of Supplementary Figures**

1512

Figure 2-figure supplement 1. FACS cell sorting and scRNA-seq data quality

check.

(A) Setting of gate parameters for sorting dTomato$^+$ cells for non-injured control and 3-

dpl samples in microglia-intact conditions (NTR–/MTZ–).

(B) Setting of gate parameters for sorting dTomato$^+$ cells for non-injured control and 3-

dpl samples in microglia-depleted conditions (NTR+/MTZ+).

(C) Percentage of target cell population for four experimental conditions after FACS

sorting.

(D, E) scRNA data quality check for non-injured and 3-dpl samples in microglia-intact

conditions (NTR–/MTZ–) (D) and microglia-depleted conditions (NTR+/MTZ+) (E).

VLN plot for the number of genes detected per cell (nFeature_RNA), the number of

UMIs per cell (nCount_RNA) and mitochondrial gene percentages per cell (percent.mt).

1525

Figure2-figure supplement 2. Cell clusters in the non-injured control dataset.

(A) UMAP of the non-injured control dataset. Six clusters are specified.

(B) Heatmap of the top 10 differentially expressed genes for each of 6 clusters.

(C) Mapping of 6 clusters of non-injured control samples on UMAP of the merged

dataset for non-injured controls and 3-dpl samples. Clusters 0–5 correspond to

qRG/NeuroEp1,   qRG/NeuroEp2,   qRG1+pRG/Neuroblast/NBN+MN/NBN,

Ependymal cells+OPC, qRG2, Pineal gland, respectively.

1533

Figure2-figure supplement 3. Cell clusters in the 3-dpl dataset.

(A) UMAP of the 3-dpl control dataset. Seven clusters are specified.

(B) Heatmap of the top 10 differentially expressed genes for each of 7 clusters.

(C) Mapping of 7 clusters of 3-dpl samples on UMAP of the merged dataset for non-

injured controls and 3-dpl samples. Clusters 0–6 correspond to qRG/NeuroEp1,

qRG1+qRG2,   qRG/NeuroEp1+OPC,   qRG/NeuroEp2,   qRG1,

pRG/Neuroblast/NBN+MN/NBN, Ependymal cells, respectively.

1541

1542

Figure2-figure supplement 4. Heatmap of the top 10 markers for clusters in the
non-injured control and 3-dpl merged dataset.
The non-injured control and 3-dpl merged dataset showed 9 clusters. The Top 10
differentially expressed genes are shown in each cluster. In accordance with
differentially expressed genes, we classified 9 clusters to 4 qRG (qRG/NeuroEp1,
qRG/NeuroEp2, qRG1, qRG2), pRG/Neuroblast/NMN, MN/NBN, OPC, Ependymal
cells, Pineal gland.
1550

Figure2-figure supplement 5. Gene module analysis of qRG markers.
(A) Dot-plot of qRG marker expression of 4 qRG and pRG. 46 qRG markers were
classified into 4 gene modules: Gene modules 1-4 are highly expressed in
qRG/NeuroEp1, qRG/NeuroEp2, qRG1 and qRG2, respectively.
(B) UMAP of the non-injured control and 3-dpl merged dataset.
(C) Feature plot analysis of Gene Module1 highly expressed in the qRG/NeuroEp1
cluster.
(D) Feature plot analysis of Gene Module2 highly expressed in the qRG/NeuroEp2
cluster.
(E) Feature plot analysis of Gene Module3 highly expressed in the qRG1 cluster.
(F) Feature plot analysis of Gene Module3 highly expressed in the qRG2 cluster.
1562

Figure2-figure supplement 6. Regional marker analysis of qRGs.
(A) Dot-plot analysis of regional markers. *zic1* (Vv) was specifically expressed in
qRG/NeuroEp2 and qRG2.
(B) UMAP of the non-injured control and 3-dpl merged dataset.
(C) Feature plot analysis of regional marker genes. *zic1* (Vv) was specifically
expressed in qRG/NeuroEp2 and qRG2.
1569

1570 **Figure2-figure supplement 7. Expression analysis of genes responsible for**
1571 **quiescence and stemness of qRGs.**
1572 (A) Dot-plot analysis of genes related to quiescence and stemness.
1573 (B) UMAP of the non-injured control and 3-dpl merged dataset.
1574 (C) Feature plot analysis of genes related to quiescence and stemness.

1575

Figure2-figure supplement 8. Gene module analysis of pRG markers.

(A) Dot-plot analysis of pRG markers. pRG markers were classified into 3 gene

modules.

(B) UMAP of the non-injured control and 3-dpl merged dataset.

(C) Feature plot analysis of Gene Module 1.

(D) Feature plot analysis of Gene Module 2, whose expression is high in qRG1.

(E) Feature plot analysis of Gene Module 3, whose expression is high in

qRG/NeuroEp1/2.

1584

Figure 4-figure supplement 1. Pseudo-time analysis using Monocle3 for non-

injured control and 3-dpl conditions.

(A) Upper: UMAP with cell trajectory generated using Monocle 3 for non-injured

control (left) and 3-dpl (right) conditions. Lower: pseudo-time transition of cell

trajectory.

(B) Gene expression analysis along pseudo-time trajectory.

1591

Figure 4-figure supplement 2. Gene expression analysis of branch1- and branch2-

associated pRG/Neuroblast/NBN cells.

(A) Left: Cell trajectory generated with Monocle 3 in 3-dpl conditions.

pRG/Neuroblast/NBN cells are indicated in light blue. Right: UMAP of extracted

pRG/Neuroblast/NBN cells associated with Branch 1 and Branch 2. Branch 1-

associated qRG/Neuroblast/NBN were separated from Branch 2-associated

qRG/Neuroblast/NBN.

(B) Dot-plot of differentially expressed genes (DEGs) in Branch1- and Branch2-

associated pRG/Neuroblast/NBN cells. Branch 1-associated pRG/Neuroblast/NBN

cells show a very similar expression profile to qRG1.

(C) Violin plot of DEGs in Branch 1- and Branch 2-associated pRG/Neuroblast/NBN

cells. qRG1-specific RG markers are shown in red squares, and highly expressed

Branch 1-associated pRG/Neuroblast/NBN cells.

1605

Figure 4-figure supplement 3. AhR signaling pathway is specifically activated in qRG/NeuorEp2.

(A) UMAP of the non-injured control and 3-dpl merged dataset.

(B) Feature plot analysis of AhR signaling pathway genes, arh2 and cyp1b1.

(C) Violin plot of AhR signaling pathway genes, arh2 and cyp1b1.

(D) Dot-plot of expression of AhR target genes in qRGs and pRG.

Figure 5-figure supplement 1. Signaling network of upregulated genes in qRG/NeuroEp1 in response to TBI.

(A) Classification of upregulated genes in regulatory pathways; ribosomal metabolism, cytokine signaling and the JAK/STAT pathway.

(B) UMAP of the non-injured control and 3-dpl merged dataset.

(C) Feature plot analysis of upregulated genes related to JAK-STAT signaling.

(D) Feature plot analysis of upregulated genes related to cytokine signaling.

(E) Feature plot analysis of upregulated genes related to ribosomal metabolism.

Figure 5-figure supplement 2. Gene expression profiles of the top 30 upregulated genes in qRG/neuroEp1 along a pseudo-time trajectory.

Top: The top 30 upregulated genes whose expression is high in both qRG1 and pRG.

Middle: The top 30 upregulated genes whose expression is high in qRG1.

Bottom: The top 30 upregulated genes whose expression is high in pRG.

qRG1 lineage (Branch 1) and pRG lineage (Branch 2) at 3 dpl are shown at the left and right columns, respectively.

Figure 5-figure supplement 3. Signaling network of downregulated genes in qRG/NeuroEp1 in response to TBI.

(A) Classification of downregulated genes in regulatory pathways. Inhibition of neurogenesis, lipid metabolism, and cholesterol synthesis.

(B) UMAP of the non-injured control and 3-dpl merged dataset.

(C) Feature plot analysis of downregulated genes related to cholesterol metabolism.

(D) Feature plot analysis of downregulated genes related to lipid metabolism.

1637   (E) Feature plot analysis of downregulated *id2b* expression indicating inhibition of

1638   neurogenesis.

1639

1640 Figure 5-figure supplement 4. Gene expression profiles of the top 12

1641 downregulated genes in qRG/neuroEp1 along a pseudo-time trajectory.

1642   (A) The top 12 downregulated genes in the 3-dpl qRG1 lineage (Branch 1) using dot-

1643   plotting graph along a pseudo-time trajectory path.

1644   (B) The top 12 downregulated genes in the 3-dpl pRG lineage (Branch 2) using a dot-

1645   plotting graph along a pseudo-time trajectory path.

1646

1647 Figure 6-figure supplement 1. Microglia are activated in lesioned hemispheres at 1

1648 dpl.

(A) Microglial morphology with different sphericity indices. Scales: 10 μm.

(B) Visualization of mpeg1.1:EGFP$^+$ microglia in non-injured control, 1 dpl, and 3 dpl

telencephalon. The Dd domain in non-injured control and lesioned hemispheres are

indicated by blue and red squares, respectively. In the case of non-injured control

conditions, left and right hemispheres were used instead of non-injured controls and

lesioned hemispheres of 1- and 3-dpl brain. Scales: 100 μm.

(C) Sphericity distribution of microglia in the Dd domain of non-injured control and

lesioned hemispheres at 1 dpl and 3 dpl.

(D) Higher magnification of blue and red squares shown in panel (B). Microglia with

high and low sphericity are shown in yellow and red, respectively. Scales: 20 μm.

(E) The percentage of the number of microglia located in the lesioned hemispheres

relative to the number of microglia located in both lesioned and non-injured control

hemispheres. Histogram for microglia with high sphericity (index > 0.6) and

histogram for all microglia are shown to the left and right, respectively. Only high-

sphericity microglia at 1dpl are significantly higher than in non-injured conditions.

p*<0.05: One-way ANOVA, Dunnett's multiple comparison test.

1665

1666 Figure 6-figure supplement 2. NTR-MTZ treatment affects microglial integrity

1667 and reduces their number.

(A) Visualization of mpeg1.1:EGFP$^+$ microglia in *Tg[mpeg1.1:NTR-eYFP]* transgenic

telencephalon with no MTZ treatment, MTZ treatment for 2 days, and MTZ

treatment for 3 days. No microglial cells were detected in MTZ treatment for 3

days. Scales: 100 µm.

(B) Number of microglia per section of no MTZ, 2d MTZ, and 3d MTZ treatments. The

number is significantly lower in 3d MTZ treatment than in no MTZ treatment.

p***<0.005: One way ANOVA, Bonferroni multiple comparison test.

(C) Sphericity index distribution of microglia with no MTZ, 2d MTZ, and 3d MTZ

treatments. The sphericity index is significantly higher in 2d MTZ treatment than in

no MTZ treatment; however, there is no significant difference between 3d MTZ

treatment and no MTZ treatment. p***<0.005: One way ANOVA, Bonferroni

multiple comparison test.

1680

Figure 6-figure supplement 3. **Microglial depletion reduces the proliferative**

response of RGs.

(A) Double labeling of 3-dpl microglia-depleted (NTR+; MTZ+) and microglia-intact

(NTR–; MTZ+) telencephalon with anti-GFAP and anti-BrdU antibodies. Lower

panels indicate higher-magnification images of dotted squares shown in the upper

panels. Scales: 70 µm (upper panels) and 30 µm (lower higher magnification image

panels).

(B) The number of BrdU$^+$; GFAP$^+$ cells, which correspond to pRGs, in the Dd domain

of the ventricular zone. The number is significantly decreased in 3-dpl microglia-

depleted telencephalon, compared with 3-dpl microglia-intact telencephalon.

p*<0.05: Unpaired t test.

1692

Figure7-figure supplement 1. Cell clusters in the 3 dpl+MTZ dataset.

(A) UMAP of the 3 dpl+MTZ dataset. Eleven clusters are specified.

(B) Heatmap of the top 10 differentially expressed genes for each of 11 clusters.

(C) UMAP of the merged dataset for 3-dpl and 3 dpl+MTZ samples.

(D) Mapping of 11 clusters of 3 dpl+MTZ samples on UMAP of the merged dataset for

3-dpl and 3 dpl+MTZ samples. Clusters 0–11 correspond to qRG/NeuroEp1+qRG3,

qRG1+qRG2, pRG/Neuroblast/NBN, qRG4, qRG1+qRG2, 3 types of MN/NBN,

Pineal gland, Ependymal cells, and Microglia (MG)/macrophage, respectively.

1701

Figure 7-figure supplement 2. Heatmap of the top 10 markers for clusters

identified in the 3-dpl and 3 dpl+MTZ merged dataset.

Eleven clusters are classified into qRG/NeuroEp1, qRG/NeuroEp2, qRG1, qRG2,

pRG/Neuroblast/NBN, MN/NBN, Ependymal cells, Pineal gland, Microglia

(MG)/Macrophage. In addition, two new qRG clusters, qRG3 and qRG4, are specified.

Genes specific to each cluster are shown in the left column.

1708

Figure 7-figure supplement 3. Clarifying origins of new qRG3 and qRG4 clusters

in the 3-dpl and 3dpl+MTZ merged dataset.

(A) Split UMAP of the 3-dpl and 3 dpl+MTZ merged dataset.

(B) UMAP of the non-injured control and 3-dpl merged dataset.

(C) Reference mapping cell clusters identified in the 3-dpl and 3 dpl+MTZ merged

dataset on UMAP of the non-injured control and 3-dpl merged dataset. qRG3 is

derived from qRG/NeuroEp1, whereas qRG4 is derived from qRG/NeuroEp1 and

OPC.

1717

Figure 7-figure supplement 4. Dot-plot analysis of pRG marker expression in

clusters of the 3-dpl and 3 dpl+MTZ merged dataset.

Dot-plot analysis of pRG marker expression in 6 qRG clusters, pRG/Neuroblast/NBN,

and MN/NBN of the 3-dpl and 3 dpl+MTZ merged dataset. pRG marker genes are

classified into new 4 gene modules, called DM-gene modules 1-4. pRG markers of

"Gene module 1" in the non-injured control and 3-dpl merged dataset are rearranged

into three MD-gene modules 1 – 3. Most pRG markers of "Gene module 3" in the non-

injured control and 3-dpl merged dataset are overlapped with MD-gene modules 4.

1726

Figure 8-figure supplement 1. qRG/NeuroEp1 downregulated genes in response to

microglial depletion.

(A) Classification of downregulated genes in regulatory pathways: ribosomal
metabolism, chaperonin-mediated protein folding, cytokine and JAK/STAT
pathway, jun, her4/her6.
(B) Split UMAP of the 3-dpl and 3dpl+MTZ merged dataset.
(C) Feature plot analysis of downregulated genes related to the JAK-STAT and
cytokine pathways.
(D) Feature plot analysis of downregulated genes related to chaperone-mediated protein
folding.
(E) Feature plot analysis of downregulated genes related to ribosomal metabolism.
(F) Feature plot analysis of downregulated genes related to her4/her6.
(G) Feature plot analysis of downregulated genes related to jun.
1740

Figure 8-figure supplement 2. qRG/NeuroEp1 upregulated genes in response to
microglial depletion.
(A) Classification of upregulated genes in the regulatory pathway: wound response and
epithelium regeneration.
(B) Split UMAP of the 3-dpl and 3dpl+MTZ merged dataset.
(C) Feature plot analysis of upregulated genes related to wound response and
epithelium regeneration.
1748
1749

List of supplementary materials

Source Data files

Figure 1-Source Data 1. Data for Figure 1FGH.

Figure 3-Source Data 1. Data for Figure 3B.

Figure 3- Source Data 2. Data for Figure 3DF.

Figure 6- Source Data 1. Data for Figure 6CDE

Figure 6- Source Data 2. Data for Figure 6-figure supplement 1B

Figure 6- Source Data 3. Data for Figure 6-figure supplement 1E

Figure 6- Source Data 4. Data for Figure 6-figure supplement 2BC

Figure 6- Source Data 5. Data for Figure 6-figure supplement 3B

1760

Supplementary figures

Figure 2-figure supplement 1. FACS cell sorting and scRNA-seq data quality check.

Figure2-figure supplement 2. Cell clusters in the non-injured control dataset.

Figure2-figure supplement 3. Cell clusters in the 3-dpl dataset.

Figure2-figure supplement 4. Heatmap of the top 10 markers for clusters in the non-
injured control and 3-dpl merged datasets.

Figure2-figure supplement 5. Gene module analysis of qRG markers.

Figure2-figure supplement 6. Regional marker analysis of qRGs.

Figure2-figure supplement 7. Expression analysis of genes responsible for quiescence
and stemness of qRGs.

Figure2-figure supplement 8. Gene module analysis of pRG markers.

Figure 4-figure supplement 1. Pseudo-time analysis using Monocle3 for non-injured
control and 3-dpl conditions.

Figure 4-figure supplement 2. Gene expression analysis of branch1- and branch2-
associated pRG/Neuroblasts/NBN cells.

Figure 4-figure supplement 3. AhR signaling pathway is specifically activated in
qRG/NeuorEp2.

Figure 5-figure supplement 1. Signaling network of upregulated genes in
qRG/NeuroEp1 in response to TBI.

Figure 5-figure supplement 2. Gene expression profiles of the TOP30 upregulated
genes in qRG/neuroEp1 along a pseudo-time trajectory.

Figure 5-figure supplement 3. Signaling network of downregulated genes in

qRG/NeuroEp1 in response to TBI.

Figure 5-figure supplement 4. Gene expression profiles of the TOP12 downregulated

genes in qRG/neuroEp1 along a pseudo-time trajectory.

Figure 6-figure supplement 1. Microglia are activated in lesioned hemispheres at 1dpl

Figure 6-figure supplement 2. NTR-MTZ treatment affects microglial integrity and

reduces their number.

Figure 6-figure supplement 3. Microglial depletion reduces the proliferative response

of RGs.

Figure 7-figure supplement 1. Cell clusters in the 3 dpl+MTZ dataset.

Figure 7-figure supplement 2. Heatmap of the top 10 markers for clusters identified in

the 3-dpl and 3 dpl+MTZ merged dataset.

Figure 7-figure supplement 3. Clarifying origins of new qRG3 and qRG4 clusters in

the 3-dpl and 3dpl+MTZ merged dataset.

Figure 7-figure supplement 4. Dot-plot analysis of pRG marker expression in clusters

of the 3-dpl and 3 dpl+MTZ merged dataset.

Figure 8-figure supplement 1. qRG/NeuroEp1 downregulated genes in response to

microglial depletion.

Figure 8-figure supplement 2. qRG/NeuroEp1 upregulated genes in response to

microglial depletion.

1802

Supplementary Tables

Supplementary Table 1. Differentially expressed genes for each cluster in the non-

injured dataset.

Supplementary Table 2. Differentially expressed genes for each cluster in the 3-dpl

dataset.

Supplementary Table 3. Differentially expressed genes for each cluster in the non-

injured control and 3-dpl merged dataset.

Supplementary Table 4. Differentially expressed genes for each cluster in the

3dpf+MTZ dataset.

Supplementary Table 5. Differentially expressed genes for each cluster in the 3-dpl

and 3dpl+MTZ merged data set.

1814

1815

1816

1817 **Appendix**

1818

Key Resources Table				
Reagent type (species) or resource	**Designation**	**Source or reference**	**Identifier**	**Additional information**
strain, strain background (zebrafish, Danio rerio)	Okinawa wild type	PMID:28 196805		
genetic reagent (zebrafish, Danio rerio)	Tg(*mpeg1*:NTR-EYFP)^{w202}	PMID: 24 961798	ZFIN: ZDB-TGCONSTRCT-140903-2	Randall T. Moon University of Washington
genetic reagent (zebrafish, Danio rerio)	Tg(mpeg1.1:EGFP) ^{oki053}	PMID: 34 872632		
genetic reagent (zebrafish, Danio rerio)	Tg(mfap4:tdTomato) ^{oki083}	PMID: 34 872632		
genetic reagent (zebrafish, Danio rerio)	Tg(GFAP:dTomato) ^{nns17Tg}	PMID: 22 302816	ZDB-TGCONSTRCT-120215-2	
genetic reagent (zebrafish, Danio rerio)	Tg(GFAP:GFP)^{mi2001}	PMID: 31 998714	ZDB-ALT-060623-4	

antibody	Anti-BrdU	Abcam	Cat# ab6326	
antibody	Anti-zrf1/Anti-GFAP	ZIRC	ZDB-ATB-081002-46	
antibody	Anti-GFP	Thermofisher	Cat# A11122	
antibody	Anti-p_rps6 (Ser235/2376)	Cell Signaling	Cat# 2211	
commercial assay or kit	Single Cell 3' Library kit v3	10X Genomics	Part No.: 1000078	
commercial assay or kit	SYTOX™ Dead Cell Stain	Thermofisher	Cat# S34862	
commercial assay or kit	ReverTra Ace□™ qPCR master mix with gDNA remover	TOYOBO	Code No FSQ-301	
chemical compound, drug	5-Bromo-2'-Deoxyuridine	Nacalai Tesque	Cat# 05650-24	
chemical compound, drug	Metronidazole	Sigma Aldrich	Cat# M1547-25G	
chemical compound, drug	Deoxyribonuclease I	Worthington	Cat# LK003170	
chemical compound, drug	Papain	Worthingt	Cat# LK	

		on	003176	
chemical compound, drug	Sepasol-RNA/Super G	Nacalai Tesque	Cat# 09379	
sequence- based reagent	gstm.1 in situ Forward primer	NM_2126 76.1	PCR primer	ATGGCA ATGAAAT TGGCTTA TTG
sequence- based reagent	gstm.1 in situ reverse primer	NM_2126 76.1	PCR primer	TCACTCC TTCTTGT TTCCCCA
sequence- based reagent	lgals2a in situ Forward primer	NM_2128 94	PCR primer	ATGGCCG GTGTGCT TATACAG A
sequence- based reagent sequence- based reagent	lgals2a in situ Reverse primer	NM_2128 94	PCR primer	CTATTTA ATTTCAA CCCCTTG GAT
sequence- based reagent	NFSβ genotyping Forward primer	Gene ID: 945778	PCR primer	AGATCA AAACGCT ACTGCAA TACA
sequence- based reagent	NFSβ genotyping Reverse primer	Gene ID: 945778	PCR primer	CTGGTCA ACAACC AGCTTCA G
software, algorithm	Seurat version 4.1.0	PMID: 34 062119		https://satij alab.org/se

software, algorithm			
			urat/
software, algorithm	Monocle3 version 0.2.3.0	PMID: 28 825705		https://cole-trapnell-lab.github.io/monocle3/
software, algorithm	dittoSeq version 4.1	PMID: 33 313640		https://github.com/dtm2451/dittoSeq
software, algorithm	Cellranger version 3.1.0	PMID: 28 091601		https://support.10xgenomics.com/single-cell-gene-expression/software/pipelines/latest/what-is-cell-ranger
software, algorithm	clusterProfiler version 4.0.5	PMID: 22 455463		https://hbctraining.github.io/DGE_workshop/lessons/09_functional_analysis.html
software, algorithm	topGO version 2.44.0	DOI: 10.1 8129/B9.		

		bioc.topGO		
software, algorithm	pathview version 1.32.0	PMID: 23 740750		
software, algorithm	IMARIS version 9.3.0	URL: http://www.bitplane.com/imaris/imaris		
software, algorithm	ImageJ. Version 2.0.0	Rasband, W.S., ImageJ, U. S. National Institutes of Health, Bethesda, Maryland, USA, https://imagej.nih.gov/ij/, 1997-2018.		
software, algorithm	Biorender	https://biorender.com		

1819

1820

1821

1822

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7

Figure 8