De-differentiation and Proliferation of Artery Endothelial Cells

Drive Coronary Collateral Development

Gauri Arolkar[1*], Sneha K.[1*], Hanjay Wang[2], Karen M. Gonzalez[3,4], Suraj Kumar[1],

Bhavnesh Bishnoi[1], Pamela E. Rios Coronado[5], Y. Joseph Woo[2], Kristy Red-

Horse[3,4,6], Soumyashree Das[1#]

[1] National Center for Biological Sciences, Tata Institute of Fundamental Research,

Bengaluru, KA 560065, India

[2] Department of Cardiothoracic Surgery, Stanford University School of Medicine,

Stanford, CA 94305, USA

[3] Department of Biology, Stanford University, Stanford, CA 94305, USA

[4] Institute for Stem Cell Biology and Regenerative Medicine, Stanford University

School of Medicine, Stanford, CA 94305, USA

[5] Department of Bioengineering, Stanford University, Stanford, CA 94305, USA

[6] Howard Hughes Medical Institute

* Equal contribution

Corresponding author (soumyashree@ncbs.res.in)

Key Words

Coronary Collateral Artery, Endothelial Cell, Vascular Endothelial Growth Factor

Receptor-2, De-differentiation, Proliferation, Cardiac Regeneration, Single Cell RNA

Sequencing, Myocardial Infarction, Ischemia

Abstract

Collateral arteries act as natural bypasses which re-route blood flow to

ischemic regions and facilitate tissue regeneration. In an injured heart, neonatal

artery endothelial cells orchestrate a systematic series of cellular events, which

includes their outward migration, proliferation, and coalescence into fully functional

collateral arteries. This process, called Artery Reassembly, aids complete cardiac

regeneration in neonatal hearts, but is absent in adults. The reason for this age-

dependent disparity in artery cell response is completely unknown. In this study, we

investigated if regenerative potential of coronary arteries, like in cardiomyocytes, is

dictated by their ability to de-differentiate. We performed single cell RNA sequencing

of coronary endothelial cells and identified differences in molecular profiles of

neonatal and adult endothelial cells. Neonates show significant increase in actively

cycling artery cells that expressed prominent de-differentiation markers. Using both,

in silico pathway analyses and *in vivo* experiments, we show that cell cycle re-entry

of pre-existing artery cells, subsequent collateral artery formation and cardiac

function is dependent on arterial VegfR2. This sub-population of de-differentiated

and proliferating artery cells is absent in non-regenerative P7 or adult hearts.

Together, our data indicate that adult artery endothelial cells fail to drive collateral

artery development due to their limited ability to de-differentiate and proliferate.

Background

Cardiovascular disease (CVD) represents 32% of all global deaths (World Health Organization) and its burden is increasing sharply in many new parts of the world. Earlier, deaths due to CVD dominated the western world, but currently, at least 58% of all CVD deaths occur in Asia (*http://ghdx.healthdata.org/gbd-results-tool*). The primary cause of CVD is coronary artery blockage, and can lead to ischemic heart disease (IHD). Despite the deadly outcomes, management of symptoms associated with IHD often include invasive procedures, which, many patients are ineligible for. Effective therapeutic approaches targeting complete cardiac regeneration, can reduce mortality and/or manage symptoms of IHD. This can be accomplished by understanding, and subsequently utilizing, the molecular mechanisms associated with different physiological aspects of IHD. One such mechanism adapted by the heart is building collateral arteries.

Large collateral arteries enhance perfusion[8], vascularize hypoxic area, improve myocardial viability[9], and are associated with better survival rate[10]. Collateral arteries are the arterial segments which connect a healthy arteriole to an occluded/non-perfused arteriole, and are sufficiently large to carry adequate blood flow to ischemic regions in need. Data from clinical studies suggest that the presence of native/pre-existing coronary collateral arteries present at the onset of myocardial infarction (MI) helps limit the infarct size, and improves prognosis[9,11]. The number of pre-existing collateral arteries as well as their capacity to undergo remodeling upon occlusion, are the key determinants of tissue recovery[12]. Genetic factors are proposed to regulate differences in collateral number and their functionality across human races and species[13,14]. However, a complete map of

molecular mechanisms/pathways remain unknown. These molecular mechanisms

could be regulated by genetic factors, and can be a useful tool to design therapeutic

interventions for IHD patients.

Brain, a highly complex organ with low regenerative ability, is known to

possess pre-existing collateral vessels in its pial layer. In mice, these collateral

connections develop at an embryonic stage, and provide protection against ischemia

in adulthood[15,16]. Pre-existing collateral arteries in mouse hind limb can grow by two

distinct processes— arterialization of capillary endothelial cells (ECs)[17] and lateral

expansion/widening of smaller pre-existing collateral arteries i.e. arteriogenesis[18].

Both, in brain and hind limb, genetic background is a major contributor towards the

variation in the number of observed collaterals[19,20]. Specifically, polymorphism in

Vascular Endothelial Growth Factor-a (*Vegfa)* locus is shown to control Vegfa

expression and pre-determine the extent of collateral artery network in mice[12,21].

Particularly in hind limb ischemia model, density of pre-existing collateral arteries is

regulated via Vegf Receptor-1 (VegfR1)[21]. However, if VegfR1 also controls post-

ischemic coronary collateral development in mice, is unknown.

In mice, complete cardiac regeneration is accomplished only when hearts are

injured within a neonatal regenerative window (P0-P6). Any injury beyond this time

point (>P7), limits cardiac regeneration[22,23]. Previously we have shown that, in

regenerative neonatal mice, coronary collateral arteries are built by the systematic

execution of 3 major cellular events driven by pre-existing artery ECs— migration,

proliferation and coalescence[24]. Artery ECs expressed C-X-C motif Chemokine

Receptor 4 (CXCR4), a receptor for a chemokine, C-X-C motif Chemokine Ligand 12

(Cxcl12), which steers the migration of artery ECs into the watershed. This process,

named Artery Reassembly, involves disassembly of artery tips into single migratory

artery cells, followed by their reassembly to build new artery segments (collateral

arteries). Interestingly, despite *Cxcl12* expression in adult injured watershed, Artery

Reassembly was absent in these hearts. The precise cause for absence of Artery

Reassembly in adult injured hearts remains unknown.

The age-dependent discrepancy in artery EC response triggered by MI, could

be because of their distinct molecular profiles[25–27]. To identify potential molecular

drivers, we performed single cell RNA sequencing of neonatal and adult ECs, at a

time point prior to when collateral arteries appear in the injured mouse hearts. Using

both bioinformatics analyses and *in vivo* experiments, we show that, upon injury,

neonatal artery ECs undergo de-differentiation and VegfR2-mediated proliferation,

which is critical for collateral artery formation by Artery Reassembly and subsequent

cardiac function. Both cellular events, arterial de-differentiation and proliferation,

were absent in older non-regenerative hearts, post-MI. We hope, in future, these

findings will aid strategizing ways to induce collateral arteries and enhance

regenerative potential of cardiac tissue, by specifically targeting artery cells.

Results

Increase in cycling neonatal artery ECs, post-MI

Molecular profiling cardiac ECs which participate in collateral artery formation,

can improve our understanding of ways in which complete cardiac reperfusion and

regeneration can be attained. We thus, performed single cell RNA sequencing of

neonatal post-MI heart cells. Specifically, *Cx40CreER; Rosa26^TdTomato* mice were

used, where a single dose of Tamoxifen to the lactating mothers induced expression

of TdTomato fluorescence in *Cx40* expressing artery endothelial cells (aECs) in

neonates. Neonates were then subjected to either sham or MI surgeries at P2, and

hearts were harvested at P4 or P5 (Figure **1A**). TdTomato+ cardiac cells were sorted

using Fluorescent Activated Cell Sorting (FACS) method, followed by sequencing

using 10X genomics platform. Sequences were processed using Cell Ranger and

the subsequent output files were analyzed using Seurat on RStudio[28].

Sorting TdTomato+ cells during FACS, enriched our samples for *Cx40CreER*+

lineage labelled/traced artery ECs. When analyzed, these TdTomato+ cells were

additionally identified to be cardiomyocytes, smooth muscle cells and cells

undergoing endoMT (data not shown). We also obtained some TdTomato- non-artery

cells. To specifically study EC biology post-injury, we thus analyzed *Cadherin5*+ ECs,

which included, both, TdTomato+ artery ECs, genetically labeled between P0-P2

(Figure **1A**), and TdTomato- artery ECs (if any), identified by expression of known

arterial EC genes. Pre-processing and quality control steps (Supplementary figure

S1A) resulted in 5432 sham and 2200 MI ECs (Figure **1B**-**D**)**,** out of which 4423

sham, and 262 MI ECs, expressed at least one copy of TdTomato (Figure **1E**, **F**). As

previously observed[29,30], compared to sham, we obtained fewer cells upon MI. This

is likely due to extensive cell death following coronary ligation and subsequent tissue

ischemia.

Unbiased grouping of *Cadherin5*+ ECs from sham and MI, resulted in 10

distinct clusters (Supplementary figure **S1B**). Merged data from sham and MI were

projected onto two-dimensional space using Uniform Manifold Approximation and

Projection (UMAP). The identified cell types were— mature artery ECs (aEC1-4),

transitioning ECs (trans-EC), cycling artery ECs (cyc-aEC), cells undergoing

endothelial to mesenchymal transition (endoMT), smooth muscle cells (SMC1,2) and

endocardial ECs (endoEC) (Figure **1G**, **H**).

We identified the cell types with known vascular and cardiac cell markers and

their relative expression levels (Figure **1I**, Supplementary figure **S1-3**). aEC1-4

expressed markers for mature artery ECs (*Cx40*, *Cxcr4*) and SMC1,2 expressed

markers for both pericytes and SMCs (*Pdgfrb*, *Notch3*). Cells undergoing endoMT

were identified by upregulated expression of mesenchymal markers such as *Col1a1*

and *Cdh2* and endo-MT associated transcription factors like *Serpine1*, *Twist1* and

Snai1 (Figure **1I** and Supplementary figure **S1C**). Some of these markers were co-

expressed by SMC clusters (Supplementary figure **1C**). An EC cluster expressed

several genes specific to G2/M (*Top2a*, *Mki67*, *Birc5*, *Cenpa*, *Kif2c*, *Bub1*, *Nuf2*,

Hmmr) (Figure **1I** and Supplementary figure **S1D-G**) and G1/S (*Tyms*, *Mcm4*, *Mcm6*,

Mcm7, *Ung*, *Chaf1b*, *Fen1*, *Gmnn*) (Figure **1I** and Supplementary figure **S1H-K**)

transition. Sham cells in this particular cluster mostly expressed genes like *Cx40*,

Gja4, *Jag1*, *Cxcr4* (Supplementary figure **S2A-D**) known to be significantly

expressed and being specific to mouse[31] and human[32] coronary arteries. The top 20

genes in this cluster included artery specific genes[33] like *Fgfr3*, *Slc45a4*, *Gja4* and

Ssu2 (data not shown). We hence, named the cluster as cycling aEC (cyc-aEC).

EndoEC expressed *Npr3*[34], *Cdh11*, *Tgfbi*[35] and *Adgrg6*[36] (Figure **1I**) and some

venous markers (Supplementary figure **S1L**). Non-endothelial cardiac cells were

mostly absent in our dataset (Supplementary figure **S1L**).

We also identified a cluster of cells (named as trans-EC in Figure **1G**, **H**) that

did not express markers for any single endothelial cell sub-type. Re-clustering this

group resulted into 4 distinct EC sub-clusters, those highly expressed markers for

arteries, capillaries, venous/capillaries and myofibrillar genes[37] (Supplementary

figure **S3A**-**B**). This cluster especially was checked for doublets. No doublets were

present in MI trans-EC cluster (data not shown). Trans-EC cluster in the sham

dataset had 3 doublets (data not shown). 42.8% of artery-like ECs, in this cluster,

were TdTomato$^+$ (Supplementary figure **3C**) indicating their origin from pre-existing

artery ECs, those labelled between P0-P2 (Figure **1A**). Additionally, this artery-like

EC sub-cluster had fewer cells expressing *Cx40* or *Cxcr4* (Supplementary figure **3B**).

The indistinct molecular nature of this entire cluster, and its proximity to both, a

proliferative (cyc-aEC) and mature (aEC1/3) artery EC clusters (Supplementary

figure **S1B**), led us to hypothesize that cells in this cluster are in an intermediate

state, i.e., between a proliferative and mature state. We hence, called this cluster,

transitioning EC (trans-EC). Bioinformatics analyses of *Pecam1$^+$* cells, instead of

Cadherin5$^+$ cells, resulted in identification of the same cell types, clustered in a

similar manner (Supplementary figure **S4**).

Artery cells rarely proliferate, and exiting cell cycle is required for artery cell

specification[24,38–40]. However, we identified an artery EC cluster (cyc-aEC) in our

dataset, expressing several genes that mark G1/S and G2/M transitions (Figure **1I**

and S1D-**K**). Cell cycle analyses of *Cadherin5$^+$* cells showed a 12.4% increase in MI

cells undergoing G2 to M transition as compared to the sham group (Figure **1J**). The

number of MI cells in S-phase increased by 3.3% and G0/G1 phase reduced by

15.6% (Figure **1J**), suggesting a higher number of MI cells were likely mitotic.

We next performed cluster-wise quantification of mitotic cells (expressing

G2/M markers), in sham and MI groups. In sham group, 2.5% of all mitotic cells were

present in artery clusters (aEC1-4 and cyc-aEC) (Figure **1K**). In MI group, we

observed 11% of all mitotic cells in artery clusters (Figure **1K**), indicating a 4.4-fold

increase in mitotic artery ECs, upon MI. 15.9% of all *Cadherin5$^+$* cells were mitotic in

the MI group, and majority of these cells (~69.2%) were artery ECs (Figure **1K**), a

stably differentiated cell type in neonatal hearts.

MI-induces new artery development, from sources independent of pre-existing

artery ECs

Post-MI artery EC proliferation is associated with Artery Reassembly and

complete cardiac regeneration[24]. Use of artery EC specific *Cx40CreER* mouse line,

allowed us to mark the lineage of pre-existing artery ECs with TdTomato fluorescent

reporter (Figure **1A**). We performed cell cycle analysis, and quantified the number of

Cadherin5+TdTomato+ ECs that expressed genes for G2/M transition, and were

likely to be mitotic. 13.4% of all *Cadherin5*+TdTomato+ MI cells were mitotic, ~63% of

which were present in artery EC clusters (aEC1-4 and cyc-aEC) (Figure **1L**).

Analyzing mitotic cell distribution within the MI group showed 69.3% of all mitotic

cells to be present in artery clusters but very few (6.3%) of them were TdTomato+

artery ECs (Figure **1M**). This suggests that MI induces expansion of an artery EC

population, not lineage labelled by, or traced from, *Cx40CreER*+ cells. This is

consistent with our previous observation that injury causes organ-wide artery growth

(especially, artery-tips), independent of Artery Reassembly, and through capillary

differentiation[24].

We next performed *in vivo* experiments to confirm if an artery EC population,

originating from non-artery sources (as seen in Figure **1M**) is indeed proliferating,

upon MI. EdU labelled P6 hearts from *Cx40CreER; Rosa26*^{TdTomato} neonates

(Tamoxifen at P0, MI at P2) were harvested and immunostained for Cx40. Cx40

immunolabelled all mature artery ECs, including *Cx40CreER* lineage-traced

TdTomato⁺ pre-existing aECs. We indeed, observed numerous Cx40⁺TdTomato⁻

arteries (Supplementary figure **S5A**-**C**), especially in the ischemic regions.

Furthermore, we identified these artery ECs to be proliferating (EdU⁺) and integrated

into Cx40⁺ vessels (Figure **1N**-**O** and Supplementary figure **S5A**-**C**). As expected, we

also observed Cx40⁺TdTomato⁺EdU⁺ artery ECs (Figure **1N**, **P** and Supplementary

figure **S5D**, **E**). Thus, ischemia can induce formation of new coronary arteries via

multiple mechanisms (Supplementary figure **S5F**). Firstly, coronary collateral artery

segments are generated via Artery Reassembly, a process driven by pre-existing

artery ECs. Secondly, an overall organ-wide artery growth such as extension of

artery tips occurs by two processes: 1) incorporation of capillary ECs as observed

earlier[24] (also suggested in Supplementary figure **S5A**-**C**) and by 2) proliferation of

pre-existing (TdTomato⁺) artery ECs (as shown in Supplementary figure **S5D**, **E**).

Neonatal artery ECs de-differentiate upon MI

Stably differentiated cells may attain a less differentiated state and re-enter

cell cycle in response to pathophysiological cues. The acquired proliferative abilities

allow them to transform into less specialized cell type. They can then, either re-

differentiate into cells within the same lineage, or into cells of a different lineage[41].

This phenomenon can often be detected through changes in gene expression

patterns. Presence of an actively cycling differentiated cell cluster in neonates (cyc-

aEC) led us to hypothesize that cells in this cluster de-differentiate and contribute to

new arteries, in response to MI.

De-differentiation is indicated by loss of differentiation markers, cell cycle re-

entry, and expression of early genes, suggesting lack of cell fate specification. We

checked if MI cells, when compared to sham cells, manifested any of these de-

differentiation indicators. In sham, *Cx40*, a mature artery EC marker, was expressed

by cyc-aECs, at levels comparable to mature artery clusters—aEC1-4 (Figure **2A**,

B). In MI group, cells in cyc-aEC cluster expressed *Cx40*, at levels much lower than

what was observed in sham cyc-aEC (Figure **2B**) or the rest of the artery ECs

(aEC1-4) (Figure **2B**, **C**). The percentage of cells expressing *Cx40*, upon MI,

reduced in all clusters (Figure **2A**, **C**), but this decrease was most prominent in trans-

EC (19% in sham, 0% in MI) and cyc-aEC (57% in sham, 3% in MI) clusters

(Supplementary figure **S6A**). A similar reduction in number of *Cxcr4*[+] MI cells was

observed specifically in cyc-aEC and trans-EC clusters (Supplementary figure **S6A**).

Thus, there is an overall reduction in number of cells expressing mature artery

markers, and their expression levels, in cyc-aEC and trans-EC cluster, upon MI.

During development, an immature coronary vessel plexus undergoes

morphogenesis to build coronary arteries[42]. We hypothesized that cyc-aECs, if de-

differentiating, would re-express some early genes expressed by this vessel plexus

that mostly consists of venous/capillary ECs and from which coronaries originate

during development. We hence, analyzed sham and MI cells for cardiac

microvascular genes such as *Aplnr*[43] and *Car4*[32,44]. There was a dramatic increase in

number of MI cells expressing *Aplnr* in artery ECs (aEC1-4) (Figure **2A**, **C and**

supplementary figure **S6A**), but the gene was only minimally expressed in mature

artery EC clusters (aEC1-4), upon MI (Figure **2B**). In contrast to this, the total

number of *Aplnr*[+] MI cells was most prominently increased in cyc-aEC (by 2.5 fold)

and trans-EC (by ~2 fold) cluster (Supplementary figure **S6A**), and so was the

expression of *Aplnr* in these clusters (Figure **2B**). *Car4* also followed the same

pattern (Figure **2A**, **C** and Supplementary figure **S6A**), indicating both cyc-aEC and

trans-EC express venous-capillary marker genes, upon injury.

Angiogenic properties of ECs facilitate vascular remodeling and subsequent

cardiac regeneration. We next investigated if cyc-aEC (or trans-EC) clusters

expressed any pro-angiogenic genes, yet another cellular feature of de-

differentiation. Upon MI, 61% of the cyc-aECs and 47% of trans-ECs expressed pro-

angiogenic *Apln* gene (Figure **2A**, **C** and Supplementary figure **S6A**), known to drive

vascular development[45,46] and regeneration[47–49] *in vivo*. Expression of other

angiogenic genes like *Foxm1*[50,51], *Nrp2*[33], *Col15a1*[33], and *Cd34*[52] were enriched in

these two clusters (Supplementary figure **S6B**-**E**). Additionally, ECs in cyc-aEC and

trans-EC clusters expressed various capillary-like/venous specific genes such as

Gpihbp1, *Fabp3*, and *Pcdh17* (Supplementary figure **S6F-H**). All changes in

expression of *Cxcr4*, *Cx40*, *Aplnr*, *Car4*, *Apln* in cyc-aEC between sham and MI

groups, were found to be statistically significant using Wilcoxon sum rank test (**Table**

S1). Thus, cyc-aEC and trans-EC clusters are likely to possess angiogenic

properties, which are absent in stably differentiated artery ECs.

Cardiac vascular cells (ECs and SMCs) and cardiomyocytes have common

precursor cells. In addition to classic angiogenic endothelial cell markers, cyc-aEC

and trans-EC cells also expressed cardiac myofibrillar genes (*Tnnt2*, *Myl3* and

Myh6) (Supplementary figure **S6I-K**). Thus, cardiac ECs maintain low, but active

transcription of these cardiomyocyte specific markers as observed earlier[37].

The differentiation status of cells and the projected pseudotime, together, can

determine the root of differentiation, and hence can predict the direction of path

taken by cell populations[53]. A package in RStudio– Cellular (Cyto) Trajectory

Reconstruction Analysis using Gene counts and Expression (CytoTRACE), uses the

number of genes and their differential expression, to determine relative differentiation

status of each cell with respect to others[53]. CytoTRACE score for cycling aEC, in MI

dataset, was highest amongst all clusters, making it the least differentiated cluster

(Figure **2D**). We also performed pseudotime analysis of sham and MI cells using

monocle3 package in RStudio. In sham, the predicted trajectory connected trans-EC

to aEC1-4. Sham cyc-aECs were excluded from the projected path (Figure **2E**). In

contrast to sham, in MI, cyc-aEC, trans-EC and aEC1 clusters were associated with

the predicted path (Figure **2F**). If multiple clusters are connected through a single

trajectory, it is likely that, a less specialized cluster differentiates into a terminal cell

type. This prediction is supported by our observations from genetic lineage tracing

experiments *in vivo*, where upon MI, *Cx40CreER* lineage-labeled, proliferating artery

ECs give rise to mature coronary collateral arteries[24] and contribute to organ-wide

artery growth (Supplementary figure **S5F**).

We next assessed the relative differentiation status of the 3 clusters

participating in the trajectory obtained for MI cells (Figure **2F**)─ cyc-aEC, tans-EC

and aEC1. We compared the cellular and molecular features indicating de-

differentiation, across these 3 clusters (Figure **2G**). As expected, cyc-aEC and aEC1

were the least and most differentiated clusters, respectively (Figure **2G**). Trans-EC

cluster showed de-differentiation features intermediate between cyc-aEC and aEC1

(Figure **2G**). Thus, it is likely that proliferating (less specialized) artery ECs in cyc-

aEC cluster are differentiating into mature artery ECs in aEC1 (Figure **2F, G**). These

results were supported by RNA velocity analyses, which, additionally predicts

directionality of projected trajectories based on the ratio of spliced and un-spliced

transcripts (Supplementary figure **S7A**, **B**). Together, we hypothesized that upon

neonatal MI, at least a subpopulation of mature artery ECs originates from de-

differentiation and proliferation of pre-existing/ resident artery cells.

We checked if artery ECs, those enter cell cycle upon MI, (as seen in

our single cell analyses in Figure **1M**), downregulate Cx40 protein expression, in an

in vivo setting (Figure **3A**). We compared Cx40 protein levels in EdU$^+$ and EdU$^-$

artery ECs labelled with Cx40 immunostaining in *Cx40CreER; Rosa26^{TdTomato}*

neonates, 4 days post-MI (Figure **3B**-**D**). Both categories of artery cells– single and

associated with lumenized artery vessels, were analyzed. Single aECs observed in

ischemic watershed are derived from resident arteries and contribute to Artery

Reassembly. We observed that, Cx40 staining was significantly reduced in these

EdU$^+$ TdTomato$^+$ proliferative single aECs (Figure **3B**, **C**), when compared to non-

proliferative single aECs or artery tips (Figure **3B**). In contrast, EdU$^+$ proliferating

aECs, associated with well-formed arteries, did not show any significant change in

their Cx40 expression (Figure **3D**). Together, this suggests that Cx40 is

downregulated in proliferating single aECs, upon MI.

We further checked if MI-induced single artery ECs express early markers

such as *Apj*, which labels capillary/venous ECs from which arteries originate during

coronary development via arterialization. We used *Cx40$^{eGFP/+}$*; *ApjCreER*;

Rosa26^{TdTomato} mice to label arteries with GFP and capillary/venous ECs with

TdTomato upon Tamoxifen administration. Transgenic neonates were subjected to

Tamoxifen and MI at P2, to lineage label and trace *ApjCreER$^+$* cells (Figure **3E**).

Hearts were harvested at P6 when both single artery ECs and coronary collateral

arteries are present (Figure **3E**) and analyzed for Apj expression in pre-existing

artery ECs, upon MI. We identified single artery cells based on their location i.e.

proximity to distal branches of arteries spanning the ischemic watershed area and

their morphology i.e. single (not incorporated into any lumenized artery vessels) and

spindle shaped as observed earlier with *Cx40CreER*-lineage trace (Figure **3B**).

Indeed, many GFP+ single artery ECs expressed TdTomato, indicative of Apj

expression at 4 days post-MI (Figure **3F** and **G**). These GFP+ ECs were not

restricted to artery tips, but also observed in watershed regions (data not shown).

Thus, several artery ECs upon MI upregulate Apj expression indicating change in

cell fate in response to MI.

Together, our data suggests that single artery ECs exiting pre-existing artery

tips which are the drivers of Artery Reassembly, show reduced Cx40 expression and

increased Apj expression and proliferate during the course of neonatal Artery

Reassembly. All three features i.e. lack of mature cell marker, increase in early

angiogenic marker and cell cycle re-entry are indicative of de-differentiation. Thus,

we propose that pre-existing artery ECs de-differentiate and proliferate to build

coronary collateral arteries during Artery Reassembly (Figure **3H**).

Adult artery ECs fail to proliferate during MI-induced artery neogenesis

New collateral arteries in adult mice, form in response to coronary occlusion

within a week of injury[54]. However, adult coronary collateral arteries are fewer in

number, have smaller diameters, and are predicted to perfuse poorly than those in

neonates[55]. The reason for this age-dependent difference in which artery ECs

respond to ischemia is unclear. One possibility could be that adult collateral arteries

do not form naturally through Artery Reassembly. Keeping this in mind, we

investigated the molecular nature of adult coronary artery ECs, and their derivative

cell-types, in sham and MI mice.

Adult hearts execute Artery Reassembly only in response to exogeneous

Cxcl12. As a result, artery EC lineage traced coronary collateral arteries are

observed only after 14 days post-MI[24]. To capture single artery EC response that

precedes Artery Reassembly, we harvested adult hearts 9 days post-MI. We

performed single cell analyses of cardiac cells sorted from adult *Cx40CreER;*

Rosa26^TdTomato ventricles (Figure **4A**). Similar to neonatal data, we obtained both,

TdTomato$^+$ and some unlabeled cells. After performing quality control steps

(Supplementary figure **S8A**), we analyzed 6256 *Cadherin5$^+$* sham and 1690

Cadherin5$^+$ MI cells (Figure **4B**, **C**). Like neonatal data, our adult dataset was

enriched for artery ECs (Figure **4D**, **E**) and we obtained fewer *Cadherin5$^+$* (Figure

3C) and TdTomato$^+$ (Figure **4F**, **G**) cells in MI group than sham.

Based on cell-specific gene expression patterns, we obtained 12 adult cell

populations (Figure **4D**); 7 of which, constituted mature artery ECs (aEC1-7), 1

cycling artery EC (cyc-aEC), 1 capillary EC (cap-EC), 1 endoMT and 2 SMC clusters

(Figure **4D**, **E**). Unlike neonates, trans-EC cluster was absent in adults. Instead, we

identified a prominent adult capillary EC population, which when further clustered

generated 3 distinct microvascular EC populations—artery-capEC, capEC, and

venous-capEC (Supplementary figure **S8B**-**D**). A significant number of art-cap EC

cluster (34%) expressed TdTomato fluorescence (Supplementary figure **S8D**),

indicating an origin from pre-existing mature arteries. Identity of cycling aECs was

confirmed with comparable expression of artery specific genes (*Gja4*, *Cx40*, *Cxcr4*,

Jag1) in cyc-aEC and aEC1-7 (Supplementary figure **S8E**). A similar cell type

distribution was also observed when bioinformatics analyses of *Pecam1$^+$* cells,

instead of *Cadherin5$^+$* cells, was performed (Supplementary figure **S8F**-**H**).

Adult cardiomyocytes have limited proliferation capacity[23]. To assess if the

same holds true for cardiac ECs, we performed cell cycle analyses on sham and MI

adult *Cadherin5*[+] ECs. Unlike neonates, we observed only a minimal increase of

2.5% in the number of ECs undergoing G2 to M transition, upon MI (Figure **4H**).

Further cluster-wise analyses revealed only 1.8% increase in the mitotic (G2/M)

artery ECs (4.5% to 6.3%, Figure **4I**), compared to 8.5% increase observed in

neonates (Figure **1K**), upon MI. 4.6% of adult sham cells were mitotic (Figure **4I**),

suggesting that the pre-injury mitotic baseline for adult *Cadherin5*[+] artery ECs was

higher than that observed in neonates (2.5%, Figure **1K)**. Thus, in comparison to

neonates, fewer adult artery ECs undergo G2/M transition in response to MI.

P7 or older hearts are non-regenerative[22,23] and do not illicit Artery

Reassembly[24]. Cell cycle analyses of *Cadherin5*[+] TdTomato[+] lineage-traced adult

ECs revealed minimal changes in G2/M ECs including in artery EC clusters (aEC1-7,

cyc-aEC) (Figure **4J**). Furthermore, we checked which cell types originate from pre-

existing artery ECs, upon MI. For this, we analyzed all TdTomato[+] cells (labelled and

traced from *Cx40CreER*[+] resident artery ECs) in adult MI dataset, and compared it

with neonatal MI dataset (Figure **4K**). While the neonatal MI group showed a distinct

cyc-aEC cluster, this population was absent in the adult MI group (Figure **4K** and

Supplementary figure **S8I**, **J**). Thus, adult hearts lack a proliferating artery EC sub-

population, known to be crucial for Artery Reassembly, and complete cardiac

regeneration[24].

We next performed *in vivo* experiments to check if lack of proliferating artery

cell population in injured adult hearts (Figure **4J**, **K**), is also observed in non-

regenerative P7 hearts. *Cx40CreER; Rosa26*[TdTomato] neonates were administered

with Tamoxifen to lineage label artery ECs at P5, through P7, followed by MI at P7

(Figure **4L**). Proliferation status of *Cx40CreER*[+] lineage traced, TdTomato[+], artery

ECs were analyzed with EdU label, prior to harvesting hearts at P11, 4 days post-MI.

As shown earlier[24], these hearts did not demonstrate Artery Reassembly (Figure **4L**,

M), which was evident from lack of lineage traced (TdTomato[+]) collateral arteries and

single artery ECs in the injured watershed regions (Figure **4M**). Very few single

artery ECs were observed adjacent to artery tips (arrowheads, Figure **4M**); we called

them 'residual' single artery ECs. EdU[+] cells in TdTomato[+] artery EC population—in

both, artery tips (Figure **4N**) and residual single artery ECs in the watershed (Figure

4O)—were rarely observed. Together, data from both bioinformatics analyses and *in*

vivo experiments indicate that artery ECs in non-regenerative P7 or adult hearts

have limited self-renewing capacities.

Adult artery ECs do not de-differentiate, upon MI

We next checked if adult MI *Cadherin5*[+] cycling aECs (Figure **3I**) demonstrate

any characteristics indicative of de-differentiation. Though adult MI cycling aEC

cluster was the least differentiated aEC cluster (Figure **5A**, **B**), its CytoTRACE score

was comparable to a mature artery cluster—aEC5 in MI (Figure **5B**). Expression of

proangiogenic marker *Apln*, and its receptor *Aplnr* were seemingly upregulated in

this cluster, but there was no change in expression of *Car4*, *Cx40*, or *Cxcr4* (Figure

5C). We analyzed the cellular and molecular features of neonatal cyc-aECs, which

were indicative of de-differentiation, and checked those indicators in adult cyc-aECs

(Figure **5D**). Some of the key de-differentiation features absent in adult cyc-aECs

were—1. their adequate number, which is representative of their ability to self-renew/

re-enter cell cycle 2. loss of mature cell markers and 3. gain of microvascular/ pro-

angiogenic markers (shown in red, Figure **5D**). For instance, in contrast to neonatal

MI cyc-aECs, adult MI cyc-aECs, showed almost no change in the number of *Cx40*

expressing cells (Figure **5D**). Adult cells expressing *Cxcr4* in this group, increased by

1.76 fold (Figure **5D**). The number of cells expressing *Apln*, *Aplnr*, and *Car4* in the

adult cyc-aECs were increased upon MI. However, these changes in cell numbers

or gene expression levels, were statistically insignificant when compared between

adult sham and MI groups, using Wilcoxon sum rank test (* in Figure **5D and Table**

S1). Furthermore, cycling aEC cluster did not contribute to the predicted path

associated with mature aECs (aEC1-7), even in MI group (Figure **5E**, **F**). Thus,

unlike neonate, adult cycling aECs were neither connected to artery ECs in trajectory

analyses (Figure **5E**, **F**), nor were lineage traced from pre-existing arteries (Figure

4J, **K**). Together, adult cycling aEC population fails to drive artery regeneration

(Figure **5G**) due to its limited number and inability to de-differentiate (Figure **5D**).

Arterial VegfR2 promotes artery EC proliferation during Artery Reassembly

Pre-existing artery ECs drive collateral artery formation in neonatal mice, only

upon MI[24,56]. We isolated all genes from *Cx40CreER*-lineage traced TdTomato[+]

injured artery ECs (Figure **1F**) and performed gene ontology enrichment analysis

and visualization (GOrilla). This allowed us to identify biological processes triggered

upon MI. Key vascular responses such as angiogenesis, migration and proliferation,

were significantly upregulated upon MI (Figure **6A**). Artery Reassembly is a

systematic step-wise cellular event where artery EC migration and proliferation are

crucial. Thus, our neonatal single cell dataset contains the artery EC population,

including those that drive Artery Reassembly, and subsequent neonatal cardiac

regeneration.

aEC1, in neonatal MI dataset, is the only artery EC cluster associated with the

predicted trajectory and connected with cycling artery ECs (Figure **2F**). We

performed pathway analyses on *Cadherin5*[+] aEC1 cells using g:Profiler and filtered

pathways from various databases, based on their statistical significance (p>0.05).

Many pathways relevant to vascular development and injury response were enriched

in aEC1 such as Apelin, TNF and Interleukin signaling[57] (Figure **6B**). Vegfa-VegfR2

signaling (p=0.0011) was of particular interest (Figure **6B**), as it has established

molecular functions in mouse coronary artery development[40,58] and

neovascularization upon injury in different tissues[59,60].

We checked the expression levels of ligands (*Vegfa*, *Vegfc*) and a major

receptor (*VegfR2* or *Kdr*) associated with coronary morphogenesis, in our dataset.

While *Vegfa* and *Vegfc* were expressed in artery ECs (aEC1-4), *VegfR2* was

specifically enriched in cycling aEC and trans-EC population (Figure **6C**). The

VegfR2 expression, was significantly different in cyc-aECs, when compared between

neonatal sham and MI groups, using Wilcoxon sum rank test (**Table S1**). When

immunostained for VegfR2, the protein was located in coronary artery ECs of small

arteries, artery tips and capillary ECs (Supplementary figure **S9A**). The levels of

VegfR2 protein in arteries was lower as compared to capillaries, and the protein was

almost always absent in large arteries (Supplementary figure **S9A**). Both EC types

(arteries and capillaries) facilitate Artery Reassembly, either by directly participating

in the process (artery ECs) or providing the Cxcl12 ligand (capillary ECs). Given

VegfR2 expression in both, capillaries and small arteries, we hypothesized that Vegf

pathway has a role in Artery Reassembly, post-MI.

We analyzed the role of arterial *VegfR2* in Artery Reassembly, *in vivo*. Control

(*Cx40CreER*; *Rosa26^{TdTomato}*) and artery EC-specific, conditional *VegfR2* knockout

(*VegfR2^{L/L}*; *Cx40CreER*; *Rosa26^{TdTomato}*) neonates received Tamoxifen at P0/P1

which allowed labelling of pre-existing artery ECs. These hearts were subjected to

MI at P2/P3 and hearts were harvested 4 days post-MI, at P6/P7 (Figure **6D**). Whole

heart imaging allowed us to visualize the entire arterial network of the heart (Figure

6E, **F**) as well as focus into regions of interest such as the watershed region where

new collaterals are expected to appear after injury (Figure **6G-I**). Control and *VegfR2*

artery specific knockout hearts did not illustrate any gross differences in their

baseline artery architecture, without MI (Supplementary figure **S9B**-**D**). Watershed

regions from sham (data not shown) or no injury (Figure **6G**) hearts, did not show

any collateral arteries lineage labelled with TdTomato, suggesting absence of Artery

Reassembly in non-ischemic hearts. Upon MI, control hearts consistently showed

several (average of 8) *Cx40CreER*-lineage traced (TdTomato[+]) collateral arteries as

observed earlier[24] (Figure **6E**, **H**). In contrast, we rarely observed any such collateral

arteries in *VegfR2* artery specific knockout hearts (Figure **6F**, **I**, **J**). This suggests an

artery specific role of VegfR2 in executing Artery Reassembly, upon MI.

We next investigated the cause for lack of TdTomato[+] collateral arteries in

VegfR2 knockout watershed regions. Artery Reassembly is a three step cellular

event—migration, proliferation and coalescence, of artery ECs. We first assessed

the ability of TdTomato[+] cells to migrate into the watershed. Qualitative and

quantitative observations suggested that the number of *Cx40CreER*-lineage traced

single artery ECs, observed in the watershed regions, were comparable in control

and knockout MI hearts (Figure **6H**, **I** and Supplementary figure **S9E-H**). This

suggests that in absence of arterial VegfR2, the migration of artery ECs from their

resident arteries into the watershed region is less likely affected. We next

investigated the proliferation step, and injected neonates with EdU, one hour prior to

harvesting the hearts. We then, quantified the number of proliferating single aECs in

the MI watershed regions. Interestingly, *VegfR2* knockout watershed regions showed

a ~2.4-fold reduction in the number of proliferating single artery ECs as compared to

control hearts (Figure **6K**-**M**). Such reduction in number of proliferating single artery

ECs was also observed in non-regenerative P7 injured hearts (Figure **4O**). Thus,

arterial VegfR2 regulates neonatal artery EC proliferation, post-MI. A defect in

VegfR2-mediated artery cell proliferation could also affect subsequent coalescence

of TdTomato$^+$ cells and successful execution of Artery Reassembly, as observed in

VegfR2 knockout hearts (Figure **6J**).

Functional coronary collateral arteries built by Artery Reassembly promote

cardiac regeneration. Since arterial deletion of *VegfR2* reduced number of collateral

arteries, we tested if this neonatal *VegfR2* deletion affects cardiac function in adults.

We performed MIs a neonatal stage (P2) and checked several parameters for

cardiac function 30 days post-MI. While parameters like Ejection Fraction and

Fractional Shortening did not change significantly (Figure **6N**, **O**), cardiac output and

stroke volume were significantly reduced (Figure **6P**, **Q**) in *VegfR2* arterial EC

specific knockouts. Thus, arterial VegfR2 facilitates neonatal Artery Reassembly

which regulates cardiac function in adult mice.

We next investigated if Vegf ligands and receptors are expressed in adult

ECs. The expression of Vegf ligands, *Vegfa* and *Vegfc*, were mostly observed in

mature artery EC clusters (Supplementary figure **S10A**) like in neonatal data (Figure

6C). The receptor *VegfR2* (*Kdr*), was upregulated in capillary ECs, as expected

(Supplementary figure **S10A**). Interestingly, at least at the mRNA level, *VegfR2* was

downregulated in ~75% of actively cycling aECs (Supplementary figure **S10A**). This

is in contrast to the upregulated expression of *VegfR2* in neonatal cyc-aECs (Figure

6C). When compared to sham group, *VegfR2* was significantly downregulated in

adult MI cyc-aEC cluster (**Table S1**). Given the functional significance of arterial

VegfR2 in neonatal aEC proliferation (Figure **6K**-**M**), the lack of *VegfR2* expression

in "residual" adult cycling aECs, could explain depleted potential for aEC proliferation

in injured adults. In summary, our data indicate that arterial VegfR2 drives,

specifically, the proliferation step of Artery Reassembly in injured neonatal mice

(Supplementary figure **S10B**).

Discussion

In this study we utilize both *in silico* analyses and *in vivo* experiments to

explain the age-dependent artery EC responses to cardiac injury. Firstly, with single

cell sequencing data analyses we show, for the first time, that coronary artery ECs,

an otherwise stably differentiated cardiac cell type, undergoes injury-induced de-

differentiation in neonates, but not in adults. In this study, we defined artery EC de-

differentiation as a biological process involving four major cellular/molecular features

1. downregulation mature artery EC cell markers, 2. expression of early and

angiogenic markers, 3. cell cycle re-entry and 4. Subsequent differentiation into more

artery ECs to build new collateral arteries by Artery Reassembly. With the

combination of bioinformatics and mouse experiments, we show that the ability of

artery ECs to de-differentiate, correlates with their ability to proliferate. Both cellular

responses —artery cell de-differentiation and proliferation— are specific to neonatal

regenerative window. Secondly, using *in vivo* experiments, we discover a second

artery-specific receptor (apart from Cxcr4, which controls artery cell migration during

Artery Reassembly[24]), which regulates proliferation of artery ECs in response to

neonatal MI. Expression of this receptor, *VegfR2*, is downregulated in adult cycling

artery ECs and could explain absence of Artery Reassembly in adults. Finally, our

data suggests that each step of Artery Reassembly is regulated by a distinct

signaling axis, and the successful execution of each step will require activation of

these molecular pathways in the right space and time. In summary, we highlight the

differences in age-dependent molecular profiles of mouse coronaries and propose a

mechanism for reduced plasticity of artery ECs and lack of Artery Reassembly in

non-regenerative P7 or adult hearts.

We used single cell RNA sequencing to investigate MI-induced artery EC

response in neonates and adults. We obtained fewer artery ECs upon MI as

compared to sham. This was observed in both neonates and adults. In our dataset,

the time points when the neonatal or adult hearts were harvested represent a state

when cardiac regeneration has not been completed yet. At the chosen timepoints,

while many vascular regenerative processes have been initiated, it is likely that cell

death would still be in progression. This phenomenon is also reflected in

Cx40CreER-lineage traced TdTomato artery ECs. Such small fraction of artery ECs

observed in our dataset have also been reported in other neonatal and adult[27]

datasets[29,30] and likely represent extensive cell death following coronary ligation and

a vascular environment prior to complete cardiac recovery.

TdTomato cells in our MI datasets were ~17 fold less in neonates and ~5 fold

less in adults as compared to their sham counterparts. We hence analyzed all ECs

which included both TdTomato+ and TdTomato- cells. The analyses of *Cadherin5+*

all ECs allowed us to build several hypotheses which were tested in vivo. Using

genetic lineage tracing and confocal imaging of whole hearts, we show that post-MI,

1) neonatal artery ECs also originate from sources other than pre-existing artery

ECs. 2) During Artery Reassembly cycling pre-existing artery ECs downregulate

mature artery EC marker (*Cx40*) and upregulate marker for capillary/vein (*Apj*) from

which artery ECs originate during development 3) artery EC proliferation is minimal

in non-regenerative P7 hearts when subjected to MI and finally, 4) injury induced

artery EC proliferation is regulated by VegfR2 mediated pathway which subsequently

impacts cardiac function of adult mice. These in vivo results were based on the

findings from our single cell dataset. Thus, despite of their low numbers in our single

cell datasets, the artery EC population is representative of post-injury vascular niche

in vivo.

De-differentiation and Proliferation are Coupled during Cardiac Regeneration

New cardiomyocytes originate from pre-existing cardiomyocytes upon injury[22,23]

by de-differentiating and re-entering cell cycle[61,62]. Cardiomyocyte de-differentiation

has been elucidated upon adult zebrafish cardiac injury via disassembly of

sarcomeric structures, detachment of cardiomyocytes from one another, and

expression of early embryonic genes (α-Smooth Muscle Actin (αSMA), embryonic

Myosin Heavy Chain, and Gata4)[61–63]. Post-MI adult cardiomyocytes also

demonstrate somewhat limited de-differentiation potential indicated by

downregulation of mature cardiomyocyte functions such as hypertrophy, contractility,

electrical conduction, and upregulation of several early genes like *αSMA*, *Gata4*,

Runx1 and *Dab2*[64]. Unlike cardiomyocytes, EC de-differentiation has not been

reported in any cardiac injury model. We show here, for the first time, that neonatal

mouse coronary artery ECs undergo de-differentiation upon MI; a process limited in

adults.

Since there are no objective indicators of artery EC pluripotency, one can argue

that changes in gene expression or phenotypes could be a reflection of their

plasticity, and not necessarily reversion of cell fate into a de-differentiated state. It is

also possible that number of de-differentiated cells observed in cyc-aEC and

transEC clusters of uninjured hearts simply increase upon MI. To address this, we

combine single cell analyses and genetic lineage tracing. Lineage labelled non-

cycling artery ECs in MI hearts can be sequentially traced into cycling single artery

ECs in the watershed, and consequently into non-cycling artery ECs on the coronary

collaterals. We analyzed the transiently cycling artery ECs and show that several

indicators for de-differentiation, were enhanced in neonates. One such indicator was

presence of Apj lineage in Cx40$^+$ artery ECs. It is possible that Apj-lineage traced

capillaries differentiate into artery ECs, a phenomenon called arterialization already

shown to contribute (though minimally) to collateral formation by Artery

Reassembly[24]. Our in vivo data cannot distinguish between differentiation of capillary

ECs into artery ECs and upregulation of capillary marker (Apj) in arteries. However,

the unique location and morphology of cells allowed us to confidently identify Cx40$^+$

single artery ECs and conclude that at least a sub-population of artery ECs, upon MI,

express Apj—an indicative of vascular de-differentiation.

The de-differentiation indicators were absent in adults. This finding positively

correlated with the ability of artery ECs to re-enter cell cycle. Neonates had much

higher number of cycling artery ECs than P7 or adults. This finding was verified via

two independent methods—single cell gene expression analyses of cyc-aECs and

presence of *Cx40CreER*$^+$ EdU$^+$ cells in an *in vivo* cardiac MI model. Thus, lack of de-

differentiation limits the number of cycling aECs in non-regenerative hearts and

consequently affects their ability to form collateral arteries by Artery Reassembly.

Reactivation of Developmental Pathways upon Injury?

Several molecular pathways relevant to vascular development regulate EC

response to injury[65]. For instance, Notch[66], Apelin[45], Cxcl12[67–70] signaling, in mice,

regulate different aspects of vascular development and remodeling during cardiac

morphogenesis; ranging from EC fate determination to regulation of different EC

responses such as migration, proliferation and tube formation. Interestingly, some of

these pathways are also associated with EC regeneration upon injury[1,24,48], making

them potential therapeutic candidates for treating vascular conditions in human

patients[71]. Further studies are required, and will elucidate if (and how) the cellular

and molecular mechanisms associated with these signaling pathways, during

development and injury responses, are similar.

Vegf signaling drives coronary development in zebrafish and mouse hearts. In

mice, Vegf signaling drives migration of coronary endothelial[43,72] and endocardial

endothelial cells[58] in developing hearts, and proliferation of coronary ECs in heart

explant cultures[73]. In zebrafish, *vegfa* mutants show reduced EC proliferation

possibly via PI3K signaling[74]. Upon cardiac injury, endocardial Vegfa signaling[1] and

Vegfc-mediated (Cxcl8a/Cxcr1-driven) coronary EC proliferation[75] execute

revascularization and cardiac regeneration of ischemic hearts. In this study, we

highlight the role of VegfR2-mediated signaling in artery EC response triggered by

neonatal cardiac injury. We show that VegfR2 in neonates is expressed in coronary

artery ECs, deletion of which, significantly reduced the number of collateral arteries

formed via Artery Reassembly.

Our data show that artery tips do not proliferate. Proliferation is rather

demonstrated by single aECs which originate from these artery tips. This suggests

that migration of pre-existing artery ECs could precede artery EC proliferation in the

watershed. Appearance of *VegfR2* depleted single artery ECs in ischemic watershed

regions suggested that artery EC migration was not affected. VegfR2 depleted single

artery ECs showed reduced proliferation potential, which could account for lack of

lineage-traced collateral arteries in the knock out hearts. Proliferation of artery ECs is

followed by their coalescence into collateral arteries. Thus, we cannot exclude the

possible role of Vegf in coalescence of artery cells, or the vascular steps that follow,

like lumenization or maturation of collateral arteries.

The precise source of Vegf ligand in injured mouse hearts could be the

endocardium[76], myocardium (Vegfa)[58,77], epicardium (Vegfc[43], or Vegfa[78]), or cardiac

endothelial cells[75], thus suggesting an autocrine or paracrine signaling as observed

in injured zebrafish hearts[75]. In our neonatal mouse single cell dataset, we found

both *Vegfa* and *Vegfc* being expressed by mature artery ECs (aEC1-4), but at lower

levels by cyc-aECs (or by trans-ECs, SMCs, endoMT ECs, endo ECs), indicating a

possible cell autonomous mechanism. However, an alternate source of Vegf ligand,

such as, cardiomyocytes, fibroblasts or immune cells, which are excluded from our

dataset, cannot be neglected.

Role of Vegf Pathway in Human Heart Disease and its Treatment

Single nucleotide polymorphisms (SNPs) in both introns and exons of *KDR*

(murine *VegfR2*) are strongly associated with coronary heart disease (CHD)[79]. These

SNPs can either modulate the affinity to ligand, or alter E2F binding to *KDR*

promoter, subsequently, affecting *KDR* activity and downstream signaling[79]. Elevated

levels of circulating VEGF ligands are positively and significantly associated with

CHD[80], coronary artery disease (CAD)[81,82] and coronary artery lesions[83,84]. Several

SNPs affect VEGF levels[80,82] and may account for the variation observed in

revascularization abilities in individuals across different genetic backgrounds. At

least three such SNPs in *VEGFA* are strongly associated with increased risk of

CHD/MI[85] and one with coronary collateral vessel perfusion[86]. How the association of

VEGF/KDR signaling axis with CHD may serve for better prognosis or indicate the

need for vascularization, remains unclear.

Preclinical studies in animals with sparse[87] to no[54] pre-existing collateral

arteries have consistently elucidated role of Vegf in ischemia-induced collateral

formation. On the contrary, numerous past and ongoing clinical trials with *VEGF*

alone, or combined with pro-angiogenic/ regenerative factors, have been concluded

with little to no success (data available at clinicaltrials.gov)[87,88]. The delivery methods

appear to be safe, but the angiogenic effects, particularly the efficacy of *VEGF* gene

therapies in stimulating reperfusion via collateral arteries is minimal[87]. Mechanistic

details of Vegf's mode of action, in orchestrating vascular cell responses that lead to

collateral formation, could aid designing clinical trials with higher efficacies.

We show that mouse coronary ECs in MI hearts, de-differentiate and then

expand in VegfR2-dependent manner. The mere presence of a significant number of

single artery ECs in a limited watershed area could be sufficient to create new

arteries[31]. Thus, it will be interesting to see if this threshold density of artery ECs is

accomplished by Vegf-mediated proliferation. Similarly, identifying the molecular

drivers of EC de-differentiation is also crucial, and when combined with Vegf, could

promote collaterals in individuals with CHD and CAD.

Authors Contribution

SD and KR conceived the idea. SD, KR, YJW provided resources. GA performed

experiments with neonatal MI model. Sneha K and Suraj K performed bioinformatics

analyses. SD, GA and Sneha K wrote the manuscript. HW performed adult MI

surgeries. PERC and BB performed some neonatal MI surgeries. SD and KMG

performed FACS.

Acknowledgements

We thank the following facilities for their technical support: Stanford Genome

Sequencing Center, Stanford shared FACS facility, Animal Care and Resource

Center at NCBS, Mouse Genome Engineering Facility at NCBS, Central Imaging and

Flow Cytometry Facility at NCBS, Dr. Dhandapany Perundurai's lab from Institute

For Stem Cell Science and Regenerative Medicine (InStem), for their technical help

with echocardiography.

Funding

This work is supported by NCBS/TIFR core funding and DBT/Wellcome Trust India

Alliance Intermediate Fellowship to SD (IA/I/20/2/505205). YJW is supported by

National Institutes of Health (5R01HL089315-11). KR is supported by National

Institutes of Health (R01-HL128503) and is a HHMI Investigator. HW is supported by

American Heart Association (18POST33990223). KMG is supported by Gabilan

Stanford Graduate Fellowship, NSF GRFP. Suraj K is supported by DBT/Wellcome

Trust India Alliance Intermediate Fellowship to SD. PERC is supported by the

NIGMS of the National Institutes of 1121 Health (NIH T32GM007276) and NSF-

GRFP (DGE-1656518).

Declaration of interests

None

Data availability:

Raw sequence files and Cell Ranger processed data are available on GEO (

accession ID: GSE210307)

Scripts used for the analysis are available at

https://github.com/Snehasrivatsa/scRNA

Bibliography

1. Marín-Juez, R. *et al.* Fast revascularization of the injured area is essential to support zebrafish heart regeneration. *Proc. Natl. Acad. Sci. U. S. A.* **113**, 11237–11242 (2016).

2. Ingason, A. B. *et al.* Angiogenesis Precedes Cardiomyocyte Migration in Regenerating Mammalian Hearts. *J. Thorac. Cardiovasc. Surg.* **155**, 1118-1127.e1 (2017).

3. He, L. *et al.* Preexisting endothelial cells mediate cardiac neovascularization after injury. *J. Clin. Invest.* **127**, 2968–2981 (2017).

4. Li, Z. *et al.* Single-cell transcriptome analyses reveal novel targets modulating cardiac neovascularization by resident endothelial cells following myocardial infarction. *Eur. Heart J.* **40**, 2507–2520 (2019).

5. Kobayashi, K. *et al.* Dynamics of angiogenesis in ischemic areas of the infarcted heart. *Sci. Rep.* **7**, 1–13 (2017).

6. Betgem, R. P. *et al.* Intramyocardial haemorrhage after acute myocardial infarction. *Nat. Rev. Cardiol.* **12**, 156–167 (2015).

7. Buschmann, I. & Schaper, W. Arteriogenesis versus angiogenesis: Two mechanisms of vessel growth. *News Physiol. Sci.* **14**, 121–125 (1999).

8. Seiler, C., Stoller, M., Pitt, B. & Meier, P. The human coronary collateral circulation: Development and clinical importance. *Eur. Heart J.* **34**, 2674–2682 (2013).

9. Sabia, P. J. *et al.* An Association between Collateral Blood Flow and Myocardial Viability in Patients with Recent Myocardial Infarction. *N. Engl. J. Med.* **327**, 1825–1831 (1992).

10. Meier, P. *et al.* Beneficial effect of recruitable collaterals: A 10-year follow-up

study in patients with stable coronary artery disease undergoing quantitative

collateral measurements. *Circulation* **116**, 975–983 (2007).

11. Habib, G. B. *et al.* Influence of coronary collateral vessels on myocardial infarct

size in humans. Results of phase I thrombolysis in myocardial infarction (TIMI)

trial. The TIMI Investigators. *Circulation* **83**, 739–746 (1991).

12. Clayton, J. A., Zhang, H., Pomp, D., Faber, J. E. & Chalothorn, D. Collateral

density, remodeling, and VEGF-A expression differ widely between mouse

strains. *Physiol. Genomics* **30**, 179–191 (2007).

13. Teunissen, P. F. A., Horrevoets, A. J. G. & van Royen, N. The coronary

collateral circulation: Genetic and environmental determinants in experimental

models and humans. *J. Mol. Cell. Cardiol.* **52**, 897–904 (2012).

14. Maxwell, M. P., Hearse, D. J. & Yellon, D. M. Species variation in the coronary

collateral circulation during regional myocardial ischaemia: A critical

determinant of the rate of evolution and extent of myocardial infarction.

Cardiovasc. Res. **21**, 737–746 (1987).

15. Chalothorn, D. & Faber, J. E. Formation and maturation of the native cerebral

collateral circulation. *J. Mol. Cell. Cardiol.* **49**, 251–259 (2010).

16. Perovic, T., Harms, C. & Gerhardt, H. Formation and Maintenance of the

Natural Bypass Vessels of the Brain. *Front. Cardiovasc. Med.* **9**, 1–6 (2022).

17. Gabhann, F. Mac & Peirce, S. M. Collateral capillary arterialization following

arteriolar ligation in murine skeletal muscle. *Microcirculation* **17**, 333–347

(2010).

18. Scholz, D. *et al.* Contribution of arteriogenesis and angiogenesis to

postocclusive hindlimb perfusion in mice. *J. Mol. Cell. Cardiol.* **34**, 775–787

(2002).

19.   Zhang, H., Prabhakar, P., Sealock, R. & Faber, J. E. Wide genetic variation in

the native pial collateral circulation is a major determinant of variation in

severity of stroke. *J. Cereb. Blood Flow Metab.* **30**, 923–934 (2010).

20.   Chalothorn, D. & Faber, J. E. Strain-dependent variation in collateral

circulatory function in mouse hindlimb. *Physiol. Genomics* **42**, 469–479 (2010).

21.   Clayton, J. A., Chalothorn, D. & Faber, J. E. Vascular endothelial growth

factor-a specifies formation of native collaterals and regulates collateral growth

in ischemia. *Circ. Res.* **103**, 1027–1036 (2008).

22.   Porrello, E. R. *et al.* Transient regenerative potential of the neonatal mouse

heart. *Science (80-.).* **331**, 1078–1080 (2011).

23.   Porrello, E. R. *et al.* Regulation of neonatal and adult mammalian heart

regeneration by the miR-15 family. *Proc. Natl. Acad. Sci.* **110**, 187–192 (2013).

24.   Das, S. *et al.* A Unique Collateral Artery Development Program Promotes

Neonatal Heart Regeneration. *Cell* **176**, 1128-1142.e18 (2019).

25.   Aurora, A. B. *et al.* Macrophages are required for neonatal heart regeneration

Find the latest version : Macrophages are required for neonatal heart

regeneration. **124**, 1382–1392 (2014).

26.   Wang, Y. *et al.* Single-cell analysis of murine fibroblasts identifies neonatal to

adult switching that regulates cardiomyocyte maturation. *Nat. Commun.* **11**,

(2020).

27.   Wang, Z. *et al.* Cell-Type-Specific Gene Regulatory Networks Underlying

Murine Neonatal Heart Regeneration at Single-Cell Resolution. *Cell Rep.* **33**,

108472 (2020).

28.   Hao, Y. *et al.* Integrated analysis of multimodal single-cell data. *Cell* **184**,

3573-3587.e29 (2021).

29. Farbehi, N. *et al.* Single-cell expression profiling reveals dynamic flux of

cardiac stromal, vascular and immune cells in health and injury. *Elife* **8**, (2019).

30. Zhuang, L., Lu, L., Zhang, R., Chen, K. & Yan, X. Comprehensive Integration

of Single-Cell Transcriptional Profiling Reveals the Heterogeneities of Non-

cardiomyocytes in Healthy and Ischemic Hearts. *Front. Cardiovasc. Med.* **7**,

(2020).

31. Raftrey, B. *et al.* Dach1 Extends Artery Networks and Protects Against Cardiac

Injury. *Circ. Res.* 702–716 (2021) doi:10.1161/circresaha.120.318271.

32. Phansalkar, R. *et al.* Coronary blood vessels from distinct origins converge to

equivalent states during mouse and human development. *Elife* **10**, 1–32

(2021).

33. Kalucka, J. *et al.* Single-Cell Transcriptome Atlas of Murine Endothelial Cells.

Cell **180**, 764-779.e20 (2020).

34. Zhang, H. *et al.* Endocardium Minimally Contributes to Coronary Endothelium

in the Embryonic Ventricular Free Walls. *Circ. Res.* **118**, 1880–1893 (2016).

35. Conway, S. & Molkentin, J. Periostin as a Heterofunctional Regulator of

Cardiac Development and Disease. *Curr. Genomics* **9**, 548–555 (2008).

36. Torregrosa-Carrión, R. *et al.* Adhesion G protein-coupled receptor

Gpr126/Adgrg6 is essential for placental development. *Sci. Adv.* **7**, (2021).

37. Yucel, N. *et al.* Cardiac endothelial cells maintain open chromatin and

expression of cardiomyocyte myofibrillar genes. *Elife* **9**, (2020).

38. Su, T. *et al.* Single-cell analysis of early progenitor cells that build coronary

arteries. *Nature* **559**, 356–362 (2018).

39. Fang, J. S. *et al.* Shear-induced Notch-Cx37-p27 axis arrests endothelial cell

cycle to enable arterial specification. *Nat. Commun.* **8**, (2017).

40. Luo, W. *et al.* Arterialization requires the timely suppression of cell growth. *Nature* **589**, 437–441 (2021).

41. Yao, Y. & Wang, C. Dedifferentiation: inspiration for devising engineering strategies for regenerative medicine. *npj Regen. Med.* **5**, (2020).

42. Sharma, B., Chang, A. & Red-Horse, K. Coronary Artery Development: Progenitor Cells and Differentiation Pathways. *Annu. Rev. Physiol.* **79**, 1–19 (2017).

43. Chen, H. I. *et al.* The sinus venosus contributes to coronary vasculature through VEGFC-stimulated angiogenesis. *Development* **141**, 4500–4512 (2014).

44. Gillich, A. *et al.* Capillary cell-type specialization in the alveolus. *Nature* **586**, 785–789 (2020).

45. Helker, C. S. M. *et al.* Apelin signaling drives vascular endothelial cells towards a pro-angiogenic state. *Elife* **9**, 1–44 (2020).

46. Tian, X. *et al.* Peritruncal coronary endothelial cells contribute to proximal coronary artery stems and their aortic orifices in the mouse heart. *PLoS One* **8**, (2013).

47. Eyries, M. *et al.* Hypoxia-induced apelin expression regulates endothelial cell proliferation and regenerative angiogenesis. *Circ. Res.* **103**, 432–440 (2008).

48. Masoud, A. G. *et al.* Apelin directs endothelial cell differentiation and vascular repair following immune-mediated injury. *J. Clin. Invest.* **130**, 94–107 (2020).

49. Godoy, R. S. *et al.* Novel Apelin-expressing gCap Endothelial Stem-like Cells Orchestrate Lung Microvascular Repair. *bioRxiv* 2021.07.12.452061 (2022) doi:10.1101/2021.07.12.452061.

50.   Mirza, M. K. *et al.* FoxM1 regulates re-annealing of endothelial adherens

junctions through transcriptional control of β-catenin expression. *J. Exp. Med.*

207, 1675–1685 (2010).

51.   Zhao, Y. Y. *et al.* Endothelial cell-restricted disruption of FoxM1 impairs

endothelial repair following LPS-induced vascular injury. *J. Clin. Invest.* **116**,

2333–2343 (2006).

52.   Siemerink, M. J. *et al.* CD34 marks angiogenic tip cells in human vascular

endothelial cell cultures. *Angiogenesis* **15**, 151–163 (2012).

53.   Gulati, G. S. *et al.* Single-cell transcriptional diversity is a hallmark of

developmental potential. *bioRxiv* **411**, 405–411 (2019).

54.   Zhang, H. & Faber, J. E. De-novo collateral formation following acute

myocardial infarction: Dependence on CCR2+bone marrow cells. *J. Mol. Cell.*

Cardiol. **87**, 4–16 (2015).

55.   Anbazhakan, S. *et al.* Blood flow modeling reveals improved collateral artery

performance during mammalian heart regeneration. *bioRxiv*

2021.09.17.460699 (2021).

56.   He, L. *et al.* Genetic lineage tracing discloses arteriogenesis as the main

mechanism for collateral growth in the mouse heart. *Cardiovasc. Res.* **109**,

419–430 (2016).

57.   Zhou, D. *et al.* Effects of IL-1β and TNF-α on the Expression of P311 in

Vascular Endothelial Cells and Wound Healing in Mice. *Front. Physiol.* **11**,

(2020).

58.   Wu, B. *et al.* Endocardial cells form the coronary arteries by angiogenesis

through myocardial-endocardial VEGF signaling. *Cell* **151**, 1083–1096 (2012).

59.   Koh, B. I. *et al.* VEGFR2 signaling drives meningeal vascular regeneration

upon head injury. *Nat. Commun.* **11**, (2020).

60.   Isner, J. M. *et al.* Clinical evidence of angiogenesis after arterial gene transfer

of phVEGF165 in patient with ischaemic limb. *Lancet* **348**, 370–374 (1996).

61.   Kikuchi, K. *et al.* Primary contribution to zebrafish heart regeneration by gata4+

cardiomyocytes. *Nature* **464**, 601–605 (2010).

62.   Jopling, C. *et al.* Zebrafish heart regeneration occurs by cardiomyocyte

dedifferentiation and proliferation. *Nature* **464**, 606–609 (2010).

63.   Bertozzi, A., Wu, C. C., Hans, S., Brand, M. & Weidinger, G. Wnt/β-catenin

signaling acts cell-autonomously to promote cardiomyocyte regeneration in the

zebrafish heart. *Dev. Biol.* **481**, 226–237 (2022).

64.   Zhang, Y. *et al.* Single-cell imaging and transcriptomic analyses of

endogenous cardiomyocyte dedifferentiation and cycling. *Cell Discov.* **5**,

(2019).

65.   Lowe, V., Wisniewski, L. & Pellet-Many, C. The zebrafish cardiac endothelial

cell—roles in development and regeneration. *J. Cardiovasc. Dev. Dis.* **8**,

(2021).

66.   Fernández-Chacón, M., García-González, I., Mühleder, S. & Benedito, R. Role

of Notch in endothelial biology. *Angiogenesis* **24**, 237–250 (2021).

67.   Phansalkar, R. *et al.* Endocardium-to-coronary artery differentiation during

heart development and regeneration involves sequential roles of Bmp2 and

Cxcl12/Cxcr4. (2021).

68.   Ivins, S. *et al.* The CXCL12/CXCR4 Axis Plays a Critical Role in Coronary

Artery Development. *Dev. Cell* **33**, 455–468 (2015).

69.   Cavallero, S. *et al.* CXCL12 Signaling is Essential for Maturation of the

Ventricular Coronary Endothelial Plexus and Establishment of Functional

945   Coronary Circulation. *Dev. Cell* **33**, 469–477 (2015).

946 70. Chang, A. H. *et al.* DACH1 stimulates shear stress-guided endothelial cell

947   migration and coronary artery growth through the CXCL12-CXCR4 signaling

948   axis. *Genes Dev.* **31**, 1308–1324 (2017).

949 71. Evans, C. E., Iruela-Arispe, M. L. & Zhao, Y. Y. Mechanisms of Endothelial

950   Regeneration and Vascular Repair and Their Application to Regenerative

951   Medicine. *Am. J. Pathol.* **191**, 52–65 (2021).

952 72. Red-Horse, K., Ueno, H., Weissman, I. L. & Krasnow, M. A. Coronary arteries

953   form by developmental reprogramming of venous cells. *Nature* **464**, 549–553

954   (2010).

955 73. Rhee, S. *et al.* Endothelial deletion of Ino80 disrupts coronary angiogenesis

956   and causes congenital heart disease. *Nat. Commun.* **9**, (2018).

957 74. Lange, M. *et al.* Zebrafish mutants in vegfab can affect endothelial cell

958   proliferation without altering ERK phosphorylation and are phenocopied by

959   loss of PI3K signaling. *Dev. Biol.* **486**, 26–43 (2022).

960 75. El-Sammak, H. *et al.* A Vegfc-Emilin2a-Cxcl8a Signaling Axis Required for

961   Zebrafish Cardiac Regeneration. *Circ. Res.* 1014–1029 (2022)

962   doi:10.1161/circresaha.121.319929.

963 76. Marín-Juez, R. *et al.* Coronary Revascularization During Heart Regeneration Is

964   Regulated by Epicardial and Endocardial Cues and Forms a Scaffold for

965   Cardiomyocyte Repopulation. *Dev. Cell* **51**, 503-515.e4 (2019).

966 77. Giordano, F. J. *et al.* A cardiac myocyte vascular endothelial growth factor

967   paracrine pathway is required to maintain cardiac function. *Proc. Natl. Acad.*

968   *Sci. U. S. A.* **98**, 5780–5785 (2001).

969 78. Karra, R. *et al.* Vegfaa instructs cardiac muscle hyperplasia in adult zebrafish.

Proc. Natl. Acad. Sci. U. S. A. **115**, 8805–8810 (2018).

79.   Wang, Y. *et al.* Polymorphisms of KDR Gene Are Associated With Coronary

Heart Disease. *J. Am. Coll. Cardiol.* **50**, 760–767 (2007).

80.   Lee, K. W., Lip, G. Y. H. & Blann, A. D. Plasma angiopoietin-1, angiopoietin-2,

angiopoietin receptor Tie-2, and vascular endothelial growth factor levels in

acute coronary syndromes. *Circulation* **110**, 2355–2360 (2004).

81.   Wada, H. *et al.* VEGF-C and mortality in patients with suspected or known

coronary artery disease. *J. Am. Heart Assoc.* **7**, 1–10 (2018).

82.   Kucukardali, Y. *et al.* The relationship between severity of coronary artery

disease and plasma level of vascular endothelial growth factor. *Cardiovasc.*

Revascularization Med. **9**, 66–70 (2008).

83.   Ohno, T. *et al.* Serum vascular endothelial growth factor: A new predictive

indicator for the occurrence of coronary artery lesions in Kawasaki disease.

Eur. J. Pediatr. **159**, 424–429 (2000).

84.   Kariyazono, H. *et al.* Association of vascular endothelial growth factor (VEGF)

and VEGF receptor gene polymorphisms with coronary artery lesions of

Kawasaki disease. *Pediatr. Res.* **56**, 953–959 (2004).

85.   Wang, Y. *et al.* Vascular endothelial growth factor A polymorphisms are

associated with increased risk of coronary heart disease: A meta-analysis.

Oncotarget **8**, 30539–30551 (2017).

86.   Palmer, B. R. *et al.* Vascular endothelial growth factor-A promoter

polymorphisms, circulating VEGF-A and survival in acute coronary syndromes.

PLoS One **16**, 1–15 (2021).

87.   Rubanyi, G. M. Mechanistic, technical, and clinical perspectives in therapeutic

stimulation of coronary collateral development by angiogenic growth factors.

Mol. Ther. **21**, 725–738 (2013).

88.   Kukuła, K. *et al.* Intramyocardial plasmid-encoding human vascular endothelial

997   growth factor A165/basic fibroblast growth factor therapy using percutaneous

998   transcatheter approach in patients with refractory coronary artery disease (VIF-

999   CAD). *Am. Heart J.* **161**, 581–589 (2011).

1000   89.   Cao, J. *et al.* The single-cell transcriptional landscape of mammalian

1001   organogenesis. *Nature* **566**, 496–502 (2019).

1002   90.   Melsted, P. *et al.* Modular, efficient and constant-memory single-cell RNA-seq

1003   preprocessing. *Nat. Biotechnol.* **39**, 813–818 (2021).

1004   91.   La Manno, G. *et al.* RNA velocity of single cells. *Nature* **560**, 494–498 (2018).

1005   92.   Eden, E., Navon, R., Steinfeld, I., Lipson, D. & Yakhini, Z. GOrilla: A tool for

1006   discovery and visualization of enriched GO terms in ranked gene lists. *BMC*

1007   *Bioinformatics* **10**, (2009).

1008   93.   Raudvere, U. *et al.* G:Profiler: A web server for functional enrichment analysis

1009   and conversions of gene lists (2019 update). *Nucleic Acids Res.* **47**, W191–

1010   W198 (2019).

1011   94.   Miquerol, L. *et al.* Endothelial plasticity drives arterial remodeling within The

1012   endocardium after myocardial infarction. *Circ. Res.* **116**, 1765–1771 (2015).

1013   95.   Mahmoud, A., Porrello, E. & Kimura, W. Surgical models for cardiac

1014   regeneration in neonatal mice. *Nat. Protoc.* **9**, 305–311 (2014).

1015   96.   Miquerol L. et al. Architectural and functional asymmetry of the His–Purkinje

1016   system of the murine heart. Cardiovascular Research. Volume 63, Issue 1, 77-

1017   86 (2004).

1018

1019

Methods

Tissue digestion and single cell sorting

Neonatal and adult mouse hearts were dissected and atria was removed from each

heart. Tissue digestion was performed only with cardiac ventricles. Adult mouse

hearts were additionally perfused before harvesting hearts. ~15-20 neonatal (for

sham and MI each) and ~10-12 adult (for sham and MI each) *Cx40CreER*;

Rosa26^{TdTomato} mice were used for tissue digestion. Post-explant, ventricles of the

hearts were minced using razor blade and fine forceps and transferred to a tissue

digestion buffer. The tissue digestion buffer was composed of 500U/ml Collagenase

IV (Worthington, Catalog# LS004186), 1.2U/ml Dispase (Worthington, Catalog#

LS02100), 32U/ml DNase I (Worthington, Catalog# LS002007) in commercial grade

sterile 1X DPBS with Ca^{2+} and Mg^{2+}. Each heart was transferred to ~300µl of

digestion buffer. Hearts bathing in digestion buffer were incubated in 37° Celsius

shaker for 40-45 minutes, with periodic resuspension using micropipettes to break

down tissue lumps. Digested tissue was then equilibrated with 4 volumes of 5% FBS

prepared in 1XDPBS, gently mixed with a micropipette and filtered using 40µm of

sterile cell strainer. The filtered tissue digest was spun down for 5 minutes at 400g at

4° Celsius, and the cell pellet was gently dissolved in 5ml of 3% FBS in 1XDPBS.

The re-suspended cell pellet was spun down for 5 minutes at 400g at 4° Celsius.

Supernatant was discarded and the cell pellet was suspended in ~500-600µl of 3%

FBS in 1XDPBS. These suspended single cells were stained with Ter119-Alexafluor-

647 (Biolegend, Catalog# 116218, 1:100 dilution) for 30 minutes at 4° Celsius. Post-

staining, cells were washed with and re-suspended in fresh 3% FBS/ 1XDPBS,

followed by single cell sorting with fluorescent activated cell sorter. DAPI was added

1045   prior to starting the cell sort. Live single endothelial cells (DAPI⁻ Ter119⁻ TdTomato⁺)

1046   were collected, in that order, using BD FACS Aria II cell sorter. Single cells were

1047   counted, and submitted to Stanford Genome Sequencing Service Center for 10X

1048   single cell V3 library preparation. Sequencing was performed using HiSeq 4000.

1049

1050   **Single Cell RNA sequencing data analyses**

1051

1052   **Processing of FASTA files**

1053   Illumina reads obtained were processed on Cell Ranger (10X Genomics). To

1054   demultiplex the BCL files and convert them to FASTQ files, the *mkfastq* function was

1055   used. The reads were aligned to mm10 (mouse genome) and TdTomato sequences

1056   with the count function.

1057

Preprocessing on R

The output files of Cell Ranger were imported into RStudio v4.05 using Seurat v4.03.

This was performed with *Read10x* feature by providing path to a folder with zip files

of expression matrix, features and barcodes files. The cells were analyzed for

nFeature RNA count and percent of mitochondrial RNA using the standard pre-

processing mentioned on the Seurat[28]. The tutorial is available at

https://satijalab.org/seurat/articles/get_started.html .

The percentage of mito genes expressed in the cells of our dataset were less than 1,

hence no cells were removed. Based on Q/C violin plot, all cells with >250 nFeature

RNA count were analyzed. These cutoffs were determined from quality control

analyses of all datasets, which include, subsets of cardiac cells filtered with

expression of *Cadherin5*, *Pecam1* or TdTomato from uninjured and injured neonates

and adults (Figure **S1A**, **S2A**, **S9A**, **E**). We obtained and analyzed the following

number of cells from each subset. *Cadherin5* >1: 5432 sham and 2200 MI neonatal

cells, *Cadherin5*>1: 6256 sham and 1690 MI adult cells, *Pecam1*>1: 5520 sham and

2152 MI neonatal cells, *Pecam1*>1: 6228 sham and 1774 MI adult cells,

TdTomato>0: 5557 sham and 497 MI neonatal cells, TdTomato>0: 3090 sham and

892 MI adult cells.

DoubletFinder, a package on RStudio was used to check if there were any doublets

incorporated in the dataset. The analysis showed that, 7.5% of all cells were

doublets in both sham and MI datasets. These doublets were mostly present in

endoEC, endoMT and SMC cell clusters which were excluded from the study. Trans-

EC, a unique cluster expressing marker genes for multiple cell types was considered

specifically for the analysis (as the chance of it being a doublet is higher). No

doublets were found in MI trans-EC cluster. In the sham trans-EC cluster, there were

3 doublets.

Normalization, scaling and clustering

Data was log normalized with scale factor set to 10000, followed by finding variable

features with n set to 2000 and finally, scaled with *scaledata*. We focused on artery

endothelial cells, by sub-setting the data using different markers. *Cadherin5* or

Pecam1 (pan endothelial markers) expression greater than 1 were analyzed. A filter

for greater than 1 was used for *Cadherin5* or *Pecam1* to exclude non-ECs

expressing these genes at a minimal level. As the cells from both filters echoed the

same clustering pattern, we used *Cadherin5* cells for further analyses. The pattern

acquired by clustering sham and MI data separately or together (merged) echoed the

same patterns. Hence, the sham and MI data were integrated using the

IntegrateData feature from the Seurat package. Normalization, finding variable

features and scaling data was repeated on the integrated data, using the same

settings as above.

1097

We performed Principal Component Analysis (PCA) using the top 30 PCs, from

integrated data. We did not observe any variations in clustering patterns when 10-

100 PCs were used. Clusters for the integrated data were obtained with Louvain

algorithm with resolution set to 0.5

(https://satijalab.org/seurat/articles/pbmc3k_tutorial.html). The cells were projected

and visualized in two-dimension using Uniform Manifold Approximation and

Projection for Dimension Reduction (UMAP); parameters were n.neighbors set to 10,

dims set to 1:20, spread set to 2, and min.dist set to 0.3. Attributes *group.by* and

split.by were used to observe sham and MI cells. This was done for optimal spread

of cells and best visualization.

TdTomato sequence was aligned with and checked in our datasets. Thus, cells with

TdTomato expression greater than 0, were analyzed. Since the MI cell number was

10 fold lesser than that of the sham, data was analyzed by sub-setting the merged

data based on its orig.ident and re-clustering the respective cell populations. The

steps used with *Cadherin* >1 cells were used for all clustering steps with TdTomato

>0.

For expression of genes, data was sub-setted for gene >0, the cell numbers per

cluster were obtained using table(object@active.ident).

1116

Cluster identification

The communities detected were identified with Violin plots (*StackedVlnPlot* from

CellChat v1.1.0 package (https://github.com/sqjin/CellChat), *DotPlots* (a feature of

Seurat package to visualize gene expression), and feature plots (Seurat argument).

Clusters were identified with established markers for a variety of cardiac cell specific

genes. Cluster identifications were confirmed using *AddModuleScore* feature from

Seurat package. A list of cell-type specific, established markers were used for this

analysis on *Cadherin5*>1, *Pecam1*>1 and TdTomato>0 datasets in both neonate and

adult datasets. Average expression of the listed genes for each cell type was

visualized using a feature plot. Neonatal trans-EC cluster did not show unique

expression of any known cardiac cell types.

1128

1129 Heatmaps were used to check the variation in expression of de-differentiation genes

1130 across sham and MI datasets. *DoHeatmap* feature from Seurat was used to plot the

1131 genes of interest. Violin plots generated from Seurat package were analyzed by

1132 using the *split* attribute.

1133

Cell cycle analysis

Updated G2M and S phase specific gene lists (from 2019) on Seurat were used. As

the genes were in human gene format, *useMart* function from biomaRt v3.12

(RStudio package) was employed to obtain hsapiens_gene_ensembl and

mmusculus_gene_ensembl from ensembl database. Homology mapping and gene

symbol conversion of genes from 2 gene-sets was performed using *getLDS* function.

Genes were converted to mouse gene format using *convertHumanGeneList*. Cell

cycle analysis was performed using standard cell cycle regression

(CellCycleScoring) on Seurat.

1143

Trajectory analysis

Monocle3[89] was used for trajectory analysis to understand possible pseudo time

relationship between the cell types. Seurat object split into sham and MI from the

integrated data, subsetted for ECs alone, was used for the analysis. *as.cell_data_set*

feature from SeuratWrappers was used to convert Seurat object to Monocle3 object.

Further, the Monocle3 object was subjected to pseudo time analysis using

learn_graph and *order_cells* functions (https://cole-trapnell-

lab.github.io/monocle3/docs/starting/). Results were visualized on UMAPs with the

aid of *plot_cells* function.

1153

RNA Velocity

Spliced and unspliced counts were generated from raw Fastq files using kallisto

bustools wrapper *kb* (kb-python v0:27:0, https://www.kallistobus.tools/)[90]. Mouse

genomic (DNA) FASTA and GTF annotations were downloaded from Ensembl and a

transcriptome index was generated by kallisto using *kb ref* command. *Kb count*

command was used to generate the count matrix in the form of h5ad file. The h5ad

file was converted into a seurat object using SeuratDisk (v0.0.0.9019). The

preprocessing, normalization, scaling and clustering were done as described earlier.

The RNA velocity maps were generated using Velocyto package (velocyto.R v6)[91].

1163

Analysis of differentiation state

R package CytoTRACE v0.3.3[53] (https://cytotrace.stanford.edu/) was used for cell

differentiation analysis. Cell counts and UMAP embeddings were obtained as data

frames from sham and MI clusters separately, for the analysis. A CytoTRACE score

was assigned to each cell with the CytoTRACE function, which was visualized on the

UMAP embeddings with *plotCytoTRACE* feature. We compared the mean

CytoTRACE scores between all EC clusters. To do so, CytoTRACE score for each

cell and its cluster number was obtained. The mean CytoTRACE score was then

calculated for each cluster. These mean values along with standard deviations were

plotted as boxplots using *ggboxplot* feature from ggpubr v0.4.0 package.

1174

Gene Ontology (GO) analysis

1175

List from *FindAllMarkers* was used to recognize processes that appear on publicly

available online tool GOrilla (Gene Ontology enRIchment anaLysis and visuaLizAtion

tool)[92] (http://cbl-gorilla.cs.technion.ac.il/). Gene Ontology analyses was performed

on MI gene list, with sham genes serving as background. Default settings with a filter

of p-value less than 10^{-3} were used. Processes with significant p-values (<0.0005),

and relevant to our study, were plotted on a bar graph.

1182

g:Profiler

1183

1184 Pathway analysis was performed using an online tool g:Profiler[93]. The gene list was

1185 obtained for neonatal cluster aEC1 (from merged *Cadherin5* >1 dataset), using

1186 *FindMarkers* feature, with test.use set to "MAST". The organism was set to *Mus*

1187 *musculus* and the pathways were obtained from databases like KEGG, reactome,

1188 WikiPathways etc.

1189

Statistical analysis on sequenced data

1190

Wilcoxon test was performed using *RunPresto* feature from SeuratWrappers

package. The statistical test was performed on cyc-aEC and aEC1 cells from

neonatal *Cadherin5*[+] MI group. Additionally, the test was performed on *Cadherin5*[+]

cyc-aECs from neonatal and adult MI datasets. For this, the data was merged using

the *merge* feature from Seurat, and retained as such, without any further

modifications. Test was done to check statistical significance for expression of *Cx40*,

Cxcr4, *Aplnr*, *Car4*, *Apln*, *Foxm1, Vegfa*, *Vegfc*, *Kdr* across different treatments

(sham and MI in Figure **4**) or clusters (cyc-aEC and aEC1 in neonates, in Figure

S7A).

1200

Experimental model

Following mouse lines were used in this study. *Rosa26^{TdTomato}* Cre reporter line (The

Jackson Laboratory, B6.Cg-Gt(ROSA)26Sortm9(CAG-TdTomato)Hze/J, Stock#

007909), *Cx40CreER*[94], *VegfR2 flox* (The Jackson Laboratory, *Kdr^{tm2Sato}*/J,

Stock#018977), *Cx40$^{eGFP/+}$*[96].

All mice were housed and bred either in Stanford (in accordance with institutional

animal care and use committee (IACUC)) or NCBS animal facility (in accordance

with Institutional Animal Ethics Committee (IAEC)). All Tamoxifen injections (6mg

dissolved in corn oil) were administered intraperitoneally to nursing mothers for Cre

activation in new born pups. Efficiency of Tamoxifen and Cre activation has been

shown earlier to be ~84% using Cx40CreER mice and ~86% using ApjCreER mice[24].

Data shown in this study is compiled from age-matched males and females from

several litters.

1214

Experimental study design

Sample size for the current experiment was determined from our previous study

conducted with the same animal model with similar treatments[24]. In any given litter,

animals were categorized as control or knockout based on their genotype, and

irrespective of the sex. Severity of MI was determined by first checking the location

of LCA ligation under fluorescence stereo-microscope (to check knot around

TdTomato+ arteries). Hearts with severe MI (where either the main LCA or the two

primary branches of LCA were captured), were included in this study. Hearts with

mild MI (where secondary, or tertiary branches were ligated), were excluded from the

study. Biological replicates from multiple litters were included in the study, and are

mentioned below for each in vivo experiment.

1226

1227 **Genotyping information**

Primer description (Forward (F); Reverse (R))	Primer sequence (5' to 3')
Cre transgene F	GCATTACCGGTCGATGCAACGAGTGATGAG
Cre transgene R	GAGTGAACGAACCTGGAAATCAGTGCG
Cre internal positive control F	CTAGGCCACAGAATTGAAAGATCT
Cre internal positive control R	GTAGGTGGAAATTCTATCATCC
TdTomato wildtype F	AAGGGAGCTGCAGTGGAGTA
TdTomato wildtype R	CCGAAAATCTGTGGGAAGTC
TdTomato mutant F	CTGTTCCTGTACGGCATGG
TdTomato mutant R	GGCATTAAAGCAGCGTATCC
VegfR2 flox F	CCACAGAACAACTCAGGGCTA
VegfR2 flox R	GGGAGCAAAGTCTCTGGAAA
GFP F	GCACGACTTCTTCAAGTCCGCCATGCC
GFP R	GCGGATCTTGAAGTTCACCTTGATGCC

1228

Adult LCA ligations

Adult mice were anesthetized in a 4% isoflurane chamber and then intubated with a

20G angiocatheter. General anesthesia was maintained with 1.5-4% isoflurane

during mechanical ventilation at a respiratory rate of 180 breaths per minute. The

mice were placed in right lateral decubitus position and the left chest was shaved

and sterilely prepped. 1 mg/kg Bupivicaine and 0.5 mg/kg Buprenorphine were

injected subcutaneously for perioperative analgesia. A 1 cm left thoracotomy incision

was made, and the pericardium was opened. The left coronary artery was identified

and permanently ligated using a 6-0 polypropylene suture. The chest was then

closed in layers using 5-0 polypropylene sutures. Isoflurane was then weaned off

and the mice were extubated after regaining physiological respirations. 5mg/kg

Carprofen was injected subcutaneously daily for the first three days after surgery for

postoperative analgesia. For adult MI experiments, 2-3 months old mice were used

(both males and females). To activate Cre, a single dose of 6mg Tamoxifen

intraperitoneal injections were administered to each mouse, 2 days prior to MI.

Echocardiograms were obtained from P32/P33 adult mice (sham or MI surgeries at

P2) ~30days post-MI using VisualSonics Vevo 3100 imaging system (FUJIFILM

VisualSonics Inc.) with an MX400 20–46 MHz transducer. Data was analyzed using

Vevo LAB version 3.1.0 (Build 13029)

1248

Neonatal LCA ligations

Technical details of the procedure are referred and modified from Mahmoud et al[95].

To induce hypothermic arrest, neonatal mice (P2 or P7) were incubated on ice.

Neonates were wrapped with a 3-layer, pre-cooled, gauze piece for ~3 minutes.

Anesthesia was confirmed using toe pinch. Surgery was performed in sterile

conditions from start to end, using sterile surgical tools at all times. Animals were

placed in a supine position, on a sterile surgical drape, wrapped around cold ice

pack. Animals were then prepped with betadine and 70% ethanol. Left anterolateral

thoracotomy was performed by making an incision on fourth intercostal space to

access heart, under a dissecting microscope. The left coronary artery (LCA) was

identified and ligated by passing a needle attached to 8-0 non-absorbable prolene

suture (Ethicon Ethilon, catalog# NW3322). Post-ligation, the 8-0 prolene suture was

used to close the ribs, intercostal muscles and skin separately. The neonate was

then transferred to warm plate (37° Celsius) to gain consciousness and recover

completely. When conscious, neonates were returned to their mother's cage and

closely observed over the next few days.

1265

Whole mount immunostaining of neonatal hearts

The procedure has been described earlier[24]. Neonatal whole hearts were dissected

and immediately fixed in 4% paraformaldehyde at 4° Celsius for 1hr on a rocker.

Hearts were then subjected to three, 15 minutes washes, with 1X PBS at 4° Celsius.

Primary antibody staining was done by incubating hearts in respective dilutions of

primary antibodies, prepared in at least 5 volumes of 1X PBS containing 0.5% Triton-

X 100 (0.5% PBT). Incubations were done overnight at 4° Celsius, on a shaker.

Hearts were washed in 20 volumes of 0.5% PBT for 12 hours, with a change in wash

buffer (0.5% PBT), every 2 hours at 4° Celsius. The secondary antibody staining was

performed by incubating hearts in 1:250 dilution prepared in 0.5% PBT. Incubation

was done on a shaker. Next, hearts were washed in 20 volumes of 0.5% PBT, for 4

days with a change in wash buffer, every two hours at 4° Celsius. All washes were

performed with continuous but gentle shaking. Hearts were cleared in at least 2

volumes of Anti-fade mounting media

(https://www.jacksonimmuno.com/technical/products/protocols/anti-fade) for 2

hours, in dark, at room temperature. Whole hearts were then stored at -20° Celsius

till imaged. Post staining washes were done in 15 and 50 ml tubes while incubation

in antibodies or mounting media were performed in 1.5ml microtubes.

1284

1285 Neonates were injected with 30µl of 10mM 5-ethynyl-2'-deoxyuridine (EdU)

1286 (prepared in 1X commercial grade PBS, catalog #10010023) for 1hour, prior to

1287 harvesting their hearts. EdU label detection was combined with post-secondary

1288 immunostaining steps as described above, and, detected using Click-iT staining kit

1289 (Invitrogen, Catalog# C10340).

1290

Confocal imaging of neonatal hearts

Whole neonatal (immunostained) hearts were mounted on a double concave

microscope slide (Sail brand, catalog# 7104) using a microscope coverslip (0.13-

0.17mm, Thermo Scientific, Catalog# 48367-059) so that the ventral watershed

regions spanning left and right coronary arteries can be imaged. Imaging was

performed using Olympus FV3000 upright confocal microscope. Whole hearts were

imaged using 4X, 10X, 20X or 60X (oil) objectives. Multiple z stacks were acquired

using FV31S-SW software. Multiple images of the whole heart were processed on

Fiji-ImageJ software, and stitched with Adobe Photoshop.

1300

1301 **Antibodies**

1302 Primary antibodies: Anti-CX40 (1:500; Alpha Diagnostics Int. Inc.; Catalog# CX40-A,

1303 RRID: AB_1616128), Anti-VEGFR2 (1:125, R&D Systems, Catalog# AF644, RRID:

1304 AB_355500). Secondary antibodies: Alexa fluor-conjugated antibodies (488, 555,

1305 633) from Invitrogen were used at 1:250 dilutions.

1306

Quantification from images

Collateral number

Number of *Cx40CreER* lineage-traced (TdTomato positive) collateral arteries were

quantified from images acquired using 10X objective from whole hearts. Number of

stacks (12-14) and step size (2µm) of the images used, were consistent across

genotypes. Data was compiled from 6-9 litters from each genotype. 8 control, 5

heterozygous and 14 knockout hearts were used for quantification. An unpaired t-

test was performed to check statistical significance.

1315

Cx40 intensity in proliferating artery ECs

Proliferating TdTomato$^+$ Cx40$^+$ cardiac artery ECs (4 days post-MI) were identified

with EdU$^+$ nuclear labelling. Cytoplasmic intensity of Cx40 in these (TdTomato$^+$

Cx40$^+$ EdU$^+$) ECs were quantified using images acquired with 60X (oil) objective.

Mean fluorescent intensity (MFI) of cytoplasmic Cx40 signal was measured using a

selection tool available with Fiji-ImageJ software. Quantification was performed with

68 single TdTomato$^+$ Cx40$^+$ EdU$^+$ artery ECs, found in the watershed of 4 hearts,

collected 4 days post-MI. Comparison between proliferative (EdU$^+$) and non-

proliferative (EdU$^-$) artery ECs were made within the same field of view. An unpaired

t-test was done to check statistical significance. Similar analysis was done on 67

artery ECs within artery tips or integrated into mature arteries, as identified by Cx40

immunostaining. Data was compiled from 5 MI hearts from different 4 litters.

1328

Single artery EC proliferation in control and *VegfR2* knockout hearts

EdU$^+$ TdTomato$^+$ single artery ECs in the watershed regions, were quantified using

images acquired with 20X objective. On an average 20-30 TdTomato$^+$ single artery

ECs were observed per watershed (each field of view), which was consistent across

genotypes. Quantification was performed on 6 control (36 field of views) and 5

VegfR2 knockout (24 field of views) hearts. A total of 784 control TdTomato$^+$ artery

ECs and 512 *VegfR2* knockout TdTomato$^+$ artery ECs were counted for this

quantification. Data was compiled from 3-5 litters from each genotype. 6 control and

5 knockout hearts were used for quantification. An unpaired t-test was performed to

check statistical significance.

1339

Single artery EC proliferation in P11 MI hearts (MI at P7)

EdU⁺ TdTomato⁺ proliferating artery ECs were counted in artery tips and saECs from

3 *Cx40CreER*; *Rosa26^{TdTomato}* hearts. ~80-120μm of distal ends of arteries were

identified as artery tips. 25 fields of view from P11 watershed regions, from images

taken with 60X (oil) objective, were used to count a total of 113 TdTomato⁺ residual

single cells.

1346

TdTomato fluorescence in control and *VegfR2* knockout watershed

Mean Fluorescence Intensity was measured using Fiji for *Cx40CreER* lineage-traced

TdTomato⁺ single artery ECs found in watershed area 4 days post-MI. Images of

watershed regions (each field of view) lacking any large or collateral arteries were

acquired with 20X objective. Watershed images only with single artery cells were

considered for this quantification. 3 hearts   from each genotype were used for this

quantification. An unpaired t-test was performed to check statistical significance.

1354

1355   **LCA width:**

1356   LCA width was measured using a line selection tool on Fiji. The analysis was done

1357   for 3 control and 2 *VegfR2* KO hearts. Width was measured for the main branch of

1358   LCA in all hearts and genotypes. An unpaired t-test was performed to check

1359   statistical significance.

1360

1361

Figures and legends

Figure 1: Single cell analyses of neonatal artery ECs

(**A**) Experimental setup to obtain neonatal cardiac cells for scRNA sequencing using

10X genomics (**B**) UMAP showing the distribution of *Cadherin5*⁺ neonatal sham and

MI cells. (**C**, **D**) Feature Plot showing the distribution of *Cadherin5*⁺ cells from (**C**)

sham and (**D**) MI group. (**E**, **F**) Distribution of TdTomato expressing cells within

Cadherin5⁺ (**E**) sham and (**F**) MI group. (**G**, **H**) UMAP showing distribution of (**G**)

sham and (**H**) MI cell clusters from the merged *Cadherin5*⁺ cells. (**I**) Dot Plot showing

average expression and percentage of cells, expressing relevant markers used for

cluster identification. (**J**) Graph showing percentage of sham and MI *Cadherin5*⁺ cells

in different cell cycle stages based on cell cycle sorting analysis on Seurat. (**K**)

Graph showing percentage of *Cadherin5*⁺ cells undergoing G2 to M transition, in

sham and MI groups, in all clusters or in only artery EC clusters (aEC1-4 and cyc-

aEC). (**L**) Graph showing percentage of *Cadherin5*⁺ and *Cx40creER*-linegae traced

(TdTomato⁺) cells undergoing G2 to M transition, in sham and MI groups, in all

clusters or in only artery EC clusters (aEC1-4 and cyc-aEC). (**M**) Pie chart showing

the percent distribution of G2/M *Cadherin5*⁺ cells, upon MI. (**N-P**) Confocal image of

an (**N**) artery tip from a post-MI heart, immunostained for Cx40, lineage traced with

TdTomato and labelled with EdU show presence of (**O**) Cx40⁺ TdTomato⁻ EdU⁺

(cyan arrowhead) artery EC and (**P**) Cx40⁺ TdTomato⁺ EdU⁺ (yellow arrowhead)

within the same artery branch. MI, myocardial infarction; *Cdh5*, *Cadherin5;* Td,

TdTomato; art, artery. Scale bar: **N-P**: 50µm

1387

1388

1389

1390

 Figure 2: In silico assessment of de-differentiation in neonatal cycling artery

 ECs

G Comparison of features indicating de-differentiation across neonatal EC clusters in MI
(cyc-aEC, trans-EC, aEC1)

Cellular/ Molecular features	Evidence	cyc-aEC	trans-EC	aEC1
Origin (from a differentiated cell-type)	Cells originating from pre-existing artery ECs (TdTomato+ cells)	8.4%	8.5%	20.8%
Loss of mature cell markers	Gja5 and Cxcr4 (av. expression)	Cx40: 0.17 Cxcr4: 0.16	Cx40: 0.002 Cxcr4: 0.27	Cx40: 1 Cxcr4:1
Gain of (early) capillary/venous EC markers	Aplnr and Car4 (av. expression)	Aplnr: 1 Car4: 1	Aplnr: 1.69 Car4: 2.24	Aplnr: 0.98 Car4: 0.88
Gain of pro-angiogenic markers	Apln, Foxm1 (av. expression)	Apln: 1 Foxm1: 1	Apln: 0.91 Foxm1: 0.1	Apln: 0.18 Foxm1: 0.04
Cell cycle re-entry	Cell cycle sorting (% cells)	G2M: 52.16% S: 47.8%	G2M: 7.02% S: 30.1%	G2M: 0.97% S: 22.2%
Cellular fate (differentiation into a specialized cell-type)	CytoTrace	least differentiated	intermediate	most differentiated
	Monocle3	origin of predicted path	connecting cyc-aEC and aEC1	end of predicted path

(**A**) Heatmap showing gene expression of *Apln*, *Aplnr*, *Car4, Cx40* and *Cxcr4* across

all *Cadherin5*+ clusters in neonatal sham group. (**B**) Violin plots comparing

expression of *Cx40* and *Aplnr* between different EC clusters in sham and MI group.

(**C**) Heatmap showing gene expression of *Apln*, *Aplnr*, *Car4, Cx40* and *Cxcr4* across

all *Cadherin5*+ clusters in neonatal MI group. (**D**) Box plots illustrating a relative

differentiation state of *Cadherin5*+ MI cells with scores obtained from CytoTRACE.

(**E**, **F**) Trajectories obtained by performing Pseudo time analysis on neonatal

Cadherin5+ (**E**) sham cells or (**F**) MI cells, using Monocle3. (**G**) Comparison of de-

differentiation features across the clusters involved in pseudo time analysis of

Cadherin5+ neonatal MI cells (cyc-aEC, trans-EC and aEC1).

1405

1406

1407

1408

1409

1410

1411

1412

1413

1414

1415

1416

1417

1418

Figure 3: In vivo assessment of de-differentiation in neonatal cycling artery

1420 **ECs**

1421

1422

(**A**) Experimental set up to check Cx40 levels in proliferating TdTomato artery ECs

post-MI (**B**) Confocal images of EdU+ TdTomato+ single artery ECs (yellow

arrowhead) and EdU⁻TdTomato⁺ artery tip EC or saEC (white arrowheads) in P6

Cx40CreER; *Rosa26^{TdTomato}* hearts, 4 days post-MI. Cyan arrowhead points to EdU⁺

Tdtomato⁺ saECs with relatively high Cx40 expression. (**C-D**) Quantification of

cytoplasmic Cx40 in P6, *Cx40CreER-* lineage traced (**C**) single aECs (p=0.0001) and

(**D**) artery ECs (p=0.09), 4 days post-MI. (**E**) Experimental set up to check presence

of *Apj* in TdTomato single artery ECs post-MI. (**F-G**) Confocal images of GFP+ artery

tips and single artery ECs showing presence (marked by *) or absence (marked by

arrowhead) of TdTomato+ Apj-lineage. (**H**) A proposed model where upon MI, pre-

existing stably differentiated artery ECs (green) exit artery tips as single artery ECs,

de-differentiate (blue), proliferate (red) and coalesce into coronary collateral arteries.

saECs are lineage traced from pre-existing artery ECs (TdTomato⁺), but

downregulate expression of mature artery marker Cx40 and upregulate cap/vein EC

marker Apj upon de-differentiation. Capillary ECs are shown in grey. P, postnatal

day; saECs, single artery endothelial cells. Scale bar: **B**: 50µm, **F-G**: 50µm

1439
1440

Figure 4: Cell cycle analyses of adult artery ECs post-MI

(**A**) Experimental setup to obtain adult cardiac cells for scRNA sequencing using 10X

genomics. (**B**) Feature Plot showing the distribution of adult *Cadherin5*⁺ cells from

both sham and MI groups. (**C**) UMAP showing the distribution of *Cadherin5*$^+$ adult

sham and MI cells. (**D**) UMAP showing distribution of both sham and MI cell clusters

from the merged adult *Cadherin5*$^+$ cells. (**E**) DotPlot showing average expression

and percentage of cells, expressing relevant markers used for cluster identification in

D. (**F**, **G**) Distribution of TdTomato expressing cells within *Cadherin5*$^+$ (**F**) sham and

(**G**) MI group. (**H**) Graph showing percentage of adult, sham and MI, *Cadherin5*$^+$

cells in different cell cycle stages based on cell sorting analysis on Seurat. (**I**) Graph

showing percentage of *Cadherin5*$^+$ cells undergoing G2 to M transition, in sham and

MI groups, in all clusters or in only artery EC clusters (aEC1-7 and cyc-aEC). (**J**)

Graph showing percentage of *Cadherin5*$^+$ and *Cx40creER*-linegae traced

(TdTomato$^+$) cells undergoing G2 to M transition, in sham and MI groups, in all

clusters or in only artery EC clusters (aEC1-7 and cyc-aEC). (**K**) Single cell data

analyses of all TdTomato$^+$ cells from neonatal and adult MI groups, showing

distribution of different cell types that originate from *Cx40CreER* lineage labelling or

subsequent tracing. (**L**, **M**) Representative image of P11 *Cx40CreER*; *Rosa26*^{TdTomato}

(**L**) whole heart and respective (**M**) watershed region, 4 days post-MI. MI was

performed at P7. Lineage traced artery ECs are shown in black. * marks ligated

branches of LCA, arrowheads point to "residual" saECs. (**N**, **O**) Graph showing

number and percentage of EdU$^+$ Tdtomato$^+$ ECs in (**N**) artery tips and (**O**) residual

saECs, respectively, at P11. MIs were performed at P7. P, postnatal day; LCA, left

coronary artery; RCA, right coronary artery; art., artery; Adt, adult; Neo, neonate;

saECs, single artery endothelial cells. Scale bars: **L**: 500µm; **M**: 200µm

1468

Figure 5: Analyses of de-differentiation features in adult MI cells

(**A-B**) Box plots illustrating a relative differentiation state of adult *Cadherin5*⁺ (**A**)

sham and (**B**) MI cells, with scores obtained from CytoTRACE. (**C**) Heatmap of adult

sham and MI cells showing the expression of *Apln*, *Aplnr*, *Car4*, *Cx40* and *Cxcr4*, in

their respective cyc-aEC cluster. (**D**) Comparison of de-differentiation features in

Cadherin5+ neonatal and adult cyc-aEC cluster. (**E-F**) Pseudo time analysis of adult

Cadherin5+ cells from (**E**) sham and (**F**) MI group. (**G**) Schematic showing post-MI

neonatal artery ECs accomplish artery regeneration by undergoing de-differentiation

and proliferation; a phenomenon rarely observed in adult. Dotted line, minimal

contribution. FC, Fold Change; av., average; diff, differentiation; Td, TdTomato; #,

number.

1482

Figure 6: Vegf pathway in neonatal Artery Reassembly and cardiac function

1484

1485

(**A**) Results from Gene Ontology analysis on TdTomato⁺ cells from neonatal MI

group (with sham as background), using the tool GOrilla. (**B**) Pathway analysis on

genes expressed in neonatal aEC1 cluster, using g:Profiler. (**C**) DotPlot showing

average expression and percentage of cells expressing *Vegfa*, *Vegfc* and *Kdr*

(*VegfR2*) in all *Cadherin5*$^+$ cells. (**D**) Experimental set up to test the role of arterial

VegfR2 in neonatal Artery Reassembly. (**E**, **F**) Representative image of (**E**) control

and (**F**) *VegfR2* artery endothelial cell specific knockout whole hearts, 4 days post-

MI. (**G**-**I**) Representative confocal images of P6 watershed regions from (**G**) un-

injured, (**H**) MI control and (I) MI *VegfR2* knockout hearts. MIs were performed at P2.

Lineage traced artery ECs are shown in black. (**J**) Quantification of *Cx40CreER*-

lineage traced collateral artery numbers observed 4 days post-MI. * represents p-

value 0.0368, **** represents p-value <0.0001 (**K**) Quantification of percentage of

EdU$^+$ Tdtomato$^+$ saECs in watershed regions from *Cx40CreER*-lineage traced

hearts, 4 days post-MI. ** represents p-value 0.0083 (**L**, **M**) Representative confocal

images of EdU$^+$ Tdtomato$^+$ saECs (white arrowheads) in watershed regions from

Cx40CreER-lineage traced hearts, (**L**) with or (**M**) without arterial *VegfR2*, 4 days

post-MI. LCA, left coronary artery; RCA, right coronary artery; WSA, watershed Area.

(**N**-**Q**) Graphs showing various parameters of cardiac function as measured in adult

wild-type (n=27) and *VegfR2* artery-EC specific knockout mice (n=10). p=0.5067

(Ejection Fraction), p=0.3401 (Fractional Shortening), p=0.0108 (Cardiac Output),

p=0.0186 (Stroke Volume). Scale bars: **E**, **F**: 500µm; **G-I:** 100 µm; **L**, **M**: 50 µm

1507
1508

1509

1510

1511

1512

1513

1514

 Supplemental Figures and legends

 Supplemental Figure 1: Identification of neonatal *Cadherin5+* cell clusters

1517

1518

(**A**) Violin plots showing quality control (analyses of nFeature counts, nCount_RNA

and mitochondrial genes) performed on neonatal *Cadherin5+* dataset. (**B**)

Visualization of different identified cell clusters, in neonatal *Cadherin5+* dataset, using

UMAP. (**C**) Violin Plot showing expression of endo-MT genes in neonatal *Cadherin5+*

cells. (**D**-**G**) Feature plots showing the expression of genes specific to G2/M phase.

(**H**-**K**) Feature plots showing the expression of S phase markers in neonatal

Cadherin5$^+$ cells. (**L**) Dot plot showing parentage of cells and average expression of

genes, specific to different cell-types.

1527

1528

1529

1530

1531

1532

1533

1534

1535

1536 **Supplemental Figure 2: Artery specific gene expression in neonatal *Cadherin5*+**

1537 **cycling aEC cluster**

1538

1539

1540 (**A-D**) Violin plots showing the expression levels of artery endothelial cell specific

1541 markers in *Cadherin5*+ neonatal sham cells.

1542

1543 **Supplemental Figure 3: Gene expression analyses of neonatal *Cadherin5*+**

1544 **trans-EC cluster**

1545

1546

1547 (**A**) Visualization of cells upon re-clustering trans-EC cluster from neonatal dataset,

1548 using UMAP. (**B**) Dot plot showing average gene expression and percentage of cells

1549 expressing various genes expressed in ECs. (**C**) Feature plot showing distribution of

1550 TdTomato expressing cells in **A**. MF, myofibrillar; cap, capillary; exp, expression.

1551

Supplemental Figure 4: Identification of neonatal *Pecam1⁺* cell clusters

(**A**) Violin plots showing quality control (analyses of nFeature counts, nCount_RNA

and mitochondrial genes) performed on neonatal *Pecam1⁺* dataset. (**B**) Visualization

of clusters in neonatal *Pecam1⁺* cells, using UMAP. (**C**) DotPlot showing percentage

of cells and average gene expression of cell-type specific genes.

 Supplemental Figure 5: Analyses of post-MI artery EC proliferation in neonatal

 hearts

1563

Representative confocal images of artery segments from neonatal P6 hearts, 4 days

post MI, immunostained for Cx40, lineage traced for *Cx40creER* (with TdTomato)

and labelled for EdU. Images are taken from ischemic regions (watershed area

below the ligation). Arrows and arrowheads show (**A**-**C**) EdU$^+$ Cx40$^+$ Tdtomato$^-$

artery ECs and (**D**, **E**) EdU$^+$ Cx40$^+$ Tdtomato$^+$ artery ECs. (**F**) Modes of post-MI

artery growth in neonatal hearts. neg., negative; aEC, artery endothelial cells; cap-

EC, capillary endothelial cells. Td, TdTomato. Scale bar: 50µm

1571

 Supplemental Figure 6: Analyses of de-differentiation features in *Cadherin5+*

 neonatal cells

(**A**) Table showing percentage of sham and MI cells expressing *Apln*, *Aplnr*, *Car4*,

Cx40 and *Cxcr4*. (**B**) DotPlot from neonatal *Cadherin5+* cells showing average

expression and percentage of cells expressing pro-angiogenic genes (*Cd34*, *Foxm1*,

Nrp2, *Col15a1*, *Apln*). (**C**-**E**) Feature Plots showing expression of pro-angiogenic

Apln, *Col15a1*, and *Nrp2* in *Cadherin5*⁺ neonatal cells, from sham and MI together.

(**F**-**H**) Feature Plots showing expression of angiogenic genes, also expressed by

capillary or venous ECs (*Gpihbp1*, *Fabp3*, *Pcdh17*), in *Cadherin5*⁺ neonatal cells,

from sham and MI together. (**I**-**K**) Feature Plots showing expression of mesenchymal

genes (*Tnnt2*, *Myl3*, *Myh6*), in *Cadherin5*⁺ neonatal cells, from sham and MI

together.

1586

1587

1588

1589

1590

1591

 Supplemental Figure 7: Comparison of de-differentiation features between

 neonatal *Cadherin5*⁺ cells in cyc-aEC, trans-EC and aEC1 clusters

1594

1595

1596 (**A**, **B**) RNA velocity analysis on *Cadherin5*⁺ (**A**) sham and (**B**) MI cells.

1597

1598

1599

1600

1601

1602

1603

1604

1605

1606

1607

 Supplemental Figure 8: Identification of adult cardiac cell clusters

1609

(**A**) Violin plots showing quality control (analyses of nFeature counts, nCount_RNA

and mitochondrial genes) performed on adult *Cadherin5*⁺ dataset. (**B**) DotPlot

showing percentage of cells and their average gene expression used for identifying

clusters, obtained from re-clustering *Cadherin5*⁺ cap-ECs. (**C**) Visualization of re-

clustered adult *Cadherin5*+ cap-ECs on a UMAP. Clusters were identified using the

DotPlot shown in **B**. (**D**) FeaturePlot showing the distribution of TdTomato

expressing cells on re-clustered cap-ECs in **C**. (**E**) Violin Plots showing expression of

artery specific genes in adult *Cadherin5*+ cells in both sham and MI group,

integrated. (**F**) Violin plots showing quality control (analyses of nFeature counts,

nCount_RNA and mitochondrial genes) performed on adult *Pecam1*+ dataset. (**G**)

Visualization of clusters in adult *Pecam1*+ cells, using UMAP. (**H**) DotPlot showing

average gene expression and percentage of cells expressing cell type specific genes

in *Pecam1*+ dataset. (**I**) Visualization of clusters in adult TdTomato+ cells, using

UMAP. (**J**) DotPlot showing average gene expression and percentage of cells

expressing cell type specific genes in TdTomato+ dataset.

1625

1626

1627

1628

1629

1630

1631

1632

1633

1634

1635

1636

1637

 Supplemental Figure 9: Analyses of arterial Vegf pathway in uninjured and

 injured neonatal hearts

VegfR2 expression in artery ECs and capillaries

No significant difference in the artery network between control and VegfR2 artery-specifc KO hearts

Wastershed regions from *VegfR2 L/L; Cx40CreER; Rosa26TdTomato* hearts

1640

(**A**) Representative confocal images from uninjured P6 *Cx40CreER*; *Rosa26^TdTomato*

neonatal hearts showing expression of VegfR2 in small arteries, artery tips and

capillaries, but not in large arteries. (**B-D**) Representative whole heart images from

uninjured *Cx40CreER*; *Rosa26^TdTomato* mice, (**B**) with or (**C**) without arterial EC

VegfR2. Tamoxifen was injected at P0, hearts were explanted at P6. (**D**)

Quantification of width of main left coronary artery (LCA) in uninjured control and

hearts deleted for *VegfR2* from arterial ECs. Tamoxifen was injected at P0, hearts

were explanted at P6. The difference was statistically insignificant (p-value 0.1881).

(**E-G**) Representative confocal images of *VegfR2* depleted watershed regions from

P6 neonatal *Cx40CreER*; *Rosa26^TdTomato* hearts, 4 days post-MI. Tamoxifen was

administered at P0/P1, MI was performed at P2/P3. (**H**) Quantification of

Cx40CreER-lineage traced (TdTomato⁺) saECs in *VegfR2* depleted watershed

regions from P6 neonatal hearts (shown in **E-G**), 4 days post-MI. * represents p-

value 0.0449. cap, capillary; KO, knockout; wsa, watershed area. Scale bars: **A**:

50µm; **B**, **C**: 500 µm; **E-G**: 100 µm

1656

1657

1658

1659

1660

1661

1662

1663

1664

1665

1666 **Supplemental Figure 10: Proposed working model**

1667

(**A**) Dot plot showing average gene expression and percentage of cells expressing

Vegfa, *Vegfc* and *VegfR2 (Kdr)* in adult *Cadherin5*[+] dataset. Blue box indicates

ligand expression by aECs, red box indicate downregulation of *VegfR2* in cyc-aEC

cluster. (**B**) Proposed working model of cellular events associated with Artery

Reassembly. In regenerative neonatal hearts, MI induces sequential migration (via

Cxcr4), de-differentiation, and proliferation (via VegfR2) of pre-existing artery ECs,

which, leads to formation of collateral arteries and efficient revascularization of

ischemic heart regions. Non-regenerative hearts show minimal migration, de-

differentiation or proliferation. While Cxcl12/Cxcr4 pathway drives migration, in this

study we show that artery EC proliferation is regulated by Vegf pathway. Green,

artery ECs; Blue, de-diff (de-differentiation); Red, prol. (proliferation); mig, migration;

? unknown molecules associated with the cellular events linked to Artery

Reassembly. Findings of this study (de-differentiation and VegfR2-regulated

proliferation of artery ECs) are underlined.

1682

1683

1684

Table S1: Summary of all p-values obtained from Wilcoxon sum rank test

Statistical significance of gene expression variation as observed using Wilcoxon test						
Statistical analysis of de-differentiation genes across neonatal *Cadherin5⁺* sham and MI cyc-aEC clusters						
	p_val	auc	avg_log2FC	pct.1	pct.2	p_val_adj
Gja5	6.80E-52	0.225561	-1.39143	0.026	0.569	1.30E-47
Cxcr4	1.71E-20	0.329778	-0.30521	0.082	0.431	3.27E-16
Aplnr	1.75E-17	0.727798	0.699507	0.673	0.257	3.36E-13
Car4	5.90E-06	0.598174	0.413379	0.312	0.125	1.13E-01
Apln	1.57E-14	0.701489	0.762624	0.608	0.222	3.02E-10
Foxm1	No significant difference					
Statistical analysis of de-differentiation genes across neonatal *Cadherin5⁺* MI cyc-aEC and trans-EC clusters						
	p_val	auc	avg_log2FC	pct.1	pct.2	p_val_adj
Gja5	No significant difference					
Cxcr4	No significant difference					
Aplnr	4.82E-12	0.377149	-0.5043	0.673	0.758	9.23E-08
Car4	1.02E-07	0.415964	-0.55117	0.312	0.436	1.95E-03
Apln	No significant difference					

Foxm1	7.04E-44	0.654272	0.367115	0.341	0.031	1.35E-39

Statistical analysis of de-differentiation genes across neonatal *Cadherin5*[+] MI trans-EC and aEC1 clusters

	p_val	auc	avg_log2FC	pct.1	pct.2	p_val_adj
Gja5	1.64E-24	0.423338	-0.30514	0.001	0.155	3.14E-20
Cxcr4	5.20E-12	0.398321	-0.40858	0.117	0.314	9.97E-08
Aplnr	2.26E-16	0.684601	0.517376	0.758	0.367	4.33E-12
Car4	1.43E-08	0.613955	0.616043	0.436	0.222	2.74E-04
Apln	7.20E-22	0.694867	0.94641	0.488	0.106	1.38E-17
Foxm1	No significant difference					

Statistical analysis of de-differentiation genes across neonatal *Cadherin5*[+] MI cyc-aEC and aEC1 clusters

	p_val	auc	avg_log2FC	pct.1	pct.2	p_val_adj
Gja5	No significant difference					
Cxcr4	1.00E-14	0.378972	-0.48813	0.082	0.314	1.92E-10
Aplnr	No significant difference					
Car4	No significant difference					
Apln	5.76E-30	0.753943	1.028363	0.608	0.106	1.10E-25
Foxm1	2.78E-19	0.663357	0.399056	0.341	0.014	5.33E-15

Statistical analysis of de-differentiation genes across adult *Cadherin5*⁺ sham and MI cyc-aEC clusters

	p_val	auc	avg_log2FC	pct.1	pct.2	p_val_adj
Gja5	No significant difference					
Cxcr4	0.037701	0.373907	-0.57171	0.327	0.5	1
Aplnr	0.015657	0.612245	0.824505	0.265	0.036	1
Car4	No significant difference					
Apln	0.091189	0.565233	0.623217	0.163	0.036	1
Foxm1	No significant difference					

Statistical analysis of *Vegf* pathway genes across neonatal *Cadherin5*⁺ sham and MI cyc-aEC clusters

	p_val	auc	avg_log2FC	pct.1	pct.2	p_val_adj
Kdr	3.49E-22	0.770358	0.799244	0.933	0.743	6.69E-18
Vegfa	1.53E-21	0.299646	-0.49758	0.144	0.528	2.93E-17
Vegfc	1.81E-51	0.176933	-1.4309	0.099	0.708	3.48E-47

Statistical analysis of Vegf pathway genes across adult *Cadherin5*⁺ sham and MI cyc-aEC clusters

	p_val	auc	avg_log2FC	pct.1	pct.2	p_val_adj
Kdr	1.45E-15	0.125061	-1.96031	0.857	0.97	2.67E-11

Vegfa	3.01E-03	0.586006	0.375123	0.286	0.107	1.00E+00
Vegfc	1.29E-02	0.586856	0.265401	0.388	0.19	1.00E+00

1687

1688

1689

1690