De-differentiation and Proliferation of Artery Endothelial Cells Drive Coronary Collateral Development

Gauri Arolkar[1*], Sneha K.[1*], Hanjay Wang[2], Karen M. Gonzalez[3,4], Suraj Kumar[1], Pamela E. Rios Coronado[5], Y. Joseph Woo[2], Kristy Red-Horse[3,4,6], Soumyashree Das[1#]

[1] National Center for Biological Sciences, Tata Institute of Fundamental Research, Bengaluru, KA 560065, India

[2] Department of Cardiothoracic Surgery, Stanford University School of Medicine, Stanford, CA 94305, USA

[3] Department of Biology, Stanford University, Stanford, CA 94305, USA

[4] Institute for Stem Cell Biology and Regenerative Medicine, Stanford University School of Medicine, Stanford, CA 94305, USA

[5] Department of Bioengineering, Stanford University, Stanford, CA 94305, USA

[6] Howard Hughes Medical Institute

[*] Equal contribution

[#] Corresponding author (soumyashree@ncbs.res.in)

Key Words

Coronary Collateral Artery, Endothelial Cell, Vascular Endothelial Growth Factor Receptor-2, De-differentiation, Proliferation, Cardiac Regeneration, Single Cell RNA Sequencing, Myocardial Infarction, Ischemia

Abstract

Collateral arteries act as natural bypasses, which re-route blood flow to ischemic regions and facilitate tissue regeneration. In the heart, injured neonatal artery endothelial cells orchestrate a systematic series of cellular events, which includes their outward migration, proliferation, and coalescence into fully functional collateral arteries. This process, called Artery Reassembly, aids complete cardiac regeneration in neonatal hearts, but is absent in adults. The reason for this age-dependent disparity in artery cell response is completely unknown. In this study, we investigated if regenerative potential of coronary arteries, like in cardiomyocytes, is dictated by their ability to de-differentiate. We performed single cell RNA sequencing of coronary endothelial cells and identified differences in molecular profiles of neonatal and adult endothelial cells. Neonates show significant increase in actively cycling artery cells that expressed prominent de-differentiation markers. Using both, *in silico* pathway analyses and *in vivo* experiments, we show that cell cycle re-entry of pre-existing artery cells and subsequent collateral artery formation, is dependent on arterial VegfR2. This sub-population of de-differentiated and proliferating artery cells is absent in non-regenerative P7 or adult hearts. Together, our data indicate that adult artery endothelial cells fail to drive collateral artery development due to their limited ability to de-differentiate and proliferate.

Background

Cardiovascular disease (CVD) represents 32% of all global deaths (World Health Organization) and its burden is increasing sharply in many new parts of the world. Earlier, deaths due to CVD dominated the western world, but currently, at least 58% of all CVD deaths occur in Asia (*http://ghdx.healthdata.org/gbd-results-tool*). The primary cause of CVD is coronary artery blockage, and can lead to ischemic heart disease (IHD). Despite the deadly outcomes, management of symptoms associated with IHD often include invasive procedures, which, many patients are ineligible for. Effective therapeutic approaches targeting complete cardiac regeneration, can reduce mortality and/or manage symptoms of IHD. This can be accomplished by understanding, and subsequently utilizing, the molecular mechanisms associated with different physiological aspects of IHD.

Complete cardiac regeneration can restore the functional integrity of an ischemic heart. In preclinical studies, fast revascularization has been crucial[1], and often, precedes expansion of cardiomyocyte population[1,2]; a process, that consequently re-establishes cardiac function. Under the influence of cardiac angiogenic signals, pre-existing coronary blood vessels facilitate formation of numerous micro-vessels near the infarct area and mitigate injury-induced tissue damage[3–5]. This *de novo* process of small vessel formation triggers early reperfusion and restores blood flow in ischemic regions[5]. Though microvascular network is critical for tissue survival, these small vessels lack mural cell coverage, do not provide fast, consistent or voluminous blood flow, and are susceptible to rupture[6,7]. However, re-perfusion via large arteries can be an efficient mode of blood-flow restoration under ischemic conditions.

Large collateral arteries enhance perfusion[8], vascularize hypoxic area, improve myocardial viability[9], and are associated with better survival rate[10]. Collateral arteries are the arterial segments which connect a healthy arteriole to an occluded/non-perfused arteriole, and are sufficiently large to carry adequate blood flow to ischemic regions in need. Data from clinical studies suggest that the presence of native/pre-existing coronary collateral arteries present at the onset of myocardial infarction (MI) helps limit the infarct size, and improves prognosis[9,11]. The number of pre-existing collateral arteries as well as their capacity to undergo remodeling upon occlusion, are the key determinants of tissue recovery[12]. Genetic factors are proposed to regulate differences in collateral number and their functionality across human races and species[13,14]. A complete map of molecular mechanisms/pathways, regulated by these genetic factors, can be a useful tool to design therapeutic interventions for IHD patients.

Brain, a highly complex organ with low regenerative ability, is known to possess pre-existing collateral vessels in its pial layer. In mice, these collateral connections develop at an embryonic stage, and provide protection against ischemia in adulthood[15,16]. Pre-existing collateral arteries in mouse hind limb can grow by two distinct processes— arterialization of capillary endothelial cells (ECs)[17] and lateral expansion/widening of smaller pre-existing collateral arteries i.e. arteriogenesis[18]. Both, in brain and hind limb, genetic background is a major contributor towards the variation in the number of observed collaterals[19,20]. Specifically, polymorphism in *Vascular Endothelial Growth Factor-a* (*Vegfa)* locus is shown to control Vegfa expression and pre-determine the extent of collateral artery network in mice[12,21]. Particularly in hind limb ischemia model, density of pre-existing collateral arteries is

regulated via Vegf Receptor-1 (VegfR1)[21]. However, if VegfR1 also controls post-ischemic coronary collateral development in mice, is unknown.

In mice, complete cardiac regeneration is accomplished only when hearts are injured within a neonatal regenerative window (P0-P6). Any injury beyond this time point (>P7), limits cardiac regeneration[22,23]. Previously we have shown that, in regenerative neonatal mice, coronary collateral arteries are built by the systematic execution of 3 major cellular events driven by pre-existing artery ECs— migration, proliferation and coalescence[24]. Artery ECs expressed C-X-C motif Chemokine Receptor 4 (CXCR4), a receptor for a chemokine, C-X-C motif Chemokine Ligand 12 (Cxcl12), which steers the migration of artery ECs into the watershed. This process, named Artery Reassembly, involves disassembly of artery tips into single migratory artery cells, followed by their reassembly to build new artery segments (collateral arteries). Interestingly, despite *Cxcl12* expression in adult injured watershed, Artery Reassembly was absent in these hearts. The precise cause for absence of Artery Reassembly in adult injured hearts remains unknown.

The age-dependent discrepancy in artery EC response triggered by MI, could be because of their distinct molecular profiles[25–27]. To identify potential molecular drivers, we performed single cell RNA sequencing of neonatal and adult ECs, at a time point prior to when collateral arteries appear in the injured mouse hearts. Using both bioinformatics analyses and *in vivo* experiments, we show that, upon injury, neonatal artery ECs undergo de-differentiation and VegfR2-mediated proliferation, which is critical for collateral artery formation by Artery Reassembly. Both cellular events, arterial de-differentiation and proliferation, were absent in older non-regenerative hearts, post-MI. We hope, in future, these findings will aid

strategizing ways to induce collateral arteries and enhance regenerative potential of cardiac tissue, by specifically targeting artery cells.

Results

Increase in cycling neonatal artery ECs, post-MI

Molecular profiling cardiac ECs which participate in collateral artery formation, can improve our understanding of ways in which complete cardiac reperfusion and regeneration can be attained. We thus, performed single cell RNA sequencing of neonatal post-MI heart cells. Specifically, *Cx40CreER; Rosa26^{TdTomato}* mice were used, where a single dose of Tamoxifen to the lactating mothers induced expression of TdTomato fluorescence in *Cx40* expressing artery endothelial cells (aECs) in neonates. Neonates were then subjected to either sham or MI surgeries at P2, and hearts were harvested at P4 or P5 (Figure **1A**). TdTomato$^+$ cardiac cells were sorted using Fluorescent Activated Cell Sorting (FACS) method, followed by sequencing using 10X genomics platform. Sequences were processed using CellRanger and the subsequent output files were analyzed using Seurat on RStudio[28].

Sorting TdTomato$^+$ cells during FACS, enriched our samples for *Cx40CreER$^+$* lineage labelled/traced artery ECs. However, we also obtained some TdTomato$^-$ non-artery cells. Thus, we analyzed *Cadherin5$^+$* ECs, which included, both, TdTomato$^+$ artery ECs, genetically labeled between P0-P2 (Figure **1A**), and TdTomato$^-$ artery ECs (if any), identified by expression of known arterial EC genes. Pre-processing and quality control steps (Figure **S1A**) resulted in 5432 sham and 2200 MI ECs (Figure **1B**-**D**), out of which 4423 sham, and 262 MI ECs, expressed at

least one copy of TdTomato (Figure **1E**, **F**). As previously observed[29,30], compared to sham, we obtained fewer cells upon MI. This is likely due to extensive cell death following coronary ligation and subsequent tissue ischemia.

Unbiased grouping of *Cadherin5*[+] ECs from sham and MI, resulted in 10 distinct clusters (Figure **S1B**). Merged data from sham and MI were projected onto two dimensional space using Uniform Manifold Approximation and Projection (UMAP). The identified cell types were— mature artery ECs (aEC1-4), transient ECs (trans-EC), cycling artery ECs (cyc-aEC), cells undergoing endothelial to mesenchymal transition (endoMT), smooth muscle cells (SMC1,2) and endocardial ECs (endoEC) (Figure **1G**, **H**).

We identified the cell types with known vascular and cardiac cell markers and their relative expression levels (Figure **1I**, **S1**-**3**). aEC1-4 expressed markers for mature artery ECs (*Cx40*, *Cxcr4*) and SMC1,2 expressed markers for both pericytes and SMCs (*Pdgfrb*, *Notch3*). Cells undergoing endoMT were identified by upregulated expression of mesenchymal markers such as *Col1a1* and *Cdh2* and endo-MT associated transcription factors like *Serpine1*, *Twist1* and *Snai1* (Figure **1I** and **S1C**). Some of these markers were co-expressed by SMC clusters (Figure **S1C**). An EC cluster expressed several genes specific to G2/M (*Top2a*, *Mki67*, *Birc5*, *Cenpa*, *Kif2c*, *Bub1*, *Nuf2*, *Hmmr*) (Figure **1I** and **S1D**-**G**) and G1/S (*Tyms*, *Mcm4*, *Mcm6*, *Mcm7*, *Ung*, *Chaf1b*, *Fen1*, *Gmnn*) (Figure **1I** and Figure **S1H**-**K**) transition. Sham cells in this particular cluster mostly expressed genes like *Cx40, Gja4, Jag1, Cxcr4* (Figure **S2A**-**D**) known to be significantly expressed and being specific to mouse[31] and human[32] coronary arteries. The top 20 genes in this cluster included artery specific genes[33] like *Fgfr3*, *Slc45a4*, *Gja4* and *Ssu2* (data not shown). We hence, named the cluster as cycling aEC (cyc-aEC). EndoEC expressed *Npr3*[34],

Cdh11, *Tgfbi*[35] and *Adgrg6*[36] (Figure **1I**) and some venous markers (Figure **S1L**). Non-endothelial cardiac cells were mostly absent in our dataset (Figure **S1L**).

We also identified a cluster of cells (named as trans-EC in Figure **1G**, **H**) that did not express markers for a single endothelial cell sub-type. Re-clustering this group resulted into 4 distinct EC sub-clusters, those highly expressed markers for arteries, capillaries, venous/capillaries and myofibrillar genes[37] (Figure **S3A**-**B**). 42.8% of artery-like ECs, in this cluster, were TdTomato$^+$ (Figure **S3C**) indicating their origin from pre-existing artery ECs, those labelled between P0-P2 (Figure **1A**). Additionally, this artery-like EC sub-cluster had fewer cells expressing *Cx40* or *Cxcr4* (Figure **S3B**). The indistinct molecular nature of this entire cluster, and its proximity to both, a proliferative (cyc-aEC) and mature (aEC1/3) artery EC clusters (Figure **S1B**), led us to hypothesize that cells in this cluster are in an intermediate state, i.e., between a proliferative and mature state. We hence, called this cluster, transitioning EC (trans-EC). Bioinformatics analyses of *Pecam1*$^+$ cells, instead of *Cadherin5*$^+$ cells, resulted in identification of the same cell types, clustered in a similar manner (Figure **S4**).

Artery cells rarely proliferate, and exiting cell cycle is required for artery cell specification[24,38–40]. However, we identified an artery EC cluster (cyc-aEC) in our dataset, expressing several genes that mark G1/S and G2/M transitions (Figure **1I** **and S1D**-**K**). Cell cycle analyses of *Cadherin5*$^+$ cells showed a 12.4% increase in MI cells undergoing G2 to M transition as compared to the sham group (Figure **1J**). The number of MI cells in S-phase increased by 3.3% and G0/G1 phase reduced by 15.6% (Figure **1J**), suggesting a higher number of MI cells were likely mitotic.

We next performed cluster-wise quantification of mitotic cells (expressing G2/M markers), in sham and MI groups. In sham group, 2.5% of all mitotic cells were

present in artery clusters (aEC1-4 and cyc-aEC) (Figure **1K**). In MI group, we observed 11% of all mitotic cells in artery clusters (Figure **1K**), indicating a 4.4-fold increase in mitotic artery ECs, upon MI. 15.9% of all *Cadherin5*[+] cells were mitotic in the MI group, and majority of these cells (~69.2%) were artery ECs (Figure **1K**), a stably differentiated cell type in neonatal hearts.

Figure 1: Single cell analyses of neonatal artery ECs

(**A**) Experimental setup to obtain neonatal cardiac cells for scRNA sequencing using 10X genomics (**B**) UMAP showing the distribution of *Cadherin5*$^+$ neonatal sham and MI cells. (**C**, **D**) Feature Plot showing the distribution of *Cadherin5*$^+$ cells from (**C**) sham and (**D**) MI group. (**E**, **F**) Distribution of TdTomato expressing cells within *Cadherin5*$^+$ (**E**) sham and (**F**) MI group. (**G**, **H**) UMAP showing distribution of (**G**) sham and (**H**) MI cell clusters from the merged *Cadherin5*$^+$ cells. (**I**) Dot Plot showing average expression and percentage of cells, expressing relevant markers used for cluster identification. (**J**) Graph showing percentage of sham and MI *Cadherin5*$^+$ cells in different cell cycle stages based on cell cycle sorting analysis on Seurat. (**K**) Graph showing percentage of *Cadherin5*$^+$ cells undergoing G2 to M transition, in sham and MI groups, in all clusters or in only artery EC clusters (aEC1-4 and cyc-aEC). (**L**) Graph showing percentage of *Cadherin5*$^+$ and *Cx40creER*-linegae traced (TdTomato$^+$) cells undergoing G2 to M transition, in sham and MI groups, in all clusters or in only artery EC clusters (aEC1-4 and cyc-aEC). (**M**) Pie chart showing the percent distribution of G2/M *Cadherin5*$^+$ cells, upon MI. (**N**-**P**) Confocal image of an (**N**) artery tip from a post-MI heart, immunostained for Cx40, lineage traced with TdTomato and labelled with EdU show presence of (**O**) Cx40$^+$ TdTomato$^-$ EdU$^+$ (cyan arrowhead) artery EC and (**P**) Cx40$^+$ TdTomato$^+$ EdU$^+$ (yellow arrowhead) within the same artery branch. MI, myocardial infarction; *Cdh5*, *Cadherin5;* Td, TdTomato; art, artery. Scale bar: **N**-**P**: 50µm

MI-induces new artery development, from sources independent of pre-existing artery ECs

Post-MI artery EC proliferation is associated with Artery Reassembly and complete cardiac regeneration[24]. Use of artery EC specific *Cx40CreER* mouse line, allowed us to mark the lineage of pre-existing artery ECs with TdTomato fluorescent reporter (Figure **1A**). We performed cell cycle analysis, and quantified the number of *Cadherin5*+TdTomato+ ECs that expressed genes for G2/M transition, and were likely to be mitotic. 13.4% of all *Cadherin5*+TdTomato+ MI cells were mitotic, ~63% of which were present in artery EC clusters (aEC1-4 and cyc-aEC) (Figure **1L**). Analyzing mitotic cell distribution within the MI group showed 69.3% of all mitotic cells to be present in artery clusters but very few (6.3%) of them were TdTomato+ artery ECs (Figure **1M**). This suggests that MI induces expansion of an artery EC population, not lineage labelled by, or traced from, *Cx40CreER*+ cells. This is consistent with our previous observation that injury causes organ-wide artery growth (especially, artery-tips), independent of Artery Reassembly, and through capillary differentiation[24].

We next performed *in vivo* experiments to confirm if an artery EC population, originating from non-artery sources (as seen in Figure **1M**) is indeed proliferating, upon MI. EdU labelled P6 hearts from *Cx40CreER; Rosa26^TdTomato* neonates (Tamoxifen at P0, MI at P2) were harvested and immunostained for Cx40. Cx40 immunolabelled all mature artery ECs, including *Cx40CreER* lineage-traced TdTomato+ pre-existing aECs. We indeed, observed numerous Cx40+TdTomato- arteries (Figure **S5A**-**C**), especially in the ischemic regions. Furthermore, we

identified these artery ECs to be proliferating (EdU⁺) and integrated into Cx40⁺ vessels (Figure **1N**-**O** and Figure **S5A**-**C**). As expected, we also observed Cx40⁺TdTomato⁺EdU⁺ artery ECs (Figure **1N**, **P** and Figure **S5D**, **E**). Thus, ischemia can induce formation of new coronary arteries via multiple mechanisms (Figure **S5F**): 1. Artery Reassembly to generate collateral arteries and 2. Organ-wide artery growth by incorporation of capillary ECs, as also observed earlier[24] and/or proliferation of pre-existing (TdTomato⁺) artery ECs.

Neonatal artery ECs de-differentiate upon MI

Stably differentiated cells may attain a less differentiated state and re-enter cell cycle in response to pathophysiological cues. The acquired proliferative abilities allow them to transform into less specialized cell type. They can then, either re-differentiate into cells, within the same lineage, or into cells of a different lineage[41]. This phenomenon can often be detected through changes in gene expression patterns. Presence of an actively cycling differentiated cell cluster in neonates (cyc-aEC) led us to hypothesize that cells in this cluster de-differentiate and contribute to new arteries, in response to MI.

De-differentiation is indicated by loss of differentiation markers, cell cycle re-entry, and expression of early genes, suggesting lack of cell fate specification. We checked if MI cells, when compared to sham cells, manifested any of these de-differentiation indicators. In sham, *Cx40*, a mature artery EC marker, was expressed by cyc-aECs, at levels comparable to mature artery clusters—aEC1-4 (Figure **2A**, **B**). In MI group, cells in cyc-aEC cluster expressed *Cx40*, at levels much lower than what was observed in sham cyc-aEC (Figure **2B**) or the rest of the artery

ECs (aEC1-4) (Figure **2B**, **C**). The percentage of cells expressing *Cx40*, upon MI, reduced in all clusters (Figure **2A**, **C**), but this decrease was most prominent in trans-EC (19% in sham, 0% in MI) and cyc-aEC (57% in sham, 3% in MI) clusters (Figure **S6A**). A similar reduction in number of *Cxcr4*[+] MI cells was observed specifically in cyc-aEC and trans-EC clusters (Figure **S6A**). Thus, there is an overall reduction in number of cells expressing mature artery markers, and their expression levels, in cyc-aEC and trans-EC cluster, upon MI.

We checked if artery ECs, those enter cell cycle upon MI, (as seen in our single cell analyses in Figure **1M**), downregulate Cx40 protein expression, in an *in vivo* setting (Figure **2D**-**F**). We compared Cx40 protein levels in EdU[+] and EdU[-] artery ECs labelled with Cx40 immunostaining in *Cx40CreER; Rosa26[TdTomato]* neonates, 4 days post-MI (Figure **2D**-**F**). Both categories of artery cells– single and associated with lumenized artery vessels, were analyzed. Single aECs observed in ischemic watershed are derived from resident arteries and contribute to Artery Reassembly. We observed that, Cx40 staining was significantly reduced in these EdU[+] TdTomato[+] proliferative single aECs (Figure **2D**, **F**), when compared to non-proliferative single aECs or artery tips (Figure **2F**). In contrast, EdU[+] proliferating aECs, associated with well-formed arteries, did not show any significant change in their Cx40 expression (Figure **2E**). Together, this suggests that Cx40 is downregulated in proliferating single aECs, which could be a prerequisite for single aECs to re-enter cell cycle, upon MI.

During development, an immature coronary vessel plexus undergoes morphogenesis to build coronary arteries[42]. We hypothesized that cyc-aECs, if de-differentiating, would re-express some early genes like markers for venous/capillary ECs, from which they originate during development. We hence,

analyzed sham and MI cells for cardiac microvascular genes such as *Aplnr*[43] and

Car4[32,44]. There was a dramatic increase in number of MI cells expressing *Aplnr* in

artery ECs (aEC1-4) (Figure **2A**, **C**), but the gene was only minimally expressed in

mature artery EC clusters (aEC1-4), upon MI (Figure **2B**). In contrast to this, the total

number of *Aplnr*[+] MI cells was most prominently increased in cyc-aEC (by 2.5 fold)

and trans-EC (by ~2 fold) cluster (Figure **S6A**), and so was the expression of *Aplnr* in

these clusters (Figure **2B**). *Car4* also followed the same pattern (Figure **2A**, **C** and

Figure **S6A**), indicating both cyc-aEC and trans-EC express venous-capillary marker

genes, upon injury.

Angiogenic properties of ECs facilitate vascular remodeling and subsequent

cardiac regeneration. We next investigated if cyc-aEC (or trans-EC) clusters

expressed any pro-angiogenic genes, yet another cellular feature of

de-differentiation. Upon MI, 61% of the cyc-aECs and 47% of trans-ECs expressed

pro-angiogenic *Apln* gene (Figure **2A**, **C** and Figure **S6A**), known to drive vascular

development[45,46] and regeneration[47–49] *in vivo*. Expression of other angiogenic genes

like *Foxm1*[50,51], *Nrp2*[33], *Col15a1*[33], and *Cd34*[52] were enriched in these two clusters

(Figure **S6B**-**E**). Additionally, ECs in cyc-aEC and trans-EC clusters expressed

various capillary-like/venous specific genes such as *Gpihbp1*, *Fabp3*, and *Pcdh17*

(Figure **S6F**-**H**). All changes in expression of *Cxcr4*, *Cx40*, *Aplnr*, *Car4*, *Apln* in

cyc-aEC between sham and MI groups, were found to be statistically significant

using Wilcoxon sum rank test (**Table S1**). Thus, cyc-aEC and trans-EC clusters are

likely to possess angiogenic properties, which are absent in stably differentiated

artery ECs.

Cardiac vascular cells (ECs and SMCs) and cardiomyocytes have common

precursor cells. In addition to classic angiogenic endothelial cell markers, cyc-aEC

and trans-EC cells also expressed cardiac myofibrillar genes (*Tnnt2*, *Myl3* and

Myh6) (Figure **S6I**-**K**). Because cardiac ECs maintain low, but active transcription of

these cardiomyocyte specific markers[37], it is possible that cyc-aECs and trans-ECs

are multi-potent, and could be activated with necessary *in vivo* cues.

The differentiation status of cells and the projected pseudotime, together, can

determine the root of differentiation, and hence can predict the direction of path

taken by cell populations[53]. A package in RStudio – Cellular (Cyto) Trajectory

Reconstruction Analysis using Gene counts and Expression (CytoTRACE), uses the

number of genes and their differential expression, to determine relative differentiation

status of each cell with respect to others[53]. CytoTRACE score for cycling aEC, in MI

dataset, was highest amongst all clusters, making it the least differentiated cluster

(Figure **2G**). We also performed pseudotime analysis of sham and MI cells using

monocle3 package in RStudio. In sham, the predicted trajectory connected trans-EC

to aEC1-4. Sham cyc-aECs were excluded from the projected path (Figure **2H**). In

contrast to sham, in MI, cyc-aEC, trans-EC and aEC1 clusters were associated with

the predicted path (Figure **2I**). If multiple clusters are connected through a single

trajectory, it is likely that, a less specialized cluster differentiates into a terminal cell

type. This prediction is supported by our observations from genetic lineage tracing

experiments *in vivo*, where upon MI, *Cx40CreER* lineage-labeled, proliferating artery

ECs give rise to mature coronary collateral arteries[24] and contribute to organ-wide

artery growth (Figure **S5F**).

We next assessed the relative differentiation status of the 3 clusters

participating in the trajectory obtained for MI cells (Figure **2I**)─ cyc-aEC, tans-EC and

aEC1. We compared the cellular and molecular features indicating de-differentiation,

across these 3 clusters (Figure **S7A**). As expected, cyc-aEC and aEC1 were the

least and most differentiated clusters, respectively (Figure **S7A**). Trans-EC cluster

showed de-differentiation features intermediate between cyc-aEC and aEC1 (Figure

S7A). Thus, it is likely that proliferating (less specialized) artery ECs in cyc-aEC

cluster are differentiating into mature artery ECs in aEC1 (Figure **2I** and **S7A**). These

results were supported by RNA velocity analyses, which, additionally predicts

directionality of projected trajectories based on the ratio of spliced and un-spliced

transcripts (Figure **S7B**, **C**). Together, upon neonatal MI, at least a subpopulation of

mature artery ECs originates from de-differentiation and proliferation of pre-existing/

resident artery cells (Figure **2J**).

A sham (4429)

B MI sham

C MI (1586)

D Low Cx40 expression in EdU+ saECs
single aECs
***

E arteries
ns

F Low Cx40 expression in EdU+ saECs (compared to EdU- artery tip)
Td Cx40 Edu   TdTomato   Cx40   EdU

Low Cx40 expression in EdU+ saECs (compared to EdU- saECs)
Td Cx40 Edu   TdTomato   Cx40   EdU

P6, Cx40CreER; Rosa26TdTomato (MI at P2)

G cyc-aEC is the least differentiated cluster
Cdh5+ MI cells

H Cycling aECs contribute to predicted cell trajectory upon MI
Sham

I MI

J Artery ECs de-differentiate and re-enter cell cycle upon MI

capillary ECs
Artery tips
proliferation
de-differentiation

capillary EC proliferation

single aEC (Td+/Cx40+/low) proliferation

collateral artery

Figure 2: Assessment of de-differentiation in neonatal cycling artery ECs

(**A**) Heatmap showing gene expression of *Apln*, *Aplnr*, *Car4*, *Cx40* and *Cxcr4* across all *Cadherin5*[+] clusters in neonatal sham group. (**B**) Violin plots comparing expression of *Cx40* and *Aplnr* between different EC clusters in sham and MI group. (**C**) Heatmap showing gene expression of *Apln*, *Aplnr*, *Car4*, *Cx40* and *Cxcr4* across all *Cadherin5*[+] clusters in neonatal MI group. (**D-E**) Quantification of cytoplasmic Cx40 in P6, *Cx40CreER*- lineage traced (**D**) single aECs (p=0.0001) and (**E**) artery ECs (p=0.09), 4 days post-MI. (**F**) Confocal images of EdU[+] TdTomato[+] single artery ECs (yellow arrowhead) and EdU[-]TdTomato[+] artery tip EC or saEC (white arrowheads) in P6 *Cx40CreER*; *Rosa26*[TdTomato] hearts, 4 days post-MI. Cyan arrowhead points to EdU[+] Tdtomato[+] saECs with relatively high Cx40 expression. (**G**) Box plots illustrating a relative differentiation state of *Cadherin5*[+] MI cells with scores obtained from CytoTRACE. (**H**, **I**) Trajectories obtained by performing Pseudo time analysis on neonatal *Cadherin5*[+] (**H**) sham cells or (**I**) MI cells, using Monocle3. (**J**) A proposed model where upon MI, pre-existing stably differentiated artery ECs (green) exit artery tips as single artery ECs, de-differentiate (blue), proliferate (red) and coalesce into coronary collateral arteries. saECs are lineage traced from pre-existing artery ECs (TdTomato[+]), but downregulate expression of mature artery marker Cx40 upon de-differentiation. Capillary ECs are shown in grey. P, postnatal day; saECs, single artery endothelial cells. Scale bar: **F**: 50µm

Adult artery ECs fail to proliferate during MI-induced artery neogenesis

New collateral arteries in adult mice, form in response to coronary occlusion within a week of injury[54]. However, adult coronary collateral arteries are fewer in number, have smaller diameters, and are predicted to perfuse poorly than those in neonates[55]. The reason for this age-dependent difference in which artery ECs respond to ischemia is unclear. One possibility could be that adult collateral arteries do not form naturally through Artery Reassembly. Keeping this in mind, we investigated the molecular nature of adult coronary artery ECs, and their derivative cell-types, in sham and MI mice. We performed single cell analyses of cardiac cells sorted from adult *Cx40CreER; Rosa26^TdTomato^* ventricles (Figure **3A**). Similar to neonatal data, we obtained both, TdTomato⁺ and some unlabeled cells. After performing quality control steps (Figure **S8A**), we analyzed 6256 *Cadherin5*⁺ sham and 1690 *Cadherin5*⁺ MI cells (Figure **3B**, **C**). Like neonatal data, our adult dataset was enriched for artery ECs (Figure **3D**, **E**) and we obtained fewer *Cadherin5*⁺ (Figure **3C**) and TdTomato⁺ (Figure **3F**, **G**) cells in MI group than sham.

Based on cell-specific gene expression patterns, we obtained 12 adult cell populations (Figure **3D**); 7 of which, constituted mature artery ECs (aEC1-7), 1 cycling artery EC (cyc-aEC), 1 capillary EC (cap-EC), 1 endoMT and 2 SMC clusters (Figure **3D**, **E**). Unlike neonates, trans-EC cluster was absent in adults. Instead, we identified a prominent adult capillary EC population, which when further clustered generated 3 distinct microvascular EC populations—artery-capEC, capEC, and venous-capEC (Figure **S8B**-**D**). A significant number of art-cap EC cluster (34%) expressed TdTomato fluorescence (Figure **S8D**), indicating an origin from pre-existing mature arteries. Identity of cycling aECs was confirmed with comparable

expression of artery specific genes (*Gja4*, *Cx40*, *Cxcr4*, *Jag1*) in cyc-aEC and aEC1-7 (Figure **S8E**). A similar cell type distribution was also observed when bioinformatics analyses of *Pecam1*+ cells, instead of *Cadherin5*+ cells, was performed (Figure **S8F**-**H**).

Adult cardiomyocytes have limited proliferation capacity[23]. To assess if the same holds true for cardiac ECs, we performed cell cycle analyses on sham and MI adult *Cadherin5*+ ECs. Unlike neonates, we observed only a minimal increase of 2.5% in the number of ECs undergoing G2 to M transition, upon MI (Figure **3H**). Further cluster-wise analyses revealed only 1.8% increase in the mitotic (G2/M) artery ECs (4.5% to 6.3%, Figure **3I**), compared to 8.5% increase observed in neonates (Figure **1K**), upon MI. 4.6% of adult sham cells were mitotic (Figure **3I**), suggesting that the pre-injury mitotic baseline for adult *Cadherin5*+ artery ECs was higher than that observed in neonates (2.5%, Figure **1K**). Thus, in comparison to neonates, fewer adult artery ECs undergo G2/M transition in response to MI.

P7 or older hearts are non-regenerative[22,23] and do not illicit Artery Reassembly[24]. Cell cycle analyses of *Cadherin5*+ TdTomato+ lineage-traced adult ECs revealed minimal changes in G2/M ECs including in artery EC clusters (aEC1-7, cyc-aEC) (Figure **3J**). Furthermore, we checked which cell types originate from pre-existing artery ECs, upon MI. For this, we analyzed all TdTomato+ cells (labelled and traced from *Cx40CreER*+ resident artery ECs) in adult MI dataset, and compared it with neonatal MI dataset (Figure **3K**). While the neonatal MI group showed a distinct cyc-aEC cluster, this population was absent in the adult MI group (Figure **3K** and Figure **S8I**, **J**). Thus, adult hearts lack a proliferating artery EC sub-population, known to be crucial for Artery Reassembly, and complete cardiac regeneration[24].

We next performed *in vivo* experiments to check if lack of proliferating artery cell population in injured adult hearts (Figure **3J**, **K**), is also observed in non-regenerative P7 hearts. *Cx40CreER; Rosa26*^{TdTomato} neonates were administered with Tamoxifen to lineage label artery ECs at P5, through P7, followed by MI at P7 (Figure **3L**). Proliferation status of *Cx40CreER*$^+$ lineage traced, TdTomato$^+$, artery ECs were analyzed with EdU label, prior to harvesting hearts at P11, 4 days post-MI. As shown earlier[24], these hearts did not demonstrate Artery Reassembly (Figure **3L**, **M**), which was evident from lack of lineage traced (TdTomato$^+$) collateral arteries and single artery ECs in the injured watershed regions (Figure **3M**). Very few single artery ECs were observed adjacent to artery tips (arrowheads, Figure **3M**); we called them 'residual' single artery ECs. EdU$^+$ cells in TdTomato$^+$ artery EC population—in both, artery tips (Figure **3N**) and residual single artery ECs in the watershed (Figure **3O**)—were rarely observed. Together, data from both bioinformatics analyses and *in vivo* experiments indicate that artery ECs in non-regenerative P7 or adult hearts have limited self-renewing capacities.

Figure panels:

A Cx40CreER; Rosa26^TdTomato — Tam d0; sham or MI d2; d11 d13 digest hearts; sort TdTomato+ cells; 10X genomics (scRNA-seq)

B Cdh5

C Cdh5⁺ cells — sham (6256 cells); MI (1690 cells)

D Merge — SMC1, SMC2, cap-EC, aEC3, aEC4, aEC2, aEC1, endo MT, cyc aEC, aEC7, aEC5, aEC6

E Av. exp; Perc. expressed

F TdTomato sham (2912 cells); cyc aEC

G MI (567 cells); cyc aEC

Minimal increase in mitotic artery ECs upon MI

H Cell cycle analyses of all Cdh5+ cells — G2/M, S, G0/G1; up 2.5%; down 10.3%; up 7.7%

I Cdh5⁺ cells — all clusters 4.6% 7.1%; art. clusters 4.5% 6.3%; mitotic, non-mitotic

J Cdh5⁺Td⁺ (lineage-traced) cells — all clusters 3.9% 4%; art. clusters 3.8% 3.7%; mitotic, non-mitotic

Absence of cyc-aEC cluster in adult MI group

K Td+ (lineage-traced) cells — aECs, cyc-aECs, cap-EC, endo-MT, SMC, endo-EC, CM; Neo, Adt

Lack of saECs in non-regenerative hearts, post-MI

L Cx40CreER; Rosa26^TdTomato 4 days post-MI (MI at P7) — ligated LCA, ligation, RCA branches, scar; *ligated branches of LCA

M watershed — residual saECs, ligated artery branch, RCA branch, residual saECs

Minimal proliferation of Td+ (lineage traced) aECs with P7 MI

N No. of EdU+ Td+ ECs per Td+ artery tip — artery tips

O % EdU+ Td+ single artery ECs — residual saECs

Figure 3: Cell cycle analyses of adult artery ECs post-MI

(**A**) Experimental setup to obtain adult cardiac cells for scRNA sequencing using 10X genomics. (**B**) Feature Plot showing the distribution of adult *Cadherin5*$^+$ cells from both sham and MI groups. (**C**) UMAP showing the distribution of *Cadherin5*$^+$ adult sham and MI cells. (**D**) UMAP showing distribution of both sham and MI cell clusters from the merged adult *Cadherin5*$^+$ cells. (**E**) DotPlot showing average expression and percentage of cells, expressing relevant markers used for cluster identification in **D**. (**F**, **G**) Distribution of TdTomato expressing cells within *Cadherin5*$^+$ (**F**) sham and (**G**) MI group. (**H**) Graph showing percentage of adult, sham and MI, *Cadherin5*$^+$ cells in different cell cycle stages based on cell sorting analysis on Seurat. (**I**) Graph showing percentage of *Cadherin5*$^+$ cells undergoing G2 to M transition, in sham and MI groups, in all clusters or in only artery EC clusters (aEC1-7 and cyc-aEC). (**J**) Graph showing percentage of *Cadherin5*$^+$ and *Cx40creER*-linegae traced (TdTomato$^+$) cells undergoing G2 to M transition, in sham and MI groups, in all clusters or in only artery EC clusters (aEC1-7 and cyc-aEC). (**K**) Single cell data analyses of all TdTomato$^+$ cells from neonatal and adult MI groups, showing distribution of different cell types that originate from *Cx40CreER* lineage labelling or subsequent tracing. (**L**, **M**) Representative image of P11 *Cx40CreER*; *Rosa26*^{TdTomato} (**L**) whole heart and respective (**M**) watershed region, 4 days post-MI. MI was performed at P7. Lineage traced artery ECs are shown in black. * marks ligated branches of LCA, arrowheads point to "residual" saECs. (**N**, **O**) Graph showing number and percentage of EdU$^+$ Tdtomato$^+$ ECs in (**N**) artery tips and (**O**) residual saECs, respectively, at P11. MIs were performed at P7. P, postnatal day; LCA, left

coronary artery; RCA, right coronary artery; art., artery; Adt, adult; Neo, neonate;

saECs, single artery endothelial cells. Scale bars: **L**: 500μm; **M**: 200μm

Adult artery ECs do not de-differentiate, upon MI

We next checked if adult injured *Cadherin5*$^+$ cycling aECs (Figure **3I**)
demonstrate any characteristics indicative of de-differentiation. Though adult MI
cycling aEC cluster was the least differentiated aEC cluster (Figure **4A**, **B**), its
CytoTRACE score was comparable to a mature artery cluster—aEC5 in MI (Figure
4B). Expression of proangiogenic marker *Apln*, and its receptor *Aplnr* were
seemingly upregulated in this cluster, but there was no change in expression of
Car4, *Cx40*, or *Cxcr4* (Figure **4C**). We analyzed the cellular and molecular features
of neonatal cyc-aECs, which were indicative of de-differentiation, and checked those
indicators in adult cyc-aECs (Figure **4D**). Some of the key de-differentiation features
absent in adult cyc-aECs were—1. their adequate number, which is representative of
their ability to self-renew/ re-enter cell cycle 2. loss of mature cell markers and 3.
gain of microvascular/ pro-angiogenic markers (shown in red, Figure **4D**). For
instance, in contrast to neonatal MI cyc-aECs, adult MI cyc-aECs, showed no
change in the number of *Cx40* expressing cells (Figure **4D**). Cells expressing *Cxcr4*
in this group, increased by 38% (Figure **4D**). The number of cells expressing *Apln*,
Aplnr, and *Car4* in the adult cyc-aECs were increased upon MI. However, these
changes in cell numbers or gene expression levels, were statistically insignificant
when compared between adult sham and MI groups, using Wilcoxon sum rank test (*
in Figure **4D and Table S1**). Furthermore, cycling aEC cluster did not contribute to
the predicted path associated with mature aECs (aEC1-7), even in MI group (Figure
4E, **F**). Thus, unlike neonate, adult cycling aECs were neither connected to artery
ECs in trajectory analyses (Figure **4E**, **F**), nor were lineage traced from pre-existing
arteries (Figure **3J**, **K**). Together, adult cycling aEC population fails to drive artery

regeneration (Figure **4G**) due to its limited number and inability to de-differentiate

(Figure **4D**).

De-differentiation features present in neonatal and adult artery ECs
Cellular and molecular features absent in adult artery ECs
* Statistically insignificant (sham vs. MI) as confirmed with Wilcoxon sum rank test (Table S1)
In vivo evidence

Figure 4: Analyses of de-differentiation features in adult MI cells

(**A**-**B**) Box plots illustrating a relative differentiation state of adult *Cadherin5*⁺ (**A**) sham and (**B**) MI cells, with scores obtained from CytoTRACE. (**C**) Heatmap of adult sham and MI cells showing the expression of *Apln*, *Aplnr*, *Car4*, *Cx40* and *Cxcr4*, in their respective cyc-aEC cluster. (**D**) Comparison of de-differentiation features in *Cadherin5*⁺ neonatal and adult cyc-aEC cluster. (**E**-**F**) Pseudo time analysis of adult *Cadherin5*⁺ cells from (**E**) sham and (**F**) MI group. (**G**) Schematic showing post-MI neonatal artery ECs accomplish artery regeneration by undergoing de-differentiation and proliferation; a phenomenon rarely observed in adult. Dotted line, minimal contribution. FC, Fold Change; av., average; diff, differentiation; Td, TdTomato; #, number**.**

Arterial VegfR2 promotes artery EC proliferation during Artery Reassembly

Pre-existing artery ECs drive collateral artery formation in neonatal mice, only upon MI[24,56]. We isolated all genes from *Cx40CreER*-lineage traced TdTomato[+] injured artery ECs (Figure **1F**) and performed gene ontology enrichment analysis and visualization (GOrilla). This allowed us to identify biological processes triggered upon MI. Key vascular responses such as angiogenesis, migration and proliferation, were significantly upregulated upon MI (Figure **5A**). Artery Reassembly is a systematic step-wise cellular event where artery EC migration and proliferation are crucial. Thus, our neonatal single cell dataset contains the artery EC population, including those that drive Artery Reassembly, and subsequent neonatal cardiac regeneration.

aEC1, in neonatal MI dataset, is the only artery EC cluster associated with the predicted trajectory and connected with cycling artery ECs (Figure **2I**). We performed pathway analyses on *Cadherin5*[+] aEC1 cells using g:Profiler and filtered pathways from various databases, based on their statistical significance (p>0.05). Many pathways relevant to vascular development and injury response were enriched in aEC1 such as Apelin, TNF and Interleukin signaling[57] (Figure **5B**). Vegfa-VegfR2 signaling (p=0.0011) was of particular interest (Figure **5B**), as it has established molecular functions in mouse coronary artery development[40,58] and neovascularization upon injury in different tissues[59,60].

We checked the expression levels of ligands (*Vegfa*, *Vegfc*) and a major receptor (*VegfR2* or *Kdr*) associated with coronary morphogenesis, in our dataset. While *Vegfa* and *Vegfc* were expressed in artery ECs (aEC1-4), *VegfR2* was specifically enriched in cycling aEC and trans-EC population (Figure **5C**). The

VegfR2 expression, was significantly different in cyc-aECs, when compared between neonatal sham and MI groups, using Wilcoxon sum rank test (**Table S1**). When immunostained for VegfR2, the protein was located in coronary artery ECs of small arteries, artery tips and capillary ECs (Figure **S9A**). The levels of VegfR2 protein in arteries was lower as compared to capillaries, and the protein was almost always absent in large arteries (Figure **S9A**). Both EC types (arteries and capillaries) facilitate Artery Reassembly, either by directly participating in the process (artery ECs) or providing the Cxcl12 ligand (capillary ECs). Given VegfR2 expression in both, capillaries and small arteries, we hypothesized that Vegf pathway has a role in Artery Reassembly, post-MI.

We analyzed the role of arterial *VegfR2* in Artery Reassembly, *in vivo*. Control (*Cx40CreER*; *Rosa26^{TdTomato}*) and artery EC-specific, conditional *VegfR2* knockout (*VegfR2$^{L/L}$*; *Cx40CreER*; *Rosa26^{TdTomato}*) neonates received Tamoxifen at P0/P1 which allowed labelling of pre-existing artery ECs. These hearts were subjected to MI at P2/P3 and hearts were harvested 4 days post-MI, at P6/P7 (Figure **5D**). Whole heart imaging allowed us to visualize the entire arterial network of the heart (Figure **5E**, **F**) as well as focus into regions of interest such as the watershed region where new collaterals are expected to appear after injury (Figure **5G**-**I**). Control and *VegfR2* artery specific knockout hearts did not illustrate any gross differences in their baseline artery architecture, without MI (Figure **S9B**-**D**). Watershed regions from sham (data not shown) or no injury (Figure **5G**) hearts, did not show any collateral arteries lineage labelled with TdTomato, suggesting absence of Artery Reassembly in non-ischemic hearts. Upon MI, control hearts consistently showed several (average of 8) *Cx40CreER*-lineage traced (TdTomato[+]) collateral arteries as observed earlier[24] (Figure **5E**, **H**). In contrast, we rarely observed any such collateral

arteries in *VegfR2* artery specific knockout hearts (Figure **5F**, **I**, **J**). This suggests an artery specific role of VegfR2 in executing Artery Reassembly, upon MI.

We next investigated the cause for lack of TdTomato$^+$ collateral arteries in *VegfR2* knockout watershed regions. Artery Reassembly is a three step cellular event—migration, proliferation and coalescence, of artery ECs. We first assessed the ability of TdTomato$^+$ cells to migrate into the watershed. Qualitative and quantitative observations suggested that the number of *Cx40CreER*-lineage traced single artery ECs, observed in the watershed regions, were comparable in control and knockout MI hearts (Figure **5H**, **I** and Figure **S9E**-**H**). This suggests that in absence of arterial VegfR2, the migration of artery ECs from their resident arteries into the watershed region is less likely affected. We next investigated the proliferation step, and injected neonates with EdU, one hour prior to harvesting the hearts. We then, quantified the number of proliferating single aECs in the MI watershed regions. Interestingly, *VegfR2* knockout watershed regions showed a ~2.4-fold reduction in the number of proliferating single artery ECs as compared to control hearts (Figure **5K**-**M**). Such reduction in number of proliferating single artery ECs was also observed in non-regenerative P7 injured hearts (Figure **3O**). Thus, arterial VegfR2 regulates neonatal artery EC proliferation, post-MI. A defect in VegfR2-mediated artery cell proliferation could also affect subsequent coalescence of TdTomato$^+$ cells and successful execution of Artery Reassembly, as observed in *VegfR2* knockout hearts (Figure **5J**).

We next investigated if Vegf ligands and receptors are expressed in adult ECs. The expression of Vegf ligands, *Vegfa* and *Vegfc*, were mostly observed in mature artery EC clusters (Figure **S10A**) like in neonatal data (Figure **5C**). The receptor *VegfR2* (*Kdr*), was upregulated in capillary ECs, as expected (Figure **S10A**).

Interestingly, at least at the mRNA level, *VegfR2* was downregulated in ~75% of actively cycling aECs (Figure **S10A**). This is in contrast to the upregulated expression of *VegfR2* in neonatal cyc-aECs (Figure **5C**). When compared to sham group, *VegfR2* was significantly downregulated in adult MI cyc-aEC cluster (**Table S1**). Given the functional significance of arterial *VegfR2* in neonatal aEC proliferation (Figure **5K**-**M**), the lack of *VegfR2* expression in "residual" adult cycling aECs, could explain depleted potential for aEC proliferation in injured adults. In summary, our data indicate that arterial VegfR2 drives, specifically, the proliferation step of Artery Reassembly in injured neonatal mice (Figure **S10B**).

A TdTomato+ pre-existing aECs express proliferation/migration genes

GOrilla analyses

- cell-cell adhesion
- extracellular structure organization
- angiogenesis
- actomyosin structure organization
- positive regulation of vasculature development
- negative regulation of cell death
- response to wounding
- cellular response to growth factor stimulus
- cell migration
- regulation of cell proliferation

p value

B VegfR2 signaling in aEC1

g:Profiler output

- IL4 and IL13 signaling
- Regulation of IGF transport
- Vegfa-VegfR2 signaling
- MAPK signaling
- Apelin signaling
- TNF signaling
- IL18 signaling

p value

C Expression of VegfR2 in cycling artery ECs

aEC1, aEC2, aEC3, aEC4, trans-EC, cyc-aEC, SMC1, SMC2, endoMT, endoEC

Vegfa, *Vegfc*, *Kdr*

Average Expression
Percent Expressed (25, 50, 75)

Artery endothelial *VegfR2* regulate Artery Reassembly

D *Cx40CreER; Rosa26^TdTomato* or *VegfR2^L/L;Cx40CreER; Rosa26^TdTomato*

lineage label pre-existing aECs
lineage trace labelled aECs in the WS

P0/P1 — P2/P3 — P6/P7

Tam — MI — assess Artery Reassembly

E *Cx40CreER; Rosa26^TdTomato*

ligated artery, ligation, RCA branches, H

F *VegfR2^L/L Cx40CreER; Rosa26^TdTomato*

ligated artery, ligation, RCA branches, I

Cx40CreER; Rosa26^TdTomato

G No MI

watershed, RCA branch, LCA branch

H MI

RCA branch, collateral, ligated artery branch

VegfR2^L/L;Cx40CreER; Rosa26^TdTomato

I MI

RCA branch, saECs, ligated artery branch

J Td+ collaterals per heart

**** * ****

VegfR2^+/+, *VegfR2^+/-*, *VegfR2^L/L*

Cx40CreER; Rosa26^TdTomato

K % EdU+Td+ saECs in ischemic WSA per heart

**

VegfR2^+/+, *VegfR2^L/L*

Cx40CreER; Rosa26^TdTomato

L *Cx40CreER; Rosa26^TdTomato*

Td+ saECs EdU

M *VegfR2^L/L Cx40CreER; Rosa26^TdTomato*

Td+ saECs EdU

Figure 5: Vegf pathway in neonatal Artery Reassembly

(**A**) Results from Gene Ontology analysis on TdTomato$^+$ cells from neonatal MI group (with sham as background), using the tool GOrilla. (**B**) Pathway analysis on genes expressed in neonatal aEC1 cluster, using g:Profiler. (**C**) DotPlot showing average expression and percentage of cells expressing *Vegfa*, *Vegfc* and *Kdr* (*VegfR2*) in all *Cadherin5$^+$* cells. (**D**) Experimental set up to test the role of arterial VegfR2 in neonatal Artery Reassembly. (**E**, **F**) Representative image of (**E**) control and (**F**) *VegfR2* artery endothelial cell specific knockout whole hearts, 4 days post-MI. (**G**-**I**) Representative confocal images of P6 watershed regions from (**G**) un-injured, (**H**) MI control and (I) MI *VegfR2* knockout hearts. MIs were performed at P2. Lineage traced artery ECs are shown in black. (**J**) Quantification of *Cx40CreER*- lineage traced collateral artery numbers observed 4 days post-MI. (**K**) Quantification of percentage of EdU$^+$ Tdtomato$^+$ saECs in watershed regions from *Cx40CreER*-lineage traced hearts, 4 days post-MI. (**L**, **M**) Representative confocal images of EdU$^+$ Tdtomato$^+$ saECs (white arrowheads) in watershed regions from *Cx40CreER*-lineage traced hearts, (**L**) with or (**M**) without arterial *VegfR2*, 4 days post-MI. LCA, left coronary artery; RCA, right coronary artery; WSA, watershed Area. Scale bars: **E**, **F**: 500µm; **G**-**I:** 100 µm; **L**, **M**: 50 µm

Discussion

In this study we utilize both *in silico* analyses and *in vivo* experiments to explain the age-dependent artery EC responses to cardiac injury. Firstly, with single cell sequencing data analyses we show, for the first time, that coronary artery ECs, an otherwise stably differentiated cardiac cell type, undergoes injury-induced de-differentiation in neonates, but not in adults. With the combination of bioinformatics and mouse experiments, we show that the ability of artery ECs to de-differentiate, correlates with their ability to proliferate. Thus, de-differentiation could be a pre-requisite for proliferation, and together, both cellular responses (which are specific to neonatal regenerative window) drive MI-induced artery regeneration. Secondly, using *in vivo* experiments, we discover a second artery-specific receptor (apart from Cxcr4, which controls artery cell migration during Artery Reassembly[24]), which regulates proliferation of artery ECs in response to neonatal MI. Expression of this receptor, *VegfR2*, is downregulated in adult cycling artery ECs and could explain absence of Artery Reassembly in adults. Finally, our data suggests that, each step of Artery Reassembly is regulated by a distinct signaling axis, and the successful execution of each step will require activation of these molecular pathways in the right space and time. In summary, we highlight the differences in age-dependent molecular profiles of mouse coronaries and propose a mechanism for reduced plasticity of artery ECs and lack of Artery Reassembly in non-regenerative P7 or adult hearts.

De-differentiation and Proliferation are Coupled during Cardiac Regeneration

New cardiomyocytes originate from pre-existing cardiomyocytes upon injury[22,23] by de-differentiating and re-entering cell cycle[61,62]. Cardiomyocyte de-differentiation has been elucidated upon adult zebrafish cardiac injury via disassembly of sarcomeric structures, detachment of cardiomyocytes from one another, and expression of early embryonic genes (α-Smooth Muscle Actin (αSMA), embryonic Myosin Heavy Chain, and Gata4)[61–63]. Post-MI adult cardiomyocytes also demonstrate somewhat limited de-differentiation potential indicated by downregulation of mature cardiomyocyte functions such as hypertrophy, contractility, electrical conduction, and upregulation of several early genes like *αSMA*, *Gata4*, *Runx1* and *Dab2*[64]. Unlike cardiomyocytes, EC de-differentiation has not been reported in any cardiac injury model. We show here, for the first time, that neonatal mouse coronary artery ECs undergo de-differentiation upon MI; a process limited in adults.

Since there are no objective indicators of artery EC pluripotency, one can argue that changes in gene expression or phenotypes could be a reflection of their plasticity, and not necessarily reversion of cell fate into a de-differentiated state. Here, we combine single cell analyses and genetic lineage tracing to address this issue. Lineage labelled non-cycling injured artery ECs can be sequentially traced into cycling single artery ECs in the watershed, and consequently into non-cycling artery ECs on the coronary collaterals. We analyzed the transiently cycling artery ECs and show that several indicators for de-differentiation, were enhanced in neonates but were absent in adults. This finding positively correlated with the ability of artery ECs to re-enter cell cycle. Neonates had much higher number of cycling artery ECs than P7 or adults. This finding was verified via two independent methods—single cell gene expression analyses of cyc-aECs and presence of TdTomato[+] EdU[+] cells in an

in vivo cardiac MI model. Thus, lack of de-differentiation limits the number of cycling aECs in non-regenerative hearts and consequently affects their ability to form collateral arteries.

Reactivation of Developmental Pathways upon Injury?

Several molecular pathways relevant to vascular development regulate EC response to injury[65]. For instance, Notch[66], Apelin[45], Cxcl12[67–70] signaling, in mice, regulate different aspects of vascular development and remodeling during cardiac morphogenesis; ranging from EC fate determination to regulation of different EC responses such as migration, proliferation and tube formation. Interestingly, some of these pathways are also associated with EC regeneration upon injury[1,24,48], making them potential therapeutic candidates for treating vascular conditions in human patients[71]. Further studies are required, and will elucidate if (and how) the cellular and molecular mechanisms associated with these signaling pathways, during development and injury responses, are similar.

Vegf signaling drives coronary development in zebrafish and mouse hearts. In mice, Vegf signaling drives migration of coronary endothelial[43,72] and endocardial endothelial cells[58] in developing hearts, and proliferation of coronary ECs in heart explant cultures[73]. In zebrafish, *vegfa* mutants show reduced EC proliferation possibly via PI3K signaling[74]. Upon cardiac injury, endocardial Vegfa signaling[1] and Vegfc-mediated (Cxcl8a/Cxcr1-driven) coronary EC proliferation[75] execute revascularization and cardiac regeneration of ischemic hearts. In this study, we highlight the role of VegfR2-mediated signaling in artery EC response triggered by neonatal cardiac injury. We show that VegfR2 in neonates is expressed in coronary

artery ECs, deletion of which, significantly reduced the number of collateral arteries formed via Artery Reassembly.

Our data show that artery tips do not proliferate. Proliferation is rather demonstrated by single aECs which originate from these artery tips. This suggests that migration of pre-existing artery ECs could precede artery EC proliferation in the watershed. Appearance of *VegfR2* depleted single artery ECs in ischemic watershed regions suggested that artery EC migration was not affected. VegfR2 depleted single artery ECs showed reduced proliferation potential, which could account for lack of lineage-traced collateral arteries in the knock out hearts. Proliferation of artery ECs is followed by their coalescence into collateral arteries. Thus, we cannot exclude the possible role of Vegf in coalescence of artery cells, or the vascular steps that follow, like lumenization or maturation of collateral arteries.

The precise source of Vegf ligand in injured mouse hearts could be the endocardium[76], myocardium (Vegfa)[58,77], epicardium (Vegfc[43], or Vegfa[78]), or cardiac endothelial cells[75], thus suggesting an autocrine or paracrine signaling as observed in injured zebrafish hearts[75]. In our neonatal mouse single cell dataset, we found both *Vegfa* and *Vegfc* being expressed by mature artery ECs (aEC1-4), but at lower levels by cyc-aECs (or by trans-ECs, SMCs, endoMT ECs, endo ECs), indicating a possible cell autonomous mechanism. However, an alternate source of Vegf ligand, such as, cardiomyocytes, fibroblasts or immune cells, which are excluded from our dataset, cannot be neglected.

Role of Vegf Pathway in Human Heart Disease and its Treatment

Single nucleotide polymorphisms (SNPs) in both introns and exons of *KDR* (murine *VegfR2*) are strongly associated with coronary heart disease (CHD)[79]. These SNPs can either modulate the affinity to ligand, or alter E2F binding to *KDR* promoter, subsequently, affecting *KDR* activity and downstream signaling[79]. Elevated levels of circulating VEGF ligands are positively and significantly associated with CHD[80], coronary artery disease (CAD)[81,82] and coronary artery lesions[83,84]. Several SNPs affect VEGF levels[80,82] and may account for the variation observed in revascularization abilities in individuals across different genetic backgrounds. At least three such SNPs in *VEGFA* are strongly associated with increased risk of CHD/MI[85] and one with coronary collateral vessel perfusion[86]. How the association of VEGF/KDR signaling axis with CHD may serve for better prognosis or indicate the need for vascularization, remains unclear.

Preclinical studies in animals with sparse[87] to no[54] pre-existing collateral arteries have consistently elucidated role of Vegf in ischemia-induced collateral formation. On the contrary, numerous past and ongoing clinical trials with *VEGF* alone, or combined with pro-angiogenic/ regenerative factors, have been concluded with little to no success (data available at clinicaltrials.gov)[87,88]. The delivery methods appear to be safe, but the angiogenic effects, particularly the efficacy of *VEGF* gene therapies in stimulating reperfusion via collateral arteries is minimal[87]. Mechanistic details of Vegf's mode of action, in orchestrating vascular cell responses that lead to collateral formation, could aid designing clinical trials with higher efficacies.

We show that injured mouse coronary ECs, de-differentiate and then expand in VegfR2-dependent manner. The mere presence of a significant number of single artery ECs in a limited watershed area could be sufficient to create new arteries[31]. Thus, it will be interesting to see if this threshold density of artery ECs is

accomplished by Vegf-mediated proliferation. Similarly, identifying the molecular driver of EC de-differentiation is also crucial, and when combined with Vegf, could promote collaterals in individuals with CHD and CAD.

Authors Contribution

SD and KR conceived the idea. SD, KR, YJW provided resources. GA performed experiments with neonatal MI model. Sneha K and Suraj K performed bioinformatics analyses. SD, GA and Sneha K wrote the manuscript. HW performed adult MI surgeries. PERC performed some neonatal MI surgeries. SD and KMG performed FACS.

Acknowledgements

We thank the following facilities for their technical support: Stanford Genome Sequencing Center, Stanford shared FACS facility, Animal Care and Resource Center at NCBS, Mouse Genome Engineering Facility at NCBS, Central Imaging and Flow Cytometry Facility at NCBS

Funding

This work is supported by NCBS/TIFR core funding and DBT/Wellcome Trust India Alliance Intermediate Fellowship to SD (IA/I/20/2/505205). YJW is supported by National Institutes of Health (5R01HL089315-11). KR is supported by National Institutes of Health (R01-HL128503) and is a HHMI Investigator. HW is supported by

American Heart Association (18POST33990223). KMG is supported by Gabilan

Stanford Graduate Fellowship, NSF GRFP. Suraj K is supported by DBT/Wellcome

Trust India Alliance Intermediate Fellowship to SD. P.E.R.C is supported by the

NIGMS of the National Institutes of 1121 Health (NIH T32GM007276) and

NSF-GRFP (DGE-1656518).

Declaration of interests

None

Bibliography

1. Marín-Juez, R. *et al.* Fast revascularization of the injured area is essential to support zebrafish heart regeneration. *Proc. Natl. Acad. Sci. U. S. A.* **113**, 11237–11242 (2016).

2. Ingason, A. B. *et al.* Angiogenesis Precedes Cardiomyocyte Migration in Regenerating Mammalian Hearts. *J. Thorac. Cardiovasc. Surg.* **155**, 1118-1127.e1 (2017).

3. He, L. *et al.* Preexisting endothelial cells mediate cardiac neovascularization after injury. *J. Clin. Invest.* **127**, 2968–2981 (2017).

4. Li, Z. *et al.* Single-cell transcriptome analyses reveal novel targets modulating cardiac neovascularization by resident endothelial cells following myocardial infarction. *Eur. Heart J.* **40**, 2507–2520 (2019).

5. Kobayashi, K. *et al.* Dynamics of angiogenesis in ischemic areas of the infarcted heart. *Sci. Rep.* **7**, 1–13 (2017).

6. Betgem, R. P. *et al.* Intramyocardial haemorrhage after acute myocardial infarction. *Nat. Rev. Cardiol.* **12**, 156–167 (2015).

7. Buschmann, I. & Schaper, W. Arteriogenesis versus angiogenesis: Two mechanisms of vessel growth. *News Physiol. Sci.* **14**, 121–125 (1999).

8. Seiler, C., Stoller, M., Pitt, B. & Meier, P. The human coronary collateral circulation: Development and clinical importance. *Eur. Heart J.* **34**, 2674–2682 (2013).

9. Sabia, P. J. *et al.* An Association between Collateral Blood Flow and Myocardial Viability in Patients with Recent Myocardial Infarction. *N. Engl. J. Med.* **327**, 1825–1831 (1992).

10.   Meier, P. *et al.* Beneficial effect of recruitable collaterals: A 10-year follow-up study in patients with stable coronary artery disease undergoing quantitative collateral measurements. *Circulation* **116**, 975–983 (2007).

11.   Habib, G. B. *et al.* Influence of coronary collateral vessels on myocardial infarct size in humans. Results of phase I thrombolysis in myocardial infarction (TIMI) trial. The TIMI Investigators. *Circulation* **83**, 739–746 (1991).

12.   Clayton, J. A., Zhang, H., Pomp, D., Faber, J. E. & Chalothorn, D. Collateral density, remodeling, and VEGF-A expression differ widely between mouse strains. *Physiol. Genomics* **30**, 179–191 (2007).

13.   Teunissen, P. F. A., Horrevoets, A. J. G. & van Royen, N. The coronary collateral circulation: Genetic and environmental determinants in experimental models and humans. *J. Mol. Cell. Cardiol.* **52**, 897–904 (2012).

14.   Maxwell, M. P., Hearse, D. J. & Yellon, D. M. Species variation in the coronary collateral circulation during regional myocardial ischaemia: A critical determinant of the rate of evolution and extent of myocardial infarction. *Cardiovasc. Res.* **21**, 737–746 (1987).

15.   Chalothorn, D. & Faber, J. E. Formation and maturation of the native cerebral collateral circulation. *J. Mol. Cell. Cardiol.* **49**, 251–259 (2010).

16.   Perovic, T., Harms, C. & Gerhardt, H. Formation and Maintenance of the Natural Bypass Vessels of the Brain. *Front. Cardiovasc. Med.* **9**, 1–6 (2022).

17.   Gabhann, F. Mac & Peirce, S. M. Collateral capillary arterialization following arteriolar ligation in murine skeletal muscle. *Microcirculation* **17**, 333–347 (2010).

18.   Scholz, D. *et al.* Contribution of arteriogenesis and angiogenesis to postocclusive hindlimb perfusion in mice. *J. Mol. Cell. Cardiol.* **34**, 775–787

(2002).

19.	Zhang, H., Prabhakar, P., Sealock, R. & Faber, J. E. Wide genetic variation in the native pial collateral circulation is a major determinant of variation in severity of stroke. *J. Cereb. Blood Flow Metab.* **30**, 923–934 (2010).

20.	Chalothorn, D. & Faber, J. E. Strain-dependent variation in collateral circulatory function in mouse hindlimb. *Physiol. Genomics* **42**, 469–479 (2010).

21.	Clayton, J. A., Chalothorn, D. & Faber, J. E. Vascular endothelial growth factor-a specifies formation of native collaterals and regulates collateral growth in ischemia. *Circ. Res.* **103**, 1027–1036 (2008).

22.	Porrello, E. R. *et al.* Transient regenerative potential of the neonatal mouse heart. *Science (80-.).* **331**, 1078–1080 (2011).

23.	Porrello, E. R. *et al.* Regulation of neonatal and adult mammalian heart regeneration by the miR-15 family. *Proc. Natl. Acad. Sci.* **110**, 187–192 (2013).

24.	Das, S. *et al.* A Unique Collateral Artery Development Program Promotes Neonatal Heart Regeneration. *Cell* **176**, 1128-1142.e18 (2019).

25.	Aurora, A. B. *et al.* Macrophages are required for neonatal heart regeneration Find the latest version : Macrophages are required for neonatal heart regeneration. **124**, 1382–1392 (2014).

26.	Wang, Y. *et al.* Single-cell analysis of murine fibroblasts identifies neonatal to adult switching that regulates cardiomyocyte maturation. *Nat. Commun.* **11**, (2020).

27.	Wang, Z. *et al.* Cell-Type-Specific Gene Regulatory Networks Underlying Murine Neonatal Heart Regeneration at Single-Cell Resolution. *Cell Rep.* **33**, 108472 (2020).

28.	Hao, Y. *et al.* Integrated analysis of multimodal single-cell data. *Cell* **184**,

3573-3587.e29 (2021).

29.   Farbehi, N. *et al.* Single-cell expression profiling reveals dynamic flux of cardiac stromal, vascular and immune cells in health and injury. *Elife* **8**, (2019).

30.   Zhuang, L., Lu, L., Zhang, R., Chen, K. & Yan, X. Comprehensive Integration of Single-Cell Transcriptional Profiling Reveals the Heterogeneities of Non-cardiomyocytes in Healthy and Ischemic Hearts. *Front. Cardiovasc. Med.* **7**, (2020).

31.   Raftrey, B. *et al.* Dach1 Extends Artery Networks and Protects Against Cardiac Injury. *Circ. Res.* 702–716 (2021) doi:10.1161/circresaha.120.318271.

32.   Phansalkar, R. *et al.* Coronary blood vessels from distinct origins converge to equivalent states during mouse and human development. *Elife* **10**, 1–32 (2021).

33.   Kalucka, J. *et al.* Single-Cell Transcriptome Atlas of Murine Endothelial Cells. *Cell* **180**, 764-779.e20 (2020).

34.   Zhang, H. *et al.* Endocardium Minimally Contributes to Coronary Endothelium in the Embryonic Ventricular Free Walls. *Circ. Res.* **118**, 1880–1893 (2016).

35.   Conway, S. & Molkentin, J. Periostin as a Heterofunctional Regulator of Cardiac Development and Disease. *Curr. Genomics* **9**, 548–555 (2008).

36.   Torregrosa-Carrión, R. *et al.* Adhesion G protein-coupled receptor Gpr126/Adgrg6 is essential for placental development. *Sci. Adv.* **7**, (2021).

37.   Yucel, N. *et al.* Cardiac endothelial cells maintain open chromatin and expression of cardiomyocyte myofibrillar genes. *Elife* **9**, (2020).

38.   Su, T. *et al.* Single-cell analysis of early progenitor cells that build coronary arteries. *Nature* **559**, 356–362 (2018).

39.   Fang, J. S. *et al.* Shear-induced Notch-Cx37-p27 axis arrests endothelial cell

cycle to enable arterial specification. *Nat. Commun.* **8**, (2017).

40. Luo, W. *et al.* Arterialization requires the timely suppression of cell growth. *Nature* **589**, 437–441 (2021).

41. Yao, Y. & Wang, C. Dedifferentiation: inspiration for devising engineering strategies for regenerative medicine. *npj Regen. Med.* **5**, (2020).

42. Sharma, B., Chang, A. & Red-Horse, K. Coronary Artery Development: Progenitor Cells and Differentiation Pathways. *Annu. Rev. Physiol.* **79**, 1–19 (2017).

43. Chen, H. I. *et al.* The sinus venosus contributes to coronary vasculature through VEGFC-stimulated angiogenesis. *Development* **141**, 4500–4512 (2014).

44. Gillich, A. *et al.* Capillary cell-type specialization in the alveolus. *Nature* **586**, 785–789 (2020).

45. Helker, C. S. M. *et al.* Apelin signaling drives vascular endothelial cells towards a pro-angiogenic state. *Elife* **9**, 1–44 (2020).

46. Tian, X. *et al.* Peritruncal coronary endothelial cells contribute to proximal coronary artery stems and their aortic orifices in the mouse heart. *PLoS One* **8**, (2013).

47. Eyries, M. *et al.* Hypoxia-induced apelin expression regulates endothelial cell proliferation and regenerative angiogenesis. *Circ. Res.* **103**, 432–440 (2008).

48. Masoud, A. G. *et al.* Apelin directs endothelial cell differentiation and vascular repair following immune-mediated injury. *J. Clin. Invest.* **130**, 94–107 (2020).

49. Godoy, R. S. *et al.* Novel Apelin-expressing gCap Endothelial Stem-like Cells Orchestrate Lung Microvascular Repair. *bioRxiv* 2021.07.12.452061 (2022) doi:10.1101/2021.07.12.452061.

50. Mirza, M. K. *et al.* FoxM1 regulates re-annealing of endothelial adherens junctions through transcriptional control of β-catenin expression. *J. Exp. Med.* **207**, 1675–1685 (2010).

51. Zhao, Y. Y. *et al.* Endothelial cell-restricted disruption of FoxM1 impairs endothelial repair following LPS-induced vascular injury. *J. Clin. Invest.* **116**, 2333–2343 (2006).

52. Siemerink, M. J. *et al.* CD34 marks angiogenic tip cells in human vascular endothelial cell cultures. *Angiogenesis* **15**, 151–163 (2012).

53. Gulati, G. S. *et al.* Single-cell transcriptional diversity is a hallmark of developmental potential. *bioRxiv* **411**, 405–411 (2019).

54. Zhang, H. & Faber, J. E. De-novo collateral formation following acute myocardial infarction: Dependence on CCR2+bone marrow cells. *J. Mol. Cell. Cardiol.* **87**, 4–16 (2015).

55. Anbazhakan, S. *et al.* Blood flow modeling reveals improved collateral artery performance during mammalian heart regeneration. *bioRxiv* 2021.09.17.460699 (2021).

56. He, L. *et al.* Genetic lineage tracing discloses arteriogenesis as the main mechanism for collateral growth in the mouse heart. *Cardiovasc. Res.* **109**, 419–430 (2016).

57. Zhou, D. *et al.* Effects of IL-1β and TNF-α on the Expression of P311 in Vascular Endothelial Cells and Wound Healing in Mice. *Front. Physiol.* **11**, (2020).

58. Wu, B. *et al.* Endocardial cells form the coronary arteries by angiogenesis through myocardial-endocardial VEGF signaling. *Cell* **151**, 1083–1096 (2012).

59. Koh, B. I. *et al.* VEGFR2 signaling drives meningeal vascular regeneration

upon head injury. *Nat. Commun.* **11**, (2020).

60. Isner, J. M. *et al.* Clinical evidence of angiogenesis after arterial gene transfer of phVEGF165 in patient with ischaemic limb. *Lancet* **348**, 370–374 (1996).

61. Kikuchi, K. *et al.* Primary contribution to zebrafish heart regeneration by gata4+ cardiomyocytes. *Nature* **464**, 601–605 (2010).

62. Jopling, C. *et al.* Zebrafish heart regeneration occurs by cardiomyocyte dedifferentiation and proliferation. *Nature* **464**, 606–609 (2010).

63. Bertozzi, A., Wu, C. C., Hans, S., Brand, M. & Weidinger, G. Wnt/β-catenin signaling acts cell-autonomously to promote cardiomyocyte regeneration in the zebrafish heart. *Dev. Biol.* **481**, 226–237 (2022).

64. Zhang, Y. *et al.* Single-cell imaging and transcriptomic analyses of endogenous cardiomyocyte dedifferentiation and cycling. *Cell Discov.* **5**, (2019).

65. Lowe, V., Wisniewski, L. & Pellet-Many, C. The zebrafish cardiac endothelial cell—roles in development and regeneration. *J. Cardiovasc. Dev. Dis.* **8**, (2021).

66. Fernández-Chacón, M., García-González, I., Mühleder, S. & Benedito, R. Role of Notch in endothelial biology. *Angiogenesis* **24**, 237–250 (2021).

67. Phansalkar, R. *et al.* Endocardium-to-coronary artery differentiation during heart development and regeneration involves sequential roles of Bmp2 and Cxcl12/Cxcr4. (2021).

68. Ivins, S. *et al.* The CXCL12/CXCR4 Axis Plays a Critical Role in Coronary Artery Development. *Dev. Cell* **33**, 455–468 (2015).

69. Cavallero, S. *et al.* CXCL12 Signaling is Essential for Maturation of the Ventricular Coronary Endothelial Plexus and Establishment of Functional Coronary Circulation. *Dev. Cell* **33**, 469–477 (2015).

70. Chang, A. H. *et al.* DACH1 stimulates shear stress-guided endothelial cell migration and coronary artery growth through the CXCL12-CXCR4 signaling axis. *Genes Dev.* **31**, 1308–1324 (2017).

71. Evans, C. E., Iruela-Arispe, M. L. & Zhao, Y. Y. Mechanisms of Endothelial Regeneration and Vascular Repair and Their Application to Regenerative Medicine. *Am. J. Pathol.* **191**, 52–65 (2021).

72. Red-Horse, K., Ueno, H., Weissman, I. L. & Krasnow, M. A. Coronary arteries form by developmental reprogramming of venous cells. *Nature* **464**, 549–553 (2010).

73. Rhee, S. *et al.* Endothelial deletion of Ino80 disrupts coronary angiogenesis and causes congenital heart disease. *Nat. Commun.* **9**, (2018).

74. Lange, M. *et al.* Zebrafish mutants in vegfab can affect endothelial cell proliferation without altering ERK phosphorylation and are phenocopied by loss of PI3K signaling. *Dev. Biol.* **486**, 26–43 (2022).

75. El-Sammak, H. *et al.* A Vegfc-Emilin2a-Cxcl8a Signaling Axis Required for Zebrafish Cardiac Regeneration. *Circ. Res.* 1014–1029 (2022) doi:10.1161/circresaha.121.319929.

76. Marín-Juez, R. *et al.* Coronary Revascularization During Heart Regeneration Is Regulated by Epicardial and Endocardial Cues and Forms a Scaffold for Cardiomyocyte Repopulation. *Dev. Cell* **51**, 503-515.e4 (2019).

77. Giordano, F. J. *et al.* A cardiac myocyte vascular endothelial growth factor paracrine pathway is required to maintain cardiac function. *Proc. Natl. Acad. Sci. U. S. A.* **98**, 5780–5785 (2001).

78. Karra, R. *et al.* Vegfaa instructs cardiac muscle hyperplasia in adult zebrafish. *Proc. Natl. Acad. Sci. U. S. A.* **115**, 8805–8810 (2018).

79. Wang, Y. *et al.* Polymorphisms of KDR Gene Are Associated With Coronary Heart Disease. *J. Am. Coll. Cardiol.* **50**, 760–767 (2007).

80. Lee, K. W., Lip, G. Y. H. & Blann, A. D. Plasma angiopoietin-1, angiopoietin-2, angiopoietin receptor Tie-2, and vascular endothelial growth factor levels in acute coronary syndromes. *Circulation* **110**, 2355–2360 (2004).

81. Wada, H. *et al.* VEGF-C and mortality in patients with suspected or known coronary artery disease. *J. Am. Heart Assoc.* **7**, 1–10 (2018).

82. Kucukardali, Y. *et al.* The relationship between severity of coronary artery disease and plasma level of vascular endothelial growth factor. *Cardiovasc. Revascularization Med.* **9**, 66–70 (2008).

83. Ohno, T. *et al.* Serum vascular endothelial growth factor: A new predictive indicator for the occurrence of coronary artery lesions in Kawasaki disease. *Eur. J. Pediatr.* **159**, 424–429 (2000).

84. Kariyazono, H. *et al.* Association of vascular endothelial growth factor (VEGF) and VEGF receptor gene polymorphisms with coronary artery lesions of Kawasaki disease. *Pediatr. Res.* **56**, 953–959 (2004).

85. Wang, Y. *et al.* Vascular endothelial growth factor A polymorphisms are associated with increased risk of coronary heart disease: A meta-analysis. *Oncotarget* **8**, 30539–30551 (2017).

86. Palmer, B. R. *et al.* Vascular endothelial growth factor-A promoter polymorphisms, circulating VEGF-A and survival in acute coronary syndromes. *PLoS One* **16**, 1–15 (2021).

87. Rubanyi, G. M. Mechanistic, technical, and clinical perspectives in therapeutic stimulation of coronary collateral development by angiogenic growth factors. *Mol. Ther.* **21**, 725–738 (2013).

88. Kukuła, K. *et al.* Intramyocardial plasmid-encoding human vascular endothelial growth factor A165/basic fibroblast growth factor therapy using percutaneous transcatheter approach in patients with refractory coronary artery disease (VIF-CAD). *Am. Heart J.* **161**, 581–589 (2011).

89. Cao, J. *et al.* The single-cell transcriptional landscape of mammalian organogenesis. *Nature* **566**, 496–502 (2019).

90. Melsted, P. *et al.* Modular, efficient and constant-memory single-cell RNA-seq preprocessing. *Nat. Biotechnol.* **39**, 813–818 (2021).

91. La Manno, G. *et al.* RNA velocity of single cells. *Nature* **560**, 494–498 (2018).

92. Eden, E., Navon, R., Steinfeld, I., Lipson, D. & Yakhini, Z. GOrilla: A tool for discovery and visualization of enriched GO terms in ranked gene lists. *BMC Bioinformatics* **10**, (2009).

93. Raudvere, U. *et al.* G:Profiler: A web server for functional enrichment analysis and conversions of gene lists (2019 update). *Nucleic Acids Res.* **47**, W191–W198 (2019).

94. Miquerol, L. *et al.* Endothelial plasticity drives arterial remodeling within The endocardium after myocardial infarction. *Circ. Res.* **116**, 1765–1771 (2015).

95. Mahmoud, A., Porrello, E. & Kimura, W. Surgical models for cardiac regeneration in neonatal mice. *Nat. Protoc.* **9**, 305–311 (2014).

Methods

Tissue digestion and single cell sorting

Neonatal and adult mouse hearts were dissected and atria was removed from each heart. Tissue digestion was performed only with cardiac ventricles. Adult mouse hearts were additionally perfused before harvesting hearts. ~15-20 neonatal (for sham and MI each) and ~10-12 adult (for sham and MI each) *Cx40CreER*; *Rosa26^{TdTomato}* mice were used for tissue digestion. Post-explant, ventricles of the hearts were minced using razor blade and fine forceps and transferred to a tissue digestion buffer. The tissue digestion buffer was composed of 500U/ml Collagenase IV (Worthington, Catalog# LS004186), 1.2U/ml Dispase (Worthington, Catalog# LS02100), 32U/ml DNase I (Worthington, Catalog# LS002007) in commercial grade sterile 1X DPBS with Ca^{2+} and Mg^{2+}. Each heart was transferred to ~300µl of digestion buffer. Hearts bathing in digestion buffer were incubated in 37° Celsius shaker for 40-45 minutes, with periodic resuspension using micropipettes to break down tissue lumps. Digested tissue was then equilibrated with 4 volumes of 5% FBS prepared in 1XDPBS, gently mixed with a micropipette and filtered using 40µm of sterile cell strainer. The filtered tissue digest was spun down for 5 minutes at 400g at 4° Celsius, and the cell pellet was gently dissolved in 5ml of 3% FBS in 1XDPBS. The re-suspended cell pellet was spun down for 5 minutes at 400g at 4° Celsius. Supernatant was discarded and the cell pellet was suspended in ~500-600µl of 3% FBS in 1XDPBS. These suspended single cells were stained with Ter119-Alexafluor-647 (Biolegend, Catalog# 116218, 1:100 dilution) for 30 minutes at 4° Celsius. Post-staining, cells were washed with and re-suspended in fresh 3% FBS/ 1XDPBS, followed by single cell sorting with fluorescent activated cell sorter. DAPI was added prior to starting the cell sort. Live single endothelial cells (DAPI⁻

Ter119⁻ TdTomato⁺) were collected, in that order, using BD FACS Aria II cell sorter. Single cells were counted, and submitted to Stanford Genome Sequencing Service Center for 10X single cell V3 library preparation. Sequencing was performed using HiSeq 4000.

Single Cell RNA sequencing data analyses

Processing of FASTA files

Illumina reads obtained were processed on Cell Ranger (10X Genomics). To demultiplex the BCL files and convert them to FASTQ files, the *mkfastq* function was used. The reads were aligned to mm10 (mouse genome) and TdTomato sequences with the count function.

Preprocessing on R

The output files of Cell Ranger were imported into RStudio v4.05 using Seurat v4.03. This was performed with *Read10x* feature by providing path to a folder with zip files of expression matrix, features and barcodes files. The cells were analyzed for nFeature RNA count and percent of mitochondrial RNA using the standard pre-processing mentioned on the Seurat[28]. The tutorial is available at https://satijalab.org/seurat/articles/get_started.html .

Cells in our datasets did not have any mitochondrial gene expression. Based on Q/C violin plot, all cells with >250 nFeature RNA count were analyzed. These cutoffs were determined from quality control analyses of all datasets, which include, subsets of cardiac cells filtered with expression of *Cadherin5*, *Pecam1* or TdTomato from uninjured and injured neonates and adults (Figure **S1A**, **S2A**, **S9A**, **E**). We obtained

and analyzed the following number of cells from each subset. *Cadherin5* >1: 5432 sham and 2200 MI neonatal cells, *Cadherin5*>1: 6256 sham and 1690 MI adult cells, *Pecam1*>1: 5520 sham and 2152 MI neonatal cells, *Pecam1*>1: 6228 sham and 1774 MI adult cells, TdTomato>0: 5557 sham and 497 MI neonatal cells, TdTomato>0: 3090 sham and 892 MI adult cells.

Normalization, scaling and clustering

Data was log normalized with scale factor set to 10000, followed by finding variable features with n set to 2000 and finally, scaled with *scaledata*. We focused on artery endothelial cells, by sub-setting the data using different markers. *Cadherin5* or *Pecam1* (pan endothelial markers) expression greater than 1 were analyzed. As the cells from both filters echoed the same clustering pattern, we used *Cadherin5* cells for further analyses. The pattern acquired by clustering sham and MI data separately or together (merged) echoed the same patterns. Hence, the sham and MI data were integrated using the *IntegrateData* feature from the Seurat package. Normalization, finding variable features and scaling data was repeated on the integrated data, using the same settings as above.

We performed Principal Component Analysis (PCA) using the top 30 PCs, from integrated data. We did not observe any variations in clustering patterns when 10-100 PCs were used. Clusters for the integrated data were obtained with Louvain algorithm with resolution set to 0.5 (https://satijalab.org/seurat/articles/pbmc3k_tutorial.html). The cells were projected and visualized in two-dimension using Uniform Manifold Approximation and Projection for Dimension Reduction (UMAP); parameters were n.neighbors set to 10,

dims set to 1:20, spread set to 2, and min.dist set to 0.3. Attributes *group.by* and *split.by* were used to observe sham and MI cells. This was done for optimal spread of cells and best visualization.

Cells with TdTomato expression greater than 0, were analyzed. Since the MI cell number was 10 fold lesser than that of the sham, data was analyzed by sub-setting the merged data based on its orig.ident and re-clustering the respective cell populations. The steps used with *Cadherin* >1 cells were used for all clustering steps with TdTomato >0.

Cluster identification

The communities detected were identified with Violin plots (*StackedVlnPlot* from CellChat v1.1.0 package (https://github.com/sqjin/CellChat), *DotPlots* (a feature of Seurat package to visualize gene expression), and feature plots (Seurat argument). Clusters were identified with established markers for a variety of cardiac cell specific genes. Cluster identifications were confirmed using *AddModuleScore* feature from Seurat package. A list of cell-type specific, established markers were used for this analysis on *Cadherin5*>1, *Pecam1*>1 and TdTomato>0 datasets in both neonate and adult datasets. Average expression of the listed genes for each cell type was visualized using a feature plot. Neonatal trans-EC cluster did not show unique expression of any known cardiac cell types.

Heatmaps were used to check the variation in expression of de-differentiation genes across sham and MI datasets. *DoHeatmap* feature from Seurat was used to plot the

genes of interest. Violin plots generated from Seurat package were analyzed by using the *split* attribute.

Cell cycle analysis

Updated G2M and S phase specific gene lists (from 2019) on Seurat were used. As the genes were in human gene format, *useMart* function from biomaRt v3.12 (RStudio package) was employed to obtain hsapiens_gene_ensembl and mmusculus_gene_ensembl from ensembl database. Homology mapping and gene symbol conversion of genes from 2 gene-sets was performed using *getLDS* function. Genes were converted to mouse gene format using *convertHumanGeneList*. Cell cycle analysis was performed using standard cell cycle regression (CellCycleScoring) on Seurat.

Trajectory analysis

Monocle3[89] was used for trajectory analysis to understand possible pseudo time relationship between the cell types. Seurat object split into sham and MI from the integrated data, subsetted for ECs alone, was used for the analysis. *as.cell_data_set* feature from SeuratWrappers was used to convert Seurat object to Monocle3 object. Further, the Monocle3 object was subjected to pseudo time analysis using *learn_graph* and *order_cells* functions (https://cole-trapnell-lab.github.io/monocle3/docs/starting/). Results were visualized on UMAPs with the aid of *plot_cells* function.

RNA Velocity

Spliced and unspliced counts were generated from raw Fastq files using kallisto bustools wrapper *kb* (kb-python v0:27:0, https://www.kallistobus.tools/)[90]. Mouse genomic (DNA) FASTA and GTF annotations were downloaded from Ensembl and a transcriptome index was generated by kallisto using *kb ref* command. *Kb count* command was used to generate the count matrix in the form of h5ad file. The h5ad file was converted into a seurat object using SeuratDisk (v0.0.0.9019). The preprocessing, normalization, scaling and clustering were done as described earlier. The RNA velocity maps were generated using Velocyto package (velocyto.R v6)[91].

Analysis of differentiation state

R package CytoTRACE v0.3.3[53] (https://cytotrace.stanford.edu/) was used for cell differentiation analysis. Cell counts and UMAP embeddings were obtained as data frames from sham and MI clusters separately, for the analysis. A CytoTRACE score was assigned to each cell with the CytoTRACE function, which was visualized on the UMAP embeddings with *plotCytoTRACE* feature. We compared the mean CytoTRACE scores between all EC clusters. To do so, CytoTRACE score for each cell and its cluster number was obtained. The mean CytoTRACE score was then calculated for each cluster. These mean values along with standard deviations were plotted as boxplots using *ggboxplot* feature from ggpubr v0.4.0 package.

Gene Ontology (GO) analysis

List from *FindAllMarkers* was used to recognize processes that appear on publicly available online tool GOrilla (Gene Ontology enRIchment anaLysis and visuaLizAtion tool)[92] (http://cbl-gorilla.cs.technion.ac.il/). Gene Ontology analyses was performed on MI gene list, with sham genes serving as background. Default settings with a filter

of p-value less than 10^{-3} were used. Processes with significant p-values (<0.0005), and relevant to our study, were plotted on a bar graph.

g:Profiler

Pathway analysis was performed using an online tool g:Profiler[93]. The gene list was obtained for neonatal cluster aEC1 (from merged *Cadherin5* >1 dataset), using *FindMarkers* feature, with test.use set to "MAST". The organism was set to *Mus musculus* and the pathways were obtained from databases like KEGG, reactome, WikiPathways etc.

Statistical analysis on sequenced data

Wilcoxon test was performed using *RunPresto* feature from SeuratWrappers package. The statistical test was performed on cyc-aEC and aEC1 cells from neonatal *Cadherin5*+ MI group. Additionally, the test was performed on *Cadherin5*+ cyc-aECs from neonatal and adult MI datasets. For this, the data was merged using the *merge* feature from Seurat, and retained as such, without any further modifications. Test was done to check statistical significance for expression of *Cx40*, *Cxcr4*, *Aplnr*, *Car4*, *Apln*, *Foxm1, Vegfa*, *Vegfc*, *Kdr* across different treatments (sham and MI in Figure **4**) or clusters (cyc-aEC and aEC1 in neonates, in Figure **S7A**).

Experimental model

Following mouse lines were used in this study. *Rosa26*^{TdTomato} Cre reporter line (The Jackson Laboratory, B6.Cg-Gt(ROSA)26Sortm9(CAG-TdTomato)Hze/J, Stock#

007909), *Cx40CreER*[94], *VegfR2 flox* (The Jackson Laboratory, *Kdr^tm2Sato*/J, Stock#018977).

All mice were housed and bred either in Stanford or NCBS animal facility according to the approved ethical guidelines. All Tamoxifen injections (6mg dissolved in corn oil) were administered intraperitoneally to nursing mothers for Cre activation in new born pups. Data shown in this study is compiled from age-matched males and females from several litters.

Adult LCA ligations

Adult mice were anesthetized in a 4% isoflurane chamber and then intubated with a 20G angiocatheter. General anesthesia was maintained with 1.5-4% isoflurane during mechanical ventilation at a respiratory rate of 180 breaths per minute. The mice were placed in right lateral decubitus position and the left chest was shaved and sterilely prepped. 1 mg/kg Bupivicaine and 0.5 mg/kg Buprenorphine were injected subcutaneously for perioperative analgesia. A 1 cm left thoracotomy incision was made, and the pericardium was opened. The left coronary artery was identified and permanently ligated using a 6-0 polypropylene suture. The chest was then closed in layers using 5-0 polypropylene sutures. Isoflurane was then weaned off and the mice were extubated after regaining physiological respirations. 5mg/kg Carprofen was injected subcutaneously daily for the first three days after surgery for postoperative analgesia. For adult MI experiments, 2-3 months old mice were used (both males and females). To activate Cre, a single dose of 6mg Tamoxifen intraperitoneal injections were administered to each mouse, 2 days prior to MI.

Neonatal LCA ligations

Technical details of the procedure are referred and modified from Mahmoud et al[95].

To induce hypothermic arrest, neonatal mice (P2 or P7) were incubated on ice.

Neonates were wrapped with a 3-layer, pre-cooled, gauze piece for ~3 minutes.

Anesthesia was confirmed using toe pinch. Surgery was performed in sterile

conditions from start to end, using sterile surgical tools at all times. Animals were

placed in a supine position, on a sterile surgical drape, wrapped around cold ice

pack. Animals were then prepped with betadine and 70% ethanol. Left anterolateral

thoracotomy was performed by making an incision on fourth intercostal space to

access heart, under a dissecting microscope. The left coronary artery (LCA) was

identified and ligated by passing a needle attached to 8-0 non-absorbable prolene

suture (Ethicon Ethilon, catalog# NW3322). Post-ligation, the 8-0 prolene suture was

used to close the ribs, intercostal muscles and skin separately. The neonate was

then transferred to warm plate (37° Celsius) to gain consciousness and recover

completely. When conscious, neonates were returned to their mother's cage and

closely observed over the next few days.

Whole mount immunostaining of neonatal hearts

The procedure has been described earlier[24]. Neonatal whole hearts were dissected

and immediately fixed in 4% paraformaldehyde at 4° Celsius for 1hr on a rocker.

Hearts were then subjected to three, 15 minutes washes, with 1X PBS at 4° Celsius.

Primary antibody staining was done by incubating hearts in respective dilutions of

primary antibodies, prepared in at least 5 volumes of 1X PBS containing 0.5%

Triton-X 100 (0.5% PBT). Incubations were done overnight at 4° Celsius, on a

shaker. Hearts were washed in 20 volumes of 0.5% PBT for 12 hours, with a change

in wash buffer (0.5% PBT), every 2 hours at 4° Celsius. The secondary antibody

staining was performed by incubating hearts in 1:250 dilution prepared in 0.5% PBT. Incubation was done on a shaker. Next, hearts were washed in 20 volumes of 0.5% PBT, for 4 days with a change in wash buffer, every two hours at 4° Celsius. All washes were performed with continuous but gentle shaking. Hearts were cleared in at least 2 volumes of Anti-fade mounting media (https://www.jacksonimmuno.com/technical/products/protocols/anti-fade) for 2 hours, in dark, at room temperature. Whole hearts were then stored at -20° Celsius till imaged. Post staining washes were done in 15 and 50 ml tubes while incubation in antibodies or mounting media were performed in 1.5ml microtubes.

Neonates were injected with 30µl of 10mM 5-ethynyl-2'-deoxyuridine (EdU) (prepared in 1X commercial grade PBS, catalog #10010023) for 1hour, prior to harvesting their hearts. EdU label detection was combined with post-secondary immunostaining steps as described above, and, detected using Click-iT staining kit (Invitrogen, Catalog# C10340).

Confocal imaging of neonatal hearts

Whole neonatal (immunostained) hearts were mounted on a double concave microscope slide (Sail brand, catalog# 7104) using a microscope coverslip (0.13-0.17mm, Thermo Scientific, Catalog# 48367-059) so that the ventral watershed regions spanning left and right coronary arteries can be imaged. Imaging was performed using Olympus FV3000 upright confocal microscope. Whole hearts were imaged using 4X, 10X, 20X or 60X (oil) objectives. Multiple z stacks were acquired using FV31S-SW software. Multiple images of the whole heart were processed on Fiji-ImageJ software, and stitched with Adobe Photoshop.

Antibodies

Primary antibodies: Anti-CX40 (1:500; Alpha Diagnostics Int. Inc.; Catalog# CX40-A), Anti-VEGFR2 (1:125, R&D Systems, Catalog# AF644). Secondary antibodies: Alexa fluor-conjugated antibodies (488, 555, 633) from Invitrogen were used at 1:250 dilutions.

Quantification from images

Collateral number

Number of *Cx40CreER* lineage-traced (TdTomato positive) collateral arteries were quantified from images acquired using 10X objective from whole hearts. Number of stacks (12-14) and step size (2µm) of the images used, were consistent across genotypes.

Cx40 intensity in proliferating artery ECs

Proliferating TdTomato$^+$ Cx40$^+$ cardiac artery ECs (4 days post-MI) were identified with EdU$^+$ nuclear labelling. Cytoplasmic intensity of Cx40 in these (TdTomato$^+$ Cx40$^+$ EdU$^+$) ECs were quantified using images acquired with 60X (oil) objective. Mean fluorescent intensity (MFI) of cytoplasmic Cx40 signal was measured using a selection tool available with Fiji-ImageJ software. Quantification was performed with 68 single TdTomato$^+$ Cx40$^+$ EdU$^+$ artery ECs, found in the watershed of 4 hearts, collected 4 days post-MI. Comparison between proliferative (EdU$^+$) and non-proliferative (EdU$^-$) artery ECs were made within the same field of view. An unpaired t-test was done to check statistical significance. Similar analysis was done

on 67 artery ECs within artery tips or integrated into mature arteries, as identified by Cx40 immunostaining.

Single artery EC proliferation in control and *VegfR2* knockout hearts

EdU$^+$ TdTomato$^+$ single artery ECs in the watershed regions, were quantified using images acquired with 20X objective. On an average 20-30 TdTomato$^+$ single artery ECs were observed per watershed (each field of view), which was consistent across genotypes. Quantification was performed on 6 control (36 field of views) and 5 *VegfR2* knockout (24 field of views) hearts. A total of 784 control TdTomato$^+$ artery ECs and 512 *VegfR2* knockout TdTomato$^+$ artery ECs were counted for this quantification.

Single artery EC proliferation in P11 MI hearts (MI at P7)

EdU$^+$ TdTomato$^+$ proliferating artery ECs were counted in artery tips and saECs from 3 *Cx40CreER*; *Rosa26^{TdTomato}* hearts. ~80-120µm of distal ends of arteries were identified as artery tips. 25 fields of view from P11 watershed regions, from images taken with 60X (oil) objective, were used to count a total of 113 TdTomato$^+$ residual single cells.

TdTomato fluorescence in control and *VegfR2* knockout watershed

Mean Fluorescence Intensity was measured using Fiji for *Cx40CreER* lineage-traced TdTomato$^+$ single artery ECs found in watershed area 4 days post-MI. Images of watershed regions (each field of view) lacking any large or collateral arteries were acquired with 20X objective. Watershed images only with single artery cells were

considered for this quantification. 3 hearts from each genotype were used for this quantification.

LCA width:

LCA width was measured using a line selection tool on Fiji. The analysis was done for 3 control and 2 *VegfR2* KO hearts. Width was measured for the main branch of LCA in all hearts and genotypes.

Supplemental Figures, Tables and Legends

FigureS1: Identification of neonatal *Cadherin5*+ cell clusters

(**A**) Violin plots showing quality control (analyses of nFeature counts, nCount_RNA and mitochondrial genes) performed on neonatal *Cadherin5*+ dataset. (**B**) Visualization of different identified cell clusters, in neonatal *Cadherin5*+ dataset, using UMAP. (**C**) Violin Plot showing expression of endo-MT genes in neonatal *Cadherin5*+ cells. (**D**-**G**) Feature plots showing the expression of genes specific to G2/M phase. (**H**-**K**) Feature plots showing the expression of S phase markers in neonatal *Cadherin5*+ cells. (**L**) Dot plot showing parentage of cells and average expression of genes, specific to different cell-types.

Figure S2: Artery specific gene expression in neonatal *Cadherin5*⁺ cycling aEC cluster

(**A**-**D**) Violin plots showing the expression levels of artery endothelial cell specific markers in *Cadherin5*⁺ neonatal sham cells.

Figure S3: Gene expression analyses of neonatal *Cadherin5⁺* trans-EC cluster

(**A**) Visualization of cells upon re-clustering trans-EC cluster from neonatal dataset, using UMAP. (**B**) Dot plot showing average gene expression and percentage of cells expressing various genes expressed in ECs. (**C**) Feature plot showing distribution of TdTomato expressing cells in **A**. MF, myofibrillar; cap, capillary; exp, expression.

Figure S4: Identification of neonatal *Pecam1* cell clusters

(**A**) Violin plots showing quality control (analyses of nFeature counts, nCount_RNA and mitochondrial genes) performed on neonatal *Pecam1*[+] dataset. (**B**) Visualization of clusters in neonatal *Pecam1*[+] cells, using UMAP. (**C**) DotPlot showing percentage of cells and average gene expression of cell-type specific genes.

Figure S5: Analyses of post-MI artery EC proliferation in neonatal hearts

Representative confocal images of artery segments from neonatal P6 hearts, 4 days post MI, immunostained for Cx40, lineage traced for *Cx40creER* (with TdTomato) and labelled for EdU. Images are taken from ischemic regions (watershed area below the ligation). Arrows and arrowheads show (**A**-**C**) EdU$^+$ Cx40$^+$ Tdtomato$^-$ artery ECs and (**D**, **E**) EdU$^+$ Cx40$^+$ Tdtomato$^+$ artery ECs. (**F**) Modes of post-MI artery growth in neonatal hearts. neg., negative; aEC, artery endothelial cells; cap-EC, capillary endothelial cells. Td, TdTomato. Scale bar: 50μm

A

% cells per sham cluster
% cells per MI cluster

	aEC1	aEC2	aEC3	aEC4	trans EC	cyc aEC
Apln	2% / 28%	1% / 44%	0% / 22%	1% / 40%	16% / 47%	22% / 61%
Aplnr	2% / 46%	1% / 51%	1% / 27%	1% / 15%	36% / 71%	26% / 67%
Car4	0% / 22%	0% / 41%	0% / 30%	0% / 19%	35% / 51%	13% / 75%
Cx40	52% / 15%	86% / 33%	88% / 29%	52% / 22%	19% / 0%	57% / 3%
Cxcr4	38% / 31%	32% / 17%	23% / 12%	24% / 30%	32% / 11%	43% / 8%

Expression of pro-angiogenic genes by cycling artery ECs and transient ECs

Expression of angiogenic capillary/vein genes by cycling artery ECs and transient ECs

Expression of mesenchymal genes by cycling artery ECs and transient ECs

Figure S6: Analyses of de-differentiation features in *Cadherin5*⁺ neonatal cells

(**A**) Table showing percentage of sham and MI cells expressing *Apln*, *Aplnr*, *Car4*, *Cx40* and *Cxcr4*. (**B**) DotPlot from neonatal *Cadherin5*⁺ cells showing average expression and percentage of cells expressing pro-angiogenic genes (*Cd34*, *Foxm1*, *Nrp2*, *Col15a1*, *Apln*). (**C**-**E**) Feature Plots showing expression of pro-angiogenic *Apln*, *Col15a1*, and *Nrp2* in *Cadherin5*⁺ neonatal cells, from sham and MI together. (**F**-**H**) Feature Plots showing expression of angiogenic genes, also expressed by capillary or venous ECs (*Gpihbp1*, *Fabp3*, *Pcdh17*), in *Cadherin5*⁺ neonatal cells, from sham and MI together. (**I**-**K**) Feature Plots showing expression of mesenchymal genes (*Tnnt2*, *Myl3*, *Myh6*), in *Cadherin5*⁺ neonatal cells, from sham and MI together.

A Comparison of features indicating de-differentiation across neonatal EC clusters (cyc-aEC, trans-EC, aEC1) in MI

Cellular/ Molecular features	Evidence	cyc-aEC	trans-EC	aEC1
Origin (from a differentiated cell-type)	Cells originating from pre-existing artery ECs (TdTomato+ cells)	8.4%	8.5%	20.8%
Loss of mature cell markers	*Gja5* and *Cxcr4* (av. expression)	*Cx40: 0.17* *Cxcr4: 0.16*	*Cx40: 0.002* *Cxcr4: 0.27*	*Cx40: 1* *Cxcr4:1*
Gain of (early) capillary/venous EC markers	*Aplnr* and *Car4* (av. expression)	*Aplnr: 1* *Car4: 1*	*Aplnr: 1.69* *Car4: 2.24*	*Aplnr: 0.98* *Car4: 0.88*
Gain of pro-angiogenic markers	*Apln*, *Foxm1* (av. expression)	*Apln: 1* *Foxm1: 1*	*Apln: 0.91* *Foxm1: 0.1*	*Apln: 0.18* *Foxm1: 0.04*
Cell cycle re-entry	Cell cycle sorting (% cells)	G2M: 52.16% S: 47.8%	G2M: 7.02% S: 30.1%	G2M: 0.97% S: 22.2%
Cellular fate (differentiation into a specialized cell-type)	CytoTrace	least differentiated	intermediate	most differentiated
	Monocle3	origin of predicted path	connecting cyc-aEC and aEC1	end of predicted path

Cycling aECs contribute to predicted cell trajectory upon MI

Figure S7: Comparison of de-differentiation features between neonatal

***Cadherin5+* cells in cyc-aEC, trans-EC and aEC1 clusters**

(**A**) Comparison of de-differentiation features across the clusters involved in pseudo time analysis of *Cadherin5+* neonatal MI cells (cyc-aEC, trans-EC and aEC1). (**B**, **C**) RNA velocity analysis on *Cadherin5+* (**B**) sham and (**C**) MI cells. av., average.

Figure S8: Identification of adult cardiac cell clusters

(**A**) Violin plots showing quality control (analyses of nFeature counts, nCount_RNA and mitochondrial genes) performed on adult *Cadherin5*[+] dataset. (**B**) DotPlot showing percentage of cells and their average gene expression used for identifying clusters, obtained from re-clustering *Cadherin5*[+] cap-ECs. (**C**) Visualization of re-clustered adult *Cadherin5*[+] cap-ECs on a UMAP. Clusters were identified using the DotPlot shown in **B**. (**D**) FeaturePlot showing the distribution of TdTomato expressing cells on re-clustered cap-ECs in **C**. (**E**) Violin Plots showing expression of artery specific genes in adult *Cadherin5*[+] cells in both sham and MI group, integrated. (**F**) Violin plots showing quality control (analyses of nFeature counts, nCount_RNA and mitochondrial genes) performed on adult *Pecam1*[+] dataset. (**G**) Visualization of clusters in adult *Pecam1*[+] cells, using UMAP. (**H**) DotPlot showing average gene expression and percentage of cells expressing cell type specific genes in *Pecam1*[+] dataset. (**I**) Visualization of clusters in adult TdTomato[+] cells, using UMAP. (**J**) DotPlot showing average gene expression and percentage of cells expressing cell type specific genes in TdTomato[+] dataset.

VegfR2 expression in artery ECs and capillaries

A

small artery   artery tip   large artery

No significant difference in the artery network between control and VegfR2 artery-specifc KO hearts

B

Cx40CreER; Rosa26^{TdTomato}

C

VegfR2$^{L/L}$
Cx40CreER; Rosa26^{TdTomato}

D

Main LCA width (um)

VegfR2$^{+/+}$ *VegfR2$^{L/L}$*

Cx40CreER; Rosa26^{TdTomato}

Wastershed regions from *VegfR2$^{L/L}$*; *Cx40CreER*; *Rosa26^{TdTomato}* hearts

E   Heart 1   F   Heart 2   G   Heart 3

H

*

TdTomato intensity
in WSA (AU)

VegfR2$^{+/+}$ *VegfR2$^{L/L}$*

Cx40CreER; Rosa26^{TdTomato}

Figure S9: Analyses of arterial Vegf pathway in uninjured and injured neonatal hearts

(**A**) Representative confocal images from uninjured P6 *Cx40CreER*; *Rosa26^{TdTomato}* neonatal hearts showing expression of VegfR2 in small arteries, artery tips and capillaries, but not in large arteries. (**B**-**D**) Representative whole heart images from uninjured *Cx40CreER*; *Rosa26^{TdTomato}* mice, (**B**) with or (**C**) without arterial EC *VegfR2*. Tamoxifen was injected at P0, hearts were explanted at P6. (D) Quantification of width of main left coronary artery (LCA) in uninjured control and hearts deleted for *VegfR2* from arterial ECs. Tamoxifen was injected at P0, hearts were explanted at P6. (**E**-**G**) Representative confocal images of *VegfR2* depleted watershed regions from P6 neonatal *Cx40CreER*; *Rosa26^{TdTomato}* hearts, 4 days post-MI. Tamoxifen was administered at P0/P1, MI was performed at P2/P3. (**H**) Quantification of *Cx40CreER*-lineage traced (TdTomato^+) saECs in *VegfR2* depleted watershed regions from P6 neonatal hearts (shown in **E**-**G**), 4 days post-MI. cap, capillary; KO, knockout; wsa, watershed area. Scale bars: **A**: 50µm; **B**, **C**: 500 µm; **E**-**G**: 100 µm

Figure S10: Proposed working model

(**A**) Dot plot showing average gene expression and percentage of cells expressing *Vegfa*, *Vegfc* and *VegfR2 (Kdr)* in adult *Cadherin5*[+] dataset. Blue box indicates ligand expression by aECs, red box indicate downregulation of *VegfR2* in cyc-aEC cluster.

(**B**) Proposed working model of cellular events associated with Artery Reassembly. In regenerative neonatal hearts, MI induces sequential migration (via Cxcr4), de-differentiation, and proliferation (via VegfR2) of pre-existing artery ECs, which, leads to formation of collateral arteries and efficient revascularization of ischemic heart regions. Non-regenerative hearts show minimal migration, de-differentiation or proliferation. While Cxcl12/Cxcr4 pathway drives migration, in this study we show that artery EC proliferation is regulated by Vegf pathway. Green, artery ECs; Blue, de-diff (de-differentiation); Red, prol. (proliferation); mig, migration; ? unknown molecules associated with the cellular events linked to Artery Reassembly. Findings of this study (de-differentiation and VegfR2-regulated proliferation of artery ECs) are underlined.

colspan						
Statistical significance of gene expression variation as observed using Wilcoxon test						
Statistical analysis of de-differentiation genes across neonatal *Cadherin5*[+] sham and MI cyc-aEC clusters						
	p_val	auc	avg_log2FC	pct.1	pct.2	p_val_adj
Gja5	6.80E-52	0.225561	-1.39143	0.026	0.569	1.30E-47
Cxcr4	1.71E-20	0.329778	-0.30521	0.082	0.431	3.27E-16
Aplnr	1.75E-17	0.727798	0.699507	0.673	0.257	3.36E-13
Car4	5.90E-06	0.598174	0.413379	0.312	0.125	1.13E-01
Apln	1.57E-14	0.701489	0.762624	0.608	0.222	3.02E-10
Foxm1	No significant difference					
Statistical analysis of de-differentiation genes across neonatal *Cadherin5*[+] MI cyc-aEC and trans-EC clusters						
	p_val	auc	avg_log2FC	pct.1	pct.2	p_val_adj
Gja5	No significant difference					
Cxcr4	No significant difference					
Aplnr	4.82E-12	0.377149	-0.5043	0.673	0.758	9.23E-08
Car4	1.02E-07	0.415964	-0.55117	0.312	0.436	1.95E-03
Apln	No significant difference					
Foxm1	7.04E-44	0.654272	0.367115	0.341	0.031	1.35E-39
Statistical analysis of de-differentiation genes across neonatal *Cadherin5*[+] MI trans-EC and aEC1 clusters						
	p_val	auc	avg_log2FC	pct.1	pct.2	p_val_adj
Gja5	1.64E-24	0.423338	-0.30514	0.001	0.155	3.14E-20
Cxcr4	5.20E-12	0.398321	-0.40858	0.117	0.314	9.97E-08
Aplnr	2.26E-16	0.684601	0.517376	0.758	0.367	4.33E-12
Car4	1.43E-08	0.613955	0.616043	0.436	0.222	2.74E-04
Apln	7.20E-22	0.694867	0.94641	0.488	0.106	1.38E-17
Foxm1	No significant difference					
Statistical analysis of de-differentiation genes across neonatal *Cadherin5*[+] MI cyc-aEC and aEC1 clusters						
	p_val	auc	avg_log2FC	pct.1	pct.2	p_val_adj
Gja5	No significant difference					
Cxcr4	1.00E-14	0.378972	-0.48813	0.082	0.314	1.92E-10
Aplnr	No significant difference					
Car4	No significant difference					
Apln	5.76E-30	0.753943	1.028363	0.608	0.106	1.10E-25
Foxm1	2.78E-19	0.663357	0.399056	0.341	0.014	5.33E-15

Statistical analysis of de-differentiation genes across adult *Cadherin5*[+] sham and MI cyc-aEC clusters						
	p_val	auc	avg_log2FC	pct.1	pct.2	p_val_adj
Gja5	No significant difference					
Cxcr4	0.037701	0.373907	-0.57171	0.327	0.5	1
Aplnr	0.015657	0.612245	0.824505	0.265	0.036	1
Car4	No significant difference					
Apln	0.091189	0.565233	0.623217	0.163	0.036	1
Foxm1	No significant difference					
Statistical analysis of *Vegf* pathway genes across neonatal *Cadherin5*[+] sham and MI cyc-aEC clusters						
	p_val	auc	avg_log2FC	pct.1	pct.2	p_val_adj
Kdr	3.49E-22	0.770358	0.799244	0.933	0.743	6.69E-18
Vegfa	1.53E-21	0.299646	-0.49758	0.144	0.528	2.93E-17
Vegfc	1.81E-51	0.176933	-1.4309	0.099	0.708	3.48E-47
Statistical analysis of Vegf pathway genes across adult *Cadherin5*[+] sham and MI cyc-aEC clusters						
	p_val	auc	avg_log2FC	pct.1	pct.2	p_val_adj
Kdr	1.45E-15	0.125061	-1.96031	0.857	0.97	2.67E-11
Vegfa	3.01E-03	0.586006	0.375123	0.286	0.107	1.00E+00
Vegfc	1.29E-02	0.586856	0.265401	0.388	0.19	1.00E+00

Table S1: Summary of all p-values obtained from Wilcoxon sum rank test