Combined SPT and FCS methods reveal a mechanism of RNAP II oversampling in cell nuclei

Marie Fournier[1,6,§], Pierre Leclerc[1,6,§], Aymeric Leray[2,6], Dorian Champelovier[1,6], Florence Agbazahou[1,6], Gabriel Bidaux[3,6], Alessandro Furlan[1,4,5,6,*], Laurent Héliot[1,6] *

1. University of Lille, CNRS, UMR 8523, PhLAM Laboratoire de Physique des Lasers, Atomes et Molécules, Lille, France
2. Laboratoire Interdisciplinaire Carnot de Bourgogne, UMR 6303 CNRS, Université de Bourgogne Franche Comte, Dijon, France
3. INSERM UMR 1060, CarMeN laboratory, Univ Lyon1, IHU OPERA, Hôpital Louis Pradel, Hospices Civils de Lyon, Lyon, France
4. Univ. Lille, CNRS, Inserm, CHU Lille, UMR9020-U1277-CANTHER—Cancer Heterogeneity Plasticity and Resistance to Therapies, F-59000 Lille, France
5. Unité Tumorigenèse et Résistance aux Traitements, Centre Oscar Lambret, F-59000 Lille, France
6. Groupement de Recherche ImaBio, CNRS Villeneuve d'Ascq, F−59655 France

§: These authors contributed equally to this work

*: Co-senior authors; for correspondence: alessandro.furlan@univ-lille.fr ; laurent.heliot@univ-lille.fr

Keywords: transcription, RNAP II, RPB1, biophotonics, molecular dynamics, diffusion, single-particle tracking

ABSTRACT

ABSTRACT 200 mots

Gene expression orchestration is a key question in fundamental and applied research. Different models for transcription regulation were proposed, yet the dynamic regulation of RNA polymerase II (RNAP II) activity remains a matter of debate. To improve our knowledge of this topic, we investigated RNAP II motility in eukaryotic cells by combining Single Particle Tracking (SPT) and Fluorescence Correlation Spectroscopy (FCS) techniques, to take advantage of their different sensitivities in order to analyze together slow and fast molecular movements. Thanks to calibrated samples, we developed a benchmark for quantitative analysis of molecular dynamics, to eliminate the main potential instrumental biases. We applied this workflow to study the diffusion of RPB1, the catalytic subunit of RNAP II. By a cross-analysis of FCS and SPT, we could highlight different RPB1 motility states and identified a stationary state, a slow diffusion state, and two different modes of subdiffusion. Interestingly, our analysis also unveiled the oversampling by RPB1 of nuclear subdomains. Based on these data, we propose a novel model of spatio-temporal transcription regulation. Altogether, our results highlight the importance of combining microscopy approaches at different time scales to get a full insight into the real complexity of molecular diffusion kinetics in cells.

INTRODUCTION

Biological activities are classically modulated through the dynamic binding and unbinding of molecular partners [1]. Depending on the process considered, molecules may exist in several distinct complexes, which differ in their motility and availability features. In the frame of gene expression modulation, diffusion and target search constitute critical features for transcription actors [2,3]. It was acknowledged for a long time that RNA Pol II was recruited on the target genes by general transcription factors during pre-initiation complex (PIC) formation [4–6]. Indeed, RNAP II was initially observed in mammalian cells in cluster-like structures by immunostaining of the polymerase active form in fixed cells and was co-localized with nascent RNAs [7,8]. This led to the interpretation of static transcription factories. Recent advancements in the field however suggest that RNA Pol II assembly is very dynamic throughout time and that several RNA pol II can be co-recruited and form transient clusters which disintegrate after a few seconds, depending on the transcriptional activity [9–11]. How these clusters match with the concepts of transcription factories or TAD (Topologically Associated Domains)[12], and how they organize over time remain elusive. In light of these results, it is interesting to revisit the investigation of RNAPII mobility by taking advantage of the progress in molecular labelling as well as in camera speed and detector sensitivity.

Various microscopy approaches have been developed in the last decades to quantify and/or track molecular motilities. Fluorescence microscopy has notably been widely used to decipher the dynamics of molecular orchestration and is a key element to push forward the understanding of spatiotemporal molecular mechanisms. The mobility of a given molecule in cells classically encompasses several diffusion modes [13,14], and retrieving quantitative data about the diffusion of both fast and slow molecules is critical to understanding the dynamics of molecular events, including transcription, in the nucleus. To this end, we took advantage of the complementarity of FCS (Fluorescence Correlation Spectroscopy) high temporal resolution and SPT (Single-Particle Tracking) accuracy in spatial measurements of slower displacements, to perform a more complete analysis of RNAP II dynamics.

The calculation of molecular scattering using FCS is based on the autocorrelation of the measurement of the fluorescence fluctuations due to the transit of fluorescent molecules through the excitation focal volume of a laser beam [15]. Although the focal volume is small (about 0.1 μm^3), it corresponds at the nucleus scale to a subdomain that can encompass several distinct molecular crowding conditions or molecular interactions, and each experimental point thus already constitutes an average of several behaviors [16]. Moreover, such fluctuation-based techniques are blind to immobile molecules [17,18], which constitutes a drawback that has to be taken into account. FCS is rather suitable for processes occurring faster than 1 $\mu m^2/s$ [21].

In parallel, super-resolution SPT techniques have been developed thanks to the improvement of fluorophores and camera sensitivity over the last decade, and allow tracking of individual molecules in the

field of view over time [19,20]. Their temporal sampling rate (classically 10 ms/frame for a 10 µm² area) allows

the measurement of molecular motilities with diffusion coefficients up to 5 µm²/s [14,21]. In addition, SPT

techniques require the analysis of a large number of individual traces (10^3-10^5 trajectories) that may not be

equally represented as a function of their length, depending on the mobility of the molecules and the

experimental set-up. These biases may result in an over-representation of slower molecules with longer

trajectories and in less accurate statistics concerning faster molecules, which is often underestimated in

SPT analyses [3].

In this study, we first evaluated the results obtained by FCS and SPT methods with reference

samples, to establish a benchmark for the use of these methods to unravel RPB1 behavior in the nucleus.

This allowed us to overcome the potential biases that could derive from the different timescales and different

analytical protocols of the two techniques. We reasoned that it was important to deal with experimental data

rather than simulations because the former describe at best the reality of experimental images and

instrumental constraints and contributions. We came up with a robust process displaying optimized

conditions, which we applied to the analysis of RPB1 dynamics. Thanks to it, we show that RNAP II

dynamics in living cells can be described by at least four different mobility modes with a significant proportion

of anomalous diffusion probably corresponding to enhanced local target search by RNAP II.

RESULTS

Before applying FCS and SPT techniques to investigate RNAP II dynamics, we first characterized
their sensitivity in our set-ups, thanks to a system based on the diffusion of fluorescent beads in different
media of controlled viscosity. To recapitulate the apparent diffusion values observed in the literature for
proteins of the transcription complex in the nucleus, i.e. between 0.1 and 15 µm²/s [3,22], we selected
appropriate parameters to perform FCS and SPT experiments (see Material and Methods for more details).

Characterization of calibrated bead diffusion by ACF analysis of FCS measurements

The FCS experiments were conducted with fluorescent microspheres in a water/glycerol mix, with
a proportion of glycerol ranging from 0 to 80 %. These combinations provided us with control solutions
displaying theoretical diffusion coefficients ranging from 1 µm²/s to 15 µm²/s (Table S1).

We chose to fit the autocorrelation curves resulting from these experiments with an anomalous
model, which is classically used when studying the molecular motility in living cells [23–25], and allowed us to
maximize the dispersion range of the results and to explore at the same time the goodness of this model
that we would use later to investigate RNAP II diffusion in cells. We globally obtained a good correlation
between the theoretical values and our FCS experimental values (Fig.1 and Table S1).

We noticed a slight overestimation of our computed results for the diffusion coefficients when
compared to the theoretical values, especially for the diffusion of beads in pure water (Fig.1A). This could
stem from a minimal error in the estimated value of viscosity inferred, as previously reported by [26] and
described in Material and methods. Altogether, we could assess diffusion coefficients ranging from 1 to 15
µm²/s in our FCS set-up.

Besides, up to a glycerol content of 50 % in water, the ACF analysis yielded an anomalous
coefficient value very close to the theoretical value of 1 (between 0.96 and 1.02), as expected for Brownian
diffusion (Fig.1B). Only the diffusion of beads in the solution containing 80 % glycerol was described as
anomalous, with an alpha of 0.70 +/- 0.02. However, it is noteworthy that this mixture might be non-perfectly
homogenous due to the high viscosity of glycerol, and that may result in anomalous diffusion, as previously
described in the literature [27,28].

Characterization of calibrated bead diffusion by SPT analysis

To compare the measurements by FCS and SPT, we applied a similar strategy to analyze the
diffusion of reference beads with SPT.

We performed SPT acquisitions in HILO (Highly Inclined and Laminated Optical sheet) microscopy
mode [29], a technique classically used for nucleus imaging because it allows a selection of the plan of
observation thanks to a very inclined optical sheet. We analyzed image series with the MTT (Multiple-Target
Tracing) algorithm tracking software [30] to detect and reconstruct trajectories of the particles. Since short
trajectories strongly penalize the stability of the adjustment function methods [31], we only selected trajectories
with more than 10 detections, which are more reliable. With this cut-off, we obtained a mean number of

detections per trajectory superior or equal to 20 for all conditions (Fig.S1). In concordance with FCS
analysis, the trajectories were analyzed with an anomalous model. We obtained the diffusion parameters
from the linear regression of the logarithmic MSD (Mean Square Displacement) of all trajectories as a
function of time, alpha being the slope of the line and D_α the value at the origin. We found a computed
anomalous coefficient (α) very close to 1, which is consistent with the fact that beads in solution should
present a Brownian movement (Fig.2A). As observed in FCS, a high content of glycerol (80%) resulted in
an alpha computed at 0.8, with this apparent anomaly probably associated with a certain heterogeneity of
the mixture.

At the same time, we computed the diffusion coefficients D_α from the linear regression of MSD, as
explained in the previous paragraph. Besides, since it is assumed that, at short time intervals, molecules
behave like Brownian particles, we also calculated the diffusion coefficient D_{60ms} for each trajectory by
considering the slope of the vector from the origin to its value at 60 ms, as previously performed in the
community [32].

Both methods yielded diffusion coefficients slightly smaller than the theoretical values (Fig.2B), as
previously described and discussed [33]. Interestingly, the standard deviation associated with D_{60ms} was two
times lower when compared to that of D_α, so we chose the D_{60ms} method for our subsequent SPT analyses
of RPB1 diffusion.

We noticed that, for theoretical values of diffusion coefficients above 4 µm²/s, the values of D_{60ms}
and D_α were lower than the expected theoretical values, suggesting that we had reached a plateau (Fig.2B).
Thus, this calibration step shows that SPT allows to study molecular diffusion below 4 µm²/s. This agrees
with the classical measurement ranges observed in the literature with this method, due to limitations in
camera frame rates, tracking algorithms and fluorophore brightness.

Consistently, the representation of measurements as a scatter plot (Fig.2C) nicely illustrated the
fact that the populations that we detected at different glycerol concentrations could be easily distinguished,
up to this plateau where they overlapped. The distribution of bead diffusion measurements still evidenced
that this method was well suited to detect particles diffusing with coefficients over several orders of
magnitude.

To sum up this characterization step, in our experimental conditions, FCS was able to detect
relatively high (in the biological field) diffusion coefficients, ranging from 1 to 15 µm²/s, while SPT allowed
us to detect slower diffusions, ranging from 10^{-2} to 4 µm²/s, thus confirming the complementarity of these
methods.

Analysis of SPT data including mixed populations

In order to evaluate how our approach could discriminate mixed subpopulations in a sample, as is
the case in cells, we constituted mixtures of bead trajectories within various glycerol contents, from the

previous data. We chose to use the histogram analysis of MSD (called h-MSD), which allows to get a

comprehensive overview of diffusion features in complex samples with a histogram representation based

on the MSD analysis of each trajectory [32].

157   Under the assumption that D_{60ms} and (α) distributions follow sums of power laws, one can

discriminate different subpopulations emerging from a heterogeneous sample, by using a Gaussian Mixture

Model (GMM) to fit the data of log(D) and log(α) distributions. Using a methodology derived from

unsupervised machine learning, we used the BIC (Bayesian Information Criteria) and AIC (Akaike

Information Criteria) to infer the proper *k* number of populations (ranging from 1 to 10) by analyzing the point

of inflection of the BIC and AIC curves as a function of *k*. Then we computed GMM with this *k* to fit our

distributions.

164   We tested this methodology by mixing batches of data originating from distinct bead acquisitions

with various glycerol contents and assessed its ability to discriminate subpopulations and retrieve both their

right proportions and accurate coefficient values. Consistently, the BIC and AIC analyses indicated that two-

population models best described our different mixtures (Fig.3A and B for a mixture of trajectories from the

0 % and 50 % glycerol conditions). We showed that the h-MSD/GMM method allowed for discriminating

distinct subpopulations, with good quality in retrieving both the proper values and proportions for each

subpopulation (Fig.S2 and Table S2). Logically, in these settings in which particles correspond to Brownian

particles, the discrimination happened in the diffusion coefficient dimension while alpha logically remained

centered around 1 (Fig.3C). This method was pushed towards its limits only when mixing subpopulations

that were very close to the upper limit of detection, namely when merging populations with diffusion

coefficients centered around 3.8 and 4.0 μm^2/s, that were considered as a single population (Table S2D).

FCS detects RPB1 anomalous diffusion in living cell nuclei.

176   Once this benchmarking of molecular dynamics analyses was validated, we applied our methods to

177   dissect the orchestration of transcription in eukaryotic cells. For that purpose, we engineered U2OS cell

178   lines to replace endogenous RPB1, the catalytic subunit of RNAPII, with an SYFP2 fluorescently-labeled

179   amanitin-resistant RPB1 version (Fig.4A), following the strategy previously used by Darzacq and

180   collaborators [9,34].

181   This allowed us to probe RPB1 diffusion by FCS. Anomalous coefficients were mostly distributed

between 0.5 and 0.8, with a mean value of 0.61, indicating a restricted diffusion of RPB1 in nuclei (Fig.4B).

Besides, we chose to compute the transit-times in the field of view (Tt, in ms/μm^3) as a common metric to

further compare the results between FCS and SPT (see material and methods for more details). Indeed, as

described in the literature, it is not possible to directly compare the diffusion coefficients associated with

different anomalous coefficients.

Most RPB1 transit times in FCS ranged from 20 to 150 ms/µm^3, with a mean value of 62 ms/µm^3 (D=5.7
µm²/s$^\alpha$). G(0) analysis determined that the number of RPB1 per focal volume is 2.6 +/-1.4 molecules.These
transit times, as well as anomalous coefficients, were spread out as evidenced in the scatter plot, which is
representative of a variety of different behaviors that could be detected by FCS. In addition, the distribution
of these diffusion values was imperfectly fitted by a Gaussian model, suggesting the existence of
subpopulations and/or the presence of some outliers.

H-MSD SPT highlights the existence of distinct RPB1 subpopulations.

We then performed SPT experiments to complete our analysis of RPB1 diffusion. SPT requires a
dedicated labeling strategy to get only a limited number of molecules that can be tracked over time in the
field of view. These molecules should be bright enough to be detected in a single particle tracking regime
and possess a high photostability to be followed over a sufficient period (at least a few seconds, and ideally
several tens of seconds). For that purpose, we took advantage of the novel Janelia fluorophores coupled to
the Halo ligand JF549 [35] and engineered a stable cell line expressing RPB1 fused to the Halo tag.

In that context, we obtained staining levels adequate to identify single RPB1 molecules (Fig.5A), on
which we were able to perform measurements of more than one thousand RPB1 trajectories (Movie 1 as
an example) that were processed as aforementioned. We observed a great variety of molecular behaviors,
as illustrated by the different kinds of tracks reported (Fig.5B), which shows the existence of various and
complex behavior of RNAPII in the nucleus.

Following this workflow, we generated a scatter plot of anomalous coefficients and transit times
corresponding to RNAP II (RPB1-Halo/JF549) trajectories (Fig.5C). The BIC and AIC analyses of the whole
data suggested the presence of four distinct subpopulations (Fig.S3), but we limited our description to two
main populations highlighted in the plot (Fig.5C). A majority one, probably encompassing several
subpopulations as suggested by the distorted shape of the Probability Density Function, represented 75%
of the trajectories, and was centered around a transit time Tt=777 s/µm^3 (D=0.02 µm^2/s) and an anomaly
coefficient α=0.31. A second one, corresponding to 25 % of the trajectories, displayed a mean Tt=4.2 s/µm^3
(D=0.5 µm^2/s) and α=0.95. In addition, around 20 % of RNAP II observed displayed a $D_{60ms} < 5 \cdot 10^{-3}$ ms/µm²
and were considered as immobile (Movie 2).

When comparing with FCS measurements (Fig.4B), one can observe that SPT allowed us to detect
distinct diffusion behaviors since the smallest transit times detected in SPT exceeded the highest values
measured by FCS (see also Fig.S4).

Spatio-temporal localization of RPB1 subpopulations.

Another intriguing observation that we made resides in the fact that no static regions with increased
RPB1 concentrations could be observed within our super-resolution experiments. To get enough detections
to infer the spatial information, we chose to stain the cells with a JF549 concentration higher than that used

for multiple particle tracking and trajectory analysis. This level of signal allowed us to image the whole

nucleus and investigate the global spatio-temporal distribution of RPB1. The analysis of detections along

the acquisition indicated that spots appeared and disappeared over time, demonstrating the highly dynamic

nature of these elements (Movies 3, 4 and 5).

When checking the mean intensity projection signal along the movie, we could observe that some

areas were highlighted, that could correspond to transcription areas, over a diffuse background, showing

that RPB1 explores the whole nucleus (except for nucleoli (Fig.6A). In this condition, tracking could be

performed on several particles at the same time (Fig.6B). Moreover, when summing all the detections

occurring throughout the acquisition, we noticed that some regions were more frequently visited by RPB1

molecules (Fig.6C), shedding light on a previously undescribed mode of dynamic spatio-temporal regulation

of gene transcription by RNAP II oversampling. Importantly, when averaging the detections from all frames,

we obtained a quite homogenous noisy background (data not shown), indicating that the regions highlighted

in Fig.6A and B did not result from a static phenomenon. When comparing the signals from Fig.6A and 6C

(merged in Fig.6D), one can see that regions with higher intensity mainly match, still, a great proportion of

the nucleoplasm was visited by RPB1 but did not lead to detections by SPT.

Altogether, these data point out a very dynamic orchestration of RPB1 diffusion and activity.

DISCUSSION

Quantifying the mobility of molecules in living cells is a very important issue for understanding the
regulation of physiological mechanisms. The FCS and SPT techniques are highly suitable for such studies.
However, because they strongly differ in acquisition methods, analysis, and experimental parameters, they
are not easily comparable. In addition, the photo-physical properties of fluorophores (i.e. blinking, lifetime,
etc.) affect the two techniques in different ways. For these two techniques to produce complementary
information, it is necessary to properly characterize their windows of sensitivity and identify their limitations.

FCS is known to be particularly suited for the quantification of fast diffusion events but the classical
ACF analysis often misses low-frequency events, whereas SPT allows the study of diffusion heterogeneities
on a molecule-by-molecule basis, and potentially gives access to the monitoring of rare molecular events.
One main limitation of SPT remains its low sensitivity to fast displacements because a large number of
photons must be detected to allow accurate localization of single molecules, which requires integration times
in the millisecond (ms) scale, while FCS is using sampling in the microsecond scale (μs). However, labeling
with bright fluorophores presently allows obtaining good frame rates and track particles for tens of time
points [36]. The optical settings and configurations used in FCS or SPT are also quite different, with each
instrument response functions (IRF) that must be considered to define their influence.

We, therefore, reasoned that a measurement-driven comparison between the two techniques was
instrumental. For that purpose, we first characterized the influence of the setups on the diffusion
measurements of calibrated beads. Whereas the use of simple models such as beads is widespread in FCS
characterization, few works have dealt with this concept in SPT. We selected this solution, over *in silico*
simulations of random behavior generation, because this approach displays the advantage of taking into
account instrumental functions and real experimental noise. Thus, we could compare the performances of
both techniques with relatively simple experimental settings. We show that SPT gives reliable results with
high accuracy for diffusion coefficients below 4 μm^2/s (Tt > 0.25 s/μm^3), while FCS allows measurements
from 1 to 15 μm^2/s (Tts < 0.33 s/μm^3).

To better control the measurement and analysis parameters of heterogeneous samples, we
developed a benchmark based on a mixture of tracks from bead populations with different diffusion
coefficients. Interestingly, our SPT analysis workflow, relying on the result distribution fitting by a Gaussian
Mixture Model (GMM), discriminated quite well between the two initial subpopulations and obtained values
very close to the expected ones (i.e., measured in the initial mono-populations). Moreover, the proportions
of populations given by the analysis were also close to those used to make the mixtures, with very low
variability. Taken together, we can consider that the proportions are reliable with a relative uncertainty as
low as 10 %.

We then applied this validated workflow coupling SPT and FCS to the elucidation of the RNA
polymerase II choreography in cell nuclei. Labelling of the RPB1 subunit of polymerase II (RNAP II) with
fluorescent proteins, or Halo tags and adequate ligands, allowed us to measure RPB1 molecular dynamics.
The analysis of RNAP II SPT trajectories with H-MSD showed that 20 % of RPB1 molecules appeared
immobile for more than 60ms. Among RPB1 trajectories, we discriminated at least two main populations by
SPT/H-MSD. The fastest population, representing approximately 20 % of RNAP II particles, displayed a
mean transit-time (Tt) of 4.2 s/μm^3 (D_{60ms} =0.5μm^2/s), and an anomalous coefficient very close to 1 (mean
α=0.95), which reflects fairly well a free Brownian diffusion. This value is in the same order of magnitude as
those found for the free diffusion of transcription factors [23,37], but it is an order of magnitude lower than the
measurements of GFP multimer diffusion in the nucleus [38]. This could indicate that free RPB1 and
transcription factors have a slowed down Brownian diffusion in the nucleoplasm. The second mobile
population that we identified by SPT/h-MSD had a mean transit-time (Tt) of 777 s/μm^3 (D_{60ms} =0.02 μm^2/s),
and a mean anomalous coefficient α=0.31, and represented 60 % of particles. This RNAP II subpopulation
thus displayed a highly constrained subdiffusion and a very slow motion, even if it was distinct from bound
immobile RNAP II.

We also established that most RPB1 diffusion measurements by FCS exceeded the speeds caught
by SPT in our benchmark, even though such molecules probably contributed to the background noise of
SPT acquisitions. RPB1 mean diffusion measured by FCS was characterized by Tt=62.10^{-3} s/μm3 (D_α=5.7
μm^2/s$^\alpha$) and α= 0.61. These values are very close to those that we previously observed for CYCT1 [39,40],
which suggests that these RPB1 mobility measurements may correspond to RNAP II molecules involved in
the transcription pause release process. FCS, which is more sensitive to fast diffusions, hardly detected the
subpopulations analyzed in SPT, with most molecules displaying transit-times 2 to 4 orders of magnitude
higher than those identified by FCS, thus highlighting the complementarity of the two approaches.

The FCS data histogram distribution was imperfectly fit by Gaussian functions and displayed an
array of relatively fast-diffusing molecules with a continuous range of diffusion parameters. We assume that
within this range, there are no strict categories of RPB1 behaviors but rather transitions between states of
interactions progressively evolving. Such behaviors may refer to the recent concepts that multivalent and
dynamic interactions can drive phase separation [41] and that the function of large complexes could be
maintained while molecules are constantly exchanged [42].

Interestingly, another spatiotemporal clustering analysis described foci predominantly occupied by
a single RNAP II at the same time point [43], in agreement with data from immunogold labelling in live cells,
concluding on the absence of significant RNAP II clustering [45]. This was also recently confirmed with an
investigation of RNA nanodomains that evidenced small RNAP II complexes mainly displaying only one
RNAP II molecule [46]. The absence of regions of high RNAP II concentration in our observations is thus
completely consistent with these highlights. Still, we observed in our SPT movies subdomains oversampled

by fast-moving of RNAPII. Quite remarkably, the population measured by FCS shown a motility with 3 to 6

magnitude order greater than which measured in other subpopulation (fig S4) and subdiffusion (α=0.61).To

understand the mobility of RNAP II involved in this process we can think of the dynamics of a ball in a pinball

machine, which presents very fast movements in a constrained space which increases the probability of

encountering its targets. When these oversampling movements are observed with a sufficient integration

time (30s-1 min) we obtain the image of a stronger concentration of fluorescence in these bumper-like

domains. We hypothesize that this could be a mechanism for RNAP II recruitment and regulation of

transcription. It is interesting to note that CYCT1, a cofactor of the transcriptional pause release, was also

described to oversample the nuclear environment[3]. Moreover, we previously highlighted that

CYCT1displays very similar diffusion properties, in subdomains whose size is in the same order of

magnitude as that of TAD (Topologically Associated Domains) [39,47]. . Taken together, our results thus

reinforce the arguments in favor of the "RNAP II clustering" concept [9]. The proper execution of RNAP II

activity would rely on the increased probability of RPB1 to visit subdomains via an oversampling without

necessarily increasing its concentration in a specific site (pinball-dumper effect). Interestingly, the study of

Herpes Simplex Virus replication recently evidenced the possibility for RNAP II to visit in a repeated manner

nearby DNA binding sites without clear boundaries between territories, which may be a common process

[50]. In that frame, highly dynamic interactions would ensure the transcription complex activity is maintained

despite the exchange of molecules [42]. Such a link between subdiffusion, topological frameworks and

recruitment/regulation steps in transcription is emerging in the field [48] and starts to be described at the

molecular level [49].

Altogether, our combined experiments suggest an RNAP II dynamic model with at least four kinetic

subpopulations: a) chromatin-bound RNAP II, b) RNAP II slowly-diffusing in a Brownian manner, and two

major subpopulations: c) subdiffusive slowly-diffusing RNAP II and d) fast-diffusing and oversampling RNAP

II molecules. Moreover, our data also suggest the potential existence of other subpopulations. Future work

will be aimed at elucidating whether we can identify and investigate these minority subpopulations to refine

our knowledge of the transcription process.

In addition, FCS gave us access to molecular concentration in focal volume and indicated the

presence in average of 2.6 +/-1.4 RPB1 molecules per focal volume, i.e. approximately 14 molecules per

μm^3. Considering the previous estimation of 80,200 RNAP II molecules per nucleus in U2OS cells, i.e. 37

molecules per μm^3 [43], this would mean that FCS detected almost one-third of RPB1 molecules. In a first

approach, we can extrapolate that the remaining two-thirds of RNAP II molecules, 23 out of 37 molecules

per μm^3, are observed by SPT. Within this category, 20% of SPT trajectories (i.e. 13 % of total RNAP II)

were immobile (7-8 molecules/μm^3). This estimate stands in agreement with the number of chromatin-bound

RNAP II proposed by Cisse et al with another microscopy method [9]. Besides, about 40% of RNAP II

displayed a strong sub-diffusive behavior (mean anomalous coefficient α=0.31). Taken together, these data

point out a global view of a majority of RNAP II molecules that are strongly restricted, and potentially involved

in regulatory steps. In SPT, only 20% of trajectories (i.e. 13 % of total RNAP II) could be considered free,

according to their anomalous coefficient α=0.95. This is again in agreement with the literature, including

recent FRAP measurements concluding on a minority fraction of freely diffusing polymerases [44]. X

At this stage, we hypothesize that the observed mobilities are associated with the mechanism of

recruitment of RNAPII to transcription regulation. We also hypothesize that the subpopulations described,

on the basis of their mobilities, are corresponding to transient states of RNAPII molecules and not to stable

molecular states. We propose two models of RNAPII recruitment that can account for the observations and

measurements of RPB1 dynamics. Both models encompass a free population slowly diffusing in an almost

Brownian walk. The first model displays a single type of cluster and two steps corresponding to successive

stages of regulation (fig 7A): in a first step, RNAP II is "trapped" in a subdomain with a strong oversampling

of this space (pinball-bumper effect), and in a second step it presents a very constrained diffusion and a

very slow speed of displacement leading to its immobilization on chromatin. We can assume that it is then

the first step corresponding to the entry into the topologically active domains (TADs) which induces the

oversampling and the first level of sub-diffusion, while the association with the cofactors causes the second

step. We can also think of a second model where RNAP II is trapped in two types of clusters via two distinct

co-existing mechanisms (fig 7 B). In some clusters, significant oversampling could correspond to "RNAP II

trapping" in gene regions controlled by transcriptional pausing (similarity with CYCT1), whereas in other

clusters "RNAP II trapping" would induce a strong and constrained diffusion that may correspond to

transcriptional regulations without transcriptional pausing.

This study paves the way for future work that will be aimed at investigating which of these two

hypotheses correspond to the reality of RNAP II recruitment to active genes, and at identifying the molecular

mechanisms responsible for RNAP II subdiffusion. In that frame, the coupling of FCS and SPT acquisitions

on the same instrument would represent an interesting instrumental development to simultaneously acquire

molecules by both methods in a spatio-temporal manner, and we are moving forward in that direction.

METHODS

Reagents and sample preparations

Standard samples preparation

Water/Glycerol solutions were prepared considering weight percentage with the use of a precision balance.
Solutions were vortexed for 30 sec before being stored at 37°C. A few minutes before acquisitions, to avoid
agglomerations of beads, we diluted our fluorescent beads stocks in two distinct manners:

- 100 nm diameter beads (Molecular Probes, Fluorescent microspheres (505/515), 2 % solid) diluted
10^4 in solutions for SPT acquisition

- 40 nm diameter beads (Molecular Probes, Fluorescent microspheres (505/515), 2 % solid) diluted
10^2 in solutions for FCS acquisition

To compare the SPT and FCS workflows, we applied a similar strategy. However, since the experimental
focal volume in FCS was in the order of 200 nm diameter in the x-y axis, the use of 100nm-diameter beads
resulted in huge bursts of fluorescence during signal acquisition, which were misinterpreted during the
Autocorrelation Function (ACF) process. We therefore chose smaller fluorescent beads (40nm-diameter)
for FCS experiments, which allowed us to measure more reliably fluorescence fluctuations and infer their
diffusion coefficients.

Bead solutions were finally deposited in 35-mm glass-bottom dishes (Ibidi, Clinisciences, France) and kept
at 37° in a thermostatic chamber during acquisitions.

The theoretical values of beads' diffusion were calculated according to the Stokes-Einstein formula:

384
$$D = \frac{kT}{6\pi r \mu}$$

385 With k Boltzmann constant, T temperature in Kelvin, r the hydrodynamical radius of the bead and µ the
386 viscosity.

387 Viscosity values of different (w:g) solutions were calculated following Takamura et al. [26] and are indexed in
388 the following table.

% glycerol	Calculated D value (µm²/s) for 40 nm beads	Calculated D value (µm²/s) for 100 nm beads
0	15.1	5,98
10	12.1	4,78
20	9.05	3,58
50	3.24	1,25
80	0.4	0,19

Biological resources

Cell line culture

Human U2OS osteosarcoma cells (HTB-96 accession number from the American Type Culture Collection,
Manassas, VA) were tested for mycoplasma contamination before establishing the RPB1 cell lines. Cells
were cultured in DMEM medium (Gibco Laboratories, Gaithersburg, MD) supplemented with 10% FCS (V/V)
and penicillin/streptomycin (100 mg/mL). Cells were grown in a humidified incubator at 37°C / 5 % CO_2.
Cells were passaged every 2–4 days before reaching confluency.

We generated plasmids encoding for human RPB1 (N792D) α-amanitin-resistant mutated fused to an N-
terminal Halo tag, by molecular cloning.

We stably transfected U2OS cells with this plasmid using Fugene 6 following the manufacturer's instruction
(Promega). α-Amanitin (SIGMA #A2263) was used to select transfected cells, at a concentration of 2 μg/mL
and was used thereafter in permanence in the culture of the cells at a concentration of 1μg/mL to avoid
endogenous RPB1 re-expression as described in (Boehning et al. 2018). For live-cell imaging, cells were
plated on 35-mm glass-bottom dishes (Ibidi, CliniSciences, Nanterre, France), filled with L-15 medium
without Phenol Red (Life Technologies, Carlsbad, CA), and incubated at 37°C in a microscope thermostatic
chamber (OKO, Italia) for imaging.

Labeling protocol

We labeled cells just before imaging. Following HaloTag technology protocol (Promega), cells were
incubated for 30 min in DMEM 10%FCS supplemented with 0.1 nM JF 549 for SPT. Then, cells were rinsed
3 times with DMEM + 10% SVF and were re-incubated for 30 min. Finally, the cell medium was replaced by
L-15 medium (Life Technologies, CA, USA) before imaging, to avoid DMEM auto-fluorescence.

SPT set up

SPT measurements were performed with an inverted microscope (Ti-Eclipse, Nikon Instrument, Japan)
adapted with an x100 NA=1,49 TIRF water immersion objective and an appropriate dichroic
405/488/561/635 nm (ref Di03-R405/488/561/635 SEMROCK, USA). The excitation laser (488nm Argon
LASER, Melles-Griot, USA and 561nm, OXXIUS, France) was injected into the TIRF Nikon module into a
mono-mode fiber and focused on the backplane of the objective. Illumination was performed in HILO mode
[29] to improve contrast in images. Perfect Focus System (PFS, Nikon) was set to avoid drift on Z-axis
(defocusing) on the objective, relative to the coverslip. Experiments were acquired under continuous
illumination (488 nm 20 kW/cm² on the sample, or 561 nm 20 kW/cm² on the sample). Emitted fluorescence
was harvested on an sCMOS camera (PCO EDGE 4.2, PCO, Germany) with a pixel size of 6,5 μm. The
acquisition format was 100x100 px, and the acquisition rate was set at 100 Hz (10 ms per frame). All devices

were driven by NIS software (Nikon Instruments, Japan). A thermostatic chamber ('on-stage-incubator',

OKOLAB, Italy) was used to keep samples at 37 Celsius degrees.

SPT data analysis:

Tracks were reconstructed (detection and tracking) with MTT algorithm tracking software [30] running on

MATLAB (MathWorks), which is well adapted for a high concentration of particles (up to 10 for a 100x100px

field of view). For each condition, we had at least 1000 trajectories lasting more than 10 detections (smaller

trajectories were not included in the analysis). Tracks were then analyzed with custom code kindly provided

by Laura Caccianini (Coppey/Hajj lab, Institut Curie, France). H-MSD (histogram Mean Square

Displacement) technique is based on the Time-Average MSD calculation $TA - MSD(\tau) = \langle r^2(\tau) \rangle_T =$

$\langle [x(t+\tau) - x(t)]^2 + [y(t+\tau) - y(t)]^2 \rangle_T \propto 4D\tau^\alpha$ where Δt , with n the number of frames considered for the

sliding average (0<n<L_{traj}, $L_{traj}$the length of the trajectory), Δt the lag time between two frames (10 ms), D

the diffusion coefficient (µm²/s), and α is the anomaly coefficient.

With H-MSD each track was treated individually as described in [32,51]. The TA-MSD curves were fitted with

a linear fit until $\tau = 60\ ms$ to obtain the instantaneous diffusion coefficient (D_{60ms}) following the previous

works of Michalet et al. describing a procedure to analyze the individual MSD curves [31]. In accordance with

this work, the first point MSD (1Δt) is not considered in the fit because it is the same order of magnitude as

the static localization error (50 nm, calculated on fixed RPB1-Halo in U2OS cells.

Anomaly coefficient alpha (α) was calculated by fitting the log-log representation of the TA-MSD with a

power-law function TA-MSD(t)$\propto 4t^\alpha$. For individual tracks, it is common that the second half of the MSD

curve is noisy because of the low statistics for each MSD(nΔt) value. We decided to consider the first third

of the curve for the linear fit because more than the second half of the MSD curve is noisy and cannot be

interpreted. The slope of the regression line gave alpha (α). In addition, the diffusion coefficient could also

be computed from the intercept of the slope, D_α , and was compared to D_{60ms} in the results.

A Gaussian mixture model (GMM) was applied to fit the data. GMM was used as a clustering algorithm

where the number k of populations is chosen from an analysis of the Bayesian Information Criterion and the

Akaike Information Criterion. This population number was chosen as small as possible, following those

criteria. By evaluating the number of discernible populations with a statistical criterion we tried to minimize

any bias that can occur when using such a supervised machine learning algorithm and create a robust and

reproducible analysis pipeline.

Mix diffusion data models

For the two sub-populations studies, we mixed batches of trajectories coming from different conditions. The

different conditions and the different fractions for each mix are indexed in the following tables.

Combined SPT and FCS methods highlight a mechanism of RNAP II oversampling in cell nuclei

Condition A		Total number of trajectories	Nb of D1 trajectories	% D1	Nb of D2 trajectories	% D2
D1=4 µm²/s (w :g =100 :0)	m1	2013	845	42	1168	58
& D2=0.1 µm²/s	m2	3118	2339	75	780	25
(w :g=20 :80)	m3	3369	842	25	2527	75

Condition B		Total number of trajectories	Nb of D1 trajectories	% D1	Nb of D2 trajectories	% D2
D1=0.9 µm²/s (w :g =50 :50) &	m1	2027	932	46	1095	54
D2=0.1 µm²/s	m2	3207	1860	58	1347	42
(w :g=20 :80)	m3	3633	908	25	2725	75

Condition C		Total number of trajectories	Nb of D1 trajectories	% D1	Nb of D2 trajectories	% D2
D1=4.0 µm²/s (w :g =100 :0) &	m1	4470	894	20	3576	80
D2=0.9 µm²/s	m2	4791	2443	51	2348	49
(w :g=50 :50)	m3	5119	3993	78	1126	22

Condition D		Total number of trajectories	Nb of D1 trajectories	% D1	Nb of D2 trajectories	% D2
D1=4.0 µm²/s (w :g =100 :0) &	m1	5293	688	13	4605	87
D2=3.8 µm²/s	m2	5310	2443	46	2867	54
(w :g=90 :10)	m3	5398	4049	75	1350	25

FCS set up

FCS measurements were performed on a Nikon A1 confocal microscope (Nikon Instrument, Tokyo, Japan)

with a 40x objective NA=1,25 (Nikon Instrument, Tokyo, Japan). Excitation Argon LASER 488 nm (Coherent,

USA) was used to scan samples in confocal illumination. To spectrally select the excitation from the emitted

light, we used a bandpass dichroic mirror (405/488 nm). Fluorescence photons were collected on a

photodiode and images were reconstructed with NIS software (Nikon Instruments, Tokyo, Japan).

To perform FCS, the CLSM was adapted with a TCSPC module (PicoQuant, Germany), and a pulsed laser

at 488 nm (LDH-D-C 488, PicoQuant), a TSCPC unit, and a single molecule counting detector (SPCM CD

3516H, Excelitas Technologies Corp., USA). To drive this photon counting system, we used SymphoTime

software (PicoQuant, Germany) in FCS mode. A high pass filter (BLP-488R, SEMROCK, USA) was set

before the detector.

FCS analysis:

FCS setup was calibrated with ATTO 488 diluted at 10 nM in Milli-Q water to define the focal volume by

best practice. In the water at 36°, the temperature of our incubator, the Atto 488 diffusion coefficient is 536

µm²/s (source: PicoQuant Application Note Kapusta and GmbH, n.d.).

The number N of molecules in this observation volume V was inferred from the autocorrelation function with

473
$$N = \frac{1}{G(0)}$$

474   And the observation volume determined with:

475
$$V_{eff} = \frac{1}{G(0). N_A. c}$$

N_A is the Avogadro number and c is the concentration of the calibration solution.

Once calibration was done, we had a good estimate of the focal volume of the microscope and could

proceed to the analysis of fluorescence fluctuations. Measurements displaying a linear regression of the

fluorescent signal characteristic of photobleaching were excluded. For data analysis, we use a home-written

MATLAB code named EASY FCS [40] which can fit the autocorrelation function (ACF).

In the case of normal diffusion the ACFs were adjusted with a Gaussian function (assuming the beam as a

Gaussian profile) :

483
$$G(\tau) = \frac{1}{N}\left(\frac{1}{1+\left(\frac{\tau}{\tau_D}\right)}\right)\left(\frac{1}{\sqrt{1+\frac{w_0^2}{z_0^2}\left(\frac{\tau}{\tau_D}\right)}}\right)$$

484   Where *N* is the number of molecules and τ_D is the characteristic residence time.

485 For anomalous diffusion, the mean square displacement is no longer linearly dependent on time, but it

486 evolves proportionally with time to the power of α. In this case, the autocorrelation function becomes:

487
$$G(\tau) = \frac{1}{N} \left(\frac{1}{1 + \left(\frac{\tau}{\tau_D}\right)^{\alpha}} \right) \left(\frac{1}{\sqrt{1 + \frac{w_0^2}{z_0^2} \left(\frac{\tau}{\tau_D}\right)^{\alpha}}} \right)$$

(4)

The fitting of the ACFs was done with the Trust-Region-Reflective Least Squares Algorithm. To evaluate

the goodness of the fit, we minimized the sum of the squared error.

Since the diffusion coefficient (D) obtained by anomalous model fit is given with time dimensions dependent

on exponent α, it is recommended to give the results in transit time [23,24,39] to compare data.

Standardization of analysis metrics between FCS and SPT

FCS and SPT data are obtained by analytical methods from microscopic approaches with different metrics.

In order to compare them and use them together to analyze molecular mobility, it is useful to express them

with the same metric and unit. The adjustment of the results by an anomalous model in both techniques

also induced a difficulty, because when the anomalous coefficient changed, the time unit was

simultaneously modified. To overcome this difficulty, it is usual in FCS to express the ACF results as

residence times in the focal volume. Here, we propose to extend this metric to SPT data by proposing a

transit time (Tt) within a reference volume (s/µm³) which could then be a common metric to both FCS and

SPT techniques. In a first approach, we make the hypothesis that the displacements measured by SPT are

isotropic in the volume at the scale of the confocal resolution (femtoliter), to pass from a surface

measurement to the exploration of the corresponding volume. This allows us to define a normalized transit

time from each measurement.

Limité a 60 ref

REFERENCES

1. Presman, D. M. *et al.* Quantifying transcription factor binding dynamics at the single-molecule level in live cells. *Methods* **123**, 76–88 (2017).

2. Halford, S. E. & Marko, J. F. How do site-specific DNA-binding proteins find their targets? *Nucleic Acids Res* **32**, 3040–3052 (2004).

3. Izeddin, I. *et al.* Single-molecule tracking in live cells reveals distinct target-search strategies of transcription factors in the nucleus. *eLife* **3**, (2014).

4. Kim, T.-K. *et al.* Trajectory of DNA in the RNA polymerase II transcription preinitiation complex. *PNAS* **94**, 12268–12273 (1997).

5. Kornberg, R. D. The molecular basis of eukaryotic transcription. *PNAS* **104**, 12955–12961 (2007).

6. Lee, T. I. & Young, R. A. Transcription of eukaryotic protein-coding genes. *Annu. Rev. Genet.* **34**, 77–137 (2000).

7. Chakalova, L. & Fraser, P. Organization of Transcription. *Cold Spring Harb Perspect Biol* **2**, a000729 (2010).

8. Jackson, D. A., Hassan, A. B., Errington, R. J. & Cook, P. R. Visualization of focal sites of transcription within human nuclei. *EMBO J* **12**, 1059–1065 (1993).

9. Cisse, I. I. *et al.* Real-Time Dynamics of RNA Polymerase II Clustering in Live Human Cells. *Science* **341**, 664–667 (2013).

10. Cho, W.-K. *et al.* RNA Polymerase II cluster dynamics predict mRNA output in living cells. *eLife* **5**, (2016).

11. Cho, W.-K. *et al.* Super-resolution imaging of fluorescently labeled, endogenous RNA Polymerase II in living cells with CRISPR/Cas9-mediated gene editing. *Sci Rep* **6**, 35949 (2016).

12. Jabbari, K., Chakraborty, M. & Wiehe, T. DNA sequence-dependent chromatin architecture and nuclear hubs formation. *Sci Rep* **9**, 14646 (2019).

13. Hager, G. L., McNally, J. G. & Misteli, T. Transcription Dynamics. *Molecular Cell* **35**, 741–753 (2009).

14. Hansen, A. S. *et al.* Robust model-based analysis of single-particle tracking experiments with Spot-On. *eLife* https://elifesciences.org/articles/33125 (2018) doi:10.7554/eLife.33125.

15. Magde, D., Elson, E. & Webb, W. W. Thermodynamic Fluctuations in a Reacting System---Measurement by Fluorescence Correlation Spectroscopy. *Phys. Rev. Lett.* **29**, 705–708 (1972).

16. Langowski, J. Protein-protein interactions determined by fluorescence correlation spectroscopy. *Methods Cell Biol.* **85**, 471–484 (2008).

17. Altan-Bonnet, N. & Altan-Bonnet, G. Fluorescence Correlation Spectroscopy in Living Cells: A Practical Approach. *Current Protocols in Cell Biology* **45**, 4.24.1-4.24.14 (2009).

18. Elson, E. L. Fluorescence correlation spectroscopy: past, present, future. *Biophys J* **101**, 2855–2870 (2011).

19. Cheezum, M. K., Walker, W. F. & Guilford, W. H. Quantitative comparison of algorithms for tracking single fluorescent particles. *Biophys J* **81**, 2378–2388 (2001).

20. Manley, S. *et al.* High-density mapping of single-molecule trajectories with photoactivated localization microscopy. *Nat Methods* **5**, 155–157 (2008).

21. Harwardt, M.-L. I. E., Dietz, M. S., Heilemann, M. & Wohland, T. SPT and Imaging FCS Provide Complementary Information on the Dynamics of Plasma Membrane Molecules. *Biophysical Journal* **114**, 2432–2443 (2018).

22. Grünwald, D. *et al.* Probing Intranuclear Environments at the Single-Molecule Level. *Biophysical*
Journal **94**, 2847–2858 (2008).
23. Mazza, D., Abernathy, A., Golob, N., Morisaki, T. & McNally, J. G. A benchmark for chromatin binding
measurements in live cells. *Nucleic Acids Research* **40**, e119–e119 (2012).
24. Stasevich, T. J. *et al.* Cross-validating FRAP and FCS to quantify the impact of photobleaching on in vivo
binding estimates. *Biophys J* **99**, 3093–3101 (2010).
25. Wachsmuth, M., Waldeck, W. & Langowski, J. Anomalous diffusion of fluorescent probes inside living
cell nuclei investigated by spatially-resolved fluorescence correlation spectroscopy. *J. Mol. Biol.* **298**,
677–689 (2000).
26. Takamura, K., Fischer, H. & Morrow, N. R. Physical properties of aqueous glycerol solutions. *Journal of*
Petroleum Science and Engineering **98–99**, 50–60 (2012).
27. Banks, D. S., Tressler, C., Peters, R. D., Höfling, F. & Fradin, C. Characterizing anomalous diffusion in
crowded polymer solutions and gels over five decades in time with variable-lengthscale fluorescence
correlation spectroscopy. *Soft Matter* **12**, 4190–4203 (2016).
28. Banks, D. S. & Fradin, C. Anomalous Diffusion of Proteins Due to Molecular Crowding. *Biophysical*
Journal **89**, 2960–2971 (2005).
29. Tokunaga, M., Imamoto, N. & Sakata-Sogawa, K. Highly inclined thin illumination enables clear single-
molecule imaging in cells. *Nat Methods* **5**, 159–161 (2008).
30. Sergé, A., Bertaux, N., Rigneault, H. & Marguet, D. Dynamic multiple-target tracing to probe
spatiotemporal cartography of cell membranes. *Nat. Methods* **5**, 687–694 (2008).
31. Michalet, X. Mean square displacement analysis of single-particle trajectories with localization error:
Brownian motion in an isotropic medium. *Physical Review E* **82**, (2010).
32. Etoc, F. *et al.* Non-specific interactions govern cytosolic diffusion of nanosized objects in mammalian
cells. *Nature Materials* **17**, 740–746 (2018).
33. Wieser, S. & Schütz, G. J. Tracking single molecules in the live cell plasma membrane—Do's and
Don't's. *Methods* **46**, 131–140 (2008).
34. Darzacq, X. *et al.* In vivo dynamics of RNA polymerase II transcription. *Nat Struct Mol Biol* **14**, 796–806
(2007).
35. Grimm, J. B. *et al.* A general method to fine-tune fluorophores for live-cell and in vivo imaging. *Nat.*
Methods **14**, 987–994 (2017).
36. Grimm, J. B. *et al.* A general method to improve fluorophores for live-cell and single-molecule
microscopy. *Nat. Methods* **12**, 244–250, 3 p following 250 (2015).
37. Jankevics, H. *et al.* Diffusion-Time Distribution Analysis Reveals Characteristic Ligand-Dependent
Interaction Patterns of Nuclear Receptors in Living Cells. *Biochemistry* **44**, 11676–11683 (2005).
38. Bancaud, A. *et al.* Molecular crowding affects diffusion and binding of nuclear proteins in
heterochromatin and reveals the fractal organization of chromatin. *EMBO J.* **28**, 3785–3798 (2009).
39. Bidaux, G. *et al.* FRET Image Correlation Spectroscopy Reveals RNAPII-Independent P-TEFb
Recruitment on Chromatin. *Biophysical Journal* **114**, 522–533 (2018).
40. Furlan, A. *et al.* HEXIM1 Diffusion in the Nucleus Is Regulated by Its Interactions with Both 7SK and P-
TEFb. *Biophys. J.* **117**, 1615–1625 (2019).
41. Lu, H. *et al.* Phase-separation mechanism for C-terminal hyperphosphorylation of RNA polymerase II.
Nature **558**, 318–323 (2018).
42. Darzacq, X. & Tjian, R. Weak multivalent biomolecular interactions: a strength versus numbers tug of
war with implications for phase partitioning. *RNA* **28**, 48–51 (2022).

43. Zhao, Z. W. *et al.* Spatial organization of RNA polymerase II inside a mammalian cell nucleus revealed
by reflected light-sheet superresolution microscopy. *Proceedings of the National Academy of Sciences*
111, 681–686 (2014).
44. Steurer, B. *et al.* Live-cell analysis of endogenous GFP-RPB1 uncovers rapid turnover of initiating and
promoter-paused RNA Polymerase II. *Proc Natl Acad Sci USA* **115**, E4368–E4376 (2018).
45. Orlov, I. *et al.* Live cell immunogold labelling of RNA polymerase II. *Sci Rep* **5**, 8324 (2015).
46. Castells-Garcia, A. *et al.* Super resolution microscopy reveals how elongating RNA polymerase II and
nascent RNA interact with nucleosome clutches. *Nucleic Acids Research* **50**, 175–190 (2022).
47. Pombo, A. & Dillon, N. Three-dimensional genome architecture: players and mechanisms. *Nat Rev Mol*
Cell Biol **16**, 245–257 (2015).
48. Cardozo Gizzi, A. M. A Shift in Paradigms: Spatial Genomics Approaches to Reveal Single-Cell Principles
of Genome Organization. *Front Genet* **12**, 780822 (2021).
49. Lee, R. *et al.* CTCF-mediated chromatin looping provides a topological framework for the formation of
phase-separated transcriptional condensates. *Nucleic Acids Research* **50**, 207–226 (2022).
50. McSwiggen, D. T. *et al.* Evidence for DNA-mediated nuclear compartmentalization distinct from phase
separation. *Elife* **8**, (2019).
51. Normanno, D., Dahan, M. & Darzacq, X. Intra-nuclear mobility and target search mechanisms of
transcription factors: A single-molecule perspective on gene expression. *Biochimica et Biophysica Acta*
(BBA) - Gene Regulatory Mechanisms **1819**, 482–493 (2012).

ACKNOWLEDGEMENTS

We thank all members of our team and GDR Imabio-CNRS, particularly Dr Hugues Berry, Dr Ignacio Izeddin

and Dr Thorsten Wohland for their help and fruitful discussion/comments. We are much grateful to Dr Raoul

Torero-Ibad for his great work and a precious help in providing us with a functional lab environment, and for

his enlightened opinion on this study. We thank Christian Hubert (Errol company) for his valuable input in

instrumental advances. We are much grateful to Dr Jonathan Grimm and Dr Luke Lavis (Janelia Farm, VA,

USA) for kindly providing us with JF dyes to perform our experiments.

AUTHOR CONTRIBUTIONS

MF, PL, GB, AF, and LH designed research. MF, PL, DC, and FA performed experiments. MF, PL, AL, GB,

AF, and LH analyzed and interpreted the data. MF, PL, AF, and LH wrote the manuscript.

DATA AVAILABILITY STATEMENT

Data and material are available on request to laurent.heliot@univ-lille.fr

SUPPLEMENTARY DATA

Supplementary Movies are available at:

https://nextcloud.univ-lille.fr/index.php/s/aqXXqPsMj93DESq

https://nextcloud.univ-lille.fr/index.php/s/Da6m2W9oGPexfcY

https://nextcloud.univ-lille.fr/index.php/s/KjXnoYcG26AnFXL

https://nextcloud.univ-lille.fr/index.php/s/HQW5rXBSg78JHMw

https://nextcloud.univ-lille.fr/index.php/s/eFTsxyXaPnApE9C

CONFLICT OF INTEREST

The authors declare that they have no competing interests.

FUNDING

This work was supported by CNRS and Ministerial Funding; Agence Nationale de la Recherche (Dynam-ic-

12-BSV5-0018-02 and ABC4M- ANR-20-CE45-0023); the LABEX CEMPI (ANR-11-LABX-0007), I-SITE

ULNE (ANR-16-IDEX-0004) as well as by the Ministry of Higher Education and Research, Hauts de France

council and European Regional Development Fund (ERDF) through the Contrat de Projets Etat-Region

(CPER) Photonics for Society (P4S), Region «Hauts de France» I-PRIMER, CNRS and GDR-ImaBio

Interdisciplinary Master's Scholarship and Nikon partnership agreement.

Funding for open access charge: ANR-20-CE45-0023

FIGURES AND LEGENDS

Figure 1. Characterization of diffusion by Fluorescence Correlation Spectroscopy.

Fluorescent microspheres were resuspended in water/glycerol mixtures, with proportions of glycerol

ranging from 0 to 80 %. The ACF curves obtained from FCS acquisitions were fit with an anomalous model

to infer the anomalous coefficients (A) and diffusion coefficients (B). Experimental values (red squares) can

be compared to theoretical values (black dots). Experimental results are expressed as means +/- SD from

at least 20 independent measurements.

Figure 2. Characterization of diffusion by Single Particle Tracking. Fluorescent microspheres were
resuspended in water/glycerol mixtures, with proportions of glycerol ranging from 0 to 80 %. SPT
acquisitions were performed at a frame rate of 100 Hz. A. The anomalous coefficient α is provided as a
function of the glycerol proportion. Experimental values (red squares) can be compared to theoretical values
(black dots). B. The diffusion coefficients were computed by two methods: D_a (blue triangles) were obtained
from the slopes of TEA-MSD (Time Ensemble Average Mean Square Displacement) expressed as log, while
D_{60ms} were calculated for each trajectory during the first 60ms of its occurrence (red squares). These
experimental values could be compared to theoretical values (black dots). Both methods yielded an
asymptotic trend for high D values. C. The distributions of D_{60ms} and α from the different experimental
conditions are represented with log scales in a scatter plot. Distinct subpopulations can be distinguished
(green, purple, and yellow dots) within a limited range of parameters, whereas some others overlap (yellow,
red, and blue dots). Data come from at least one hundred measurements per condition.

Figure 3. Discrimination of merged subpopulations thanks to h-MSD. SPT trajectories from experiments with fluorescent microspheres diffusing in 20 % and 50 % glycerol solutions respectively, were merged and submitted to the h-MSD analysis workflow to compute for each trajectory the alpha and log(D). The number of Gaussian functions to be found in our distribution was inferred from the AIC (Akaike Information Criteria) (panel A) and the BIC (Bayesian Information Criteria) (panel B) respectively. In this example, the inflection point occurs for a number of populations k=2, consistently with the mixture of two experiments. Using the results of the h-MSD analysis, we represented in panel C the distributions of alpha (upper graph) and D (lower graph). The distribution was then fitted with a GMM (Gaussian Mixture Model). The details of all combinations and the comparison of calculated values with the initial values are provided in the Supplemental Material.

Figure 4. FCS detects RPB1 anomalous diffusion in living cell nuclei. A. Representative picture of a
U20S RPB1-SYFP2 cell. B. Scatter plot of normalized residence times versus anomalous coefficients
computed from 138 FCS measurements performed in U2OS cells expressing RBP1-SYFP2 (n=6). 80% of
the transit times range from 23.7 and 89.1 ms/μm^3 with an average of 61.9 ms/μm^3 (D=5.7 μm^2/s$^\alpha$). And the
anomalous coefficients α range from 0.5 to 0.8 with a mean value of 0.61.

Figure 5. H-MSD SPT highlights the existence of distinct RPB1 subpopulations. A. Picture

representative of the signal-to-noise ratio obtained with JF549-halo tag at 0.1nM in U2OS RPB1-Halo cells.

B. Examples of trajectory obtained with the tracking pipeline. C. 1051 trajectories from SPT experiments

carried out in the U2OS RPB1-Halo cell line (n=10), stained with JF549 at 0.1nM, were analyzed by h-MSD.

Tt and α values from these experiments are represented in a scatter plot, in which concentric lines show the

Probability Density Function given by the Gaussian Mixture Model. At least two distinct populations are

clearly visible.

Figure 6. Spatio-temporal localization of RPB1 subpopulations. These images were acquired as a separate experiment within a 10s movie, after staining with JF549 at 1 nM. Panel A displays the mean intensity projection throughout the movie. Panel B illustrates the trajectories tracked at a given frame within the movie. Panel C represents the projection of all detected localizations using MTT in the whole film. Picture D merges the information from pictures A (in purple) and C (in cyan). The movie is also provided in supplementary data (Movies S3, S4 and S5), with raw data (S3), ongoing tracking superimposed (S4), or detections summed over time (S5).

Figure 7. Models for RNAP II recruitment. This scheme displays the two hypotheses proposed to explain
RNAP II mobilities and oversampling. The blue disks and arrows symbolize RNAP II molecules and their
displacement vectors, respectively. A. Model with successive steps: 1) A free RNAP II (α=0.95) enters and
oversamples a large domain (blue), potentially a TAD, where it is confined (α=0.61); 2) it is then trapped in
a subdomain (green) of higher confinement (α= 0.31), possibly through interactions with transcription
factors, and binds to chromatin, 3) before being stabilized and starting transcription (fixed RNAP II).

B. Model with distinct confinement domains (a and b) : 1a) and b) A free RNAPII (α=0.95) can enter two
types of confined domains, leading to fast oversampling or not; 2a) and b) RNAPII is then recruited and
binds to chromatin (fixed RNAPII). This second model helps to account for the existence of different
mechanisms of RNAPII recruitment and in particular the existence of the proximal transcriptional pause for
only half of genes.

SUPPLEMENTAL MATERIAL
709

Supplemental Figure 1. Mean number +/- standard deviation of detections obtained with Single Particle Tracking of fluorescent beads diffusing in aqueous solutions with different glycerol proportions.

A

Anomaly coefficient	Experimental values		Theory
% glycerol	mean α	SD	α
0	0.96	0.01	1
10	0.94	0.02	1
20	0.97	0.02	1
50	1.02	0.02	1
80	0.7	0.02	1

B

Diffusion coefficient	Experimental values		Theory
% glycerol	mean D_α (µm²/s)	SD	D (µm²/s)
0	15.5	1.7	15.1
10	12.1	1.8	12.1
20	9.2	1.5	9.0
50	4.5	0.8	3.2
80	1.1	0.2	0.4

Supplemental Table 1. Comparison of experimental vs theoretical values of anomaly coefficients (A) and diffusion coefficients (B) computed from the ACF of FCS measurements performed with fluorescent beads diffusing in aqueous solutions with different glycerol proportions**.**

712
713

Supplemental Figure 2. Representation of subpopulations identified through the GMM (Gaussian Mixture Model) after mixing batches of trajectories obtained in different conditions, as detailed within the graphics legend.

A

		Input	Calculated by H-MSD analysis
Mix 1 (m1)	D1 (µm²/s)	4.0	3.76 (SD=0.65)
	% pop 1	42%	40%
	D2 (µm²/s)	0.1	0.09 (SD=0.02)
	% pop 2	58%	60%
Mix 2 (m2)	D1 (µm²/s)	4.0	3.68 (SD=0.61)
	% pop 1	75%	73%
	D2 (µm²/s)	0.1	0.09 (SD=0.06)
	% pop 2	25%	27%
Mix 3 (m3)	D1 (µm²/s)	4.0	3.76 (SD=0.63)
	% pop 1	25%	24%
	D2 (µm²/s)	0.1	0.09 (SD=0.04)
	% pop 2	75%	76%

B

		Input	Calculated by H-MSD analysis
Mix 1 (m1)	D1 (µm²/s)	0.9	0.88 (SD=0.07)
	% pop 1	46%	43%
	D2(µm²/s)	0.1	0.10 (SD=0.05)
	% pop 2	54%	57%
Mix 2 (m2)	D1 (µm²/s)	0.9	0.84 (SD=0.06)
	% pop 1	58%	66%
	D2 (µm²/s)	0.1	0.11 (SD=0.04)
	% pop 2	42%	34%
Mix 3 (m3)	D1(µm²/s)	0.9	0.90 (SD=0.07)
	% pop 1	25%	23%
	D2 (µm²/s)	0.1	0.10 (SD=0.04)
	% pop 2	75%	77%

C

		Input	Calculated by H-MSD analysis
Mix 1 (m1)	D1 (µm²/s)	4.0	3.95 (SD=0.93)
	% pop 1	20%	15%
	D2 (µm²/s)	0.9	0.81 (SD=0.04)
	% pop 2	80%	85%
Mix 2 (m2)	D1(µm²/s)	4.0	4.01 (SD=0.45)
	% pop 1	51%	41%
	D2(µm²/s)	0.9	0.82 (SD=0.74)
	% pop 2	49%	59%
Mix 3 (m3)	D1(µm²/s)	4.0	4.10 (SD=0.45)
	% pop 1	78%	67%
	D2 (µm²/s)	0.9	0.87 (SD=0.42)
	% pop 2	22%	33%

D

		Input	Calculated by H-MSD analysis
Mix 1 (m1)	D1 (µm²/s)	4.0	1 population detected D=3.5µm²/s (SD=0.7)
	% pop 1	13%	
	D2 (µm²/s)	3.8	
	% pop 2	87%	
Mix 2 (m2)	D1 (µm²/s)	4.0	1 population detected D=3.6µm²/s (SD=0.9)
	% pop 1	46%	
	D2 (µm²/s)	3.8	
	% pop 2	54%	
Mix 3 (m3)	D1(µm²/s)	4.0	1 population detected D=3.8µm²/s (SD=0.8)
	% pop 1	75%	
	D2 (µm²/s)	3.8	
	% pop 2	25%	

Supplemental Table 2. Comparison of input values vs h-MSD computed values after mixing batches of trajectories obtained in different conditions.

Supplemental Figure 3. AIC (Akaike Information Criteria) (panel A) and BIC (Bayesian Information Criteria) (panel B) scores corresponding to the n-population analysis of RPB1 SPT trajectories

Supplemental Figure 4. Diffusion parameters for RPB1 measured by FCS (blue dots) and SPT (green dots and gray dots, respectively corresponding to the main and minority subpopulations detected) are gathered in this scatter plot.