High intraspecific growth variability despite strong evolutionary heritage in a neotropical forest

Sylvain Schmitt[1,*] (iD), Bruno Hérault[2,3,4] (iD), Géraldine Derroire[5] (iD)

[1] CNRS, UMR EcoFoG (Agroparistech, Cirad, INRAE, Université des Antilles, Université de la Guyane), Campus Agronomique, 97310 Kourou, French Guiana

[2] CIRAD, UPR Forêts et Sociétés, Yamoussoukro, Côte d'Ivoire

[3] Forêts et Sociétés, Université de Montpellier, CIRAD, Montpellier, France

[4] Institut National Polytechnique Félix Houphouët-Boigny, INP-HB, Yamoussoukro, Côte d'Ivoire

[5] Cirad, UMR EcoFoG (Agroparistech, CNRS, INRAE, Université des Antilles, Université de la Guyane), Campus Agronomique, 97310 Kourou, French Guiana

Correspondence: Sylvain Schmitt, UMR EcoFoG, Campus Agronomique, 97310 Kourou, French Guiana. Email: sylvain.m.schmitt@gmail.com

Abstract

Individual tree growth is a key determinant of species performance and a driver of forest

dynamics and composition. Previous studies on tree growth unravelled the variation in

species growth as a function of demographic trade-offs that are partially predicted using

functional traits. They have explored the environmental determinants of species growth

potential and the variation of intraspecific growth over space and time due to environment

and biotic factors. However, variation in individual growth within species remains

underexplored for a whole community and the relative role of species' evolutionary heritage

and of local environments remains unquantified. Here, based on 36 years of diameter

records for thousands of mapped individuals belonging to 138 species, we assessed

individual tree growth potential in a local neotropical forest community in the Amazon basin.

We further related variation in individual growth potential with taxonomic levels, local

topography, and neighbourhood crowding, before exploring species growth potential link to

functional traits and distribution along the phylogeny. We found that most of the variation in

growth potential was individual, and that taxonomic structure explained a third of the

observed variation. Species growth potential was phylogenetically conserved with positive

conservatism up to the genus level in the vast majority of species. Functional traits of roots,

wood and leaves together predicted species growth potential. Phylogeny suggested joint

selection of species' growth strategies and associated functional traits during convergent

evolutions. Finally, neighbourhood crowding had a significant effect on individual growth

potential, although much of this inter-individual variation remains largely unexplained and the

underlying ecological and evolutionary factors are still little explored. The high intraspecific

variation observed could allow individuals in these hyperdiverse ecosystems to respond to

the variable light and competitive conditions offered by successional niches during forest gap

dynamics.

Introduction

Individual growth is a key determinant of species performance (Violle et al., 2007) and a driver of forest dynamics (Hérault et al., 2010) and composition (Russo et al., 2008). Thus, a comprehensive understanding of the determinants of tree growth is of primary importance for predicting the fate of tropical forests, especially in the face of increasing anthropogenic disturbances. Tree growth plays a major role in the ecological strategies of tree species through demographic trade-offs (Rüger et al., 2019) including a growth-mortality trade-off (Aubry-Kientz et al., 2015; Philipson et al., 2014; Wright et al., 2010). This trade-off opposes fast-growing species to slow-growing species (Aubry-Kientz et al., 2015; Phillipson et al., 2014;), with lower support costs but at greater risk of damage and increased mortality (King et al., 2006). To forecast the dynamics of tropical forests, efforts have been made to predict the growth of species. To this end, functional traits, have been widely explored and used (Hérault et al., 2011, Visser et al., 2016, Osazuwa-Peters et al., 2017, Poorter et al., 2010). Functional traits are defined as phenotypic traits that have an impact on fitness through their effect on individual performance, which is defined as the ability to recruit, grow, survive and reproduce (Violle et al. 2007). Functional traits are therefore expected to play a role in species growth (but see Yang et al., 2018). For instance, wood density has been shown to be an important determinant of species growth potential (King et al., 2005; King et al., 2006; Hérault et al., 2011; Visser et al., 2016), as well as species maximum diameter and height, $\delta^{13}C$ of leaf (Hérault et al., 2011), wood anatomical features (Osazuwa-Peters et al., 2017; Poorter et al., 2010) and hydraulic conductance (Poorter et al., 2010).

Multiple environmental determinants of species growth have already been identified, which in turn affect species distribution and shape community structure. Light interception was recognised early on as a key factor in the growth of species (King et al., 2005). The role of light has notably been demonstrated experimentally in seedlings with species showing a

trade-off between shade tolerance and establishment in gaps (Baraloto et al., 2005). Fast-growing species are found in environments with high access to light, where shade-tolerant species show reduced growth (Wright et al., 2010). Treefalls open gaps in the forest creating bright environments and reduced competition from large trees. During forest succession, gaps are filled by vegetation, resulting in shadier environments and increased competition from large trees. This process, called forest gap dynamics (Martinez-Ramos *et al.,* 1989), produces successional niches, which favour species with a variety of survival and growth strategies (Hérault *et al.,* 2010; Rüger *et al.,* 2009). Fast-growing pioneer species quickly colonise treefall gaps, while slower-growing late-successional species gradually establish themselves in shadier environments (Craven *et al.,* 2015). Species distribution is also driven by soil (Kupers et al., 2019). Nutrient availability together with water shape species performance with fast-growing species dying more on the poorest habitat and slow-growing species being outcompeted in resource-rich habitats (Russo et al., 2008).

Nevertheless, access to light and nutrients, modulated by forest gap dynamics, topography and competition, also determines variation in individual growth within species over space and time. Indeed, species-specific functional traits still poorly predict individual tree rates (Yang et al., 2018), partly because they ignore individual variation, and because critical aspects of demographic rates are not captured by most measured functional traits. Individual- and species-based approaches to functional traits are conceptually fundamentally different (Poorter et al., 2018): the species-based approach focuses on potential and the individual-based approach focuses on realised traits and growth rates. Forest gap dynamics, notably through light interception variation, is a key factor shaping the growth of individuals within species with increased tree growth near canopy openings (Hérault et al., 2010; Schmitt *et al.,* 2022). Similarly, individuals with greater investment in biomass achieved higher growth rates with a suggested allocation in light capture (Umaña et al., 2018). Topography is driving both water and nutrient availability in tropical forests through the dissolution of iron oxides, litter- and tree-fall transfers and waterlogging (Ferry et al. 2010;

John et al. 2007). Therefore, individual growth within species is faster in valleys than on

ridges in neotropical forests (Fortunel et al., 2018; but see O'Brien Escudero 2021). The

effects of abiotic determinants on individual growth within species can be mediated by biotic

interactions and tree ontogeny. Neighbourhood crowding through competition for resources,

including light, nutrients and water, but also potentially through facilitation, is known to

modulate tree growth (Uriarte et al., 2004; Lewis and Tanner 2000). Individuals' responses

to neighbourhood crowding depend on the species identity of the focal tree and the

composition of its neighbours (Uriarte et al., 2004; Lewis and Tanner 2000). Neighbourhoods

interact with other abiotic factors, e.g. growth is faster in bottomland than on higher ground,

but neighbourhood crowding has a stronger negative effect on growth in bottomland than in

higher ground for one in ten species (Fortunel et al., 2018). Finally, individual tree growth

varies also during its life with ontogeny (Hérault *et al.*, 2011). Tree size, as a proxy for

ontogeny, is a major determinant of individual growth within species (Rees et al., 2010;

Hérault *et al.*, 2011; Gibert et al., 2016), even though tree size was shown to be less

important for growth than light (Rüger et al., 2011).

However, most of the studies of intraspecific variation in growth were carried out for a limited

number of species of a community, and the consistency of the observed patterns across

species of a community remains underexplored. This prevents assessing the relative

contribution to individual growth variability of species evolutionary heritage linked to

phylogenetic background and individual adaptation and plasticity in response to local

environments. Individual trees however differ in many dimensions within and among species

with genetic heritage and life history that may influence their growth response to local abiotic

environments and biotic interactions (Le Bec et al., 2015). Genetic processes within and

among species hold promises for understanding tree growth (Grattapaglia et al., 2009).

Evolutionary heritage resulted in convergent species growth within tree genera from the

Amazon Basin (de Souza *et al.*, 2016). A recent study further suggested individual

adaptation of tree growth within species to the successional niches generated by tropical

forest gap dynamics in the tropical species from the *Symphonia* genus (Schmitt et al., 2022).

The topographic origin of parent trees was also shown to determine the individual's growth

response of the offspring to increasing abundance of neighbours (O'Brien Escudero 2022).

Altogether, these results plead for a better integration of species and individuals in tree

growth studies, in order to better link individual tree growth to genetic, phylogenetic and

environmental variations.

Here, we assessed tree growth potential for most individuals in a local neotropical forest

community in the Amazon basin, and related variation in individual growth potential to

taxonomic levels, local topography and neighbourhood crowding, in addition to the species

phylogeny. Locally, neighbourhood crowding is an indirect measure of light access and

forest gap dynamics, while topography reflects the distribution of water and soil nutrients.

Based on 36 years of diameter records for 7,961 of mapped individuals belonging to 138

species, we constructed individual ontogenetic growth trajectories to infer individual growth

potential using a hierarchical Bayesian model. We used individual growth potential in a linear

mixed model with topography, neighbourhood crowding and taxonomic level to explore the

environmental drivers and evolutionary heritage of tree growth. We further explored the

phylogenetic signal of species growth potential and the role of previously published

functional traits in explaining species growth potential. We specifically addressed the

following questions:

1. What is the importance of evolutionary history in the growth potential of tropical

trees?

2. Can functional traits help predict species growth potential?

3. How important is the variability of individual growth potential and how does the local

environment influence it?

We hypothesised species growth to have a phylogenetic signal with conserved interspecific

growth in part influenced by species functional traits. We expected high intraspecific

variation in tree growth influenced by topography and neighbourhood crowding.

Material and Methods

Study site

The study was conducted in the Guiana Shield, in the coastal region of French Guiana, at the Paracou field station (5°18′N, 52°53′W). The site is characterised by an average annual rainfall of 3041 mm and an average air temperature of 25.71°C (Aguilos et al. 2018). A rich tropical forest occupies this lowland area characterised by heterogeneous microtopographic conditions with numerous small hills generally not exceeding 45 m in altitude (Gourlet-Fleury et al. 2004). The site includes fifteen 6.25 ha plots and a 25 ha plot with trees mapped to the nearest metre and censused (diameter at breast height >10 cm) every 1 to 5 years since 1985. Nine of the plots were intentionally manipulated in 1986 with a range of disturbance intensities that created a variety of biotic environments (details of the experiment in Hérault and Piponiot 2018).

Species and individuals

We focused on trees located 20 metres from any plot edge for neighbourhood analyses. We used only (i) trees recruited since the beginning of censuses to exclude large diameter trees that show little to no variation in their growth trajectories, (ii) trees with at least 10 measurements to better assess their complete growth trajectories and (iii) species with at least 10 trees meeting the previous requirements for a good representation of intraspecific variability. We did include disturbed plots in our study in order to explore a greater variety of neighbourhood crowding.

Individual growth

We first used a reduced dataset to explore the best model shape (Tab. S1) and the best hierarchical integration of individual and species effects (Tab. S2) to infer individual growth

potential. Based on goodness of fit (likelihood), cross-validation (leave-one-out estimate of

the expected log pointwise predictive density, see Vehtari *et al.*, 2007) and prediction quality

(root mean square error of prediction), we chose to model tree diameter over time using the

sum of the arithmetic series of the annual tree growth rate observed at the census year for

each individual with a Gaussian distribution (model 1). At any year t since tree recruitment,

individual annual growth rate $AGR_{i,t}$ can be defined following a Gompertz model (Hérault *et*

al., 2011) based on individual diameter at breast height from previous census $DBH_{i,t-1}$:

184
$$AGR_{i,t} = Gmax_i \times \exp\left(\frac{-1}{2} \left[\frac{\log\left(DBH_{i,t-1} / Dopt_i\right)}{Ks_i} \right]^2 \right)$$

with $Dopt_i \ N\left(Dopt_s, \sigma D\right)$ and $Ks_i \sim N\left(Ks_s, \sigma K\right)$

where $Gmax_i$ is the individual maximum growth potential, $Dopt_i$ is the optimal diameter at

which the individual reaches its maximum growth potential, and Ks_i is the kurtosis defining

the width of the bell-shaped growth-trajectory (see figure 1 in Hérault *et al.*, 2011). $Dopt_i$

and Ks_i are random effects centred on species parameters $Dopt_s$ and Ks_s with associated

variances σD and σK. Thus, annual individual diameter $DBH_{i,t}$ can be calculated as

previous year individual diameter $DBH_{i,t-1}$ plus individual annual growth rate $AGR_{i,t}$:

192
$$DBH_{i,t} = DBH_{i,t-1} + AGR_{i,t}$$

193   This arithmetic serie can be integrated from individual recruitment to build individual growth

194   trajectory with annual individual diameter $DBH_{i,t}$:

195
$$DBH_{i,t} = DBH0_i + \sum_{y=0}^{y=t} AGR_{i,y}$$

196   where $DBH0_i$ is individual diameter at recruitment. Finally, a model can be fitted to predict

197   annual individual diameter $DBH_{i,t}$ with observed diameter from censuses using a Gaussian

198   distribution:

199
$$DBH_{i,t} \sim N\left(DBH0_i + \sum_{y=0}^{y=t} AGR_{i,y}, \sigma\right)$$

200   This model was fitted for every census t at which the individual was measured. A Bayesian

201   method was employed to infer parameters using the stan language (Carpenter *et al.*, 2017)

202   and the *rstan* package (Stan Development Team, 2018) in the R environment (R Core

203   Team, 2020).

204 Descriptors of individual growth potential

We used the mean neighbourhood crowding index (NCI, Uriarte *et al.,* 2004) over the last 30

years, an indirect measurement of access to light and forest gap dynamics for each

individual. The mean neighbourhood crowding index NCI_i from tree individual i was

calculated as follows:

$$NCI_i = \overline{\sum_{(j|\delta_{i,j}<20m)} DBH_{j,t}^2 \times \exp\left(\frac{-1}{4}\delta_{i,j}\right)}$$

with $DBH_{j,t}$ the diameter of the neighbouring tree j in year t and $\delta_{i,j}$ its distance to the

individual tree i. NCI_i is computed for all neighbours at a distance $\delta_{i,j}$ inferior to the

maximum neighbouring distance of 20 metres. The power of neighbours $DBH_{j,t}$ effect was

set to 2 to represent an area. The decrease of neighbours' diameter effect with distance was

set to -¼ to represent trees at 20 metres of the focal trees having less than 1% of the effect

of the same tree at 0 metres (*i.e.* exp(-¼ x 20) < 1%). NCI_i is computed as the mean of

$NCI_{i,t}$ per census over the last 30 years denoted by the overline in the equation.

We used the topographic wetness index (TWI) as proxy of the distribution of soil water and

nutrients in Paracou (Schmitt et al. 2021). Waterlogging and topography have been

highlighted as crucial for forest dynamics (Ferry et al. 2010), species-habitat relationships

(Engelbrecht et al. 2007), and phenotypic variation (Schmitt et al. 2020). TWI was derived

from a 1-m-resolution digital elevation model using SAGA-GIS (Conrad et al. 2015) based on

a LiDAR campaign of the whole Paracou field station done in 2015.

We also tested the link between functional traits and species growth using the mean trait
values of the 120 species that Vleminckx et al. (2021) shared with our study (over 138, Fig.
S10). Vleminckx et al. (2021) functional data included 19 leaf, stem and root traits. Leaf traits
included carbon, nitrogen, potassium, phosphorus, calcium, and chlorophyll content, carbon
13 isotopic ratio, toughness, thickness, area and specific area. Stem traits included sapwood
specific gravity and trunk bark thickness. Root traits included specific tip abundance, specific
length, wood specific gravity and fine roots tissue density and diameter.

232 Analyses

To study the effect of phylogeny and environment, we investigated the effects of family,
genus, species, topography (TWI) and neighbourhood (NCI) indices on individual growth
potential (Gmax) with a linear mixed model (model 2). Environmentals variables (TWI and
NCI) were used as fixed effects, while taxonomic variables (family, genus, and species) were
used as random effects. We propagated the uncertainty in the estimation of individual growth
(model 1) to the model testing the effect of phylogeny and environment (model 2). To do so,
we considered the individual growth potentials obtained in 500 iterations of model 1 as 500
different dataset on which we fitted a Bayesian linear mixed model (model 2). For each of
these dataset, we kept the 1000th iteration after a warm up of 999 iterations. We pooled all
these last iterations to obtain a posterior distribution of the effect of phylogeny and
environment that takes into account the uncertainty in individual growth potential. We
reported the resulting marginal (fixed effects alone) and conditional (fixed and random
effects) goodness of fit (R², Nakagawa & Schielzeth 2013).

We further investigated species growth potential across the phylogeny. To account for the
lognormal distribution of individual growth potential within species (Schmitt et al. 2022), we
used the median as a measure of species growth potential ($Gmax_s = Median(Gmax_i)$). We
tested the phylogenetic signal of species growth potential with multiple measures including

Pagel's λ (Keck *et al.*, 2016). We further computed the phylogenetic correlogram of species

growth potential using partistic phylogenetic distance and tested its significance with 999

bootstrap assuming brownian motion. We explored the phylogenetic signal structure across

the phylogeny with the local indicator of phylogenetic association.

We finally explored the link between species functional traits and species growth potential.

We used a linear model with a step selection of the best model explaining species log-

transformed growth potential with log-transformed functional traits to meet the normality

assumptions (model 3). We measured functional traits relative importance in explaining

species growth potential averaging over orderings of regressors (Groemping 2007). We

finally compared the taxonomic and phylogenetic structures of functional traits with those of

species growth potential using linear mixed models with random taxonomic levels (family,

genus, and species) and phylogenetic correlograms. We used the *tidyverse* (Wickham *et*

al., 2019), *lme4* (Bates *et al.,* 2015), *phylosignal* (Keck *et al.,* 2016)*,* and *ggtree* (Yu 2020)

packages in the R environment (R Core Team, 2020) for all analyses.

Results

We recorded 117,688 diameter measurements in the local community across 7,961

individuals belonging to 138 species, 95 genera and 38 families. Bayesian inference of the

individual growth model converged correctly (*e.g.,* Fig. S1) with a good mixture of chains for

most of the individual growth potentials ($Gmax_i$) shown with an $\hat{R}<1.1$ (99.2%, Fig. S2). The

posteriors of the parameters did not show any correlation problems (*e.g.,* Fig. S3). Posterior

distributions of individual growth potentials showed limited uncertainty (*e.g.,* Fig. S4). The

resulting predicted diameters showed very good goodness of fit with a root mean square

error of 0.43 centimetres (Fig. 1, illustrated for 90 individuals in 9 species in Fig. S5).

Based on the inferred individual growth potentials, we found most of the variation in growth

potential to be among individuals within species shown with residuals ($\sigma=0.72$, Tab. 1), then

among genera within families ($\sigma=0.12$), then among species within genera ($\sigma=0.04$) and

among families ($\sigma=0.03$). The taxonomic structure explained about a third of the observed

variation in individual growth potential (Conditional $R^2=0.347$ against Marginal $R^2=0.019$).

The neighbourhood crowding index (NCI) had a marked negative significant effect on

individual growth potential ($\beta=-0.51, CI=[-0.61;-0.42]$, Tab. 1, Fig. 2A, Fig. S6) which

explained 2% of the observed variation knowing taxonomy (Marginal $R^2=0.019$, Tab. 1). The

topographic wetness index had no effect on individual growth potential (

$\beta=-0.03, CI=[-0.13;-0.07]$, Tab. 1, Fig. S6).

Species growth potential was significantly structured in the phylogeny (Pagel's

$\lambda=0.78, p<0.001$, Fig. 2B, Tab. S3). Phylogenetic autocorrelograms revealed a short

distance significant positive association and a long distant significant negative association of

species growth potential in the phylogeny (Fig. S7). The local indicator of phylogenetic

association highlighted the conservation of species growth potential at the genus level (Fig.

S8), as illustrated for instance with fast growing species from the *Cecropia* genus opposed to

slow growing species from the *Eschweilera* genus (Fig. 2C). However, a few species have

different growth potential in the same genus, such as slow growing *Drypetes variabilis*

opposed to fast growing *Drypetes fanshawei*.

Functional traits explained 40% of the observed variation of species growth potential among

120 tropical tree species ($R^2=0.394$). Leaf potassium content, sapwood specific gravity, fine

roots diameter and specific length had significant negative effects on species growth

potential, while leaf nitrogen content had a significant positive effect (Tab. 2). Sapwood

specific gravity and leaf nitrogen content had the largest relative importance (resp. $R^2=0.18$

and $R^2=0.11$), while other traits had small relative importance ($R^2\leq0.04$). Moreover, these

six functional traits had a phylogenetic autocorrelogram similar to species growth potential

with a short distance significant positive association, stronger for sapwood specific gravity

than other traits (Fig. S9).

Figure 1: Goodness of fit for Bayesian inference of the individual growth model, illustrated by

predicted versus observed diameters at breast height. The dots represent the median of the

posterior distributions of predicted diameters while the error bars show the 90% credibility

intervals. The red dashed line represents the expected value for a perfect prediction (1:1).

The X and Y axes have been log-transformed to better represent the many small diameter

values. An example of corresponding diameter trajectories can be found in Figure S4.

Table 1: Effects of phylogeny and environment on individual growth potential. We

investigated the effects of family, genus, species, topography (TWI) and neighbourhood

crowding (NCI) indices on individual growth potential (Gmax) with a linear mixed model. First

rows show estimates and credible intervals (CI) for environmental fixed effects. Middle rows

show estimates for variance components from phylogenetic random effects. The last row

shows marginal (fixed effects alone) and conditional (fixed and random effects) goodness of

fit.

$\log\left(Gmax_i\right)$		
Predictors	*Estimates*	*CI (95%)*
Intercept	3.88	3.05 – 4.68
$\log\left(NCI_i\right)$	-0.51	-0.61 – -0.42
$\log\left(TWI_i+1\right)$	-0.03	-0.13 – 0.07
Random effects		
Residual (Individual)	0.72	
Species | Genus | Family	0.04	
Genus | Family	0.12	
Family	0.03	
Marginal R² / Conditional R²	0.019 / 0.347	

322
323

*Figure 2: Variation in the growth potential of individuals and species as a function of neighbourhood crowding, taxonomic levels or across phylogeny. **A.** Individual growth potential ($Gmax_i$) is significantly decreasing with neighbourhood crowding index (NCI, $\beta = -0.51$, $Marginal\ R^2 = 0.019$, see Tab. 1). **B.** The partitioning of the variation of individual growth potential ($Gmax_i$) across taxonomy shows that most of the variation occurs at the individual ($\sigma = 0.72$, Tab. 1), then genus ($\sigma = 0.12$) before species ($\sigma = 0.04$) and family ($\sigma = 0.03$) levels. **C.** The distribution of species growth potential (Median[$Gmax_i$]) across the phylogeny from slow growing species in dark blue to fast growing species in yellow (log-scale) is phylogenetically structured (Pagel's $\lambda = 0.78$, $p < 0.001$) with a significant positive autocorrelation to a phylogenetic partistic distance below 100 (Fig. S1), corresponding to the genus level (Tab. 1).*

Table 2: Effects of functional traits on species growth potential. We extracted traits from
Vleminckx et al. (2021) for the 120 species shared with our study, and used a linear model
with a step selection of the best model explaining species growth potential with log-
transformed growth potential and functional traits. We thus selected 6 root, stem and leaf
functional traits. Columns show estimates, confidence intervals (CI), significance (p-value),
and relative importance (R²) for selected functional traits. The last row shows raw and
adjusted goodness of fit.

	$\log\left(Gmax_s\right)$			
Predictors	*Estimates*	*CI*	*p-value*	*R²*
Intercept	-0.01	-0.15 – 0.13	0.882	
leaf toughness	0.18	-0.01 – 0.37	0.063	0.02
leaf nitrogen content	0.38	0.22 – 0.55	**<0.001**	0.11
leaf potassium content	-0.24	-0.41 – -0.07	**0.005**	0.04
sapwood specific gravity	-0.57	-0.76 – -0.39	**<0.001**	0.18
fine roots diameter	-0.49	-0.82 – -0.17	**0.003**	0.04
specific root length	-0.35	-0.67 – -0.02	**0.035**	0.03
R² / R² adjusted	0.424 / 0.394			

Discussion

Using 36 years of diameter records for thousands of mapped individuals belonging to 138 species, we found that the evolutionary heritage of species shaped individual growth trajectories, explaining up to one third of the observed variation in growth potential among the eight thousand individuals studied. Functional traits of wood, leaves and roots together predicted species growth potential, showing that multiple functional dimensions determine the performance of tropical tree species in their environments. Phylogenetic correlograms suggested joint selection of species' growth strategies and associated functional traits during convergent evolutions. Nevertheless, at the individual tree level, the observed effect of neighbourhood crowding suggests that forest gap resulted in fast-growing trees in high light conditions with reduced competition, as opposed to slow-growing trees in low light conditions with strong competition. But the ecological and evolutionary drivers of the high variability in individual growth potential remain largely undetermined, and the underlying factors, which include phenotypic plasticity and genetic adaptations, are little explored. The high intraspecific variation observed could allow individuals in hyperdiverse ecosystems such as tropical forests to respond to the variable light and competitive conditions offered by successional niches during forest gap dynamics, in addition to the many other potential environmental dimensions that shape the coexistence of species.

Evolutionary history shapes the growth of tropical trees

We found that evolutionary history shaped individual growth potential. Previous studies already highlighted regionally a phylogenetic signal in species growth potential in the Amazon Basin, with evolutionarily related genera having more similar growth values than expected by chance (de Souza *et al.*, 2016). Regionally, woody biomass and tree size also showed a strong phylogenetic signal in tropical dry forests (de Aguiar-Campos *et al.*, 2021),

besides longevity rather than growth determined biomass (Körner 2017). Similarly, Cadotte

et al., (2009) found evolutionary relationships explaining grasslands productivity. Thus,

evolutionary history shapes observed phylogenetic and functional diversity at the regional

scale, with for instance lasting influences from the Paleoclimate (Svenning *et al.*, 2015;

Bosela *et al.*, 2016). Our results are in agreement as we also identified a strong phylogenetic

signal in species growth potential with positive phylogenetic correlation up to the genus level,

showing that previous regional scale results hold locally. To our knowledge, our study is the

first attempt to link individual growth potential to species growth potential. Using taxonomy

as a proxy for evolutionary heritage of species, we found that the evolutionary history of

species explained up to one third of the observed variation in growth potential among the

eight thousand individuals studied. Thus, evolutionary history is an important determinant of

individual variation in growth that is stronger than the direct effect of the environment, at

least for the topography and neighbourhood crowding tested here.

Several convergent evolution events resulted in similar growth patterns of species within

genera with, for instance, repeated evolution of fast-growing species in *Urticaceae*,

Fabaceae, *Hypericaceae*, and *Melastomataceae* families with species respectively

belonging to the *Cecropia*, *Inga*, *Vismia* and *Miconia* genera. This pattern might be explained

by repeated evolution constrained by forest gap dynamics, leading to shade-tolerance under

closed canopies as opposed to fast-growing pioneer species in light gaps (de Souza *et al.*,

2016). Indeed, *Cecropia, Visimia* and *Miconia* are for instance recognized as pioneer

species colonising first light gaps after a treefall (Dalling *et al.*, 1998). Convergent evolutions

of habitat specialisation could also explain the divergent growth of genera within families. For

instance at the species level, Fine *et al.,* (2014) evidenced specialisation to white sands or

flooded soils within *Protieae*. Nevertheless, the fact that topography and forest gap

dynamics, two major determinants of the local community (e.g. Ferry *et al.*, 2010; Molino and

Sabatier 2001), have less explanatory power than taxonomy reveals the importance of past

evolutionary constraints on the growth trajectories of individual trees through species

heritage compared to the direct effect of the current environment to which individual trees
could respond through microadaptations and phenotypic plasticity (*e.g.* Schmitt *et al*., 2021;
2022).

Taken together, our results therefore partly support species-averaged community ecology
approaches, at least for growth trajectories (but see last paragraph). Swenson *et al*., (2013)
suggested the use of phylogenetic approaches to understand community assembly.
However, phylogeny alone is not sufficient to predict demographic rates (Che-Castaldo *et*
al., 2018). Indeed, phylogenetic approaches must be conducted with caution given the high
remaining intraspecific variability (see last paragraph) and the differences observed among
closely-related species.

Multiple functional dimensions together predict the species growth potential

Using the 120 species in common with Vleminckx *et al.* (2021), we found six functional traits
explaining an important part of variation in species growth potential (40%). Sapwood specific
gravity was already shown to be a major predictor of species growth (King et al., 2005; King
et al., 2006; Hérault et al., 2011; Visser et al., 2016), with fast-growing species investing less
in wood resulting in a low density as opposed to slow-growing ones. Leaf nitrogen content
was the second most important predictor in our study, with nitrogen-rich species growing
faster, as already observed for pioneer species (Aidar *et al.*, 2003), and in rich environments
(Russo et al., 2008). Leaf potassium and toughness were also small predictors of species
growth potential. Finally, root traits also predicted species growth potential with fast-growing
species investing less in their root systems with decreased fine roots diameters and higher
specific root length, as already shown for temperate species (Comas, Bouma and Eissenstat
2002, Comas and Eissenstat 2004) but never in the tropics, in the limit of our knowledge.

Soil heterogeneity can be expected to play a key role in shaping root traits (Vleminckx *et al.,*

2021), while the dynamics of forest gaps cannot be excluded either (Xiang *et al.*, 2013).

Future research on the role of root traits in tropical tree growth is thus particularly promising

given that tropical forests harbour the greatest diversity of root characteristics (Ma *et al.*,

2018). Interestingly, species growth potential is better predicted by the combination of traits

from roots, wood and leaf, in accordance with the idea of multiple functional dimensions that

allow tropical tree species to optimise their performance in a given environment Vleminckx *et*

al. (2021), while participating together in a whole plant economic spectrum (Reich 2014).

Finally, we found a similar phylogenetic structure between the 6 functional traits and species

growth potential, with significant and positive phylogenetic correlation at short distance, in

particular in sapwood specific gravity (de Souza *et al.*, 2016). The traits explaining the most

species growth had the strongest positive phylogenetic correlation (*e.g.* sapwood specific

gravity), while traits such as leaf potassium content and toughness had smaller positive

phylogenetic correlation only significant to a short phylogenetic distance. The similarity

between phylogenetic correlograms suggests a joint selection of species growth strategies

and associated functional traits during convergent evolutions. This result further supports the

use of phylogeny and related traits to predict species growth trajectories and potential

demography (Swenson 2014; Tucker *et al.*, 2018; Paquette *et al.,* 2015).

442 Individual growth potential is influenced by forest gap dynamics

443 but remains largely unexplained

We found evolutionary heritage shaping the individual growth of tropical trees, with functional

traits as important predictors of species growth potential. However, we still observe a huge

intraspecific variation with individual growth potential varying widely within species

(logarithmic coefficient of variation of 98% [35-430%]). The growth trajectories of individual

trees are strongly and negatively affected by the average neighbourhood crowding over the

last 30 years, which can be related to the mosaic of light and competition environments

shaped by forest gap dynamics (Schmitt *et al.,* 2022). In a nutshell, we observe fast-growing

trees in high-light conditions with decreased competition opposed to slow-growing trees in

low-light conditions with strong competition within and among species. Fast-growing pioneer

species opposed to slow-growing species under closed-canopies are widely known and

expected at the interspecific level (Dalling*et al.*, 1998; Molino and Sabatier 2001; King et al.,

2005; Hérault et al., 2011), but we also evidenced fast-growing individual opposed to slow-

growing individual within species along successional niches, advocating for a wide breadth

of successional niches in tropical tree species (e.g. Schmitt *et al.,* 2022 for *Symphonia*

species). The effect of competition and its reduction has already been suggested as a factor

increasing tree radial growth within species in the context of selective logging, especially

within shade-tolerant species (Peña-Claros *et al.,* 2008). Indeed, logging gaps, with

increased light access, results in increased tree growth at short distance from the gap,

especially within slow-growing species (Hérault *et al.,* 2010). Rare species have been also

shown to be more sensitive to light variation (Rüger *et al.,* 2011). The effect of crowding can

be a direct limit to light access expected in successional niches, but the effect of neighbours

identity found in tropical trees (Potvin *et al.,* 2008) also suggest other above- and below-

ground competition and facilitation processes. In short, forest gap dynamics with tree fall

results in successional niches ranging from high light and low competition after recent tree

fall to low light and high competition in closed canopies. We found along this gradient fast-

growing species and individuals within species in early-succession niches as opposed to

slow-growing species and individuals within species in late-succession niches.

Nevertheless, forest gap dynamics only explained a tenth of the high variation of individual

growth potential within species. Topographic position, proxied with topographic wetness

index, did not influence the individual growth variation within species in our study (but see

O'Brien and Escudero 2021), despite its known importance in tropical forest dynamics (Ferry

et al., 2010). The role of topography on individual growth potential was weak, but we can

assume that topography shapes species growth through species evolutionary heritage, as

several species studied showed locally pervasive habitat preferences along topography

(Allié *et al.,* 2018; Schmitt *et al.,* 2021b) and microgeographic adaptations to topography

(Schmitt *et al.,* 2021). Consequently, ecological and evolutionary factors of individual growth

potential remain largely undetermined. The process through which forest gap dynamics and

undetermined factors affect individual growth potential also remains underexplored. The

spatio-temporal variation of forest gap dynamics led to assume growth potential variation

within species to be due to phenotypic plasticity (dos Santos *et al.,* 2020). However, recent

studies revealed local adaptation (O'Brien and Escudero 2021) and microadaptation

(Schmitt *et al.,* 2022) of individual trees within species to neighbourhood crowding and

competition resulting in varying individual growth potential. Thus forest gap dynamics could

have both a strong evolutionary heritage on species and may shape genotypes within

species with strong spatio-temporal variations (Schmitt *et al.,* 2022).

Intraspecific variability in performance can have strong implications for species coexistence

(Chesson 2000, Clark 2010, see also Stump *et al.,* 2022 for a synthesis), although this view

remains an open question (*e.g.* Clark *et al.,* 2022). Clark (2010) suggested that intraspecific

variability allows species to differ in the distribution of their responses to the environment

and thus to pass environmental filtering: an individual may persist in a given environment

with a suitable phenotype while the same environment would have filtered out the average

phenotype of the species. This hypothesis is consistent with theories that predict the

coexistence of a greater number of species, with competition being stronger within species,

among individuals, than among species (Chesson 2000). Modelling approaches further

support the hypothesis that intraspecific genetic and phenotypic variability promotes species

coexistence (Lichstein *et al.,* 2007). In the case of forest gap dynamics, late-successional

species have been shown to have more variation in response to competition and light

variation than early-successional species (Peña-Claros *et al.,* 2008; Hérault *et al.,* 2010),

which could be linked to a greater diversity of light and competitive environments. The high

intraspecific variation observed (logarithmic coefficient of variation of 98%) could therefore

allow individuals in these hyperdiverse ecosystems to adjust to the variable conditions of

light and competition offered by the successional niches during the closure of the forest

gaps, in addition to the other numerous potential niches shaping the high-dimensional

coexistence of species (Clark 2010). Consequently, the methodology used in our study

paves the way to future research on determinants and processes shaping tree growth within

and among species. The combination of studies from forest censuses (e.g. this study),

modelling approaches (e.g. Schmitt et al., 2020b), and experimental studies (e.g. O'Brien

and Escudero 2021) holds promise for a better understanding of tree performance within

and among species, with the potential to better explain and predict species coexistence and

forest dynamics. In particular, linking individual genotypes within species to forest gap

dynamics, topography and individual growth potential (Schmitt *et al.,* 2022) among several

species to extrapolate to the community level seems a promising approach to elucidate the

determinants of tree growth.

Acknowledgements

520 We are grateful to Pascal Petronelli, Julien Engel, Christopher Baraloto and the CIRAD

521 inventory team for their tremendous work in tree inventory and botanical identification,

522 without which this entire study would simply not have been possible.

Funding

524 This research received no specific grant.

Conflict of interest disclosure

526 The authors declare they have no conflict of interest relating to the content of this article.

Data, script and code availability

DBH and spatial positions of individuals were extracted from the Paracou Station database, for which access is available at https://dataverse.cirad.fr/dataverse/paracou with corresponding DOIs given in table S4. Scripts used for analyses can be found on GitHub (https://github.com/sylvainschmitt/treegrowth).

Authors contribution

SS conceived the ideas and designed the methodology. SS, BH, and GD analysed model outputs. SS led the writing of the manuscript. All authors contributed critically to the drafts and gave final approval for publication.

References

Aguilos, M., Stahl, C., Burban, B., Hérault, B., Courtois, E., Coste, S., … Bonal, D. (2018). Interannual and Seasonal Variations in Ecosystem Transpiration and Water Use Efficiency in a Tropical Rainforest. *Forests*, *10*(1), 14. https://doi.org/10.3390/f10010014

Aidar, M. P. M., Schmidt, S., Moss, G., Stewart, G. R., & Joly, C. A. (2003). Nitrogen use strategies of neotropical rainforest trees in threatened Atlantic Forest. *Plant, Cell and Environment*, *26*(3), 389–399. https://doi.org/10.1046/j.1365-3040.2003.00970.x

Allié, E., Pélissier, R., Engel, J., Petronelli, P., Freycon, V., Deblauwe, V., … Baraloto, C. (2015). Pervasive local-scale tree-soil habitat association in a tropical forest community. *PLoS ONE*, *10*(11), e0141488. https://doi.org/10.1371/journal.pone.0141488

Aubry-Kientz, M., Rossi, V., Boreux, J.-J., & Hérault, B. (2015). A joint individual-based model coupling growth and mortality reveals that tree vigor is a key component of tropical forest dynamics. *Ecology and Evolution*, *5*(12), 2457–65. https://doi.org/10.1002/ece3.1532

Aubry-Kientz, M., Rossi, V., Wagner, F., & Hérault, B. (2015). Identifying climatic drivers of tropical forest dynamics. *Biogeosciences*, *12*(19), 5583–5596. https://doi.org/10.5194/bg-12-5583-2015

Baraloto, C., Goldberg, D. E., & Bonal, D. (2005). Performance trade-offs among tropical tree seedlings in contrasting microhabitats. *Ecology*, *86*(9), 2461–2472. https://doi.org/10.1890/04-1956

Bates, D., Mächler, M., Bolker, B. M., & Walker, S. C. (2015). Fitting linear mixed-effects models using lme4. *Journal of Statistical Software*, *67*(1). https://doi.org/10.18637/jss.v067.i01

Bauman, D., Fortunel, C., Cernusak, L. A., Bentley, L. P., McMahon, S. M., Rifai, S. W., … Malhi, Y. (2021). Tropical tree growth sensitivity to climate is driven by species intrinsic growth rate and leaf traits. *Global Change Biology*, 1–19. https://doi.org/10.1111/gcb.15982

Bosela, M., Popa, I., Gömöry, D., Longauer, R., Tobin, B., Kyncl, J., … Büntgen, U. (2016). Effects of post-glacial phylogeny and genetic diversity on the growth variability and climate sensitivity of European silver fir. *Journal of Ecology, 104*(3), 716–724. https://doi.org/10.1111/1365-2745.12561

Cadotte, M. W., Cavender-Bares, J., Tilman, D., & Oakley, T. H. (2009). Using phylogenetic, functional and trait diversity to understand patterns of plant community productivity. *PLoS ONE, 4*(5), 1–9. https://doi.org/10.1371/journal.pone.0005695

Carpenter, B., Gelman, A., Hoffman, M. D., Lee, D., Goodrich, B., Betancourt, M., … Riddell, A. (2017). Stan : A Probabilistic Programming Language. *Journal of Statistical Software*, *76*(1). https://doi.org/10.18637/jss.v076.i01

Che-Castaldo, J., Che-Castaldo, C., & Neel, M. C. (2018). Predictability of demographic rates based on phylogeny and biological similarity. *Conservation Biology, 32*(6), 1290–1300. https://doi.org/10.1111/cobi.13135

Chesson, P. (2000). Mechanisms of Maintenance of Species Diversity. *Annual Review of Ecology and Systematics*, *31*(1), 343–366. https://doi.org/10.1146/annurev.ecolsys.31.1.343

Clark, D. A., Piper, S. C., Keeling, C. D., & Clark, D. B. (2003). Tropical rain forest tree growth and atmospheric carbon dynamics linked to interannual temperature variation during 1984-2000. *Proceedings of the National Academy of Sciences of the United States of America*, *100*(10), 5852–5857. https://doi.org/10.1073/pnas.0935903100

Clark, J. S. (2010). Individuals and the Variation Needed for High Species Diversity in Forest Trees. *Science*, *327*(5969), 1129–1132. https://doi.org/10.1126/science.1183506

James Clark, Adam Clark, Benoit Courbaud, et al. A universal coexistence hypothesis resolves the biodiversity paradox: Species differences that generate diverse forests. *Authorea.* August 12, 2022. https://doi.org/10.22541/au.166029509.96486640/v1

Comas, L. H., Bouma, T. J., & Eissenstat, D. M. (2002). Linking root traits to potential
growth rate in six temperate tree species. *Oecologia*, *132*(1), 34–43.
https://doi.org/10.1007/s00442-002-0922-8

Comas, LH and Eissenstat, D. (2004). Linking Fine Root Traits to Maximum Potential
Growth Rate among 11 Mature Temperate Tree Species. *Functional Ecology*,
388–397.

Conrad, O., Bechtel, B., Bock, M., Dietrich, H., Fischer, E., Gerlitz, L., … Böhner, J.
(2015). System for Automated Geoscientific Analyses (SAGA) v. 2.1.4.
Geoscientific Model Development, *8*(7), 1991–2007. https://doi.org/10.5194/gmd-
8-1991-2015

Craven, D., Hall, J. S., Berlyn, G. P., Ashton, M. S., & van Breugel, M. (2015).
Changing gears during succession: shifting functional strategies in young tropical
secondary forests. *Oecologia*, *179*(1), 293–305. https://doi.org/10.1007/s00442-
015-3339-x

Dalling, J. W., Swaine, M. D., & Garwood, N. C. (1998). Dispersal patterns and seed
bank dynamics of pioneer trees in moist tropical forest. *Ecology*, *79*(2), 564–578.
https://doi.org/10.1890/0012-9658(1998)079[0564:DPASBD]2.0.CO;2

de Aguiar-Campos, N., Coelho de Souza, F., Maia, V. A., Rezende, V. L., de Souza, C.
R., de Paula, G. G. P., … dos Santos, R. M. (2021). Evolutionary constraints on
tree size and above-ground biomass in tropical dry forests. *Journal of Ecology,*
109(4), 1690–1702. https://doi.org/10.1111/1365-2745.13589

de Souza, C. F., Dexter, K. G., Phillips, O. L., Brienen, R. J. W., Chave, J., Galbraith, D.
R., … Baker, T. R. (2016). Evolutionary heritage influences Amazon tree ecology.
Proceedings. Biological Sciences, 283(1844)

dos Santos, V. A. H. F., & Ferreira, M. J. (2020). Are photosynthetic leaf traits related to
the first-year growth of tropical tree seedlings? A light-induced plasticity test in a
secondary forest enrichment planting. *Forest Ecology and Management*, *460*,
117900. https://doi.org/10.1016/j.foreco.2020.117900

Engelbrecht, B. M. J., Comita, L. S., Condit, R., Kursar, T. A., Tyree, M. T., Turner, B.
L., & Hubbell, S. P. (2007). Drought sensitivity shapes species distribution patterns
in tropical forests. *Nature*, *447*(7140), 80–82. https://doi.org/10.1038/nature05747

Ferry, B., Morneau, F., Bontemps, J.-D., Blanc, L., & Freycon, V. (2010). Higher treefall
rates on slopes and waterlogged soils result in lower stand biomass and
productivity in a tropical rain forest. *Journal of Ecology*, *98*(1), 106–116.
https://doi.org/10.1111/j.1365-2745.2009.01604.x

Fine, P. V. A., Zapata, F., & Daly, D. C. (2014). Investigating processes of neotropical
rain forest tree diversification by examining the evolution and historical
biogeography of the protieae (Burseraceae). *Evolution*, *68*(7), 1988–2004.
https://doi.org/10.1111/evo.12414

Fortunel, C., Lasky, J. R., Uriarte, M., Valencia, R., Wright, S. J., Garwood, N. C., &
Kraft, N. J. B. (2018). Topography and neighborhood crowding can interact to
shape species growth and distribution in a diverse Amazonian forest. *Ecology*.
https://doi.org/10.1002/ecy.2441

Grattapaglia, D., Plomion, C., Kirst, M., & Sederoff, R. R. (2009, April 1). Genomics of
growth traits in forest trees. *Current Opinion in Plant Biology*. Elsevier Current
Trends. https://doi.org/10.1016/j.pbi.2008.12.008

Grömping, U. (2006). Relative importance for linear regression in R: The package
relaimpo. *Journal of Statistical Software*, *17*(1), 1–27.
https://doi.org/10.18637/jss.v017.i01

Gibert, A., Gray, E. F., Westoby, M., Wright, I. J., & Falster, D. S. (2016). On the link
between functional traits and growth rate: meta-analysis shows effects change
with plant size, as predicted. *Journal of Ecology*, *104*(5), 1488–1503.
https://doi.org/10.1111/1365-2745.12594

Gourlet-Fleury, S., Guehl, J.-M., & Laroussine, O. (2004). *Ecology and management of*
a neotropical rainforest : lessons drawn from Paracou, a long-term experimental
research site in French Guiana Ecology and management of a neotropical
rainforest : lessons drawn from Paracou, a long-term experimental research sit.
Paris: Elsevier. Retrieved from http://agritrop.cirad.fr/522004/

Herault, B., Ouallet, J., Blanc, L., Wagner, F., & Baraloto, C. (2010). Growth responses
of neotropical trees to logging gaps. *Journal of Applied Ecology*, *47*(4), 821–831.
https://doi.org/10.1111/j.1365-2664.2010.01826.x

Hérault, B., Bachelot, B., Poorter, L., Rossi, V., Bongers, F., Chave, J., … Baraloto, C.
(2011). Functional traits shape ontogenetic growth trajectories of rain forest tree
species: Plant traits shape growth trajectory. *Journal of Ecology*, *99*(6), 1431–
1440. https://doi.org/10.1111/j.1365-2745.2011.01883.x

Hérault, B., & Piponiot, C. (2018). Key drivers of ecosystem recovery after disturbance
in a neotropical forest. *Forest Ecosystems*, *5*(1), 2. https://doi.org/10.1186/s40663-
017-0126-7

John, R., Dalling, J. W., Harms, K. E., Yavitt, J. B., Stallard, R. F., Mirabello, M., …
Foster, R. B. (2007). Soil nutrients influence spatial distributions of tropical tree
species. *Proceedings of the National Academy of Sciences of the United States of*
America, *104*(3), 864–9. https://doi.org/10.1073/pnas.0604666104

Keck, F., Rimet, F., Bouchez, A., & Franc, A. (2016). phylosignal: an R package to
measure, test, and explore the phylogenetic signal. *Ecology and Evolution*, *6*(9),
2774–2780. https://doi.org/10.1002/ece3.2051

King, D. A., Davies, S. J., Nur Supardi, M. N., & Tan, S. (2005). Tree growth is related
to light interception and wood density in two mixed dipterocarp forests of Malaysia.

Functional Ecology, *19*(3), 445–453. https://doi.org/10.1111/j.1365-
2435.2005.00982.x

Körner, C. (2017). A matter of tree longevity. *Science*, *355*(6321), 130–131.
https://doi.org/10.1126/science.aal2449

Kupers, S. J., Engelbrecht, B. M. J., Hernández, A., Wright, S. J., Wirth, C., & Rüger, N.
(2019). Growth responses to soil water potential indirectly shape local species
distributions of tropical forest seedlings. *Journal of Ecology*, *107*(2), 860–874.
https://doi.org/10.1111/1365-2745.13096

Le Bec, J, Courbaud, B., Moguédec, G. Le, & Pélissier, R. (2015). Characterizing
tropical tree species growth strategies: Learning from inter-individual variability and
scale invariance. *PLoS ONE*, *10*(3), e0117028.
https://doi.org/10.1371/journal.pone.0117028

Lewis, S. L., & Tanner, E. V. J. (2000). Effects of above- and belowground competition
on growth and survival of rain forest tree seedlings. *Ecology*, *81*(9), 2525–2538.
https://doi.org/10.1890/0012-9658(2000)081[2525:EOAABC]2.0.CO;2

Lichstein, J. W., Dushoff, J., Levin, S. A., & Pacala, S. W. (2007). Intraspecific Variation
and Species Coexistence. *The American Naturalist*, *170*(6), 807–818.
https://doi.org/10.1086/522937

Martinez-Ramos, M., Alvarez-Buylla, E., & Sarukhan, J. (1989). Tree Demography and
Gap Dynamics in a Tropical Rain Forest. *Ecology*, *70*(3), 555–558.
https://doi.org/10.2307/1940203

Molino, J.-F., & Sabatier, D. (2001). Tree Diversity in Tropical Rain Forests: A
Validation of the Intermediate Disturbance Hypothesis. *Science*, *294*(5547), 1702–
1704. https://doi.org/10.1126/science.1060284

Nakagawa, S., & Schielzeth, H. (2013). A general and simple method for obtaining R2
from generalized linear mixed-effects models. *Methods in Ecology and Evolution*,
4(2), 133–142. https://doi.org/10.1111/j.2041-210x.2012.00261.x

O'Brien, M. J., & Escudero, A. (2021). Topography in tropical forests enhances growth
and survival differences within and among species via water availability and biotic
interactions. *Functional Ecology*, 1–13. https://doi.org/10.1111/1365-2435.13977

Osazuwa-Peters, O. L., Wright, S. J., & Zanne, A. E. (2017). Linking wood traits to vital
rates in tropical rainforest trees: Insights from comparing sapling and adult wood.
American Journal of Botany, *104*(10), 1464–1473.
https://doi.org/10.3732/ajb.1700242

Paquette, A., Joly, S., & Messier, C. (2015). Explaining forest productivity using tree
functional traits and phylogenetic information: Two sides of the same coin over
evolutionary scale? *Ecology and Evolution, 5*(9), 1774–1783.
https://doi.org/10.1002/ece3.1456

Peña-Claros, M., Fredericksen, T. S., Alarcón, A., Blate, G. M., Choque, U., Leaño, C.,
… Putz, F. E. (2008). Beyond reduced-impact logging: Silvicultural treatments to
increase growth rates of tropical trees. *Forest Ecology and Management*, *256*(7),
1458–1467. https://doi.org/10.1016/j.foreco.2007.11.013

Philipson, C. D., Dent, D. H., O'Brien, M. J., Chamagne, J., Dzulkifli, D., Nilus, R., …
Hector, A. (2014). A trait-based trade-off between growth and mortality: Evidence
from 15 tropical tree species using size-specific relative growth rates. *Ecology and*
Evolution, *4*(18), 3675–3688. https://doi.org/10.1002/ece3.1186

Poorter, L., McDonald, I., Alarcón, A., Fichtler, E., Licona, J. C., Peña-Claros, M., …
Sass-Klaassen, U. (2010). The importance of wood traits and hydraulic
conductance for the performance and life history strategies of 42 rainforest tree
species. *New Phytologist*, *185*(2), 481–492. https://doi.org/10.1111/j.1469-
8137.2009.03092.x

Poorter, L., Castilho, C. V., Schietti, J., Oliveira, R. S., & Costa, F. R. C. (2018). Can
traits predict individual growth performance? A test in a hyperdiverse tropical
forest. *New Phytologist*, *219*(1), 109–121. https://doi.org/10.1111/nph.15206

Potvin, C., & Gotelli, N. J. (2008). Biodiversity enhances individual performance but
does not affect survivorship in tropical trees. *Ecology Letters*, *11*(3), 217–223.
https://doi.org/10.1111/j.1461-0248.2007.01148.x

R Core Team. (2020). R: A Language and Environment for Statistical Computing.
Vienna, Austria.

Rees, M., Osborne, C. P., Woodward, F. I., Hulme, S. P., Turnbull, L. A., & Taylor, S. H.
(2010). Partitioning the components of relative growth rate: How important is plant
size variation? *American Naturalist*, *176*(6). https://doi.org/10.1086/657037

Reich, P. B. (2014). The world-wide "fast–slow" plant economics spectrum: a traits
manifesto. *Journal of Ecology*, *102*(2), 275–301. https://doi.org/10.1111/1365-
2745.12211

Rüger, N., Berger, U., Hubbell, S. P., Vieilledent, G., & Condit, R. (2011). Growth
strategies of tropical tree species: Disentangling light and size effects. *PLoS ONE*,
6(9), e25330. https://doi.org/10.1371/journal.pone.0025330

Rüger, N., Condit, R., Dent, D. H., DeWalt, S. J., Hubbell, S. P., Lichstein, J. W., …
Farrior, C. E. (2020). Demographic trade-offs predict tropical forest dynamics.
Science, *368*(6487), 165–168. https://doi.org/10.1126/science.aaz4797

Russo, S. E., Brown, P., Tan, S., & Davies, S. J. (2008). Interspecific demographic
trade-offs and soil-related habitat associations of tree species along resource
gradients. *Journal of Ecology*, *96*(1), 192–203. https://doi.org/10.1111/j.1365-
2745.2007.01330.x

Schmitt, S., Hérault, B., Ducouret, É., Baranger, A., Tysklind, N., Heuertz, M., …
Derroire, G. (2020). Topography consistently drives intra- and inter-specific leaf

trait variation within tree species complexes in a Neotropical forest. *Oikos*,
129(10), 1521–1530. https://doi.org/10.1111/oik.07488

Schmitt, S., Maréchaux, I., Chave, J., Fischer, F. J., Piponiot, C., Traissac, S., &
Hérault, B. (2020b). Functional diversity improves tropical forest resilience:
Insights from a long-term virtual experiment. *Journal of Ecology*, *108*(3), 831–843.
https://doi.org/10.1111/1365-2745.13320

Schmitt, S., Tysklind, N., Hérault, B., & Heuertz, M. (2021). Topography drives
microgeographic adaptations of closely related species in two tropical tree species
complexes. *Molecular Ecology*, mec.16116. https://doi.org/10.1111/mec.16116

Schmitt, S., Tysklind, N., Derroire, G., Heuertz, M., & Hérault, B. (2021b). Topography
shapes the local coexistence of tree species within species complexes of
Neotropical forests. *Oecologia*, *196*(2), 389–398. https://doi.org/10.1007/s00442-
021-04939-2

Schmitt, S., Tysklind, N., Heuertz, M., & Hérault, B. (2022). Selection in space and time:
Individual tree growth is adapted to tropical forest gap dynamics. *Molecular
Ecology*, 1–11. https://doi.org/10.1111/mec.16392

Stan Development Team. (2018). Rstan: the R interface to Stan. Retrieved from
http://mc-stan.org/

Stump, S. M., Song, C., Saavedra, S., Levine, J. M., & Vasseur, D. A. (2022).
Synthesizing the effects of individual-level variation on coexistence. *Ecological
Monographs*, *92*(1), e01493. https://doi.org/10.1002/ECM.1493

Svenning, J. C., Eiserhardt, W. L., Normand, S., Ordonez, A., & Sandel, B. (2015). The
Influence of Paleoclimate on Present-Day Patterns in Biodiversity and
Ecosystems. *Annual Review of Ecology, Evolution, and Systematics, 46*, 551–572.
https://doi.org/10.1146/annurev-ecolsys-112414-054314

Swenson, N. G. (2013). The assembly of tropical tree communities - the advances and
shortcomings of phylogenetic and functional trait analyses. *Ecography, 36*(3), 264–
276. https://doi.org/10.1111/j.1600-0587.2012.00121.x

Swenson, N. G. (2014). Phylogenetic imputation of plant functional trait databases.
Ecography, 37(2), 105–110. https://doi.org/10.1111/j.1600-0587.2013.00528.x

Tucker, C. M., Davies, T. J., Cadotte, M. W., & Pearse, W. D. (2018). On the
relationship between phylogenetic diversity and trait diversity. *Ecology, 99*(6),
1473–1479. https://doi.org/10.1002/ecy.2349

Umaña, M. N., Zhang, C., Cao, M., Lin, L., & Swenson, N. G. (2015). Commonness,
rarity, and intraspecific variation in traits and performance in tropical tree
seedlings. *Ecology Letters*, *18*(12), 1329–1337. https://doi.org/10.1111/ele.12527

Umaña, M. N., Zipkin, E. F., Zhang, C., Cao, M., Lin, L., & Swenson, N. G. (2018).
Individual-level trait variation and negative density dependence affect growth in

787 tropical tree seedlings. *Journal of Ecology*, *106*(6), 2446–2455.
788  https://doi.org/10.1111/1365-2745.13001

789 Uriarte, M., Canham, C. D., Thompson, J., & Zimmerman, J. K. (2004). A neighborhood
790  analysis of tree growth and survival in a hurricane-driven tropical forest. *Ecological*
791  *Monographs*, *74*(4), 591–614. https://doi.org/10.1890/03-4031

792 Uriarte, M., Condit, R., Canham, C. D., & Hubbell, S. P. (2004). A spatially explicit
793  model of sapling growth in a tropical forest: Does the identity of neighbours
794  matter? *Journal of Ecology*, *92*(2), 348–360. https://doi.org/10.1111/j.0022-
795  0477.2004.00867.x

796 Vehtari, A., Gelman, A., & Gabry, J. (2017). Practical Bayesian model evaluation using
797  leave-one-out cross-validation and WAIC. *Statistics and Computing*, *27*(5), 1413–
798  1432. https://doi.org/10.1007/s11222-016-9696-4

799 Violle, C., Navas, M.-L., Vile, D., Kazakou, E., Fortunel, C., Hummel, I., & Garnier, E.
800  (2007). Let the concept of trait be functional! *Oikos*, *116*(5), 882–892.
801  https://doi.org/10.1111/j.0030-1299.2007.15559.x

802 Visser, M. D., Bruijning, M., Wright, S. J., Muller-Landau, H. C., Jongejans, E., Comita,
803  L. S., & de Kroon, H. (2016). Functional traits as predictors of vital rates across the
804  life cycle of tropical trees. *Functional Ecology*, *30*(2), 168–180.
805  https://doi.org/10.1111/1365-2435.12621

806 Vleminckx, J., Fortunel, C., Valverde-Barrantes, O., Timothy Paine, C. E., Engel, J.,
807  Petronelli, P., … Baraloto, C. (2021). Resolving whole-plant economics from leaf,
808  stem and root traits of 1467 Amazonian tree species. *Oikos*, *130*(7), 1193–1208.
809  https://doi.org/10.1111/oik.08284

810 Wagner, F., Rossi, V., Stahl, C., Bonal, D., & Hérault, B. (2012). Water availability is the
811  main climate driver of neotropical tree growth. *PLoS ONE*, *7*(4), e34074.
812  https://doi.org/10.1371/journal.pone.0034074

813 Wickham, H., Averick, M., Bryan, J., Chang, W., McGowan, L., François, R., … Yutani,
814  H. (2019). Welcome to the Tidyverse. *Journal of Open Source Software*, *4*(43),
815  1686. https://doi.org/10.21105/joss.01686

816 Wright, S. J., Kitajima, K., Kraft, N. J. B., Reich, P. B., Wright, I. J., Bunker, D. E., …
817  Zanne, A. E. (2010). Functional traits and the growth-mortality trade-off in tropical
818  trees. *Ecology*, *91*(12), 3664–3674. https://doi.org/10.1890/09-2335.1

819 Xiang, W., Wu, W., Tong, J., Deng, X., Tian, D., Zhang, L., … Peng, C. (2013).
820  Differences in fine root traits between early and late-successional tree species in a
821  Chinese subtropical forest. *Forestry*, *86*(3), 343–351.
822  https://doi.org/10.1093/forestry/cpt003

823 Yang, J., Cao, M., & Swenson, N. G. (2018). Why Functional Traits Do Not Predict Tree
824  Demographic Rates. *Trends in Ecology and Evolution*, *33*(5), 326–336.
825  https://doi.org/10.1016/j.tree.2018.03.003

826   Yu, G. (2020). Using ggtree to Visualize Data on Tree-Like Structures. *Current*
827   *Protocols in Bioinformatics*, *69*(1), 1–18. https://doi.org/10.1002/cpbi.96