Subtilases turn on Pectin Methylesterase activity for a robust apoplastic immunity against

pathogens

Daniele Coculo[1], Daniele Del Corpo[1], Miguel Ozáez Martínez[2], Pablo Vera[2], Gabriella Piro[3], Monica

De Caroli[3] and Vincenzo Lionetti[1, 4]*.

* Address correspondence to Vincenzo Lionetti (vincenzo.lionetti@uniroma1.it)

Keywords: Pectin Methylesterases, Subtilases, Plant Immunity, *Botrytis cinerea*, *Arabidopsis thaliana*,

Cell Wall Integrity

1 Dipartimento di Biologia e Biotecnologie Charles Darwin, Sapienza Università di Roma, Rome, Italy

2 Instituto de Biologia Molecular y Celular de Plantas, Universidad Politecnica de Valencia-C.S.I.C,

Ciudad Politecnica de la Innovacion, Valencia, Spain.

3 Dipartimento di Scienze e Tecnologie Biologiche e Ambientali, Università del Salento, Lecce, 73100

Italy

4 CIABC, Sapienza Università di Roma, Rome, Italy

Running title: SBTs trigger PME activity in plant immunity

Abstract

Plants involve a fine modulation of pectin methylesterase (PME) activity against microbes. PME activity can promote the cell wall stiffening and the production of damage signals able to induce defense responses. However, to date, the knowledge about the molecular mechanisms triggering PME activity during disease remains largely unknown. In this study, we explored the role of subtilases (SBTs), serine proteases consisting of 56 isoforms in *Arabidopsis thaliana,* as activators of PME activity in plant immunity. By using biochemical and reverse genetic approaches, we found that SBT3.3 and SBT3.5 are required to control PME activity and resistance to the fungus *Botrytis cinerea. Arabidopsis sbt3.3 and sbt3.5* knockout mutants showed a reduced induction of PME activity and an increased susceptibility to *B. cinerea*. The SBT3.3 overexpression overactivates PME activity only during fungal infection resulting in an increased expression of the defence-related genes and in an enhanced resistance to *B. cinerea*. We revealed that SBT3.3 and the Pro-PME17 isoforms are both secreted in the cell wall using a conventional protein secretion pathway and their colocalization in the apoplast increase over time. Our findings point to SBTs as a mechanism to switch on pectin integrity signaling and to reinforce plant immunity at the right time to avoid the growth-defense trade-off.

Introduction

Plants continuously perceive attacks by microorganisms activating a timely and effective immune response. Part of this recognition is exerted by plasma membrane Pattern Recognition Receptors (PRRs), which can detect Microbe-, Pathogen-, or endogenous Damage associated molecular patterns (MAMPs, PAMPs or DAMPs)(Zhou and Zhang, 2020). Pattern-triggered immunity (PTI) constitutes the first line of plant inducible immunity that restricts pathogen proliferation. Since the immunity efforts often result in a growth defence trade-offs, a precise regulation of immune pathways becomes critical to optimize fitness (Lionetti et al., 2010; Huot et al., 2014; Ha et al., 2021). A pivotal role in this homeostasis is played by proteolysis mediated by proteases, widely implicated in plant stresses, although their action mechanisms remain largely unknown (Schaller, 2004; Hou et al., 2018). In particular, Subtilases (SBTs, Pfam00082), serine peptidases belong to family S8, (MEROPS database; https://www.ebi.ac.uk/merops/cgi-bin/famsum?family=S8) plays key roles in defense responses against microbes, plant parasitism and adaptation to the environmental changes (Figueiredo et al., 2018; Schaller et al., 2018; Ogawa et al., 2021). In *A. thaliana,* 56 SBT isoforms are divided into 6 subfamilies: SBT1, SBT2, SBT3, SBT4, SBT5, SBT6 (Rautengarten et al., 2005).

Specific SBTs are targeted to the plant Cell Wall (CW), the foremost interface at which interactions between plants and microbes take place. In this compartment, SBTs can process larger precursor zymogens (Pro-Proteins), that require specific activation processes for enzyme maturation (Khan and James, 1998). SBT6.1 (S1P) can process the pro-peptide form of RAPID ALKALINIZATION FACTOR 23 (RALF23) modulating immune response against *Pseudomonas syringae* (Stegmann et al., 2017). In tomato, specific SBTs contributes to Jasmonic Acid-mediated resistance against *Manduca sexta* larvae, by processing the wound pro-hormone Prosystemin, releasing the bioactive peptide called Systemin (Meyer et al., 2016). The six genes of the tomato cluster SBT P69 (P69A–P69F) express proteins particularly involved in plant defence. P69B can cleave the *Phytophthora infestans* PC2, a cysteine-rich secreted apoplastic protein releasing immunogenic peptides that activates host immunity (Wang et al., 2021). In particular, P69C processes an extracellular matrix-associated Leucine-Rich Repeat protein (LRP), the first physiological target of an extracellular SBT identified in plants (Tornero et al., 1996). SBT3.3, the ortholog of tomato P69C in *Arabidopsis*, was discovered as regulator of *Arabidopsis* primed immunity (Ramirez et al., 2013). Although SBT3.3 was postulated to cleave an extracellular larger protein, the identification of its specific substrate and the mechanism by which SBT3.3 can trigger immunity remains to be identified.

During plant development, specific SBTs can process Pectin Methylesterases (PME; Pfam01095), enzymes regulating the methylesterification of Homogalacturonan (HG), the major constituent of pectin in the CW (Wormit and Usadel, 2018; Coculo and Lionetti, 2022). PMEs catalyse the hydrolysis of methylester bonds at C-6 of Galacturonic Acid (Gal A) residues, producing acidic pectins with negatively charged carboxyl groups, releasing methanol (MeOH) and protons into the apoplast. In *Arabidopsis*, 66 PME isoforms have been identified which can be classified into 2 groups on the basis of their structures (Pelloux et al., 2007; Dedeurwaerder et al., 2008). Group 1 comprises 21 PME isoforms, containing only the catalytic domain. The 45 PME isoforms of Group 2 (hereafter named Pro-PMEs) are organized as a polycistronic messenger RNA which resembles an operon-like gene cluster (Coculo and Lionetti, 2022). In Pro-PMEs, the PME catalytic domain is preceded by an N-terminal region (also referred as Pro region or PMEI-like domain) which shares structural similarities with functionally characterized PME Inhibitors (PMEIs) and it can acts as auto-inhibitor of PME activity (Bosch et al., 2005; Wolf et al., 2009; Del Corpo et al., 2020). This domain could impede a premature pectin de-methylesterification, during specific phases of their secretion, assembly, or modification contributing to a correct pectin functionality.

A local and strong induction of PME activity is exerted in *Arabidopsis thaliana* and other plants, when challenged with different microbial pathogens (Bethke et al., 2014; Lionetti, 2015; Lionetti et al., 2017). *Pro-PME17* isoform is strongly upregulated in response to several pathogens, defense hormones, elicitors or abiotic stress and its activity strongly contribute to the induction of defence-related PME activity (Bethke et al., 2014; Del Corpo et al., 2020). A finely tuned PME activity can play multiple functions in plant immunity. PMEs can produces de-methylesterified negatively charged carboxyl groups that can form consecutive calcium bonds with other HG molecules and thus strengthen the CW through the formation of "egg-boxes" structures (Senechal et al., 2014b; Del Corpo et al., 2020). Plant PME activity can also favour DAMP related immunity. In particular, the PME activity could favour the release and perception of de-methylesterified oligogalacturonides (OGs), the well characterized DAMP, locally released in cooperation with endo-polygalacturonases (PGs), and able to trigger plant immunity (Osorio et al., 2008; Osorio et al., 2011; Ferrari et al., 2013; Kohorn et al., 2014). OGs are perceived in *Arabidopsis* by the pathogen recognition receptor WALL-ASSOCIATED KINASE1 (WAK1) to activate plant immune responses (Brutus et al., 2010; Kohorn and Kohorn, 2012). WAK2 requires PME-mediated pectin de-methylesterification to activate OG-dependent stress responses (Kohorn et al., 2014). Interestingly, WAK1, WAK2 and another PRR,

FERONIA (FER), preferentially bind to de-methylesterified pectins (Decreux and Messiaen, 2005;

Kohorn et al., 2014; Feng et al., 2018; Guo et al., 2018; Lin et al., 2021). The de-methylesterification of

pectin by PMEs represents the main mechanism to generate plant-derived methanol (MeOH), a DAMP-

like alarm signal (Hann et al., 2014). OGs and MeOH are able to trigger defensive priming effect

setting the stage for the intra and inter-plant immunity (Dorokhov et al., 2012; Komarova et al., 2014;

Gamir et al., 2021; Giovannoni et al., 2021).

Precise switch on/off mechanisms regulating PME related immunity are required because they must be

timely triggered for an efficient immune response and they cannot be persistent given the well-known

defense growth trade-off (He et al., 2022). The need of a post-transcriptional control of PME activity

by specific PME inhibitors (PMEIs) during plant invasion by microbes was previously demonstrated

(An et al., 2008; Lionetti et al., 2017; Liu et al., 2018). The involvement of SBTs as posttranscriptional

processing of Pro-PMEs and regulation of PME activity in plant immunity was still not investigated.

Here, we found that all the PME isoforms induced in *A. thaliana* against the necrotrophic fungus

Botrytis cinerea possess a Pro-region. Exploiting reverse genetic approaches and biochemistry, we

demonstrated that SBT3.3 and SBT3.5 contribute to the induction of PME activity in *Arabidopsis*

immunity against *Botrytis*. Notably, we found that SBT3.3 processes defence-related Pro-PMEs as

substrate and trigger specific defence related genes. SBT3.3 follows a conventional secretion pathway

to reach the apoplast where colocalizes with Pro-PME17. Our results provide strong evidence that the

processing of Pro-PMEs by SBTs represent a mechanism to switch on pectin integrity signaling only in

plant immunity, most likely, to avoid a growth-defense trade-offs.

RESULTS

The processing of Pro-PME zymogens is required in *Arabidopsis-Botrytis* interaction

PMEPCRA, *PME3, PME17*, *PME20*, *PME21*, and *PME41* are the PME isoforms exhibiting a

significant altered timing and level of expression in *A. thaliana - B. cinerea* interaction and in several

other pathosystems (Lionetti et al., 2012; Bethke et al., 2014; Lionetti et al., 2017). Interestingly, after a

careful comparison of the sequences we realized that all these isoforms are zymogens, expressed as

Pro-PME inactive precursors (Figure S1). This strongly suggest that a proteolytic cleavage, exerted by

specific proteases, is requested for PME activation in *Arabidopsis* immunity against *Botrytis*. While the

PME catalytic domain is highly conserved, we intriguingly noticed here a variability in the processing

motif between the different PME isoforms. Pro-PMEPCRA, Pro-PME17 and Pro-PME21 showed a

RKLL processing motif, while the RRKL motif characterized the Pro-PME20 and Pro-PME41 isoforms. Instead, two processing motifs, RKLK and RRLL were present in the Pro-PME3. Other motifs with the consensus sequence FPSW and LPLKMTERARAV are partially conserved between the isoforms. All these observations could imply the involvement of multiple proteases-PME pairs.

SBT3.3 and SBT3.5 are induced in *Arabidopsis* during *B. cinerea* as part of PTI

We explored the possibility that SBTs are involved in the processing and activation of Pro-PMEs against microbes. Exploiting publicly available microarray data, we selected two genes, *SBT3.3* and *SBT3.5* showing a significative induction during *B. cinerea* infection (Figure S2A). Interestingly, both genes are co-expressed with *Pro-PME17,* a functional PME involved in the resistance to *Botrytis cinerea* (Figure S2B) (Lionetti et al., 2012; Del Corpo et al., 2020). To confirm the transcriptomic analysis and to gain more insight in the kinetic of expression of these genes during the disease, the expression levels of *SBT3.3* and *SBT3.5* were quantified at different times in *B. cinerea*-infected and mock-inoculated *A. thaliana* wild type (WT) leaves by qPCR (Figure 1B and Figure S2C-D). *SBT3.3* is expressed only in infected tissues while *SBT3.5* shows a basal expression also in mock-inoculated leaves. *SBT3.3* and *SBT3.5* were significantly induced at 24 hpi and further increased at 48 hpi*,* when they showed about a significant 15- and 5-fold induction, respectively, in infected leaves compared to mock inoculated tissues. These data clearly reveal an involvement of *SBT3.3* and *SBT3.5* in *Arabidopsis-Botrytis* interaction. Interestingly, the expression of both *SBTs* is also triggered in *Arabidopsis* against several pathogens and by different bacterial and fungal elicitors of plant defence responses (Figure S3A). We next investigated whether pectin-related DAMPs can impact the expression of the selected SBTs. *Arabidopsis* WT seedlings were treated with OGs or water, and the expression of *SBT3.3* and *SBT3.5* was evaluated at 1 h post treatment. A 7-fold induction of *SBT3.3* was revealed after treatment with OGs (Figure S3A and B), while the expression of *SBT3.5* was not induced (Fig. S3A and S3B). All these results indicate that *SBT3.3* and *SBT3.5* are part of *Arabidopsis* immune responses against *B. cinerea* and several pathogens.

SBT3.3 and SBT3.5 contribute to *Arabidopsis* PME activation and resistance to *B. cinerea*

To explore on the possibility that SBT3.3 and SBT3.5 may be involved in the activation of Pro-PMEs in plant immunity we exploited a reverse genetic approach. Two T-DNA insertional mutants for the selected genes were isolated (Figure 1A). The insertions were localized in the fifth and ninth exons, respectively, for *sbt3.3-1* (SALK_086092) and *sbt3.3-2* (SALK_107470.45.45) mutants and in the first

exon and in the second intron, respectively for *sbt3.5-1* (SAIL_400_F09) and *sbt3.5-2* (GABI_672C08) mutants. The level of *SBT3.3* and *SBT3.5* expression was compared in leaves of mutants and WT plants after *B. cinerea* and mock inoculation by qPCR (Figure 1B). All mutants fail to induce gene expression in fungus-challenged leaves, resulting in *SBT3.3* and *SBT3.5* KO lines. No significant defects were observed in the rosette growth and seed germination in *sbts* mutants which showed the growth parameters like WT plants (Figure 1C and S4A-B). To understand if the lack of SBTs expression can influence pectin composition we compared the monosaccharide composition of matrix polysaccharides extracted from rosette leaves of SBTs mutants and WT plants. All mutants analysed showed a molar percentage of monosaccharides not significantly different from that observed in WT plants (Figure 1D).

After establishing that the two subtilases are involved in *Arabidopsis-Botrytis* interaction we evaluated their possible contribution in the control of PME activity during fungal infection. The level of PME activity was quantified in leaves of WT, *sbt3.3-1*, *sbt3.3-2, sbt3.5-1* and *sbt3.5-2* plants at 48 hours post *Botrytis* and mock inoculation using histochemical and biochemical assays based on Ruthenium Red (RR), a cationic dye with six positive charges able to bonds with the acidic groups of HG (Lionetti, 2015)(Figure 2A). A significant lower induction of PME activity was observed in infected *sbt3.3-1*, *sbt3.3-2*, *sbt3.5-1* and *sbt3.5-2* mutants (respectively of 3.3-, 2-, 1.4- and 1.3-fold reduction) compared to WT (Figure 2B and C). These results unveil that Pro-PMEs are substrates for SBT3.3 and indicate that SBT3.3 and SBT3.5 contribute to control the PME activity in *Arabidopsis* immunity to *Botrytis.* Interestingly, the significant greater effect found in SBT3.3 respect to SBT3.5 mutants suggest that the two proteases can process different Pro-PMEs isoforms during disease. No significant differences in PME activity were observed between mock-inoculated WT plants and *sbt* mutants. This was expected for SBT3.3 mutants, as SBT3.3 is not basally expressed in leaves (Figure 1B). In *sbt3.5* mutants, the lack of the basal SBT3.5 expression observed mock-inoculated leaves does not significantly affect neither the total PME activity (Figure 2C) nor the development of the leaves (Figure 1C and 2C). A possible compensation effect for the loss of SBT3.5 expression in leaves of *sbt3.5* mutants can be exerted based on activation of other PME isoforms untargeted by SBT3.5. Therefore, we evaluated the susceptibility of *sbt3.3* and *sbt3.5* mutants to *B. cinerea* (Figure 3A and B). The local symptoms of the fungus were significantly higher in both *sbt3.3* and *sbt3.5* mutants respect to WT. In particular, the lesions produced by the fungus were larger in *sbt3.3-1*, *sbt3.3-2, sbt3.5-1*, *sbt3.5-2* mutants (respectively, of 2.2-, 2-, 1.6- and 1.7-fold) respect to WT. A correlation between the level of resistance and the amount of PME activity expressed in the tissues were observed between the different genotypes

analysed (Figure 2 and 3). A greater development of *Botrytis* mycelium was detected in *sbt* mutants

respect to WT plants using trypan blue staining (Figure 3B). All our results indicate that, although at

different extent, both SBTs contribute to trigger PME activity and the resistance to *Botrytis*.

SBT3.3 overexpression over activates defence related PME activity restricting *Botrytis* infection

Given its greater effect in activating PME, we deepen our study on SBT3.3 exploring the effect of its

overexpression in *Arabidopsis* on PME activity and *Botrytis* resistance. Two transgenic *Arabidopsis*

lines, named *SBT3.3-OE1* and *SBT3.3-OE2* exhibiting high level of SBT3.3 expression in leaves were

characterized (Figure 4A). Interestingly, a 40- and 20-fold overexpression was revealed, respectively,

in leaves of *SBT3.3-OE1* and *SBT3.3-OE2* lines respect to WT leaves when infected with *B. cinerea*.

No growth abnormalities were observed in SBT3.3 overexpressing plants (Figure 4B and S4). The CW

monosaccharide composition in rosette leaves of SBT3.3-OE lines is not significantly different from

that quantified in the WT plants (Figure 4C). These results are consistent with those obtained with the

mutants and corroborate the conclusion that the expression of SBT3.3 is dedicated to the Arabidopsis

immune response. The level of PME activity was quantified in leaves of WT, *SBT3.3OE-1 and*

SBT3.3OE-2 plants at 48 hours post *Botrytis-* and mock- inoculation (Figure 5A and B). A significant

higher induction of defence-related PME activity was observed in the infected SBT3.3 overexpressing

lines compared to control. SBT3.3OE-1 and SBT3.3OE-2 lines showed, respectively, a 1.7- and 1.5-

fold increase of PME activity in the *B. cinerea* infected leaves compared to WT. Despite the

overexpression of SBT3.3 in transgenic mock-inoculated plants (Figure 4A), the level of PME activity

were not significantly different from that measured in WT plants (Figure 5A). This result highlights

that SBT3.3 trigger PME activity exclusively in *Arabidopsis* defense against the fungus. The levels of

methylesters remained in plant cell wall after 48h of *B. cinerea* inoculation were compared between

SBT3.3-OE lines and WT plants. A significant lower amount of methylesters was revealed in the CW

extracted from *B. cinerea* infected leaves of *SBT3-3-OE1* and *SBT3.3-OE2* lines respect the WT plants

(Figure 5C). This result corroborates that SBT3.3 process Pro-PMEs in plant defence and that a higher

PME activity was released by SBT3.3 overexpression in transgenic challenged with the fungus respect

the control. Next, we evaluated the susceptibility of *SBT3.3-OE* lines to *B. cinerea*. The local

symptoms of the fungus were significantly lower in both *SBT3.3-OE* plants when compared with the

WT (Figure 5D and E). In particular, the lesion area produced by the fungus in *SBT3.3-OE1* and

SBT3.3-OE2 mutants are 2.1- and 1.9-fold lower compared to WT. A minor development of *Botrytis*

mycelium was detected in SBT3.3-OE1 mutants respect to WT plants using trypan blue staining

(Figure 5E). All these results support an important role of SBT3.3 as trigger of PME activity in
Arabidopsis immunity to Botrytis and indicate that the SBT3.3 overexpression confers to *Arabidopsis* a
higher resistance to *B. cinerea* respect to WT.

SBT3.3* overexpression enhances immune responses against *Botrytis
The next goal was to verify whether the improved resistance observed in the SBT3.3 overexpressing
lines could be linked to a more robust immune response exerted by the expression of specific defence
genes. The induction of an array of defence related genes important for *Arabidopsis* defence against
Botrytis was compared between WT and *SBT3.3-OE1* plants inoculated with Mock and *Botrytis*.
WRKY33 is a key transcriptional regulator of hormonal and metabolic responses toward *B. cinerea*
infection (Birkenbihl et al., 2012). WRKY33 can targets the promoter of PHYTOALEXIN
DEFICIENT3 (PAD3) encoding an enzyme required for the synthesis of antimicrobial camalexin
(Zhou et al., 1999). The cytochrome P450 enzyme CYP81F2 is essential for the pathogen-induced
accumulation of 4-methoxy-indol-3-ylmethyl glucosinolate (Clay et al., 2009; AbuQamar et al., 2017))
WAK2 is a wall associated kinase mediating pectin related immunity and activated by de-methyl
esterified pectin to trigger defence responses (Kohorn et al., 2014). PME17 and PMEI10 are two
regulators of PME activity in plant immunity to *B.cinerea* (Lionetti et al., 2017; Del Corpo et al.,
2020). Interestingly, a higher induction of WRKY33, PAD3, and WAK2 were observed in *SBT3.3-OE1*
plants respect the WT only when challenged with the fungus (Figure 6). CYP81F2 expression is also
overexpressed in transgenic plants considering the low level of *B.cinerea β-tubulin2* revealed. The
expression of Pro-PME17 and PMEI10 follows the extent of fungal infection as revealed by *B.cinerea*
β-tubulin2, indicating that the induction of their gene expression is independent from SBT3.3 activity.
These data indicate that SBT3.3 contribute to enhance Arabidopsis immunity, most likely, improving
the production of antimicrobial compounds and reinforcing cell wall integrity surveillance.
We next explored if SBTs expression can influence Hydrogen Peroxide production, an important
defence response to *Botrytis* infestation (Pogorelko et al., 2013; Siegmund and Viefhues, 2016). Leaves
of WT, *sbt3.3-1*, *sbt3.5-1* and *SBT3.3-OE1* plants, inoculated with Mock or with *B. cinerea*, were
stained with 3,3′-diaminobenzidine, a chromogen able to reveal the H_2O_2 accumulation in the tissue
(Reem et al., 2016). No accumulation of H_2O_2 was observed in mock-inoculated leaves of all
genotypes analyzed and a similar DAB staining was revealed in all the leaves challenged with *Botrytis*.
These results indicate that H_2O_2 production are not dependent from SBTs mediated signaling (Figure
S5).

SBT3.3 and Pro-PME17 are delivered through a conventional protein secretion pathway and colocalize in the apoplast

SBT3.3 was previously demonstrated to be an extracellular subtilase, but its secretion pathway was never explored (Ramirez et al., 2013). Moreover, Pro-PME17 was predicted as an apoplastic zymogen but its localization was never demonstrated. With the aim to deepen our knowledge on the secretion patterns and the final subcellular destination of the two proteins, a fluorescent variant of SBT3.3 and of Pro-PME17 were constructed by fusing the respective encoding cDNA to *yfp* and *gfp,* under the control of *35S* promoters. The fluorescent tag was linked to the C-terminus, as none of the two protein is predicted to have the ω site of the glycosylphosphatidylinositol (GPI) anchor addition (predGPI software; http://gpcr2.biocomp.unibo.it/predgpi/) and previously found for AtPMEI1 (De Caroli et al., 2011). *SBT3.3-YFP* and *Pro-PME17-GFP* constructs, were transiently expressed in the epidermal cells of tobacco leaves and observed by laser scanning confocal microscopy (Figure 7). The fluorescence signals relative to both proteins, labelled the ER and the Golgi stacks as revealed by colocalization experiments using ER and Golgi markers *RFP-HDEL* and *ST52-mCherry* (Nelson et al., 2007) (Figure 7 A-F and K-P). Both proteins accumulate in the apoplast as confirmed by co-localization with the CW marker XTH11-RFP (De Caroli et al., 2021) (Figure 7 G-I, Q-S) and plasmolysis of epidermal cells co-expressing both *pm-rk*, marker of plasma membrane (Nelson et al., 2007) and *SBT3.3-YFP* or *Pro-PME17-GFP* (Figure 7 J, T). Intriguingly, the PME17 reaches the apoplast before the SBT. At 48 h post transformation, the CW was fluorescently labelled by Pro-PME17-GFP and after 52 h by SBT3.3-YFP protein fusion (Figure 7, B and E). These results indicate that that SBT3.3 and Pro-PME17 are correctly secreted into the apoplast, with a different rate, using conventional secretion pathway. We explore on the compartment and timing of SBT3.3 and Pro-PME17 colocalization. A new version of *Pro-PME17* tagged with *rfp,* under the control of *35S* promoter was prepared and transiently co-expressed with *SBT3.3-YFP* in the epidermal cells of tobacco leaves. The Pro-PME17 and SBT3.3 co-localize at the CW, and this is favoured over time as revealed by colocalization parameters (Figure 8). All these results strongly suggest that Pro-PME17 is released in the CW in an unprocessed and inactive form where, later when required, is activated by SBT3.3.

DISCUSSION

Plants exploit a great and local increase of PME activity in response to pathogens with different lifestyles (Bethke et al., 2014; Lionetti et al., 2014; Lionetti, 2015; Lionetti et al., 2017). *A. thaliana*

induces the expression of specific PME isoforms during the attack of different pathogens. Among them, Pro-PME17 is induced after the infection of a wide range of plant pathogens and can be considered a general biomarker of pathogenesis (Lionetti et al., 2012). *Botrytis cinerea* is an important fungal plant necrotrophic pathogen, that kills plant tissue prior to feeding using a range of toxic molecules to destroy host cells (Williamson et al., 2007; Dean et al., 2012). Pro-PME17 was recently involved in triggering PME activity and in resistance against *B. cinerea* (Del Corpo et al., 2020). The Pro region acts as an intramolecular inhibitor of Pro-PME17 activity. Here, intriguingly we note that all the PME isoforms induced in *A. thaliana* during *B. cinerea* infection are organized as zymogens in which the catalytic region is preceded by an N-terminal Pro region. The structural similarity revealed between the amino acid sequences of the Pro regions of these PMEs and the independent immunity-related PMEIs (AtPMEI10 and AtPMEI11) (Lionetti et al., 2017; Del Corpo et al., 2020) support their possible role as intramolecular inhibitors of the specific PMEs in each specific Pro-PME cluster. We therefore attempted to identify which possible proteases could be involved in the activation of PME activity against microbes. Different SBTs were associated to the processing of PMEs in plant physiology (Schaller et al., 2018). Most SBTs are targeted to the CW, where they contribute to the control of CW properties during growth and development. The *Arabidopsis* SBT3.5 was previously found to process Pro-PME17 releasing PME activity during root growth in *Arabidopsis* seedlings (Senechal et al., 2014a). SBT1.7 modulates PME activity in the mucilage release from *Arabidopsis* seed coats (Rautengarten et al., 2008). AtSBT6.1 is likely to be involved in maturation of a pectin methylesterase (Wolf et al., 2009). The evidence that the precursors of bZIP transcription factors, RALF23 and GOLVEN1 are further substrates of AtSBT6.1 and that most SBTs are synthesized as inactive precursors could suggest that SBTs isoform could be part of an activation cascade, similar to that observed for caspases during plant programmed cell death events in mammals (Vartapetian et al., 2011; Paulus and Van der Hoorn, 2019).

Although SBTs were also associated to plant immunity their molecular substrates and their function remains largely unknowns (Figueiredo et al., 2014; Schaller et al., 2018; Godson and van der Hoorn, 2021). The tomato subtilisin-like protease SlSBT3 contributes to insect resistance in tomato. Although SlSBT3 was found to affect PME activity and the level of pectin methylesterification, these effects were unrelated to its role in herbivore defence (Meyer et al., 2016). Also, AtSBT5.2 and AtSBT3.14 were involved in different plant-pathogen interactions but their functions appear unrelated to PME activity (Lozano-Torres et al., 2014; Serrano et al., 2016). Here, we explored on the possible role of

SBTs as post-transcriptional regulators of PME activity in *Arabidopsis* immunity against the necrotrophic fungus *B. cinerea*. Exploiting publicly available microarray data, we selected two isoforms, SBT3.3 and SBT3.5 co-expressed with Pro-PME17 in *Arabidopsis* during *Botrytis* infection. SBT3.3 was previously discovered as a extracellular SBT3.3 involved in the activation of *Arabidopsis* immunity against *Pseudomonas syringae* and *Hyaloperonospora arabidopsidis* (Ramirez et al., 2013) although its substrate remained unknown. Here, we demonstrated that SBT3.3 and SBT3.5 are induced in *Arabidopsis* leaves challenged with *Botrytis*. SBT3.3 exhibited a higher fold-induction respect to SBT3.5. Interestingly, the expression of both SBTs is also triggered in *Arabidopsis* against several pathogens. Moreover, SBT3.3 and SBT3.5 are induced by different bacterial and fungal elicitors of plant defence responses. SBT3.3 is specifically dedicated to plant defence because it is induced only during the infection process. However previous evidence indicates that SBT3.3 expression respond very rapidly to H_2O_2 and it enhances the activation of mitogen-activated protein kinases (MAPKs) following microbial attack (Ramirez et al., 2013). We detected SBT3.3 as a component of OGs signaling. All these evidence points to the involvement of SBT3.5 and, particularly of SBT3.3, as contributors in the activation of the earliest signaling events in PTI against Botrytis and several other pathogens.

The possible role of SBT3.3 and SBT3.5 as PME activators in *Arabidopsis* immunity against *B. cinerea* infection was explored. *sbt3.3-1*, *sbt3.3-2, sbt3.5-1*, *sbt3.5-2* mutants showed a significant higher susceptibility to the fungus when compared with WT. The significant lower induction of PME activity observed in *B. cinerea*-infected *sbt3.3* and *sbt3.5* mutants respect to WT plants indicates that SBT3.3 and SBT3.5 are involved in the release of mature active PMEs to contrast the microbial invasion. The susceptibility- and PME-related effects are more marked in the *sbt3.3* mutant respect to *sbt3.5* mutants indicating that SBT3.3 can have a major contribution on the total PME activity expressed in *Arabidopsis* against the fungus. This also suggests that the two SBTs analyzed in this study can target different Pro-PME isoforms. The involvement of multiple SBT-Pro-PME pairs is also suggested by the observed variability in the processing motifs of all the PME isoforms involved in *Arabidopsis-Botrytis* interaction. We revealed a significant higher induction of PME activity and a higher resistance to *Botrytis* in infected SBT3.3-OE1 and SBT3.3-OE2 leaves respect to untransformed plants. Overall, our evidence indicates that SBT3.3 and SBT3.5, by processing different Pro-PME isoforms, contribute to trigger PME activity to contrast *B. cinerea* infection. The absence of growth

abnormalities and alteration of monosaccharide composition in the *sbt* mutants and SBT3.3OE lines indicate that these SBTs are mostly involved in plant immunity processes in adult plants.

Moreover, we firstly demonstrated that SBT3.3 process Pro-PMEs as extracellular substrate to reinforce plant immunity. This is also corroborated by the low level of CW methylesters detected in SBT3.3OE lines. By timely activating PMEs early at the point of fungal penetration, SBTs can favor the increase of charged carboxyl groups forming a higher amount of Ca^{2+}-mediated pectin crosslinks to reinforce the CW structure and the resistance to the microbe. The SBT3.3-related pectin reinforcement is also accompanied by the stimulation of an immune response signaling. The higher induction of *WRKY33*, *PAD3*, *CYP81F2* and *WAK2* genes detected in *SBT3.3-OE1* plants respect to WT plants identify a more robust immune response as mechanism contributing to improved resistance observed in the transgenic plants. We cannot exclude that the SBTs here analysed can also process other substrates and therefore that SBT-induced PME activity could also partially participate in the activation of these genes. The activation of *WRKY33*, *PAD3* and *CYP81F2* by SBT3.3 suggest the involvement of camalexin and indole glucosinolate accumulation. The evidence that WAK2 is highly expressed in transgenic plants suggest an improvement of cell wall surveillance (Kohorn et al., 2014). Interestingly, WAK2 was demonstrated to require PME-mediated pectin de-methylesterification to activate OGs-dependent stress responses. We can hypothesize that SBT3.3 activates PME activity to favour WAK2 binding, and WAK2-OGs mediated stress signalling. Considering that OGs induce the expression of SBT3.3, WRKY33, PAD3 and CYP81F2 and that a SBT3.3 self-activation was previously reported (Denoux et al., 2008; Ramirez et al., 2013), a positive feedback loop circuit based on PME activity, could be mounted to strengthen the activation of OG-mediated defence responses. In addition, SBTs-dependent PME activity could also favour the binding and the functions of other PRR, like WAK1 and FERONIA, known to preferentially bind de-methylesterified crosslinked pectins (Decreux and Messiaen, 2005; Feng et al., 2018; Guo et al., 2018; Lin et al., 2018). Future efforts will be needed to explore on this hypothesis. All these observations indicate SBTs and PME activity exploit a cooperation between CW integrity maintenance and PTI signalling to regulate defence responses after a CW damage (Engelsdorf et al., 2018). Although the specific pairs of SBT3.3, SBT3.5 and Pro-PMEs involved in plant immunity remain to be identified, we found that Pro-PME17 and SBT3.3 are both delivered in the apoplast using a conventional secretion pathway. Pro-PME17 reaches the cell wall before SBTs and their colocalization in the apoplast increase over time. This temporal separation could avoid a premature PME activation and pectin de-methylesterifiation in the secretory compartments. Later in the apoplast, plants can accumulate Pro-PMEs clusters as soluble and inactive receptors. At the

right time, SBTs could be delivered in an apoplast already enriched in the inactive Pro-PMEs to rapidly derepress the inhibitory activity exerted by the Pro region. This mechanism could lead to a faster induction of PME activity to amplify the early immune signalling preceding defence response activation. In a similar mechanism, known as ectodomain shedding, the proteolytic cleavage of extracellular domains of cell surface proteins resulting in the activation of a variety of normal and pathological signalling processes in animals including growth factor signalling, cell adhesion, inflammation and cell survival (Lichtenthaler et al., 2018). This proposed SBTs function would be also in line with the negative regulation that speeds the response times of hormonal transcription networks in plants (Rosenfeld et al., 2002). The speed of a response, particularly to biotic stress, can be crucial for the survival of the sessile plant. Interestingly, SBT3.3 was involved in the physiological state called "priming", a sensory status able to induce a resistance faster and stronger respect to the same stimulus received before (Ramirez et al., 2013). It was previously hypothesized that priming could involve the accumulation of inactive proteins that becomes operative to initiate a signal amplification leading to a faster and stronger activation of defence responses (Bruce et al., 2007; Conrath, 2011; Ramirez et al., 2013). The processing of Pro-PMEs by SBT3.3 at a precise time and in the correct subcellular compartment could be part of the immune priming processes. Two PME-related DAMPs, OGs and MeOH are known to trigger defensive priming in plant immunity (Dorokhov et al., 2012; Komarova et al., 2014; Gamir et al., 2021; Giovannoni et al., 2021). The possibility that SBTs, trigger PMEs to assist and reinforce the OGs and MeOH signalling in priming is an interesting hypothesis to be investigated.

In conclusion, this study brings new insights into the molecular mechanisms underlying the activation of PME related immunity against necrotrophs increasing our understanding of the complexity of the interactions between plants and microbes. A switch on mechanism, mediated by SBTs for fine tuning PME activity and pectin integrity signaling, emerges providing knowledge about how plants modulate immunity. We identify Pro-PMEs as the target substrate processed by SBT3.3 indicating how this host factor mediates immune prime activation. Future studies will be useful to identify the specific SBT-Pro-PME pairs and to quantify the contribution of PME activity to priming. Activation of Pro-PMEs and plant immunity by Subtilases processing becomes an effective genomic tool to engineer durable, broad-spectrum plant disease resistance and to avoid a growth-defense trade-offs.

EXPERIMENTAL PROCEDURES

Plant, Bacteria and Fungus growth conditions

Arabidopsis thaliana Wild-type Columbia (Col-0), *sbt* mutants and SBT3.3-OE seedlings were grown

in solid MS/2 medium [Murashige and Skoog Medium including vitamins (2.2 g L^{-1}), 1% sucrose,

0.8% plant agar, pH 5.7] in a controlled environmental chamber maintained at 22°C and 70% relative

humidity, with a 16h/8h day/night cycle (PAR level of 100 μmol m^{-2}s^{-1}). Adult plants were transferred

on soil in a growth chamber at 22°C and 70% relative humidity under a 12 h/12h day/night cycle (PAR

level of 120 μmol m^{-2}s^{-1}). *Sbt* mutants used include SALK_086092 (At1g32960; *sbt3.3-1*),

SALK_107460 (At1g32960; *sbt3.3-2*), SAIL_400_F09 (At1g32940; *sbt3.5-1*) and GABI_672C08

(At1g32940; *sbt3.5-2*). All mutants are in Col-0 background and are T-DNA insertion lines obtained

from the Nottingham *Arabidopsis* Stock Centre (NASC). *Nicotiana tabacum cv* SR1 seeds were

sterilized using commercial grade bleach solution in the presence of 0.1% (v/v) for 15 min at room

temperature and then washed three times for 5 min with distilled water before sowing. The seeds were

germinated in a growth room at 25°C and 30% relative humidity under a 16h/8h day/night cycle

(photosynthetic photon flux of 50 μmol m^{-2}s^{-1}). Three- to four-week-old plants from MS medium were

transferred on soil and grown in a controlled-environment at 25°C and 40% relative humidity under a

16h/8h day/night cycle (photosynthetic photon flux of 120 μmol m^{-2}s^{-1}). *Escherichia coli* TOP10 cells

was grown in LB agar plate (Luria-Bertani: tryptone 1%, yeast exctract 0.5%, NaCl 1%) containing

ampicillin 50μg/ml and incubated overnight at 37°C. *Agrobacterium tumefaciens* (Strain GV3110) was

grown in LB medium (Luria-Bertani: tryptone 1%, yeast exctract 0.5%, NaCl 1%) containing

appropriate antibiotics (Rifampycin 100μg/ml and Spectomycin 100μg/ml or Kanamycin 50μg/ml) and

incubated for 2 days at 28°C. *Botrytis cinerea* strain SF1 (Lionetti et al., 2007) was grown for 20 days

on Malt Extract Agar at 20 g L^{-1} with Mycological Peptone at 10 g L^{-1} (MEP) at 23°C and 70% relative

humidity in the dark before conidia collection.

For seed germination analysis, *A. thaliana* WT, sbt mutants and SBT3.3-OE seeds were washed in 2

mL of isopropanol for 30 seconds followed by washing in 2 mL of sterile water for 3 min in slow

agitation. Seeds were treated with 2 mL of sterilization solution (400 μl NaClO, 1.6 ml sterile water)

for 5 min in slow agitation, followed by 4 washing steps in 2 ml of sterile water. For seed germination

assay, triplicate sets of 60–70 sterilized seeds for each genotype were sown in Petri dishes on solid

MS/2 medium in a controlled environmental chamber maintained at 22°C and 70% relative humidity,

with a 16h/8h day/night cycle (PAR level of 100 μmol m-2s-1). Seeds germination rate, as the number

of germinated seeds divided by the total amount of total seeds, were scored at 24, 48 and 72 hours.

Genotyping of T-DNA insertion mutants

Genomic DNA was extracted using RBC Bioscience Genomic DNA Extraction Kits, according to the manufacturer's instructions, from leaves of 4-week-old *Arabidopsis* wild-type and mutant plants and subjected to a PCR-based screening using sequences of primer, described in Supplemental Table S2, generated with Primer3 (https://primer3.ut.ee/). Taq Kapa DNA Polymerase (Kapa Biosystem) was used at 1 units/30 µl of assay. Amplification was carried out in presence of dNTPs (0.2 mM), 2.5 mM $MgCl_2$ and specific primers (10 µM) in the 1x buffer provide by the supplier. Conditions for amplification were: 95°C for 3 minutes; 36 cycles of amplification: 95°C for 30 sec, 60°C for 30 sec, 72°C for 60 sec, and a final extension of 72°C for 1 min. PCR products were separated by agarose gel electrophoresis and visualized by ethidium bromide staining. Plants homozygous for the mutations were propagated and used for all subsequent experiments.

Stable transformation of SBT3.3-OE *Arabidopsis* plants

Flowering *Arabidopsis* plants were transformed by floral dipping for 15 seconds in 200 mL of the culture medium with the inoculum of *Agrobacterium tumefaciens* C58 carrying the construction of interest. The plants were subsequently covered for 1 day to promote the infection. The culture medium for immersion was composed of MS 2.2 g/L and sucrose 5% (p/v), supplemented with Silwet 0.005% and BAP 0.05 ug/mL. Subsequently, the selection of homozygous transgenic plants for the *SBT3.3-YFP* construction was carried out by means of selection with BASTA herbicide.

Elicitor Treatment and Gene expression analysis

B. cinerea-infected or mock-inoculated leaves were frozen in liquid nitrogen and homogenized using the mixer mill MM301 (RETSCH) and using inox beads (5 mm diameter) for about 1 min at 30 Hz, and total RNA was extracted with Isol-RNA Lysis Reagent (5′-Prime) according to the manufacturer's instructions. RNA was treated with RQ1 DNase (Promega), and first-strand complementary DNA (cDNA) was synthesized using ImProm-II reverse transcriptase (Promega). Quantitative Reverse Transcription PCR analysis was performed using a CFX96 Real-Time System (Bio-Rad). One microliter of cDNA (corresponding to 50 ng of total RNA) was amplified in 20 μL of reaction mix containing 1× Go Taq qPCR Master Mix (Promega) and 0.5 μM of each primer described in Supplemental TableS1. Conditions for amplification were: 95°C for 2 minutes; 46 cycles of amplification: 95°C for 15 sec, 58°C for 15 sec, and a final extension of 72°C for 15 sec. Expression

levels of each gene, relative to the UBIQUITIN5 (UBQ5) gene, were determined using a modification of the Pfaffl method (Pfaffl, 2001). Primer sequences were generated with Primer3 software (https://primer3.ut.ee/). Seedlings of 10-days-old *Arabidopsis* cultured in liquid MS/2 medium were treated for 1h with OGs (40 µg mL^{-1}) or water as a control, collected and processed for RNA extraction as described above.

Arabidopsis thaliana infection with *Botrytis cinerea*

Conidia of *B. cinerea* were harvested by washing the surface of the mycelium with sterile distilled water. Conidia suspensions were filtered to remove residual mycelium and the conidia concentration was determined using a Thoma chamber. Fully developed leaves from 4-week-old *A. thaliana* plants were infected with $1x10^6$ conidia mL^{-1} incubated in Potato Dextrose Broth (PDB) at 24 g L^{-1}. Six droplets of spore suspension (5 µL each) were placed on the surface of each leaf. Mock-inoculation was performed using PDB. Plants were incubated at 24°C with a 12h/12h day/night photoperiod. The lesion size produced by *B. cinerea* was evaluated as an indicator of susceptibility to the fungus.

Determination of PME activity, fungal diffusion and H$_2$O$_2$ accumulation

Total protein extracts were obtained by homogenizing uninfected and infected *Arabidopsis* leaves in the presence of 1M NaCl, 12.5 mM Citric Acid, 50 mM Na$_2$HPO$_4$, 0.02 % Sodium Azide, protease inhibitor 1:200 v/v (P8849, Sigma), pH 6.5. Each sample was resuspended in 500 µL of protein Extraction Buffer. The homogenates were shaken for 3 h at 4°C, centrifuged at 13.000 x g for 15 min, and the supernatant (400 µL) collected. Protein concentration was determined in the supernatants using Bradford protein assay method (Bradford reagent, Sigma–Aldrich), a standard curve was performed using BSA (bovin serum albumine) at 1 mg mL^{-1}. PME activity was evaluated by using PECTOPLATE assay (Lionetti, 2015): PECTOPLATE was prepared with 0.1% (w/v) of apple pectin (molecular weight range 30,000–100,000 Da; 70–75% esterification; 76282, Sigma-Aldrich, St. Louis), 1% (w/v) SeaKem® LE agarose (Lonza, Basel, Switzerland, Catalog no: 50004E), 12.5 mM citric acid and 50 mM Na$_2$HPO$_4$, pH 6.5. Equal amounts of protein samples (2 µg of total protein in 20 µL) were loaded in each well of PECTOPLATE. Plates were incubated at 30°C for 16 h, and stained with 0.05% (w/v) RR (R2751; Sigma-Aldrich, St. Louis) for 30 min. The plates were de-stained by several washes with water and the area of the fuchsia-stained haloes, resulting from de-methylesterification of pectin and the area of inner unstained haloes, resulting from the hydrolysis of pectin in the gel, were measured with ImageJ software (Abramoff, Magalhes, and am 2004). Known amounts of commercially available

PME from orange (Citrus spp.) peel (P5400; Sigma-Aldrich) was used in the PECTOPLATE to generate a standard curve used to calculate PME activity in the protein extracts.

For histochemical assay, leaves of *Arabidopsis* WT mutants and SBT3.3-OE after 48 h post *Botrytis cinerea* inoculation, were dipped in a solution with 0.05% (w/v) Ruthenium Red (RR) and subjected to vacuum infiltration for 10 min. Excess of dye was removed by several washes of stained leaves with water. Chlorophyll was extracted with hot bleaching solution (ethanol: acid acetic: glycerol = 3:1:1) at 90-95°C for 15 min. Leaves were mounted in 50% glycerol and examined by light microscopy using Nikon eclipse E200 microscope. Images were taken with a Nikon Digital Sight DS-Fi1c camera. For the detection of *B. cinerea* hyphae in infected plant tissues, detached *Arabidopsis* leaves were incubated for 30 min in lactophenol Trypan Blue solution (water: glycerol: lactic acid: phenol [1:1:1:1] + Trypan Blue solution [1 mg mL^{-1}; Sigma-Aldrich]). Thereafter, leaves were washed for overnight with gentle agitation in absolute ethanol to remove chlorophyll. Finally, leaves were transferred into 50% (v/v) glycerol solution for microscopy. Leaves of 4-week-old *Arabidopsis* wild-type, *sbt* mutants and SBT3.3-OE line were assayed for H_2O_2 accumulation employing 3,3′-diaminobenzidine staining as previously described (Daudi and O'Brien, 2012).

Determination of Monosaccharide Composition of CW and MEOH quantification

Extraction of alcohol-insoluble residue, CW monosaccharide composition and the determination of CW methylesters were performed as previously described (Lionetti et al., 2017).

SBT3.3-YFP and PME17-GFP/RFP Gene Constructs

For the SBT3.3-YFP overexpressing construct, a full-length cDNA for SBT3.3 was amplified by PCR using Pfu DNA polymerase (Stratagene, San Diego, CA) and specific primers, described in Table S3, including Gateway adapters: BP SBT3.3 FW and BP SBT3.3 RV and recombined into pDONR207 using BP ClonaseMixII kit (Invitrogen). After sequencing, the construct was recombined with pB7FWG2 destination vector (YFP in C-terminal) using LR ClonaseMixII kit (Invitrogen) and introduced into *Arabidopsis* (*sbt3.3*) via Agrobacterium transformation.

For PME17-GFP/RFP overexpressing constructs, the full-length cDNA for PME17 was amplified by PCR using *TakaRa* Ex Taq DNA polymerase (TAKARA, Kyoto), and specific primers flanked by Gateway adapters, attB sequences, described in Table S3. PCR amplified attB-flanked DNA was recombined into pDONR221 using BP ClonaseMixII kit (Invitrogen). Specific pDESTs, pK7FWG2 and pK7RWG2,0, were used for GFP and RFP fusion in C-terminus to obtain Pro-PME17-GFP and

Pro-PME17-RFP, respectively. All constructs were checked by sequencing (Eurofins Genomics, https://eurofinsgenomics.eu/). The Gateway constructs, PME17-GFP, PME17-RFP, SBT3.3-YFP, SBT3.3-mCherry, RFP-HDEL, ST52-mCherry, XTH11-RFP and pm-rk were introduced into *Agrobacterium tumefaciens* (Strain GV3110) and the agroinfiltration was performed in *Nicotiana tabacum* leaves as previously described (De Caroli et al., 2011).

Transient transformation of *Nicotiana tabacum* leaves and confocal Laser Scanning Microscopy

Almost fully expanded Nicotiana tabacum 6-week-old leaves were infiltrated with a suspension of Agrobacterium tumefaciens bearing the relevant construct in 2mM Na3PO4; 50mM 2-(N-morpholine)-ethanesulphonic acid (MES); glucose 0.5%; acetosyringone 100μM; pH= 5.6 at an OD_{600}=0.5. After 48 and 52 hours post infiltration (hpi) fluorescence was analyzed in infiltrated leaves by confocal microscopy. For co-infiltration, Agrobacterium cultures grown separately were adjusted to an OD_{600}=0.5, and mixed prior to infiltration. Transiently transformed tobacco leaves were observed using a confocal laser scanning microscope LSM710 Zeiss (http://www.zeiss.com) mounting material in water as previously described (De Caroli et al., 2011). GFP and YFP were detected within the short 505–530 nm wavelength range, assigning the green color; RFP and mCherry were detected within 560–615 nm, assigning the red color. Excitation wavelengths of 488 and 543 nm were used. The laser power was set to a minimum and appropriate controls were made to ensure there was no bleed-through from one channel to the other. Images were processed using PHOTOSHOP7.0 (Adobe, https://www.adobe.com). For quantitative colocalization analyses, the overlap coefficient (OC, after Manders) and the Pearson's correlation coefficient (R) were evaluated as previously reported (De Caroli et al., 2011).

Bioinformatics tools

The Expression Angler tool of the Bio-Analytic Resource for Plant Biology (BAR, http://bar.utoronto.ca/welcome.htm) was used for co-expression analysis. E-plant (https://bar.utoronto.ca/eplant/) was used for to study developmental map and biotic stresses. Chromas was used to manage both nucleotide sequences and amino acids (https://chromas.software.informer.com/2.5/). Primer3 (https://primer3.ut.ee/) was used to design primers respectively for Gene expression analysis and Genotyping analysis. ImageJ was used to calculate the area of the haloes in PECTOPLATE assay (http://imagej.nih.gov/ij/download.html). DNAMAN software (Lynnon, BioSoft, Quebec, Canada, https://www.lynnon.com/) was used for

amino acid sequence analyses. Genevestigator Meta-Analyzer Tools (www.genevestigator.ethz.ch/at/)

was used to analyze *SBT3.3* and *SBT3.5* gene expression during infection of *Arabidopsis* with different

pathogens and DAMPs and PAMPs treatments.

QUANTIFICATION AND STATISTICAL ANALYSIS

The statistical significance of the difference in gene expression between specific samples was analyzed

through ANOVA followed by Tukey's test (P < 0.05), for all other statistical analyses the Student's t

test was used. The error bars represent standard deviation of mean.

REFERENCES

Abramoff MD, Magalhes PJ, am SJ (2004) Image processing with ImageJ. Biophotonics international **11**: 36–42

AbuQamar S, Moustafa K, Tran LS (2017) Mechanisms and strategies of plant defense against Botrytis cinerea. Critical Reviews in Biotechnology **37**: 262–274

An SH, Sohn KH, Choi HW, Hwang IS, Lee SC, Hwang BK (2008) Pepper pectin methylesterase inhibitor protein CaPMEI1 is required for antifungal activity, basal disease resistance and abiotic stress tolerance. Planta **228**: 61–78

Bethke G, Grundman RE, Sreekanta S, Truman W, Katagiri F, Glazebrook J (2014) Arabidopsis PECTIN METHYLESTERASES Contribute to Immunity Against Pseudomonas syringae. Plant Physiol **164**: 1093–1107

Birkenbihl RP, Diezel C, Somssich IE (2012) Arabidopsis WRKY33 is a key transcriptional regulator of hormonal and metabolic responses toward Botrytis cinerea infection. Plant Physiol **159**: 266–285

Bosch M, Cheung AY, Hepler PK (2005) Pectin Methylesterase, a Regulator of Pollen Tube Growth. Plant Physiol **138**: 1334–1346

Bruce TJA, Matthes MC, Napier JA, Pickett JA (2007) Stressful "memories" of plants: Evidence and possible mechanisms. Plant Science **173**: 603–608

Brutus A, Sicilia F, Macone A, Cervone F, De Lorenzo G (2010) A domain swap approach reveals a role of the plant wall-associated kinase 1 (WAK1) as a receptor of oligogalacturonides. Proc Natl Acad Sci USA **107**: 9452–9457

Clay NK, Adio AM, Denoux C, Jander G, Ausubel FM (2009) Glucosinolate metabolites required for an Arabidopsis innate immune response. Science **323**: 95–101

Coculo D, Lionetti V (2022) The Plant Invertase/Pectin Methylesterase Inhibitor Superfamily. Frontiers in Plant Science **13**: 863892

Conrath U (2011) Molecular aspects of defence priming. Trends in Plant Science **16**: 524–531

Daudi A, O'Brien JA (2012) Detection of Hydrogen Peroxide by DAB Staining in Arabidopsis Leaves. Bio Protoc **2**: e263

De Caroli M, Lenucci MS, Di Sansebastiano GP, Dalessandro G, De Lorenzo G, Piro G (2011) Protein trafficking to the cell wall occurs through mechanisms distinguishable from default sorting in tobacco. Plant J **65**: 295–308

De Caroli M, Lenucci MS, Manualdi F, Dalessandro G, De Lorenzo G, Piro G (2015) Molecular dissection of Phaseolus vulgaris polygalacturonase-inhibiting protein 2 reveals the presence of hold/release domains affecting protein trafficking toward the cell wall. Front.Plant Sci. 6:

De Caroli M, Manno E, Piro G, Lenucci MS (2021) Ride to cell wall: Arabidopsis XTH11, XTH29
and XTH33 exhibit different secretion pathways and responses to heat and drought stress. The
Plant Journal **107**: 448–466

Dean R, van Kan JAL, Pretorius ZA, Hammond-Kosack KE, Di Pietro A, Spanu PD, Rudd JJ,
Dickman M, Kahmann R, Ellis J, et al (2012) The Top 10 fungal pathogens in molecular
plant pathology. MolPlant Pathol **13**: 414–430

Decreux A, Messiaen J (2005) Wall-associated kinase WAK1 interacts with cell wall pectins in a
calcium-induced conformation. Plant Cell Physiol **46**: 268–278

Dedeurwaerder S, Menu-Bouaouiche L, Mareck A, Lerouge P, Guerineau F (2008) Activity of an
atypical Arabidopsis thaliana pectin methylesterase. Planta **229**: 311–21

Del Corpo D, Fullone MR, Miele R, Lafond M, Pontiggia D, Grisel S, Kieffer-Jaquinod S,
Giardina T, Bellincampi D, Lionetti V (2020) AtPME17 is a functional Arabidopsis thaliana
pectin methylesterase regulated by its PRO region that triggers PME activity in the resistance to
Botrytis cinerea. Molecular Plant Pathology **21**: 1620–1633

Denoux C, Galletti R, Mammarella N, Gopalan S, Werck D, De Lorenzo G, Ferrari S, Ausubel
FM, Dewdney J (2008) Activation of defense response pathways by OGs and Flg22 elicitors in
Arabidopsis seedlings. Mol Plant **1**: 423–445

Dorokhov YL, Komarova TV, Petrunia IV, Frolova OY, Pozdyshev DV, Gleba YY (2012)
Airborne Signals from a Wounded Leaf Facilitate Viral Spreading and Induce Antibacterial
Resistance in Neighboring Plants. Plos Pathogens 8:

Engelsdorf T, Gigli-Bisceglia N, Veerabagu M, McKenna JF, Vaahtera L, Augstein F, Does DV
der, Zipfel C, Hamann T (2018) The plant cell wall integrity maintenance and immune
signaling systems cooperate to control stress responses in Arabidopsis thaliana. Sci Signal **11**:
eaao3070

Feng W, Kita D, Peaucelle A, Cartwright HN, Doan V, Duan Q, Liu M-C, Maman J, Steinhorst
L, Schmitz-Thom I, et al (2018) The FERONIA Receptor Kinase Maintains Cell-Wall
Integrity during Salt Stress through Ca2+ Signaling. Curr Biol **28**: 666-675.e5

Ferrari S, Savatin DV, Sicilia F, Gramegna G, Cervone F, De Lorenzo G (2013)
Oligogalacturonides: plant damage-associated molecular patterns and regulators of growth and
development. Front.Plant Sci. doi: 10.3389/fpls.2013.00049:

Figueiredo A, Monteiro F, Sebastiana M (2014) Subtilisin-like proteases in plant-pathogen
recognition and immune priming: a perspective. FrontPlant Sci **5**: 739.

Figueiredo J, Sousa Silva M, Figueiredo A (2018) Subtilisin-like proteases in plant defence: the past,
the present and beyond: Subtilases in plant defence. Molecular Plant Pathology **19**: 1017–1028

Gamir J, Minchev Z, Berrio E, García JM, De Lorenzo G, Pozo MJ (2021) Roots drive
oligogalacturonide-induced systemic immunity in tomato. Plant, Cell & Environment **44**: 275–
289

Giovannoni M, Lironi D, Marti L, Paparella C, Vecchi V, Gust AA, De Lorenzo G, Nürnberger T, Ferrari S (2021) The Arabidopsis thaliana LysM-containing Receptor-Like Kinase 2 is required for elicitor-induced resistance to pathogens. Plant, Cell & Environment **44**: 3775–3792

Godson A, van der Hoorn RAL (2021) The front line of defence: a meta-analysis of apoplastic proteases in plant immunity. Journal of Experimental Botany **72**: 3381–3394

Guo H, Nolan TM, Song G, Liu S, Xie Z, Chen J, Schnable PS, Walley JW, Yin Y (2018) FERONIA Receptor Kinase Contributes to Plant Immunity by Suppressing Jasmonic Acid Signaling in Arabidopsis thaliana. Curr Biol **28**: 3316-3324.e6

Ha CM, Rao X, Saxena G, Dixon RA (2021) Growth–defense trade-offs and yield loss in plants with engineered cell walls. New Phytologist **231**: 60–74

Hann CT, Bequette CJ, Dombrowski JE, Stratmann JW (2014) Methanol and ethanol modulate responses to danger- and microbe-associated molecular patterns. Front.Plant Sci. 5:

He Z, Webster S, He SY (2022) Growth-defense trade-offs in plants. Curr Biol **32**: R634–R639

Hou S, Jamieson P, He P (2018) The cloak, dagger, and shield: proteases in plant–pathogen interactions. Biochemical Journal **475**: 2491–2509

Huot B, Yao J, Montgomery BL, He SY (2014) Growth–Defense Tradeoffs in Plants: A Balancing Act to Optimize Fitness. Molecular Plant **7**: 1267–1287

Khan AR, James MN (1998) Molecular mechanisms for the conversion of zymogens to active proteolytic enzymes. Protein Sci **7**: 815–836

Kohorn BD, Kohorn SL (2012) The cell wall-associated kinases, WAKs, as pectin receptors. FrontPlant Sci **3**: 88

Kohorn BD, Kohorn SL, Saba NJ, Martinez VM (2014) Requirement for Pectin Methyl Esterase and Preference for Fragmented over Native Pectins for Wall-associated Kinase-activated, EDS1/PAD4-dependent Stress Response in Arabidopsis. JBiolChem **289**: 18978–18986

Komarova TV, Sheshukova EV, Dorokhov YL (2014) Cell wall methanol as a signal in plant immunity. Front.Plant Sci. 5:

Lichtenthaler SF, Lemberg MK, Fluhrer R (2018) Proteolytic ectodomain shedding of membrane proteins in mammals—hardware, concepts, and recent developments. EMBO J **37**: e99456

Lin W, Tang W, Anderson CT, Yang Z (2018) FERONIA's sensing of cell wall pectin activates ROP GTPase signaling in Arabidopsis. bioRxiv 269647

Lin W, Tang W, Pan X, Huang A, Gao X, Anderson CT, Yang Z (2021) Arabidopsis pavement cell morphogenesis requires FERONIA binding to pectin for activation of ROP GTPase signaling. Current Biology S096098222101589X

Lionetti V (2015) PECTOPLATE: the simultaneous phenotyping of pectin methylesterases, pectinases, and oligogalacturonides in plants during biotic stresses. FrontPlant Sci **6**: 1–8

Lionetti V, Cervone F, Bellincampi D (2012) Methyl esterification of pectin plays a role during plant-pathogen interactions and affects plant resistance to diseases. JPlant Physiol **169**: 1623–1630

Lionetti V, Fabri E, De Caroli M, Hansen AR, Willats WG, Piro G, Bellincampi D (2017) Three pectin methyl esterase inhibitors protect cell wall integrity for immunity to Botrytis. Plant Physiol **Jan 12.**: pii: pp.01185.2016.

Lionetti V, Francocci F, Ferrari S, Volpi C, Bellincampi D, Galletti R, D'Ovidio R, De Lorenzo G, Cervone F (2010) Engineering the cell wall by reducing de-methyl-esterified homogalacturonan improves saccharification of plant tissues for bioconversion. Proc Natl Acad Sci USA **107**: 616–621

Lionetti V, Raiola A, Camardella L, Giovane A, Obel N, Pauly M, Favaron F, Cervone F, Bellincampi D (2007) Overexpression of Pectin Methylesterase Inhibitors in Arabidopsis Restricts Fungal Infection by Botrytis cinerea. Plant Physiol **143**: 1871–1880

Lionetti V, Raiola A, Cervone F, Bellincampi D (2014) Transgenic expression of pectin methylesterase inhibitors limits tobamovirus spread in tobacco and Arabidopsis. MolPlant Pathol **15**: 265–274

Liu N, Sun Y, Pei Y, Zhang X, Wang P, Li X, Li F, Hou Y (2018) A Pectin Methylesterase Inhibitor Enhances Resistance to Verticillium Wilt1[OPEN]. Plant Physiol **176**: 2202–2220

Lozano-Torres JL, Wilbers RHP, Warmerdam S, Finkers-Tomczak A, Diaz-Granados A, Schaik CC van, Helder J, Bakker J, Goverse A, Schots A, et al (2014) Apoplastic Venom Allergen-like Proteins of Cyst Nematodes Modulate the Activation of Basal Plant Innate Immunity by Cell Surface Receptors. PLOS Pathogens **10**: e1004569

Meyer M, Huttenlocher F, Cedzich A, Procopio S, Stroeder J, Pau-Roblot C, Lequart-Pillon M, Pelloux J, Stintzi A, Schaller A (2016) The subtilisin-like protease SBT3 contributes to insect resistance in tomato. EXBOTJ **67**: 4325–4338

Nelson BK, Cai X, Nebenfuhr A (2007) A multicolored set of in vivo organelle markers for co-localization studies in Arabidopsis and other plants. Plant J **51**: 1126–1136

Ogawa S, Wakatake T, Spallek T, Ishida JK, Sano R, Kurata T, Demura T, Yoshida S, Ichihashi Y, Schaller A, et al (2021) Subtilase activity in intrusive cells mediates haustorium maturation in parasitic plants. Plant Physiology **185**: 1381–1394

Osorio S, Bombarely A, Giavalisco P, Usadel B, Stephens C, Araguez I, Medina-Escobar N, Botella MA, Fernie AR, Valpuesta V (2011) Demethylation of oligogalacturonides by FaPE1 in the fruits of the wild strawberry Fragaria vesca triggers metabolic and transcriptional changes associated with defence and development of the fruit. JExpBot **62**: 2855–2873

Osorio S, Castillejo C, Quesada MA, Medina-Escobar N, Brownsey GJ, Suau R, Heredia A, Botella MA, Valpuesta V (2008) Partial demethylation of oligogalacturonides by pectin methyl esterase 1 is required for eliciting defence responses in wild strawberry (Fragaria vesca). Plant J **54**: 43–55

Paulus JK, Van der Hoorn RAL (2019) Do proteolytic cascades exist in plants? Journal of
Experimental Botany **70**: 1997–2002

Pelloux J, Rustérucci C, Mellerowicz EJ (2007) New insights into pectin methylesterase structure
and function. Trends Plant Sci **12**: 267–277

Pogorelko G, Lionetti V, Fursova O, Sundaram RM, Qi MS, Whitham SA, Bogdanove AJ,
Bellincampi D, Zabotina OA (2013) Arabidopsis and Brachypodium distachyon Transgenic
Plants Expressing Aspergillus nidulans Acetylesterases Have Decreased Degree of
Polysaccharide Acetylation and Increased Resistance to Pathogens. Plant Physiol **162**: 9–23

Ramirez V, Lopez A, Mauch-Mani B, Gil MJ, Vera P (2013) An Extracellular Subtilase Switch for
Immune Priming in Arabidopsis. Plos Pathogens 9:

Rautengarten C, Steinhauser D, Bussis D, Stintzi A, Schaller A, Kopka J, Altmann T (2005)
Inferring Hypotheses on Functional Relationships of Genes: Analysis of the Arabidopsis
thaliana Subtilase Gene Family. PLoS.Comput.Biol. 1:

Rautengarten C, Usadel B, Neumetzler L, Hartmann J, Büssis D, Altmann T (2008) A subtilisin-
like serine protease essential for mucilage release from Arabidopsis seed coats. The Plant
Journal **54**: 466–480

Reem N, Pogorelko G, Lionetti V, Chambers L, Held MA, Bellincampi D, Zabotina OA (2016)
Decreased polysaccharide feruloylation compromises plant cell wall integrity and increases
susceptibility to necrotrophic fungal pathogens. FrontPlant Sci **7**: 630, doi:
10.3389/fpls.2016.00630

Rosenfeld N, Elowitz MB, Alon U (2002) Negative autoregulation speeds the response times of
transcription networks. J Mol Biol **323**: 785–793

Schaller A (2004) A cut above the rest: the regulatory function of plant proteases. Planta **220**: 183–197

Schaller A, Stintzi A, Rivas S, Serrano I, Chichkova NV, Vartapetian AB, Martínez D, Guiamét
JJ, Sueldo DJ, van der Hoorn RAL, et al (2018) From structure to function - a family portrait
of plant subtilases. New Phytol **218**: 901–915

Senechal F, Graff L, Surcouf O, Marcelo P, Rayon C, Bouton S, Mareck A, Mouille G, Stintzi A,
Hofte H, et al (2014a) Arabidopsis PECTIN METHYLESTERASE17 is co-expressed with and
processed by SBT3.5, a subtilisin-like serine protease. AnnBot **114**: 1161–1175

Senechal F, Wattier C, Rusterucci C, Pelloux J (2014b) Homogalacturonan-modifying enzymes:
structure, expression, and roles in plants. JExpBot **65**: 5125–5160

Serrano I, Buscaill P, Audran C, Pouzet C, Jauneau A, Rivas S (2016) A non canonical subtilase
attenuates the transcriptional activation of defence responses in Arabidopsis thaliana. eLife **5**:
e19755

Siegmund U, Viefhues A (2016) Reactive Oxygen Species in the Botrytis – Host Interaction. *In* S
Fillinger, Y Elad, eds, Botrytis – the Fungus, the Pathogen and its Management in Agricultural
Systems. Springer International Publishing, Cham, pp 269–289

Stegmann M, Monaghan J, Smakowska-Luzan E, Rovenich H, Lehner A, Holton N, Belkhadir Y,
Zipfel C (2017) The receptor kinase FER is a RALF-regulated scaffold controlling plant
immune signaling. Science **355**: 287–289

Tornero P, Conejero V, Vera P (1996) Primary structure and expression of a pathogen-induced
protease (PR-P69) in tomato plants: Similarity of functional domains to subtilisin-like
endoproteases. Proceedings of the National Academy of Sciences **93**: 6332–6337

Vartapetian AB, Tuzhikov AI, Chichkova NV, Taliansky M, Wolpert TJ (2011) A plant alternative
to animal caspases: subtilisin-like proteases. Cell Death Differ **18**: 1289–1297

Wang S, Xing R, Wang Y, Shu H, Fu S, Huang J, Paulus JK, Schuster M, Saunders DGO, Win J,
et al (2021) Cleavage of a pathogen apoplastic protein by plant subtilases activates host
immunity. New Phytol **229**: 3424–3439

Williamson B, Tudzynski B, Tudzynski P, Kan J a. LV (2007) Botrytis cinerea: the cause of grey
mould disease. Molecular Plant Pathology **8**: 561–580

Wolf S, Rausch T, Greiner S (2009) The N-terminal pro region mediates retention of unprocessed
type-I PME in the Golgi apparatus. The Plant Journal **58**: 361–375

Wormit A, Usadel B (2018) The Multifaceted Role of Pectin Methylesterase Inhibitors (PMEIs). Int J
Mol Sci. doi: 10.3390/ijms19102878

Zhou J-M, Zhang Y (2020) Plant Immunity: Danger Perception and Signaling. Cell **181**: 978–989

Zhou N, Tootle TL, Glazebrook J (1999) Arabidopsis PAD3, a gene required for camalexin
biosynthesis, encodes a putative cytochrome P450 monooxygenase. Plant Cell **11**: 2419–28

    Figure legends

    Figure 1. The lack of SBT3.3 and SBT3.5 expression in Arabidopsis does not alter rosette growth
    and cell wall composition. A) Schematic representation of SBT3.3 and SBT3.5 gene structures. The
    localization of T-DNA insertions in the gene sequences are shown (black arrows). 5′-UTR and 3′-UTR
    are represented in grey, exons in white and introns as a black line. **B)** The expression of *SBT3.3* and
    SBT3.5 was analyzed in the respective mutants and WT plants at 48-hour post inoculation with *B.*
    cinerea. The expression levels were normalized to *Ubiquitin 5* expression. Data represent mean ± SD
    of three independent experiments (n=3). Asterisks indicate significant differences in mutants respect to
    WT plants using a *Student's t–test* ($P < 0.05$). **C)** Representative pictures illustrating the morphology of
    vegetative rosettes (four-weeks old plants) of *sbt3.3* and *sbt3.5* mutants compared with WT. **D)** The
    monosaccharide composition of cell wall matrix polysaccharides was compared in leaves of *sbt3.3* and
    s*bt3.5* mutants with WT plants. The molar percentages of Fucose (Fuc), Rhamnose (Rha), Arabinose
    (Ara), Galactose (Gal), Glucose (Glu), Xylose (Xyl), Mannose (Man), Galacturonic Acid (Gal A) and
    Glucuronic Acid (Glc A) were quantified. The results represent the mean ± SD (n≥3). Monosaccharide
    datasets are not significantly different according to analysis of variance (ANOVA) followed by
    Tukey's test ($P < 0.05$).

    Figure 2. SBT3.3 and SBT3.5 regulate PME activity during disease. A) Schematic representation of
    the binding of ruthenium red to de-methylesterified pectin. **B)** Histochemical evidence indicating the
    induction of PME activity in WT, *sbt3.3* and *sbt3.5* mutants during *Botrytis* infection at the point of
    fungal penetration by staining leaves with ruthenium red. Bars = 1mm. **C)** Leaves of Arabidopsis WT
    and of *sbt* mutants were inoculated with *Botrytis* and mock and PME activity were quantified at 48-
    hour post inoculation (hpi). The results represent the mean ± SD (n = 3). The different letters on the
    bars indicate datasets significantly different according to ANOVA followed by Tukey's test ($P < 0.05$).
    The experiments were repeated three times with similar results.

    Figure 3. The lack of *SBT3.3 and SBT3.5* expression reduce Arabidopsis resistance to *Botrytis*
    ***cinerea*. A)** Quantification of lesion areas produced by the spreading of the fungus at 48 hours post
    infections. Data represent mean ± SD (n≥10 lesions for each genotype). The experiments were
    repeated three times with similar results. Different letters indicate statistically significant differences,
    according to ANOVA followed by Tukey's significance test ($P < 0.05$). **B)** Photographs showing lesion

areas produced by *Botrytis* on leaves of Arabidopsis WT, *sbt3.3-1 and sbt3.5-1* mutants (left) and

microphotographs showing *Botrytis* colonization revealed by trypan blue staining (right). Bars = 1mm.

Figure 4. The SBT3.3 overexpression in Arabidopsis does not alter rosette growth and cell wall

composition. A) The expression of *SBT3.3* was analyzed in the *SBT3.3-OE1* and *SBT3.3-OE2*

transgenic plants at 48-hour post inoculation with Mock and *B. cinerea* spores. The expression levels

were normalized to *Ubiquitin5* expression. Data represent mean ± SD. The experiments were repeated

three times with similar results. **B)** Representative pictures illustrating the morphology of vegetative

rosettes of SBT3.3 overexpressing plants compared with WT plants. **C)** The molar percentages of

Fucose (Fuc), Rhamnose (Rha), Arabinose (Ara), Galactose (Gal), Glucose (Glu), Xylose (Xyl),

Mannose (Man), Galacturonic Acid (Gal A) and Glucuronic Acid (Glc A) were quantified. The results

represent the mean ± SD (n ≥ 3). Monosaccharide datasets are not significantly different according to

analysis of variance (ANOVA) followed by Tukey's test ($P < 0.05$).

Figure 5. The SBT3.3 overexpression overactivates PME activity restricting *Botrytis* infection. A)

Leaves of Arabidopsis WT and of SBT3.3-OE plants were inoculated with *Botrytis* and mock and PME

activity were quantified at 48-hour post inoculation. The results represent the mean ± SD (n = 3). **B)**

Histochemical evidence indicating the induction of PME activity in WT and SBT3.3-OE1 plants during

Botrytis infection at the point of fungal penetration by staining leaves with ruthenium red. Bars = 1mm.

C) The amount of methylesters in the cell wall were quantified at 48-hour post inoculation. **D)**

Quantification of lesion areas produced by the spreading of the fungus at 48 hours post infections. Data

represent mean ± SD (n≥10 lesions for each genotype). The experiments were repeated three times

with similar results. **E)** Photographs showing lesion areas produced by *Botrytis* on leaves of WT and

SBT3.3-OE1 plants (left) and microphotographs showing *Botrytis* colonization revealed by trypan blue

staining. Bars = 1mm. The different letters in A , C and D indicate datasets significantly different

according to ANOVA followed by Tukey's test ($P < 0.05$). The experiments were repeated three times

with similar results.

Figure 6. *SBT3.3* overexpression enhances immune responces. The induction of the expression of

defence genes *WRKY33, PAD3, WAK2, CYP81F2, PME17, PMEI10* were compared between *SBT3.3-*

OE1 line and WT plants challenged with *Botrytis*. The mRNA expression was analysed at 48 hpi in *B.*

cinerea- or mock-inoculated leaves of 6-week-old *Arabidopsis* by qPCR. Botrytis *β-tubulin 2*

expression was also quantified. The expression levels were normalized to *Ubiquitin5* expression. Data

represent mean ± SD. The experiments were repeated three times with similar results.

Figure 7. SBT3.3 and Pro-PME17 are delivered in the apoplast through a conventional protein

secretion pathway. Confocal microscopy images of epidermal cells of *Nicotiana tabacum* adult leaves

showing the co-expression of SBT3.3-YFP **(A-J)** or Pro-PME17-GFP **(K-T)** with different markers of

subcellular compartments (**B, L:** RFP-HDEL, endoplasmic reticulum; **E, O**: ST52-mCherry, Golgi

stacks; **H, R**: XTH11-RFP, cell wall) at the indicated hours post infiltration. Plasmolyzed cells (1 M

NaCl) expressing SBT3.3-YFP or Pro-PME17-GFP and the plasma membrane marker pm-rk show the

green fluorescent cell wall and the retracted red fluorescent plasma membrane (**J, T**). Scale bars = 10

µm **(A-F)**, 20 µm **(G-I, M-S)**, and 5 µm **(J, T).**

Figure 8. Pro-PME17 and SBT3.3 colocalize in the cell wall and their colocalization is favoured

over time. Confocal microscopy images of epidermal cells of tobacco adult leaves co-expressing Pro-

PME17-RFP **(A and B)** and SBT3.3-YFP **(C and D)** at the indicated hours post transformation.

Merged images showing the co-localization of SBT3.3-YFP and Pro-PME17-RFP in the cell wall.

Overlap coefficient (OC, after Manders) and R (Pearson's correlation coefficient) values are reported in

each merged image. Chlorophyll auto-fluorescence within the chloroplasts is depicted in blue. Scale

bars = 20 µm

Figure 1. The lack of SBT3.3 and SBT3.5 expression in Arabidopsis does not alter rosette growth and cell wall composition.

A Ruthenium Red

De-methylesterified Pectin

B WT *sbt3.3-1* *sbt3.5-1*

C

PME activity
(mU/µg of total proteins)

□ WT
□ *sbt3.3-1*
□ *sbt3.3-2*
■ *sbt3.5-1*
■ *sbt3.5-2*

Mock   *Botrytis*

Figure 2. SBT3.3 and SBT3.5 regulate PME activity during disease

Figure 3. The lack of *SBT3.3 and SBT3.5* expression reduce Arabidopsis resistance to *Botrytis cinerea*

Figure 4. The SBT3.3 overexpression in Arabidopsis does not alter rosette growth and cell wall composition.

Figure 5. The SBT3.3 overexpression overactivates PME activity restricting *Botrytis* infection.

Figure 6. *SBT3.3* **overexpression enhances immune responces**

Figure 7. SBT3.3 and Pro-PME17 are delivered in the apoplast through a conventional protein secretion pathway.

Figure 8. Pro-PME17 and SBT3.3 colocalize in the cell wall and their colocalization is favoured over time.