Hitchhiking of bactericidal/permeability-increasing protein-like gene with the fibromelanosis
locus in Kadaknath black-bone-chicken

Short title: Kadaknath-specific hitchhiking of BPIL with the FM locus

Sagar Sharad Shinde[1], Ashutosh Sharma[1], Nagarjun Vijay[1]

[1]Computational Evolutionary Genomics Lab, Department of Biological Sciences, IISER

Bhopal, Bhauri, Madhya Pradesh, India

*Correspondence: nagarjun@iiserb.ac.in

42 **Keywords**: Kadaknath, black-bone chicken, genetic linkage, fibromelanosis, *Fm* locus

Abstract

Black-bone-chicken (BBC) meat is popular for its distinctive taste and texture. A complex chromosomal rearrangement at the fibromelanosis (*Fm*) locus on the 20th chromosome results in increased expression of the endothelin-3 (*EDN3*) gene and is responsible for melanin hyperpigmentation in BBC. We use public long-read sequencing data of the silkie breed to resolve high-confidence haplotypes at the *Fm* locus spanning both Dup1 and Dup2 regions and establish that the *Fm_2* scenario is correct of the three possible scenarios of the complex chromosomal rearrangement. The relationship between Chinese and Korean BBC breeds with Kadaknath native to India is underexplored. Our data from whole-genome re-sequencing establishes that all BBC breeds, including Kadaknath, share the complex chromosomal rearrangement junctions at the fibromelanosis (*Fm*) locus. We also identify two *Fm* locus proximal regions (~70Kb and ~300 Kb) with signatures of selection unique to Kadaknath. These regions harbor several genes with protein-coding changes, with the Bactericidal/permeability-increasing-protein-like (*BPIL*) gene having two Kadaknath-specific changes within protein domains. Our results indicate that protein-coding changes in the *BPIL* gene hitchhiked with the *Fm* locus in Kadaknath due to close physical linkage. Identifying this *Fm* locus proximal selective sweep sheds light on the genetic distinctiveness of Kadaknath compared to other BBC.

Introduction

Domestication of chicken during the Neolithic period involved a complex pattern of interbreeding with various jungle fowl species[1–4]. After domestication, chicken has spread worldwide and occurs as commercial, exotic, and indigenous village breeds. Humans use chicken as a research model due to their physiology and behaviour[5], as game fowl and for religious reasons, or more commonly for egg or meat production[6]. Chicken is the most preferred source of meat for humans due to its easy availability and affordability[7]. Hence, understanding the genetics of meat features is commercially relevant[8]. Although understudied, indigenous village chicken breeds with unique properties provide an opportunity to understand the genetics of meat. For instance, the black-bone chicken (BBC) is a delicacy due to its texture, color, firmness, flavor, and use in traditional medicine[9,10]. The black color of BBC results from melanin deposition throughout the body, i.e., melanin hyperpigmentation or fibromelanosis caused by the *Fm* allele[11]. Bateson and Punnet were the pioneers in identifying the autosomal dominant *Fm* allele[12]. Modern studies found a

chromosomal rearrangement on chromosome 20 is involved in the *Fm* locus[13,14].
The overexpression of the endothelin-3 (*EDN3*) gene located within the *Fm* locus is
responsible for hyperpigmentation seen in BBC[13,15,16].

BBC breeds have anti-fatigue and anti-hypoxic abilities, with its meat having antioxidant
properties[9,17], high carnosine[18,19], and lower fat and cholesterol content[10,20]. Some
BBC breeds also have local adaptations. For instance, the Korean Ogye has improved fetal
viability and innate immunity against microbial and viral infections[21]. BBC breeds occur
globally and have distinct names, such as Ayam Cemani (Indonesia), Black H'Mong
(Vietnam), Tuzo (Argentina), Svarthöna (Sweden)[14,22], Yeonsan Ogye (Korea)[23],
and Thai BBC (Thailand)[24]. China has a high diversity of BBC breeds, including Silkie,
Jiangshan, Lueyang, Sichuan, Xingwen, Yugan, Dehua, Jinhu, Muchuan, Wumeng, Yanjin,
Xichuan, Tuanfu, Wuliangshan, Emei and Miyi fowl[9,25–28]. It is unclear whether all BBC
breeds share the same rearrangement at the *Fm* locus, as independent structural variants can
produce similar phenotypes[29]. India has a single breed of BBC, commonly known as
Kadaknath[19].

Kadaknath is considered native to the Jhabua, Alirajpur, and Dhar districts of Madhya
Pradesh, but its farming has recently spread across India [30,31]. The Kadaknath breed has
long been documented as a distinctive indigenous Indian breed and is also called Kali Masi or
Karaknath [32]. At least three distinctive phenotypes (i.e., Jetblack, Pencil, and Golden)
occur within the Kadaknath breed[33]. In Jetblack, all the body parts like plumage, comb,
internal organs, eyes, skin, beak, shank, and claw are entirely black, whereas Pencil and
Golden have white and golden color patches on the plumage, respectively[33]. Several
unique characteristics, such as earlier egg-laying maturity, high protein content, better disease
resistance, and adaptation to the local environment, are attributed to Kadaknath chicken
[19,30,33–41]. Understanding the genetics behind these traits will help establish the
uniqueness of Kadaknath and guide de-extinction efforts and breeding programs.

Despite its immense popularity and commercial value, the genomics of the Kadaknath
chicken breed has received limited attention. Therefore, further research and genomic
analyses are required to understand adaptations in this breed and its genetic history. The aims
of this study are : (1) Perform whole-genome re-sequencing of Kadaknath and evaluate its
relationship with other black-bone and non-black-bone chicken breeds. (2) Assess whether

Kadaknath and other BBC breeds share a common origin for the *Fm* locus by comparing the

chromosomal rearrangement junction. (3) Evaluate how the BBC breeds dispersed to various

parts of Southeast Asia. (4) Use population genetic statistics to identify signatures of

selection in Kadaknath compared to other BBC breeds. Our results offer an example of the

co-selection of traits regulating immune response, meat quality, and plumage color providing

new approaches to improve breed quality and productivity in chicken simultaneously.

Materials and Methods

Population sampling

The study was approved by the Institutional Ethics Committee (IEC) of the Indian Institute of

Science   Education   and   Research,   Bhopal,   vide   reference   number

IISERB/IEC/Certificate/2018-11/03 dated 8[th] June 2018. We purchased the meat of six

individuals (two individuals each from Jetblack, Pencil, and Golden morphs (**Fig. S1**)) of

Kadaknath from an FSSAI (Food Safety and Standards Authority of India) licensed shop in

Bhopal, Madhya Pradesh, India. We procured two other individuals with a black-bone

phenotype from the same FSSAI-licensed shop to examine hybrids. One had a Golden-Pencil

external appearance, and another was completely white on the exterior. We also sampled two

non-black-bone chicken from the same shop to determine the genetic relationship of

Kadaknath with native-village chicken and broiler reared in the same poultry (see **Fig. S1**).

To reduce the chances of inter-breeding and have relatively pure Kadaknath samples, we

obtained three (one individual each from Jetblack, Pencil, and Golden morphs) additional

Kadaknath breed chickens from Jhabua, Madhya Pradesh. Whole-genomic DNA with high

purity and quality was extracted from the tissue samples using DNeasy Blood & Tissue Kits

(QIAGEN). We generated >25x coverage whole-genome short-read paired-end data (using

Illumina Novaseq) for all 13 individuals sampled (see **Table S1**).

To evaluate the relationship of Kadaknath chicken globally, we compared our dataset with

publicly available chicken re-sequencing data. High-quality re-sequencing datasets of 88

chicken individuals from other BBC, commercial chicken lines, and other distinctive chicken

breeds with a coverage >20x were selected and obtained from ENA (Europen Nucleotide

Archive) and (KNABIC) Korean National Agricultural Biotechnology Information Center

(https://nabic.rda.go.kr/). Out of 88 individuals, 22 are BBC, which we considered for further

analysis. Hence, we analyze a dataset of 101 individuals (88 public + 13 sequenced as part of

this study) from different breeds (see **Table S2** for more details)[21,23,27,42–47].

Read mapping, variant calling, and phylogeny

We mapped the paired-end raw reads of 101 individuals to the chicken genome assembly

(genome assembly version is Gallus_gallus.GRCg6a) using the BWA[48] (Burrows-Wheeler

Aligner) v0.7.17-r1188 read mapper with default parameters. We added the read group

information using Picard tools and removed duplicate reads in all 101 individual bam files

(https://github.com/broadinstitute/picard). We performed the variant calling using

FreeBayes[49]with different quality control flags such as --min-alternate-count -C 10, --min-

mapping-quality -m 20, --min-base-quality -q 20, and --min-coverage 10. Similarly, we used

the bcftools[50],[51] with mapping quality flags such as --mapping quality -C 50, --min base

quality -Q 20, and --min mapping quality -q 20 for robust variant calling. We removed the

indels from variant calls using vcftools[52] with the -remove-indels flag and extracted the

common SNPs using both SNP callers for retaining reliable SNP calls. The Single Nucleotide

Polymorphisms (SNPs) identified by both variant callers (bcftools v1.9 and FreeBayes

v1.0.0) were used for subsequent analysis. To identify the effect of genetic variants, we used

the snpEff[53]. Common SNPs from both vcf files were used to construct phylogeny for 34

and 101 individuals using SNPhylo[54] v20180901, based on a maximum-likelihood tree

with 1000 bootstrap values. We excluded the scaffolds, Z, W, and MT chromosomes in

phylogeny analysis. The local phylogeny for Dup1 and Dup2 regions was generated using the

vk phylo command (using both NJ and UPGMA methods) implemented in VCF-kit[55]. The

Mt genome haplotype median-joining networks were constructed using PopART[56] and

SplitsTree[57].

Fm **locus junction identification**

The *Fm* locus consists of a complex chromosomal rearrangement composed of two different

non-paralogous regions (Dup1 (~127 Kb) and Dup2 (~170 Kb)) separated by an intermediate

(Int) region. Dup1 and Dup2 regions are both duplicated and are involved in a complex

rearrangement consisting of two junctions (**A**) Dup1 + (inverted Dup2) and (**B**) (inverted

Dup1) + Dup2. To identify the base-pair level positions of Dup1, Dup2, and Int regions, we

compared the short-read coverage of black and non-black chicken in 1Kb windows along

chromosome 20. We used the makewindows command of bedtools[58] (v2.26.0) to create

1Kb non-overlapping windows along chromosome 20. The number of reads in each 1Kb

window was calculated using the bedtools coverage command. We shortlisted adjacent

windows with drastically different read coverage in black individuals but not in non-black

individuals. The base-pair level coordinates of Dup1 and Dup2 in the Gallus_gallus.GRCg6a

genome was narrowed down further using coverage estimates in 1bp windows. Dup1 starts at

20:10766772 and ends at 20:10894151. Dup2 occurs further along the chromosome and starts

at 20:11306686 and ends at 20:11477501. The read coverage at these junctions consistently

differed between all black-bone and non-black-bone chicken. In addition to the Dup1, Dup2,

and Int regions, we defined ~500 Kb flanking regions as Flank1 (20:10263555-10766771)

and Flank2 (20:11477502-11980000).

Black-bone-specific *Fm* locus junction

The junction between the rearranged regions in the BBC does not occur in the

Gallus_gallus.GRCg6a genome (**Fig. S2, 3**). Hence, we searched for the *Fm* locus junction in

the Korean BBC PacBio data from ENA (SRR6189090). Based on our search of PacBio

reads that mapped to the Gallus_gallus.GRCg6a genome, we shortlisted reads that

simultaneously aligned to two of the five (Flank1, Dup1, Int, Dup2, and Flank2) genomic

regions we have defined. We found several reads spanning Flank1-Dup1, Dup1-Int, Int-

Dup2, and Dup2-Flank2, which is unsurprising because these regions are adjacent.

We also found five reads (SRR6189090.111279, SRR6189090.56386, SRR6189090.880702,

and SRR6189090.387043 and SRR6189090.54824) that span across both Dup1 and Dup2.

Dup1 and Dup2 are far apart in their genomic location in Gallus_gallus.GRCg6a assembly.

Hence, these reads support a rearrangement that leads to junctions between these two regions.

Three of these five reads (SRR6189090.111279, SRR6189090.880702, and

SRR6189090.387043) support the junction Dup1 + (Inverted Dup2) (i.e., START-DUP1-

END-END-DUP2-START), and the other two reads (SRR6189090.54824 and

SRR6189090.56386) support the junction (inverted Dup1) + Dup2 (i.e., END-DUP1-

START-START-DUP2-END). The read coverage at these junctions consistently differed

between all black-bone and non-black-bone chicken. We identified 34 out of 101 samples

analyzed as BBC based on the read coverage across the *Fm* locus junctions.

Long-read-based resolution of haplotypes at Dup1 and Dup2 regions

Recently, a new high-coverage multi-platform genomic public dataset for the silkie black-bone chicken became available on the ENA (Europen Nucleotide Archive) as part of the Bioproject# PRJNA805080 (we thank China Agricultural University for publishing this data). With a ~65X (Nanopore) and >660X coverage (PacBio) of the same silkie individual, this dataset is well suited to resolve the haplotypes at the *Fm* locus and identify the exact order of rearrangement. Upon visual inspection of the long-read alignments at the Flank-1-Dup1 and Dup2-Flank2 junctions region, three heterozygous sites were recognized at the Dup1 start region, and Dup2 end region, potentially separating the two haplotypes of Dup1 and Dup2. A detailed blastn-based search of reads from both haplotypes anchored one haplotype to the Flank1 + Dup1 junction and another haplotype to the (inverted Dup2) + Dup1 junction. The Flank1 + Dup1 junction containing haplotype is referred to as Dup1 haplotype-1 (D1H1) and contains the allele T at position 10766895. The (inverted Dup2) + Dup1 junction containing haplotype is referred to as Dup1 haplotype-2 (D2H2) and contains the allele A at position 10766895. We extended these haplotypes toward the end of Dup1 by relying upon overlapping long-reads at haplotype-defining sites. To ensure the robustness of our approach, we required that more than ten sequencing reads should support haplotype-specific alleles at each adjacent site (see **Table S3**).

In the Dup2 region, one haplotype was anchored to the Dup2+ Flank2 junction and another haplotype to the Dup2 + (inverted Dup1) junction. The Dup2 + Flank2 junction containing haplotype is Dup2 haplotype-1 (D2H1) and contains the allele A at position 11476819. The Dup2 + (inverted Dup1) junction containing haplotype is Dup2 haplotype-2 (D2H2) and contains the allele G at position 11476819. We extended these haplotypes toward the start of Dup2 by relying upon overlapping long-reads at haplotype-defining sites. We identified 25 well-resolved, high-confidence haplotype-defining sites spanning from the end of Dup2 to the beginning of Dup2 (see **Table S3**). Haplotype-1 of Dup2 contains reads spanning from the end of the Int region to the start of Flank2, and Haplotype-2 of the Dup2 region spans from the end of the Haplotype-1 of Dup1 (D1H1) region to the beginning of Haplotype-2 of Dup1 (D1H2) region.

Principal component analysis and admixture

Principal components analysis (PCA) was performed using PCAngsd[59] based on genotype likelihood estimates from ANGSD[60] (Analysis of Next Generation Sequencing Data) v0.935. PCA was done both genome-wide and *Fm* locus region-wise (Flank1, Dup1, Int,

Dup2, and Flank2) for the 34 black-bone and all 101 chicken individuals**.** We used several

flags in ANGSD for population structure analysis as follows: -GL 2, -doMaf 1, -minMapQ

30, -minQ 20, -doGlf 2, and -SNP_pval < 1e-6. Genotype likelihood values from ANGSD

were used to identify principal components using PCAngsd and genome-wide admixture

proportions using NGSadmix[61]. NGSadmix was run for different values of K from K1 to

K10 using each K with 15 iterations with flags -minMaf 0.05 and -minInd as specified. We

used the online Clumpak[62] web server to find the best K.

Population genetic analysis

Chinese black-bone breeds have diverged to differing extents from Kadaknath and have

limited sample sizes. Hence, we combined the individuals from closely related XBBC,

LCEM, and LCMY breeds into a single population representative of Chinese black-bone

(CHIN) chicken and JETB, PENC, and GOLD into another population representative of

Kadaknath (KADK) (see **Table S4**). We calculated population genetic statistics genome-wide

to identify signatures of selection. However, to avoid false positives, we excluded genomic

regions (50 Kb windows with <80 percent callable sites) with poor callability. We used the

CallableLoci walker of GATK[63] on the bam files to quantify callability. For identifying

callable regions from mapped bam files for CHIN and KADK individuals, we used

GATK with different flags such as --minMappingQuality 20, --minBaseQuality 20, --

minDepth 10, --minDepthForLowMAPQ 20, and --maxFractionOfReadsWithLowMAPQ 20.

Furthermore, we filtered 50Kb windows with >0.1 repeat element fraction to rule out the

possibility of false positives (see **Fig. S4-6**). We used stringent coverage and quality criteria

(-GL 2, -dosaf 1, -baq 1, -C 50, -setMinDepthInd 6, -minInd 3, 4 or 9, -minMapQ 30, -minQ

20, and -doCounts 1) in ANGSD to calculate all population genetic statistics. The list of

individuals in each population and the population pairs compared is in **Table S4**. We found

that at least 20,000 of the 21,659 50Kb windows had sufficient high-quality data in all

populations.

We used the folded site frequency spectrum (SFS) approach implemented in ANGSD to

calculate genome-wide population-specific estimates of π (the average pairwise differences),

Watterson's θ (the average number of segregating sites), τ (Tajima's D), and Fu and Li's D.

We estimated inter-population genomic differentiation (F_{ST}) and divergence (D_{xy}) using

ANGSD   and   popgenWindows   python   script

(https://github.com/simonhmartin/genomics_general) for each population pair. The F_{ST}

estimates from the two methods were highly correlated (Pearson's ρ= 0.98, p-value < 2.2 e-

16) (see **Fig. S7, 8**). We defined F_{ST} outlier regions as 50Kb windows in the top 1 percent of

the genome-wide estimates and merged adjoining windows using bedtools. Similarly, the

50Kb windows in the top 10 percent of the genome-wide estimates of D_{xy} were deemed to

have high levels of divergence. We identified fixed sites as those sites with $F_{ST} > 0.9$. We

estimate haplotype-based statistics iHS and XP-EHH in the rehh[64] R package with option

polarised=FALSE using genotype data phased with SHAPEIT[65].

Results

Whole-genome re-sequencing

We generated >25X coverage whole-genome re-sequencing Illumina data for 13 chicken

individuals from India. Our collection includes 9 Kadaknath samples as representatives of all

three extant morphs Jetblack (JETB n=3), Golden (GOLD n=3), and Pencil (PENC n=3). We

also sampled one individual with a golden-pencil-like phenotype (GOPE n=1), one with

complete white plumage with black-bone (CROS n=1), one non-black-bone individual where

plumage was black (NONB), and one individual of broiler breed (**Fig. S1** and **Table S1**). We

compared these Kadaknath chicken genomes with public re-sequencing datasets of chicken

breeds. The sampling location (see **Fig. 1a** for black-bone chicken) of individuals analyzed in

this study is spread across Asia (for more detail, see **Table S2,** which includes the complete

list of all chicken breeds examined). A detailed map of China depicts the locations of all the

Chinese black-bone breeds (**Fig. S9**).

Kadaknath is a distinct breed of black-bone fowl

The major axes of genetic variation in BBC (PC1:18.9% and PC2:13%) separated the

Kadaknath (KADK), Yeonsan Ogye (YOSK), and Chinese BBC (see **Fig. 1b** and **Fig. S10**).

We observed that all Kadaknath individuals form a single cluster distinct from other black-

bone (from Indonesia, China, and South Korea) (**Fig. 1b** and **Fig. S10**). Similarly, in the

genome-wide PCA analysis of 101 individuals, Kadaknath forms a separate cluster from

other commercial, native, and BBC breeds in different PC comparisons (see **Fig. S11**).

Admixture analysis suggests Kadaknath has sub-structure but remains distinct from the other

black-bone breeds at best K=7 (see **Fig. 1c** and **Fig. S12, 13**). In Chinese BBC, the admixture

analysis identified three groups that support the clustering pattern in PCA, while YOSK

forms a separate cluster (see **Fig. 1c** and **Fig. S10**). We observed lower genetic diversity (π =

0.002, θ = 0.002) in YOSK, while Kadaknath and Chinese BBC have comparable genetic

diversity (π = 0.004 and 0.003, θ = 0.003 in both) (**Fig. S14a** and **b**). In contrast to the nuclear

genome, the mitochondrial genome haplotype network did not separate Kadaknath from other

BBC breeds (**Fig. S15, 16**). The *Fm* locus region on chromosome 20, which codes for the

black-bone phenotype, is the defining feature of all BBC. Even after excluding chromosome

20, the PCA of the remaining chromosomes finds that Kadaknath is genetically distinct from

other BBC breeds (**Fig. S17**). Hence, the genetic distinctiveness of Kadaknath is spread

across the entire genome.

All black-bone chicken breeds share the same rearrangement junctions at *Fm* locus

In the non-black chicken, the Dup1 (~127 Kb), Int (~412 Kb), and Dup2 (~170 Kb) regions

occur in a single copy, are arranged sequentially, and are flanked by Flank1 (~500 Kb) and

Flank2 (~500 Kb) regions (see **Fig. 2a**). While the Dup1 region contains five protein-coding

genes (*EDN3*, *ZNF831*, *SLMO2*, *ATP5E*, *TUBB1*), the Dup2 region probably consists of only

long-non-coding RNA genes. All non-black chickens have a single copy of this region,

referred to as **N* locus (see **Fig. 2a**). The corresponding locus in black chickens is known as

the *Fm* locus. The *Fm* locus consists of three different non-paralogous regions (Dup1, Int, and

Dup2) that form a complex chromosomal rearrangement in which both the Dup1 and Dup2

regions are duplicated, giving rise to two junctions (**A**) Dup1 + (inverted Dup2) and (**B**)

(inverted Dup1) + Dup2. Although the exact ordering of these regions in the rearrangement is

not conclusively established, both regions are known to be duplicated due to these regions

having a sequencing coverage that is twice the genomic average [13]. The presence of Dup1

+ (inverted Dup2) and (inverted Dup1) + Dup2 junctions has been verified in several black

chicken breeds [13,14,23]. Based on this information, three possible scenarios have been

proposed by earlier studies (see **Fig. 2b**). The **Fm_2* scenario is supported based on crosses

between the black bone and non-black bone chicken [13]. However, both **Fm_2* and **Fm_3*

scenarios require two rearrangement events compared to a single rearrangement event needed

for the **Fm_1* scenario [23].

Distinguishing between these three (**Fm_1*, **Fm_2*, and **Fm_3*) scenarios requires long-

range connectivity information such as long-read sequencing data (PacBio, Nanopore,

Synthetic long reads, etc.), Hi-C (high-resolution chromosome conformation capture)[66,67],

or Optical Mapping[68]. Both **Fm_1* and **Fm_2* contain the same adjacent regions for Dup1

(defined as Haplotype-1 and Haplotype-2 in **Fig. 2b**) inconsistent with **Fm_3*. Hence, long-

range information that can span the entire ~127 Kb Dup1 region in a single read will be able

to distinguish between (*Fm_1* and *Fm_2*) vs. *Fm_3*. However, to differentiate between

Fm_1 and *Fm_2* scenarios, we need to span the entire ~170 Kb Dup2 region in a single

read. Spanning the Dup1 or Dup2 regions with PacBio (average read lengths of 10 to 25 Kb)

or Nanopore (average read lengths of 10 to 30 Kb) technologies with a single read is

challenging but not impossible as reads longer than 1 Mb are possible [69,70]. Without such

individual reads that can span the entire region, it is possible to perform read-based phasing

to infer haplotypes that extend to distinct junctions [71].

Using a public long-read dataset, we have inferred distinct high-confidence Dup1 and Dup2

haplotypes (supporting *Fm_2* scenario). Dup1 haplotypes are anchored by (Flank-1

end)/(Dup2 start) at the Dup1 start and (Int start)/(Dup2 end) at the Dup1 end (**Fig 3**). Each

haplotype can be distinguished at 24 sites (see **Table S3**) along the Dup1 region. Each

haplotype-defining site shares more than ten long reads with haplotype-specific alleles at

adjacent positions. Haplotype-1 (see **Fig. 3a** and **Fig. S18a**, **b**) spans Dup1 from Flank-1 to

haplotype-2 Dup2 (D2H2) end, and haplotype-2 (see **Fig. 3b** and **Fig. S18a**, **b**) spans Dup1

from haplotype-2 Dup2 (D2H2) start to Int start. Our inference of Dup1 haplotype-1 (i.e.,

Flank-1 + Dup1 + (inverted Dup2)) and haplotype-2 (i.e., (inverted Dup2) + Dup1 + Int)

sequences rule out the *Fm_3* scenario, suggesting that either *Fm_1* and *Fm_2* scenario is

possible. To verify the correct arrangement, we used the read phasing approach on the Dup2

region, and we could distinguish the two haplotypes at the Dup2 region based on 25 sites (see

Table S3). Haplotype-1 of the Dup2 region (see **Fig. 4a** and **S19a, b**) spans Dup2 from the

end of Int to that start of Flank2, and haplotype-2 (see **Fig. 4b** and **S19a, b**) spans Dup2 from

haplotype-2 Dup1 (D1H2) start to haplotype-1 Dup1 (D1H1) end. Our inference of

haplotype-1 Of Dup2 (i.e., Int+ Dup2 + Flank2) and haplotype-2 (i.e., start of Dup1 +

(inverted Dup2) + Dup1 end) sequences rule out the *Fm_1* and *Fm_3* scenarios, suggesting

that *Fm_2* scenario is correct (see **Fig. S20**).

Conclusive inference regarding which scenario is present in BBC has been extremely

challenging due to the large (~1Mb) size and complexity of the rearrangement. However, in

the case of Kadaknath, it has not even been established whether the Dup1 + (inverted Dup2)

and (inverted Dup1) + Dup2 Junctions identified in other BBC are present. We have

compared the normalized short-read coverage at these junctions to evaluate whether all black-

bone chicken breeds share the same rearrangement junctions at the *Fm* locus. The normalized

short-read coverage in black-bone chicken (**Fig. 5a-e**) abruptly increases at Dup1 and Dup2

regions, while no such increase occurs in non-black-bone (**Fig. 5f**) chicken. The drastic change in coverage at Dup1 and Dup2 boundaries occurs at the same base in all black-bone chickens (see **Fig. S21-22a-d**). In the case of non-black-bone chickens, no change in coverage occurs at this position (see **Fig. S23**). We also identified the black-bone-specific *Fm* locus junctions Dup1 + (inverted Dup2) and (inverted Dup1) + Dup2 using published PacBio data (see **Fig. 6a, b,** and **Methods**). Short-read coverage spanning these black-bone-specific *Fm* locus junctions is present in black-bone and missing in non-black-bone chicken (**Fig. S24-26**). We can infer that black-bone chicken defining *Fm* locus originated through a single event based on coverage and rearranged junction sequence.

Crossbreeding with native and commercial breeds affects the genome-wide patterns of genetic variation in black-bone breeds (see **Fig. S27-29**). Hence, we focussed on the *Fm* locus to investigate the history of the BBC breeds. The F_{ST} between black-bone and non-black-bone chicken along chromosome 20 is elevated at the Dup1 and Dup2 regions (see **Fig. S30**). However, other population genetic parameters, such as genetic diversity (π and θ) and divergence (D_{xy}), lack any prominent signatures (**Fig. S31-33**). As evident from the elevated F_{ST}, the major axis of genetic variation (assessed using local PCA) in Dup1 and Dup2 regions separates the BBC from non-black-bone chickens (**Fig. S34**). A phylogenetic tree of the SNPs from Dup1 and Dup2 regions also largely separates the BBC breeds from non-BBC breeds (**Fig. S35a** and **35b**). Our evaluation of the genetic differentiation (F_{ST}) landscape between BBC breeds found reduced F_{ST} at Dup1 and Dup2 regions compared to the genomic background (see **Fig. S36**). However, comparing individual BBC breeds with non-black chicken breeds showed the opposite pattern with elevated F_{ST} at Dup1 and Dup2 regions (see **Fig. S37, 38**). Hence, the patterns of genetic differentiation (F_{ST}), local PCA, and phylogenetic tree also support a common origin of the *Fm* locus in all BBC breeds.

Isolation by distance pattern suggests dispersal between India and China

Our analysis discovered that the rearrangement junctions in all BBC are identical to the one in Kadaknath and strongly support a common origin for the *Fm* locus. An independent origin for the *Fm* locus would mean separate rearrangement events have created identical junction sequences. Given the lack of repeat sequences at the junctions, such independent origins seem highly unlikely and are not consistent with the genetic relatedness of BBC and non-BBC breeds at the *Fm* locus. Hence, the current distribution of BBC breeds across Southeast Asia needs an explanation. We hypothesized (see a schematic of the proposed scenario in

Fig. S39) that the dispersal of BBC occurred after a common origin for the *Fm* locus,
followed by recent crossbreeding with native and commercial breeds.

To test our hypothesis and delineate the dispersal route of different black-bone breeds, we
first evaluated if a pattern of isolation by distance (IBD) is prevalent. The IBD plot shows a
consistent increase in genetic distance with an increase in geographic distance (see **Fig. S40**,
mantel's r=0.64, p-value=0.0002). The IBD pattern persisted even after we repeated the
analysis using ANGSD/variant-call-based estimates of nucleotide differentiation (F_{ST}) after
excluding various populations to avoid errors due to auto-correlation and biased estimates
from isolated populations (see **Fig. S41-47**). Among the black-bone breeds, Tibetan black-
bone (TBTC) and Sichuan black-bone (LCTMJ) chickens are genetically closest to
Kadaknath (mean F_{ST}~0.17). The other black-bone breeds that are geographically more
distant from Jhabua (LCEM, mean F_{ST}=0.21; XBBC, mean F_{ST}=0.22; LCMY, mean F_{ST}=0.3
and YOSK, mean F_{ST}=0.36) occur at increasing genetic distances from Kadaknath (see **Table**
S5). The pattern of IBD spanning India and China suggests potentially human-mediated
dispersal. We lack conclusive evidence to identify the direction of dispersal. However,
analysis of our dataset, even after excluding alleles found in dbSNP[72] (**Fig. S48, 49**), found
that Kadaknath has almost twice the number of private alleles than Chinese and Korean BBC
breeds and suggests an India-to-China dispersal. More extensive fine-scale sampling may
provide a definitive answer regarding the direction and timing of the dispersal.

Genome-wide signatures of selection in Kadaknath chicken

We screened the genome for selection signatures to identify all Kadaknath-specific regions
by comparing Kadaknath with Chinese BBC. We also compared Kadaknath and Chinese
BBC (XBBC) with the Korean Yeonsan Ogye (YOSK) to identify population-specific
signatures. Despite the sizeable geographic separation, the genome-wide mean F_{ST} between
KADK and CHIN is 0.16, making it amenable to identifying selection signatures (see **Fig. 7a**
and **Table S6**). A pairwise comparison between KADK and CHIN revealed 137 genic
regions in the top 1% F_{ST} windows (see **Table S7** and **Fig. S50-82**). We further shortlisted
candidates using the additional criteria that genetic diversity (π and θ) and integrated
haplotype score (iHS) should be strikingly different between KADK and CHIN. Two
prominent signatures from these shortlisted regions indicative of selective sweeps in the
KADK breed are evident on the 20th chromosome in the vicinity of the Dup1 region. The first
region (R1) of ~300 Kb is ~160 Kb before Dup1, and the second region (R2) of ~70 Kb is
~0.7 Mb before Dup1. In both regions, the genetic diversity (π and θ) and Tajima's D (τ) are

strongly reduced in KADK compared to CHIN (see **Fig. 7b** and **Fig. S69**). The elevated iHS

in KADK compared to CHIN and the positive value of extended haplotype homozygosity

(XP-EHH) supports a selective sweep in KADK. This region also occurs in the top 10%

genome-wide genetic divergence (D_{xy}) windows (see **Fig. S83**).

In the first region (R1), Deoxynucleotidyltransferase Terminal Interacting Protein 1

(*DNTTIP1*), Acyl-CoA Thioesterase 8 (*ACOT8*), and solute carrier family 12 member 5

(*SLC12A5*) genes contain non-synonymous changes. The TELO2-interacting protein one

homolog (*TTI1*), which occurs near the first sweep region (R1), also has a non-synonymous

change (**Fig. 7c** and **Table S8, 9**). The second sweep region (R2) contains three genes with

non-synonymous changes: Histocompatibility Minor 13 (*HM13*), Bactericidal/permeability-

increasing protein-like (*BPIL*), and Targeting protein for Xklp2 (*TPX2*). The most striking

differences were in the *BPIL* gene, which has three fixed sites with non-synonymous

changes. Of these changes, two alleles are unique to the KADK population (see **Fig. S84**).

The first non-synonymous change (C**G**G->C**A**G, R->Q) occurs at position 69 in exon-2. The

second non-synonymous change (**G**AG->**A**AG, E->K) is within the first BPI superfamily

protein domain at position 159 in exon-4. The third and last non-synonymous change (C**G**C-

>C**A**C, R->H) is in the second domain at position 405 in exon-11. We also found that *HM13,*

SLC12A5, and *DNTTIP1* have novel Kadaknath-specific changes.

The Korean Yeonsan Ogye (YOSK) breed is closer to the Chinese XBBC (mean F_{ST} = 0.26)

than to Kadakanth (mean F_{ST}=0.3). However, the genome-wide mean F_{ST} is still low enough

to distinguish selection signatures. A comparison of YOSK with KADK confirmed that the

R1 and R2 regions are specific to Kadaknath (see **Fig. S85** and **Table S6**). Surprisingly, our

comparisons with YOSK identified a third region (R3) of ~570 Kb, which is ~2.16 Mb before

Dup1 with a sweep specific to YOSK (see **Fig. S86** and **Table S9**). Local PCA of the *Fm*

locus, R1, and R2 regions demonstrates that Flank1, R1, and R2 regions have changed only

in Kadaknath and not in other BBC (**Fig. S87-89**). A careful examination of the entire

chromosome 20 failed to identify any other sweep regions (see **Fig. S90**). Similar to the

sweep in KADK, we found two genomic regions with signatures of a sweep in the CHIN

population (see **Table S9**). The sweep region on Chr 4 contains the *PCDH7* gene, and the

region on Chr 9 contains the *COL4A3* and *MFF* genes (see **Fig. S53, 58**). However, none of

these genes have any non-synonymous fixed differences.

Discussion

All BBC breeds share a common origin of the *Fm* locus

As part of this study, we have generated the first whole-genome dataset of the Kadaknath breed spanning all three morphs. Hence, by comparing our data with public datasets, we can evaluate the relationship between all BBC breeds. Our comparison of the *Fm* locus junction region in genome sequencing data from BBC breeds conclusively establishes a common origin for the complex chromosomal rearrangement[14]. The following four lines of evidence support this conclusion: (1) The short-read coverage along the *Fm* locus, (2) the high sequence identity of the rearranged junctions across the BBC breeds, (3) the local and phylogenetic relationship between chicken breeds at the *Fm* locus, and (4) the patterns of pairwise genetic differentiation between BBC and non-BBC breeds.

Spread of BBC across Southeast Asia

The earliest records of black-bone chicken have been found in the writings of Marco Polo, the Venetian explorer-writer who traveled through Asia [13]. The Compendium of Materia Medica or Bencao Gangmu, compiled and edited by Li Shizhen and published in the late 16th century, attributes various medicinal properties to BBC [73]. In Korea, black-bone chicken is thought to have had a royal connection, and "Dongui Bogam", a traditional Korean medical encyclopedia compiled & edited by Heo Jun in 1613, records the medicinal use of BBC [23]. While an earlier study notes that "In 1635 AD, the finding of chickens with black meat (typical of fibromelanosis, FM mutation) in Mozambique suggested direct introductions from India."[2] we could not find literature on how Kadaknath ended up in Jhabua, India.

Jhabua is close to the ancient port cities of Bharuch (260 Km) and Lothal (300 Km) on the west coast of India. Given the proximity to port cities and the prevalence of BBC only in Western India (and the spread of BBC to Africa), we suspected that Kadaknath may have traveled through a marine route. Ancient sea trade between India, Korea, and other parts of Asia are well documented [74]. Moreover, the movement of domesticated breeds during the period of colonialism was also facilitated by the common rule of large parts of Southeast Asia by various European powers [75].

Historical records are patchy, prone to error/obfuscation, and fail to provide conclusive information about the spread of BBC across Asia. Hence, we evaluated the relationship between BBC breeds and patterns of isolation by distance to infer the ancient dispersal route. Although each BBC breed has considerable genetic distinctiveness, population genetic analyses support the common heritage of all BBC breeds and reveal a trend of isolation by

distance (IBD). While IBD patterns are well established in Chinese native chicken
breeds[76], our dataset spans BBC breeds from India, China, Indonesia, and Korea. The
sample size of some BBC breeds in our analysis is limited, and the sampling does not cover
the entire geographic distribution of some breeds. However, our analysis achieves genome-
wide pan-BBC breed sampling by including all the major breeds from Asia.

The BBC chicken from Tibet is genetically closest to the Kadaknath breed and suggests that
the old Tibet-Nepal salt trade route or the maritime silk route may have facilitated the spread
of BBC. Interestingly, the Tibetan BBC is genetically more similar to Kadaknath than some
Chinese BBC breeds. Unfortunately, we lack BBC samples from Nepal. However, we believe
BBC poultry in Nepal is Kadaknath chicken, recently imported from India. Hence, the
ancestral stock of the BBC spread across Asia may not be currently available. Changes in
trade routes and the introduction of commercial poultry breeds limit our ability to trace the
historical prevalence of BBC breeds.

Did black-bone chicken originate in India?

The lack of data from India and the considerable interest in BBC breeds in Europe, China,
and Korea has meant that studies have focussed mainly on non-Indian BBC. Hence, Southern
China and Tibet are considered the source of all BBC breeds[25,77]. However, despite a
traditionally restricted geographic distribution, the Kadaknath breed has nucleotide diversity
comparable to Chinese BBC breeds and much higher than the Korean BBC. The number of
private alleles identified in Kadakanth is also greater than in Chinese and Korean BBC
breeds. Moreover, the mean genome-wide F_{ST} of 0.11 between Jhabua and Bhopal is
comparable to the differentiation between some Chinese BBC breeds. The presence of three
distinct morphs within Kadaknath chicken suggests phenotypic diversity derived either from
commercial breeds or existing variations within Kadaknath. The extent of phenotypic
diversity within BBC breeds from China, and Korea is not documented to allow a fair
comparison. The high genetic diversity in Kadaknath supports the potential origin of all BBC
in Jhabua, India. The export of black chicken from India to Africa in ~1600 AD also supports
that BBC was present in ancient India[2].

While we cannot conclude whether BBC had an Indian, Chinese, or more Southeast Asian
origin, our data suggest that any of these sources are plausible. More widespread geographic
sampling and analysis of allele-sharing patterns may provide a definitive answer regarding
the origin of BBC. Unlike the initial domestication of the chicken, which may have occurred

independently[2,47] in several locations, the BBC has a single origin linked to the

rearrangement at the *Fm* locus. Irrespective of the source of all BBC, we identify several

Kadaknath-specific genetic changes. Hence, the genetic distinctiveness of the Kadaknath

breed has long diverged from other BBC breeds and is a result of its unique heritage

sustained by the Bhil and Bhilala tribal communities of Madhya Pradesh. The beliefs and

practices of tribal communities may have contributed to the domestication and conservation

of Kadaknath[78].

Conclusive resolution of the complex chromosomal rearrangement

Earlier studies of the *Fm* locus have proposed three possible scenarios for the complex

chromosomal rearrangement. Although the *Fm_2* scenario was favored based on crosses

between BBC and non-BBC breeds, the correct scenario was not established by previous

studies as the genome assembly of this region is challenging. We use a haplotype phasing

approach relying upon published long-read datasets to conclusively resolve the correct

arrangement to be *Fm_2* scenario at the *Fm* locus. In the *Fm_2* scenario, the distal region

(Dup1+Int+Dup2+Flank2) is similar to the *N* arrangement found in non-BBC breeds.

However, the proximal region (Flank1+Dup1+(inverted Dup2)) has a very different

arrangement than the non-BBC one. In contrast to the *Fm_2* scenario, the proximal region

would be similar to the *N* arrangement in *Fm_3* scenario. In the *Fm_1* scenario, only the

first occurrence of Dup1 and last occurrence of Dup2 are similar to the *N* arrangement.

Hence, in the *Fm_2* and *Fm_3* scenarios recombination with *N* arrangement may be easier

(i.e., recombination is supressed only in the inverted ~127Kb (Dup1) or 170Kb (Dup2)

region) compared to * *Fm_1* scenario (i.e., recombination is supressed in ~709Kb (inverted

(Dup2+Int+Dup1)) region). The ease of recombination with *N* arrangement could explain

the prevalence of *Fm_2* scenario which requires two rearrangement events.

Selective sweep near *Fm* locus may be a consequence of linkage

Our analyses identified two genomic regions (R1 and R2) in the vicinity of the *Fm* locus with

prominent signatures of a selective sweep in the KADK population. The most promising

candidate gene in this region that may have been the focus of selection is the *BPIL* gene

which has accumulated three non-synonymous changes. The *BPIL* gene is part of the innate

immune defense system, which binds and neutralizes lipopolysaccharides (LPS) from the

outer membrane of Gram-negative bacteria[79–81]. In the chicken genome, five *BPI*-like

genes occur near *BPI*[82]*,* which help arrest bacterial growth, are prominent in phagocytosis

by neutrophils, and work as a bactericidal protein[83,84]. In chickens, defense against gram-
negative bacteria may be especially important as they lack the complement *C9* gene required
to effectively eliminate pathogens through the Membrane Attack Complex (MAC)
formation[85]. Notably, another BBC breed from Korea, YOSK, has signatures of selection
at the TLR4 gene involved in detecting Gram-negative bacteria[21]. Hence, various immune
genes may have been selected in different chicken breeds to protect against gram-negative
bacteria.

Several other genes with non-synonymous changes occur in R1 and R2 genomic regions.
Although we cannot identify the functional consequences of these changes, the strong
signature of selection and KADK-specific protein-coding alterations suggest phenotype-
altering ability. The proximity of R1 and R2 regions to the *Fm* locus indicates that selective
sweep may be due to close physical linkage with the *Fm* locus. Such co-selection of traits due
to linkage and pleiotropy has occurred during animal domestication, including in
chicken[6,86,87]. Hitchhiking of alleles can increase the frequency of both beneficial and
mildly deleterious alleles [88–95]. Whether the changes in KADK are beneficial or
deleterious will need to be investigated in future studies.

The genomic co-occurrence of agronomically important traits in domesticated plants and
animals is also known[96–105]. The alleles causing phenotypic changes can occur in closely
linked genes and will undergo selection for any of the phenotypes being favored by the
breeding process. Notably, the R1 and R2 selective sweeps are found only in KADK, R3 in
YOSK, and none in Chinese BBC breeds. Hence, the genetic variants and the selective sweep
may represent recent events after the BBC breeds separated from each other. Comparing
different BBC breeds provides snapshots of the selection process and the build-up of
genetically linked co-selected allelic changes and may help understand the hitchhiking
process during domestication.

Genome-wide data generated as part of this study and comparative analysis with other BBC
breeds establishes the genetic uniqueness of Kadaknath that extends beyond the *Fm* locus.
We also identify specific genes with signatures of selection that are likely to be responsible
for the Kadaknath-specific phenotypes. Co-selection of genes that are in linkage, as shown in
Kadaknath, is widespread in domestic species[6,86,96–99]. Such clustering of linked alleles
under selection is favored in low recombination regions and near chromosomal

rearrangements. Our work exemplifies the interaction of artificial selection and chromosomal
rearrangement linked traits in domesticated species.

References

1.   Wang MS, Thakur M, Peng MS, Jiang Y, Frantz LAF, Li M, et al. 863 genomes reveal
the origin and domestication of chicken. Cell Res. 2020;30: 693. doi:10.1038/S41422-
020-0349-Y

2.   Tixier-Boichard M, Bed'Hom B, Rognon X. Chicken domestication: from archeology
to genomics. C R Biol. 2011;334: 197–204. doi:10.1016/J.CRVI.2010.12.012

3.   Lawal RA, Hanotte O. Domestic chicken diversity: Origin, distribution, and
adaptation. Anim Genet. 2021;52: 385–394. doi:10.1111/AGE.13091

4.   Eriksson J, Larson G, Gunnarsson U, Bed'hom B, Tixier-Boichard M, Strömstedt L, et
al. Identification of the yellow skin gene reveals a hybrid origin of the domestic
chicken. PLoS Genet. 2008;4. doi:10.1371/JOURNAL.PGEN.1000010

5.   Cogburn LA, Porter TE, Duclos MJ, Simon J, Burgess SC, Zhu JJ, et al. Functional
genomics of the chicken--a model organism. Poult Sci. 2007;86: 2059–2094.
doi:10.1093/PS/86.10.2059

6.   Rubin CJ, Zody MC, Eriksson J, Meadows JRS, Sherwood E, Webster MT, et al.
Whole-genome resequencing reveals loci under selection during chicken
domestication. Nature. 2010;464: 587–591. doi:10.1038/nature08832

7.   Kralik G, Kralik Z, Grčević M, Hanžek D. Quality of Chicken Meat. Anim Husb Nutr.
2016; 63–94. doi:10.5772/intechopen.72865

8.   Mir NA, Rafiq A, Kumar F, Singh V, Shukla V. Determinants of broiler chicken meat
quality and factors affecting them: a review. J Food Sci Technol. 2017;54: 2997–3009.
doi:10.1007/S13197-017-2789-Z

9.   Dou T, Yan S, Liu L, Wang K, Jian Z, Xu Z, et al. Integrative analysis of
transcriptomics and metabolomics to reveal the melanogenesis pathway of muscle and
related meat characters in Wuliangshan black-boned chickens. BMC Genomics.
2022;23: 1–15. doi:10.1186/s12864-022-08388-w

10.   Jaturasitha S, Srikanchai T, Kreuzer M, Wicke M. Differences in Carcass and Meat
Characteristics Between Chicken Indigenous to Northern Thailand (Black-Boned and
Thai Native) and Imported Extensive Breeds (Bresse and Rhode Island Red). Poult
Sci. 2008;87: 160–169. doi:10.3382/PS.2006-00398

11.   Muroya S, Tanabe RI, Nakajima I, Chikuni K. Molecular characteristics and site
specific distribution of the pigment of the silky fowl. J Vet Med Sci. 2000;62: 391–
395. doi:10.1292/JVMS.62.391

12.   Bateson BW, -a F, Punnett RC, Contents M. The inheritance of the peculiar
pigmentation of the silky fowl. J Genet 1911 13. 1911;1: 185–203.
doi:10.1007/BF02981551

13.   Dorshorst B, Molin AM, Rubin CJ, Johansson AM, Strömstedt L, Pham MH, et al. A
complex genomic rearrangement involving the Endothelin 3 locus causes dermal
hyperpigmentation in the chicken. PLoS Genet. 2011;7.

14.   Dharmayanthi AB, Terai Y, Sulandari S, Zein MSA, Akiyama T, Satta Y. The origin
and evolution of fibromelanosis in domesticated chickens: Genomic comparison of
Indonesian Cemani and Chinese Silkie breeds. PLoS One. 2017;12.
doi:10.1371/JOURNAL.PONE.0173147

15.   Shinomiya A, Kayashima Y, Kinoshita K, Mizutani M, Namikawa T, Matsuda Y, et al.
Gene duplication of endothelin 3 is closely correlated with the hyperpigmentation of
the internal organs (fibromelanosis) in silky chickens. Genetics. 2012.

16.   Zhang P, Cao Y, Fu Y, Zhu H, Xu S, Zhang Y, et al. Revealing the Regulatory
Mechanism of lncRNA-LMEP on Melanin Deposition Based on High-Throughput
Sequencing in Xichuan Chicken Skin. Genes (Basel). 2022;13: 2143.
doi:10.3390/GENES13112143/S1

17.   Tu Y gang, Sun Y zhen, Tian Y gang, Xie M yong, Chen J. Physicochemical
characterisation and antioxidant activity of melanin from the muscles of Taihe Black-
bone silky fowl (Gallus gallus domesticus Brisson). Food Chem. 2009;114: 1345–
1350. doi:10.1016/J.FOODCHEM.2008.11.015

18.   Tian Y, Xie M, Wang W, Wu H, Fu Z, Lin L. Determination of carnosine in Black-
Bone Silky Fowl (Gallus gallus domesticus Brisson) and common chicken by HPLC.
Eur Food Res Technol. 2007;226: 311–314. doi:10.1007/S00217-006-0528-
1/FIGURES/1

19.   Sharma R, Sehrawat R, Ahlawat S, Sharma V, Parmar A, Thakur MS, et al. An attempt
to valorize the only black meat chicken breed of India by delineating superior
functional attributes of its meat. Sci Reports 2022 121. 2022;12: 1–12.
doi:10.1038/s41598-022-07575-9

20.   Tian Y, Zhu S, Xie M, Wang W, Wu H, Gong D. Composition of fatty acids in the
muscle of black-bone silky chicken (Gallus gellus demesticus brissen) and its
bioactivity in mice. Food Chem. 2011;126: 479–483.
doi:10.1016/J.FOODCHEM.2010.11.024

21.   Cho Y, Kim JY, Kim N. Comparative genomics and selection analysis of Yeonsan
Ogye black chicken with whole-genome sequencing. Genomics. 2022;114.
doi:10.1016/J.YGENO.2022.110298

22.   Johansson AM, Nelson RM. Characterization of genetic diversity and gene mapping in
two Swedish local chicken breeds. Front Genet. 2015;6: 44.
doi:10.3389/fgene.2015.00044

23.   Sohn J Il, Nam K, Hong H, Kim JM, Lim D, Lee KT, et al. Whole genome and
transcriptome maps of the entirely black native Korean chicken breed Yeonsan Ogye.
Gigascience. 2018;7: 1–14. doi:10.1093/GIGASCIENCE/GIY086

24.   Buranawit K, Chailungka C, Wongsunsri C, Laenoi W. Phenotypic characterization of
Thai native black-bone chickens indigenous to northern Thailand. Thai J Vet Med.
2016;46: 547–554.

25.   Zhu WQ, Li HF, Wang JY, Shu JT, Zhu CH, Song WT, et al. Molecular genetic
diversity and maternal origin of Chinese black-bone chicken breeds. Genet Mol Res.
2014;13: 3275–3282. doi:10.4238/2014.APRIL.29.5

26.   Kriangwanich W, Piboon P, Sakorn W, Buddhachat K, Kochagul V, Pringproa K, et

al. Consistency of dark skeletal muscles in Thai native black-bone chickens (Gallus
gallus domesticus). PeerJ. 2021;9. doi:10.7717/PEERJ.10728

27. Li D, Sun G, Zhang M, Cao Y, Zhang C, Fu Y, et al. Breeding history and candidate
genes responsible for black skin of Xichuan black-bone chicken. BMC Genomics.
2020;21. doi:10.1186/S12864-020-06900-8

28. Huang X, Otecko NO, Peng M, Weng Z, Li W, Chen J, et al. Genome-wide genetic
structure and selection signatures for color in 10 traditional Chinese yellow-feathered
chicken breeds. BMC Genomics. 2020;21. doi:10.1186/S12864-020-6736-4

29. Wang Z, Qu L, Yao J, Yang X, Li G, Zhang Y, et al. An EAV-HP Insertion in 5′
Flanking Region of SLCO1B3 Causes Blue Eggshell in the Chicken. PLOS Genet.
2013;9: e1003183. doi:10.1371/JOURNAL.PGEN.1003183

30. Jadhao RG, Kumbhar DD, Wakode S. Understanding the histology and
hematobiochemical aspects of Kadaknath an Indian black fowl with an appealing taste
native to Dhar district Madhya Pradesh. Br Poult Sci. 2022 [cited 22 Dec 2022].
doi:10.1080/00071668.2022.2154637

31. Tripathi SB, Jatav G, Malik AA, Joshi S, Singh VK, Negi MS, et al. AFLP markers
based genetic diversity and population structure analysis of Kadaknath: an indigenous
black meat poultry breed of India. Anim Biotechnol. 2022;33: 1045–1055.
doi:10.1080/10495398.2020.1865390

32. Slater AE. The Poultry Industry in India. Worlds Poult Sci J. 1945;1: 46–51.
doi:10.1079/WPS19450020

33. Haunshi S, Prince LLL. Kadaknath: a popular native chicken breed of India with
unique black colour characteristics. https://doi.org/101080/0043933920211897918.
2021;77: 427–440. doi:10.1080/00439339.2021.1897918

34. Sehrawat R, Sharma R, Ahlawat S, Sharma V, Thakur MS, Kaur M, et al. First report
on better functional property of black chicken meat from India. Indian J Anim Res.
2021;55: 727–733. doi:10.18805/IJAR.B-4014

35. Haunshi S, Niranjan M, Shanmugam M, Padhi MK, Reddy MR, Sunitha R, et al.
Characterization of two Indian native chicken breeds for production, egg and semen
quality, and welfare traits. Poult Sci. 2011;90: 314–320. doi:10.3382/PS.2010-01013

36. Sahu Senior Scientist N, Vigyan Kendra K, Virendra Kumar Junior I, Sahu N, Kumar
V, Dhruw S, et al. Woman empowerment and uplifting of tribal farmers through
Kadaknath poultry farming in Dantewada district of Chhattisgarh. ~ 315 ~ J Entomol
Zool Stud. 2019;7: 315–319.

37. Thakur MS, Parmar SS, Kumar A, Tomar SS, Sarkhel BC, Das G, et al. Evaluation of
Kadaknath Chicken for Coccidial Resistance by Oocyst Count, Lesion Scoring and
Oocyst Index in Eimeria Tenella Infection . J Anim Res. 2015;5: 579.
doi:10.5958/2277-940X.2015.00097.2

38. Jena P, Panigrahi B, Panda N, Mohapatra L, Mallik B, Bagh J. Reproductive
Performance and Egg Quality Traits of Kadaknath in Intensive Managemental
Condition under Hot and Humid Climate. Int J Livest Res. 2018; 1.
doi:10.5455/IJLR.20180131054034

39. Shanmathy M, Tyagi JS, Gopi M, Mohan J, Beulah P, Ravi Kumar D. Comparative

Assessment on Performance of Aseel and Kadaknath in Hot and Humid Conditions in Tropics. Int J Curr Microbiol Appl Sci. 2018;7: 2156–2165. doi:10.20546/ijcmas.2018.705.251

40. Rout PK, Pani PK, Naithani S. Genetic susceptibility of indigenous chicks to subgroup A Rous sarcoma virus inoculated via the chorioallantoic membrane. Vet Immunol Immunopathol. 1992;33: 89–102. doi:10.1016/0165-2427(92)90037-Q

41. Haunshi S, Devatkal S, Prince LLL, Ullengala R, Ramasamy K, Chatterjee R. Carcass Characteristics, Meat Quality and Nutritional Composition of Kadaknath, a Native Chicken Breed of India. Foods (Basel, Switzerland). 2022;11: 3603. doi:10.3390/FOODS11223603

42. Wang MS, Li Y, Peng MS, Zhong L, Wang ZJ, Li QY, et al. Genomic Analyses Reveal Potential Independent Adaptation to High Altitude in Tibetan Chickens. Mol Biol Evol. 2015;32: 1880–1889. doi:10.1093/MOLBEV/MSV071

43. Ulfah M, Kawahara-Miki R, Farajalllah A, Muladno M, Dorshorst B, Martin A, et al. Genetic features of red and green junglefowls and relationship with Indonesian native chickens Sumatera and Kedu Hitam. BMC Genomics. 2016;17. doi:10.1186/S12864-016-2652-Z

44. Roux PF, Boutin M, Désert C, Djari A, Esquerré D, Klopp C, et al. Re-sequencing data for refining candidate genes and polymorphisms in QTL regions affecting adiposity in chicken. PLoS One. 2014;9. doi:10.1371/JOURNAL.PONE.0111299

45. Qanbari S, Rubin CJ, Maqbool K, Weigend S, Weigend A, Geibel J, et al. Genetics of adaptation in modern chicken. PLoS Genet. 2019;15. doi:10.1371/JOURNAL.PGEN.1007989

46. Fan WL, Ng CS, Chen CF, Lu MYJ, Chen YH, Liu CJ, et al. Genome-wide patterns of genetic variation in two domestic chickens. Genome Biol Evol. 2013;5: 1376–1392. doi:10.1093/GBE/EVT097

47. Mariadassou M, Suez M, Sathyakumar S, Vignal A, Arca M, Nicolas P, et al. Unraveling the history of the genus Gallus through whole genome sequencing. Mol Phylogenet Evol. 2021;158. doi:10.1016/J.YMPEV.2020.107044

48. Li H, Durbin R. Fast and accurate short read alignment with Burrows-Wheeler transform. Bioinformatics. 2009;25: 1754–1760. doi:10.1093/bioinformatics/btp324

49. Garrison E, Marth G. Haplotype-based variant detection from short-read sequencing. 2012.

50. Li H, Barrett J. A statistical framework for SNP calling, mutation discovery, association mapping and population genetical parameter estimation from sequencing data. Bioinformatics. 2011;27: 2987–2993. doi:10.1093/BIOINFORMATICS/BTR509

51. Danecek P, Bonfield JK, Liddle J, Marshall J, Ohan V, Pollard MO, et al. Twelve years of SAMtools and BCFtools. Gigascience. 2021;10: 1–4. doi:10.1093/GIGASCIENCE/GIAB008

52. Danecek P, Auton A, Abecasis G, Albers CA, Banks E, DePristo MA, et al. The variant call format and VCFtools. Bioinformatics. 2011;27: 2156–2158. doi:10.1093/BIOINFORMATICS/BTR330

53. Cingolani P, Platts A, Wang LL, Coon M, Nguyen T, Wang L, et al. A program for
annotating and predicting the effects of single nucleotide polymorphisms, SnpEff:
SNPs in the genome of Drosophila melanogaster strain w1118; iso-2; iso-3. Fly
(Austin). 2012;6: 80–92.
doi:10.4161/FLY.19695/SUPPL_FILE/KFLY_A_10919695_SM0001.ZIP

54. Lee TH, Guo H, Wang X, Kim C, Paterson AH. SNPhylo: a pipeline to construct a
phylogenetic tree from huge SNP data. BMC Genomics. 2014;15: 162.
doi:10.1186/1471-2164-15-162

55. Cook DE, Andersen EC. VCF-kit: assorted utilities for the variant call format.
Bioinformatics. 2017;33: 1581–1582. doi:10.1093/BIOINFORMATICS/BTX011

56. Leigh JW, Bryant D. popart: full-feature software for haplotype network construction.
Methods Ecol Evol. 2015;6: 1110–1116. doi:10.1111/2041-210X.12410

57. Huson DH. SplitsTree: analyzing and visualizing evolutionary data. Bioinformatics.
1998;14: 68–73. doi:10.1093/BIOINFORMATICS/14.1.68

58. Quinlan AR, Hall IM. BEDTools: a flexible suite of utilities for comparing genomic
features. Bioinformatics. 2010;26: 841–2. Available:
http://bioinformatics.oxfordjournals.org/content/26/6/841.short

59. Meisner J, Albrechtsen A. Inferring Population Structure and Admixture Proportions
in Low-Depth NGS Data. Genetics. 2018;210: 719–731.
doi:10.1534/GENETICS.118.301336

60. Korneliussen TS, Albrechtsen A, Nielsen R. ANGSD: Analysis of Next Generation
Sequencing Data. BMC Bioinformatics. 2014;15: 356.

61. Skotte L, Korneliussen TS, Albrechtsen A. Estimating individual admixture
proportions from next generation sequencing data. Genetics. 2013;195: 693–702.
doi:10.1534/GENETICS.113.154138

62. Kopelman NM, Mayzel J, Jakobsson M, Rosenberg NA, Mayrose I. Clumpak: a
program for identifying clustering modes and packaging population structure
inferences across K. Mol Ecol Resour. 2015;15: 1179–1191. doi:10.1111/1755-
0998.12387

63. McKenna A, Hanna M, Banks E, Sivachenko A, Cibulskis K, Kernytsky A, et al. The
genome analysis toolkit: A MapReduce framework for analyzing next-generation
DNA sequencing data. Genome Res. 2010;20: 1297–1303. doi:10.1101/gr.107524.110

64. Gautier M, Vitalis R. rehh: an R package to detect footprints of selection in genome-
wide SNP data from haplotype structure. Bioinformatics. 2012;28: 1176–1177.
doi:10.1093/BIOINFORMATICS/BTS115

65. Delaneau O, Marchini J, Zagury JF. A linear complexity phasing method for thousands
of genomes. Nat Methods 2011 92. 2011;9: 179–181. doi:10.1038/nmeth.1785

66. Kronenberg ZN, Rhie A, Koren S, Concepcion GT, Peluso P, Munson KM, et al.
Extended haplotype-phasing of long-read de novo genome assemblies using Hi-C. Nat
Commun. 2021;12. doi:10.1038/S41467-020-20536-Y

67. Wang X, Luan Y, Yue F. EagleC: A deep-learning framework for detecting a full
range of structural variations from bulk and single-cell contact maps. Sci Adv. 2022;8.

815 doi:10.1126/SCIADV.ABN9215

816 68. Weissensteiner MH, Bunikis I, Catalán A, Francoijs KJ, Knief U, Heim W, et al.
817   Discovery and population genomics of structural variation in a songbird genus. Nat
818   Commun 2020 111. 2020;11: 1–11. doi:10.1038/s41467-020-17195-4

819 69. Amarasinghe SL, Su S, Dong X, Zappia L, Ritchie ME, Gouil Q. Opportunities and
820   challenges in long-read sequencing data analysis. Genome Biol 2020 211. 2020;21: 1–
821   16. doi:10.1186/S13059-020-1935-5

822 70. Hon T, Mars K, Young G, Tsai YC, Karalius JW, Landolin JM, et al. Highly accurate
823   long-read HiFi sequencing data for five complex genomes. Sci Data 2020 71. 2020;7:
824   1–11. doi:10.1038/s41597-020-00743-4

825 71. Patterson MD, Marschall T, Pisanti N, Van Iersel L, Stougie L, Klau GW, et al.
826   WhatsHap: Weighted Haplotype Assembly for Future-Generation Sequencing Reads. J
827   Comput Biol. 2015;22: 498–509. doi:10.1089/CMB.2014.0157

828 72. Sherry ST, Ward MH, Kholodov M, Baker J, Phan L, Smigielski EM, et al. dbSNP:
829   the NCBI database of genetic variation. Nucleic Acids Res. 2001;29: 308–311.
830   doi:10.1093/NAR/29.1.308

831 73. Wang H, Cahaner A, Lou L, Zhang L, Ge Y, Li Q, et al. Genetics and breeding of a
832   black-bone and blue eggshell chicken line. 1. Body weight, skin color, and their
833   combined selection. Poult Sci. 2021;100. doi:10.1016/J.PSJ.2021.101035

834 74. Acri A. Maritime Buddhism. Oxford Res Encycl Relig. 2018 [cited 22 Dec 2022].
835   doi:10.1093/ACREFORE/9780199340378.013.638

836 75. Sudrajad P, Subiharta S, Adinata Y, Lathifah A, Lee JH, Lenstra JA, et al. An insight
837   into the evolutionary history of Indonesian cattle assessed by whole genome data
838   analysis. PLoS One. 2020;15: e0241038. doi:10.1371/JOURNAL.PONE.0241038

839 76. Nie C, Almeida P, Jia Y, Bao H, Ning Z, Qu L, et al. Genome-Wide Single-Nucleotide
840   Polymorphism Data Unveil Admixture of Chinese Indigenous Chicken Breeds with
841   Commercial Breeds. Genome Biol Evol. 2019;11: 1847–1856.
842   doi:10.1093/GBE/EVZ128

843 77. Zhang T, Du W, Lu H, Wang L. Genetic Diversity of Mitochondrial DNA of Chinese
844   Black-bone Chicken. Brazilian J Poult Sci. 2018;20: 565–572. doi:10.1590/1806-9061-
845   2018-0739

846 78. Dar JA, Subashree K, Raha D, Kumar A, Khare PK, Khan ML. Tree diversity,
847   biomass and carbon storage in sacred groves of Central India. Environ Sci Pollut Res
848   2019 2636. 2019;26: 37212–37227. doi:10.1007/S11356-019-06854-9

849 79. Sun YY, Sun L. A Teleost Bactericidal Permeability-Increasing Protein Kills Gram-
850   Negative Bacteria, Modulates Innate Immune Response, and Enhances Resistance
851   against Bacterial and Viral Infection. PLoS One. 2016;11.
852   doi:10.1371/JOURNAL.PONE.0154045

853 80. Holweg A, Schnare M, Gessner A. The bactericidal/permeability-increasing protein
854   (BPI) in the innate defence of the lower airways. Biochem Soc Trans. 2011;39: 1045–
855   1050. doi:10.1042/BST0391045

856 81. Monson MS, Van Goor AG, Persia ME, Rothschild MF, Schmidt CJ, Lamont SJ.

Genetic lines respond uniquely within the chicken thymic transcriptome to acute heat
stress and low dose lipopolysaccharide. Sci Rep. 2019;9. doi:10.1038/S41598-019-
50051-0

82. Chiang SC, Veldhuizen EJA, Barnes FA, Craven CJ, Haagsman HP, Bingle CD.
Identification and characterisation of the BPI/LBP/PLUNC-like gene repertoire in
chickens reveals the absence of a LBP gene. Dev Comp Immunol. 2011;35: 285–295.
doi:10.1016/J.DCI.2010.09.013

83. Elsbach P, Weiss J. Role of the bactericidal/permeability-increasing protein in host
defence. Curr Opin Immunol. 1998;10: 45–49. doi:10.1016/S0952-7915(98)80030-7

84. Weiss J. Bactericidal/permeability-increasing protein (BPI) and lipopolysaccharide-
binding protein (LBP): structure, function and regulation in host defence against
Gram-negative bacteria. Biochem Soc Trans. 2003;31: 785–790.
doi:10.1042/BST0310785

85. Sharma A, Gupta S, Patil AB, Vijay N. Birth and death in terminal complement
pathway. Mol Immunol. 2022;149: 174–187. doi:10.1016/J.MOLIMM.2022.07.006

86. Webster MT, Kamgari N, Perloski M, Hoeppner MP, Axelsson E, Hedhammar Å, et
al. Linked genetic variants on chromosome 10 control ear morphology and body mass
among dog breeds. BMC Genomics. 2015;16: 1–17. doi:10.1186/S12864-015-1702-
2/FIGURES/4

87. Feder JL, Nosil P, Flaxman SM. Assessing when chromosomal rearrangements affect
the dynamics of speciation: implications from computer simulations. Front Genet.
2014;5: 295. doi:10.3389/fgene.2014.00295

88. Kirkpatrick M, Barton N. Chromosome inversions, local adaptation and speciation.
Genetics. 2006;173: 419–34. doi:10.1534/genetics.105.047985

89. Flaxman SM, Feder JL, Nosil P. Genetic hitchhiking and the dynamic buildup of
genomic divergence during speciation with gene flow. Evolution. 2013;67: 2577–91.
doi:10.1111/evo.12055

90. Martin SH, Singh KS, Gordon IJ, Omufwoko KS, Collins S, Warren IA, et al. Whole-
chromosome hitchhiking driven by a male-killing endosymbiont. PLoS Biol. 2020;18.
doi:10.1371/JOURNAL.PBIO.3000610

91. Huff CD, Witherspoon DJ, Zhang Y, Gatenbee C, Denson LA, Kugathasan S, et al.
Crohn's Disease and Genetic Hitchhiking at IBD5. Mol Biol Evol. 2012;29: 101.
doi:10.1093/MOLBEV/MSR151

92. Weist P, Jentoft S, Tørresen OK, Schade FM, Pampoulie C, Krumme U, et al. The role
of genomic signatures of directional selection and demographic history in the
population structure of a marine teleost with high gene flow. Ecol Evol. 2022;12.
doi:10.1002/ECE3.9602

93. Hale MC, Campbell MA, McKinney GJ. A candidate chromosome inversion in Arctic
charr (Salvelinus alpinus) identified by population genetic analysis techniques. G3
(Bethesda). 2021;11. doi:10.1093/G3JOURNAL/JKAB267

94. Franssen SU, Nolte V, Tobler R, Schlotterer C. Patterns of linkage disequilibrium and
long range hitchhiking in evolving experimental Drosophila melanogaster populations.
Mol Biol Evol. 2015;32: 495–509. doi:10.1093/MOLBEV/MSU320

95.   Bierne N. The distinctive footprints of local hitchhiking in a varied environment and
901   global hitchhiking in a subdivided population. Evolution. 2010;64: 3254–3272.
902   doi:10.1111/J.1558-5646.2010.01050.X

903   96.   Li J, Zhang Y, Ma R, Huang W, Hou J, Fang C, et al. Identification of ST1 reveals a
904   selection involving hitchhiking of seed morphology and oil content during soybean
905   domestication. Plant Biotechnol J. 2022;20: 1110–1121. doi:10.1111/PBI.13791

906   97.   Fang C, Ma Y, Wu S, Liu Z, Wang Z, Yang R, et al. Genome-wide association studies
907   dissect the genetic networks underlying agronomical traits in soybean. Genome Biol.
908   2017;18. doi:10.1186/S13059-017-1289-9

909   98.   Zhang D, Liu C, Cheng H, Kan G, Cui S, Meng Q, et al. Quantitative trait loci
910   associated with soybean tolerance to low phosphorus stress based on flower and pod
911   abscission. Plant Breed. 2010;129: 243–249. doi:10.1111/J.1439-0523.2009.01682.X

912   99.   Kong L, Lu S, Wang Y, Fang C, Wang F, Nan H, et al. Quantitative Trait Locus
913   Mapping of Flowering Time and Maturity in Soybean Using Next-Generation
914   Sequencing-Based Analysis. Front Plant Sci. 2018;9. doi:10.3389/FPLS.2018.00995

915   100.   Guan J, Xu Y, Yu Y, Fu J, Ren F, Guo J, et al. Genome structure variation analyses of
916   peach reveal population dynamics and a 1.67 Mb causal inversion for fruit shape.
917   Genome Biol. 2021;22. doi:10.1186/S13059-020-02239-1

918   101.   Huang K, Andrew RL, Owens GL, Ostevik KL, Rieseberg LH. Multiple chromosomal
919   inversions contribute to adaptive divergence of a dune sunflower ecotype. Mol Ecol.
920   2020;29: 2535–2549. doi:10.1111/MEC.15428

921   102.   Wang X, Gao L, Jiao C, Stravoravdis S, Hosmani PS, Saha S, et al. Genome of
922   Solanum pimpinellifolium provides insights into structural variants during tomato
923   breeding. Nat Commun. 2020;11. doi:10.1038/S41467-020-19682-0

924   103.   Jayakodi M, Padmarasu S, Haberer G, Bonthala VS, Gundlach H, Monat C, et al. The
925   barley pan-genome reveals the hidden legacy of mutation breeding. Nature. 2020;588:
926   284–289. doi:10.1038/S41586-020-2947-8

927   104.   Beissinger TM, Hirsch CN, Vaillancourt B, Deshpande S, Barry K, Robin Buell C, et
928   al. A genome-wide scan for evidence of selection in a maize population under long-
929   term artificial selection for ear number. Genetics. 2014;196: 829–840.
930   doi:10.1534/GENETICS.113.160655

931   105.   Kijas JW, Lenstra JA, Hayes B, Boitard S, Neto LR, Cristobal MS, et al. Genome-
932   wide analysis of the world's sheep breeds reveals high levels of historic mixture and
933   strong recent selection. PLoS Biol. 2012;10. doi:10.1371/JOURNAL.PBIO.1001258

Acknowledgments

Funding:

Council of Scientific and Industrial Research Fellowship (SSS)

University Grants Commission Ph.D. scholarship (AS)

The Department of Biotechnology, Ministry of Science and Technology, India (Grant no.
BT/11/IYBA/2018/03) (NV)

Science and Engineering Research Board (Grant no. ECR/2017/001430) (NV)

Author contributions:

Conceptualization: SSS, AS, NV

Methodology: SSS, AS

Investigation: SSS, AS, NV

Visualization: SSS, AS

Supervision: NV

Writing—original draft: SSS, AS

Writing—review & editing: SSS, AS, NV

Competing interest statement

None to declare

Data and materials availability:

All primary sequencing data are available from the SRA under BioProject# PRJEB51457 and
PRJEB58678. All **Supplementary information**, including other associated data and scripts
used for analysis, are provided in an easy-to-browse format:
https://github.com/ceglabsagarshinde/Kadaknath_Project.

Ethics statement

The study was approved by the Institutional Ethics Committee (IEC) of the Indian Institute of

Science Education and Research, Bhopal, vide reference number

IISERB/IEC/Certificate/2018-11/03 dated 8th June 2018.

Contribution to field

Domestication of chicken occurred in Asia before spreading across the world. Large-scale

population genomics has helped unravel the history of domestication and the genetic

contribution from the wild relatives of chickens. Yet, different chicken breeds' distinctiveness

and adaptation to local environments have received limited attention, especially in India.

Hence, population genomic datasets of indigenous breeds from India will help understand the

spread of domesticated chicken and breed formation. Some indigenous breeds are

experiencing a decline in numbers and need conservation. We focussed on one of the most

distinctive indigenous chicken breeds, the Kadaknath chicken, which is well known for its

black bone and meat. Commonly known as Kali masi or karaknath, the unique taste and use

in traditional medicine have made this breed extremely popular across India.

Black-bone chicken breeds are found in several other countries and have fascinated breeders
and geneticists alike. We compared the Kadaknath genome with other black-bone chicken
breeds and found that all black-bone breeds share the same rearrangement junctions at the *Fm*
locus, responsible for the intense black pigmentation throughout the body. Notably, we use
public long-read high-coverage datasets to conclusively resolve the correct arrangement of
the complex chromosomal rearrangement at the *Fm* locus.

Figures

Figures

Fig. 1: Population structure analysis : **(a)** Geographical locations of different black-bone
chicken breeds used in this study are shown on the map using different shapes. The map was
generated using rworldmap, map, and mapdata R packages. **(b)** Genome-wide principal
component analysis (PCA) reveals the genetic relationship of 34   black-bone chicken
individuals. PC1 and PC2 explained 18.9% and 13% variance, respectively. **(c)** Population
genetic structure and individual ancestry were estimated using NGSadmix for 34 black-bone
chickens from different breeds based on best K=7.

Fig. 2: Arrangement of *Fm* locus region in non-BBC and BBC breeds: Two different
non-paralogous regions on chromosome 20 are referred to as Duplication 1 (Dup1) and
Duplication 2 (Dup2), shown in Gold and light green colors, respectively. The length of the
Dup1 region is ~127 Kb, the Intermediate (Int) is ~412 Kb, and Dup2 is ~170 Kb in size.
(**a**) These regions are neither duplicated nor rearranged in nonblack bone chicken (*N). The
Dup1, Int, and Dup2 regions are collectively referred to as the *Fm* locus. Dup1 contains five
genes, *EDN3*, *ZNF831*, *SLMO2*, *ATP5E*, and *TUBB1*, whereas the Dup2 region does not have
any protein-coding genes. (**b**) Three possible scenarios (*Fm_1*, *Fm_2*, and *Fm_3*)
for *Fm* locus have been proposed in BBC (earlier described in Dorshorst et al. 2011,
Dharmayanthi et al. 2017, and Shon et al. 2018). The dark solid purple line represents
haplotype-1, which spans across Dup1 from Flank1 to inverted Dup2 (i.e., Flank-1 + Dup1 +
(inverted Dup2)). The blue line represents haplotype-2, which spans Dup1 from inverted
Dup2 to Int (i.e., (inverted Dup2) + Dup1 + Int).

Fig. 3: Long-read-based haplotypes resolve the sequence spanning the Dup1 region at
***Fm* locus**: The two haplotypes spanning the Dup1 region are distinguished by distinct alleles
at 24 positions using long sequencing reads. Red dotted vertical boxes highlight the alleles
that differ between haplotype-1 and haplotype-2 at the same positions. The alleles at 24 sites

that separate these two haplotypes at various positions along Dup1 are presented sequentially

between the red dotted boxes. (**a**) Haplotype-1 of Dup1 spans from Flank1 to inverted Dup2

(i.e., Flank-1 + Dup1 + (inverted Dup2)). The purple lines represent overlapping reads of

Nanopore containing Dup1 haplotype-1 alleles. The light blue dotted line represents the span

of the same read from the end of haplotype-1 of Dup1 to the end of haplotype-2 of Dup2. (**b**)

Haplotype-2 spans Dup1 from haplotype-2 of Dup2 to the Int region (i.e., (inverted Dup2) +

Dup1 + Int). The blue lines represent the overlapping reads of Nanopore containing

haplotype-2 alleles. The light blue dotted line represents the span of the same read from the

start of haplotype-2 of Dup1 to the start of haplotype-2 of Dup2. Both haplotypes of Dup1 are

connected to the single haplotype of Dup2 (i.e., haplotype-2 of Dup2) but at different ends.

Fig. 4: Long-read-based haplotypes resolve the sequence spanning the Dup2 region at

Fm **locus:** The two haplotypes spanning the Dup2 region are distinguished by distinct alleles

at 25 positions using long sequencing reads. Red dotted vertical boxes highlight the alleles

that differ between haplotype-1 and haplotype-2 at the same positions. The alleles at 25 sites

that separate these two haplotypes at various positions along Dup2 are presented sequentially

between the red dotted boxes. (**a**) Haplotype-1 of Dup2 spans from Int to Flank2 (i.e., Int +

Dup2 + Flank2). The grey lines represent overlapping reads of Nanopore containing Dup2

haplotype-1 alleles. (**b**) Haplotype-2 of Dup2 spans from the start of haplotype-2 of Dup1 to

the end of haplotype-1 of Dup1 region (i.e., Dup1 +(inverted Dup2)+ Dup1). The red lines

represent the overlapping reads of Nanopore containing Dup2 haplotype-2 alleles. The light

blue dotted line represents the span of the same reads from the start of haplotype-2 of Dup1

to the start of haplotype-2 of Dup2. Similarly, the same reads span the end of haplotype-1 of

Dup1 to the end of haplotype-2 of Dup2.

Fig. 5: Comparison of *Fm* **locus read coverage in black-bone vs. non-black-bone chicken**

breeds: Read coverage along chromosome 20 at the *Fm* locus is shown in 1Kb sliding

windows. Two duplicated genomic loci, Dup1 and Dup2, are denoted by vertical dotted red

lines and have a higher coverage in black-bone chicken breeds: (**a**) Kadaknath, (**b**)

Xichuan, (**c**) Yeonsan   Ogye,   (**d**) Tianfu, (**e**) Ayam   Cemani   than   non-black   bone

chicken: (**f**) Red junglefowl (RJF).

Fig 6: The nucleotide sequence of rearranged junctions at the *Fm* **locus in black-bone**

vs. non-black-bone chicken: (**a**) Dup1 + (inverted Dup2) (**b**) (Inverted Dup1) + Dup2.

Brown arrows represent the direction of rearranged junction sequence, which is present as a

continuous sequence (verified by their presence in a single read) in BBC breeds. In contrast,
non-BBC breeds did not contain these junctions. Sequences highlighted in red and blue
represent the Dup1 and Dup2 sequences, respectively. Sequences in black color represent the
subsequent nucleotides.

Fig. 7: Within and between population comparison of KADK with CHIN: (a) Genome-
wide landscape of pairwise genetic differentiation (F_{ST}) between KADK and CHIN in 50Kb
non-overlapping windows. The dark slate gray and deep sky blue colors represent the
alternative chromosomes. The dotted horizontal black line marks the 99th percentile outlier
of estimated F_{ST} for autosome and the Z chromosome, respectively. **(b)** The highest F_{ST}
region with a prominent pattern occurs on chromosome 20. The panels from top to bottom
show estimates of F_{ST} in 10Kb non-overlapping windows within population pairwise
nucleotide diversity (π) and integrated haplotype score (iHS) with major allele as ancestral
and minor allele as derived. The genes are represented as black boxes with names. The
horizontal red dotted line marks the 99th percentile 10Kb F_{ST} outlier. The transparent grey
color highlights the Dup1 region in all panes in the B panel, while R1 and R2 represent the
selective sweep regions. **(c)** Non-synonymous changes (shown above the exon) within genes
in the regions R1 and R2 are denoted by vertical orange lines. Blue boxes are the exons
connected by black lines representing introns.

1055

(a) Arrangement at *Fm* locus in non-BBC (*N)

EDN3 | ZNF831 | SLMO2 | ATP5E | TUBB1

~127Kb

Duplication1(Dup1)

Intermediate (Int) ~412 Kb

~170Kb

Duplication2 (Dup2)

Fm locus

(b) Rearrangement scenarios at *Fm* locus in BBC

Dup1 Haplotype1
Dup1 Haplotype2
Dup2 Haplotype1
Dup2 Haplotype2

Scenario 1 (**Fm_1*)

Dup1 | Dup2 | ~412 Kb | Dup1 | Dup2

Scenario 2 (**Fm_2*)

Dup1 | Dup2 | Dup1 | ~412 Kb | Dup2

Scenario 3 (**Fm_3*)

Dup1 | ~412 Kb | Dup2 | Dup1 | Dup2

(a)   Haplotype 1

Flank1   Dup1   Dup2

Flank1	Dup1 (~127Kb)	Int (~412 Kb)	Dup2 (~170Kb)	Flank2

T
T
T

T
T
T

TCCTGTCATCGGATAGCGCAGGCT

GACTCCAGATTCGGGTCGCGTAGGG

G
G

(b)   Haplotype 2

Dup2   Dup1   Int

Flank1	Dup1 (~127Kb)	Int (~412 Kb)	Dup2 (~170Kb)	Flank2

A
A
A
A

G
G
G
G

C
C
C
C

AATCTCTGATAAGATATATGAATC

GACTCCAGATTCGGGTCGCGTAGGG