Lipid Desaturation Regulates the Balance between Self-renewal and Differentiation in Mouse Blastocyst-derived Stem Cells

Chanchal Thomas Mannully[1], Reut Bruck-Haimson[1], Anish Zacharia[1], Paul Orih[1], Alaa Shehadeh[1], Daniel Saidemberg[1], Natalya M Kogan[1], Sivan Alfandary[1], Raphael Serruya[1], Arie Dagan[1], Isabelle Petit[2]; Arieh Moussaieff[1]*

1. The Institute for Drug Research, The Hebrew University of Jerusalem, Jerusalem 9112102, Israel.
2. Sorbonne Université, Inserm, Centre de Recherche Saint-Antoine, CRSA, F-75012 Paris, France.

* To whom correspondence should be addressed at: ariehm@ekmd.huji.ac.il.

ABSTRACT

Stem cells are defined by their ability to self-renew and to differentiate, both shown in multiple studies to be regulated by metabolic processes. To decipher metabolic signatures of self-renewal in blastocyst-derived stem cells, we compared early differentiating embryonic stem cells (ESCs) and their extra-embryonic counterparts - trophoblast (T)SCs to their self-renewing counterparts. A metabolomics analysis pointed to the desaturation of fatty acyl chains as a metabolic signature of differentiating blastocyst-derived SCs via the upregulation of delta-6 desaturase (D6D; FADS2) and delta-5 desaturase (D5D; FADS1), key enzymes in the biosynthesis of polyunsaturated fatty acids (PUFAs). The inhibition of D6D or D5D by specific inhibitors or SiRNA retained stemness in ESCs and TSCs, and attenuated endoplasmic reticulum (ER) stress-related apoptosis. D6D inhibition upregulated stearoyl-CoA desaturase-1 (Scd1) in ESCs, essential to maintain ER homeostasis. In TSCs, however, D6D inhibition downregulated Scd1. TSCs show higher *Scd1* mRNA expression and high levels of monounsaturated fatty acyl chain products in comparison to ESCs. Addition of oleic acid – the product of Scd1 (essential for ESCs), to culture medium, was detrimental to TSCs. Interestingly, TSCs express a high molecular mass variant of Scd1 protein, hardly expressed by ESCs. Taken together, our data point to lipid desaturation as a metabolic regulator of the balance between differentiation and self-renewal of ESCs and TSCs. They point to lipid polydesaturation as a driver of differentiation in both cell types. In contrast, mono unsaturated fatty acids (MUFAs), known to be essential for ESCs is detrimental to TSCs.

KEYWORDS

Embryonic stem cells; Pluripotent stem cells; Trophoblast stem cells; Metabolomics; Lipid desaturation; D6D; FADS2; D5D; FADS1; Scd1; Apoptosis; Proliferation; ER stress; Self-renewal; MUFAs; PUFAs

INTRODUCTION

The regulation of cell viability by metabolic perturbations has been well-documented for a century. The set of small molecules ("metabolome") has accordingly emerged as an important means for a detailed examination of cell state. While most studies describing the metabolic regulation of cell state and cell viability, have been carried out on cancer cell lines, recent studies strongly suggest that metabolic perturbations also play major roles in stem cell (SC) biology.

The balance between self-renewal and differentiation of SCs of the blastocyst is critical for their function *in vitro* and *in vivo*, and enables the concerted development of the embryo. The blastocyst comprises an inner cell mass (ICM; consisting of the pluripotent epiblast, which generates the future embryo, and the extra-embryonic primitive endoderm), and the trophectoderm (1). During morula compaction and the formation of the blastocyst, cells undergo the first differentiation into ICM and trophoblast (2). Cells that are isolated from the ICM can be cultivated as embryonic (E)SCs, defined by their ability to differentiate into all adult cell types (3), hence pluripotent (P)SCs. Cells isolated from the trophectoderm give rise to the placenta (4, 5), and can be cultivated as trophoblast (T)SCs. Importantly, the low accessibility of early embryos, the paucity of embryonic material, and the technical difficulties in direct experimental manipulation *in vivo*, underscore the necessity of cell models for the study of early development (6). This is especially true for metabolic analyses, as there is no amplification of signal in such analyses, thereby requiring a relatively large amount of cells per sample.

Research during the last decade by several groups, including ours, has uncovered metabolic signatures of SC potency state (7-12), and roles played by metabolic shifts in the molecular circuitry that maintains stemness (7, 10, 11, 13-34). Metabolic perturbations also regulate self-renewal in pluripotent and adult SCs (13, 31, 35-39). Taken together, these studies suggest that the metabolic status of SCs is critical for the balance between self-renewal and differentiation in SCs.

We previously demonstrated a shift in glycolytic flux in PSCs upon their exit from pluripotency, driving differentiation via modification of histone acetylation (7). While shifts in central metabolism (7, 14, 19, 20, 24) and amino acid metabolism (11, 27) have been suggested to regulate the turnover and differentiation of SCs, information on the roles of

lipid pathways in SC biology is still lacking. Nevertheless, lipids represent a major energy

source during early development (40), and their content was reported to regulate the self-

renewal of human ESCs (41). In particular, phospholipids are involved in PSC

differentiation, as CDP-ethanolamine pathway synthesis of phosphatidylethanolamine is

required at the early stage of reprogramming to PSCs (42).

In a seminal study that provided an unbiased view of ESC metabolism vs

differentiated cells, Yanes and colleagues found highly unsaturated metabolite structures

in undifferentiated ESCs (9). Importantly, however, inhibition of polyunsaturated fatty acid

(PUFA) synthesis promoted pluripotency, suggesting that lipid polydesaturation may have

a unique non-redox role in the differentiation of ESCs. In mammalian cells, the rate limiting

enzyme in the conversion of saturated fatty acids (SFAs) to monounsaturated fatty acids

(MUFAs) is stearoyl-CoA desaturase (SCD). The two human isoforms of SCD are SCD1

and SCD5. SCD1 is the ubiquitous human orthologue of the mouse Scd1, whereas SCD5

is expressed in the brain and pancreas. Further support to the notion that the desaturation

of fatty acyl chains plays critical roles in SC metabolism was given by Ben-David and

colleagues, who demonstrated that SCD-1, also called Δ-9-desaturase, is essential for

ESC survival (37).

Here, we sought to reveal the metabolic events that take place upon early

differentiation of blastocyst-derived SCs, and in particular, the ICM-derived ESCs and

their extra-embryonic equivalents, the trophectoderm-derived TSCs, by comparing the

metabolic signatures of early differentiating SCs to their undifferentiated counterparts. We

found higher abundance of polyunsaturated acyl chains in both cell types upon early

differentiation. The inhibition of Δ6 desaturase (D6D, or FADS2) or Δ5 desaturase (D5D,

or FADS1) attenuated differentiation in both cell types, resulting in increased self-renewal.

Interestingly, Scd1 inhibition showed opposite effects on cell viability in ESCs and TSCs,

revealing distinct metabolic requirements between these two embryonic SC populations.

RESULTS

Our previous work demonstrated a metabolic shift in ESCs upon the exclusion of

growth factors that retain pluripotency, causing early differentiation (7). Following the

same approach, we now sought to unveil common and distinct metabolic regulatory

mechanisms of the balance between differentiation and self-renewal in ESCs and TSCs. For this aim, we allowed ESCs or TSCs to differentiate, and compared their metabolic profiles to these of their self-renewing counterparts.

ESCs were cultivated in feeder free conditions for at least 3 passages. 24 hours after seeding, we initiated spontaneous differentiation, by exclusion of LIF and 2i from feeder free culture medium, and collected cells for a metabolomics analysis 48 hours after medium change. The medium of ESCs of the same experimental lot was replaced with fresh ESC culture medium that contained LIF and 2i, and they were used as undifferentiated control. We then compared the early differentiating ESCs to their undifferentiated counterparts. The colonies of early differentiating ESCs showed a non-condensed morphology with unclear borderlines (**Figure 1A**), suggesting their exit from pluripotency. This observation was corroborated by decreased expression of the pluripotency markers *Nanog* and *Sox2* (**Figure 1B**). No significant change was noted in the expression of *Oct4*, known to be downregulated more slowly upon differentiation (43). Partial least squares regression discriminant analyses (PLS-DAs) of the LC-MS-derived metabolic profiles of the early differentiating vs the undifferentiated ESCs, was performed following quantile normalization, and showed discrete metabolome of the early differentiating ESCs (**Figure 1C**; R^2=0.89, Q^2=0.69 for blastocyst derived ESCs; R^2=0.93, Q^2=0.84 for TNGA *Nanog* ESCs).

We cultivated TSCs in feeder free medium (See Materials and methods) for 3 passages, and then excluded FGF4, TGFβ and heparin from cultivation medium, to initiate spontaneous differentiation. The medium of TSCs of the same experimental lot was replaced with fresh TSC culture medium that contained FGF4, TGFβ and heparin, and they were used as undifferentiated control. Early differentiation was confirmed by morphology (**Figure 1D**) and the expression of TSC marker genes. Specifically, the TSC marker gene *Elf5* was downregulated and the trophoblast lineage marker *Tfap2c* was upregulated (**Figure 1E**) (44). The expression of *Eomes* and *Cdx2*, two more TSC markers, did not significantly change. We compared the metabolome of TSCs after 48 hours of differentiation to that of their undifferentiated counterparts, and found a full separation: *Oct4*-GFP M2rtTA homo BL6 TSCs (blastocyst derived TSC1; "BD TSC 1";

R^2=0.99, Q^2=0.98), and *Oct4*-GFP M2rtTA hetero agouti ("BD TSC 2"; R^2=0.99, Q^2=0.91)

(**Figure 1F**).

After demonstrating the metabolic shift in ESCs and TSCs in standard cultivation

media, we wished to study these shifts in ESCs and TSCs, maintained in similar defined

cultivation medium. As TSCs have strict requirements for medium composition, and

cannot be maintained in conditions suitable for ESCs, we maintained ESCs and TSCs in

Tx basic medium, suitable for TSC cultivation (See Materials and Methods). We

supplemented Tx medium with FGF4, TGFβ and heparin for TSCs, or with LIF and 2i for

cultivation of ESCs. 24 hours following seeding, we allowed SCs to differentiate by

excluding the above factors from cultivation medium for 48 hours. The medium of SCs of

the same experimental lot was replaced with fresh ESC/TSC culture medium as above,

and they were used as undifferentiated control. We then compared the metabolic

composition of early differentiating SCs to that of undifferentiated cells. A PLS-DA

demonstrated a distinct common metabolic shift along the second principal component

(PC2), separating self-renewing SCs from early differentiating SCs (**Figure 2A**). To reveal

the metabolites responsible for the separation, we performed a variable importance in

projection (VIP) analysis. Of the 132 VIP>1 list of metabolic features, we noted a large

number of phospholipids (39). Of these, MS/MS experiments confirmed the putative

identifications of 19 phospholipids with more than two double bonds (polyunsaturated) in

their acyl chains. Notably, phospholipids with polyunsaturated acyl chains showed higher

abundances in early-differentiated TSCs and ESCs (**Figure 2B; Table S1**). The acyl

chain composition of phospholipids is an accurate and sensitive measure of fatty acid

incorporation into the cells (45). Accordingly, we assumed that the total content of

polyunsaturated lipid species in the early differentiating SCs is increased. In agreement

with this premise, we measured higher concentrations of total arachidonic acid in the early

differentiating ESCs (**Figure 2C-D**). We next wished to determine the enzymes

responsible for the shift in lipid desaturation in the early differentiating SCs.

Polydesaturation in animals requires sources for lipids with two double bonds, primarily

linoleic and α-linolenic essential fatty acids. Delta 6 desaturase (D6D; fatty acid

desaturase 2; FADS2) and delta 5 desaturase (D5D; FADS1) catalyze the synthesis of

PUFAs (**Figure 2E**). Aligned with the increased abundance of PUFAs, the expression of

D6D and *D5D* was upregulated upon early differentiation of ESCs (**Figure 2F**) and TSCs

(**Figure 2G**).

To examine the expression of markers of stemness in differentiation-inducing

culture conditions, we used SC-26196, a specific inhibitor of D6D. D6D inhibition retained

the expression of pluripotency marker genes in early differentiating ESCs (**Figure 2H)**,

and TSC marker genes in differentiating TSCs (**Figure 2I**).

Taken together, these results suggest that differentiated ESCs and TSCs share

increased levels of polyunsaturated lipids. This shift can be attributed to D6D and D5D

expression, and is allocated at the exit from a self-renewing state.

We wished to better define the influence of polydesaturation on blastocyst derived

SCs. For this, we inhibited D6D or D5D in ESCs or TSCs of the same (129 / black 6)

background (main Figures), and validated the data from these experiments using

independent ESC and TSC lines (Supplemental Figures).

Microscopic examination of the cells suggested a higher number of colonies and

greater colony size in D6D inhibited ESCs. Indeed, D6D or D5D inhibition caused an

increase in ESC numbers (**Figure 3A-B)**, concomitant to an increase in clonogenicity

(**Figure 3C-D).** D6D inhibition in ESCs in a defined Tx medium also resulted in increased

proliferation of ESCs (**Figure 3E).** The effect of D6D inhibition on self-renewal was dose-

dependent (**Figure 3F)**, and specific, as confirmed by the inhibition of D6D by SiRNA

(**Figure 3G**).

The ability of cells to respond to perturbations in endoplasmic reticulum (ER)

function is critical for cell survival (46). Alterations in ER lipid composition result in ER

stress (47). In particular, ER function is associated with the degree of desaturation of

cellular fatty acyl chains (48, 49). We assumed that polydesaturation may decrease the

viability of ESCs by driving ER stress-related apoptosis. In line with this possibility, we

found reduced expression of the ER stress activators inositol-requiring transmembrane

kinase/endoribonuclease 1α (*IRE1α*) and activating transcription factor 6 (*ATF6*) upon

D6D inhibition (**Figure 3H**). D6D inhibition reduced the percentage of dead (**Figure 3I**)

and increased the percentage of pre-apoptotic and necrotic (**Figure 3J-K**) ESCs,

confirming an anti-apoptotic effect. To further confirm the pro-apoptotic influence of D6D

in ESCs, we tested its effect on the viability of further ESC lines (**Figure S1**). We also

wished to define the effect of D6D inhibition on ESC cell cycle, and found no effect (**Figure**
S1), suggesting that the higher proliferation rate is not due to cell cycle rate, and may be
attributed to less ER stress-related apoptosis, leading to higher viability of ESCs following
downregulation of D6D.

Similarly to ESCs, the inhibition of polydesaturation by D6D or D5D increased TSC
proliferation (**Figure 4A**), and the number of colonies (**Figure 4B**), concomitantly with
decreased expression of IRE1α (**Figure 4C**), and attenuation of apoptosis (**Figure 4D-**
E). Unlike in ESCs, the inhibition of D6D also increased the percentage of TSCs in S
phase (**Figure 4F-G**), suggesting that the increase in proliferation rate in TSCs is also
partially mediated by altered cell cycle kinetics. We validated the increased proliferation
of TSCs using independent TSC lines (**Figure S2**).

The cultivation of ESCs and TSCs is mostly done on feeder cells that release
factors that support SC cultivation. To evaluate the influence of D6D on SC proliferation
in feeder cell-based culture, we carried out FACS analyses of the ratio of SCs / mouse
embryonic fibroblasts (MEFs) in culture. We used *Oct4*-GFP labeled ESCs for the
quantification of ESCs. For the quantification of TSC proliferation in co-culture, we labeled
TSCs using an anti-CD40 (a TSC marker) antibody. ESC relative proliferation in culture
with feeder cells increased following the inhibition of D6D, whereas the number of D6D
inhibited TSCs only slightly and insignificantly increased in culture with the MEF feeder
cells (**Figure S3**).

A balance between mono- and poly-desaturation of lipids is maintained in the liver,
brain, lymphocytes and adipocytes, as PUFAs represses Scd1 activity and mRNA stability
(50-52). Scd1 inhibition activates the IRE1/XBP1s signaling arm of the unfolded protein
response, causing ER stress (53, 54). Interestingly, the stress response following Scd1
inhibition causes massive apoptosis in human ESCs, but not somatic cells; mouse ESCs
showed partial elimination (37). Accordingly, we reasoned that the increase in cell viability
in D6D inhibited ESCs may result from upregulation of Scd1 activity, leading to lower
apoptosis rates. In line with this hypothesis, *Scd1* was upregulated in ESCs following
inhibition of D6D (**Figure 5A**). Surprisingly, in contrast to the inverse regulation of *Scd1*
by D6D in ESCs, *Scd1* was downregulated upon D6D inhibition and downregulation in
TSCs, suggesting a reciprocal regulation of Scd1 in the two cell types by PUFAs (**Figure**

5B). Scd1 mRNA expression was slightly downregulated in early differentiating ESCs,
whereas a non-significant upregulation noted in TSCs (**Figure 5C**). We wished to further
examine possible functional implications of Scd1 expression in blastocyst-derived SCs.
The expression of the pluripotent markers *Oct4* and *Sox2* was downregulated in early
differentiating ESCs (**Figure 5D**), whereas TSC markers were upregulated in early
differentiating TSCs (**Figure 5E**) following inhibition of Scd1. The inhibition of Scd1 by
PluriSIn 1 was reported by Ben-David and colleagues to partially eliminate mouse ESCs
(37). Aligned with the results of this study, we saw partial elimination of mouse ESCs by
PluriSIn 1 or A-939572 - another Scd1 specific inhibitor (**Figure 5F-G**). Remarkably, the
inhibition of Scd1 in TSCs increased their viability (**Figure 5H-I**). The addition of
conjugated oleate to Scd1-inhibited TSCs attenuated their proliferation, reducing it to the
level seen in control cells (**Figure 5J**). As expected from the higher viability of TSCs upon
Scd1 inhibition, they showed decrease in the expression of ER stress genes **(Figure 5K)**,
and reduced apoptosis **(Figure 5L-M)**. We tested the viability of ESCs and SCs under
Scd1 inhabitation and confirmed that the increase in viability of TSCs, whereas decrease
of ESC viability by Scd1 inhibition is not cell line specific (**Figure S4**). Inhibition of Scd1
in feeder cell-based culture resulted in increased proliferation in TSCs, whereas lower
ESC proliferation (**Figure S5**).

We assumed that the detrimental influence of Scd1 on ESCs may reflect a
differential expression of the enzyme and its products. We found approximately one order
of magnitude higher gene expression of *Scd1* in TSCs (**Figure 5N**), and a corresponding
accumulation of MUFAs (**Figure 5O**). The examination of Scd1 protein expression
suggested a surprising alteration in the molecular weight of Scd1 in ESCs (molecular
mass ~35 kDa) and TSCs (molecular mass ~70 kDa) (**Figure 5P; Figure S6**). Upon early
differentiation, Scd1 protein expression is decreased in ESCs, with no change noted in
early differentiating TSCs (**Figure 5P; Figure S6**). MEFs (differentiated embryonic cells),
show no preference to either of the Scd1 variants (**Figure 5P; Figure S6**). Scd1 is
regulated by the availability of fatty acids (50). To better understand the shift in Scd1
variants and address a possible regulation of their expression by the availability of
external fatty acids, we incubated ESCs in 0-10% FBS (the source of external fatty acids).
Although we noted an upregulation of Scd1 with increasing percentage of serum, no

change in the high molecular mass variant of Scd1 was noted (**Figure S6**). Nevertheless,

we then examined the expression of Scd1 in ESCs and TSCs, cultivated in the same

defined (Tx) medium, and saw the same alteration in the Scd1 variant (**Figure 5Q**).

Notably, oleic acid – the product of Scd1, blocked the Scd1-dependent increase in viability

in D6D inhibited TSCs (**Figure 5R**), suggesting that this increase is mediated by the

downregulation of the basal high MUFA levels in TSCs.

DISCUSSION

In the current study, we characterized the metabolic profiles of early differentiating

ESCs and extra-embryonic TSCs vis-à-vis their self-renewing counterparts. Our

metabolomics analyses clearly show a metabolic shift in both blastocyst-derived SCs as

they exit self-renewal. This shift results in higher levels of PUFAs, following increased

expression of D6D and D5D. No change was seen in the levels of linoleic acid. The

inhibition of lipid polydesaturation by D6D or D5D retains self-renewal in ESCs and TSCs,

suggesting that PUFA levels are driving differentiation in both cell types, and that

controlling lipid desaturation is essential for the balance between self-renewal and

differentiation, and for the maintenance of ESCs and TSCs in culture.

Previous work suggests that the global metabolome of ESCs is characterized by

abundant metabolites with highly unsaturated structures, of which the levels decrease

upon differentiation (9), whereas the results of our analyses demonstrate upregulation of

lipid desaturation upon differentiation. This seeming discrepancy may be a result of our

focus on desaturation of lipids. It may also be attributed to variations in the differentiation

stage studied (early differentiated vs fully differentiated neurons or cardiomyocytes).

Despite this seeming discrepancy, our data on the retention of pluripotency by the

inhibition of lipid polydesaturation in ESCs are well aligned with the results of Yanes and

colleagues on ESC differentiation (9). Our results further link the shift in lipid

polydesaturation to lower SC self-renewal by exerting ER stress-related apoptosis.

The regulation of the ER status by lipid desaturation has been reported in multiple

studies in somatic cells. In ESCs, downregulation of Scd1 causes ER stress and

apoptosis, as demonstrated by Ben-David and colleagues (37), and by our results. Given

the regulation of Scd1 by PUFAs in the liver, lymphocyte, brain, and adipocytes cells (50),

we reasoned that lipid polydesaturation inversely regulates Scd1 in ESCs. Our results are

in concurrence with this assumption, suggesting that polydesaturation – a metabolic

signature of stemness, may modulate the ER status via regulation of Scd1 expression.

Surprisingly, we saw a downregulation of Scd1 following D6D inhibition in TSCs.

Despite the reciprocal regulation of Scd1 by D6D in ESCs vs TSCs, inhibition of

polydesaturation decreases ER stress and apoptosis in both cell types. The fact that Scd1

inhibition increases ESC death while increasing the viability of TSCs provides an

explanation for this, and underscores a unique essentiality of MUFAs in ESCs when

compared to their extra-embryonic counterparts.

The finding that TSCs express higher mRNA levels and a different protein variant

of Scd1, provides mechanistic insights into the differences in its essentiality and function.

A high molecular Scd1 variant, such as the one expressed In TSCs, was previously

attributed to the formation of dimers (55, 56). It was suggested to have potentially altered

function or stability (57), but it has not been isolated, and its function has so far not been

studied .

Our results show that while ESCs are addicted to MUFAs, TSCs have high levels

of MUFAs, high mRNA expression of Scd1, and an altered variant of Scd1 protein. The

addition of conjugated oleate to D6D-inhibited TSCs rescues apoptosis, supporting the

notion that increasing MUFA levels may be detrimental for TSCs, and linking the pro-

apoptotic effect of D6D to the regulation of MUFA levels.

Further research is required to establish the function of Scd1 variants and their

regulation by PUFAs in the early embryo. Seeing that the high molecular variant of Scd1

is expressed in the extraembryonic TSCs, it would be interesting to study its expression

and function in the placenta.

The downregulation of the expression of pluripotency markers by Scd1 inhibition

in ESCs may be a nonspecific toxic effect. Still, the upregulation of TSC marker genes by

Scd1 inhibition suggests that the regulation of the blastocyst-derived SC populations by

Scd1 is also mediated by modulation of the potency of the cells to differentiate, and further

underscores its pivotal role in the regulation of the two cell populations and their

progenies.

ESCs or TSCs are mostly cultivated in co-cultures with feeder cells that provide
them with nutrients and paracrine signals required for their cultivation. Our analyses show
that the regulation of ESC viability by lipid desaturases is less pronounced in co-culture
with MEFs than in feeder free culture, but still significant. This is likely due to the
differences in the media lipid composition, conferred by the feeder cells. We saw no effect
of D6D inhibition in TSCs in co-culture, implying that the advantage offered by inhibition
of polydesaturation is limited to their cultivation in feeder free culture.

Taken together, our data suggest that lipid polydesaturation is a common
metabolic signature of differentiating blastocyst-derived SCs, marking a shift from self-
renewal to differentiation. This shift is pro-apoptotic in both ESCs and TSCs via
downregulation of Scd1 in ESCs (dominant variant~35 kDa) whereas upregulation of
Scd1 (showing a ~63-70 kDa variant) in TSCs. Our results expand the concept of
metabolic regulation of stem cells fate decisions to the extra-embryonic stem cells, and
make a clear functional distinction between mono-and poly-desaturation of lipids in these
cells. Our work on blastocyst-derived cells suggests that the availability of MUFAs and
PUFAs in the microenvironment of the blastocyst, and the expression of D6D, D5D and
Scd1 in the blastocyst cell types are critical regulators of the concerted differentiation and
expansion of the blastocyst cell populations. Such premise requires further studies, and
especially in mammalian embryos.

MATERIALS AND METHODS

Cell lines

R1 *Oct4*-GFP ESCs were kindly provided by Prof. Andras Nagy of the Lunenfeld
Research Institute. V6.5 ESCs were kindly provided by Rudolf Jaenisch, MIT. TNGA
Nanog ESCs (with a background of 129 / black 6) and the blastocyst-derived TSC clones:
BD TSC 1 with a background of 129 / black 6, and BD TSC 2 with Homo BL6 background
were kindly provided by Dr. Yosef Buganim, Hebrew University of Jerusalem.

Cell culture

Cells were cultured in standard conditions. In brief: Cells were maintained in a humidified
incubator at 5% CO_2 and 37 °C (Thermo Fisher Scientific, USA).

ESCs in maintenance were cultured as previously described by us (17, 58), on a

confluent adhesive layer of mitomycin-inactivated MEF feeder cells in 2i ESC culture

DMEM medium supplemented with 15% Fetal bovine serum (FBS; Gibco, USA), 1.5%

Pen/Strep (Biological Industries,USA), 1% sodium pyruvate (Biological Industries, USA),

1% L-glutamine (Biological Industries, USA), 1% non-essential amino acids (Gibco, USA),

0.1 mM 2-mercaptoethanol (Gibco, USA), 2.8 µM CHIR99021 (Biogems, Peprotech), 1

µM PD0325901 (Biogems, Peprotech), mouse Leukemia inhibitory factor (LIF; in-house

production)]. Every two days, cells were passaged. Before experiments, ESCs were

cultured in feeder free conditions (in LIF/2i ESC culture medium) on 0.2% gelatin coated

culture plates for 3 passages.

TSCs in maintenance were cultivated as previously described (4), on gelatin

coated mitomycin inactivated MEFs in 30% TSC medium [RPMI 1640 (Gibco, USA)

supplemented with 20% FBS (Gibco, USA), 1% non-essential amino acids (Gibco, USA),

1% L-glutamine (Biological Industries, USA), 1% sodium pyruvate (Biological Industries,

USA), 1% pen/strep (Biological Industries, USA), 50 µM β-mercaptoethanol (Gibco,

USA)] and 70% MEF conditioned medium (medium collected from mitomycin inactivated

MEFs, containing the same supplements) freshly supplemented with 25 ng/mL of human

recombinant FGF4 (prepared in house) and 1 µg/mL heparin (Sigma-Aldrich, USA). TSCs

were passaged every 3-5 days, using trypsin EDTA. For feeder free experiments, TSCs

were cultured in Matrigel ® Matrix (Corning Life Sciences, USA) coated [1:30 dilution in

DMEM/F-12 medium (Gibco, USA)] tissue culture plates in Tx defined medium (59) with

freshly added 25 ng/mL of mouse recombinant FGF4 (In-house production), 1 µg/mL

heparin (Sigma-Aldrich, USA), 1 µg/mL of human TGFβ1 (Biogems, Peprotech).

Fetal bovine serum used throughout the study was from one batch, to avoid

possible batch-dependent changes in the lipid composition of the cultivation medium.

ESC and TSC early differentiation

Cells were cultivated in a feeder free medium: ESCs in 2i culture medium on 0.2% gelatin

coated plates and TSCs on Matrigel coated plates in TX defined medium (59). Cells were

maintained in feeder free conditions for at least 3 passages prior to experiments. For

initiation of differentiation, growth factors and cytokines that maintain cells in pluripotency

(LIF and 2i) or TSC state (heparin, recombinant human TGF-β and FGF-4) were excluded

from medium. For the infliction of differentiation, 24 hours following seeding, we allowed

SCs to differentiate by excluding the above factors from cultivation medium for 48 hours.

The medium of SCs of the same experimental lot was replaced with fresh ESC/TSC

culture medium as above, and they were used as undifferentiated control. We then

compared the metabolic composition of early differentiating SCs to that of undifferentiated

cells.

Metabolomics analyses

Following 0-48 hours of spontaneous differentiation, ESCs or TSCs were harvested and

centrifuged at 200 g, 4 °C, 5 min. Cell pellets were washed twice with Dulbecco's PBS

without Ca and Mg (Biological Industries, USA), snap frozen in liquid nitrogen, and kept

at -80 °C. Cell pellets were extracted in an ice cold solvent containing 86.5% methanol

(J. T. Baker, USA), 12.5% LC-MS grade water (J. T. Baker, USA), 1% formic acid (Tokyo

Chemical Industry Co., Ltd., Japan). Samples were vortexed and extracted by

ultrasonication followed by freeze-thaw cycles. Ultrasonication was carried out using

Bioruptor® Plus sonicator (Diagenode, USA) at 30 second on / off pulsation cycles for 5

min. Following centrifugation, ultra-sonication and freezing cycles were repeated for two

additional cycles and pooled for each sample in a fresh pre-labeled glass tube.

Supernatant was transferred to clean glass tubes, and concentrated in a SC210A speed

vacuum concentrator (SAVANT SPD 121P, Thermo scientific, USA) at 30 °C. The

remaining pellet was extracted in 400 µL acetone, ultra-sonicated for 30 seconds, and

centrifuged for 10 min at 4 °C. Supernatant was transferred to the respective glass tubes

containing the methanol/water concentrated extracts and concentrated to dryness.

Samples were reconstituted in 200 µL of reconstitution solvent: 95% acetonitrile (J. T

Baker, USA; LC-MS grade), 5% water (LC-MS grade; J. T Baker, USA), 0.1% formic acid

(Tokyo Chemical Industry Co., Ltd., Japan), and transferred to 96 well 700 µl sample plate

(Waters®Acquity UPLC plate, USA).

Ten microliters of samples were injected to a Waters Acquity UPLC H-Class apparatus

(Waters, Milford, MA, USA) equipped with UPLC CSH C18 2.1 x100 mm, 1.7 µm column

(Waters, Ireland). The column was equilibrated with 0.4 mL/min flow of 40% of mobile

phase A (ACN: Water 2:98% V/V) and 60% of mobile phase B (ACN). The linear gradient

program was as follows: 60% mobile phase A (0.1% formic acid in water) and 40% mobile

phase B for 1 min. The mobile phase B proportion was increased to 70% (v/v) in 5 min. From 5 to 8 min, mobile phase consisted of 24% A, 40% B, and 36% C [(isopropanol (Chemsolute, T.Geyer)]; from 8 to 9 min, 20% A, 35% B, and 45% C; from 9 to 12 min, 18.4% A, 33% B, and 48.6% C; from 12 to 17 min, 12% A, 25% B, and 63% C; and up to 25 min, 0.4% A, 10.5% B, and 89.1% C. Aquity UPLC system was coupled to a Xevo G2-XS, high resolution and high mass accuracy QTOF equipped with ESI. Negative ion mode was used for downstream analyses. Masses of 30-2000 Dalton underwent MS^E analysis with ramp collision energies 30-60eV. Mass range calibration was performed using Leucine enkephalin (Waters, USA) infused every 0.3 seconds. A quality control (QC) mix prepared from pooled samples was injected every 10 runs. A blank sample was injected every 10 samples. A minimum intensity cutoff mass was set to 100 m/z. Fold change greater than 100 from blank and lowest mean abundance in blank was set as another threshold. Data acquisition and visualization was performed using MassLynx 4.1 (Waters Co., UK). Spectral deconvolution, alignment, and normalization were performed using Progenesis QI (Nonlinear dynamics, Waters Corporation, UK). For metabolite identification, exact mass, isotope patterns and fragmentation patterns were compared to 18 metabolite libraries compatible with Progenesis QI (60). Multivariate statistical analyses, masses were carried out using Progenesis QI and Metaboanalyst 4.0 (Waters Corporation, UK). Feature intensities were quantile normalized.

Evaluation of total arachidonic acid levels using GC-MS. ESCs or TSCs were harvested and washed twice with PBS. Cell pellet (1×10^6 cells) was snap frozen in liquid nitrogen and lyophilized (Labconco Corporation, USA). Lyophilized cell pellets were hydrolyzed with 1mL of 1M KOH (Sigma Aldrich, USA) in 70% Ethanol (JT Baker, USA; HPLC grade) at 90 °C for 1 hour. The reaction mixture was acidified with 0.2 mL of 6M HCl (Sigma Aldrich, USA) and then 1 mL of water was added. The hydrolyzed fatty acids were extracted with 1mL of hexane (GC grade, Sigma Aldrich, USA) and further concentrated using vacuum concentrator (Savant SPD121P speed-vac concentrator, Thermo Scientific, USA). Fatty acids were then methylated with 2 mL of Boron Trifluoride (BF3) in Methanol (14% W/V) (Sigma Aldrich, USA) and incubated at 55 °C for 1.5 hours with intermittent shaking every 20 min. 2 mL of saturated solution of $NaHCO_3$ (Sigma Aldrich, USA) and 3 mL of hexane were then added, and the tubes were vortex-mixed for

a minute. The hexane layer was recovered carefully and concentrated using vacuum
concentrator. Samples were reconstituted in 100 µL hexane (GC grade, Sigma Aldrich,
USA). GC-MS analysis was performed using an Agilent 8860 GC system equipped with
an Agilent 7683 N auto sampler and a Agilent J& W GC columns
(30 m × 0.25 mm × 0.2 µm, Agilent technologies USA). The injector was set at 220 °C and
1 µL injections were made with helium as carrier gas, maintained at flow rate of 1 mL/min.
GC gradient is given as **Table S2**. The transfer line from GC column to MS (Agilent 5977B
GC/MSD) was set to 250 °C, ionization energy of 70 eV. A standard for arachidonic acid
(Sigma Aldrich, USA) was used for validation of identification.

Scd1 or D6D enzyme inhibition assay
Selective inhibitors for Scd1 PluriSln-1 and A-939572, D6D (SC-26196) and D5D
(sesamin) were purchased from Cayman Chemicals, USA. The effect of the
pharmacological inhibitors was evaluated for ESCs and TSCs following 48 hours
incubation. Oleate conjugated to albumin (Sigma Aldrich, USA) was also tested for its
influence on Scd1 or D6D-inhibited TSCs.

SiRNA transfection
ESCs were maintained at a density of 2.5 x 10^4 cells/well in 24 well plate for 24 hours
following passage, in ESC LIF/2i medium. Cells were transfected with MISSION® SiRNA
oligomers or MISSION® siRNA Universal Negative Control #1 (Sigma Aldrich, Merck,
USA) at a final concentration 20 nM, mixed with 1.5 µL X-tremeGENE transfection
reagent (Sigma Aldrich, Merck, USA). The complex was incubated at room temperature
for 30 min in serum free medium and transferred to the cells. Following 6 hours post
transfection, the medium was replaced with ESC LIF/2i medium, and the cells were
harvested after 48 hours.

Cell counts
Cells were seeded at a density of 15,000 (ESCs) or 50,000 (TSCs) cells per well in 24
well plates, in feeder free culture. Cell count was carried out using Cyto Smart cell counter
(Corning Incorporated Life Sciences, USA). Counts were validated by manual counting.

Fluorescent Detection
Fluorescence of *Oct4*-GFP ESCs was further quantified using CYTATION 3 plate reader
(LUMITRON, Israel) at an excitation wavelength of 395 nm and emission peak of 509 nm.

Microscopy

Microscopic imaging was taken at 10 x magnification using OLYMPUS IX73 Inverted Microscope (OLYMPUS Life Science, USA).

Quantitative real-time Polymerase Chain Reaction (RT qPCR)

Total RNA for the experiments was extracted from the cells using GENEzol™ TriRNA Pure Kit + DNAse (Geneaid HyLabs, Taiwan) according to the manufacturer's protocol. Quantification of total RNA concentration was done using NanoDrop ND-1000 spectrophotometer (Version 3.8.1, Thermo Fisher). First strand cDNA synthesis was done from 1000 ng of RNA using qScript cDNA Synthesis kit (Quantabio, USA). RT qPCR was performed using Fast SYBR™ Green Master Mix (Applied Biosystems™, Thermo Fisher) and a CFX Connect™ Real-Time PCR Detection System. Primer sequences (HyLabs and IDT) are listed in **Table S3**. Relative expression of the transcripts was normalized to Hypoxanthine Phosphoribosyltransferase (HPRT)1 or Ubiquitin C (UBC), and the gene expression was quantified based on comparative $2^{-\Delta\Delta CT}$.

Flow cytometry

Following dissociation, cells were centrifuged at 200 g for 5 min, and cell pellets were washed twice in PBS containing 1% FBS, and filtered through a cell strainer to remove aggregates. Cells incubated with secondary and no primary antibody (for stained cells) or unlabeled cells (for GFP-labeled cells) were used as negative controls. Flow cytometry was carried out on CytoFLEX Flow Cytometer (BECKMAN COULTER, Life Sciences) using a 100 µm nozzle. Analysis was carried out using CytExpert 2.4 software (BECKMAN COULTER, Life Sciences). For TSCs in co-culture with feeder cells, cells were incubated with anti-CD40 antibody (Ab22469, 1:300; Abcam), and then with Alexa Flour-conjugated anti Rat 488 secondary antibody (Ab150153, 1:500; Abcam).

Apoptosis assay

Cells were seeded on 0.2% gelatin (ESCs) or matrigel (1:30 dilution; TSCs) coated culture plates. Cell pellets were washed with ice cold PBS and resuspended in 1X Annexin V binding buffer (Cell Signaling Technology, USA). Cells were then incubated with Annexin V-FITC conjugate (Cell Signaling Technology, USA) for 10 min on ice protected from light. Propidium Iodide (20 µg/mL; Fisher Scientific, USA) was added to cell suspension immediately before FACS analysis.

Immunoblotting

Cells were collected by trypsinization, then washed with PBS (Biological Industries, Israel). Proteins were extracted with 100 µL in-house made RIPA (1% NP-40, 0.1%SDS, 1M Tris-HCL pH 7.4, NaCl, Sodium deoxycholate, EDTA, and water) + Protease Inhibitors (Roche, USA) and quantified using Bradford reagents (Sigma Aldrich, USA). Extracted proteins were reconstituted in an in-house Protein Sample Buffer (Glycerol 100%, 1M Tris pH6.8, 10% SDS, β- mercaptoethanol, bromophenol blue) and heated for 10 min at 95 °C. Proteins were separated on 10% SDS-PAGE gels and transferred to 0.22 µm PVDF membrane (FroggaBio, USA) by an electric current at 300 mA for 70 min. The membrane was blocked in 5% BSA/TBST for 2 hours, then incubated with primary antibody overnight. Scd1 antibody was purchased from Cell Signaling Technology USA, (#2794, 1:1,000). Membrane was washed, then incubated with HRP conjugated secondary antibody (Abcam, USA, Ab 6721, 1:10,000). eIF4G2/p97 antibody from Cell Signaling Technology (#2182, 1: 3000), or mouse monoclonal anti α-tubulin antibody (T5168; Sigma Aldrich, USA) were used as a loading control. Chemiluminescence was detected by ECL (Advansta, USA), using Molecular Imager ChemiDoc™ XRS (Bio-Rad, USA).

Cell cycle analysis

Cells were seeded at a density of 30,000 (ESCs) or 100,000 (TSCs) cells per well in a 6 well culture plate and treated with D6D inhibitor or DMSO vehicle for 24 hours. Upon 70% cell confluence, cells were treated with 10 µM of Edu (5-ethynyl-2′-deoxyuridine; Invitrogen™, Thermo Fisher Scientific, USA), and incubated for 30 min. After labeling, cells were harvested and washed twice with PBS. Cell pellets were resuspended in cold PBS and fixed by adding cold absolute ethanol dropwise while mixing. Cells were incubated overnight at 4 °C. Following centrifugation, cell pellets were permeabilized by adding 0.1% Triton® X-100 (Sigma Aldrich, USA) in PBS [1% BSA (Sigma Aldrich, USA)], and stained with Click-iT reaction mix [50 mM Trizma® Hydrochloride Buffer pH 7.5 (Sigma Aldrich, USA), 150 mM NaCL (Sigma Aldrich, USA), 2 mM CuSO4 (Sigma Aldrich), 2.5 µM Alexa Fluor 647 azide triethyl ammonium salt in DMSO (Thermo Fisher Scientific, USA) and 10 mM Sodium ascorbate (Sigma Aldrich, USA)] for 30 min. The cells were again washed with 0.1% Triton® X-100, 1% BSA in PBS, and stained with propidium iodide (Thermo Fisher Scientific, USA), 50 µg/mL of RNase (Thermo Fisher

Scientific, USA) in PBS. EdU-analysis was performed using CytoFLEX Flow Cytometer

and CytExpert 2.4 software (BECKMAN COULTER, Life Sciences).

REFERENCES

.1   Sozen B, Cox AL, De Jonghe J, Bao M, Hollfelder F, Glover DM, et al. Self-Organization

of Mouse Stem Cells into an Extended Potential Blastoid. Dev Cell. 2019;51(6):698-712 e8.

.2   Shyh-Chang N, Daley GQ, Cantley LC. Stem cell metabolism in tissue development and

aging. Development. 2013;140(12):2535-47.

.3   Blair K, Wray J, Smith A. The liberation of embryonic stem cells. PLoS Genet.

2011;7(4):e1002019.

.4   Benchetrit H, Herman S, van Wietmarschen N, Wu T, Makedonski K, Maoz N, et al.

Extensive Nuclear Reprogramming Underlies Lineage Conversion into Functional Trophoblast

Stem-like Cells. Cell Stem Cell. 2015;17(5):543-56.

.5   Shahbazi MN, Zernicka-Goetz M. Deconstructing and reconstructing the mouse and

human early embryo. Nat Cell Biol. 2018;20(8):878-8.7

.6   Rossant J, Tam PPL. Opportunities and challenges with stem cell-based embryo models.

Stem Cell Reports. 2021;16(5):1031-8.

.7   Moussaieff A, Rouleau M, Kitsberg D, Cohen M, Levy G, Barasch D, et al. Glycolysis-

mediated changes in acetyl-CoA and histone acetylation control the early differentiation of

embryonic stem cells. Cell Metab. 2015;21(3):392-402.

.8   Panopoulos AD, Yanes O, Ruiz S, Kida YS, Diep D, Tautenhahn R, et al. The metabolome

of induced pluripotent stem cells reveals metabolic changes occurring in somatic cell

reprogramming. Cell Res. 2012;22(1):168-77.

.9   Yanes O, Clark J, Wong DM, Patti GJ, Sanchez-Ruiz A, Benton HP, et al. Metabolic

oxidation regulates embryonic stem cell differentiation. Nat Chem Biol. 2010;6(6):411-7.

.10   Sperber H ,Mathieu J, Wang Y, Ferreccio A, Hesson J, Xu Z, et al. The metabolome

regulates the epigenetic landscape during naive-to-primed human embryonic stem cell transition.

Nat Cell Biol. 2015;17(12):1523-35.

.11   Shiraki N, Shiraki Y, Tsuyama T, Obata F, Miura M, Nagae G, et al. Methionine

metabolism regulates maintenance and differentiation of human pluripotent stem cells. Cell Metab.

2014;19(5):780-94.

.12   Park H, Haynes CA, Nairn AV, Kulik M, Dalton S, Moremen K, et al. Transcript profiling

and lipidomic analysis of ceramide subspecies in mouse embryonic stem cells and embryoid

bodies. J Lipid Res. 2010;51(3):480-9.

.13   Ito K, Suda T. Metabolic requirements for the maintenance of self-renewing stem cells.

Nat Rev Mol Cell Biol. 2014;15(4):243-56.

.14   Carey BW, Finley LW, Cross JR, Allis CD, Thompson CB. Intracellular alpha-

ketoglutarate maintains the pluripotency of embryonic stem cells. Nature. 2015;518(7539):413-6.

.15   Cezar GG, Quam JA, Smith AM, Rosa GJ, Piekarczyk MS, Brown JF, et al. Identification

of small molecules from human embryonic stem cells using metabolomics. Stem Cells Dev.

2007;16(6):869-82.

.16   Chung S, Dzeja PP, Faustino RS, Perez-Terzic C, Behfar A, Terzic A. Mitochondrial

oxidative metabolism is required for the cardiac differentiation of stem cells. Nat Clin Pract

Cardiovasc Med. 2007;4 Suppl 1:S60-7.

.17   Elena-Herrmann B, Montellier E, Fages A, Bruck-Haimson R, Moussaieff A. Multi-

platform NMR Study of Pluripotent Stem Cells Unveils Complementary Metabolic Signatures

towards Differentiation. Sci Rep. 2020;10(1):1622.

.18   Folmes CD, Dzeja PP, Nelson TJ, Terzic A. Metabolic plasticity in stem cell homeostasis

and differentiation. Cell Stem Cell. 2012;11(5):596-606.

.19   Folmes CD, Nelson TJ, Martinez-Fernandez A, Arrell DK, Lindor JZ, Dzeja PP, et al.

Somatic oxidative bioenergetics transitions into pluripotency-dependent glycolysis to facilitate

nuclear reprogramming. Cell Metab. 2011;14(2):264-71.

.20   Gu W, Gaeta X, Sahakyan A, Chan AB, Hong CS, Kim R, et al. Glycolytic Metabolism

Plays a Functional Role in Regulating Human Pluripotent Stem Cell State. Cell Stem Cell.

2016;19(4):476-90.

.21   Han S, Auger C, Thomas SC, Beites CL, Appanna VD. Mitochondrial biogenesis and

energy production in differentiating murine stem cells: a functional metabolic study. Cell

Reprogram. 2014;16(1):84-90.

.22 Maury JJ, Chan KK, Zheng L, Bardor M, Choo AB. Excess of O-linked N-
acetylglucosamine modifies human pluripotent stem cell differentiation. Stem Cell Res.
2013;11(2):926-37.

.23 Moussaieff A, Kogan NM, Aberdam D. Concise Review: Energy Metabolites: Key
Mediators of the Epigenetic State of Pluripotency. Stem Cells. 2015;33(8):2374-80.

.24 TeSlaa T, Chaikovsky AC, Lipchina I, Escobar SL, Hochedlinger K, Huang J, et al. alpha-
Ketoglutarate Accelerates the Initial Differentiation of Primed Human Pluripotent Stem Cells. Cell
Metab. 2016;24(3):485-93.

.25 Teslaa T, Teitell MA. Pluripotent stem cell energy metabolism: an update. EMBO J.
2015;34(2):138-53.

.26 Prigione A, Fauler B, Lurz R, Lehrach H, Adjaye J. The senescence-related
mitochondrial/oxidative stress pathway is repressed in human induced pluripotent stem cells. Stem
Cells. 2010;28(4):721-33.

.27 Shyh-Chang N, Locasale JW, Lyssiotis CA, Zheng Y, Teo RY, Ratanasirintrawoot S, et
al. Influence of threonine metabolism on S-adenosylmethionine and histone methylation. Science.
2013;339(6116):222-6.

.28 Teslaa T, Teitell MA. Pluripotent stem cell energy metabolism: an update. EMBO J. 2014.

.29 Wang J, Alexander P, Wu L, Hammer R, Cleaver O, McKnight SL .Dependence of mouse
embryonic stem cells on threonine catabolism. Science. 2009;325(5939):435-9.

.30 Lu C, Thompson CB. Metabolic regulation of epigenetics. Cell Metab. 2012;16(1):9-17.

.31 Zhang H, Badur MG, Divakaruni AS, Parker SJ, Jager C, Hiller K ,et al. Distinct Metabolic
States Can Support Self-Renewal and Lipogenesis in Human Pluripotent Stem Cells under
Different Culture Conditions. Cell Rep. 2016;16(6):1536-47.

.32 Jang H, Kim TW, Yoon S, Choi SY, Kang TW, Kim SY, et al. O-GlcNAc regulates
pluripotency and reprogramming by directly acting on core components of the pluripotency
network. Cell Stem Cell. 2012;11(1):62-74.

.33 Kleger A, Busch T, Liebau S, Prelle K, Paschke S, Beil M, et al. The bioactive lipid
sphingosylphosphorylcholine induces differentiation of mouse embryonic stem cells and human
promyelocytic leukaemia cells. Cell Signal. 2007;19(2):367-77.

.34   Prigione A, Adjaye J. Modulation of mitochondrial biogenesis and bioenergetic
metabolism upon in vitro and in vivo differentiation of human ES and iPS cells. Int J Dev Biol.
2010;54(11-12):1729-41.

.35   Lin N, Qin S, Luo S, Cui S, Huang G, Zhang X. Homocysteine induces cytotoxicity and
proliferation inhibition in neural stem cells via DNA methylation in vitro. FEBS J.
2014;281(8):2088.96-

.36   Kondoh H, Lleonart ME, Nakashima Y, Yokode M, Tanaka M, Bernard D, et al. A high
glycolytic flux supports the proliferative potential of murine embryonic stem cells. Antioxid Redox
Signal. 2007;9(3):293-9.

.37   Ben-David U, Gan QF, Golan-Lev T, Arora P, Yanuka O, Oren YS, et al. Selective
elimination of human pluripotent stem cells by an oleate synthesis inhibitor discovered in a high-
throughput screen. Cell Stem Cell. 2013;12(2):167-79.

.38   Zhang J, Nuebel E, Daley GQ, Koehler CM, Teitell MA. Metabolic regulation in
pluripotent stem cells during reprogramming and self-renewal. Cell Stem Cell. 2012;11(5):589-
95.

.39   Zhu S, Li W, Zhou H, Wei W, Ambasudhan R, Lin T, et al. Reprogramming of human
primary somatic cells by OCT4 and chemical compounds. Cell Stem Cell. 2010;7(6):651-5.

.40   Sturmey RG, Reis A, Leese HJ, McEvoy TG. Role of fatty acids in energy provision during
oocyte maturation and early embryo development. Reprod Domest Anim. 2009;44 Suppl 3:50-8.

.41   Garcia-Gonzalo FR, Izpisua Belmonte JC. Albumin-associated lipids regulate human
embryonic stem cell self-renewal. PLoS One. 2008;3(1):e1384.

.42   Wu Y, Chen K, Xing G, Li L, Ma B, Hu Z, et al. Phospholipid remodeling is critical for
stem cell pluripotency by facilitating mesenchymal-to-epithelial transition. Sci Adv.
2019;5(11):eaax7525.

.43   Radzisheuskaya A, Chia Gle B, dos Santos RL, Theunissen TW, Castro LF, Nichols J, et
al. A defined Oct4 level governs cell state transitions of pluripotency entry and differentiation into
all embryonic lineages. Nat Cell Biol. 2013;15(6):579-90.

.44   Latos PA, Sienerth AR, Murray A, Senner CE, Muto M, Ikawa M, et al. Elf5-centered
transcription factor hub controls trophoblast stem cell self-renewal and differentiation through
stoichiometry-sensitive shifts in target gene networks. Genes Dev. 2015;29(23):2435-48.

.45   Stanley EG, Jenkins BJ, Walker CG, Koulman A, Browning L, West AL, et al. Lipidomics
Profiling of Human Adipose Tissue Identifies a Pattern of Lipids Associated with Fish Oil
Supplementation .J Proteome Res. 2017;16(9):3168-79.

.46   Tabas I, Ron D. Integrating the mechanisms of apoptosis induced by endoplasmic
reticulum stress. Nat Cell Biol. 2011;13(3):184-90.

.47   Kim SJ, Zhang Z, Saha A, Sarkar C, Zhao Z, Xu Y, et al. Omega-3 and omega-6 fatty acids
suppress ER- and oxidative stress in cultured neurons and neuronal progenitor cells from mice
lacking PPT1. Neurosci Lett. 2010;479(3):292-6.

.48   Jakobsen CH, Storvold GL, Bremseth H, Follestad T, Sand K, Mack M, et al. DHA induces
ER stress and growth arrest in human colon cancer cells: associations with cholesterol and calcium
homeostasis. J Lipid Res. 2008;49(10):2089-100.

.49   Serini S, Piccioni E, Merendino N, Calviello G. Dietary polyunsaturated fatty acids as
inducers of apoptosis: implications for cancer. Apoptosis. 2009;14(2):135-52.

.50   Ntambi JM. Regulation of stearoyl-CoA desaturase by polyunsaturated fatty acids and
cholesterol. J Lipid Res. 1999;40(9):1549-58.

.51   Sampath H, Ntambi JM. Polyunsaturated fatty acid regulation of gene expression. Nutr
Rev. 2004;62(9):333-9.

.52   Sessler AM, Kaur N, Palta JP, Ntambi JM. Regulation of stearoyl-CoA desaturase 1 mRNA
stability by polyunsaturated fatty acids in 3T3-L1 adipocytes. J Biol Chem. 1996;271(47):29854-
8.

.53   Grandjean JMD, Madhavan A ,Cech L, Seguinot BO, Paxman RJ, Smith E, et al.
Pharmacologic IRE1/XBP1s activation confers targeted ER proteostasis reprogramming. Nat
Chem Biol. 2020;16(10):1052-61.

.54   Volmer R, van der Ploeg K, Ron D. Membrane lipid saturation activates endoplasmic
reticulum unfolded protein response transducers through their transmembrane domains. Proc Natl
Acad Sci U S A. 2013;110(12):4628-33.

.55   Zhang S, Yang Y, Shi Y. Characterization of human SCD2, an oligomeric desaturase with
improved stability and enzyme activity by cross-linking in intact cells. Biochem J. 2005;388(Pt
1):135-42.

.56   Fernandez Gianotti T, Burgueno A, Gonzales Mansilla N, Pirola CJ, Sookoian S. Fatty

liver is associated with transcriptional downregulation of stearoyl-CoA desaturase and impaired

protein dimerization. PLoS One. 2013;8(9):e76912.

.57   Flowers MT, Ntambi JM. Role of stearoyl-coenzyme A desaturase in regulating lipid

metabolism. Curr Opin Lipidol. 2008;19(3):248-56.

.58   Moussaieff A, Rogachev I, Brodsky L, Malitsky S, Toal TW, Belcher H, et al. High-

resolution metabolic mapping of cell types in plant roots. Proc Natl Acad Sci U S A.

2013;110(13):E1232-41.

.59   Kubaczka C, Senner C, Arauzo-Bravo MJ, Sharma N, Kuckenberg P, Becker A, et al.

Derivation and maintenance of murine trophoblast stem cells under defined conditions. Stem Cell

Reports. 2014;2(2):232-42.

.60   Shehadeh A, Bruck-Haimson R, Saidemberg D, Zacharia A, Herzberg S, Ben-Meir A, et

al. A shift in follicular fluid from triacylglycerols to membrane lipids is associated with positive

pregnancy outcome. FASEB J. 2019;33(9):10291-9.

ACKNOWLEDGEMENTS

700 We thank Dr. Yosef (Yossi) Buganim for the kind gift of blastocyst isolated TSCs and

701 ESCs and for providing the knowhow of TSC maintenance, and Abraham Greenberg and

702 Prof. Itamar Simon for their kind help with the cell cycle analysis.

CONFLICT OF INTERESTS

705 The authors declare no competing interests.

AUTHOR CONTRIBUTIONS

A.M. conceived the project, supervised the project, analyzed data, secured funding, and

wrote the manuscript. C.T.M. and R.B.H. maintained cell cultures and performed

experiments, A.S. maintained cell cultures, P.O. performed experiments. A.D. assisted in

preparing samples for analyses. A.Z., D.S., N.M.K., R.S. and S.A. analyzed data, I.P.

validated the effect of D6D inhibition on a second line of ESCs. A.M. and C.T.M. prepared

the manuscript for submission.

ETHICS APPROVAL AND CONSENT TO PARTICIPATE

This study did not require ethical approval.

FUNDING

This study was supported by The Dr. Adolf and Klara Brettler Center for Research in Molecular Pharmacology and Therapeutics.

DATA AVAILABILITY STATEMENT

The datasets used and/or analyzed during the current study are available from the corresponding author on reasonable request.

FIGURE LEGENDS

Figure 1.

The metabolism of blastocyst-derived SCs shifts upon early differentiation. ESCs and TSCs (0.5×10^6 cells per sample) were seeded in the culture media commonly used for their maintenance (Materials and Methods). After 24 hours, medium was changed to fresh medium with ("Undiff") or without ("Early Diff") factors that support self-renewal. ESCs were allowed to differentiate by the exclusion of LIF and 2i from cell medium. The colonies of early differentiating ESCs show a non-condensed, non-organized morphology (A). Early differentiating ESCs of the experimental lot used for the metabolomics analysis were evaluated for the expression of pluripotency markers by RT q-PCR (B). Following quantile normalization of the data, partial least squares discriminant analyses (PLS-DA) of early differentiating vs undifferentiated ESCs was performed. $R^2=0.89$, $Q^2 = 0.69$ for hetero agouti (HA) blastocyst derived ESCs (BD ESCs); $R^2 = 0.93$, $Q^2 = 0.84$ for TNGA *Nanog* ESCs (C). TSCs were allowed to differentiate by the exclusion of FGF4 and heparin from cell medium. The colonies of early differentiating TSCs show a non-condensed non-organized morphology (D). The expression of TSC marker genes upon early differentiation was evaluated by RT q-PCR (E). Following quantile normalization of the data, PLS-DA of early differentiating vs undifferentiated TSCs. $R^2 = 0.99$, $Q^2 = 0.98$ for blastcyst derived (BD TSC 1, $R^2 = 0.99$, $Q^2 = 0.91$ for BD TSC 2 (F). Values are the mean ± SEM. n=6. *, P < 0.05; *, P < 0.01; **.

Figure 2. Polyunsaturated lipids are a metabolic signature of early differentiating ESCs and TSCs. ESCs and TSCs of the same background (129 / black 6) were seeded in Tx defined culture medium. After 24 hours, medium was changed to fresh medium with ("Undiff") or without ("Early Diff") factors that support self-renewal. Their metabolome was examined after 48 hours, using 0.5×10^6 cells per sample. The second principal component (PC2) of a partial least squares discriminant analysis (PLS-DA; A) shows discrete metabolome of the differentiating SCs. $R^2 = 0.93$, $Q^2 = 0.83$ (n = 6). A heat map representation of the abundance of identified differential polyunsaturated phospholipids of the dataset shown in A, following variable importance in projection (VIP; > 1) analysis (B). The abundance of each metabolic feature was normalized to the sum intensity of peaks of each chromatogram. The colors of quadrants in the heat maps represent the relative abundance of the metabolites, after normalization to the mean abundance of each metabolite across all samples (n = 6). Following hydrolysis of lipids in ESC extracts, total abundance of linoleic acid (C) and arachidonic acid (D) was measured by GC-MS and normalized to the sum of peaks area. The biosynthesis of PUFAs starts from essential fatty acids with two double bonds, and is catalyzed by D6D and D5D (E). *D6D* and *D5D* expression is upregulated in early differentiating ESCs (F) and TSCs (G), as measured by RT-PCR (n = 6). ESCs and TSCs (129 / black 6 background) were treated by the D6D inhibitor SC-26196 (0.2 µM) or DMSO control for 48 hours in feeder free culture without 2i/LIF or FGF4/TGFβ/heparin and the expression of stem cell markers was measured by RT-qPCR (H-I; n = 4). Values are the mean ± SEM. *, P < 0.05; **, P < 0.01; ***, P < 0.001.

Figure 3. Polydesaturation of lipids causes ER stress-related apoptosis in ESCs. ESCs (129 / black 6 background in A-I; V6.5 in J-K) were treated with the specific D6D inhibitor SC-26196 (0.2 µM), D5D inhibitor sesamin (10 µM), or DMSO control for 48 hours in feeder free culture. Cell counting was carried out using CytoSmart cell counter and validated by manual counting (A; n = 4). Fluorescence was quantified in *Oct4*-GFP labeled 129 / black 6 ESCs using a plate reader to verify the increase in proliferation (B; n = 4). Colony numbers per field were estimated under the microscope (C; 5 fields taken

for each well, for 4 wells). Cell survival was further evaluated by a clonogenicity assay (D;

(n = 48). Cell count was further carried out in defined Tx medium (E; n = 4). Dose response

evaluation of the influence of D6D inhibition on cell proliferation rate was performed with

0.1-3 μM SC-26196 (F; n = 4). For validation of the specific effect on D6D, siRNA was

applied (G; n = 4). The expression of the ER stress activators *IRE1*, *ATF6*, and *ASP* was

assessed by RT-qPCR upon D6D inhibition for 24 hours (H; n = 4). The percentage of

dead cells in D6D inhibited ESCs was further assessed by propidium iodide (PI) staining

(I; n = 4). To evaluate apoptosis in V6.5 ESCs (as they have no GFP label) following 24

hours of D6D inhibition, we used a kit based on annexin V-FITC staining. A representative

dot plot and the corresponding bar graph of one of four experiments (n = 4 for each) is

presented (J-K). Data are presented as mean ± SEM. *, $P < 0.05$; **, $P < 0.01$; ***, $P <$

0.001.

Figure 4. D6D causes ER stress-related apoptosis and modifies cell cycle in TSCs.

TSCs (129 / black 6) were treated by SC-26196 (0.2 μM) or DMSO control for 48 hours

in feeder free culture. Cell counting was carried out using CytoSmart cell counter and

validated by manual counting (A; n = 4). Colony numbers per field were estimated under

the microscope (B; 5 fields taken for each well, for 4 wells). The expression of *IRE1*, *ATF6*

and *ASP* was examined by RT-qPCR following D6D inhibition for 24 h (C; n = 4).

Apoptosis was assessed in TSCs following D6D inhibition for 24 hours using Annexin

V/PI staining (D-E; n = 4). The numbers in each quadrant of dot plot represent the

percentage of cells. Cell cycle was analyzed following inhibition of D6D by SC-26196 for

24 hours. The numbers in each quadrant of dot plot represent the percentage of cells (F-

G; n = 4). Data are presented as mean ± SEM. *, $P < 0.05$; ***, $P < 0.001$.

Figure 5. D6D regulates Scd1 expression, essential for ESCs, but deleterious for

TSCs. The expression of *Scd1* in feeder free culture following 48 hours of inhibition of

D6D (0.2 μM SC-26196), or in DMSO-treated control cells, was evaluated by RT-qPCR

in undifferentiated ESCs (A) and TSCs (B) (both 129 / black 6 background). *Scd1*

expression was also assessed in early differentiating ESCs and TSCs, compared to their

undifferentiated counterparts (C). The influence of Scd1 on pluripotency (D) or TSC (E)

markers in early differentiating SCs, as well as on cell viability (F-J) was evaluated following treatment with Scd1 inhibitors [PluriSIn 1 (20 µM) or A939571 (300 nM)] or DMSO vehicle for 48 hours. Conjugated oleate (100 µM) was added to PluriSin 1-treated TSC culture (J). The expression of *IRE1α*, *ATF3*, *ATF6*, and *ASP* was assessed by RT-qPCR following inhibition of Scd1 by PluriSIn 1 for 48 hours (K), and concomitant apoptosis was evaluated using an Annexin V-FITC-based kit (L-M). The relative expression of *Scd1* in self-renewing ESCs and TSCs was evaluated by RT-qPCR (N). Monounsaturated fatty acyl chain phospholipids were identified and quantified in self-renewing TSCs and ESCs by LC-MS. The abundance of each metabolic feature was normalized to the sum intensity of peaks of each chromatogram. The colors of quadrants in the heat maps represent the relative abundance of the metabolites, after normalization to the mean abundance of each metabolite across all samples (n = 6; O). Scd1 protein expression was determined by Western blot analysis (10% acrylamide gel) in self-renewing ESCs and TSCs in ESC cultivation medium or Tx medium, respectively (P), or both cell types in the same defined Tx medium (Q). TSC proliferation was evaluated in TSCs, as above, following incubation with PluriSIn 1 or PluriSIn 1+conjugated oleate (100 µM) (n = 4; R). Data are presented as mean ± SEM. n=4. *, $P < 0.05$; **, $P < 0.01$; ***, $P < 0.001$.

A

Undiff   Early Diff

100 µm   100 µm

B

Relative mRNA expression

■ Undiff
■ Early Diff

Oct 4   *Nanog* ***   *Sox2* **

C

BD ESCs

● Undiff
● Early Diff

Component 2 (17.8%)

Component 1 (20.5%)

TNGA Nanog ESCs

Component 2 (38.3%)

Component 1 (35.5%)

D

Undiff   Early Diff

100 µm   100 µm

E

Relative mRNA expression

■ Undiff
■ Early Diff

Eomes   *Cdx2*   *Elf5* **   *Tfap2c* *

F

BD TSCs 1

● Undiff
● Early Diff

Component 2 (33.4%)

Component 1 (51.4%)

BD TSCs 2

Component 2 (11.2%)

Component 1 (58.7%)

A

BD TSC cell
numbers

B

BD TSC colony
count

C

Expression of ER stress markers
in D6D inhibited BD TSCs

Apoptosis analysis

D

E

Cell cycle analysis

F

G