A Tale of Two Trees: Modeling Apical and Basal Tree Contribution to L2/3 V1

Pyramidal Cell Orientation Selectivity

Konstantinos-Evangelos Petousakis[1,2,*], Jiyoung Park[3,*], Athanasia Papoutsi[1], Stelios Smirnakis[3,+],

Panayiota Poirazi[1,+]

[1]IMBB, FORTH, Heraklion, Crete, 70013, Greece

[2]Department of Biology, University of Crete, Heraklion, Crete, 70013, Greece

[3]Department of Neurology, Brigham and Women's Hospital and Jamaica Plain Veterans Administration

Hospital, Harvard Medical School, Boston, MA, USA

+Corresponding authors: poirazi@imbb.forth.gr, msmirnakis@bwh.harvard.edu

[*]These authors contributed equally to this work

Pyramidal neurons, a mainstay of cortical regions, receive a plethora of inputs from various

areas onto their morphologically distinct apical and basal trees. Both trees differentially

contribute to the somatic response, defining distinct anatomical and possibly functional sub-

units. To elucidate the contribution of each tree to the encoding of visual stimuli at the

somatic level, we modeled the response pattern of a L2/3 V1 pyramidal neuron to orientation

tuned synaptic input. Towards this goal, we used a morphologically detailed computational

model of a single cell that replicates electrophysiological and two-photon imaging data. Our

simulations predict a synergistic effect of apical and basal trees on somatic action potential

generation: basal tree activity, in the form of either depolarization or dendritic spiking, is

necessary for producing somatic activity, despite the fact that most somatic spikes are heavily

driven by apical dendritic spikes. This model provides evidence for synergistic computations

taking place in the basal and apical trees of the L2/3 V1 neuron along with mechanistic

explanations for tree-specific contributions and emphasizes the potential role of predictive

and attentional feedback input in these cells.

26
27

28
29
30 ***Abstract Figure. Graphical Representation of AP Generation in a L2/3 V1 Pyramidal Neuron Model****. Action*
31 *potential generation requires the spatiotemporal coincidence of apical sodium spikes with either basal sodium*
32 *spikes or significant basal depolarizations, allowing the neuron to respond to salient stimuli that may or may not*
33 *be affected by attentional and/or predictive signals from higher-order areas.*
34
35
1. Introduction

In the primary visual cortex (V1), layer 2 and 3 (L2/3) pyramidal neurons conform to a

stereotypical morphology characterized by separate apical and basal dendritic arbors. The

apical tree consists of a thick apical trunk that extends into L1 and splits into an apical tuft.

The basal tree consists of numerous dendritic segments, the majority of which sprout directly

from the base of the soma, indicating that they can directly influence somatic output

(Spruston, 2008). This compartmentalization of V1 neurons is also evident in the wiring

diagram describing each tree: layer 4 (L4) and L2/3 pyramidal neurons synapse with basal

dendrites of L2/3 pyramidal neurons, providing them with feedforward input (Coogan &

Burkhalter, 1990; Larkum, 2013). The apical dendrites of the L2/3 pyramidal neurons instead

receive feedback input from higher order areas of the cortex (such as the secondary and

tertiary visual cortices, lateromedial area and prefrontal cortex (Coogan & Burkhalter, 1990;

Larkum, 2013), as well as orientation-tuned thalamocortical input (Chen et al., 2013; Cruz-

Martín et al., 2014; Jia et al., 2010; Roth et al., 2015). These anatomical and connectivity

features of the apical and basal trees, complemented by their distinct biophysical properties

(Cho et al., 2008), shape both the local dendritic processing and the integration of the two

input streams at the soma.

Few studies have undertaken the technically difficult task of measuring single spine tuning

properties in the mouse visual cortex (Chen et al., 2013; Iacaruso et al., 2017; Jia et al., 2010).

These studies have shown that single spines of L2/3 pyramidal neurons exhibit orientation

selectivity, with synaptic orientation preferences varying even on the same branch (Iacaruso

et al., 2017; Jia et al., 2010). These findings raise the question of how dendritic integration of

sparse synaptic inputs along the apical and basal trees shape orientation tuning in L2/3 V1

pyramidal neurons.

The effect of synaptic input on neuronal output depends on the biophysical properties of

dendritic branches. It has been found that individual dendritic segments of L2/3 neurons

generate nonlinear regenerative responses, termed dendritic spikes, independently of the

soma, making their exact contribution to somatic output unclear (Palmer et al., 2014; Smith

et al., 2013). This could indicate that multiple dendritic segments need to be activated in order

to generate a spike at the soma. On the other hand, it was recently proposed that strong,

sparse inputs to the dendrites are sufficient to drive somatic output (Goetz et al., 2021).

Interestingly, although dendritic spikes in the apical tuft of L2/3 V1 pyramidal neurons

influence orientation selectivity (Goetz et al., 2021; Smith et al., 2013), this orientation

selectivity is robust to ablation of the apical tree (Park et al., 2019). Altogether, the above

studies highlight the need to scrutinize the functional effect that synaptic inputs along the

basal and apical dendritic arbors have on the soma towards a realistic theory of visual

processing.

In this work, we used a detailed biophysical model of a L2/3 V1 pyramidal cell (Park et al.,

2019), to tease apart the relative contributions of apical vs. basal dendritic trees in somatic

orientation tuning. We start by examining the threshold and strength of dendritic non-

linearities, which are found to vary along the apical and basal dendrites. By comparing model

output to *in vivo* two-photon calcium fluorescence imaging data, we find that dendritic and

somatic spontaneous activity patterns in the model adequately reproduce experimental data.

We then simulate orientation tuning in our model and evaluate its responses to specific

combinations of dendritic disparity and synaptic distributions. We find that the tuned output

of the model does not always match the linear summation of tuned input, instead deviating

towards orientation preferences more closely matching either the apical or basal mean

orientation preference.

We continue by evaluating the dependence of somatic spikes on dendritic voltage-gated

sodium channel conductances by performing just-in-time interventions that nullify sodium

conductance on either the apical or basal tree prior to a somatic spike. We find that the

majority of stimulus-driven somatic spiking activity is dependent on sodium spikes generated

in the apical tree, suggesting that apical and basal trees contribute via different mechanisms

to tuned somatic output. We investigate this further by examining the role of both specific

ionic (sodium channels) and synaptic (AMPA/NMDA) mechanisms in shaping orientation

tuning. Results indicate that apical sodium channels are critical for proper neuronal tuning,

while basal sodium channels do not play as significant a role. Conversely, AMPA and NMDA

activity of both the apical and basal trees is required for orientation tuning at the level of the

soma. A potential reason for this is that basal dendritic branches feature higher sodium

spiking thresholds for similar length-normalized electrotonic length constant values

compared to apical branches, reducing their propensity to generate sodium spikes compared

to apical dendrites.

Overall, our simulations suggest a synergistic effect between apical and basal trees, as somatic

spikes are only reliably produced when apical dendritic sodium spikes coincide with basal

synaptically-driven depolarizations and/or spikes. In this manner, apical and basal dendrites

contribute to neuronal function using different mechanisms, ionic or synaptic, which together

drive action potential generation in this neuronal type.

2. Summary Methods

2.1. Model Description

The single neuron model is based on the L2/3 V1 pyramidal cell model of (Park et al., 2019)

created in the NEURON simulation environment (Hines & Carnevale, 2001). As the model

makes use of a variety of passive and active mechanisms (*Supplementary Tables 1-4*), its

electrophysiological properties were validated against experimental data (Park et al., 2019).

Neither the morphology nor the biophysical properties of the source model were altered,

barring specific ion channel or synaptic mechanism conductance changes wherever specified

(see Extended Methods for details). Apical stimulus-driven synaptic inputs have had a 10 ms

delay added, to better reflect experimental data on feedback synaptic inputs (Ju et al., 2020).

Simulations with synchronous activation of the apical and basal synaptic inputs resulted in

qualitatively similar results (data not shown). For a more detailed description of the model,

see (Park et al., 2019) and Extended Methods, section 8.1.

2.2. Simulation Information

All simulations were performed on our High-Performance Computing Cluster (Rocks 7.0) with

624 cores and 3.328 TB of shared RAM under CentOS 7.4 OS (Papadopoulos et al., 2003),

through the NEURON simulation environment (Hines & Carnevale, 2001). Model neuron

output measurements were obtained at a sampling rate of 10 KHz (dt = 0.1 ms). Unless

specified otherwise, voltage recordings from any section of the model (soma / dendrites)

were performed at the midpoint of the respective section. Simulation duration was 2500 ms,

with stimulus onset (where present) at t = 500 ms. For information on manipulations of

biophysical properties for specific experiments, see Extended Methods, section 8.2. For

details on specific simulation protocols, see Extended Methods, section 8.3.

2.3 Two-Photon Imaging

All experimental protocols were approved by Brigham and Women's Hospital (BWH)

Institution Animal Care and Use Committee. Male and female wild-type (C57BL/6) mice were

purchased from The Jackson Laboratory and bred for experiments. A total of 2 mice were

used in these experiments, with recordings from 11 neurons used for data analysis. For details

on chronic window implantation, sparse labeling and *in vivo* imaging methodology, see

Extended Methods, section 8.4.

2.4. Data Analysis

Data analysis of both modeling and two-photon imaging experiments was performed through Python 3.8+, using publicly available libraries as well as custom code. Analysis procedures include methods for dendritic spike detection, quantification of dendritic non-linearities, calculation of dendritic electrotonic length constants, analysis of two-photon imaging recordings, processing of calcium fluorescence traces, detection of calcium events, conversion of voltage traces into approximate calcium fluorescence traces, and comparison of *in vivo* fluorescence data with converted model voltage traces. For specifics, see Extended Methods, section 8.5.

3. Results

3.1. The degree of non-linearity varies across basal and apical model dendrites

To ensure the validity of our modeling approach, we first examined dendritic integration by characterizing the sodium spike properties of all dendritic segments (*Figure 1A*), wherein a specific number of synapses is synchronously activated by two pulses at 50 Hz while all other compartments are modeled as passive ('I3P'; see Extended Methods, section 8.3.1), and the response of each dendritic segment is assessed. Results show that the model can produce realistic dendritic spikes (*Supplementary Figures 1C&D*), while the properties of the spiking events vary between dendritic segments (*Figures 1B-D, Supplementary Figures 1A&B*). Both apical and basal dendritic segments exhibit a large degree of variation in their spiking thresholds (*Figures 1E&F;* threshold synapse count mean and standard deviations: basal 12.286±6.649 sodium, 51.429±13.179 NMDA; apical 14.442±13.825 sodium, 40.721±23.675

NMDA). Dendritic segments with high sodium spiking thresholds also tended to exhibit higher

NMDA spiking thresholds. To quantitatively characterize the nonlinear behavior of each

dendritic segment we used the Nonlinearity Relative to Linear Extrapolation (NRLE) metric

(Behabadi et al., 2012), which is an estimate of the degree of non-linear dendritic integration

within a specific branch (see Extended Methods, section 8.5.2). Results indicate that all

dendritic segments of the model neuron exhibit non-linear input-output functions (*Figure 1D,*

NRLE=1 denotes linear, NRLE<1 sublinear, and NRLE>1 supralinear), in line with experimental

observations (Smith et al., 2013; Wilson et al., 2016). In addition, the NRLE values are

relatively widely dispersed (values ranging from >1 to ~3.5), providing a large dynamic range

of possible dendrite-level computations. The above results show that there is diversity in

apical and basal dendritic integration that can be exploited to modulate orientation tuning.

The above differences are, at least in part, due to the morphological and electrophysiological

features of dendritic segments, as we found that the sodium spiking propensity varies as a

function of dendritic segment volume and electrotonic constant (see *Figures 5C&D*).

Figure 1. Dendritic Properties of a L2/3 V1 Neuron Model. *A. Graphical representation of the iterative paired-*

pulse protocol. Green dots represent allocated synapses. Synapses were activated only within the designated

(red) dendritic segment, with all other segments being passive. We record the voltage response of the dendritic

segment with NMDA channels disabled or enabled, acquiring information on dendritic sodium and NMDA spikes,

respectively. B. Example of an "expected vs. actual plot", showing the response produced by a single basal

dendritic segment (solid black line) as a result of voltage-gated sodium channel activity elicited through

simultaneous activation of 1 to 200 AMPA synapses. The dashed black line indicates the expected dendritic

response assuming a linear dendrite. C. Same as B, but for an apical dendritic segment. D. NRLE value histogram

for all apical (red) and basal (black) dendritic segments. E. Bar plot of per-dendrite nonlinearity thresholds for all

basal dendritic segments, sorted by the sodium spike thresholds. Note that NMDA spike thresholds largely follow

the sodium spike threshold increase. F. As E, for all apical dendritic segments.

3.2. Model neuron features robust orientation tuning

Given the diverse integration properties of basal and apical dendrites in our model cell, we

sought to assess their functional cooperation when receiving a more realistic set of

background- and stimulus-driven inputs. Stimulus driven synapses were modeled with

individual orientation preferences (Jia et al., 2010; Park et al., 2019), leading them to exhibit

different firing rates depending on the orientation of the presented stimulus (for details, see

Extended Methods, section 8.1). As dictated by experimental data (Chen et al., 2013),

background-driven synapses comprise the majority of inputs to the model, although they are

activated at lower rates than stimulus-driven synapses (see Extended Methods, section 8.1).

To validate their impact on somatic output, we compared the simulated spontaneous activity

with *in vivo* calcium two-photon fluorescence imaging data (see Extended Methods, section

8.5.7 and *Supplementary Figure 2*). Although this approach has its limits, it allows us to use

two-photon imaging data to increase model reliability. We found that spontaneous activity

patterns in the model are very similar to those seen in two-photon calcium imaging data

(*Supplementary Figures 2E&F*).

Next, we combined stimulus-driven and background-driven activity to derive the orientation

tuning curve of the model neuron (see Extended Methods, section 8.3.3). We focused on the

biologically plausible model (see Extended Methods, section 8.1), whereby the distribution of

tuned synaptic inputs onto the basal and apical trees is in line with experimentally reported

values (60% vs 40%, respectively; DeFelipe & Fariñas, 1992). The resulting orientation tuning

curve (preferred orientation firing rate: 1.55±0.3 Hz, orthogonal orientation firing rate:

0.21±0.12 Hz, OSI 0.77±0.11, tuning width 44±9.17°; *Figure 2A*) indicates that the model

neuron is well-tuned (OSI>0.2, tuning width<80°; see Extended Methods, section 8.3.3), in

agreement with our prior work using the original version of this model (Park et al., 2019).

Having established the orientation tuning of our model cell, we tried to disentangle the effects

of the apical and basal trees by introducing dendritic disparity (*Figure 2B;* see Extended

Methods, section 8.3.4). We used the disparity protocol, in which the apical and basal trees

are tuned to different orientations: the apical tree features synaptic orientation preferences

sampled from a Gaussian distribution with a mean of 0°, whereas the basal tree uses the same

method, but with a different mean (0° - 90°, in steps of 10°). The expected orientation

preference of the model neuron is calculated using Euler's Formula, which assumes that

synapses contribute to overall orientation preference in a linear fashion (for details, see

Extended Methods, section 8.3.4). We found that the model neuron diverges from the

expected orientation preference (linear summation of synaptic inputs) and exhibits a bias

towards the preferred orientation of the basal tree. This is an example of *basal dominance*

(*Figure 2D*) and was previously observed using the original version of the model used here

(Park et al., 2019). Of note, the model neuron did not exhibit adequate tuning (i.e. 20% or

more of neurons featured OSI<0.2 or tuning width >80°; see Extended Methods, section 8.3.3)

beyond a dendritic disparity of 50°.

To assess whether this result depends on the particular synaptic distribution used, we

repeated the experiment using two different synaptic distribution profiles – the "inverse

biologically plausible" (40% on the basal tree, 60% on the apical tree) and "even" (50% on

both trees) models (see Extended Methods, section 8.1). We found that, indeed, the deviation

from expectation displayed by the neuron appears to be decreased in the "even" model, and

is clearly reversed in the "inverse biologically plausible" model. Similar to the "biologically

plausible" model, these two models did not feature adequate tuning past 50° of dendritic

disparity. Overall, the observed deviation from expectation indicates that apical and basal

synaptic inputs are not linearly combined, but rather interact in a non-linear manner to

produce somatic output. This interaction is influenced by the synaptic distribution along the

two dendritic trees, as well as their mean synaptic orientation preferences.

Figure 2. Neuronal Orientation Preference is Robust and Resists Dendritic Tuning Disparity. *A. Orientation tuning curve for the "biologically plausible" model. B: Example from a configuration of a model neuron set to a disparity of 40°. Right-side polar plots display the distribution of synaptic orientation preferences for the indicated dendritic segments. Left-side polar plots display the distribution of synaptic orientation preferences for the apical (top) and basal (bottom) dendritic trees. C: Example voltage traces recorded at the soma of a model neuron from (A). Note the clearly different responses between the preferred orientation (0°) and all others. D: Orientation preference plot for the "biologically plausible" model featuring various degrees of disparity. Grey dashed line denotes the expected orientation preference per degree of disparity. Black and red arrows point in the direction of basal and apical dominance, respectively. E: As D, but for the "even" model. Note that dominance has shifted from the basal domain towards the apical domain. F: As D, but for the "inverted biologically plausible" model.*

This configuration clearly features apical dominance. Error bars: Standard error of the mean, for all panels.

3.3. Apical Sodium Spikes are the Primary Driver of Neuronal Output

The previous results show that orientation tuning emerges from non-linear interactions

between apical and basal trees. To elucidate the contribution of each tree in orientation

tuning, we selectively interrupt specific aspects of their operation and assess the effect of the

intervention on somatic tuning. We start by selectively and precisely modifying the sodium

spiking activity of each dendritic tree, while controlling for the type of input received

(stimulus-driven vs background-driven). Specifically, we use the ionic intervention protocol

(see Extended Methods, section 8.3.6) to remove the sodium-induced depolarization that is

provided by either the apical or the basal tree, just before the generation of a somatic spike.

We start by using a control version of the "biologically plausible" model ("Control

background") and a preferred (0°) stimulus (case #1). In this condition, both stimulus-driven

and background-driven synapses are active, and most synapses (60%; both stimulus-driven

and background-driven) are allocated on the basal tree. For every somatic spike, we perform

the intervention protocol and categorize the spike as either: "apically-driven" (dependent on

the presence of apical sodium conductances), "basally-driven (dependent on the presence of

basal sodium conductances) or "cooperative" (dependent on the presence of both apical and

basal sodium conductances) (*Figure 3A*). We find that the majority of somatic spikes are

dependent on apical sodium conductance (N = 131 spikes; apically-driven 76.51±10.21%,

basally-driven 9.61±5.03%, cooperative 13.89±11.22%; *Figure 3B)*, which is surprising,

considering the proximity of the basal dendrites to the soma, as well as the salient nature of

the feedforward, visually-evoked inputs received by the basal tree. These somatic events are

a mix of stimulus-driven and background-driven input. In order to separate the two, we

changed the orientation of the stimulus being presented to 90° (i.e. orthogonal to the

orientation preference of the neuron), rendering the vast majority of stimulus-driven

synapses inactive, while keeping background-driven synapses active as normal (case #2). We

found that the resulting percentages are similar to the previous condition (N = 25 spikes;

apically-driven 71.07±31.56%, basally-driven 17.5±22.5%, cooperative 1.43±4.29%; *Figure*

3D), indicating that the dependence of somatic spiking activity on dendritic sodium

conductance does not significantly differ between instances of stimulus-driven and

background-driven activity.

However, it is possible that background-driven synapses are actually the ones that determine

somatic spiking. While firing at low rates (0.11 Hz vs 0.3 Hz for stimulus-driven; see Extended

Methods, section 8.1), these synapses are vastly more numerous compared to stimulus-

driven synapses (75% versus 25% of total synapses). To test this possibility, we equalized the

contribution of background-driven synapses to apical and basal tree outputs, by introducing

a new condition ("Evenly distributed background") whereby the model is configured as in case

#1 (i.e. "biologically plausible" model), but the distribution of *background-driven synapses* is

the same on both trees (i.e. excitatory: 50% apical, 50% basal; inhibitory: 46.5% apical, 46.5%

basal, 7% soma). In this way, we ensure that the greater proportion of background-driven

synapses on the basal tree is no longer a confounding factor. Presenting a stimulus of the

preferred orientation (0°) under these conditions (case #3), we find that the resulting

percentages (N = 156 spikes; apically-driven 85.08±7.34%, basally-driven 5.15±6.72%,

cooperative 9.78±8.17%; *Figure 3C)* are similar to those seen in case #1, indicating that the

distribution of background-driven synapses does not significantly affect somatic spike

generation.

By changing the orientation of the presented stimulus to 90° for this altered model

configuration, we can also evaluate whether the same conclusion holds for background-

driven spikes (case #4). Indeed, the results (N = 42 spikes; apically-driven 85.83±29.83%,

basally-driven 2.5±7.5%, cooperative 1.67±5.0%; *Figure 3E)* are not significantly different

from those seen in case #2. These simulations predict that somatic spike generation reliably

depends on apical sodium conductance, a seemingly counterintuitive finding, as feedforward

stimulus-driven input reaches mostly the basal tree and not the apical tree.

To gain more insight into these findings, we looked at the spatiotemporal evolution of voltage

through different apical and basal paths during the occurrence of an apically- or basally-driven

spike (*Supplementary Figure 6*). For heavily branching apical dendrites, input summation

and/or voltage propagation in one segment along the path exceeds the sodium spiking

threshold, leading to a cascade of dendritic sodium spikes both in the direction of the soma

as well as towards any terminal dendrites that did not yet generate sodium spikes of their

own (e.g. *Supplementary Figure 6H;* dendritic sodium spiking does not originate from the

terminal segment, but propagates to it). The lack of extensive branching on the basal

dendrites, on the other hand, renders them less predictable in terms of sodium spiking activity

(e.g. *Supplementary Figure 6C*). Although the intervention experiments are required in order

to decisively determine the origin of each somatic spike, the temporal precedence of dendritic

sodium spiking reliably indicates which tree drives somatic spiking activity (e.g. compare

Supplementary Figure 6B vs 6C and 6E vs 6F).

Figure 3. Neuronal Output Relies Disproportionately on Apical Sodium Conductance. *A: Diagram describing the ionic intervention protocol, with example traces on the right. Shaded areas on the traces denote the pre-spike (grey) and post-spike (red) intervention time windows. B: Somatic spike dependence on dendritic sodium conductance for the "biologically plausible" model with a stimulus of 0° (preferred orientation). C: Same as B, but background synapses are evenly distributed across the apical and basal dendritic trees, while stimulus-driven synapses follow the "biologically plausible" (60% basal, 40% apical) model. D: Somatic spike dependence on dendritic sodium conductance for the "biologically plausible" model with a stimulus of 90° (orthogonal/non-preferred orientation). E: Same as D, but background synapses follow the "even" model (evenly distributed), while*

3.4. Apical and Basal Dendrites Influence Somatic Tuning via Different Mechanisms

The previous results are counter-intuitive: although we would expect both apical and basal

dendritic segments to contribute to somatic spiking, the contribution of the apical tree

overshadows that of the basal tree, at least with respect to the role of sodium conductances.

However, as indicated by our results in *Figure 2D*, the basal tree, being the bearer of the

majority of feedforward inputs, has a dominant role in orientation tuning. Thus, we assume

that the basal tree exerts its influence via a different mechanism. To assess the veracity of

this assumption, we performed a series of experiments aimed at evaluating the impact of

ionic (see Extended Methods, sections 8.2.1 and 8.3.6) and synaptic (see Extended Methods,

sections 8.2.2, 8.2.3 and 8.3.7) mechanisms on neuronal orientation tuning, using the

biologically plausible model (see Extended Methods, section 8.3.2).

First, we assess the tuning characteristics of the neuron when stimulus-driven input is

removed from either the apical or the basal tree (see Extended Methods, section 8.2.3), and

compare it with the control case. We find that the neuron remains tuned in all cases, albeit

with significantly decreased OSI values (dependent two-sample t-test, control vs basal: $p \approx$

0.000037; control vs apical: $p \approx 0.000007$) (*Figure 4A*). Removal of apical stimulus-driven input

has a much smaller effect on tuning than removal of basal input (dependent two-sample t-

test, apical vs basal: $p \approx 0.0095$), which is in line with our previous findings (Park et al., 2019).

Next, we use the synaptic intervention protocol to selectively reduce synaptic (AMPA and

NMDA) weights by 50% on either the apical or basal tree for a brief time only, and only when

a somatic spike is about to occur (see Extended Methods, section 8.3.7). We find that somatic

tuning is completely abolished when manipulating the basal but not the apical synaptic

weights, although tuning is significantly affected by loss of apical inputs as well (*Figure 4B*)

(dependent two-sample t-test, control vs apical: $p \approx 0.00065$). These results indicate that

synaptic input from the basal tree is critical for orientation tuning, while apical tree synaptic

inputs play a less important role.

Finally, we assess the impact of sodium conductance nullification (as in Fig. 3), this time on

orientation tuning rather than somatic spiking. In line with our previous results (Fig. 3), we

find that sodium conductance nullification in the apical tree completely abolishes orientation

tuning (*Figure 4C*) while the same manipulation in the basal tree has no impact on tuning

(dependent two-sample t-test, control vs basal: $p \approx 0.24125$).

3.4.1 Sensitivity Analysis

To assess whether the above findings are robust to biologically-relevant variability in the

conductance values of sodium channels, we performed a sensitivity analysis whereby we

reduced the sodium conductance only in the apical tree by either 5% or 10% (*Supplementary*

Figure 7). For a decrease of 5%, we found that elimination of the apical but not the basal

sodium component results in total loss of tuning (dependent two-sample t-test, control vs

basal p-value: 0.32549). Importantly, in this scenario the apical tree remains the primary

driver of somatic activity as the majority of somatic spikes are apically driven (*Supplementary*

Figure 7A, N = 345 spikes; 60±15% of spikes are apically-driven, 30±18% are basally-driven,

and 10±8% of spikes are cooperatively-driven, for details on the characterization of spiking

activity, see Results, section 3.3). However, a 10% decrease in the apical sodium conductance

results in total loss of tuning when either the apical or the basal sodium component is

eliminated (*Supplementary Figure 7B*). In this case, a large number of spikes are completely

lost (N = 190 spikes vs 345), and out of the remainder, 41±25% are apically-driven, 48±28%

are basally-driven, and 11±12% are cooperatively-driven. We further compare these two

conditions (5% and 10% reduction) with one where the apical sodium conductance is

unchanged (*Supplementary Figure 7C,* identical to *Figure 4C*). Similarly to the results seen in

the 5% reduction, elimination of the apical - but not the basal - sodium component results in

total loss of tuning (dependent two-sample t-test, control vs basal p-value: 0.24125). In this

case, 80±9% of spikes are apically-driven, 13±8% are basally-driven, and 7±4% of spikes are

cooperatively-driven. Notably, the number of spikes is drastically higher in this condition (N =

885 spikes in the unchanged condition vs. N = 345 for the 5% reduction and N = 190 for the

10% reduction).

The above simulations show that, as expected, the reduction in apical sodium conductance

results in a decrease of the overall spiking rate of the neuron (number of spikes: 885 in the

unchanged condition, 345 for the 5% reduction, and 190 for the 10% reduction). Even so,

decreasing apical sodium conductance by 5% failed to negate its role in somatic spike

generation, with the apical tree remaining the primary contributor (60±15% of spikes were

apically-driven vs 30±18% basally-driven; *Supplementary Figure 7A*). A decrease of 10%,

however, exerts greater influence as an ever-larger number of apically-driven spikes are lost,

shifting the percentages to a more even distribution between apically-driven and basally-

driven (41±25% vs 48±28%, respectively; *Supplementary Figure 7B*). Of note, in this particular

case the neuronal firing rate is well below the experimentally-relevant range, suggesting that

such a reduction is not physiological (see *Supplementary Figure* 7A vs. 7B).   Overall, our

sensitivity analysis suggests a greater-than-expected contribution of apical sodium non-

linearities in somatic spike generation. This finding is in line with experimental work showing

that L2/3 V1 neurons can produce spikes even in the absence of feedforward (i.e. basal

stimulus-driven) activity (Keller et al., 2020).

Overall, our simulations predict that apical sodium channels are necessary for both spike

generation in general (*Figure 3)* as well as for orientation tuning in particular (*Figure 4C).*

Additionally, AMPA and NMDA activity is necessary for both apical and basal trees (*Figures*

4A, 4B). This means that the apical tree most likely exerts its influence on somatic spiking via

sodium-mediated spiking, whereas the basal tree mostly utilizes AMPA- and NMDA-driven

depolarizations, rather than sodium spikes.

Figure 4. Orientation Tuning Critically Depends on Apical Sodium Conductance and AMPA/NMDA Activity. A:

Orientation tuning curves (left) and mean OSI values (right) for the experiment where stimulus-driven synapses

are completely absent on either the apical ("apical target"; red) or basal ("basal target"; green) tree. B: Selective

reduction of stim-driven synaptic (AMPA/NMDA) weights by 50% 30 ms prior to somatic spike occurrence (lasting

up to 10 ms after the pre-recorded somatic spike timing), on either the apical (apical target; red) or basal (basal

target; green) tree. C: Nullification of sodium conductances 3 ms prior to somatic spike occurrence (lasting up to

1 ms after the pre-recorded somatic spike timing), on either the apical (apical target; red) or basal (basal target;

green) tree. Error bars: Standard error of the mean, for all panels. If present, "n/a" denotes a lack of orientation

3.5. Tree-Specific Morphological and Electrophysiological Properties Influence Dendritic and Somatic Behavior

Our simulations predict that apical and basal dendritic segments exert causal influence on

somatic spike generation via separate mechanisms. In an attempt to explain these differences

in contribution, we looked at the anatomical and electrophysiological properties of the two

types of dendrites (apical/basal). Specifically, we searched for links between

anatomical/electrophysiological properties of dendrites and (a) signal attenuation or (b)

dendritic excitability (number of synapses required to drive a local sodium spike, or

"threshold"; see also *Extended Methods,* section 8.5.9 for details).

To ensure a fair comparison among different dendritic branches, we induced a 20 mV local

EPSP (actual amplitudes: 19.78 ± 0.98 mV) and assessed its attenuation at the soma (see

Extended Methods, section 8.5.9). We found that EPSP attenuation is greater in apical versus

basal dendritic segments of similar length (*Figure 5A;* p-value 0.000002), diameter (*Figure 5B;*

p-value 0.005864) or volume (data not shown; p-value 0.000001). When normalized by

distance from the soma (*data not shown*; p-value 0.42941), dendritic path volume (data not

shown; p-value 0.027934), or electrotonic constant (data not shown; p-value 0.666959), EPSP

attenuation did not significantly differ between apical and basal dendritic segments. We also

examined the relationship between apical vs. basal EPSP attenuation in respect to high apical

dendrite branching order (>3), as well as high apical dendrite distance from the soma (>182.7

μm). We found no significant difference between apical and basal dendrite EPSP attenuation

in respect to high apical dendrite branching order (*Supplementary Figure 8D*; apical

attenuation: 16.59±3.18 mV, basal attenuation: 13.96±2.8 mV, p-value 0.0956). A statistically

significant difference did emerge in the case of EPSP attenuation in respect to high apical

dendrite distance from the soma (*Supplementary Figure 8C*; apical attenuation: 17.64±1.84

mV, basal attenuation: 13.96±2.8 mV, p-value 0.0312) if an outlier was removed from the

basal dendrite attenuation data (see *Extended Methods,* section 8.5.10 for details).

We repeated this analysis for dendritic sodium spikes (*Supplementary Figure 8*) and found

similar results: spike attenuation is greater in apical versus basal dendritic segments, when

comparing apical and basal dendritic segments of approximately equivalent length

(*Supplementary Figure 8B;* p-value 0.000097), diameter (*Supplementary Figure 8A;* p-value

0.00646) or volume (data not shown; p-value 0.000002). There was no significant difference

in spike attenuation between apical and basal dendritic segments when controlling for

distance from the soma (data not shown; p-value 0.768547), dendritic path volume (data not

shown; p-value 0.061389), or electrotonic constant (data not shown; p-value 0.396982). As

above, we also examined dendritic spike attenuation at the soma in respect to high apical

dendrite branching order and high apical dendrite distance from the soma. There was no

significant difference between attenuation for apical vs basal sodium spikes in respect to

branching order (*Supplementary Figure 8F*; apical attenuation: 64.93±26.28 mV, basal

attenuation: 59.39±5.8 mV, p-value 0.2894), but a significant difference was present when

comparing attenuation in respect to distance from the soma (*Supplementary Figure 8E*; apical

attenuation: 70.94±22.85 mV, basal attenuation: 59.39±5.8 mV, p-value 0.0336), excluding an

outlier from the basal tree (see *Extended Methods,* section 8.5.10 for details).

The above suggest that signals originating in apical vs. basal dendritic segments with similar

morphological characteristics (length, diameter, volume), attenuate more than those

originating in basal dendritic segments. This is partly because most apical dendrites are

located further away from the soma compared to basal dendrites: comparison of EPSP and

sodium spike attenuation in respect to distance from the soma yields statistically significant

differences between basal and distant apical dendrites (*Supplementary Figure 8C&E*).

Finally, we also examined the relationship between dendritic sodium spiking thresholds and

the morphological and electrophysiological parameters stated previously. We found that

dendritic sodium spiking thresholds are significantly lower for apical versus basal dendritic

segments with approximately equivalent values of the electrotonic constant (*Figure 5C;* p-

value 0.001585), but this relationship is reversed for dendritic segments with similar volume

(*Figure 5D*; p-value 0.000002). On the contrary, apical and basal dendritic segments featuring

similar distances from the soma (data not shown; p-value 0.320436), dendritic path volume

(data not shown; p-value 0.418108), length (data not shown; p-value 0.407686) or diameter

(data not shown*;* p-value 0.592490) did not exhibit significant differences in their sodium

spiking thresholds.

Taken together, this analysis shows that - as expected - the attenuation of signals originating

in morphologically and electrophysiologically equivalent dendritic segments is significantly

larger for apical vs. basal dendrites, likely because most apical dendrites are located further

away from the soma. However, the threshold for inducing sodium spikes in apical dendritic

segments is significantly smaller compared to basal dendritic segments of similar electrotonic

isolation (as indicated by their electrotonic constant), making it easier to drive such spikes in

the apical compared to the basal tree. These findings provide a more intuitive explanation of

the greater role of sodium conductances in the apical vs. basal trees of our model neuron.

 Figure 5. Dendritic Morphological and Electrophysiological Properties Affect both Local and Somatic Behavior.

A: Left: Attenuation of a 20 mV dendritic EPSP (measured at the soma) as a function of dendritic segment length.

Right: Comparison of length-normalized EPSP attenuation at the soma between apical and basal dendritic

segments. Apical dendritic segments feature greater attenuation. B: Left: Attenuation of a 20 mV dendritic EPSP

(measured at the soma) as a function of dendritic segment diameter. Right: Comparison of diameter-normalized

EPSP attenuation at the soma between apical and basal dendritic segments. Apical dendritic segments feature

greater attenuation. C: Left: Dendritic sodium spiking thresholds as a function of the dendritic electrotonic

constant (ELC). Right: Comparison of ELC-normalized sodium spiking thresholds between apical and basal

dendritic segments. Apical dendritic segments feature lower thresholds. D: Left: Dendritic sodium spiking

thresholds as a function of dendritic volume. Right: Comparison of volume-normalized sodium spiking thresholds

between apical and basal dendritic segments. Apical dendritic segments feature greater thresholds, but also

much greater variability of thresholds. Stars denote statistical significance at a significance level (α) of 0.05 (*),

0.01 (**) or 0.001 (***), whereas "n.s." stands for "no statistically significant difference", i.e. p-value ≥ α.

497 **4. Discussion**

In this work, we used a detailed computational model to delineate the dendritic constituents

of neuronal computation in L2/3 V1 pyramidal cells. We found that the model exhibits a wide

range of non-linear behavior at the dendritic segment level (*Figure 1*) and can mimic the

activity patterns of neurons under spontaneous conditions *in vivo* (*Supplementary Figure 2*).

The model neuron exhibits robust orientation tuning in response to tuned input (*Figure 2A*),

and under conditions of dendritic disparity, the model still remains robustly tuned (OSI < 0.2

and tuning width > 80°) for non-extreme dendritic disparity conditions (disparity ≤ 50°).

Importantly, the neuronal orientation preference deviates from the expected one (*Figure 2D-*

F), indicating that apical and basal inputs interact in non-linear ways. This bias is dependent

on the distribution of synaptic inputs along the two trees (*Figure 2D-F*), it follows the

preference of basal tree inputs under physiological conditions (*Figure 2D*) and is in line with

prior work, whereby apical tree ablation had no impact on somatic orientation preference

(*Supplementary Figure 9;* Park et al., 2019). Interestingly, while the above would seem to

suggest a minor role for apical inputs in the orientation preference of our model cell (*Figure*

2D), we found that the orientation tuning of the neuron is greatly affected by sodium

conductances in the apical - but not the basal - tree (*Figure 4*). Interestingly, while basal

sodium channel conductances are not critically important (*Figure 4C*), AMPA and NMDA

conductances are necessary in both dendritic trees for orientation tuning (*Figures 4A&B*). This

indicates that depolarizations from the basal tree play a key role in shaping neuronal output,

even when basal sodium spikes do not. As such, neuronal output is a synergistic phenomenon

mostly between apical sodium spikes and basal depolarizations or sodium spikes. Our

simulations suggest that a form of *intra-tree dendritic cooperativity -* a synergistic effect

between apical and basal dendritic segments that exhibit activity in relative synchrony - might

be present in these neurons. This is supported by the existence of extinction-prone

"cooperative" somatic spikes that are lost when either of their dendritic components (apical

or basal, be that depolarization or spiking) is lost (*Figure 3*). Additional evidence is provided

by the experiments in which removal or suppression (decrease in conductance) of basal

stimulus-driven synapses (i.e. only background-driven synapses are fully active) significantly

decreases the quality of orientation tuning (i.e. OSI values) in the model neuron, or abolishes

tuning altogether (*Figures 4A&B*).

A potential explanation for the differential contributions of apical and basal dendrites lies

with their morphological and electrophysiological properties. Despite featuring a uniform

sodium conductance throughout both dendritic trees, basal dendrites have higher sodium

spiking thresholds than apical dendrites with similar electrotonic constant values. Moreover,

apical dendrites have highly variable spiking thresholds (*Figure 5D*). This is likely because the

soma acts as a current sink, effectively increasing the sodium spiking threshold of all soma-

proximal branches (mostly basal dendrites). Meanwhile, the majority of apical dendrites

feature much better electrical isolation from the soma due to distance and branching, and

are as such exempt from such current sink effects, instead only suffering a loss of current from

other connected apical dendritic branches. As such, they exhibit a much greater range of

possible thresholds compared to basal dendrites of similar volume (*Figure 5D*), while

simultaneously featuring greater signal attenuation (*Supplementary Figure 8C&E*). Put

together, these results indicate that apical dendrites overcome their signal attenuation deficit

(compared to basal dendrites) by having a greater propensity for dendritic sodium spike

generation for different amounts of input. This suggests that sodium spikes are more difficult

to induce in basal compared to apical dendrites, offering a potential explanation as to why

apical sodium conductances contribute more to somatic output.

Taken together, our model predicts that somatic output in L2/3 V1 pyramidal neurons is likely

to be determined through *Bimodal Input Coincidence*, a subcategory of coincidence detection

(*Abstract Figure*). Specifically, we propose that the basal tree receives visual feedforward

input, representing the information therein as a series of hyperpolarizations, depolarizations,

and occasional dendritic spikes. Thus, visual input creates a basal "backdrop" of

depolarizations that represents visual information. At the same time, predictive and

attentional signals from higher-order cortical areas reach the apical tree of the neuron,

causing the generation of dendritic spikes that propagate to the soma and are temporally

summed with any concurrent basal depolarization. In the event that visual input is non-

existent, or of a non-preferred orientation, the depolarization is minimal to none. Thus, the

apical dendritic spike will not be significantly augmented through summation and will fail to

produce a somatic response in the majority of cases. If there is visual input, however,

especially of an orientation matching the preferred orientation of the basal tree, the

aforementioned "backdrop" will include multiple subthreshold depolarizations, perhaps even

dendritic spikes. The apical dendritic spikes are thus likely to be temporally summed with

these depolarizations and generate a somatic spike. As such, even though most somatic spikes

will be generated through an apical tree dendritic spike, the cases in which this is possible in

the first place will be dictated by the backdrop of depolarizations (and/or spikes) provided by

the basal tree.

The findings presented here propose a more central role for the apical tree than previously

suggested. Specifically, in Park et al. (2019), we found that orientation selectivity was not

abolished by *in vivo* laser ablation of the apical tree in L2/3 mouse V1 pyramidal neurons,

remaining essentially unchanged following a recovery period of ~1 day. Importantly, the

model presented here reproduces the *in vivo* ablation findings of the aforementioned study

(see Extended Methods, section 8.3.8; *Supplementary Figure 9*). What is seen as a

contradiction is resolved by considering that in both Park et al. (2019) and the simulations

presented here, apical dendrite ablation is assumed to be accompanied by homeostatic

alterations that counteract the increase in input resistance from the loss of the apical tree.

These homeostatic mechanisms keep the firing rate of the neuron the same pre- and post-

ablation, as seen experimentally (Park et al., 2019). Thus, the contribution of the apical tree

to somatic spiking in the intact neuron can be obviated by homeostatic mechanisms following

apical tree ablation.

Our findings suggest that L2/3 V1 pyramidal cells may perform salient feature extraction from

visual input compatible with predictive coding (Rao & Ballard, 1999). When top-down signals

reflecting attentional or predictive processes reach the apical tree, they "highlight"

appropriate parts of the signal "landscape" generated by the visual input reaching the basal

trees through the generation of apical dendritic sodium spikes. This could result in a neuron

that is activated only when specific features are present in its receptive field, facilitating the

processing of salient stimuli. Such phenomena have been observed experimentally as a result

of attention (Simons & Chabris, 1999).

However, this work also suffers from a number of limitations. Our findings are inherently

bound to the particular characteristics of the simulated morphology. To ensure their

generality, further exploration of different neuronal morphologies of L2/3 V1 neurons is

required. However, given that our findings correlate with specific anatomical and

electrophysiological features of the model neuron (*Figure 5, Supplementary Figure 8*), it

would be possible to infer whether the behavior we observe in our model could be present

in other neuronal morphologies as well. In addition, while repeating our analysis using other

morphologies is beyond the scope of this study (validating model function and performing

the extensive parameter exploration reported here is highly resource-intensive), our code and

all analysis scripts are made publicly available, allowing interested scientists to apply our

approach to other morphologies. The sensitivity analysis that we added in Section 3.4.1

(Supplementary Figure 7) also further establishes the robustness of our findings within

biologically relevant ranges of ionic conductances. We have taken all these steps in order to

more conclusively demonstrate that our findings are likely to generalize to other L2/3 V1

pyramidal neurons. Furthermore, our model is limited by the lack of data on dendritic features

such as signal attenuation and distribution of ion channel conductances. However, this has

been counteracted at least to some extent by the use of experimental data (e.g. see

Supplementary Figure 2) and the sensitivity analysis, which demonstrates that our conclusions

are robust to biologically relevant variations of the key conductances (*Supplementary Figure*

7). Additionally, the single-neuron model sacrifices much of the complexity found in the actual

L2/3 network, especially concerning inhibitory inputs, which in our model are not patterned

according to different interneuron subtypes and are background-driven, lacking tuning and

featuring firing rates lower than stimulus-driven excitatory synapses. A similar complexity loss

is incurred by using a relatively simple stimulation protocol, compared to the alternating

stimulation of the "drifting gradient" protocol that is more widely used. Finally, the model

does not account for retinotopically shifted inhibitory inputs (e.g. Rossi et al., 2020).

Retinotopic shift in inhibitory inputs would not affect our results, since stimulation is implicitly

assumed to only take place at the center of the receptive field of the model neuron, a fact

which would render any retinotopically offset neurons quiescent. Additionally, although

inhibitory inputs could be biased towards one dendritic tree over the other, the overall effect

of such an imbalance would be a change in the net amount of excitation received by each tree

- a scenario which we already explore in part (*Figure 2* and *Figure 3*), finding tuning and

somatic spike generation to be generally unaffected. The omission of these features that were

unlikely to influence the presence or absence of orientation tuning was performed in an

attempt to reduce the complexity (and thus computational requirements) of the model.

In terms of questions that still need to be answered, the exact nature of the attentional and

predictive signals received by the apical tree remains to be deciphered. Secondly, the

formation of the visual "backdrop" as a result of basal tree depolarization merits further

study. Encoding of visual information using mostly subthreshold depolarizations in the

presence of noise is an interesting problem, possibly resolved through the effects of intra-

tree dendritic cooperativity, where apical spikes sharpen responses to stimulus-driven input

rather than noise, improving the signal-to-noise ratio (Poleg-Polsky, 2019). In addition, a

number of differences emerge when comparing the visual system across different model

species. In rodents, orientation-selective cells akin to V1 simple cells have been discovered in

the thalamus itself (Scholl et al., 2013), and direct thalamocortical projections to L1 of V1 have

also been observed (Roth et al., 2016). Furthermore, rodent V1 organization follows a

dispersed salt-and-pepper motif, unlike the highly structured organization typical of higher

mammals like primates and cats (Ohki & Reid, 2007). These differences are important to take

into account when trying to infer general rules of function from information derived from

rules of information processing in different animal models. Finally, perhaps the most

interesting unanswered question is whether these computations take place elsewhere in the

cortex in addition to the visual system. Predictive coding as a means of stimulus compression

has already been proposed as a way to simplify visual perception (Rao & Ballard, 1999), and

this might also be the case for other sensory or cognitive tasks. Regardless, further

investigation is required in order to unravel the Gordian knot that is visual perception.

5. Acknowledgments

This work was supported by the ERC Starting Grant "dEMORY: Dissecting the role of dendrites

in memory", ERC-2012-StG-311435, the Fondation Santé, a FORTH-Synergy Grant (EVO-

NMDA) and the NEUREKA grant H2020-FETOPEN-2018- 2019-2020-01 to YP, the NIH NEI R01

grant EY-024019 and NINDS R21 grant NS088457 to SS, a Hellenic Foundation for Research

and Innovation (HFRI) PhD Scholarship (no. 6731) to K-E.P, and the FORTH-Synergy Grant

(FlexBe) to AP.

6. References

1. Adesnik, H., Bruns, W., Taniguchi, H., Huang, Z. J., & Scanziani, M. (2012). A neural circuit for spatial summation in visual cortex. *Nature*, *490*(7419), 226–231. https://doi.org/10.1038/nature11526

2. Behabadi, B. F., Polsky, A., Jadi, M., Schiller, J., & Mel, B. W. (2012). Location-dependent excitatory synaptic interactions in pyramidal neuron dendrites. *PLoS Computational Biology*, *8*(7), 40–42. https://doi.org/10.1371/journal.pcbi.1002599

3. Chen, T.-W., Wardill, T. J., Sun, Y., Pulver, S. R., Renninger, S. L., Baohan, A., Schreiter, E. R., Kerr, R. A., Orger, M. B., Jayaraman, V., Looger, L. L., Svoboda, K., & Kim, D. S. (2013). Ultrasensitive fluorescent proteins for imaging neuronal activity. *Nature*, *499*(7458), 295–300. https://doi.org/10.1038/nature12354

4. Cho, K.-H., Jang, H.-J., Lee, E.-H., Yoon, S. H., Hahn, S. J., Jo, Y.-H., Kim, M.-S., & Rhie, D.-J. (2008). Differential Cholinergic Modulation of Ca2+ Transients Evoked by Backpropagating Action Potentials in Apical and Basal Dendrites of Cortical Pyramidal Neurons. *Journal of Neurophysiology*, *99*(6), 2833–2843. https://doi.org/10.1152/jn.00063.2008

5. Coogan, T. A., & Burkhalter, A. (1990). Conserved patterns of cortico-cortical connections define areal hierarchy in rat visual cortex. *Experimental Brain Research*, *80*(1), 49–53. https://doi.org/10.1007/BF00228846

6. Cruz-Martín, A., El-Danaf, R. N., Osakada, F., Sriram, B., Dhande, O. S., Nguyen, P. L., Callaway, E. M., Ghosh, A., & Huberman, A. D. (2014). A dedicated circuit links direction-selective retinal ganglion cells to the primary visual cortex. *Nature*, *507*(7492), 358–361. https://doi.org/10.1038/nature12989

7. Defelipe, J., Elston, G. N., Fujita, I., Fuster, J., Harrison, K. H., Hof, P. R., Kawaguchi, Y., Martin, K. A. C., Rockland, K. S., Thomson, A. M., Wang, S. S. H., White, E. L., & Yuste, R. (2002). Neocortical circuits: Evolutionary aspects and specificity versus non-specificity of synaptic connections. Remarks, main conclusions and general comments and discussion. *Journal of Neurocytology*, *31*(3-5 SPEC. ISS.), 387–416. https://doi.org/10.1023/A:1024142513991

8. DeFelipe, J., & Fariñas, I. (1992). The pyramidal neuron of the cerebral cortex: morphological and chemical characteristics of the synaptic inputs. *Prog Neurobiol*, *39*(6), 563–607. https://doi.org/10.1016/0301-0082(92)90015-7

9. Goetz, L., Roth, A., & Häusser, M. (2021). Active dendrites enable strong but sparse inputs to determine orientation selectivity. *Proceedings of the National Academy of Sciences of the United States of America*, *118*(30). https://doi.org/10.1073/pnas.2017339118

10. Haider, B., Häusser, M., & Carandini, M. (2013). Inhibition dominates sensory responses in the awake cortex. *Nature*, *493*(7430), 97–100. https://doi.org/10.1038/nature11665

11. Hines, M. L., & Carnevale, N. T. (2001). Neuron: A Tool for Neuroscientists. *The Neuroscientist*, *7*(2), 123–135. https://doi.org/10.1177/107385840100700207

12. Iacaruso, M. F., Gasler, I. T., & Hofer, S. B. (2017). Synaptic organization of visual space in primary visual cortex. *Nature*. https://doi.org/10.1038/nature23019

13. Jia, H., Rochefort, N. L., Chen, X., & Konnerth, A. (2010). Dendritic organization of

sensory input to cortical neurons in vivo. *Nature*, *16*(3), 236–237.
https://doi.org/10.1038/nature08947

14. Ju, N., Li, Y., Liu, F., Jiang, H., Macknik, S. L., Martinez-Conde, S., & Tang, S. (2020).
Spatiotemporal functional organization of excitatory synaptic inputs onto macaque
V1 neurons. *Nature Communications*, *11*(1). https://doi.org/10.1038/s41467-020-
14501-y

15. Keller, A. J., Roth, M. M., & Scanziani, M. (2020). Feedback generates a second
receptive field in neurons of the visual cortex. *Nature*, *582*(7813), 545–549.
https://doi.org/10.1038/s41586-020-2319-4

16. Ko, H., Hofer, S. B., Pichler, B., Buchanan, K. A., Sjöström, P. J., & Mrsic-Flogel, T. D.
(2011). Functional specificity of local synaptic connections in neocortical networks.
Nature, *473*(7345), 87–91. https://doi.org/10.1038/nature09880

17. Larkum, M. (2013). A cellular mechanism for cortical associations: An organizing
principle for the cerebral cortex. *Trends in Neurosciences*, *36*(3), 141–151.
https://doi.org/10.1016/j.tins.2012.11.006

18. Mrsic-Flogel, T. D., Hofer, S. B., Ohki, K., Reid, R. C., Bonhoeffer, T., & Hübener, M.
(2007). Homeostatic Regulation of Eye-Specific Responses in Visual Cortex during
Ocular Dominance Plasticity. *Neuron*, *54*(6), 961–972.
https://doi.org/10.1016/j.neuron.2007.05.028

19. Ohki, K., & Reid, R. C. (2007). Specificity and randomness in the visual cortex. *Current
Opinion in Neurobiology*, *17*(4), 401–407.
https://doi.org/10.1016/j.conb.2007.07.007

20. Pachitariu, M., Stringer, C., Dipoppa, M., Schröder, S., Rossi, L. F., Dalgleish, H.,
Carandini, M., & Harris, K. (2016). Suite2p: beyond 10,000 neurons with standard
two-photon microscopy. *BioRxiv*, *July*, 061507. https://doi.org/10.1101/061507

21. Palmer, L. M., Shai, A. S., Reeve, J. E., Anderson, H. L., Paulsen, O., & Larkum, M. E.
(2014). NMDA spikes enhance action potential generation during sensory input.
Nature Neuroscience, *17*(3), 383–390. https://doi.org/10.1038/nn.3646

22. Papadopoulos, P. M., Katz, M. J., & Bruno, G. (2003). NPACI rocks: Tools and
techniques for easily deploying manageable linux clusters. *Concurrency and
Computation: Practice and Experience*, *15*(7–8), 707–725.
https://doi.org/10.1002/cpe.722

23. Park, J., Papoutsi, A., Ash, R. T., Marin, M. A., Poirazi, P., & Smirnakis, S. M. (2019).
Contribution of apical and basal dendrites to orientation encoding in mouse V1 L2/3
pyramidal neurons. *Nature Communications*, *10*(1), 1–11.
https://doi.org/10.1038/s41467-019-13029-0

24. Poleg-Polsky, A. (2019). Dendritic spikes expand the range of well tolerated
population noise structures. *Journal of Neuroscience*, *39*(46), 9173–9184.
https://doi.org/10.1523/JNEUROSCI.0638-19.2019

25. Rao, R. P. N., & Ballard, D. H. (1999). Predictive coding in the visual cortex: A
functional interpretation of some extra-classical receptive-field effects. *Nature
Neuroscience*, *2*(1), 79–87. https://doi.org/10.1038/4580

26. Rossi, L. F., Harris, K. D., & Carandini, M. (2020). Spatial connectivity matches
direction selectivity in visual cortex. *Nature*, *588*(7839), 648–652.
https://doi.org/10.1038/s41586-020-2894-4

27. Roth, M. M., Dahmen, J. C., Muir, D. R., Imhof, F., Martini, F. J., & Hofer, S. B. (2015).
Thalamic nuclei convey diverse contextual information to layer 1 of visual cortex.

Nature Neuroscience, *19*(2), 299–307. https://doi.org/10.1038/nn.4197

28. Roth, M. M., Dahmen, J. C., Muir, D. R., Imhof, F., Martini, F. J., & Hofer, S. B. (2016). Thalamic nuclei convey diverse contextual information to layer 1 of visual cortex. *Nature Neuroscience*, *19*(2), 299–307. https://doi.org/10.1038/nn.4197

29. Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T., Preibisch, S., Rueden, C., Saalfeld, S., Schmid, B., Tinevez, J. Y., White, D. J., Hartenstein, V., Eliceiri, K., Tomancak, P., & Cardona, A. (2012). Fiji: An open-source platform for biological-image analysis. *Nature Methods*, *9*(7), 676–682. https://doi.org/10.1038/nmeth.2019

30. Scholl, B., Tan, A. Y. Y., Corey, J., & Priebe, N. J. (2013). Emergence of Orientation Selectivity in the Mammalian Visual Pathway. *Journal of Neuroscience*, *33*(26), 10616–10624. https://doi.org/10.1523/JNEUROSCI.0404-13.2013

31. Schuz, A., & Palm, G. (1989). Density of Neurons and Synapses. *Journal of Comparative Neurology*, *455*, 442–455.

32. Simons, D. J., & Chabris, C. F. (1999). Gorillas in our midst: sustained inattentional blindness for dynamic events. *Perception*, *28*(28), 1059–1074.

33. Smith, S. L., Smith, I. T., Branco, T., & Häusser, M. (2013). Dendritic spikes enhance stimulus selectivity in cortical neurons in vivo. *Nature*, *503*(7474), 115–120. https://doi.org/10.1038/nature12600

34. Spruston, N. (2008). Pyramidal neurons: dendritic structure and synaptic integration. *Nature Reviews Neuroscience*, *9*(3), 206–221. https://doi.org/10.1038/nrn2286

35. van Elburg, R. A. J., & van Ooyen, A. (2010). Impact of dendritic size and dendritic topology on burst firing in pyramidal cells. *PLoS Computational Biology*, *6*(5), 1–19. https://doi.org/10.1371/journal.pcbi.1000781

36. Wilson, D. E., Whitney, D. E., Scholl, B., & Fitzpatrick, D. (2016). Orientation selectivity and the functional clustering of synaptic inputs in primary visual cortex. *Nature Neuroscience*, *19*(8), 1003–1009. https://doi.org/10.1038/nn.4323

 7. Supplementary Information

Compartment Type	Passive/Active Mechanisms	Synaptic Mechanisms
Soma	Hodgkin/Huxley voltage-gated Na$^+$ channels	GABA$_A$ (background-driven)
Basal Dendrites	Hodgkin/Huxley voltage-gated K$^+$ channels Muscarinic voltage-gated K$^+$ channels A-Type voltage-gated K$^+$ channels	AMPA (background-driven) NMDA (background-driven) GABA$_A$ (background-driven)
Apical Dendrites	T-Type Ca^{++} channels High voltage activated (HVA) Ca^{++} channels Calcium-dependent K$^+$ channels Active ATP Ca^{++} pumps	AMPA (stimulus-driven) NMDA (stimulus-driven) GABA$_A$ (stimulus-driven)

787 ***Supplementary Table 1***. *Outline of passive, active and synaptic mechanisms present in the model neuron.*

Conductance (mS/cm^2)	Soma	Apical	Basal
g$_{Na}$	0.505	0.303	0.303
g$_{Kdr}$	0.05	1.5*10^{-3}	1.5*10^{-3}
g$_{Km}$	2.8*10^{-3}	1.27*10^{-3}	1.27*10^{-3}
g$_A$	5.4	Diameter ≤ 0.8 μm: 108 Diameter > 0.8 μm: 10.8	Diameter ≤ 0.8 μm: 108 Diameter > 0.8 μm: 10.8
g$_T$	0.03	x ≤ 260 μm: 0.029*sin(0.009*x+0.88) x > 260μm: 0.012	0.03+6*10^{-5}*x
g$_{HVA}$	0.05*10^{-3}	x ≤ 260 μm: 0.049*10^{-3}*sin(0.009*x+0.88) x > 260μm: 0.02*10^{-3}	0.05*10^{-3}+10^{-7}*x
g$_{KCa}$	2.1*10^{-3}	2.1*10^{-3}	2.1*10^{-3}

788 ***Supplementary Table 2***. *Outline of membrane mechanism conductances (not synaptic). Reproduced from* (Park

789 et al., 2019).

	Model	Cho et al., 2010
RMP, mV	-79	-78.56 ± 1.34
IR, MΩ	123.6	125.2 ± 8.2
τ, ms	17.3	16 ± 0.7
AP amplitude, mV	66.1	67.8 ± 1.8
AP threshold, mV	-41.8	-37.7 ± 1.3
AHP, mV	17.9	13.3 ± 0.5
P-T time, ms	38.6	55.3 ± 2.7
AP adaptation	1.16	1.18 ± 0.02

 Supplementary Table 3. *Outline of model electrophysiological properties. RMP: resting membrane potential, IR:*

 Input Resistance measured at hyperpolarizing current (-0.04 nA), AP: action potential, AHP: after

 hyperpolarization measured at depolarizing current (0.16 nA), P-T peak-trough. Reproduced from (Park et al.,

 2019).

	Conductance (nS)	τ_1, ms	τ_2, ms
NMDA	1.15	2	30
AMPA	0.84	0.1	2.5
GABA$_A$	1.25	0.2	1.4

 Supplementary Table 4. *Outline of synaptic mechanism conductances and time constants. Reproduced from*

 (Park et al., 2019).

 Supplementary Figure 1. *Visualization of dendritic non-linear properties. A: Expected vs Actual plot for all basal*

 dendrites undergoing the I3P protocol for 1-200 synapses. Blue line represents the dendrite in Figure 1B (basal

 dendrite #0). B: Same as A, but for all apical dendrites. Blue line represents dendrite in Figure 1C (apical dendrite

 #0). C: Example traces from the I3P protocol for NMDA spike generation from basal dendrite #0. Traces are 10

Supplementary Figure 2. Comparing Model Output with Two-Photon Imaging Data. *Simultaneous recordings*

from basal dendritic segments and somata of L2/3 V1 pyramidal neurons were performed in anesthetized mice

(spontaneous activity of 11 neurons from 2 mice, see Extended Methods, section 8.5; panel A). These imaging

data were compared to modeling data generated using the spontaneous activity protocol, whereby simulations

feature only background activity, i.e. noise, in the absence of a visual stimulus (see Extended Methods, section

8.3.5). A: Image from one of the neurons in the dataset. Note the separate, multiple ROIs per dendritic segment,

as well as the 4 circular ROIs used for background fluorescence calculation. B: Example traces from the model

data conversion pipeline. Voltage spiking data is binarized and convolved with a GcaMP6s kernel function

(Supplementary Figure 3). Resulting data is downsampled to two-photon sampling rates (~7.25 Hz) from the

original sampling rate (10 KHz), and processed exactly like in vivo *two-photon fluorescence data. Red vertical*

lines show the timing of the original spikes. C: Example somatic (top) and distal dendritic (bottom) $\Delta F/F_0$ traces

from the neuron in (A). Black dots at the origin of colored dashed lines denote detected events. Note that somatic

events (green dashed lines/arrows) are not necessarily coincident with dendritic events (red dashed

lines/arrows). D: Event-triggered averages for all somatic (blue) and basal dendritic segment ROIs (1 dt $\simeq$ 0.138

s; error bars: Standard error of the mean). E: Event pair timing difference histogram for the two-photon data.

48.12% of event pairs feature basal dendritic events that are decoupled from the soma (red; dendritic events

occur at least 1 dt before the onset of a somatic spike), with 27.8% being decoupled if the 4 bins (~552 ms) closest

to zero are excluded (excluded data marked as lightly shaded bars). F: Event pair timing difference histogram for

the converted model data. 48.79% of event pairs feature basal dendritic events that are decoupled from the soma

(red; dendritic events occur at least 1 dt before the onset of a somatic spike), with 24.3% being decoupled if the

4 bins (~552 ms) closest to zero are excluded (excluded data marked as lightly shaded bars). Note that in both D

and E, the zero-difference bin was removed, as it is assumed to contain the majority of backpropagating events.

The middle two bars in F have been truncated for visualization purposes; the corresponding values are (left to

right) 0.15958 and 0.16533, respectively. Some of the detected somatic events are not closely preceded by any

detected basal dendritic events (panel F, green bars), suggesting that other events, perhaps arising in the apical

tree or basal dendritic segments not in the recording field of view, contribute to somatic activity in such cases.

Supplementary Figure 3. *Properties of the GcaMP6s kernel. A: Illustration of the output of the kernel function*

for a single event (AP) occurring at t_0 = 500 ms. (marked as a red line). B: Response intensity as a function of the

number of events (APs) within a 250ms window. Compare with (Chen et al., 2013).

***Supplementary Figure 4.** Waveforms of detected calcium fluorescence events. Somatic waveforms in blue, basal*

dendritic waveforms in red. Spike onset detected at t = 5 dt. A: Waveforms from all detected events from both

mice. B: Waveforms from all detected events in mouse 1. C: Waveforms from all detected events from mouse 2.

Note: 1 dt is equal to approximately 0.138 s, and exactly equal to the microscope recording frame time.

Supplementary Figure 5. *A: Event-triggered averages for somatic (blue) and basal dendrite (red) events in mouse*

1. B: Event-triggered averages for somatic (blue) and basal dendrite (red) events in mouse 2. Note the difference

in fluorescence intensity compared to mouse 1. Error bars: Standard error of the mean. C: Comparison of the true

(two-photon) data distribution (blue) with the null distribution (light red). Distributions are different (K.S. two-

sample test; test value ~0.105, p << 0.05). Note that the middle bar for the true data has been truncated (actual

value 0.12267).

Supplementary Figure 6. *Example traces showing the spatiotemporal evolution of somatodendritic voltage before, during and after the generation of a dendritic-sodium-mediated somatic spike, across different dendritic paths. A: Visualization of basal dendrites. Colors correspond to traces in panels B and C. B: Voltage traces from the basal dendrites and soma (blue) around the occurrence of an apically-driven somatic spike. C: Voltage traces from the basal dendrites and soma (blue) around the occurrence of a basally-driven somatic spike. D:*

Visualization of a specific apical path. Colors correspond, in the same sequence (proximal to distal), to the traces

in panels E and F. E: Voltage traces from the soma (blue) and the apical dendritic segments shown in D around

the occurrence of an apically-driven somatic spike. F: Voltage traces from the soma (blue) and the apical dendritic

segments shown in D around the occurrence of a basally-driven somatic spike. G: As in D, but for a different apical

path. H: As in E, but for a different apical path. I: As in F, but for a different apical path. In all panels, traces are

ordered by distance of the recording point from the soma (closest to farthest). Red dots denote the maximum

voltage attained by the corresponding dendritic segment. Magenta dashed line indicates the timing of the

somatic spike. The soma (and corresponding somatic trace) is always marked in blue. Dendritic segments colored

grey are not inactive during these events, but their activity is not shown directly in these traces. Dashed arrows

indicate propagation of potential towards (dark green) or away from (dark red) the soma.

Supplementary Figure 7. *Results from experiments similar to Figure 4C, only with a reduction in apical sodium*

conductance. A. Apical g_{Na} decreased by 5%. Elimination of the apical sodium component results in total loss of

tuning. B. Apical g_{Na} decreased by 10%. Unlike A, elimination of the apical or basal sodium component results in

total loss of tuning. C. Apical g_{Na} is unchanged (control condition, identical to Figure 4C). Similar to A, elimination

of the apical sodium component results in total loss of tuning. If present, "n/a" denotes a lack of orientation

tuning for that specific experimental setup. Stars denote statistical significance at a significance level (α) of 0.05

(*), 0.01 (**) or 0.001 (***), whereas "n.s." stands for "no statistically significant difference", i.e. p-value ≥ α.

Dependent two-sample t-tests were used in all cases.

Supplementary Figure 8. *Results from analyses similar to Figure 5. Dendritic segments were stimulated with a*

number of synapses sufficient to elicit dendritic sodium spiking, and the attenuation at the soma was measured.

A: Somatic sodium spike attenuation as a function of dendritic segment diameter (left panel). Differences

between apical and basal dendrites can be evaluated by normalizing spike attenuation by the diameter of the

dendritic segment (right panel), showing significantly greater attenuation per μm for apical over basal dendrites

(p-value 0.006459). B: Somatic sodium spike attenuation as a function of dendritic segment length (left panel).

Differences between apical and basal dendrites can be evaluated by normalizing spike attenuation by the length

of the dendritic segment (right panel), showing significantly greater attenuation per μm for apical over basal

dendrites (p-value 0.000097). C: 20 mV EPSP attenuation as a function of distance of the dendritic segment from

the soma. D: 20 mV EPSP attenuation as a function of the branch order of the dendritic segment. E: As C, but for

sodium spikes rather than 20 mV EPSPs. F: As D, but for sodium spikes rather than 20 mV EPSPs. Stars denote

statistical significance at a significance level (α) of 0.05 (*), 0.01 (**) or 0.001 (***), whereas "n.s." stands for

"no statistically significant difference", i.e. p-value ≥ α.

Supplementary Figure 9. *Comparison of orientation tuning behavior seen in the model under control (blue) and*

apical dendrite ablation (red) configurations. Note that the y-axis shows the normalized response (firing rate

values divided by their corresponding maximum). Error bars: standard error of the mean.

8. Extended Methods

8.1. Model Description

8. Extended Methods

8.1. Model Description

The single neuron model was derived from the L2/3 V1 pyramidal cell model of (Park et al.,

2019) created in the NEURON simulation environment (Hines & Carnevale, 2001). It is a

morphologically detailed reconstruction, featuring 43 apical dendritic segments, 7 basal

dendritic segments and an axon. As the model makes use of a variety of passive and active

mechanisms (*Supplementary Tables 1-4*), its electrophysiological properties were validated

against experimental data (Park et al., 2019). Neither the morphology nor the biophysical

properties of the source model were altered, barring specific ion channel or synaptic changes

where explicitly specified.

The model features both excitatory and inhibitory synaptic input. Excitatory input was

subdivided into background-driven (noise) and stimulus-driven, the latter consisting of

orientation-tuned synapses. Inhibitory synapses are modeled as background-driven

(inhibitory noise) which alongside background-driven excitatory synapses (excitatory noise)

comprise the background-driven inputs. Stimulus-driven and background-driven synaptic

input to the neuron is distributed uniformly on the target compartments using a density of 2

synapses per μm (DeFelipe & Fariñas, 1992; Schuz & Palm, 1989). Activation of synapses

occurs via pseudo-randomly generated Poisson spike trains. Background-driven and stimulus-

driven synapses feature different peak Poisson activation frequencies (0.11 Hz and 0.3 Hz,

respectively) to produce the electrophysiologically-recorded firing rates (Adesnik et al., 2012;

Park et al., 2019). Stimulus-driven synapses feature variable activation frequencies, with a

maximum of 0.3 Hz, based on the angular distance of the stimulus from the preferred

orientation of each individual synapse. Unlike the original model by (Park et al., 2019), apical

stimulus-driven synaptic inputs have had a 10 ms delay added, to better reflect experimental

data on feedback synaptic inputs in V1 (Ju et al., 2020).

We modulate the proportion of stimulus-driven excitatory synapses on the apical versus the

basal tree in various experiments in order to examine the effects of synaptic input distribution

on somatic output. In all cases, the amount of inhibition at the soma is 7% of the total

inhibitory synapses (Defelipe et al., 2002). The default synapse distribution was 40% apical to

60% basal ("biologically plausible model"), in adherence with experimental data (excitation:

40% apical, 60% basal; inhibition: 33% apical, 60% basal, 7% soma) (Defelipe et al., 2002). The

"inverse biologically plausible" model, conversely, features a 60% apical to 40% basal

distribution (excitation: 60% apical, 40% basal; inhibition: 55.8% apical, 37.2% basal, 7%

soma). The "even" model features a balanced distribution of synaptic input across both trees

(even: excitation: 50% apical, 50% basal; inhibition: 46.5% apical, 46.5% basal, 7% soma). The

synaptic distribution of the "even" and "inverse biologically relevant" models are not derived

by experimental data, yet they allow for the investigation of apical and basal tree interactions

under conditions of balanced or imbalanced synaptic input. We also make use of other

distributions biased in either the apical or basal direction (referring to stimulus-driven

excitatory input; from 10% basal – 90% apical to 90% basal – 10% apical, in steps of 10%).

Similar to above, background-driven synapses closely follow the stimulus-driven synapse

distribution in these cases, except for background inhibition, which always features 7%

allocated to the soma.

In our model, orientation preference was assigned on two levels: synaptic (input/single

synapse) and dendritic (whole tree). Single synapse orientation preference is modeled as a

synaptic weight vector biased towards a predefined (preferred) orientation, which also

factors into the activation frequency of orientation-tuned synapses (Chen et al., 2013). Each

synapse is assigned an orientation preference, with values being 0° to 360° in steps of 10°.

This distribution of orientation preferences to the stimulus-driven synapses at the dendritic

tree level followed a Gaussian probability density function wrapped around the unit circle:

945
$$f(s) = \frac{1}{W\sqrt{2\pi}} \sum_{k=-1}^{1} e^{-\frac{(s-b+2\pi k)^2}{2W^2}}$$

where W is the standard deviation (30°; Chen et al., 2013), s is the synaptic orientation

preference and b is the mean of the distribution (apical or basal tree mean orientation

preference, ranging from 0° to 90° in steps of 10°). The parameter k controls the wrapping

around the unit circle, and takes the values [-1, -0.5, 0, 0.5, 1]. As a result, the mean

orientation preference of each dendritic tree is biased towards the chosen mean orientation

preference. By implementing excitatory stimulus-driven connectivity in this way, we also

emulate the "like-to-like" connectivity motifs seen *in vivo* (as seen in e.g. Ko et al., 2011; Rossi

et al., 2020), as (virtual) presynaptic neurons with orientation preferences close to the mean

orientation preference of the model neuron are more likely to connect to it, due to being

over-represented in the overall pool of possible synapses. Somatic/neuronal orientation

preference was evaluated using the approach described in section 2.3.3.

8.2. Biophysical Manipulations

The following manipulations aimed to evaluate somatic output under different sets of

conditions:

(1) Sodium (channel) blockage: selectively nullify (set to zero) sodium conductance (g_{Na}) in a

designated compartment (e.g. apical tree, specific basal dendritic segments, or soma).

(2) <u>Synaptic modulation</u>: selectively alter synaptic conductances (g_{AMPA}/g_{NMDA}) in a designated

compartment or dendritic tree.

(3) <u>Input manipulation</u>: selectively deactivate specific types of input (excitatory stimulus-

driven and background-driven).

8.3. Simulation Protocols

All simulations were performed on our High-Performance Computing Cluster (Rocks 7.0) with

624 cores and 3.328 TB of shared RAM under CentOS 7.4 OS (Papadopoulos et al., 2003),

through the NEURON simulation environment (Hines & Carnevale, 2001). Model neuron

output measurements were obtained at a sampling rate of 10 KHz (dt = 0.1 ms).

The following simulation protocols were implemented:

(1) <u>Iterative paired-pulse protocol (I3P)</u>: Used for the evaluation of dendritic non-linearities.

Following synaptic silencing and g_{Na} nullification in the entire model neuron excluding the

dendritic segment under investigation, a number (1-200, with step 1) of evenly spaced

(uniformly distributed) excitatory synapses are synchronously activated (2 pulses at 50 Hz).

Voltage at the midpoint of each dendritic segment was recorded to evaluate dendritic non-

linear behavior caused by sodium and NMDA spikes.

(2) <u>Stimulation protocol</u>: the model is simulated for 2500 ms (2.5 s), with oriented stimulus

presentation onset at 500 ms (0.5 s), lasting until the end of the simulation (duration 2 s).

(3) <u>Orientation tuning validation protocol</u>: Using the above stimulation protocol, we simulate

10 model "neurons" that feature different synaptic location patterns, defining 10 distinct

neuron IDs. For each one of these neuron IDs, we use 10 different permutations, with

different instances of the input spike trains, resulting in 10 different *simulation IDs* for each

neuron ID. Stimuli of 18 different orientations (0° to 170° in steps of 10°) are then presented

in separate simulations. The tuning curve is extrapolated by averaging firing rates across

simulation IDs for each neuron ID. The overall tuning properties (width, OSI) were then

computed through averaging responses across neuron IDs for each stimulus orientation.

Neuron IDs with an OSI of less than 0.2 or a tuning width greater than 80° (as determined by

a Gaussian fit) are excluded. If more than 20% of neuron IDs do not meet this criterion, the

specific parameter combination is considered to not result in orientation tuning. No neurons

were excluded based on the preceding criteria in our simulations.

(4) Orientation disparity protocol: To evaluate the contribution of the apical and basal trees

in the somatic orientation preference, we keep the mean apical tree orientation preference

fixed at 0°, and set the mean basal tree orientation preference to any one of 10 different

values (0° to 90° in steps of 10°) (see also section 8.1). Thus, we introduce orientation tuning

disparity between the two trees, ranging from 0° to 90°. Akin to the orientation tuning

protocol, we independently generate 10 different neuron IDs with 10 different simulation IDs

for each one and present stimuli as above (section 8.3.3). Orientation preference is then

calculated as described above (section 8.3.3). If more than 20% of the neuron IDs are rejected

(OSI < 0.2 or tuning width > 80°) for a given degree of orientation tuning disparity, then the

model as a whole is deemed to exhibit no orientation tuning for that level of disparity.

In order to determine whether the somatic response follows the mean apical or basal tree

preferred orientation most closely, we compute the *expected* orientation preference of each

neuron ID; that is, the orientation preference assuming a linear summation of the synaptic

inputs. Specifically, we record the assigned orientation preference of all synapses per neuron

ID, and then we use Euler's Formula to calculate the expected orientation preference for this

neuron ID:

1011
$$E_d = arctan\left(\frac{\sum_i sin(T_i^A) + \sum_j sin(T_j^B)}{\sum_i cos(T_i^A) + \sum_j cos(T_j^B)}\right)$$

1012

Where E_d is the expected orientation preference for a degree of disparity d, i and j are the

numbers of stimulus-driven apical and basal synapses respectively, and T^A and T^B are lists of

synaptic orientation preferences for the apical and basal trees, respectively. In this particular

protocol, a remapping of preferences is required in specific cases (e.g. 180° → 0°) since we

are not simulating direction selectivity, projecting orientation preferences into the [-90°, 90°)

interval.

(5) Spontaneous activity protocol: Similarly to the orientation tuning validation protocol, we

use 50 different neuron IDs with 100 different simulation IDs for each one. Simulations

feature no stimulus-driven synaptic activity, leaving only background-driven synapses active.

(6) Ionic intervention protocol: Using 50 neuron IDs with 10 different simulation IDs for each

one, prior knowledge of action potential occurrence is obtained via the stimulation protocol

(see section 8.3.2; presented stimulus: 0°, corresponding also to the mean apical and basal

tree orientation preference), using the 'biological plausible' synaptic distribution (section 8.1).

This protocol proceeds similarly to the stimulation protocol (with the notable exception of

somatic g_{Na} being set to 0 to prevent backpropagation) until a time before point t_i, which is

the time of somatic spike occurrence. At $t_i - 3$ ms, g_{Na} is set to 0 on either the apical or basal

tree dendrites in two separate simulations, until $t_i + 1$ ms. This manipulation selectively

eliminates the contribution of apical or basal dendritic sodium conductances (seen either as

depolarization or as dendritic spikes) to the somatic spike. This protocol is repeated for all

somatic spikes and the ensuing voltage traces are analyzed to ascertain whether the somatic

spikes survived the manipulation. Spikes rendered extinct only when intervening on the apical

tree are labeled as "apically-driven", while those rendered extinct when intervening on the

basal tree are labeled as "basally-driven". The spikes rendered extinct by either intervention

are labeled as "cooperative". We observed no cases of somatic spikes failing to become

extinct in at least one of the two interventions while using this protocol.

(7) <u>Synaptic intervention protocol</u>: This protocol proceeds similarly to the ionic intervention

protocol (note that somatic g_{Na} is not set to 0 in this protocol), with two main differences: (a)

the intervention reduces stim-driven AMPA and NMDA synapse conductances on either the

apical or basal tree by a specific percentage (e.g. 50%, 90% etc.), rather than setting g_{Na} to 0,

(b) the window of the intervention changes from $[t_i - 3$ ms until $t_i + 1$ ms] to $[t_i - 30$ ms until

$t_i +10$ ms] (10x increase in window size). This was done to compensate for the relatively low

peak activation frequencies of stim-driven synapses (0.3 Hz).

(8) <u>Ablation protocol</u>: Apical dendrites and their corresponding synapses are not generated

under this protocol, simulating a neuron 1 day after laser ablation of the apical dendrites (Park

et al., 2019). As firing rates greatly increase (more than 10-fold increase) under these

conditions, basal AMPA and NMDA synaptic conductances are reduced by 45% to bring the

firing rate back down to expected levels.

1050

8.4 Two-Photon Microscopy

(1) <u>Animals in Experiments:</u> All experimental protocols were approved by Brigham and

Women's Hospital (BWH) Institution Animal Care and Use Committee. Male and female wild-

type (C57BL/6) mice were purchased from The Jackson Laboratory and bred for experiments.

A total of 2 mice were used in these experiments.

(2) <u>Chronic Window Implantation and Sparse Labeling:</u> Viral vectors were purchased from

Addgene. For viral injections and chronic window implantation, 6-10 week old wild type

C57BL/6 mice (both male and female) were anesthetized with isoflurane (1-1.5%). The depth

of anesthesia was assessed via monitoring breathing rates. Meloxicam (2.5mg/kg) and

Buprenorphine (0.1mg/kg) were administered subcutaneously every 12 hours following the

surgical procedure for 2 days. A headpost was implanted on the skull and a 3mm diameter

craniotomy was made over the visual cortex of the left hemisphere. The craniotomy was

centered 2.7mm lateral to the midline and 1.5mm posterior to the bregma. To sparsely label

pyramidal neurons with GCaMP6s, a mix (~90 nl per site, up to 2 sites) of diluted CamKII-CRE

(AAV1, diluted 80,000X) and flex-GCaMP6s (AAV5, diluted up to 2X) was injected slowly over

5 minutes per penetration using a Drummond Nanoject III. Up to two penetrations ~0.5mm

apart on average were performed per craniotomy. After the viral injection, a round coverslip

was fitted to the craniotomy and sealed with vetbond and dental cement. Most chronic

windows in our hands remain clear for 2-3 months. 2-photon imaging was performed on week

3-4 following viral injection, at which time GCaMP6s expression was optimal.

(3) <u>In vivo calcium imaging of sparsely labeled neurons:</u> Three to four weeks after the

injection, each GCaMP6s expressing mouse was sedated with Fentanyl (0.5 mg kg^{-1}) and

Dexmedetomidine (0.5 mg kg^{-1}) (Mrsic-Flogel et al., 2007) for imaging experiments. A stable

level of anesthesia was confirmed by stable breathing rate and lack of movement. Neuronal

and dendritic calcium activity was recorded from L2/3 pyramidal neurons in V1. We chose

neurons with 1 to 3 primary basal dendritic segments clearly visible up to their primary

branching point on the same plane with the soma. Apical dendrites were not present in the

focal plane during recording. Mice were in the dark during the imaging process and their

spontaneous firing rate was recorded. Images were acquired at ~7.3 frames/sec using an

Ultima IV microscope in spiral scanning mode with a 25X 1.0 NA Nikon objective (5-30mW

laser power, 900 nm). Recordings lasted for 283 s (2750 frames). The size of the recording

FoV was 103 μm^2, and the pixel size was 0.201 μm^2. 12 pyramidal neurons were imaged, and

11 of these were selected for analysis. The final neuron was not selected due to an insufficient

number of basal dendritic segments being positively identified as belonging to said neuron

following visual inspection of a z-stack image.

1086

8.5. Data Analysis

Data analysis of both modeling and two-photon imaging experiments was performed through

Python 3.8+, using publicly available libraries as well as custom-made scripts and functions.

The latter include:

(1) <u>Dendritic Spike Detection</u>: Dendritic sodium spikes were identified via a detection

algorithm that identified depolarizations exceeding a -20 mV threshold. For the identification

of NMDA spikes in the I3P experiments (see section 8.3.1) we located the inflection points

(points where the second derivative is zero) of the voltage trace after the second pulse. As

the NMDA spike exhibits a voltage plateau, 2 or more inflection points indicate an NMDA

spike. As our voltage measurements are not continuous and the I3P protocol does not feature

noise, it is unlikely to encounter a measurement point at which the second derivative is

exactly zero. Thus, we assume an inflection point exists at some recording point P_n if and only

if $(P_{n-1} \cdot P_{n+1}) < 0$, where P_{n-1} and P_{n+1} refer to existing points in the second derivative of the

voltage trace, immediately preceding and following the theorized inflection point.

(2) <u>Quantification of Dendritic Nonlinearities</u>: Using data from the I3P experiments, we

calculated the maximum amplitude of dendritic segment depolarization per synapse count. A

linear input-output curve is then generated, extrapolating from the response of the dendritic

segment to a single synaptic input (unitary response). This curve is compared to the actual

response of the dendritic segment to increasing input in an "Expected vs. Actual" plot, thus

coarsely characterizing the integration mode of the dendritic segment. The integration mode

of the dendritic segments was quantified in detail using the Nonlinearity Relative to Linear

Extrapolation (NRLE) metric (Behabadi et al., 2012), which is defined as the maximum ratio of

actual to expected neuronal output signal. An NRLE value of less than 1 denotes a sub-linear

dendritic segment, NRLE of exactly 1 indicates a linear dendritic segment and NRLE values

over 1 characterize a dendritic segment as supra-linear. The non-linearity threshold of all

dendritic segments was also calculated, measured as the minimum number of synapses

required to elicit the corresponding electrogenic event (sodium or NMDA dendritic spike).

(3) <u>Two-Photon Imaging Data Analysis:</u> Recordings are motion corrected via the suite2p

software (Pachitariu et al., 2016) to accurately extract fluorescence values from dendritic

segments. In the case of somatic activity data, we use the original video, as motion correction

is not necessary at the level of magnification used. Using the Fiji ImageJ software (Schindelin

et al., 2012), ROIs are manually defined, using a maximum intensity z-stack as a reference

image. Dendritic segments are identified as belonging to the cell of interest via visual

inspection of a z-stack recording. Each dendritic segment is broken up into multiple ROIs. Only

one ROI is defined for the soma. We completely avoid areas where there is an overlap of the

cell of interest with nearby dendrites/axons of cells from outside the field of view. Dendritic

ROIs are slightly wider than the dendritic segments themselves, to partially account for

misplacement of the ROI or uncorrected movement (drift) of the dendrite. 4 equal-sized

circular ROIs are also defined in areas where no visible dendritic segments or somata can be

found to calculate the background fluorescence level.

(4) <u>Calcium Trace Processing:</u> Raw calcium fluorescence data is normalized via a moving

window average of the fluorescence trace (window size of 20 s), subtracting F_0 from it

(defined as the average of the 20% lowest fluorescence values of said moving 20s window),

then dividing by the same value (F0), finally producing a normalized $\Delta F/F_0$ value. These $\Delta F/F_0$

values were significantly different from the background $\Delta F/F_0$ as calculated from the 4 circular

reference ROIs (cross-correlation test; $p < 0.05$; no manually-defined ROIs were excluded

through this process).

(5) <u>Calcium Event Detection:</u> Calcium events are detected via a simple dual-threshold

approach: a calcium event is registered when the $\Delta F/F$ value exceeds the 93.5^{th} percentile at

the same time as the derivative of the $\Delta F/F$ value exceeds the 67^{th} percentile. These percentile

values were calculated via trial-and-error, attempting to match the expected firing frequency

of a neuron (soma) under spontaneous activity conditions (approximately 0.2 Hz; Haider et

al., 2013). To ensure that the algorithm returns true events rather than random noise, we

performed an Event-Triggered Average (ETA) analysis. We extracted all of the detected event

waveforms, centered them on the time of detection, and averaged across waveforms. We did

this separately for somatic and dendritic events (*Supplementary Figure 2C)*, as well as for each

animal separately (*Supplementary Figures 4A, 4B*). Calcium event-triggered averaging of the

processed fluorescence signal yields the expected temporal profile, indicating that these

events are not likely to represent noise.

(6) <u>Generation of "fluorescence-like" traces from model voltage traces:</u> 50 neuron IDs were

simulated with 100 simulation IDs each, using the "biologically plausible" configuration under

the spontaneous activity protocol, in order to closely match the properties of the real neurons

under spontaneous activity conditions. Next, we ran a spike detection algorithm to find all

somatic and dendritic spiking events. We binarized the traces, using 1 for spike peaks and 0

otherwise. We then convolved these binarized traces with a GcaMP6s kernel function of our

own design:

$$F(t) = -s \left(e^{-\frac{t-t_0}{t_r}} - e^{-\frac{t-t_0}{t_d}} \right)$$

1153

1154

Where t is time (in ms), s is a scaling factor (s ≈ 0.5714), t0 is the time of the event, t_r is the

rise time (t_r = 74 ms) and t_d is the decay time (t_d ≈ 401.45 ms). The output of the function is

unitless, representing fluorescence. With these parameters, our kernel function has an actual

rise time constant of 194.225 ms, with output values decaying by ~52.11% within 0.6 s

(*Supplementary Figure 3A*). These values closely match experimentally observed values for

GCaMP6s (*Supplementary Figure 3B;* Chen et al., 2013). The ensuing fluorescence traces were

then downsampled to a dt of 0.138 s (~7.25 Hz) from the original dt of 0.1 ms (10 KHz), to

emulate the sampling rate of our two-photon calcium fluorescence imaging. The transformed

traces were then processed by the same pipeline as true calcium fluorescence data (see

sections 8.5.4 and 8.5.5).

(7) Analysis and comparison of true vs "generated" fluorescence data: Voltage traces from

the soma and basal dendritic segments were converted into approximations of two-photon

fluorescence traces (see section 8.5.6) to allow a direct comparison with in vivo calcium

imaging data that were collected under similar conditions. Significant fluorescence increase

"events" were identified for both datasets using a custom detection algorithm (see section

8.5.5). To ensure that the algorithm detects true events rather than noise in the two-photon

signal, we performed an event-triggered averaging (ETA) analysis of trace segments centered

on the event onset timepoint (see section 8.5.5; 1131 dendritic events, 96 somatic events

across all 11 neurons and their respective dendritic ROIs). The results of this analysis indicate

that random noise is unlikely to be the cause of the detected events, as there is a marked

increase in fluorescence intensity at the time of event onset for all detected events

(*Supplementary Figure 2D, Supplementary Figures 4&5*) which persists over time.

Next, in order to infer the relationship between basal and somatic events, we looked at their

relative timing differences. Using dendritic events as the starting point, we paired each

dendritic event with the most temporally proximal somatic event, and computed the time

difference between the two for each event pair (measured in imaging frames, each lasting

approximately 138 ms). Using the same approach, we measured the fraction of somatic-

dendritic event pairs in our model and compared them to the in vivo data. Event pairs

featuring a temporal distance of zero (occurring within the same frame) were excluded as

they are most likely to represent backpropagating events (excluded event pairs: model:

219,291/245,717 event pairs, 89.25%; in vivo: 29,143/237,572 event pairs, 12.27%; difference

attributed to a lack of noise-related fluctuations in the model data compared with the in vivo

data). The resulting distributions (*Supplementary Figures 2E&F*) show that in both the model

and in vivo data local dendritic events can be generated independently of somatic events.

Paired events separated by more than 4 recording frames (~552 ms) are highly unlikely to be

the result of backpropagation, even when considering the slow kinetics of the calcium sensor.

Moreover, the somatic events that are preceded by basal dendritic events are similar in

proportion between datasets (*Supplementary Figures 2E&F*, red bins: 48.79% model, 48.12%

actual when including all event pairs with a non-zero timing difference; 24.3% model, 27.8%

actual when including only event pairs with an absolute timing difference >4 recording

frames).

To mitigate the risk that the in vivo findings are the result of noise or random chance, we

constructed a null distribution of timing differences for the in vivo data by independently

circularly shuffling the time-series data from the somatic and various dendritic segment ROIs

1000 times, and then re-computed the timing differences. We then compared the actual data

with the null distribution data using the two-sample Kolmogorov-Smirnov (KS) test, finding

that the two distributions of timing differences were significantly different (*Supplementary*

Figure 4C; test statistic ~0.105, p << 0.05), whereas a similar analysis of completely randomly

generated event timings would results in identical (not statistically different) distributions for

event pair timing differences. It is important to note that the analyses herein are inherently

subject to the noisy nature of the recordings, the limitations of the slow kinetics of the calcium

indicator, the low temporal resolution of the recording (frame time of ~138 ms), as well as

the possibility of undetected events.

(8) <u>Electrotonic constant calculation:</u> The electrotonic constants (mentioned also in text as

"ELC") of all dendritic segments were calculated using the formula from van Elburg & van

Ooyen (2010):

1211
$$\lambda_i = \sqrt{\frac{b_i r_m}{2 r_a}}$$

1212

Where i refers to the dendritic segment, λ_i is the electrotonic length constant, b_i is the radius

of the branch, r_m is the specific membrane resistance and r_a is the intracellular resistivity. To

derive the final value (electrotonic constant or "ELC"), we divide the length of each dendritic

section by its corresponding λ_i .

(9) <u>Quantification of Dendritic Impact on Somatic Output:</u> To ensure relative equivalence

between the levels of activity in each dendritic branch, we start by providing each dendritic

branch the minimum amount of synaptic input required such that each one features either

an EPSP of approximately 20 mV (19.78 ± 0.98 mV; *Figure 5A&B, Supplementary Figure 8C&D*),

or a dendritic sodium spike (*Supplementary Figure 8A&B, Supplementary Figure 8E&F*). Only

one dendritic branch received inputs each time, and somatic (but not dendritic) sodium

channels had their conductance set to zero in order to accurately measure signal attenuation.

After measuring the attenuation of this EPSP or spike at the soma, we attempted to describe

it as a function of different morphological and electrophysiological parameters by dividing the

attenuation observed at the soma by the corresponding value of the parameter for the given

dendritic branch. The parameters used were dendritic length, dendritic diameter, distance of

dendritic segment from the soma, dendritic volume, dendritic path volume (defined as the

sum of the volumes of all dendritic segments linking the given dendrite with the soma,

including the volume of the given dendrite) and dendritic electrotonic constant (see section

8.5.8). Values can then be compared across dendrite types (apical vs basal) for each

parameter by performing Welch's t-test. We also performed a similar analysis using the

dendritic sodium spiking threshold (*Figure 1E&F*) instead of dendritic signal attenuation, with

the same morphological and electrophysiological parameters as above (*Figure 5C&D*). To

ensure fairness of comparisons, we excluded from these analyses any dendrites belonging to

the apical trunk (4/43 apical dendritic segments), any apical dendrites with atypical properties

(4/43 apical dendritic segments featuring a relatively high sodium spiking threshold for their

morphology, produced as a side-effect of the 3d reconstruction; lengths: 16.2±10.5 µm;

diameters: 0.711±0.027 µm), as well as any apical dendrites whose value for the parameter

under examination was outside 3 standard deviations of the basal dendrite mean for the same

parameter. Note that the qualitative results of the analysis did not change even if no dendrites

were excluded.

(10) <u>Relationship of Branch Order and Distance with Signal Attenuation:</u> To check whether

apical and basal dendrites feature different levels of attenuation for 20 mV EPSPs or dendritic

sodium spikes (see section 8.5.9), we compared the levels of attenuation for these events

between basal dendrites and apical dendrites of high branch order (> 3) or high distance from

the soma (> 182.7 µm, equivalent to the maximum distance of any basal dendritic segment

from the soma). We excluded basal dendritic segment #3 (root branch of the bifurcating basal

dendrite) from the analysis, since it featured extremely low attenuation values because of

additional current received from its child branches. After isolating the EPSP and spike

attenuation data from the dendrites that matched the above criteria, we compared the

resulting sets of values using Welch's t-test.