Spatiotemporal Distribution of Vector Mosquito Species in Maricopa County, Arizona: The highest West Nile virus disease incidence area in the United States in 2021

André B. B. Wilke[1,*], Dan Damian[2], Agnese Zardini[3], Piero Poletti[3], Stefano Merler[3], John-Paul Mutebi[4], John Townsend[2], Marco Ajelli[1,*]

[1]Laboratory for Computational Epidemiology and Public Health, Department of Epidemiology and Biostatistics, Indiana University School of Public Health, Bloomington, IN, USA; [2]Maricopa County Environmental Services Department Vector Control Division, Phoenix, AZ, USA; [3]Center for Health Emergencies, Bruno Kessler Foundation, Trento, Italy; [4]Arboviral Diseases Branch (ADB), Division of Vector-Borne Diseases (DVBD), Centers for Disease Control and Prevention (CDC), Fort Collins, CO, USA.

*Corresponding authors: andwilke@iu.edu; majelli@iu.edu.

Abstract

Mosquito-borne diseases are a major global public health concern and mosquito surveillance systems are essential for the implementation of effective mosquito control strategies. The objective of our study is to determine the spatiotemporal distribution of vector mosquito species in Maricopa County from 2011 to 2021. The Maricopa County Mosquito Control surveillance system utilizes BG-Sentinel and EVS-CDC traps throughout the entire urban and suburban areas of the county. We estimated specific mosquito species relative abundance per unit area using the Kernel density estimator in ArcGIS 10.2. Our results show that vector mosquito species are widely distributed and abundant in the urban areas of Maricopa County. A total of 691,170 *Culex quinquefasciatus*, 542,733 *Culex tarsalis*, and 292,305 *Aedes aegypti* were collected from 2011 to 2022. The relative abundance of *Ae. aegypti* was highly seasonal peaking in the third and fourth quarters of the year. *Culex quinquefasciatus*, on the other hand, was abundant throughout the year with several regions consistently yielding high numbers of mosquitoes. *Culex tarsalis* was abundant but it only reached high numbers in well-defined areas bordering natural and rural areas. The well-defined species-specific spatiotemporal patterns found in this study can be used to develop and/or refine vector control strategies.

Keywords: *Aedes aegypti*, *Culex quinquefasciatus*, *Culex tarsalis*, Arbovirus

Introduction

Mosquito-borne diseases are a major global public health concern [1–4]. The incidence of dengue has increased substantially in the last decades, from 500,000 reported cases in 2000 to 2.4 million in 2010 and 5.2 million in 2019 [5]. The current estimate is 390 million dengue virus (DENV) infections worldwide every year [6]. Between 2015 and 2018, there were more than 1 million confirmed cases of Zika in the Americas followed by a significant increase in fetus malformation as a direct consequence of Zika virus (ZIKV) infection during pregnancy [3,4].

Albeit on a smaller scale, arboviral infections have also increased in the United States. The ZIKV was introduced multiple times in the United States in 2016 leading to 224 locally transmitted human cases in Florida and Texas [9–11]. Local transmission of the DENV has also been reported in Florida. In 2020, Monroe County experienced a major dengue outbreak with 65 confirmed locally transmitted cases. In the same year, 6 locally transmitted cases of dengue were reported in Miami-Dade County [12–14].

While local outbreaks of dengue and Zika were reported in the United States, the West Nile virus (WNV) has become endemic and the most widespread mosquito-borne arbovirus in North America [15]. WNV has been detected in more than 60 mosquito species in the continental United States (CONUS) [16], however, selected species in the genus *Culex* such as *Culex quinquefasciatus*, *Culex pipiens*, *Culex tarsalis*, and *Culex nigripalpus* are responsible for driving most of the epizootic and epidemic WNV transmission. Birds are the natural reservoir hosts of WNV as they are able to develop viremia high enough to infect mosquito vectors. Humans, horses, and other mammals

are not able to amplify the virus to viremic levels high enough to infect mosquitoes and

are therefore considered dead-end hosts [16,17].

Since it was first detected in New York in 1999 [18], the WNV has progressively

become endemic in all states in the contiguous United States and has been sporadically

detected in Alaska, Hawaii, and Puerto Rico [19]. From 1999 to 2020 a total of 52,532

human cases were reported in the United States leading to 2,456 deaths [20].

Maricopa County, Arizona is located in the Sonoran Desert ecoregion spreading

through 23,890 km^2. It has a hot desert climate with long hot summers and short winters

[21]. According to the 2020 census [22], Maricopa County has a population of 4,420,568

people, being the most densely populated county in Arizona as well as in the Sonoran

Desert ecoregion. There are three major cities in Maricopa County, Phoenix (the

Arizona state capital, and the fifth-most populous city in the United States), Mesa, and

Chandler. It has historically been one of the counties most affected by West Nile virus

disease in the United States with an average of 70 cases reported every year in the last

decade. However, 1,427 cases of West Nile were reported in this county in 2021, the

highest ever reported in a single county in the United States [12].

Controlling mosquito populations is widely considered the most effective way of

preventing the spread of mosquito-borne diseases [23]. However, the development and

implementation of effective mosquito control strategies in urban areas is complex and

difficult to put into action [24]. They rely on a framework consisting of multiple

successive actions that need to be implemented for a successful intervention. The first

step in the implementation of effective strategies is the development of an extensive

and reliable mosquito surveillance system [25]. Such a system is key to assessing the

mosquito community composition, identifying factors influencing the population

dynamics of mosquito species, estimating the relative abundance of populations of

different vector mosquito species, and providing evidence for ascertaining and

anticipating patterns in mosquito movement and demographics that are essential for the

development of mosquito control strategies. The aim of this study is to determine the

spatiotemporal distribution of vector mosquito species in Maricopa County, Arizona, by

analyzing mosquito surveillance data collected from 2011 to 2021.

Results

The Maricopa County Environmental Services Department Vector Control

Division Surveillance System was comprised of 478 traps in 2011 and new traps were

gradually included over the years as the surveillance system expanded (Figure 2). From

2016 to 2021 the number of traps remained similar with 781 and 833 traps, respectively.

A total of 691,170 *Cx. quinquefasciatus*, 542,733 *Cx. tarsalis*, and 292,305 *Aedes*

aegypti were collected from 2011 to 2022 by the traps in the Surveillance System

(Figure 1, Supplementary Table 1).

Figure 1. Number of active traps in the Maricopa County Environmental Services Department Vector Control Division Surveillance System (black line) and the number of mosquitoes collected by year from 2011 to 2021.

Over the study period, *Ae. aegypti* gradually expanded its range and relative abundance in the urban areas of Maricopa County. In particular, a substantial increase in the range of *Ae. aegypti* was observed in 2021 in relation to 2020, when a total of 88,187 specimens were collected by the traps in the surveillance system (Figure 2A). In contrast to *Ae. aegypti*, *Cx. quinquefasciatus* was already commonly found in the urban areas in Maricopa County in 2011. However, since 2011, *Cx. quinquefasciatus* became

more confined to specific areas of the county and is particularly abundant in those

areas. Since 2016, *Cx. quinquefasciatus* started to increase in abundance, especially in

the southeastern region of Maricopa County (Figure 2B). Although *Cx. tarsalis* could be

detected in urban areas, this species was mainly confined to specific areas of Maricopa

County, primarily to areas inside or bordering natural or rural areas. By 2021, *Cx.*

tarsalis had gradually expanded its range, but it remained highly abundant in this

ecological niche (Figure 2C, Supplementary Tables 2 and 3).

Figure 2. Kernel density-based heat map of the relative abundance of vector mosquito species collected by the Maricopa County Environmental Services Department Vector Control Division Surveillance System from 2012 to 2021. The color gradient represents the sum of mosquitoes collected by each trap in a 12-month period; (A) *Aedes aegypti*; (B) *Culex quinquefasciatus*; and (C) *Culex tarsalis*. The figure was produced using ArcGIS 10.2 (Esri, Redlands, CA), using layers available at

the   Maricopa   County   GIS   Mapping   Applications   available   at:

https://www.maricopa.gov/3942/GIS-Mapping-Applications.

Our results show great seasonal variations in the mean numbers of mosquitoes

collected in Maricopa County. Rainfall is one of the most important drivers for the

proliferation of mosquitoes in the region, and the peaks in abundance of *Cx.*

quinquefasciatus and *Cx. tarsalis* overlap with the peak of the rainy season in the

Spring and later in early Fall. Furthermore, rainfall is potentially an important driver for

the increase in the abundance of *Ae. aegypti* over the year until the beginning of the dry

season, when the abundance of *Ae. aegypti* populations is drastically reduced. Each of

the three vector mosquito species, *Ae. aegypti*, *Cx. tarsalis*, and *Cx. quinquefasciatus*,

displayed different population dynamic patterns peaking in abundance at different times

of the year indicating that even though rainfall is an important driver for their

proliferation, each species responded differently to the rainfall (Figure 3).

Figure 3. Mean number of *Aedes aegypti*, *Culex quinquefasciatus*, and *Culex tarsalis* collected by the Maricopa County Environmental Services Department Vector Control Division Surveillance System from 2016 to 2021. Solid line represents the monthly mean average temperature (Fahrenheit) between 2016 and 2021; Dashed line represents the monthly mean average rainfall (inches) between 2016 and 2021.

The mean number of collected *Ae. aegypti* varied greatly within each year. It was virtually absent during the first quarter of the year yielding a low number of mosquitoes collected by the traps in the surveillance system. It then gradually increased in abundance and range during the second quarter, showing the highest abundance in the southeastern part of the county. This species peaked in abundance during the third

quarter of the year covering most of the urban areas of Maricopa County and reaching

high abundance in several geographical areas. The mean number of *Ae. aegypti*

collected in the fourth quarter of the year was smaller than in the third quarter albeit

areas with high abundance were still present (Figure 4A).

Culex quinquefasciatus was abundantly found in Maricopa County year-round.

The mean number of *Cx. quinquefasciatus* was the highest during the second and third

quarters of the year with many areas of the county yielding an average higher than 10

mosquitoes per trap per day. One region bordering rural and leisure areas in the

southeastern part of the county yielded more than 10 mosquitoes per trap per day on

average, the whole year round (Figure 4B). *Culex tarsalis* was moderately affected by

seasonality. The mean number of *Cx. tarsalis* remained constant during the second,

third, and fourth quarters of the year with specific areas yielding high numbers of

mosquitoes. During the first quarter collections of *Cx. tarsalis* were not as abundant as

in the other quarters of the year, but for one region bordering rural and leisure areas in

the southeastern part of the county. Furthermore, *Cx. tarsalis* yielded the highest

average number of mosquitoes collected in a selected group of traps bordering natural

and rural areas. Even though the mean number of *Cx. tarsalis* reached high levels in

those specific areas, it was not abundantly found in urban areas. (Figure 4C).

Figure 4. Kernel density-based heat map of the mean number of *Aedes aegypti*,

Culex quinquefasciatus, and *Culex tarsalis* collected by the Maricopa County

Environmental Services Department Vector Control Division Surveillance System

from 2016 to 2021. The color gradient represents the mean number of mosquitoes

collected by each trap in a 3-month period. The figure was produced using ArcGIS 10.2

(Esri, Redlands, CA), using layers available at the Maricopa County GIS Mapping

Applications available at: https://www.maricopa.gov/3942/GIS-Mapping-Applications.

Discussion

Anthropogenic changes in the environment create favorable conditions for the

development of vector mosquito species that can exploit the resources available in

urban areas [26]. The decreased biodiversity in those urban areas increases the contact

rate between mosquito vector species and human hosts, increasing the risk of arbovirus

transmission to humans [14,27,28]. Our results show that vector mosquito species are

widely distributed and abundantly found in the urban areas of Maricopa County,

Arizona. *Aedes aegypti* gradually become more abundant and expanded its range since

only one specimen was collected in 2011. In 2012, 1,189 *Ae. aegypti* were collected; by

2014 that number increased to 27,208, and by 2021 a total of 88,187 specimens were

collected indicating a substantial increase in its relative abundance over the years. A

similar phenomenon could be observed for *Cx. tarsalis* and *Cx. quinquefasciatus*, and

although they were already present in great numbers since 2011, with more than 20,000

specimens from each species collected that year, reaching 89,401 collected specimens

of Cx. tarsalis and 94,774 of *Cx. quinquefasciatus* in 2021.

The high abundance of vector mosquitoes in Maricopa County represents a

potentially dangerous scenario. *Aedes aegypti* is the primary vector of dengue,

chikungunya, yellow fever, Zika viruses, and *Cx. quinquefasciatus* and *Cx. tarsalis* are

vectors of the WNV, Saint Louis Encephalitis, Eastern Equine encephalitis, and

Japanese Encephalitis viruses [29]. The close proximity of these mosquito vector

species to human populations increases the risk of arbovirus transmission as observed

in the WNV disease outbreak in Maricopa County resulting in approximately 1,500

cases of WNV disease in 2021 [12].

The relative abundance of *Ae. aegypti* was highly seasonal with fewer

mosquitoes collected in the first 3 months of the year followed by a moderate increase

in numbers in the second quarter indicating that the environmental conditions were not

conducive to the proliferation of *Ae. aegypti* during the first quarter of the year. However, as the year progressed a substantial increase in the abundance of *Ae. aegypti* was observed in the third and fourth quarters. The observed seasonal variation in the abundance of *Ae. aegypti* can potentially be an important driver for arbovirus transmission in Maricopa County as its abundance and range increased as the year progressed, therefore, potentially increasing the contact rate between *Ae. aegypti* and human hosts.

Culex quinquefasciatus, on the other hand, was abundantly found during all seasons of the year. It was not as abundant in the first quarter as it was during the remaining of the year, but it could be found throughout the urban areas during that time of the year. Several regions consistently yielded high numbers of *Cx. quinquefasciatus* mosquitoes indicating that those areas were hot spots and were able to support high numbers of *Cx. quinquefasciatus* the whole year-round. *Culex tarsalis* was also abundantly found in Maricopa County but in contrast to *Ae. aegypti* and *Cx. quinquefasciatus*, it only reached high numbers in well-defined areas bordering natural and rural areas. However, this is not surprising because *Cx. tarsalis* larval habitats include irrigation agricultural runoff, wetlands, sewage effluent, oil field run-off, and lake beds [30], all these habitat types are rarely associated with urban areas. Human cases of WNV disease have been consistently reported in Arizona, being one of the most affected states in the United States. The high abundance of *Culex* vector species, especially the year-round presence of *Cx. quinquefasciatus*, has major implications for the likelihood of WNV transmission in Maricopa County as was the case observed in

2021, when approximately 1,500 human WNV disease cases were reported in the
county [12].

Our results showed well-defined spatiotemporal patterns for each of the species
included in this study. However, the main drivers for the variations in the population
dynamics of each mosquito vector species are still not well understood and determining
how these species are driven by the availability of resources at the microgeographic
scale is essential to improving the effectiveness of mosquito control operations [31].
This is especially important for the development of control strategies targeted to reduce
the abundance of *Cx. tarsalis*, and to a lesser extent *Cx. quinquefasciatus*, since the
analysis of their population dynamics revealed a high abundance of both species in
well-defined areas. Furthermore, there is a need to assess and validate the
effectiveness of different control methods and strategies aimed to reduce mosquito
populations in urban areas focusing on how the proliferation of vector mosquito species
can be locally reduced by source reduction, community engagement, and by modifying
specific features of the urban built environment that facilitate the proliferation of vector
mosquito species [31–34]. Developing and implementing effective mosquito control
strategies is a difficult task with many factors that can potentially hinder the
effectiveness of mosquito control interventions, such as levels of insecticide resistance,
diel activity of target mosquito species, and outdated and ineffective mosquito control
tools [35–37].

This study is not without limitations. Mosquitoes were collected using BG-
Sentinel and CDC traps. Even though these traps are the gold standard for collecting
mosquitoes in urban areas, they may have underestimated the presence of less

anthropophilic mosquito species. Furthermore, new traps were gradually added during

the period of this study increasing the sampling effort over time, and even though we

were able to show an increase in the relative abundance of mosquitoes it is a limitation

in our study.

Effective mosquito control operations rely on multiple factors, with the most

important being consistent and reliable mosquito surveillance systems, community

engagement, and targeted source reduction. Now, more than ever, effective mosquito

control strategies are needed to curb the increase of mosquito-borne disease

outbreaks. Effective mosquito surveillance systems are key for guiding and

implementing mosquito control strategies and should be considered a priority for local,

state, and federal public health professionals.

Methods

The Maricopa County Environmental Services Department Vector Control

Division surveillance grid covers the entire urban and peri-urban areas of the county

with approximately one trap per square mile. The surveillance system is comprised of

BG-Sentinel traps baited with BG-Lure (Biogents AG, Regensburg, Germany) (ranging

from 13 traps in 2016 to 8 traps in 2021) and EVS-CDC traps baited with CO_2 (ranging

from 478 traps in 2011 to 825 traps in 2021). Each trap is deployed once a week for 24

hours for 50 weeks a year (Figure 5). All collected mosquitoes are transported to the

Maricopa County Environmental Services Department Vector Control Division

Laboratory and morphologically identified to species using taxonomic keys [38]. BG-

Sentinel traps baited with BG-Lure and EVS-CDC traps baited with CO_2 attract female

mosquitoes seeking hosts for blood-feeding [39]. Even though male mosquitoes were

sometimes collected, male data was not included in the proposed analyses.

We used the geodesic method for densities in square kilometers to calculate the

magnitude-per-unit area (number of inputs per point - i.e., mosquito relative abundance)

using the Kernel density estimator in ArcGIS 10.2 (Esri, Redlands, CA). Figures were

created with layers available at the Maricopa County GIS Mapping Applications

available at: https://www.maricopa.gov/3942/GIS-Mapping-Applications.

Figure 5. Map of Maricopa Mosquito Control Surveillance System. BG-Sentinel

traps are shown in black; EVS-CDC traps are shown in red.

Supplementary Table 1. Number of traps and mosquitoes collected by the

Maricopa County Environmental Services Department Vector Control Division

Surveillance System from 2011 to 2021.

Supplementary Table 2. Number of mosquitoes (*Aedes aegypti*, *Culex*

***quinquefasciatus*, and *Culex tarsalis*) collected per trap.**

Supplementary Table 3. Number of mosquitoes (*Aedes aegypti*, *Culex*

***quinquefasciatus*, and *Culex tarsalis*) collected by the 478 traps in use since 2016.**

Acknowledgments

References

References

1.   Messina JP, Brady OJ, Pigott DM, Brownstein JS, Hoen AG, Hay SI. A global
compendium of human dengue virus occurrence. Sci Data. 2014;1: 140004.

2.   Brady OJ, Hay SI. The global expansion of dengue: How *Aedes aegypti*
mosquitoes enabled the first pandemic arbovirus. Annu Rev Entomol. 2020;65:
191–208.

3.   Keyel AC, Raghavendra A, Ciota AT, Elison Timm O. West Nile virus is predicted
to be more geographically widespread in New York State and Connecticut under
future climate change. Glob Chang Biol. 2021;27: 5430–5445.

4.   Dorigatti I, Morrison S, Donnelly CA, Garske T, Bowden S, Grills A. Risk of
yellow fever virus importation into the United States from Brazil, outbreak years
2016–2017 and 2017–2018. Sci Rep. 2019;9: 1–11.

5.   World Health Organization. Dengue and severe dengue. Available at:
https://www.who.int/news-room/fact-sheets/detail/dengue-and-severe-dengue

6.   Bhatt S, Gething PW, Brady OJ, Messina JP, Farlow AW, Moyes CL, et al. The
global distribution and burden of dengue. Nature. 2013;496: 504–7.

7.   World Health Organization. Zika cases and congenital syndrome associated with
Zika virus reported by countries and territories in the Americas (Cumulative
Cases), 2015–2017. Available at: https://www.paho.org/hq/index.php?option =
com_content&view = article&id = 12390:zika-cumulative-cases&Itemid =
42090&lang = en.

8. Shapiro-Mendoza CK, Rice ME, Galang RR, Fulton AC, VanMaldeghem K,

Prado MV, et al. Pregnancy outcomes after maternal Zika virus infection during

pregnancy — U.S. Territories, January 1, 2016–April 25, 2017. Morb Mortal Wkly

Rep. 2017;66: 615–621.

9. Grubaugh ND, Ladner JT, Kraemer MUG, Dudas G, Tan AL, Gangavarapu K, et

al. Genomic epidemiology reveals multiple introductions of Zika virus into the

United States. Nature. 2017;546: 401–405.

10. Likos A, Griffin I, Bingham AM, Stanek D, Fischer M, White S, et al. Local

mosquito-borne transmission of Zika virus — Miami-Dade and Broward Counties,

Florida, June–August 2016. Morb Mortal Wkly Rep. 2016;65: 1032–1038.

11. Centers for Disease Control and Prevention. Zika cases in the United States.

Available at: https://www.cdc.gov/zika/reporting/index.html.

12. Centers for Disease Control and Prevention. Imported human disease cases

reported to CDC by county of residence. Available at:

https://wwwn.cdc.gov/arbonet/Maps/ADB_Diseases_Map/index.html

13. Florida Department of Health. Available at: http:// www. flori dahea lth. gov/ disea

ses- and- conditions/ mosqu ito- borne- disea ses/_docum ents/ alert- dade- wnv-

human- 10- 19- 20. pdf.

14. Coatsworth H, Lippi CA, Vasquez C, Ayers JB, Stephenson CJ, Waits C, et al. A

molecular surveillance-guided vector control response to concurrent dengue and

West Nile virus outbreaks in a COVID-19 hotspot of Florida. The Lancet Regional

Health – Americas. 2022;11: 100231

15. Ronca SE, Ruff JC, Murray KO. A 20-year historical review of West Nile virus

since its initial emergence in North America: Has West Nile virus become a

neglected tropical disease? PLoS Negl Trop Dis. 2021;15: e0009190

16. Centers for Disease Control and Prevention. Mosquito species in which West

Nile virus has been detected. Available at:

https://www.cdc.gov/westnile/resources/pdfs/Mosquito%20Species%201999-

2012.pdf

17. Centers for Disease Control and Prevention. West Nile Virus in the United

States: Guidelines for surveillance, prevention, and control. Available:

https://www.cdc.gov/westnile/resources/pdfs/wnvGuidelines.pdf

18. Hile JA, Mangiafico JA, Clippinger TL, Raphael BL, Smith JF, Danner DK, et al.

An outbreak of West Nile virus in a New York City captive wildlife population. Am

J Trop Med Hyg. 2002;67: 67–75.

19. Centers for Disease Control and Prevention. West Nile virus disease cases

reported to CDC by state of residence, 1999-2020. Available at:

https://www.cdc.gov/westnile/statsmaps/cumMapsData.html.

20. Centers for Disease Control and Prevention. West Nile virus neuroinvasive

disease incidence reported to CDC by year, 1999-2020. Available at:

https://www.cdc.gov/westnile/statsmaps/cumMapsData.html#seven

21. Kottek M, Grieser J, Beck C, Rudolf B, Rubel F. World Map of the Köppen-

Geiger climate classification updated. Meteorol Z. 2006;15: 259–263.

22. United States Census Bureau. Quick Facts Maricopa County, Arizona. Available
at: https://www.census.gov/quickfacts/maricopacountyarizona

23. World Health Organization. Handbook for Integrated Vector Management (World
Health Organization, 2012).

24. Lizzi KM, Qualls WA, Brown SC, Beier JC. Expanding Integrated Vector
Management to promote healthy environments. Trends Parasitol. 2014;30: 394–
400.

25. Wilke ABB, Vasquez C, Medina J, Carvajal A, Petrie W, Beier JC. Community
composition and year-round abundance of vector species of mosquitoes make
Miami-Dade County, Florida a receptive gateway for arbovirus entry to the United
States. Sci Rep. 2019;9: 8732.

26. Wilke ABB, Vasquez C, Carvajal A, Moreno M, Fuller DO, Cardenas G, et al.
Urbanization favors the proliferation of *Aedes aegypti* and *Culex*
quinquefasciatus in urban areas of Miami-Dade County, Florida. Sci Rep.
2021;11: 22989.

27. Ajelli M, Moise IK, Hutchings TCSG, Brown SC, Kumar N, Johnson NF, et al.
Host outdoor exposure variability affects the transmission and spread of Zika
virus: Insights for epidemic control. PLoS Negl Trop Dis. 2017;11: e0005851.

28. Zohdy S, Schwartz TS, Oaks JR. The coevolution effect as a driver of spillover.
Trends Parasitol. 2019;35: 399–408.

29. Centers for Disease Control and Prevention. Arbovirus Catalog. Available at:
https://wwwn.cdc.gov/Arbocat/Default.aspx

30.   Reisen W. The western encephalitis mosquito, *Culex tarsalis*. Wing Beats.
391   1993;4: 16

392   31.   Wilke ABB, Vasquez C, Carvajal A, Medina J, Chase C, Cardenas G, et al.
393   Proliferation of *Aedes aegypti* in urban environments mediated by the availability
394   of key aquatic habitats. Sci Rep. 2020;10: 12925.

395   32.   Paploski IAD, Rodrigues MS, Mugabe VA, Kikuti M, Tavares AS, Reis MG, et al.
396   Storm drains as larval development and adult resting sites for *Aedes aegypti* and
397   *Aedes albopictus* in Salvador, Brazil. Parasit Vectors. 2016;9: 1–8.

398   33.   Souza RL, Mugabe VA, Paploski IAD, Rodrigues MS, Moreira PSDS,
399   Nascimento LCJ, et al. Effect of an intervention in storm drains to prevent *Aedes*
400   *aegypti* reproduction in Salvador, Brazil. Parasit Vectors. 2017;10: 1–6.

401   34.   Wilke ABB, Chase C, Vasquez C, Carvajal A, Medina J, Petrie WD, et al.
402   Urbanization creates diverse aquatic habitats for immature mosquitoes in urban
403   areas. Sci Rep. 2019;9: 15335.

404   35.   Mundis SJ, Estep AS, Waits CM, Ryan SJ. Spatial variation in the frequency of
405   knockdown resistance genotypes in Florida *Aedes aegypti* populations. Parasit
406   Vectors. 2020;13: 241.

407   36.   Wilke ABB, Beier JC, Benelli G. Transgenic mosquitoes – Fact or fiction? Trends
408   in Parasitology. 2018;34: 456–465.

409   37.   Mutebi J-P, Wilke ABB, Vasquez C, Cardenas G, Carvajal A, et al. Diel Activity
410   patterns of two distinct populations of *Aedes aegypti* in Miami, FL and
411   Brownsville, TX. Sci Rep. 2022;12: 5315

38. Darsie, R. F. Jr. & Morris, C. D. Keys to the adult females and fourth-instar larvae

of the mosquitoes of Florida (Diptera, Culicidae). 1st ed. Vol. 1. Tech Bull Florida

Mosq Cont Assoc. 2000.

39. Wilke ABB, Vasquez C, Carvajal A, Moreno M, Petrie WD, Beier JC. Evaluation

of the effectiveness of BG-Sentinel and CDC light traps in assessing the

abundance, richness, and community composition of mosquitoes in rural and

natural areas. Parasit Vectors. 2022;15: 51.

419