Spatiotemporal Distribution of Vector Mosquito Species and Areas at Risk for Arbovirus Transmission in Maricopa County, Arizona

André B. B. Wilke[1,*], Dan Damian[2], Maria Litvinova[1], Thomas Byrne[3,4], Agnese Zardini[5], Piero Poletti[5], Stefano Merler[5], John-Paul Mutebi[6], John Townsend[2], Marco Ajelli[1,*]

[1]Laboratory for Computational Epidemiology and Public Health, Department of Epidemiology and Biostatistics, Indiana University School of Public Health, Bloomington, IN, USA; [2]Maricopa County Environmental Services Department Vector Control Division, Phoenix, AZ, USA; [3]Center for Healthcare Organization and Implementation Research, VA Bedford Healthcare System, Bedford, MA, USA. [4]Boston University School of Social Work, Boston, MA, USA. [5]Center for Health Emergencies, Bruno Kessler Foundation, Trento, Italy; [6]Arboviral Diseases Branch (ADB), Division of Vector-Borne Diseases (DVBD), Centers for Disease Control and Prevention (CDC), Fort Collins, CO, USA.

*Corresponding authors: andwilke@iu.edu; majelli@iu.edu.

Abstract

Mosquito-borne diseases are a major global public health concern and mosquito surveillance systems are essential for the implementation of effective mosquito control strategies. The objective of our study is to determine the spatiotemporal distribution of vector mosquito species in Maricopa County, AZ from 2011 to 2021, and to identify the hotspot areas for West Nile virus (WNV) and St. Louis Encephalitis virus (SLEV) transmission in 2021. The Maricopa County Mosquito Control surveillance system utilizes BG-Sentinel and EVS-CDC traps throughout the entire urban and suburban areas of the county. We estimated specific mosquito species relative abundance per unit area using the Kernel density estimator in ArcGIS 10.2. We calculated the distance between all traps in the surveillance system and created a 4 km buffer radius around each trap to calculate the extent to which each trap deviated from the mean number of *Cx. quinquefasciatus* and *Cx. tarsalis* collected in 2021. Our results show that vector mosquito species are widely distributed and abundant in the urban areas of Maricopa County. A total of 691,170 *Culex quinquefasciatus*, 542,733 *Culex tarsalis*, and 292,305 *Aedes aegypti* were collected from 2011 to 2022. The relative abundance of *Ae. aegypti* was highly seasonal peaking in the third and fourth quarters of the year. *Culex quinquefasciatus*, on the other hand, was abundant throughout the year with several regions consistently yielding high numbers of mosquitoes. *Culex tarsalis* was abundant but it only reached high numbers in well-defined areas bordering natural and rural areas. We also detected high levels of heterogeneity in the risk of WNV and SLEV transmission to humans disregarding traps geographical proximity. The well-defined

43   species-specific spatiotemporal and geographical patterns found in this study can be

44   used to inform vector control operations.

45   **Keywords:** *Aedes aegypti*, *Culex quinquefasciatus*, *Culex tarsalis*, Arbovirus

Introduction

Mosquito-borne diseases are a major global public health concern [1–4]. The incidence of dengue has substantially increased globally in the last decades, from 500,000 reported cases in 2000 to 2.4 million in 2010 and 5.2 million in 2019 [5]. The current estimate is 390 million dengue virus (DENV) infections worldwide every year [6]. Between 2015 and 2018, there were more than 1 million confirmed cases of Zika in the Americas followed by a significant increase in fetus malformation as a direct consequence of Zika virus (ZIKV) infection during pregnancy [3,4].

Albeit on a smaller scale, arboviral infections have also increased in the United States. The ZIKV was introduced multiple times in the United States in 2016 leading to 224 locally transmitted human cases in Florida and Texas [9–11]. Local transmission of the DENV has also been reported in Florida. In 2020, Monroe County experienced a major dengue outbreak with 65 confirmed locally transmitted cases [12–14]. In 2022, 33 locally transmitted cases of dengue were reported in Miami-Dade County [12].

While local outbreaks of dengue and Zika were reported in the United States, the West Nile virus (WNV) has become endemic and the most widespread mosquito-borne arbovirus in North America [15]. WNV has been detected in more than 60 mosquito species in the continental United States (CONUS) [16]; however, selected species in the genus *Culex* such as *Culex quinquefasciatus*, *Culex pipiens*, *Culex tarsalis*, and *Culex nigripalpus* are responsible for driving most of the epizootic and epidemic WNV transmission. Birds are the natural reservoir hosts of WNV as they are able to develop viremia high enough to infect mosquito vectors. Humans, horses, and most mammals

are not able to amplify the virus to viremic levels high enough to infect mosquitoes and

are therefore considered dead-end hosts [16–18].

Since it was first detected in New York in 1999 [19], the WNV has progressively

become endemic in all states in the contiguous United States and has been sporadically

detected in Alaska, Hawaii, and Puerto Rico [20]. From 1999 to 2020 a total of 52,532

human cases were reported in the United States leading to 2,456 deaths [21].

Maricopa County, Arizona, is located in the Sonoran Desert ecoregion spreading

through 23,890 km^2. It has a hot desert climate with long hot summers and short winters

[22]. According to the 2020 census [23], Maricopa County has a population of 4,420,568

people, being the most densely populated county in Arizona as well as in the Sonoran

Desert ecoregion. There are three major cities in Maricopa County, Phoenix (the

Arizona state capital, and the fifth-most populous city in the United States), Mesa, and

Chandler. It has historically been one of the counties most affected by West Nile virus

disease in the United States with an average of 70 neuroinvasive cases reported every

year in the last decade (2010 to 2020). However, 1,427 neuroinvasive cases of West

Nile were reported in this county in 2021, the highest ever reported in a single county in

the United States [12].

Controlling mosquito populations is widely considered the most effective way of

preventing the spread of mosquito-borne diseases [24]. However, the development and

implementation of effective mosquito control strategies in urban areas are complex and

difficult to put into action [25]. They rely on a framework consisting of multiple

successive actions that need to be implemented for a successful intervention. The first

step in the implementation of effective strategies is the development of an extensive

and reliable mosquito surveillance system [26]. Such a system is key to assessing the

mosquito community composition, identifying factors influencing the population

dynamics of mosquito species, estimating the relative abundance of populations of

different vector mosquito species, and providing evidence for ascertaining and

anticipating patterns in mosquito population dynamics and demographics that are

essential for the development of mosquito control strategies. The aim of this study is

twofold: (i) determine the spatiotemporal distribution of mosquito species vectoring

different arboviruses (including WNV, St. Louis Encephalitis virus – SLEV, DENV, ZIKV)

in Maricopa County, Arizona, from 2011 to 2021; and (ii) investigate the hotspot areas

for WNV and SLEV transmission in 2021.

Methods

The Maricopa County Environmental Services Department Vector Control

Division surveillance grid covers the entire urban and peri-urban areas of the county

with approximately one trap per square mile. The surveillance system is comprised of

BG-Sentinel traps baited with BG-Lure (Biogents AG, Regensburg, Germany) (ranging

from 13 traps in 2016 to 8 traps in 2021) and EVS-CDC traps baited with CO_2 (ranging

from 478 traps in 2011 to 825 traps in 2021). Each trap is deployed once a week for 24

hours for 50 weeks a year (Figure 1). All collected mosquitoes are transported to the

Maricopa County Environmental Services Department Vector Control Division

Laboratory and morphologically identified to species using taxonomic keys [27]. BG-

Sentinel traps baited with BG-Lure and EVS-CDC traps baited with CO_2 attract female

mosquitoes seeking hosts for blood-feeding [28]. Even though male mosquitoes were

sometimes collected, male data was not included in the proposed analyses.

Figure 1. Map of Maricopa Mosquito Control Surveillance System. BG-Sentinel

traps are shown in black; EVS-CDC traps are shown in red.

For 2021, virological surveillance was done for all collected *Cx. quinquefasciatus*

and *Cx. tarsalis* mosquitoes in pools of up to 50 female mosquitoes. RNA was extracted

using MagMAX extraction kit (Thermo Fisher Scientific). Samples were prepared in a

PCR hood (Air Clean 600 PCR workstation "Reagent Prep Hood"). Each PCR reaction

included WNV and SLEV positive and negative controls (Table 1).

Table 1. West Nile and St. Louis Virus Real-Time RT-qPCR primers.

Sequence (5' to 3')	Sequence name	Concentration (nM)
CAGACCACGCTACGGCG	WN3'NC - forward	600
CTAGGGCCGCGTGGG	WN3'NC - reverse	600
TCTGCGGAGAGTGCAGTCTGCCGAT	WN3'NC-probe FAM	200
CTGGCTGTCGGAGGGATTCT	SLE2420 - forward	500
TAGGTCAATTGCACATCCCG	SLE2487c - reverse	500
TCTGGCGACCAGCGTGCAAGCCG	SLE2444-CAL FI Org	150

Reactions were carried out using Taqman Master Mix (Thermo Fisher Scientific)

on a QuantStudio 5 at 50°C for 5 minutes, 95°C for 20 seconds, and 40 cycles of: 95°C

for 3 seconds and 60°C for 30 seconds. Samples were considered positive when Ct

values were below 34 and negative when qPCR reaction resulted in no detectable Ct

value after 40 cycles.

We used the geodesic method for densities in square kilometers to calculate the

magnitude-per-unit area (number of inputs per point - i.e., mosquito relative abundance)

using the Kernel density estimator in ArcGIS 10.2 (Esri, Redlands, CA). Figures were

created with layers available at the Maricopa County GIS Mapping Applications

available at: https://www.maricopa.gov/3942/GIS-Mapping-Applications.

We calculated the distance between all traps in the surveillance system (linear

distance in meters) and created a 4 km buffer radius around each target trap. Traps

within each buffer with no data were removed from the analysis. We also removed

outliers in each buffer by excluding observations that lie outside the expected range of

the variability of the mean ($median \pm 1.58 \times \frac{IQR}{\sqrt{n}}$) to calculate mean buffer values. If the

target trap was among the removed outliers, it was identified as a hot (cold) spot if the

number of collected mosquitoes was above (below) the buffer mean (Figure 5). Hot/cold

spot traps were defined to highlight unexpectedly high/low number of collected

mosquitoes relative to the traps in the same area - information useful for the design of

tailored mosquito control interventions. This analysis was conducted in R (version

4.2.2).

To assess the risk of arbovirus transmission posed by the abundance of vector

mosquito species, we used the 2021 data on mosquito vector species presence,

density, and infection rate to calculate the Vector Index (VI: average number of

mosquitoes collected by trap night multiplied by the infection rate) [29]. The infection

rate was estimated as the ratio of the mosquito pools that tested positive for WNV or

SLEV over the total number of tested pools. Then, we used ArcGIS to create a 1-km

buffer around each individual trap and extracted the 2020 total population from the U.S.

Census using the weighted centroid geographic retrieval methodology [30].

Results

Spatiotemporal Distribution of Vector Mosquito Species in Maricopa County, Arizona

The Maricopa County Environmental Services Department Vector Control
Division Surveillance System was comprised of 478 traps in 2011 and new traps were
gradually included over the years as the surveillance system expanded to 833 traps in
2021. From 2016 to 2021 the number of traps remained nearly constant. A total of
691,170 *Cx. quinquefasciatus*, 542,733 *Cx. tarsalis*, and 292,305 *Aedes aegypti* were
collected from 2011 to 2022 by the traps in the Surveillance System (Figures 2 and 3,
Supplementary Table 1).

Figure 2. Number of active traps in the Maricopa County Environmental Services Department Vector Control Division Surveillance System (black line) and the number of mosquitoes collected by year from 2011 to 2021.

Over the study period, *Ae. aegypti* gradually expanded its range and relative

abundance in the urban areas of Maricopa County. In particular, compared to 2020, a

substantial increase in the range of *Ae. aegypti* was observed in 2021 when a total of

88,187 specimens were collected by the traps in the surveillance system (Figure 3A and

Supplementary Figure 1A). In contrast to *Ae. aegypti*, *Cx. quinquefasciatus* was already

commonly found in the urban areas in Maricopa County in 2011. However, since 2011,

Cx. quinquefasciatus became more confined to specific areas of the county and was

particularly abundant in those areas. Since 2016, *Cx. quinquefasciatus* started to

increase in abundance, especially in the southeastern region of Maricopa County

(Figure 3B and Supplementary Figure 1B). Although *Cx. tarsalis* could be detected in

urban areas, this species was mainly confined to specific areas of Maricopa County,

primarily to areas inside or bordering natural or rural areas. By 2021, *Cx. tarsalis* had

gradually expanded its range, but it remained highly abundant in this ecological niche

(Figure 3C, Supplementary Figure 1, Supplementary Tables 2 and 3).

Figure 3. Kernel density-based heat map of the relative abundance of vector

mosquito species collected by the Maricopa County Environmental Services

Department Vector Control Division Surveillance System in 2012, 2017, and 2021. The color gradient represents the sum of mosquitoes collected by each trap in a 12-month period; (A) *Aedes aegypti*; (B) *Culex quinquefasciatus*; and (C) *Culex tarsalis*. The figure was produced using ArcGIS 10.2 (Esri, Redlands, CA), using layers available at the Maricopa County GIS Mapping Applications available at: https://www.maricopa.gov/3942/GIS-Mapping-Applications.

Our results show great seasonal variations in the mean numbers of mosquitoes collected in Maricopa County. Rainfall is one of the possible drivers for the proliferation of mosquitoes in the region, and the peaks in abundance of *Cx. quinquefasciatus* and *Cx. tarsalis* overlap with the peak of the rainy season in the Spring and later in early Fall. Furthermore, rainfall is potentially an important driver for the increase in the abundance of *Ae. aegypti* over the year until the beginning of the dry season, when the abundance of *Ae. aegypti* populations is drastically reduced. Each of the three vector mosquito species, *Ae. aegypti*, *Cx. tarsalis*, and *Cx. quinquefasciatus*, displayed different population dynamic patterns peaking in abundance at different times of the year. This indicates that even though rainfall is an important driver for their proliferation, each species responded differently to the rainfall (Figure 4).

Figure 4. Mean number of *Aedes aegypti*, *Culex quinquefasciatus*, and *Culex*

tarsalis* collected by the Maricopa County Environmental Services Department

Vector Control Division Surveillance System from 2016 to 2021. Solid line

represents the monthly mean average temperature (Celsius) between 2016 and 2021;

Dashed line represents the monthly mean average rainfall (centimeters) between 2016

and 2021.

The mean number of collected *Ae. aegypti* varied greatly within each year. It was

virtually absent during the first quarter of the year yielding a low number of mosquitoes

collected by the traps in the surveillance system. It then gradually increased in

abundance and range during the second quarter, showing the highest abundance in the

southeastern part of the county. This species peaked in abundance during the third

quarter of the year covering most of the urban areas of Maricopa County and reaching

high abundance in several geographical areas. The mean number of *Ae. aegypti*

collected in the fourth quarter of the year was smaller than in the third quarter, albeit

areas with high abundance were still present (Figure 5A).

Culex quinquefasciatus was abundantly found in Maricopa County year-round.

The mean number of *Cx. quinquefasciatus* was the highest during the second and third

quarters of the year with many areas of the county yielding an average higher than 10

mosquitoes per trap per day. One region bordering rural and leisure areas in the

southeastern part of the county yielded more than 10 mosquitoes per trap per day on

average year-round (Figure 5B). *Culex tarsalis* was moderately affected by seasonality.

The mean number of *Cx. tarsalis* remained constant during the second, third, and fourth

quarters of the year with specific areas yielding high numbers of mosquitoes. During the

first quarter, collections of *Cx. tarsalis* were not as abundant as in the other quarters of

the year, but for one region bordering rural and leisure areas in the southeastern part of

the county. Furthermore, *Cx. tarsalis* yielded the highest average number of mosquitoes

collected in a selected group of traps bordering natural and rural areas. Even though the

mean number of *Cx. tarsalis* reached high levels in those specific areas, it was not

abundantly found in urban areas. (Figure 5C).

Figure 5. Kernel density-based heat map of the mean number of *Aedes aegypti*, *Culex quinquefasciatus*, and *Culex tarsalis* collected by the Maricopa County Environmental Services Department Vector Control Division Surveillance System from 2016 to 2021. The color gradient represents the mean number of mosquitoes collected by each trap in a 3-month period. The figure was produced using ArcGIS 10.2 (Esri, Redlands, CA), using layers available at the Maricopa County GIS Mapping Applications available at: https://www.maricopa.gov/3942/GIS-Mapping-Applications.

Hotspot Areas for the transmission of WNV and SLEV in Maricopa County, Arizona

To further our understanding and provide operational input in case of a WNV and SLEV arbovirus outbreak in Maricopa County, we focused on the mosquito and

arbovirus surveillance data from 2021, when Maricopa County reported a record-

breaking number of neuroinvasive WNV cases. To identify highly conducive

environments for the proliferation of *Cx. quinquefasciatus* and *Cx. tarsalis*, we created a

4 km buffer radius around each trap (corresponding to 15 traps on average, range: 1-

34). Then, we calculated the number of traps in each of the buffers, and the total and

mean number of *Cx. quinquefasciatus* and *Cx. tarsalis* collected by the traps in the

cluster. Our results show that from the 820 traps that collected *Cx. quinquefasciatus*

female mosquitoes in 2021, 90 were above their respective buffer average, with 21

influential points (defined as traps yielding an average number of mosquitoes at the

97.5th percentile of each of the clusters). From the 754 traps that collected *Cx. tarsalis*

female mosquitoes in 2021, 76 were above their respective buffer average, with 19

influential points (Figure 6).

Figure 6. Hot spot and cold spot traps that collected a higher (red) or lower (gray) number of female mosquitoes *Cx. quinquefasciatus* and *Cx. tarsalis* relative to the traps in their respective 4 km buffer radius in Maricopa County, Arizona in 2021.

To assess the locations in Maricopa County, Arizona, with the highest WNV and SLEV disease transmission risk, we calculated the VI. We then created a map and overlaid the VI with human population density. Identifying locations with high human population densities and high levels of infected mosquitoes serves as a powerful indicator of the highest-risk locations for WNV and SLEV transmission.

Our results show high levels of heterogeneity in the risk of WNV and SLEV transmission to humans disregarding the geographical proximity of the traps (i.e., the risk of arbovirus transmission has been driven locally at the microgeographic scale and varied greatly between closely located traps). In 2021, WNV was detected in *Cx. quinquefasciatus* mosquito pools at least once in 304 different traps, with VI values ranging from 0.02 to 0.12, and in *Cx. tarsalis* mosquito pools at least once in 222 different traps, with VI values ranging from 0.02 to 0.1. SLEV was detected in *Cx. quinquefasciatus* mosquito pools at least once in 15 different traps. It was also detected at least once in 24 different traps in *Cx. tarsalis* mosquito pools. In both cases, the VI values ranged from 0.02 to 0.04 (Figure 7).

Figure 7. High-risk areas for the transmission of WNV and SLEV in Maricopa County, Arizona in 2021. Relationship between the estimated average number of WNV and SLEV infected *Cx. quinquefasciatus* and *Cx. tarsalis* by trap night (Vector Index) in Maricopa County, Arizona, in 2021, and the 2020 total population from the U.S. Census in a 1-km buffer around each individual trap. White dots represent areas with high Vector Index and high human population density, blue dots represent areas with high Vector Index and low human population density, yellow dots represent areas with low Vector Index and high human population density, and dark red dots represent areas with low Vector Index and low human population. The overall heterogeneity of the

Vector Index for WNV and SLEV circulating in *Cx. quinquefasciatus* and *Cx. tarsalis*

populations are represented using a heatmap.

Discussion

Anthropogenic changes in the environment create favorable conditions for the

development of vector mosquito species that can exploit the resources available in

urban areas [31]. The decreased biodiversity in those urban areas increases the contact

rate between mosquito vector species and human hosts, increasing the risk of arbovirus

transmission to humans [14,32,33]. Our results show that vector mosquito species are

widely distributed and abundantly found in the urban areas of Maricopa County,

Arizona.

Aedes aegypti gradually become more abundant and expanded its range since

only one specimen was collected in Maricopa County in 2011. In 2012, 1,189 *Ae.*

aegypti were collected; by 2014 that number increased to 27,208, and by 2021 a total of

88,187 specimens were collected indicating a substantial increase in its relative

abundance over the years. A similar phenomenon could be observed for *Cx. tarsalis*

and *Cx. quinquefasciatus*, and although they were already present in great numbers

since 2011, with more than 20,000 specimens from each species collected that year,

reaching 89,401 collected specimens *of Cx. tarsalis* and 94,774 of *Cx. quinquefasciatus*

in 2021. We should stress that mosquitoes were collected using BG-Sentinel and CDC

traps, and new traps were gradually added during the period of this study increasing the

sampling effort over time. Even though our results show an increase in the relative

abundance of mosquitoes, the different sampling effort over the years remains a study

limitation.

The high abundance of vector mosquitoes in Maricopa County represents a

potentially dangerous scenario. *Aedes aegypti* is the primary vector of dengue,

chikungunya, yellow fever, and Zika viruses, and *Cx. quinquefasciatus* and *Cx. tarsalis*

are vectors of the WNV, Saint Louis Encephalitis, Eastern Equine encephalitis, and

Japanese Encephalitis viruses [34]. Moreover, differently from dengue and other *Aedes*-

borne diseases where humans do not represent dead-end hosts, for WNV and SLEV

the risk of human cases directly depends on the number of individuals exposed to bites

from infected mosquito vectors. Thus, the high concentration of human population in

urban areas of Maricopa County living in close proximity to infected mosquito vectors

has the potential to lead to major disease burden, as it was the case in 2021, when

1,427 neuroinvasive WNV cases were reported in the county [12].

The relative abundance of *Ae. aegypti* was highly seasonal with fewer

mosquitoes collected in the first 3 months of the year followed by a moderate increase

in numbers in the second quarter indicating that the environmental conditions were not

conducive to the proliferation of *Ae. aegypti* during the first quarter of the year.

However, as the year progresses, a substantial increase in the abundance of *Ae.*

aegypti was observed in the third and fourth quarters. The observed seasonal variation

in the abundance of *Ae. aegypti* can potentially be an important driver for arbovirus

transmission in Maricopa County as its abundance and range increase as the year

progresses, thus, potentially increasing the contact rate between *Ae. aegypti* and

human hosts. *Culex quinquefasciatus*, on the other hand, was abundantly found during

all seasons of the year. It was not as abundant in the first quarter as it was during the

remaining of the year, but it could be found throughout the urban areas during that time

of the year. Several regions consistently yielded high numbers of *Cx. quinquefasciatus*

mosquitoes indicating that those areas were hot spots and were able to support high

numbers of *Cx. quinquefasciatus* year-round. *Culex tarsalis* was also abundantly found

in Maricopa County but in contrast to *Ae. aegypti* and *Cx. quinquefasciatus*, it only

reached high numbers in well-defined areas bordering natural and rural areas.

However, this is not surprising because *Cx. tarsalis* larval habitats include irrigation

agricultural runoff, wetlands, sewage effluent, oil field run-off, and lake beds [35], and all

these habitat types are rarely associated with urban areas. Human cases of WNV

disease have been consistently reported in Arizona, being one of the most affected

states in the United States. The high abundance of *Culex* vector species, especially the

year-round presence of *Cx. quinquefasciatus*, has major implications for the likelihood

of WNV transmission in Maricopa County [12]. In sum, our results showed well-defined

spatiotemporal patterns for each of the species included in this study. However, further

analyses are needed to investigate the main drivers for the variations in the population

dynamics of each mosquito vector species that are essential for the development and

implementation of effective source reduction and other mosquito control strategies [36].

The cluster analysis results aimed to detect influential point traps that collected a

significantly higher number of either *Cx. quinquefasciatus* or *Cx. tarsalis* as compared to

other traps in their proximity indicate that resources available at the microgeographic

scale were responsible for their proliferation. Further analysis is needed to understand

which local geographic features (e.g., urban parks, fishing ponds, sports complexes, as

appears to be the case in our data) may have contributed to the observed patterns.

Identifying such factors is instrumental to guide vector control operations.

The identification of high-risk areas for the transmission of WNV and SLEV is

vital for the development of vector control strategies aiming at reducing the abundance

of *Cx. tarsalis* and *Cx. quinquefasciatus* and thus the risk of infection. Our results show

the overlap between areas with high human population densities and high levels of

infected mosquitoes. These areas are priority targets for the development and

implementation of vector mosquito control strategies. Furthermore, there is a need to

assess and validate the effectiveness of different control methods and strategies aimed

to reduce mosquito populations in urban areas. These activities could focus on how the

proliferation of vector mosquito species can be locally reduced by source reduction,

community engagement, and by modifying specific features of the urban built

environment [36–39]. Developing and implementing effective mosquito control

strategies is a difficult task with many factors that can potentially hinder the

effectiveness of mosquito control interventions, such as levels of insecticide resistance,

diel activity of target mosquito species, and outdated and ineffective mosquito control

tools [40–42].

Our analysis of high-risk areas for WNV and SLEV infection considers only the

density of the human population, disregarding other characteristics such as the

population's, age structure and underlying health status, as well as other social

determinants including individuals' socioeconomic status, and characteristics of the

neighborhood in which they reside [43]. Different segments of the population could

indeed be exposed to a heterogeneous risk of infection and disease depending on their

age, occupation (e.g., construction workers vs. office workers), adequacy of their

housing arrangement (e.g., presence of mosquito window screenings and A.C. units),

and the built environment in their neighborhood (e.g. presence of abandoned

lots/buildings that may provide more breeding areas). A segment of the population that

is at particular risk of arbovirus infection is represented by people experiencing

unsheltered homelessness as they are exposed nearly 24/7 to mosquito vector bites. It

has been shown that the prevalence of WNV in Houston, Texas, is higher in people that

have been homeless for more than one year when compared with the general

population [44]. In other metropolitan areas in the United States, including Maricopa

County, Arizona, it is so far unknown to what extent vulnerable and underserved

populations are being disproportionally exposed to arboviruses. These groups are

already affected by large disparities in terms of mortality and a broad array of other

health outcomes [45], and arriving at a better understanding of the nature and extent of

their disproportionate exposure to arboviral infections is crucial for designing policy

interventions that can mitigate such a preventable health disparity and advance health

equity.

Effective mosquito control operations rely on multiple factors, with the most

important being consistent and reliable mosquito surveillance systems, community

engagement, and targeted source reduction. Now, more than ever, effective mosquito

control strategies are needed to deal with a flux of importation of travelers carrying

arboviruses to prevent local outbreaks, and to curtail them when they start. Effective

mosquito surveillance systems are key for guiding and implementing mosquito control

strategies and should be considered a priority for local, state, and federal public health professionals.

Supplementary Table 1. Number of traps and mosquitoes collected by the Maricopa County Environmental Services Department Vector Control Division Surveillance System from 2011 to 2021.

Supplementary Table 2. Number of mosquitoes (*Aedes aegypti*, *Culex quinquefasciatus*, and *Culex tarsalis*) collected per trap.

Supplementary Table 3. Number of mosquitoes (*Aedes aegypti*, *Culex quinquefasciatus*, and *Culex tarsalis*) collected by the same set of 478 traps in use since 2016.

Supplementary Figure 1. Kernel density-based heat map of the relative abundance of vector mosquito species collected by the Maricopa County Environmental Services Department Vector Control Division Surveillance System from 2012 to 2021. The color gradient represents the sum of mosquitoes collected by each trap in a 12-month period; (A) *Aedes aegypti*; (B) *Culex quinquefasciatus*; and (C) *Culex tarsalis*. The figure was produced using ArcGIS 10.2 (Esri, Redlands, CA), using layers available at the Maricopa County GIS Mapping Applications available at: https://www.maricopa.gov/3942/GIS-Mapping-Applications.

Acknowledgments

Disclaimer: The information provided and views expressed in the publication do not necessarily reflect the official views of the Centers for Disease Control and Prevention (CDC).

References

1. Messina JP, Brady OJ, Pigott DM, Brownstein JS, Hoen AG, Hay SI. A global compendium of human dengue virus occurrence. Sci Data. 2014;1: 140004.

2. Brady OJ, Hay SI. The global expansion of dengue: How *Aedes aegypti* mosquitoes enabled the first pandemic arbovirus. Annu Rev Entomol. 2020;65: 191–208.

3. Keyel AC, Raghavendra A, Ciota AT, Elison Timm O. West Nile virus is predicted to be more geographically widespread in New York State and Connecticut under future climate change. Glob Chang Biol. 2021;27: 5430–5445.

4. Dorigatti I, Morrison S, Donnelly CA, Garske T, Bowden S, Grills A. Risk of yellow fever virus importation into the United States from Brazil, outbreak years 2016–2017 and 2017–2018. Sci Rep. 2019;9: 1–11.

5. World Health Organization. Dengue and severe dengue. Available at: https://www.who.int/news-room/fact-sheets/detail/dengue-and-severe-dengue

472   6.   Bhatt S, Gething PW, Brady OJ, Messina JP, Farlow AW, Moyes CL, et al. The

473   global distribution and burden of dengue. Nature. 2013;496: 504–7.

474   7.   World Health Organization. Zika cases and congenital syndrome associated with

475   Zika virus reported by countries and territories in the Americas (Cumulative

476   Cases), 2015–2017. Available at: https://www.paho.org/hq/index.php?option =

477   com_content&view = article&id = 12390:zika-cumulative-cases&Itemid =

478   42090&lang = en.

479   8.   Shapiro-Mendoza CK, Rice ME, Galang RR, Fulton AC, VanMaldeghem K,

480   Prado MV, et al. Pregnancy outcomes after maternal Zika virus infection during

481   pregnancy — U.S. Territories, January 1, 2016–April 25, 2017. Morb Mortal Wkly

482   Rep. 2017;66: 615–621.

483   9.   Grubaugh ND, Ladner JT, Kraemer MUG, Dudas G, Tan AL, Gangavarapu K, et

484   al. Genomic epidemiology reveals multiple introductions of Zika virus into the

485   United States. Nature. 2017;546: 401–405.

486   10.   Likos A, Griffin I, Bingham AM, Stanek D, Fischer M, White S, et al. Local

487   mosquito-borne transmission of Zika virus — Miami-Dade and Broward Counties,

488   Florida, June–August 2016. Morb Mortal Wkly Rep. 2016;65: 1032–1038.

489   11.   Centers for Disease Control and Prevention. Zika cases in the United States.

490   Available at: https://www.cdc.gov/zika/reporting/index.html.

491   12.   Centers for Disease Control and Prevention. Imported human disease cases

492   reported to CDC by county of residence. Available at:

493   https://wwwn.cdc.gov/arbonet/Maps/ADB_Diseases_Map/index.html

494   13.   Florida Department of Health. Available at: http:// www. flori dahea lth. gov/ disea

495   ses- and- conditions/ mosqu ito- borne- disea ses/_docum ents/ alert- dade- wnv-

496   human- 10- 19- 20. pdf.

497   14.   Coatsworth H, Lippi CA, Vasquez C, Ayers JB, Stephenson CJ, Waits C, et al. A

498   molecular surveillance-guided vector control response to concurrent dengue and

499   West Nile virus outbreaks in a COVID-19 hotspot of Florida. The Lancet Regional

500   Health – Americas. 2022;11: 100231

501   15.   Ronca SE, Ruff JC, Murray KO. A 20-year historical review of West Nile virus

502   since its initial emergence in North America: Has West Nile virus become a

503   neglected tropical disease? PLoS Negl Trop Dis. 2021;15: e0009190

504   16.   Centers for Disease Control and Prevention. Mosquito species in which West

505   Nile virus has been detected. Available at:

506   https://www.cdc.gov/westnile/resources/pdfs/Mosquito%20Species%201999-

507   2012.pdf

508   17.   Centers for Disease Control and Prevention. West Nile Virus in the United

509   States: Guidelines for surveillance, prevention, and control. Available:

510   https://www.cdc.gov/westnile/resources/pdfs/wnvGuidelines.pdf

511   18.   Jeffrey Root J. West Nile virus associations in wild mammals: A synthesis.

512   Archives of Virology. Springer-Verlag Wien; 2013. pp. 735–752.

513   19.   Hile JA, Mangiafico JA, Clippinger TL, Raphael BL, Smith JF, Danner DK, et al.

514   An outbreak of West Nile virus in a New York City captive wildlife population. Am

515   J Trop Med Hyg. 2002;67: 67–75.

20. Centers for Disease Control and Prevention. West Nile virus disease cases

reported to CDC by state of residence, 1999-2020. Available at:

https://www.cdc.gov/westnile/statsmaps/cumMapsData.html.

21. Centers for Disease Control and Prevention. West Nile virus neuroinvasive

disease incidence reported to CDC by year, 1999-2020. Available at:

https://www.cdc.gov/westnile/statsmaps/cumMapsData.html#seven

22. Kottek M, Grieser J, Beck C, Rudolf B, Rubel F. World Map of the Köppen-

Geiger climate classification updated. Meteorol Z. 2006;15: 259–263.

23. United States Census Bureau. Quick Facts Maricopa County, Arizona. Available

at: https://www.census.gov/quickfacts/maricopacountyarizona

24. World Health Organization. Handbook for Integrated Vector Management (World

Health Organization, 2012).

25. Lizzi KM, Qualls WA, Brown SC, Beier JC. Expanding Integrated Vector

Management to promote healthy environments. Trends Parasitol. 2014;30: 394–

400.

26. Wilke ABB, Vasquez C, Medina J, Carvajal A, Petrie W, Beier JC. Community

composition and year-round abundance of vector species of mosquitoes make

Miami-Dade County, Florida a receptive gateway for arbovirus entry to the United

States. Sci Rep. 2019;9: 8732.

27. Darsie, R. F. Jr. & Morris, C. D. Keys to the adult females and fourth-instar larvae

of the mosquitoes of Florida (Diptera, Culicidae). 1st ed. Vol. 1. Tech Bull Florida

Mosq Cont Assoc. 2000.

28.   Wilke ABB, Vasquez C, Carvajal A, Moreno M, Petrie WD, Beier JC. Evaluation

of the effectiveness of BG-Sentinel and CDC light traps in assessing the

abundance, richness, and community composition of mosquitoes in rural and

natural areas. Parasit Vectors. 2022;15: 51.

29.   Centers for Disease and Control. West Nile Virus in the United States: Guidelines

for   Surveillance,   Prevention,   and   Control.   Available   at:

https://www.cdc.gov/westnile/resources/pdfs/wnvGuidelines.pdf.

30.   ArcGIS.   Data   apportionment.   Available   at:

https://developers.arcgis.com/rest/geoenrichment/api-reference/data-

apportionment.htm#ESRI_SECTION1_206F094B25E540B3A01DB46E499EB08

C

31.   Wilke ABB, Vasquez C, Carvajal A, Moreno M, Fuller DO, Cardenas G, et al.

Urbanization   favors   the   proliferation   of   *Aedes   aegypti*   and   *Culex*

quinquefasciatus in urban areas of Miami-Dade County, Florida. Sci Rep.

2021;11: 22989.

32.   Ajelli M, Moise IK, Hutchings TCSG, Brown SC, Kumar N, Johnson NF, et al.

Host outdoor exposure variability affects the transmission and spread of Zika

virus: Insights for epidemic control. PLoS Negl Trop Dis. 2017;11: e0005851.

33.   Zohdy S, Schwartz TS, Oaks JR. The coevolution effect as a driver of spillover.

Trends Parasitol. 2019;35: 399–408.

34.   Centers   for   Disease   and   Control.   Arbovirus   Catalog.   Available   at:

https://wwwn.cdc.gov/Arbocat/Default.aspx

35. Reisen W. The western encephalitis mosquito, *Culex tarsalis*. Wing Beats.

1993;4.

36. Wilke ABB, Vasquez C, Carvajal A, Medina J, Chase C, Cardenas G, et al.

Proliferation of *Aedes aegypti* in urban environments mediated by the availability

of key aquatic habitats. Sci Rep. 2020;10: 12925.

37. Paploski IAD, Rodrigues MS, Mugabe VA, Kikuti M, Tavares AS, Reis MG, et al.

Storm drains as larval development and adult resting sites for *Aedes aegypti* and

Aedes albopictus in Salvador, Brazil. Parasit Vectors. 2016;9: 1–8.

38. Souza RL, Mugabe VA, Paploski IAD, Rodrigues MS, Moreira PSDS,

Nascimento LCJ, et al. Effect of an intervention in storm drains to prevent *Aedes*

aegypti reproduction in Salvador, Brazil. Parasit Vectors. 2017;10: 1–6.

39. Wilke ABB, Chase C, Vasquez C, Carvajal A, Medina J, Petrie WD, et al.

Urbanization creates diverse aquatic habitats for immature mosquitoes in urban

areas. Sci Rep. 2019;9: 15335.

40. Mundis SJ, Estep AS, Waits CM, Ryan SJ. Spatial variation in the frequency of

knockdown resistance genotypes in Florida *Aedes aegypti* populations. Parasit

Vectors. 2020;13: 241.

41. Wilke ABB, Beier JC, Benelli G. Transgenic mosquitoes – Fact or fiction? Trends

Parasitol. 2018;34: 456–465.

42. Mutebi J-P, Wilke ABB, Vasquez C, Cardenas G, Carvajal A, et al. Diel Activity

patterns of two distinct populations of *Aedes aegypti* in Miami, FL and

Brownsville, TX. Sci Rep. 2022;12: 5315

43.   Braveman P, Gottlieb L. The social determinants of health: It's time to consider

the causes of the causes. Public Health Reports. 2014;129: 19–31.

44.   Meyer TE, Bull LM, Holmes KC, Pascua RF, Travassos da Rosa A, Gutierrez

CR, et al. West Nile virus infection among the homeless, Houston, Texas. Emerg

Infect Dis. 2007;13: 1500–1503.

45.   Fazel S, Geddes JR, Kushel M. The health of homeless people in high-income

countries: descriptive epidemiology, health consequences, and clinical and policy

recommendations. The Lancet. 2014;384: 1529–1540.