The spectral underpinnings of pathogen spread on animal networks

Authors: Nicholas M. Fountain-Jones[1*], Mathew Silk[2,3], Raima Carol Appaw[1], Rodrigo Hamede[1], Julie Rushmore[4], Kimberly VanderWaal[5], Meggan E Craft[6], Scott Carver[1] & Michael Charleston[1]

Affiliations:

[1] School of Natural Sciences, University of Tasmania, Hobart Australia 7001.

[2] CEFE, University of Montpellier, CNRS, EPHE, IRD, University of Paul Valéry Montpellier 3, Montpellier, France

[3] Centre for Ecology and Conservation, University of Exeter, Penryn Campus, Penryn, UK

[4] Odum School of Ecology, University of Georgia, Athens, GA USA.

[5] Department of Veterinary Population Medicine, University of Minnesota, St Paul, MN, USA

[6] Department of Ecology, Evolution, and Behavior, University of Minnesota, St Paul, MN, USA

*Nick.FountainJones@utas.edu.au

Number of words in the main text/abstract:

Number of figures/tables/boxes: 5 figures and 1 box

Keywords: Disease simulation models, wildlife, graph theory, machine learning.

Abstract

Predicting what factors promote or protect populations from infectious disease is a

fundamental epidemiological challenge. Social networks, where nodes represent hosts and

edges represent direct or indirect contacts between them, are important in quantifying these

aspects of infectious disease dynamics. However, how network structure and epidemic

parameters interact in empirical networks to promote or protect animal populations from

infectious disease remains a challenge. Here we draw on advances in spectral graph theory

and machine learning to build predictive models of pathogen spread on a large collection of

empirical networks from across the animal kingdom. We show that the spectral features of an

animal network are powerful predictors of pathogen spread for a variety of hosts and

pathogens and can be a valuable proxy for the vulnerability of animal networks to pathogen

spread. We validate our findings using interpretable machine learning techniques and provide

a flexible web application for animal health practitioners to assess the vulnerability of a

particular network to pathogen spread.

Introduction

Capturing patterns of direct or indirect contacts between hosts is crucial to model pathogen spread in populations (Newman 2002; Craft 2015; Pastor-Satorras *et al.* 2015; Sah *et al.* 2018, 2021). Increasingly, contact or social network approaches, where hosts are nodes and edges reflect interactions between hosts, play a central role in epidemiology and disease ecology (e.g., Meyers *et al.* 2005; Bansal *et al.* 2007; Eames *et al.* 2015; White *et al.* 2017). When there is heterogeneity in contacts between individuals, network approaches can provide more nuanced and reliable estimates of pathogen spread, including in wildlife populations (e.g., Meyers *et al.* 2006; Bansal *et al.* 2010; Craft *et al.* 2011). Formulating general rules for how easy-to-calculate network structure properties may promote or protect populations from pathogen spread can reveal important insights into how host behaviour can mediate epidemic outcomes (Sah *et al.* 2017), and provide practitioners with a proxy for how vulnerable a population is to disease without extensive simulations (Silk *et al.* 2017; Sah *et al.* 2018). Further, structural properties of networks can be incorporated into traditional susceptible–infected–recovered (SIR) models to account for contact heterogeneity when predicting pathogen dynamics across populations (e.g., Meyers *et al.* 2005; Bansal *et al.* 2007; Pastor-Satorras *et al.* 2015). For example, approaches such as degree-based mean field approximation (DBMF), can estimate outbreak size using the degree distribution of uncorrelated networks (i.e., the degree of one node is uncorrelated with neighbouring nodes, (Newman 2002; Pastor-Satorras *et al.* 2015)).

However, it remains unclear whether one structural characteristic or a combination of network characteristics can reliably predict pathogen dynamics across empirical networks in different biological systems (Ames *et al.* 2011; Sah *et al.* 2018). For example, species that are more social tend to have more clustered or "modular" networks, and this modularity has been

found to increase (Lentz *et al.* 2012), reduce (Salathé & Jones 2010), or have little effect (Sah

et al. 2018) on outbreak size across different systems. The average number of contacts

between hosts can be identical across networks and yet still result in substantially different

outbreak patterns (Ames *et al.* 2011). Even the apparent size (number of nodes) of the

network, often constrained by limitations of sampling, can impact estimates of pathogen

spread, particularly in wildlife populations (McCabe & Nunn 2018). As network

characteristics such as network size and modularity are often correlated (Newman 2006; Silk

et al. 2017) and can have complex impacts on spread (Sah *et al.* 2017; McCabe & Nunn

2018; Porter 2020), determining those characteristics that affect outbreaks remains a

fundamental question in infectious disease biology (Sah *et al.* 2018).

Searching for general relationships between network structure and pathogen spread in animal

populations is further challenged, as the relationship is also affected by host-pathogen traits,

such as infectiousness and recovery rate. For example, modularity appears to make no

difference to pathogen prevalence for highly infectious pathogens (Sah *et al.* 2017).

Pathogens with long recovery rates can increase outbreak size across networks (Shu *et al.*

2016). Given that we rarely have reliable estimates of pathogen traits in wild populations

(e.g., for different probabilities of infection per contact, or recovery rates), any predictive

model of the relationship between spread and network structure would ideally be

generalizable across pathogens.

Advances in spectral graph theory have the potential to resolve this challenge, offering an

additional set of measures based on the **spectrum** of a network rather than average node or

edge level attributes (see Text Box 1 for further definitions of terminology in bold). A graph

spectrum is the set of **eigenvalues** (often denoted with a Greek lambda λ) of a matrix

representation of a network. Theoretical studies have shown relationships between particular

eigenvalues and connectivity across networks are independent of pathogen propagation

models (Prakash *et al.* 2010). For example, networks with a high algebraic connectivity, also

known as the **Fiedler value** (the second smallest eigenvalue of the network's **Laplacian**

matrix) are "more connected" than those with low values. It has been found that, in

ecological networks for example, if the Fiedler value is sufficiently large, removing edges

will have little effect on overall network connectivity (Kumar *et al.* 2019), but whether this

lack of effect is mirrored by pathogen dynamics is not yet clear. Another quantity of interest

is the **spectral radius** – the largest absolute value of the eigenvalues of its **adjacency matrix**.

The link between the spectral radius and epidemiological dynamics is better understood, with

theoretical work showing that this value closely mirrors both epidemic behaviour and

network connectivity (Prakash *et al.* 2010) and has been used to understand vulnerability of

cattle networks to disease (Darbon *et al.* 2018). For example, networks with the same number

of edges and nodes but higher spectral radius (λ_1) are more vulnerable to outbreaks than

networks with low spectral radius ($\lambda_1 \rightarrow 1$). We hypothesize that spectral measures such as

these have great potential to improve our ability to predict dynamics of pathogen spread on

networks, where previous methods such as modularity have proved inadequate (Sah *et al.*

2017).

We assess the predictive capability of spectral values compared to other structural attributes

such as **modularity** (Newman's Q; (Newman 2006)) using advances in machine learning to

construct non-linear models of simulated pathogen spread across a large collection of

empirical animal networks, including those from the Animal Social Network Repository

(ASNR) (Sah *et al.* 2019). The ASNR is a large repository of empirical social networks that

provides novel opportunities to test the utility of spectral values in predicting spread across a

wide variety of, mostly animal, taxa across a diversity of social systems -- from eusocial ants

(Arthropoda: Formicidae) to more solitary species such as the desert tortoise (*Gopherus*

agassizii). We combined networks from this resource with other published networks,

including badgers (*Meles meles*) (Weber *et al.* 2013), giraffes (*Giraffa camelopardalis*)

(VanderWaal *et al.* 2014), and chimpanzees (*Pan troglodytes*) (Rushmore *et al.* 2013) to

generate a dataset of over 600 unweighted networks from 51 species. We treated each

network as static (i.e., edges are fixed for the duration of the simulated outbreak). While

dynamic networks where contact patterns fluctuate over time can provide more realistic

estimates of pathogen spread (Bansal *et al.* 2010), they are rare, particularly for wild

populations. Therefore, we were restricted to analysing static networks, though further study

of dynamic networks would be valuable. We then simulated pathogen spread using a variety

of SIR parameters and harnessed recent advances in multivariate interpretable machine

learning models (*MrIML;* (Fountain-Jones *et al.* 2021)) to construct predictive models. As

many species were represented by multiple networks, often over different populations and or

timepoints and constructed in different ways (e.g., some edges reflected spatial proximity

rather than direct contact), we included species and network construction variables in our

models to account for these correlations in addition to exploring the diversity of network

structures across the animal kingdom.

We demonstrate that using spectral measures of network structure alone can provide a useful

proxy for network vulnerability. Our interpretable machine learning models identify putative

threshold values for the vulnerability of a network to pathogen spread. Further, we provide a

user-friendly application that utilizes our models to provide practitioners with predictions, for

example, of the prevalence of a pathogen across a variety of spread scenarios using a user-

supplied network without the need for lengthy simulation (see

https://spreadpredictr.shinyapps.io/spreadpredictr/).

Methods

Networks

We downloaded all animal social networks from the ASNR on 12[th] January 2022 (Sah *et al.*

2019) and combined these with other comparable published animal networks (Rushmore *et*

al. 2013; Weber *et al.* 2013; VanderWaal *et al.* 2014). These networks included those

reflecting direct and indirect (such as burrow sharing) contacts between hosts (hereafter

'contact networks'). Farmed domestic animals were not included in our analyses. We

binarized each network, extracted the largest connected component, and excluded networks

with fewer than 10 individuals. This left us with 603 networks from 43 species.

From each network we calculated a variety of network structure variables using the R

package *igraph* (Csárdi & Nepusz 2006) (see Table S1 for a summary of the variables

measured). As these networks were constructed using different techniques, we also extracted

metadata from the ASNR or the publication associated with the network (Table S2). These

variables were also added to the models. We used Principal Components Analysis (PCA)

biplots to examine the drivers of variation in network structure and visualise how networks

clustered by taxonomic class. We removed eight networks with missing metadata and

screened for correlations between variables. We used a pairwise correlation threshold of 0.7

to remove the variable from each such pair with the highest overall correlation. We used this

relatively high threshold as many of the machine learning algorithms are less sensitive to

variable collinearity (Fountain-Jones *et al.* 2019)

Simulations

To simulate the spread of infection on each network we used our R package "EpicR"

(Epidemics by computers in R; available on GitHub at https://github.com/mcharleston/epicr).

The simulations use a standard discretisation of the SIR model, in which time proceeds in

"ticks," for example representing days. Initially one individual was chosen at uniform random

to be infected (I) and all others were susceptible (S). At each time step, one of two changes of

state can happen to each individual (represented by a node), depending on its current state.

An 'S' individual will become infected (I) with a probability $(1 - (1 - \beta)^k)$, where k is the

number of currently infected neighbours it has, or otherwise stay as S; an 'I' individual will

recover (R) with probability γ or remain as I. Recovered (R) individuals stay as R.

In classic deterministic SIR models as a set of differential equations, β and γ are

instantaneous rates; here, they are probabilities per time step, so at a coarse level, they are

comparable.

On each network, we performed 1000 simulations using different combinations of

transmission ($\beta = 0.01, 0.025, 0.05, 0.1, 0.2$) and recovery probabilities ($\gamma = 0.02, 0.04, 0.2$,

0.4). We chose these SIR parameter values to broadly reflect a range of scenarios from high

to low transmissibility and slow to fast recovery (Leung 2021) and ensure large outbreaks

(>10% of individuals infected, (Sah *et al.* 2017)). One randomly chosen individual was

infected at the beginning of the simulation and we ran the simulation for 100 time steps to

ensure that the epidemic ended and there were no remaining infected nodes. For each

simulation we recorded two complementary epidemic measures to capture disease burden and

speed of spread: a) the maximum prevalence reached, or the maximum proportion of

individuals infected in the network after 100 time steps (hereafter 'proportion infected') and

b) time to outbreak peak (i.e., which time step had the maximum number of infections,

hereafter 'time to peak'). The epidemic measures estimated in the simulations were used as

the response variables in the machine learning models.

Machine learning pipeline

We used a recently developed multi-response interpretable machine learning approach

(*MrIML*) (Fountain-Jones *et al.* 2021) to predict outbreak characteristics using network

structure variables. Our *MrIML* approach had the advantage of allowing us to rapidly

construct and compare models across a variety of machine-learning algorithms for each of

our response variables as well as assess generalized predictive surfaces across epidemic

parameters.

To test the robustness of our results, we compared the performance of four different

underlying supervised regression algorithms in our *MrIML* models. We compared linear

models (LMs), support vector machines (SVMs), random forests (RF) and gradient boosted

models (GBMs) as they operate in markedly different ways that can affect predictive

performance (Fountain-Jones *et al.* 2019; Machado *et al.* 2019). For the best performing

model, we also tested algorithm performance using just the spectral radius and Fiedler value

for each network. Categorical predictors such as 'species' were hot-encoded for some models

as needed (i.e., representing the variable as a binary vector, see Table S4). As proportion

infected and time to peak are continuous response variables, we compared the performance

of each algorithm using the average R^2 and root mean squared error (RMSE) across all

responses (hereafter, the 'global model'). As we included models that were not fitted using

sums of squares, our R^2 estimate depended on the squared correlation between the observed

and predicted values (Kvålseth 1985). As ants (Insecta: Formicinae) were over-represented,

we compared model performance and interpretation with and without these networks using

10-fold cross validation, to prevent overfitting. We tuned hyperparameters for each model

(where appropriate) using 100 different hyper-parameter combinations (a 10×10 grid search)

and selected the combination with the lowest RMSE. The underlying algorithm with the

highest predictive performance was interrogated further.

We interpreted this final model using a variety of model-agnostic techniques within the

MrIML framework. We assessed overall and model-specific variable importance using a

variance-based method (Greenwell *et al.* 2018). We quantified how each variable alters

epidemic outcomes using accumulated local effects (ALEs) (Apley & Zhu 2016). In brief,

ALEs isolate the effect of each network characteristic on epidemic outcomes using a sliding

window approach, calculating the average change in prediction across the values' range

(while holding all other variables constant) (Molnar 2018). ALEs are less sensitive to

correlations and straightforward to interpret as points on the ALE curve are the difference in

the prediction of each observation from the mean prediction (Apley & Zhu 2016; Molnar

2018; Fountain-Jones *et al.* 2021).

To further examine the predictive performance of our black-box models (SVM, RF and

GBM) we calculated a global surrogate decision tree (hereafter 'global surrogate') to

approximate the predictions of our more complex trained models. Global surrogates are

generated by training a simpler decision tree to the *predictions* (instead of observations) of

the more complex 'black box' models using the network structure data. How well the

surrogate model performed compared to the complex model is then estimated using R^2. See

Molnar (2018) for details.

We gained more insight into model behaviour and how network structure impacted epidemic

outcomes on individual networks, including by calculating Shapley values (Štrumbelj &

Kononenko 2014). Shapley values use a game theoretic approach to play off variables in the

model with each other based on their contribution to the prediction (Shapley 1953).

Calculating Shapley values involves solving a system of linear equations to assign each feature

a unique weight based on its contribution to the predicted output and its interaction with other

features (Štrumbelj & Kononenko 2014). For example, negative Shapley values indicate that

the observed value 'contributed to the prediction' by reducing the proportion infected or time

to peak in an outbreak for a particular network. See Molnar (2018) for a more detailed

description and Fountain-Jones *et al.* (2019) and Worsley-Tonks *et al.* (2020) for how

Shapley values can be interpreted in epidemiological settings. See

https://github.com/nfj1380/igraphEpi for our complete analytical pipeline.

Lastly, we also compared our machine learning estimates of proportion infected to those

derived from the DBMF SIR approximations for each network. Briefly, the DBMF

approximation calculates the proportion infected using the following equation:

$$p_r = \sum_k P(k)\left(1 - e^{-\lambda k \phi}\right)$$

255 Where P(k) is the degree distribution of the network, λ is the epidemic threshold and k is the

256 first moment of the degree distribution. The auxiliary function (Φ) cannot be solved

257 analytically, but can be approximated using the network epidemic transition threshold (λ_c):

$$\lambda_c = \frac{\langle k \rangle}{\langle k^2 - k \rangle}$$

258 where k^2 is the second moment of the degree distribution. See Pastor-Satorras *et al.* (2015)

259 for more details.

260 **Results**
261
Diversity of network structures

We identified substantial variation in network structure across animal taxa. The static

unweighted animal social networks in our database ranged from nearly completely

unconnected (Spectral radius $\lambda_1 \sim 1$, Fiedler value ~0, not included in our predictive models)

to highly connected (Spectral radius $\lambda_1 \sim 160$, Fiedler value ~ 140, Fig. 1). Similarly, the

networks ranged from homogeneous (i.e., not modular, $Q_{rel} = 0$, see Text Box 1) to highly

modular and subdivided ($Q_{rel} > 0.8$). Our principal component analysis (PCA) identified key

axes of structural variation across empirical networks (Fig. 2). The first principal component

(PC1) distinguished networks that had a large diameter and mean path length and were highly

modular (PC1 < 0) from networks with a high mean degree and transitivity (PC1 > 0, Fig. 2,

see Table S3). The second principal component (PC2) separated networks based on network

size (number of nodes), maximum degree and the network duration (i.e., the time period over

which the network data was collected, Fig. 2). The eusocial ant networks (*Camponotus fellah*,

Insecta: Hymenoptera) and mammal networks tended to cluster separately (Fig. 2), with the

other taxonomic classes dispersed between these groups (Fig. 2) or species (see Fig. S1 for

clustering by species). The networks' spectral properties (the Fiedler value and spectral

radius) explained a unique portion of structural variance that did not covary with other

variables (see Table S3 for vector loadings and Fig S2 for all pair-wise correlations). We

found variables such as mean degree and transitivity the most correlated with the other

variables and were excluded from further analysis (Tables S2, Fig S2).

Spectral properties predict pathogen spread across epidemic scenarios

Across all SIR parameter combinations, spectral measures played a dominant role in our

predictive models (Fig. 3a, Fig. S3). We could predict the proportion infected in a network

and time to peak using both spectral measures and species identity alone (Table S4). To aid

interpretation we present the analyses from two recovery probabilities ($\gamma = 0.04, 0.4$ see Fig

S4 for the analysis with a wider variety of recovery probabilities). Network size, relative

modularity and centralization, for example, were less important in predicting proportion

infected across all SIR model parameter combinations tested (Fig. 3a). Nonlinear

relationships were likely important for prediction of proportion infected, as random forests

(RF) had the highest predictive performance overall (Table S4) and substantially

outperformed linear regression in the *MrIML* framework (root mean square error (RMSE)

0.03 vs 0.13). Our RF model just including the spectral radius and Fiedler value had very

similar predictive performance (R^2, RMSE) compared to the models with all variables

included (Table S4). Our models also substantially outperformed DBMF approximations with

a ~ten-fold decrease in RMSE across a range of β and γ values (Table S5).

Variable importance and predictor conditional effects were consistent between the machine

learning algorithms, so we subsequently analysed the best performing RF model. We found a

nonlinear relationship between proportion infected and spectral radius across each SIR

parameter combination, with the average prediction of proportion infected increasing by

~30% across the range of spectral radius values (holding all other variables constant in the

model, Fig. 3b). In contrast we found a more modest effect of the Fiedler value, with the

proportion of infected only increasing on average ~3% across the observed range of values

for all SIR parameters (β & γ, Fig 3c). When the Fiedler value was greater than about 15, we

did detect a minor increase in the proportion infected in networks (Fig. 3c). However, there

was variation in the relationship between the proportion infected and these spectral values

across transmission (β) and recovery probabilities (γ, Figs. 3d-e). For example, when the

probability of transmission was relatively high ($\beta = 0.2$) and recovery low ($\gamma = 0.04$) the

proportion infected across networks was ~80% and spectral radius had a relatively minor

effect (Fig. 3d). A network's spectral radius had a stronger effect when the probability of

recovery was higher ($\gamma = 0.4$) across all values of β. The increase in proportion infected when

the Fiedler value was low (< 15) was not apparent when spread was slower and chances of

recovery higher (e.g., $\beta = 0.025$ or 0.01, $\gamma = 0.4$; Fig 3e). The spectral radius and Fiedler

value patterns overall were similar, with larger values reducing the time-to-peak (Fig. S3).

However, modularity played a greater role in our time to peak models, with the time to peak

being longer for more modular networks above a Q_{rel} threshold of ~ 0.75 (Fig. S5).

Simplifying our models with global surrogates

When we further interrogated our moderate ($\beta = 0.05$) transmission models, we found that the

spectral radius and Fiedler value overall also played a dominant role in our predictions of

spread. To quantify the putative mechanisms that underlie our model predictions – 'to

decloak the black box' – and gain insight into possible interactions between predictors, we

constructed surrogate decision trees as a proxy for our more complex RF model. We trained

our surrogate decision tree on the predictions of the RF model rather than the network

observations directly. In each case, the surrogate decision tree approximated the predictions

of our models (thousands of decision trees) remarkably well (global $R^2 > 0.95$, see (Molnar

2018) for details). The spectral radius and, to a lesser extent, the Fiedler value and modularity

values dominated surrogate trees for all SIR parameter sets (Fig. 5, Figs. S6 & S7). For

example, for networks with a Fiedler value ≥ 0.86 and a spectral radius ≥ 20 (as was the case

for 51% of our networks, Fig. 4b) the estimated maximum proportion of the network infected

was 0.92 (Fig. 4b). The duration over which the data was collected also was included in the

surrogate model, with networks collected over > 6.5 days having higher estimates of

proportion infected (Fig. 4).

Individual networks

To further validate our predictions, we zoomed into how our models predicted spread across

individual networks using Shapley values (assuming $\beta = 0.05$, $\gamma=0.04$). Our models predicted

that the European badger network was less vulnerable to disease spread with a much lower

prediction of proportion infected compared to the average across all networks (Fig. 5a). This

prediction was largely driven by the Fiedler value (0.096, much lower than the mean of 7.31

across all networks) and, to a lesser degree, by the small spectral radius (8.10 compared to a

mean of 34.8 across all networks, Fig. 5a). In contrast, our model predicted a faster time to

peak for the giraffe network (2.04 timesteps on average compared to the global average of

3.08) and this prediction was driven by the high spectral value (106.02, Fig. 5b) even though

the Fiedler value was also high (21.07, Fig. 5b). The Fiedler value was more important in

predicting spread across an ant colony which was more vulnerable to disease (Fig. 5c, 93% of

the nodes were predicted to be infected on average).

Taken together, our findings show how the spectral values of contact networks offer a

valuable and informative "shorthand" for how vulnerable different animal networks are to

outbreaks.

Discussion
358

Here, we show that the spectral radius and Fiedler value of a network can be a remarkably

strong predictor for population vulnerability to diverse epidemics varying in key

epidemiological parameters. We demonstrate how a powerful machine learning and

simulation approach can effectively predict pathogen outbreak characteristics on a large

collection of empirical animal contact networks. We not only demonstrate the high predictive

power of a network's spectral properties but also show that our predictions can help

approximate the vulnerability of populations to infectious disease across outbreak scenarios.

Further, our findings offer insights into how nuances in social organisation translate into

differences in pathogen spread across the animal kingdom.

Across real-world contact networks, we found that the networks' spectral properties (Fiedler

distance and spectral radius) were powerful proxies for pathogen spread. The strong

relationship between spectral radius and epidemic threshold has been demonstrated for

theoretical networks (Prakash *et al.* 2010) and has been used to assess vulnerability of cattle

movement networks to spread of bovine brucellosis (Darbon *et al.* 2018). We expand these

findings to show that the spectral radius is the most important predictor in our models of

epidemic behaviour across diverse animal social systems. While we examined only SIR

propagation through our networks, theoretical results suggest that our findings will extend to

other propagation mechanics such as SIS, (susceptible-infected-susceptible) and SEIR

(susceptible, exposed, infected, recovered) (Prakash *et al.* 2010).

For some networks and epidemiological parameters, spectral radius alone was not sufficient

to predict spread, and the Fiedler value and modularity still played an important role. The

Fiedler value and spectral radius of the networks were correlated, but below our $\rho = 0.7$

threshold (Fig. S2). One potential reason for this is that the Fiedler value seems to be less

sensitive to nodes with high connectivity compared to the spectral radius (Fig. 1); however,

the mathematical relationship between these two algebraic measures of connectivity is poorly

understood (Tang & Priebe 2016). Combined, our global surrogate models and accumulated

effects plots pointed to networks with spectral radii > ~8 and Fiedler values > 1 being more

vulnerable to pathogen spread (the effect of the Fiedler value on spread was much weaker

overall). The spectral properties were dominant for the fast-spreading pathogen models (e.g.,

$\beta > 0.1$, $\gamma = 0.04$), whereas network size and modularity played a more important role in our

models for more slowly spreading pathogens (e.g., Figs. S6 & S7).

When modular structure played a role in disease spread in our study, we detected similar

patterns to those found by Sah *et al.* (2017). As in Sah *et al.* (2017), we found that epidemic

progression was only slowed in highly modular networks ($Q_{rel} > \sim 0.7$) when the probability

of transmission between nodes was low ($\beta > 0.025$). Such subdivided networks were rare in

our data and are commonly associated with high fragmentation (small groups or sub-groups)

and high subgroup cohesion (Sah *et al.* 2017). The reduced importance of modularity relative

to spectral radius is due to within-group connections being crucial for epidemic outcomes in

many contexts (Sah *et al.* 2017). Spectral values may have higher predictive performance, as

they summarize connectivity across the networks including between- and within-group

connections. Interpreting how modularity alone impacted epidemic outcomes was difficult on

these empirical networks, as all modularity measures were strongly correlated with mean

degree, diameter and transitivity (Fig. 2, Fig. S2). The extent of these correlations can vary

substantially based on other aspects of network structure and they all have interacting effects

on disease dynamics (Zhang & Zhang 2009; Ames *et al.* 2011). However, the spectral radius

captures epidemiologically important aspects of network structure on its own, without having

to untangle whether different aspects of network structure are correlated.

More broadly, our study provides a framework for how interpretable machine learning can

predict spread across networks for a wide variety of epidemic parameters. While our RF

MrIML model had much higher predictive performance compared to the corresponding linear

models, further investigation of these models provided critical insight into how network

structure impacted pathogen spread. This framework could identify general trends of disease

vulnerability, specific thresholds for pathogens with certain characteristics, as well as the

drivers of spread for individual networks.

To help practitioners apply our model to different host-pathogen systems, we developed an

R-Shiny app (https://spreadpredictr.shinyapps.io/spreadpredictr/). While network data is hard

to collect, a wide variety of scientists from behavioural ecologists to livestock and wildlife

managers collate network data to address a variety of questions regarding, for example,

health and sociality (Craft 2015). Our web application allows users to take a contact network

of interest and make predictions of pathogen spread for diverse transmission and recovery

probabilities without the need for simulation. While currently limited to pathogens with SIR

transmission dynamics, future versions of the application will include, for example, SI and

SEIR mechanics. We stress that for practitioners to make accurate predictions for a particular

pathogen, contact definitions and the duration of data should be calibrated or multiple

thresholds for what constitutes a transmission contact assessed (see Craft 2015). For

example, for the giraffe network we included edges that represented individuals seen once

together over a period of a year, and predictions of pathogen spread on this network would

likely be inflated for pathogens requiring more sustained contact (VanderWaal *et al.* 2014).

Nonetheless, this study shows the utility of linking network simulation and interpretable

machine learning approaches to tease apart the drivers of spread across empirical wildlife

networks.

As this is a broad comparative study of simulated pathogen spread on 603 empirical networks

across taxonomic groups, we made important simplifying assumptions. For example, as there

were large differences in how the empirical network edges were weighted across taxa (e.g.,

some networks were weighted by contact duration and others by contact frequency) our

approach treated all contacts as equal in unweighted networks, as is done in similar studies

(Ames *et al.* 2011; Sah *et al.* 2017). How or if our spectral-based predictions of spread hold

on networks with weighted edges remains an open research question. Further, while this

study demonstrates the power of repositories such as the ASNR, there are large biases in the

taxa covered that must be accounted for in model structure. For example, a large proportion

of the networks (~150) came from one taxon (*C. fellah*). However, removing this one taxon

did not qualitatively change our findings. While species identity was important to include in

our models to account for the nonindependence of our data, removing species as a predictor

in our models also did not alter our findings. Nonetheless, starting to fill in these taxonomic

gaps in a systematic way will increase the utility of comparative approaches such as ours and

make them generalizable across taxa and populations.

We also simulated spread across static networks, making the assumptions (i) that aggregated

networks are representative of social patterns at epidemiologically relevant timescales and (ii)

that network change happens more slowly than pathogen spread. While this is reasonable for

our study as we are aiming only to identify general rules for how spectral properties influence

epidemiological dynamics in animal social networks, it could be problematic when making

system-level predictions (Bansal *et al.* 2010). Consequently, including predictions of spread

that account for the dynamic nature of contact structure and pathogen-mediated changes in

behaviour is an important future extension of this work. However, applying dynamic network

models such as temporal exponential random graph models (Krivitsky & Handcock 2014) to

estimate spread is computationally demanding and challenging in a comparative setting due

to idiosyncrasies in the model-fitting process. While of high predictive value, our models did

not capture all aspects of uncertainty. For example, we assumed each network was fully

described, with no missing nodes or edges, which is almost always not the case for wildlife

studies. How sensitive spectral properties are to missing data is an open question. However,

interestingly, removing some edges from ecological networks with high Fiedler values does

not appear to strongly impact the stability of the network (Kumar *et al.* 2019).

This paper provides a significant step towards a spectral understanding of pathogen spread in

animal networks. We show that the spectral radius of an animal network is a powerful

predictor of spread for diverse hosts and pathogens that can be a valuable shortcut for

stakeholders to understand the vulnerability of animal networks to disease. We also

demonstrate how multivariate interpretable machine learning models can provide novel

insights into spread across scales. Moreover, this study has identified the key axes of network

structural variation across the animal kingdom that can inform future comparative network

research. As rapid advances in location-based tracking and bio-logging (Katzner & Arlettaz

2020) make network data more readily available to wildlife managers, approaches like this

one will be of increasing value.

481 **Acknowledgements**

482 This project was supported by an Australian Research Council Discovery Project Grant

483 (DP190102020). We would also like to thank Prof. Menna Jones and Prof. Sue VandeWoude

484 for their support and comments on this manuscript.

485

486 **Data statement**

487 All data and code to perform the analysis can be found at

488 https://github.com/nfj1380/igraphEpi

References

Ames, G.M., George, D.B., Hampson, C.P., Kanarek, A.R., McBee, C.D., Lockwood, D.R., *et al.* (2011). Using network properties to predict disease dynamics on human contact networks. *Proceedings of the Royal Society B: Biological Sciences*, 278, 3544–3550.

Apley, D.W. & Zhu, J. (2016). Visualizing the effects of predictor variables in black box supervised learning models. *Journal of the Royal Statistical Society. Series B: Statistical Methodology*, 82, 1059–1086.

Bansal, S., Grenfell, B.T. & Meyers, L.A. (2007). When individual behaviour matters: homogeneous and network models in epidemiology. *J R Soc Interface*, 4, 879–891.

Bansal, S., Read, J., Pourbohloul, B. & Meyers, L.A. (2010). The dynamic nature of contact networks in infectious disease epidemiology. *J Biol Dyn*, 4, 478–489.

Buzdugan, S.N., Vergne, T., Grosbois, V., Delahay, R.J. & Drewe, J.A. (2017). Inference of the infection status of individuals using longitudinal testing data from cryptic populations: Towards a probabilistic approach to diagnosis. *Sci Rep*, 7, 1111.

Corner, L.A.L., Murphy, D. & Gormley, E. (2011). *Mycobacterium bovis* Infection in the Eurasian Badger (*Meles meles*): the disease, pathogenesis, epidemiology and control. *Journal of Comparative Pathology*, 144, 1–24.

Craft, M.E. (2015). Infectious disease transmission and contact networks in wildlife and livestock. *Philosophical transactions of the Royal Society of London. Series B, Biological sciences*, 370, 20140107-.

Craft, M.E., Volz, E., Packer, C. & Meyers, L.A. (2011). Disease transmission in territorial populations: the small-world network of Serengeti lions. *Journal of the Royal Society Interface*, 8, 776–786.

Csárdi, G. & Nepusz, T. (2006). The igraph software package for complex network research. *InterJournal Complex Systems*, 1695.

Darbon, A., Valdano, E., Poletto, C., Giovannini, A., Savini, L., Candeloro, L., *et al.* (2018). Network-based assessment of the vulnerability of Italian regions to bovine brucellosis. *Preventive Veterinary Medicine*, 158, 25–34.

Delahay, R.J., Walker, N., Smith, G.S., Wilkinson, D., CLIFTON-HADLEY, R.S., Cheeseman, C.L., *et al.* (2013). Long-term temporal trends and estimated transmission rates for Mycobacterium bovis infection in an undisturbed high-density badger (Meles meles) population. *Epidemiology and Infection*, 141, 1445–1456.

Eames, K., Bansal, S., Frost, S. & Riley, S. (2015). Six challenges in measuring contact networks for use in modelling. *Epidemics*, 10, 72–77.

Fountain-Jones, N., Kozakiewicz, C., Forester, B., Landguth, E., Carver, S., Charleston, M., *et al.* (2021). MrIML: Multi-response interpretable machine learning to map genomic landscapes. *Molecular Ecology Resources*, 21, 2766–2781.

Fountain-Jones, N.M., Machado, G., Carver, S., Packer, C., Recamonde-Mendoza, M. & Craft, M.E. (2019). How to make more from exposure data? An integrated machine learning pipeline to predict pathogen exposure. *Journal of Animal Ecology*, 88.

Greenwell, B.M., Boehmke, B.C. & Mccarthy, A.J. (2018). A simple and effective model-based variable importance measure. *arXiv Preprint arXiv:1805.04755*.

Hamede, R., Bashford, J., Jones, M. & McCallum, H. (2012). Simulating devil facial tumour disease outbreaks across empirically derived contact networks. *Journal of Applied Ecology*, 49, 447–456.

Hamede, R.K., Bashford, J., McCallum, H. & Jones, M. (2009). Contact networks in a wild Tasmanian devil (*Sarcophilus harrisii*) population: using social network analysis to

reveal seasonal variability in social behaviour and its implications for transmission of
devil facial tumour disease. *Ecology Letters*, 12, 1147–1157.
Katzner, T.E. & Arlettaz, R. (2020). Evaluating contributions of recent tracking-based animal
movement ecology to conservation management. *Frontiers in Ecology and Evolution*,
7.
Krivitsky, P.N. & Handcock, M.S. (2014). A Separable Model for Dynamic Networks. *J R*
Stat Soc Series B Stat Methodol, 76, 29–46.
Kumar, D., Gupta, J. & Raha, S. (2019). Partitioning a reaction–diffusion ecological network
for dynamic stability. *Proc Math Phys Eng Sci*, 475, 20180524.
Kvålseth, T.O. (1985). Cautionary Note about R 2. *The American Statistician*, 39, 279–285.
Lentz, H.H.K., Selhorst, T. & Sokolov, I.M. (2012). Spread of infectious diseases in directed
and modular metapopulation networks. *Phys Rev E Stat Nonlin Soft Matter Phys*, 85,
066111.
Machado, G., Vilalta, C., Recamonde-Mendoza, M., Corzo, C., Torremorell, M., Perez, A., *et*
al. (2019). Identifying outbreaks of Porcine Epidemic Diarrhea virus through animal
movements and spatial neighborhoods. *Scientific Reports*, 9, 457.
McCabe, C.M. & Nunn, C.L. (2018). Effective network size predicted from simulations of
pathogen outbreaks through social networks provides a novel measure of structure-
standardized group size. *Frontiers in Veterinary Science*, 5.
McCallum, H., Jones, M., Hawkins, C., Hamede, R., Lachish, S., Sinn, D.L., *et al.* (2009).
Transmission dynamics of Tasmanian devil facial tumor disease may lead to disease-
induced extinction. *Ecology*, 90, 3379–3392.
McDonald, J.L., Robertson, A. & Silk, M.J. (2018). Wildlife disease ecology from the
individual to the population: Insights from a long-term study of a naturally infected
European badger population. *Journal of Animal Ecology*, 87, 101–112.
Meyers, L.A., Newman, M.E.J. & Pourbohloul, B. (2006). Predicting epidemics on directed
contact networks. *J Theor Biol*, 240, 400–418.
Meyers, L.A., Pourbohloul, B., Newman, M.E.J., Skowronski, D.M. & Brunham, R.C.
(2005). Network theory and SARS: predicting outbreak diversity. *J Theor Biol*, 232,
71–81.
Molnar, C. (2018). *Interpretable machine learning*. Retrieved from
https://christophm.github.io/interpretable-ml-book/.
Newman, M.E.J. (2002). The spread of epidemic disease on networks. *Physical Review E*, 66.
Newman, M.E.J. (2006). Community structure in social and biological networks.
Proceedings of the National Academy of Sciences of the United States of America, 99,
7821–6.
Pastor-Satorras, R., Castellano, C., Van Mieghem, P. & Vespignani, A. (2015). Epidemic
processes in complex networks. *Rev. Mod. Phys.*, 87, 925–979.
Porter, M.A. (2020). Nonlinearity + Networks: A 2020 Vision. *Emerging Frontiers in*
Nonlinear Science, 32, 131–159.
Prakash, B.A., Chakrabarti, D., Faloutsos, M., Valler, N. & Faloutsos, C. (2010). Got the Flu
(or Mumps)? Check the Eigenvalue! *arXiv:1004.0060 [cond-mat, physics:physics, q-*
bio].
Rushmore, J., Caillaud, D., Matamba, L., Stumpf, R.M., Borgatti, S.P. & Altizer, S. (2013).
Social network analysis of wild chimpanzees provides insights for predicting
infectious disease risk. *Journal of Animal Ecology*, 82, 976–986.
Sah, P., Leu, S.T., Cross, P.C., Hudson, P.J. & Bansal, S. (2017). Unraveling the disease
consequences and mechanisms of modular structure in animal social networks. *PNAS*,
114, 4165–4170.

Sah, P., Mann, J. & Bansal, S. (2018). Disease implications of animal social network
structure: A synthesis across social systems. *Journal of Animal Ecology*, 87, 546–558.
Sah, P., Méndez, J.D. & Bansal, S. (2019). A multi-species repository of social networks. *Sci*
Data, 6, 44.
Sah, P., Otterstatter, M., Leu, S.T., Leviyang, S. & Bansal, S. (2021). Revealing mechanisms
of infectious disease spread through empirical contact networks. *PLOS Computational*
Biology, 17, e1009604.
Salathé, M. & Jones, J.H. (2010). Dynamics and Control of Diseases in Networks with
Community Structure. *PLOS Computational Biology*, 6, e1000736.
Shapley, L.S. (1953). Stochastic games. *Proceedings of the National Academy of Sciences of*
the United States of America, 39, 1095–1100.
Shu, P., Wang, W., Tang, M., Zhao, P. & Zhang, Y.-C. (2016). Recovery rate affects the
effective epidemic threshold with synchronous updating. *Chaos*, 26, 063108.
Silk, M.J., Croft, D.P., Delahay, R.J., Hodgson, D.J., Boots, M., Weber, N., *et al.* (2017).
Using Social Network Measures in Wildlife Disease Ecology, Epidemiology, and
Management. *Bioscience*, 67, 245–257.
Štrumbelj, E. & Kononenko, I. (2014). Explaining prediction models and individual
predictions with feature contributions. *Knowledge and Information Systems*, 41, 647–
665.
Tang, M. & Priebe, C.E. (2016). Limit theorems for eigenvectors of the normalized Laplacian
for random graphs. *arXiv:1607.08601 [stat]*.
VanderWaal, K.L., Atwill, E.R., Isbell, L.A. & McCowan, B. (2014). Linking social and
pathogen transmission networks using microbial genetics in giraffe (*Giraffa*
camelopardalis). *The Journal of Animal Ecology*, 83, 406–14.
Weber, N., Carter, S.P., Dall, S.R.X., Delahay, R.J., McDonald, J.L., Bearhop, S., *et al.*
(2013). Badger social networks correlate with tuberculosis infection. *Current Biology*,
23, R915–R916.
White, L.A., Forester, J.D. & Craft, M.E. (2017). Using contact networks to explore
mechanisms of parasite transmission in wildlife. *Biological Reviews*, 92, 389–409.
Worsley-Tonks, K.E.L., Escobar, L.E., Biek, R., Castaneda-Guzman, M., Craft, M.E.,
Streicker, D.G., *et al.* (2020). Using host traits to predict reservoir host species of
rabies virus. *PLOS Neglected Tropical Diseases*, 14, e0008940.
Zhang, Z. & Zhang, J. (2009). A big world inside small-world networks. *PLoS One*, 4, e5686.
618
619

Text Box 1: Terminology used in this paper.

A *graph* (or "*network*") is a collection of *nodes* and a collection of *edges* connecting the

nodes in pairs, e.g., nodes *x*, *y* joined by edge (*x*,*y*). We define the *size* of the network –

usually *n*, as the number of nodes (this usage differs from other strict mathematical

definitions, but we feel this is more intuitive). Two nodes are said to be *adjacent* if they are

connected by an edge, and the number of vertices adjacent to a given vertex *x* is called its

degree, deg(*x*). Edges may be directed, in which case edge (*x*,*y*) is different from edge (*y*,*x*),

but in our analyses we treat them as *undirected*, so (*x*,*y*)=(*y*,*x*). Graphs can be represented

naturally by matrices whose rows and columns are indexed by the nodes (1, 2, …, *n*): the

obvious one is the *adjacency matrix A*, whose (*i,j*)-th entry A_{ij} is 1 if nodes *i* and *j* are

adjacent, and 0 otherwise. *A* is symmetric and $n \times n$, as are all the matrices in this work.

Another useful matrix is the *degree* matrix *D*, in which D_{ij} is the degree of node *i* if *i=j*, and 0

otherwise. The *Laplacian* matrix *L* is the most complex one we use herein, but is easily

calculated using $L_{ij} = D_{ij} - A_{ij}$.

The *eigenvalues* of a matrix are solutions to the matrix equation $Mv = \lambda v$, where *M* is a

matrix and *v* a vector of the appropriate size. Solving for *v* yields λ. These eigenvalues,

ordered by their size, form the *spectrum* of a graph, as derived using any of the matrices just

described. The *Fiedler value* of a graph is the second-smallest eigenvalue of *L*, and the

s*pectral radius* is the largest eigenvalue of *A*.

Measures of *Modularity* such as the Newman *Q* coefficient capture the strength of division

within a network by quantifying the density of edges within and between subgroups. When

there is no division within the network as the density of edges is the same between and within

subgroups $Q = 0$, whereas higher values of *Q* indicate stronger divisions (Newman 2006). As

Q scales with network size (small networks being generally less modular), relative

modularity (Q_{rel}) allows for comparison across network sizes by normalizing Q using the

maximum possible modularity for the network (Q_{max}) (Sah *et al.* 2017).

Fig. 1: Examples of networks analysed in this study with a) the lowest spectral radius
(baboon *Papio cynocephalus* contact network), b) the lowest Fiedler value (vole *Microtus
agrestis* trap sharing network), c) intermediate spectral radius values but high Fiedler value
(Chimpanzee *Pan troglodytes* contact network), d) high spectral radius/intermediate Fiedler
value (*Camponotus fellah* colony contact network) and e) high values of both measures
(another *C. fellah* colony contact network). The mean values across all networks were 34.80
and 7.31 for the spectral radius and Fiedler value respectively. f) summary of values across
networks (a-e). Silhouettes were sourced from phylopic (http://phylopic.org/). Note that
disconnected nodes were not included in the analysis.

Fig. 2: Principal components analysis (PCA) biplot showing that network structure largely clusters by taxonomic class. Points are coloured by taxa. Points closer together in Euclidean space have networks more similar in structure. Points are scaled by network size. The length and direction of vectors (black arrows) shows how each variable relates to each principal component with larger vectors having higher loadings on that axis. The PCA was constructed just using continuous network characteristics. Percentages next to PC scores indicate how much variability in the data is accounted for by each axis. See Table S1 for axis loadings and Fig. S1 for the species-level clustering. See Tables S2 & S3 for variable definitions. Silhouettes for some of the outlying networks were sourced from PhyloPic (http://phylopic.org/). s = scaled. Cent = Centralization.

Fig. 3: Plots showing the predictive performance, variable importance and the functional form of relationships for our best-performing *MrIML* proportion infected model. See Table S4 for model performance estimates across algorithms. The colour of the labels indicates what type of predictor it is (blue = spectral, red = non-spectral network structural variables, gold = network metadata, see Tables S1 & S2). a) Spectral radius and the Fiedler value (followed by species) are the most important predictors of proportion of individuals infected across all simulations (importance threshold >0.1) and overall model performance was high (average $R^2 = 0.96$ and root mean square error (RMSE) = 0.027). b-c) Average predictive surface showing the relationship between spectral properties and proportion infected across all epidemic values (95% confidence intervals in grey). Rug plot on the x axis of the panels on the right shows the distribution of each characteristic across empirical networks. d-e) The accumulated local effects (ALE) plot revealed that the strongly non-linear relationships between both spectral properties and proportion infected were mediated by transmission and recovery probabilities. We chose these SIR parameter values (β = transmission probability, γ = recovery probability) to ensure major outbreaks occurred on the empirical networks. Net construct = Network construction method.

Fig. 4 Global surrogate decision trees for our moderate transmission (β = 0.05) proportion infected with a) high and b) low recovery probability (γ = 0.4 and 0.04 respectively). Threshold values of each variable are included in each tree. The boxes at the tips of the trees indicate the estimates of average peak time or proportion of the network infected across simulations (top value) and percentage of networks in our dataset to be assigned to this tip. For example, 50% of our empirical networks had spectral radius values ≥ 26 and for these networks we found on average, a maximum of 0.76 of the network infected after 100 time steps. Tip boxes are coloured light to dark blue based on network vulnerability to pathogen spread (e.g., longer time to peak = light blue). Global fit = R^2 for how well the surrogate model replicates the predictions of the trained model. See Figs. S6 for the complete list of global surrogate models and Fig. S7 for 'time to peak' surrogates. Colour of the label indicates what type of predictor it is (blue = spectral, red = non-spectral structural variables, gold = network metadata, see Tables S1 & S2).

Fig. 5: Shapely values (phi) that quantify how each variable shaped a) simulated proportion infected in an empirical badger network, b) simulated time to peak in the empirical giraffe network (VanderWaal *et al.* 2014) and c) simulated proportion infected in an ant network (*C. fellah*, note nodes overlap) (β = 0.05, γ=0.04). Panels on the right or below are the corresponding networks. Negative Shapely values indicate that the variable reduced time to peak or proportion infection relative to other variables included in the model. Positive Shapely values indicate that the variables increased the proportion infected relative to other variables included in the model. Colour of the labels indicates what type of predictor it is (blue = spectral, red = non-spectral structural variables, light brown = network metadata, see Tables S2/S3). Corresponding values for each network are next to the label

708