The spectral underpinnings of pathogen spread on animal networks

Authors: Nicholas M. Fountain-Jones[1*], Mathew Silk[2,3], Raima Appaw[1], Rodrigo Hamede[1],
Julie Rushmore[4], Kimberly VanderWaal[5], Meggan Craft[6], Scott Carver[1] & Michael
Charleston[1]

Affiliations:

[1] School of Natural Sciences, University of Tasmania, Hobart Australia 7001.

[2]CEFE, University of Montpellier, CNRS, EPHE, IRD, University of Paul Valéry
Montpellier 3, Montpellier, France

[3]Centre for Ecology and Conservation, University of Exeter, Penryn Campus, Penryn, UK

[4] Odum School of Ecology, University of Georgia, Athens, GA USA.

[5] Department of Veterinary Population Medicine, University of Minnesota, St Paul, MN,
USA

[6] Department of Ecology, Evolution, and Behavior, University of Minnesota, St Paul, MN,
USA

*Nick.FountainJones@utas.edu.au

17
18 **Abstract**
19

Predicting what factors promote or protect populations from infectious disease is a

fundamental epidemiological challenge. Social networks, where nodes represent hosts and

edges represent direct or indirect contacts between them, are key to quantifying these aspects

of infectious disease dynamics. However, understanding the complex relationships between

network structure and epidemic parameters in predicting spread has been out of reach. Here

we draw on advances in spectral graph theory and interpretable machine learning, to build

predictive models of pathogen spread on a large collection of empirical networks from across

the animal kingdom. Using a small set of network spectral properties, we were able to predict

pathogen spread with remarkable accuracy for a wide range of transmissibility and recovery

rates. We validate our findings using well studied host-pathogen systems and provide a

flexible framework for animal health practitioners to assess the vulnerability of a particular

network to pathogen spread.

Introduction

Capturing patterns of direct or indirect contacts between hosts is crucial to model pathogen

spread in populations (Newman 2002; Craft 2015; Sah *et al.* 2018, 2021). Increasingly,

contact network approaches, where hosts are nodes and edges reflect interactions between

hosts, play a central role in epidemiology and disease ecology (e.g., Meyers *et al.* 2005;

Bansal *et al.* 2007; Eames *et al.* 2015; White *et al.* 2017). Incorporating networks allows

models to capture the heterogeneity of contacts between individuals that can provide more

nuanced and reliable estimates of pathogen spread, including in wildlife populations (e.g.,

Meyers *et al.* 2006; Bansal *et al.* 2010; Craft *et al.* 2011). Formulating general rules for how

easy-to-calculate network structure properties may promote or restrict pathogen spread can

reveal important insights into how host behaviour can mediate epidemic outcomes (Sah *et al.*

2017), and provide practitioners with a proxy for how vulnerable a population is to disease

without extensive simulations (Silk *et al.* 2017; Sah *et al.* 2018). Further, network structural

properties can be incorporated into traditional susceptible–infected–recovered (SIR) models

to account for contact heterogeneity when predicting pathogen dynamics across populations

(e.g., Meyers *et al.* 2005; Bansal *et al.* 2007).

However, it remains unclear whether one structural characteristic or a combination of

characteristics can reliably predict pathogen dynamics across systems (Ames *et al.* 2011; Sah

et al. 2018). For example, species that are more social tend to have more clustered or

"modular" networks, and this modularity has been found to increase (Lentz *et al.* 2012),

reduce (Salathé & Jones 2010) or have little effect (Sah *et al.* 2018) on outbreak size across

different biological systems. The average number of contacts between hosts can be identical

across networks and yet still result in substantially different outbreak patterns (Ames *et al.*

2011). Even the apparent size of the network, often constrained by limitations of sampling,

can impact estimates of pathogen spread, particularly in wildlife populations (McCabe &

Nunn 2018). As network characteristics, such as network size and modularity, are often

correlated (Newman 2006; Silk *et al.* 2017) and can have complex impacts on spread (Sah *et*

al. 2017; McCabe & Nunn 2018; Porter 2020), determining network characteristics that

promote large outbreaks, for example, remains a fundamental question in infectious disease

biology (Sah *et al.* 2018).

Searching for general relationships between network structure and pathogen spread in animal

populations is further challenged, as the relationship is also affected by pathogen traits, such

as infectiousness and recovery rate. For example, modularity appears to make no difference

to disease outcomes for highly infectious pathogens (Sah *et al.* 2017). Diseases with long

recovery rates can increase outbreak size across networks as well (Shu *et al.* 2016). Given

that we rarely have reliable estimates of pathogen traits in wild populations (e.g., for different

probabilities of infection per contact, or recovery rates) anyway, any predictive model of the

relationship between spread and network structure would ideally be generalizable across

pathogens.

Advances in spectral graph theory offer an additional set of measures based on the **spectrum**

of a network rather than average node or edge level attributes. A graph spectrum is the set of

eigenvalues (often denoted with a Greek lambda λ) of a matrix representation of a network

(see Text Box 1 for further definitions for terminology in bold). Theoretical studies have

shown relationships between particular eigenvalues and connectivity across networks are

independent of pathogen propagation models (Prakash *et al.* 2010). For example, networks

with a high **Fiedler value** (the second smallest eigenvalue of the network's **Laplacian**

matrix) are "more connected" than those with low values. It has been found that, in

ecological networks for example, if the Fiedler value is sufficiently large, removing edges

will have little effect on overall network connectivity (Kumar *et al.* 2019), but whether this

lack of effect is mirrored by pathogen dynamics is not yet clear. Another quantity of interest

is **spectral radius** – the largest absolute value of the eigenvalues of its **adjacency matrix**.

The link between the spectral radius and epidemiological dynamics is better understood, with

theoretical work showing that this value closely mirrors both epidemic behaviour and

network connectivity (Prakash *et al.* 2010) and has been used to understand vulnerability of

cattle networks to disease (Darbon *et al.* 2018). For example, networks with the same number

of edges and nodes but higher spectral radius (λ_1) are more vulnerable to outbreaks than

networks with low spectral radius ($\lambda_1 \rightarrow 1$). We hypothesize that spectral measures such as

these have great potential to improve our ability to predict dynamics of pathogen spread on

networks, where previous methods such as modularity have proved inadequate (Sah *et al.*

2017).

We assess the predictive capability of spectral values compared to other structural attributes

such as **modularity** (Newmans' Q; Newman 2006)) using advances in machine learning to

construct non-linear models of simulated pathogen spread across a large collection of

empirical animal networks including those from the Animal Social Network Repository

(ASNR) (Sah *et al.* 2019). The ASNR is a large repository of empirical contact networks that

provides novel opportunities to test the utility of spectral values in predicting spread across a

wide variety of, mostly animal, taxa across a spectrum of social systems -- from eusocial ants

(Arthropoda: Formicidae) to more solitary species such as the desert tortoise (*Gopherus*

agassizii). Farmed domestic animals were not included in our analyses. We combined

networks from this resource with other published networks, including badgers (*Meles meles*)

(Weber *et al.* 2013), giraffes (*Giraffa camelopardalis*) (VanderWaal *et al.* 2014) and

chimpanzees (*Pan troglodytes*) (Rushmore *et al.* 2013) to generate a dataset of over 600

unweighted networks from 51 species. We then simulated pathogen spread using a variety of

SIR parameters and harnessed recent advances in multivariate interpretable machine learning

models (*MrIML;* (Fountain-Jones *et al.* 2021)) to construct predictive models across SIR

parameter space. As many species were represented by multiple networks, often over

different populations and or timepoints and constructed in different ways (e.g., some edges

reflected spatial proximity rather than direct contact), we included species and network

construction variables in our models to account for these correlations in addition to exploring

the diversity of network structures across the animal kingdom. Our interpretable machine

learning models identify putative threshold values for the vulnerability of a network to

pathogen spread that can be used by practitioners to understand outbreak risk across systems.

We test how well our network structure estimates of pathogen spread, trained on SIR

simulation results, generalize to more complex pathogen dynamics in the wild We utilize

two well studied wildlife-pathogen systems to assess how our predictions compare to

empirical estimates of spread; *Mycobacterium bovis* (the bacterium that causes bovine

tuberculosis (bTB)) in badger populations and devil facial tumour disease (DFTD) in

Tasmanian devil (*Sarcophilus harrisii*) populations (Hamede *et al.* 2009). We demonstrate

that using spectral measures of network structure alone can provide a useful proxy for disease

vulnerability with estimates of prevalence comparable to those empirically derived. Further,

we provide a user-friendly app that utilizes our models to provide practitioners with

predictions, for example, of the prevalence of a pathogen across a variety of spread scenarios

131 using a user-supplied network without the need for lengthy simulation. The url for this

132 "Shiny" app is https://spreadpredictr.shinyapps.io/spreadpredictr/.

Text Box 1: Terminology used in this paper.

A *graph* (or "*network*") is a collection of *nodes* and a collection of *edges* connecting the

nodes in pairs, e.g., nodes *x*, *y* joined by edge (*x,y*). We define the *size* of the network –

usually *n*, as the number of nodes (this usage differs from other strict mathematical

definitions, but we feel this is more intuitive). Two nodes are said to be *adjacent* if they are

connected by an edge, and the number of vertices adjacent to a given vertex *x* is called its

degree, deg(*x*). Edges may be directed, in which case edge (*x,y*) is different from edge (*y,x*),

but in our analyses we treat them as *undirected*, so (*x,y*)=(*y,x*). Graphs can be represented

naturally by matrices whose rows and columns are indexed by the nodes (1,2,…,*n*): the

obvious one is the *adjacency matrix A*, whose (*i,j*)-th entry A_{ij} is 1 if nodes *i* and *j* are

adjacent, and 0 otherwise. *A* is symmetric and $n \times n$, as are all the matrices in this work.

Another useful matrix is the *degree* matrix *D*, in which D_{ij} is the degree of node *i* if *i=j*, and 0

otherwise. The *Laplacian* matrix *L* is the most complex one we use herein, but is easily

calculated using $L_{ij} = D_{ij} - A_{ij}$.

The *eigenvalues* of a matrix are solutions to the matrix equation $M\mathbf{v} = \lambda\mathbf{v}$, where *M* is a

matrix and $\mathbf{v}$ a vector of the appropriate size. Solving for $\mathbf{v}$ yields λ. These eigenvalues,

ordered by their size, form the *spectrum* of a graph, as derived using any of the matrices just

described. The *Fiedler value* of a graph is the second-smallest eigenvalue of *L*, and the

s*pectral radius* is the largest eigenvalue of *A*.

Measures of *Modularity* such as the Newman *Q* coefficient capture the strength of division

within a network by quantifying the density of edges within and between subgroups. When

there is no division within the network as the density of edges is the same between and within

subgroups $Q = 0$, whereas higher values of *Q* indicate stronger divisions (Newman 2006). As

Q scales with network size (small networks being generally less modular), relative

modularity (Q_{rel}) allows for comparison across network sizes by normalizing Q using the

maximum possible modularity for the network (Q_{max}) (Sah *et al.* 2017).

Results

Diversity of network structures

We identified substantial variation in network structure across animal taxa. The static

unweighted animal social networks in our database ranged from nearly completely

unconnected (Spectral radius $\lambda_1 \sim 1$, Fiedler value ~0, not included in our predictive models)

to highly connected (Spectral radius $\lambda_1 \sim 160$, Fiedler value ~ 140, Fig. 1). Similarly, the

networks ranged from homogeneous (i.e., not modular, $Q_{rel} = 0$, see the Text box for a

definition) to highly modular and subdivided ($Q_{rel} > 0.8$,). Our principal component analysis

(PCA) identified key axes of structural variation across empirical networks (Fig. 2). The first

principal component (PC1) distinguished networks that had a large diameter and mean path

length and were highly modular (negative values), from networks with a high mean degree

and transitivity (positive values, Fig. 2, see Table S2). The second principal component (PC2)

separated networks based on network size (number of nodes), maximum degree and the

network duration (i.e., the time period over which the network data was collected, Fig. 2).

The eusocial ant networks (*Camponotus fellah*, Insecta: Hymenoptera) and mammal

networks tended to cluster separately (Fig. 1), with the other taxonomic classes dispersed

between these groups (Fig. 1) or species (see Fig. S1 for clustering by species). The

networks' spectral properties (the Fiedler value and spectral radius) explained a unique

portion of structural variance that did not covary with other variables (see Table S1 for vector

loadings and Fig S2 for all pair-wise correlations). We found variables such as mean degree

and transitivity the most correlated with the other variables and were excluded from further

analysis (Tables S2, Fig S2).

Fig.1: Examples of networks analysed in this study with a) the lowest spectral radius
(baboons *Papio cynocephalus* contact network), b) the lowest Fiedler value (voles *Microtus
agrestis* trap sharing network), c) intermediate spectral radius values but high Fiedler value
(Chimpanzee *Pan troglodytes* contact network), d) high spectral radius/intermediate Fiedler
value (*Camponotus fellah* colony contact network) and e) high values of both measures
(another *C. fellah* colony contact network). The mean values across all networks were 34.80
and 7.31 for the spectral radius and Fiedler value respectively. f) summary of values across
networks (a-e). Silhouettes were sourced from phylopic (http://phylopic.org/). Note that
disconnected nodes were not included in the analysis.

Fig. 2: Principal components analysis (PCA) biplot showing that network structure largely clusters by taxonomic class. Points are coloured by taxa. Points closer together in Euclidean space have networks more similar in structure. Points are scaled by network size. The length and direction of vectors (black arrows) shows how each variable relates to each principal component with larger vectors having higher loadings on that axis. The PCA was constructed just using continuous network characteristics. Percentages next to PC scores indicate how much variability in the data is accounted for by each axis. Cent*: Centralization. See Table S1 for axis loadings and Fig. S1 for the species-level clustering. See Tables S2 & S3 for variable definitions. Silhouettes for some of the outlying networks were sourced from phylopic (http://phylopic.org/). s = scaled. Cent = Centralization.

Spectral properties predict pathogen spread across epidemic scenarios

We found that network characteristics alone could predict pathogen transmission dynamics

remarkably well (Figs. 3 & Fig S3). We constructed models in *MrIML* to predict the

maximum proportion of nodes infected in the network over 100 time steps (hereafter

'proportion infected'). With these models we could predict the proportion infected in a

network using both spectral measures and species identity alone (Fig. 3a). Network size,

relative modularity and centralization, for example, were less important in predicting

proportion infected across all SIR model parameter combinations tested (Fig. 2a). Nonlinear

relationships were likely important for prediction of proportion infected, as random forests

(RF) had the highest predictive performance overall (Table S4) and substantially

outperformed linear regression in the *MrIML* framework (root mean square error (RMSE)

0.13 vs 0.03). Variable importance and predictor conditional effects were consistent between

the machine learning algorithms, so we subsequently analysed the best performing RF model.

Across all SIR parameter combinations, we found a nonlinear relationship between

proportion infected and spectral radius, with the average prediction of proportion infected

increasing by ~30% across the range of spectral radius values (holding all other variables

constant in the model, Fig. 3b). In contrast we found a more modest effect of the Fiedler

value, with the proportion of infected only increasing on average ~3% across the observed

range of values for all SIR parameters (Fig 3c). We did find a sharp increase in the proportion

infected in networks when the Fiedler value was less than about 15 (Fig. 3c). However, there

was variation in the relationship between the proportion infected and these spectral values

across transmission (β) and recovery probabilities (γ, Figs. 3d-e). For example, when the

probability of transmission was relatively high ($\beta = 0.2$) and recovery low ($\gamma = 0.04$) the

proportion infected across networks was ~80% and spectral radius had a relatively minor

effect (Fig. 3d). A network's spectral radius had a stronger effect when the probability of

recovery was higher ($\gamma = 0.4$) across all values of β. The increase in proportion infected when

the Fiedler value was low (< 15) was not apparent when spread was slower and chances of

recovery higher (e.g., $\beta = 0.025$ or 0.01, $\gamma = 0.4$; Fig 3e). The spectral radius and Fiedler

value patterns overall were similar, with larger values reducing the time-to-peak prevalence

(hereafter 'time to peak', Fig. S3). However, modularity played a greater role in our time to

peak models, with the time to peak being longer for more modular networks above a Q_{rel}

threshold of ~ 0.75 (Fig. S4).

Fig. 3: Plots showing the predictive performance, variable importance and the functional form of relationships for our best-performing *MrIML* proportion infected model. See Table S4 for model performance estimates across algorithms. The colour of the labels indicates what type of predictor it is (blue = spectral, red = non-spectral network structural variables, gold = network metadata, see Tables S2 & S3). a) Spectral radius and the Fiedler value (followed by species) are the most important predictors of proportion of individuals infected across all simulations (importance threshold >0.1) and overall model performance was high (average R^2 = 0.96 and root mean square error (RMSE) = 0.027). b-c) Average predictive surface showing the relationship between spectral properties and proportion infected across all epidemic values (95% confidence intervals in grey). Rug plot on the x axis of the panels on the right shows the distribution of each characteristic across empirical networks. d-e) The accumulated local effects (ALE) plot revealed that the strongly non-linear relationships between both spectral properties and proportion infected were mediated by transmission and recovery probabilities. We chose these SIR parameter values (β = transmission probability, γ = recovery probability) to ensure major outbreaks occurred on the empirical networks. Net construct = Network construction method.

Simplifying our models with global surrogates

When we further interrogated our moderate ($\beta = 0.05$) transmission models, we found that the

spectral radius and Fiedler value overall also played a dominant role in our predictions of

spread. To quantify the putative mechanisms that underlie our model predictions – 'to

decloak the black box' – and gain insight into possible interactions between predictors, we

constructed surrogate decision trees as a proxy for our more complex RF model. We trained

our surrogate decision tree on the predictions of the RF model rather than the network

observations directly. In each case, the surrogate decision tree approximated the predictions

of our models (thousands of decision trees) remarkably well (Global $R^2 > 0.95$, see (Molnar

2018) for details). The spectral radius and, to a lesser extent, the Fiedler value and modularity

values dominated surrogate trees for all SIR parameter sets (Fig. 5, Figs. S5 & S6). For

example, for networks with a Fiedler value ≥ 0.86 and a spectral radius ≥ 20 (as was the case

for 51% of our networks, Fig. 4b) the estimated maximum proportion of the network infected

was 0.92 (Fig. 4b). The duration over which the data was collected also was included in the

surrogate model, with networks collected over > 6.5 days having higher estimates of

proportion infected (Fig. 4b).

Fig. 4 Global surrogate decision trees for our moderate transmission (β = 0.05) proportion infected with a) high and b) low recovery probability (γ = 0.4 and 0.04 respectively). Threshold values of each variable are included in each tree. The boxes at the tips of the trees indicate the estimates of average peak time or proportion of the network infected across simulations (top value) and percentage of networks in our dataset to be assigned to this tip. For example, 50% of our empirical networks had spectral radius values ≥ 26 and for these networks we found on average, a maximum of 0.76 of the network infected after 100 time steps. Tip boxes are coloured light to dark blue based on network vulnerability to pathogen spread (e.g., longer time to peak = light blue). Global fit = R^2 for how well the surrogate model replicates the predictions of the trained model. See Figs. S5 for the complete list of global surrogate models and Fig. S6 for 'time to peak' surrogates. Colour of the label indicates what type of predictor it is (blue = spectral, red = non-spectral structural variables, gold = network metadata, see Tables S2 & S3).

Do our structural estimates generalize to more complex spread scenarios?

To further validate our predictions, we examined how our models predicted *M. bovis* spread across badger networks with empirical estimates using Shapley values (Shapley, 1951). Shapley values are a game-theoretic approach to explore the relative contribution of each predictor on individual networks (see *Methods*). While *M. bovis* in badgers often has a prolonged latent period and individuals do not typically recover, generally *M. bovis* is a slow-spreading infection, with an R_0 of between 1.1 and 1.3 (Delahay *et al.* 2013). Thus, we interrogated our most similar model (β = 0.05, γ = 0.04, R_0 = 1.25). Our model predicted the proportion of infected badgers in the network to be 0.45, which was much lower than the

average proportion infected across all networks included in our study (0.71, Fig. 5a). This

difference was largely driven by the badger network's low Fiedler value (0.096, much lower

than the mean of 7.31 across all networks) and, to a lesser degree, by the small spectral radius

(8.10 compared to a mean of 34.8 across all networks, Fig. 5a). This is comparable to

contemporaneous estimates of *M. bovis* prevalence in this population, e.g., 41% of badgers

tested in the network study tested positive (Weber *et al.* 2013).

We further validated our approach using two Tasmanian devil contact networks (calibrated to

reflect potential DFTD transmission) not included in our training data (Fig. 5) and compared

to model estimate of spread to empirical observations in similar populations. Based on our

model that most closely mirrored devil facial tumour disease DFTD dynamics ($\beta = 0.2$, $\gamma =$

0.04, $R_0 = 5$, see Hamede *et al.* (2012)) we estimated the proportion infected to be 0.85-0.88

for mating and non-mating seasons respectively. Inputting the devil networks' Fiedler value

and spectral radius into the corresponding global surrogate model provides an estimate of

0.89 of individuals in the network infected (Fig. 5b). The spectral values were the most

important predictors in this model (Fig. 5c). Even though our simulations were not

formulated to model DFTD (e.g., devils rarely recover from DFTD), our machine-learning

estimates closely predicted the empirical findings for this disease. In comparable populations

across the island where the disease was monitored before the onset of the disease, maximum

prevalence estimates ranged from 0.7-1.0 in for sexually matured devils (≥ 2 y.o.) ~100

weeks after disease arrival (McCallum *et al.* 2009). Our predictions of proportion infected

were not particularly sensitive to transmissibility estimates as in our model. For example,

with a 50% reduction in the probability of transmission ($\beta = 0.1$) our estimate of proportion

infected was still similar to empirical estimates (0.83, Fig. S5a). Taken together, our findings

show how the spectral values of contact networks offer a valuable and informative

"shorthand" for how vulnerable different animal networks are to outbreaks.

Fig. 5: The spectral radius and the Fiedler underpinned our in-sample prediction of the proportion infected estimates in our a) badger and b/c) out-of-sample Tasmanian devil contact networks. a) Shapley values (φ) that quantify how each variable shaped simulated proportion infected (β = 0.05, γ = 0.04) in an empirical badger network. Negative Shapley values indicate that the variable reduced the proportion infection relative to other variables included in the model. See Fig S7 for other Shapley value analyses of other contrasting networks. b) Surrogate decision tree for the model that best approximated Tasmanian devil facial tumour disease (DFTD, β = 0.2, γ = 0.04). Red lines indicate the branches of the tree

corresponding to the spectral values from the left panels. The red outlined box is the
estimated proportion infected for both networks. c) Corresponding variable importance plot
showing the spectral radius and Fiedler value followed by data duration were the most
important predictors in the model. Colour of the labels indicates what type of predictor it is
(blue = spectral, red = non-spectral structural variables, gold = network metadata, see Tables
S2/S3). Panels on the left are the corresponding networks. Net construct = Network
construction method. *: for sexually mature individuals in comparable populations over
similar time scales to the simulations (McCallum *et al.* 2009).

Discussion

Here, we show that the spectral radius and Fiedler value of a network can be a remarkably

strong predictor for population vulnerability to diverse epidemics varying in key

epidemiological parameters. We demonstrate how a powerful machine learning and

simulation approach can effectively predict pathogen outbreak dynamics on a large collection

of empirical animal contact networks. We not only demonstrate the high predictive power of

a network's spectral properties but also show that our predictions can be a useful tool for

estimating spread in systems with complex disease dynamics. Our findings offer insights into

how nuances in social organisation translate into differences in pathogen spread across the

animal kingdom. Furthermore, our global surrogate models provide animal health

practitioners with an intuitive framework to gain rapid insights into the vulnerability of

populations to the spread of emerging infectious diseases.

Across real-world contact networks, we found that the networks' spectral properties (Fiedler

distance and spectral radius) were powerful proxies for pathogen spread. The strong

relationship between spectral radius and epidemic threshold has been demonstrated for

theoretical networks (Prakash *et al.* 2010) and has been used to assess vulnerability of cattle

movement networks to spread of bovine brucellosis (Darbon *et al.* 2018). We expand these

findings to show that the spectral radius is the most important predictor in our models of

epidemic behaviour across diverse animal social systems. While we examined only SIR

propagation through our networks, theoretical results suggest that our findings will extend to

other propagation mechanics such as SIS, (susceptible-infected-susceptible) and SEIR

(susceptible, exposed, infected, recovered) (Prakash *et al.* 2010). Given that both the badger

M. bovis and DFTD systems have more complex propagation mechanics compared to SIR,

our models could still predict disease dynamics of both disease systems reasonably well. We

that note that for DFTD, disease simulation models that assume homogeneous mixing of

hosts provide similar estimates of disease dynamics to network-based simulations (Hamede *et*

al. 2012). However, Hamede *et al.* (2012) found the outcome of simulated DFTD epidemics

sensitive to estimates of latent period and transmissibility parameters, whereas our network

structure approach provided realistic estimates of prevalence with minimal reliance on

parameter values.

For some networks and epidemiological parameters, spectral radius alone was not sufficient

to predict spread, and the Fiedler value and modularity still played an important role. The

Fiedler value and spectral radius of the networks were correlated, but below our $\rho = 0.7$

threshold (Fig. S2). One potential reason for this is that the Fiedler value seems to be less

sensitive to nodes with high connectivity compared to the spectral radius (Fig. 1); however,

the mathematical relationship between these two algebraic measures of connectivity is poorly

understood (Tang & Priebe 2016). Combined, our global surrogate models and accumulated

effects plots pointed to networks such as the devil networks with spectral radii > ~8 and

Fiedler values > 1 being more vulnerable to pathogen spread (the effect of the Fiedler value

on spread was much weaker overall). The spectral properties were dominant for the fast-

spreading pathogen models (e.g., example system), whereas network size and modularity

played a more important role in our models for more slowly spreading pathogens (e.g., Figs.

S5 & S6).

When modular structure played a role in disease spread in our study, we detected similar

patterns to those found by Sah *et al.* (2017). As in Sah *et al.* (2017), we found that epidemic

progression was only slowed in highly modular networks ($Q_{rel} > {\sim}0.7$) when the probability

of transmission between nodes was low ($\beta > 0.025$). Such subdivided networks were rare in

our data and are commonly associated with high fragmentation (small groups or sub-groups)

and high subgroup cohesion (Sah *et al.* 2017). The reduced importance of modularity relative

to spectral radius is due to within-group connections being crucial for epidemic outcomes in

many contexts (Sah *et al.* 2017). Spectral values may have higher predictive performance, as

they summarize connectivity across the networks including between- and within-group

connections. Interpreting how modularity alone impacted epidemic outcomes was difficult on

these empirical networks, as all modularity measures were strongly correlated with mean

degree, diameter and transitivity (Fig. 2, Fig. S2). The extent of these correlations can vary

wildly based on other aspects of network structure and they all have interacting effects on

disease dynamics (Zhang & Zhang 2009; Ames *et al.* 2011). However, the spectral radius

captures epidemiologically important aspects of network structure on its own without having

to untangle whether different aspects of network structure are correlated.

More broadly, our study provides a framework for how interpretable machine learning can

predict spread across networks for a wide variety of epidemic parameters. While our RF

MrIML model had much higher predictive performance compared to the corresponding linear

models, further investigation of these models provided critical insight into how network

structure impacted pathogen spread. This framework could identify general trends of disease

vulnerability, specific thresholds for pathogens with certain characteristics, as well as the

drivers of spread for individual networks.

To help practitioners apply our model to different host-pathogen systems, we developed an

R-Shiny app (https://spreadpredictr.shinyapps.io/spreadpredictr/). Our web app allows users

to make predictions of spread for diverse transmission and recovery probabilities on a contact

network of interest without the need for simulation. Even when the underlying mechanism of

spread was mis-specified, as with our case studies, our model could provide reasonable

estimates of the proportion of the population infected that align closely with empirical data.

While currently limited to pathogens with SIR transmission dynamics, future versions of the

app will include, for example, SI and SEIR mechanics. We stress that for practitioners to

make accurate predictions for a particular pathogen, contact definitions and the duration of

data should be calibrated or multiple thresholds for what constitutes a transmission contact

assessed (see Craft 2015). For example, for the giraffe network we included edges that

represented individuals seen once together over a period of a year, and predictions of

pathogen spread on this network would likely be inflated for pathogens requiring more

sustained contact (VanderWaal *et al.* 2014). Nonetheless, this study shows the utility of

linking network simulation and interpretable machine learning approaches to tease apart the

drivers of spread across empirical wildlife networks

As this is a broad, comparative study of simulated pathogen spread on 603 empirical

networks across taxonomic groups, we made important simplifying assumptions. For

example, as there were large differences in how the empirical network edges were weighted

across taxa (e.g., some networks were weighted by contact duration and others by contact

frequency) our approach treated all contacts as equal in unweighted networks, as is done in

similar studies (Ames *et al.* 2011; Sah *et al.* 2017). We also simulated spread across static

networks, making the assumptions (i) that aggregated networks are representative or social

patterns at epidemiologically-relevant timescales and (ii) that network change happens more

slowly than pathogen spread. Including predictions of spread that account for the dynamic

nature of contact structure and pathogen-mediated changes in behaviour is an important

future extension of this work. However, applying dynamic network models such as temporal

exponential random graph models (Krivitsky & Handcock 2014) to estimate spread is

computationally demanding and challenging in a comparative setting due to idiosyncrasies in

the model-fitting process. While of high predictive value, our models did not capture all

aspects of uncertainty. For example, we assumed each network was fully described, with no

missing nodes or edges, which is almost always not the case for wildlife studies. How

sensitive spectral properties are to missing data is an open question. However, promisingly,

removing edges from ecological networks with high Fiedler values does not appear to

strongly impact the stability of the network (Kumar *et al.* 2019).

Another limitation of this study is that our models did not account for uncertainty in

predictions. Currently, more probabilistic models such as BART (Bayesian Additive

Regression Trees) (Carlson 2020) are not available in the *MrIML* framework, but future

extensions may allow for methods such as BART to be incorporated (Fountain-Jones *et al.*

2021). However, one advantage of our approach is that for the RF model (proportion

infected), host species (and the other categorical variables, see Table S3) could be added as a

categorical predictor rather than hot-encoded set of 43 predictors (one binary predictor for

each species (-1)). This simplified interpretations about how host species affect pathogen

spread differently, while accounting for nonindependence of intra-species networks (e.g.,

networks for host species A from different populations of that species or from different

timepoints) (Sah *et al.* 2019). A large proportion of the networks (~150) came from one taxon

(*C. fellah*); removing this one taxon did not qualitatively change our findings. While this

study demonstrates the power of repositories such as the ASNR, there are large biases in the

taxa covered that must be accounted for in model structure. Starting to fill in these taxonomic

gaps in a systematic way will increase the utility of comparative approaches such as ours and

make them generalizable across taxa and populations.

This paper provides a significant step towards a spectral understanding of pathogen spread in

animal networks. In particular, we show that the spectral radius of an animal network is a

powerful predictor of spread for diverse hosts and pathogens that can be a valuable shortcut

for stakeholders to understand the vulnerability of animal networks to disease. We also

demonstrate how multivariate interpretable machine learning models can provide novel

insights into spread across scales. Moreover, this study identified the key axes of network

structural variation across the animal kingdom that can inform future comparative network

research. As rapid advances in location-based tracking and bio-logging (Katzner & Arlettaz

2020) make network data more readily available to wildlife managers, approaches like this

one will be of increasing value.

Methods

Networks

We downloaded all animal contact networks from the ASNR on 12[th] January 2022 (Sah *et al.*

2019) and combined these with other comparable published animal contact networks

(Rushmore *et al.* 2013; Weber *et al.* 2013; VanderWaal *et al.* 2014). We binarized each

network, extracted the largest connected component, and excluded networks with fewer than

10 individuals. This left us with 603 networks from 43 species.

From each network we calculated a variety of network structure variables using the R

package *igraph* (Csárdi & Nepusz 2006) (see Table S2). As these networks were constructed

using a wide variety of techniques, we also extracted metadata from the ASNR or the

publication associated with the network (Table S3). These variables were also added to the

models. We used Principal Components Analysis (PCA) biplots to examine the drivers of

variation in network structure and visualise how networks clustered by taxonomic class. We

removed networks with missing metadata (8 networks) and screened for correlations between

variables. As many of the machine learning variables are less sensitive to collinearity

(Fountain-Jones *et al.* 2019) we used a pairwise correlation threshold of 0.7 and removed

variables from the pair with the highest overall correlation (Table S2).

Simulations

To simulate the spread of infection on each network we used our R package "EpicR"

(Epidemics by computers in R; available on GitHub at https://github.com/mcharleston/epicr).

The simulations use a standard discretisation of the SIR model, in which time proceeds in

"ticks," for example representing days. Initially one individual was chosen at uniform random

to be infected (I) and all others were susceptible (S). At each time step, one of two changes of

state can happen to each individual (represented by a node), depending on its current state.

An 'S' individual will become infected (I) with a probability $(1 - (1 - \beta)^k)$, where k is the

number of currently infected neighbours it has, or otherwise stay as S; an 'I' individual will

recover (R) with probability γ or remain as I. Recovered (R) individuals stay as R.

In classical deterministic SIR models as a set of differential equations, β and γ are

instantaneous rates; here, they are probabilities per time step, so at a coarse level, they are

comparable.

On each network, we performed 1000 simulations using different combinations of

transmission (β = 0.01, 0.025, 0.05, 0.1, 0.2) and recovery probabilities (γ = 0.04, 0.4). We

chose these values to broadly reflect a range of scenarios from high to low transmissibility

and slow to fast recovery (Leung 2021) and ensure large outbreaks (>10% on individuals

infected, see Fig S8 for the analysis with a wider variety of recovery rates) (Sah *et al.* 2017).

For each simulation we recorded two complementary epidemic measures to capture disease

burden and speed of spread: a) the maximum prevalence reached, or the maximum proportion

of individuals infected in the network after 100 time steps and b) time to outbreak peak (i.e.,

which time step had the maximum number of infections). We chose 100 time steps to ensure

that the epidemic ended and there were no remaining infected nodes. One randomly chosen

individual was infected at the beginning of the simulation. The average maximum proportion

infected and time to outbreak across all simulations for each parameter combination were

used as the response variables in the machine learning models,

Machine learning pipeline

We used a recently developed multi-response interpretable machine learning approach *(Mr*

IML, Fountain-Jones *et al.* 2020) to predict outbreak characteristics using network structure

variables. Our *MrIML* approach had the advantage of allowing us to rapidly construct and

compare models across a variety of machine-learning algorithms for each of our response

variables as well as assess generalized predictive surfaces across epidemic parameters.

To test the robustness of our results, we compared the performance of four different

underlying supervised regression algorithms in our *MrIML* models. We compared linear

models (LMs), support vector machines (SVMs), random forests (RF) and gradient boosted

models (GBMs) as they operate in markedly different ways that can affect predictive

performance (Fountain-Jones *et al.* 2019; Machado *et al.* 2019). Categorical predictors such

as 'species' were hot-encoded for some models as needed (see Table S4). As both types of

responses in our models were continuous, we compared the performance of each algorithm

using the average R^2 and root mean squared error (RMSE) across all responses (hereafter, the

'global model'). As we included models that were not fit using sums of squares, our R^2

estimate depended on the squared correlation between the observed and predicted values

(Kvålseth 1985). As ants (Insecta: Formicinae) were over-represented, we compared model

performance and interpretation with and without these networks. To calculate each

performance metric, we used 10-fold cross validation to prevent overfitting each model. We

tuned hyperparameters for each model (where appropriate) using 100 different hyper-

parameter combinations (a 10×10 grid search) and selected the combination with the lowest

RMSE. The underlying algorithm with the highest predictive performance was interrogated

further.

We interpreted this final model using a variety of model-agnostic techniques within the

MrIML framework. We assessed overall and model-specific variable importance using a

variance-based method (Greenwell *et al.* 2018). We quantified how each variables alters

epidemic outcomes using accumulated local effects (ALEs) (Apley & Zhu 2016). In brief,

ALEs isolate the effect of each network characteristic on epidemic outcomes using a sliding

window approach calculating the average change in prediction across the values range (while

holding all other variables constant) (Molnar 2018). ALEs are less sensitive to correlations

and straightforward to interpret as points on the ALE curve are the difference from the mean

prediction (Apley & Zhu 2016; Molnar 2018; Fountain-Jones *et al.* 2021).

To further examine the predictive performance of our black-box models (SVM, RF and

GBM) we calculated a global surrogate decision tree (hereafter 'global surrogate') to

approximate the predictions of our more complex trained models. Global surrogates are

generated by training a simpler decision tree to the *predictions* (instead of observations) of

the more complex 'black box' models using the network structure data. How well the

surrogate model performed compared to the complex model is then estimated using R^2. See

Molnar (2018) for details.

Lastly, we gained more insight into model behaviour and how network structure impacted

epidemic outcomes on individual networks, including by calculating Shapley values

(Štrumbelj & Kononenko 2014). Shapley values use a game theoretic approach to play off

variables in the model with each other based on their contribution to the prediction (Shapley

1953). For example, negative Shapley values indicate that the observed value 'contributed to

the prediction' by reducing the proportion infected or time to peak in an outbreak for a

particular network. See Molnar (2018) for a more detailed description and (Fountain-Jones *et*

al. 2019; Worsley-Tonks *et al.* 2020) for how they can be interpreted in epidemiological

settings.

We validated our results using networks with well-documented disease dynamics. The

European badger network was included in our training data, and we selected the propagation

model with a slow recovery rate ($\gamma = 0.04$) and intermediate transmissibility ($\beta = 0.05$) that

provided an equivalent/similar R_0 (1.1-1.3) to *M. bovis* in the studied badger population

(Delahay *et al.* 2013). It should be noted here that *M. bovis* infection has SEI(R)(D)

dynamics, being frequently latent in badgers for long periods with infection only resolving in

some individuals (the most infectious individuals with progressed disease have elevated

mortality (Corner *et al.* 2011)). We compared the proportion infected returned by our model

to various contemporaneous estimates of *M. bovis* prevalence (Delahay *et al.* 2013; Buzdugan

et al. 2017) in the long-term study that contact network data were collected in (McDonald *et*

al. 2018).

The Tasmanian devil networks were not included in the training data. To compare

predictions, we extracted the predict function from the model that was the most similar to

estimates of DFTD dynamics based on empirical data ($\beta = 0.2$, $\gamma = 0.04$, $R_0 = 5$) (McCallum

et al. 2009; Hamede *et al.* 2012). DFTD has SEI(D) dynamics in devil populations, however,

accurately estimating the latent period is impossible, as there is (as of May 2022) no

diagnostic tool to detect DFTD prior to visual detection of the tumours (Hamede *et al.* 2012).

As we wanted to make predictions on a species not included in our dataset, we reran the

models excluding the species predictor and the model performance, and results were very

similar. See https://github.com/nfj1380/igraphEpi for our complete analytical pipeline.

Acknowledgements

This project was supported by an Australian Research Council Discovery Project Grant

(DP190102020). We would also like to thank Prof. Menna Jones and Prof. Sue VandeWoude

for their support and comments on this manuscript.

References

Ames, G.M., George, D.B., Hampson, C.P., Kanarek, A.R., McBee, C.D., Lockwood, D.R., *et al.* (2011). Using network properties to predict disease dynamics on human contact networks. *Proceedings of the Royal Society B: Biological Sciences*, 278, 3544–3550.

Apley, D.W. & Zhu, J. (2016). Visualizing the effects of predictor variables in black box supervised learning models. *Journal of the Royal Statistical Society. Series B: Statistical Methodology*, 82, 1059–1086.

Bansal, S., Grenfell, B.T. & Meyers, L.A. (2007). When individual behaviour matters: homogeneous and network models in epidemiology. *J R Soc Interface*, 4, 879–891.

Bansal, S., Read, J., Pourbohloul, B. & Meyers, L.A. (2010). The dynamic nature of contact networks in infectious disease epidemiology. *J Biol Dyn*, 4, 478–489.

Buzdugan, S.N., Vergne, T., Grosbois, V., Delahay, R.J. & Drewe, J.A. (2017). Inference of the infection status of individuals using longitudinal testing data from cryptic populations: Towards a probabilistic approach to diagnosis. *Sci Rep*, 7, 1111.

Carlson, C.J. (2020). embarcadero: Species distribution modelling with Bayesian additive regression trees in R. *Methods in Ecology and Evolution*, 11, 850–858.

Corner, L.A.L., Murphy, D. & Gormley, E. (2011). *Mycobacterium bovis* Infection in the Eurasian Badger (*Meles meles*): the disease, pathogenesis, epidemiology and control. *Journal of Comparative Pathology*, 144, 1–24.

Craft, M.E. (2015). Infectious disease transmission and contact networks in wildlife and livestock. *Philosophical transactions of the Royal Society of London. Series B, Biological sciences*, 370, 20140107-.

Craft, M.E., Volz, E., Packer, C. & Meyers, L.A. (2011). Disease transmission in territorial populations: the small-world network of Serengeti lions. *Journal of the Royal Society Interface*, 8, 776–786.

Darbon, A., Valdano, E., Poletto, C., Giovannini, A., Savini, L., Candeloro, L., *et al.* (2018). Network-based assessment of the vulnerability of Italian regions to bovine brucellosis. *Preventive Veterinary Medicine*, 158, 25–34.

Delahay, R.J., Walker, N., Smith, G.S., Wilkinson, D., CLIFTON-HADLEY, R.S., Cheeseman, C.L., *et al.* (2013). Long-term temporal trends and estimated transmission rates for Mycobacterium bovis infection in an undisturbed high-density badger (Meles meles) population. *Epidemiology and Infection*, 141, 1445–1456.

Eames, K., Bansal, S., Frost, S. & Riley, S. (2015). Six challenges in measuring contact networks for use in modelling. *Epidemics*, 10, 72–77.

Fountain-Jones, N., Kozakiewicz, C., Forester, B., Landguth, E., Carver, S., Charleston, M., *et al.* (2021). MrIML: Multi-response interpretable machine learning to map genomic landscapes. *Molecular Ecology Resources*, 21, 2766–2781.

Fountain-Jones, N.M., Machado, G., Carver, S., Packer, C., Recamonde-Mendoza, M. &
Craft, M.E. (2019). How to make more from exposure data? An integrated machine learning
pipeline to predict pathogen exposure. *Journal of Animal Ecology*, 88.

Greenwell, B.M., Boehmke, B.C. & Mccarthy, A.J. (2018). A simple and effective model-
based variable importance measure. *arXiv Preprint arXiv:1805.04755*.

Hamede, R., Bashford, J., Jones, M. & McCallum, H. (2012). Simulating devil facial tumour
disease outbreaks across empirically derived contact networks. *Journal of Applied Ecology*,
49, 447–456.

Hamede, R.K., Bashford, J., McCallum, H. & Jones, M. (2009). Contact networks in a wild
Tasmanian devil (*Sarcophilus harrisii*) population: using social network analysis to reveal
seasonal variability in social behaviour and its implications for transmission of devil facial
tumour disease. *Ecology Letters*, 12, 1147–1157.

Katzner, T.E. & Arlettaz, R. (2020). Evaluating contributions of recent tracking-based animal
movement ecology to conservation management. *Frontiers in Ecology and Evolution*, 7.

Krivitsky, P.N. & Handcock, M.S. (2014). A Separable Model for Dynamic Networks. *J R
Stat Soc Series B Stat Methodol*, 76, 29–46.

Kumar, D., Gupta, J. & Raha, S. (2019). Partitioning a reaction–diffusion ecological network
for dynamic stability. *Proc Math Phys Eng Sci*, 475, 20180524.

Kvålseth, T.O. (1985). Cautionary Note about R 2. *The American Statistician*, 39, 279–285.

Lentz, H.H.K., Selhorst, T. & Sokolov, I.M. (2012). Spread of infectious diseases in directed
and modular metapopulation networks. *Phys Rev E Stat Nonlin Soft Matter Phys*, 85, 066111.

Machado, G., Vilalta, C., Recamonde-Mendoza, M., Corzo, C., Torremorell, M., Perez, A., *et
al.* (2019). Identifying outbreaks of Porcine Epidemic Diarrhea virus through animal
movements and spatial neighborhoods. *Scientific Reports*, 9, 457.

McCabe, C.M. & Nunn, C.L. (2018). Effective network size predicted from simulations of
pathogen outbreaks through social networks provides a novel measure of structure-
standardized group size. *Frontiers in Veterinary Science*, 5.

McCallum, H., Jones, M., Hawkins, C., Hamede, R., Lachish, S., Sinn, D.L., *et al.* (2009).
Transmission dynamics of Tasmanian devil facial tumor disease may lead to disease-induced
extinction. *Ecology*, 90, 3379–3392.

McDonald, J.L., Robertson, A. & Silk, M.J. (2018). Wildlife disease ecology from the
individual to the population: Insights from a long-term study of a naturally infected European
badger population. *Journal of Animal Ecology*, 87, 101–112.

Meyers, L.A., Newman, M.E.J. & Pourbohloul, B. (2006). Predicting epidemics on directed
contact networks. *J Theor Biol*, 240, 400–418.

Meyers, L.A., Pourbohloul, B., Newman, M.E.J., Skowronski, D.M. & Brunham, R.C.
(2005). Network theory and SARS: predicting outbreak diversity. *J Theor Biol*, 232, 71–81.

Molnar, C. (2018). *Interpretable machine learning*. Retrieved from
https://christophm.github.io/interpretable-ml-book/.

Newman, M.E.J. (2002). The spread of epidemic disease on networks. *Physical Review E*, 66.

Newman, M.E.J. (2006). Community structure in social and biological networks.
Proceedings of the National Academy of Sciences of the United States of America, 99, 7821–
6.

Porter, M.A. (2020). Nonlinearity + Networks: A 2020 Vision. *Emerging Frontiers in*
Nonlinear Science, 32, 131–159.

Prakash, B.A., Chakrabarti, D., Faloutsos, M., Valler, N. & Faloutsos, C. (2010). Got the Flu
(or Mumps)? Check the Eigenvalue! *arXiv:1004.0060 [cond-mat, physics:physics, q-bio]*.

Rushmore, J., Caillaud, D., Matamba, L., Stumpf, R.M., Borgatti, S.P. & Altizer, S. (2013).
Social network analysis of wild chimpanzees provides insights for predicting infectious
disease risk. *Journal of Animal Ecology*, 82, 976–986.

Sah, P., Leu, S.T., Cross, P.C., Hudson, P.J. & Bansal, S. (2017). Unraveling the disease
consequences and mechanisms of modular structure in animal social networks. *PNAS*, 114,
4165–4170.

Sah, P., Mann, J. & Bansal, S. (2018). Disease implications of animal social network
structure: A synthesis across social systems. *Journal of Animal Ecology*, 87, 546–558.

Sah, P., Méndez, J.D. & Bansal, S. (2019). A multi-species repository of social networks. *Sci*
Data, 6, 44.

Sah, P., Otterstatter, M., Leu, S.T., Leviyang, S. & Bansal, S. (2021). Revealing mechanisms
of infectious disease spread through empirical contact networks. *PLOS Computational*
Biology, 17, e1009604.

Salathé, M. & Jones, J.H. (2010). Dynamics and Control of Diseases in Networks with
Community Structure. *PLOS Computational Biology*, 6, e1000736.

Shapley, L.S. (1953). Stochastic games. *Proceedings of the National Academy of Sciences of*
the United States of America, 39, 1095–1100.

Shu, P., Wang, W., Tang, M., Zhao, P. & Zhang, Y.-C. (2016). Recovery rate affects the
effective epidemic threshold with synchronous updating. *Chaos*, 26, 063108.

Silk, M.J., Croft, D.P., Delahay, R.J., Hodgson, D.J., Boots, M., Weber, N., *et al.* (2017).
Using Social Network Measures in Wildlife Disease Ecology, Epidemiology, and
Management. *Bioscience*, 67, 245–257.

Štrumbelj, E. & Kononenko, I. (2014). Explaining prediction models and individual
predictions with feature contributions. *Knowledge and Information Systems*, 41, 647–665.

Tang, M. & Priebe, C.E. (2016). Limit theorems for eigenvectors of the normalized Laplacian
for random graphs. *arXiv:1607.08601 [stat]*.

VanderWaal, K.L., Atwill, E.R., Isbell, L.A. & McCowan, B. (2014). Linking social and
pathogen transmission networks using microbial genetics in giraffe (*Giraffa camelopardalis*).
The Journal of Animal Ecology, 83, 406–14.

Weber, N., Carter, S.P., Dall, S.R.X., Delahay, R.J., McDonald, J.L., Bearhop, S., *et al.*
(2013). Badger social networks correlate with tuberculosis infection. *Current Biology*, 23,
R915–R916.

White, L.A., Forester, J.D. & Craft, M.E. (2017). Using contact networks to explore
mechanisms of parasite transmission in wildlife. *Biological Reviews*, 92, 389–409.

Worsley-Tonks, K.E.L., Escobar, L.E., Biek, R., Castaneda-Guzman, M., Craft, M.E.,
Streicker, D.G., *et al.* (2020). Using host traits to predict reservoir host species of rabies
virus. *PLOS Neglected Tropical Diseases*, 14, e0008940.

Zhang, Z. & Zhang, J. (2009). A big world inside small-world networks. *PLoS One*, 4, e5686.

731