Pharmacodynamic model of the dynamic response of *Pseudomonas aeruginosa* biofilms to

drug treatments

Swarnima Roychowdhury[1], Charles M. Roth[1,2*]

[1]Department of Biomedical Engineering, Rutgers, The State University of New Jersey,

Piscataway, NJ 08854, USA

[2]Department of Chemical and Biochemical Engineering, Rutgers, The State University of New

Jersey, Piscataway, NJ 08854, USA

*Corresponding author:

Email: cmroth@rutgers.edu (CMR)

Abstract

Chronic infection by gram-negative bacteria such as *Pseudomonas aeruginosa* is a leading cause of morbidity and mortality in cystic fibrosis patients in whom overabundant mucus and the formation of bacterial biofilms pose barriers to drug delivery and effectiveness. Accurate pharmacokinetic-pharmacodynamic (PK-PD) models of biofilm treatment could be used to guide formulation and administration strategies to better control bacterial lung infections. To this end, we have developed a detailed pharmacodynamic model of *P. aeruginosa* treatment with the front-line antibiotics, tobramycin and colistin, and validated it on a detailed dataset of killing dynamics. A compartmental model structure was developed in which the key features are diffusion of drug through a boundary layer to the bacteria, concentration dependent interactions with bacteria, and passage of the bacteria through successive transit states before death. The number of transit states employed was greater for tobramycin, which is a ribosomal inhibitor, than for colistin, which disrupts bacterial membranes. For both drugs, the experimentally observed delay in killing of bacteria following drug exposure was replicated and was consistent with the diffusion time, though for tobramycin, there was an additional delay reflected in the model by passage through the transit states. For each drug, the PD model with a single set of parameters described data across a ten-fold range of concentrations and for both continuous and transient exposure protocols. Furthermore, the parameters fit for each drug individually were used to model the response of biofilms to combined treatment with tobramycin and colistin. The ability to predict drug response over a range of administration protocols allows this PD model to be integrated with PK descriptions to describe in vivo antibiotic response dynamics and to predict drug delivery strategies for improved control of bacterial lung infections.

Author Summary

Biofilms are self-assembling bacterial communities that adhere to a surface and encase themselves in a protective coating. Biofilm infections are notoriously difficult to treat with conventional antibiotic administrations. To understand better the dynamics of bacterial biofilm killing in response to antibiotic treatment, we developed a mathematical model that integrates several features: drug diffusion through a boundary layer that includes the biofilm casing, concentration

dependent cell damage, and passage of the cell through damaged states to eventual death. We
validated the model by comparison with an extensive published dataset of biofilm response to
treatment with the antibiotics, tobramycin and colistin. The model fits to these datasets were able
to capture the observed trends for several antibiotic administration protocols, with model
parameters reflecting the differences in mechanism of action between the two drugs. This validated
model can be integrated with pharmacokinetic descriptions of drug distribution in the body over
time to predict dosing and administration protocols for preclinical and clinical studies.

Introduction

Bacterial biofilms contain cells that adhere to each other to produce a colony of microorganisms,
which is additionally adherent to a surface that may be living or nonliving [1]. The cells within the
biofilm secrete an extracellular polymeric substance (EPS) that encases and protects this colony
from host responses and potential drug treatments [2]. Biofilms occur on a wide range of artificial
and natural surfaces. Biofilm formation has been found in a variety of anatomic settings including
wounds, the ear, and lungs; it accounts for greater than 80% of human microbial infections [3].

In some cases, altered pathophysiology may provide a favorable setting for biofilm formation,
such as the altered mucus composition in patients with cystic fibrosis (CF). Mucin, the
glycoprotein responsible for viscoelastic properties of mucus, is overproduced, and abnormal
glycosylation patterns are observed within CF patients [4]. The mucus-filled environment gives
rise to a breeding ground of bacterial development. Chronic infection by *Pseudomonas*
aeruginosa, a gram-negative bacterium notorious for its antibiotic resistance due to biofilm
formation, is common within 80% of CF patients [3]. Medical devices and instruments may also
be contaminated with *P. aeruginosa*; thus, hospital-acquired infections are not uncommon [5].
Patients infected with *P. aeruginosa* are given antibiotic treatments, such as tobramycin and
colistin, that are only effective in high doses to treat biofilms. These high dosages, in turn, induce
systemic toxicities [6], and their prolonged use can lead to antibiotic resistance [7].

Pharmacokinetic-pharmacodynamic (PK-PD) models are frequently used as tools to design dosing
and administration protocols and as frameworks to interpret experimental results in preclinical
studies. For antibiotic treatments of infection, this is often implemented using static parameters,

such as minimum inhibitory concentration (MIC) for the pharmacodynamics and maximum drug

concentration (C_{max}) or drug area under the curve (AUC) for the pharmacokinetics [8]. The

physical barriers posed and community nature of a bacterial biofilm are such that it may be

necessary to incorporate additional factors, such as the dynamics of drug transport and delayed,

cooperative effects of drugs on biofilm bacteria in order to better describe drug response. Better

experimental quantification of the dynamics of biofilm response to various drug treatments and

their incorporation into pharmacodynamic (PD) models are crucial in understanding and

incorporating the concentration dependent and dynamic effects involved in overcoming biofilm

infections. For example, recent developments in the application of confocal laser scanning

microscopy with flow chambers has enabled monitoring of the real-time killing of bacterial

biofilms [9-11].

In the present work, a rich data set was used to validate a novel PD model for the killing of *P.*

aeruginosa in biofilms by tobramycin and colistin. The proposed model incorporates three

essential components: drug diffusion to the biofilm, nonlinear drug concentration effects on

cellular damage, and a passage through multiple transit states by which the cells eventually become

nonviable. This model was applied to various drug administration experiments that reflect the

dynamic nature of biofilm as well as the cellular mechanisms involved in response to the drug.

Specifically, the model was based on experiments in which *P. aeruginosa* biofilms were exposed

to varying concentrations of drugs. It was validated in its ability to capture the dynamics of killing

in response to transient exposure to one drug and continuous exposure to drug combinations.

Methods

Experimental dataset

A pharmacodynamic model, which captures the effects of drug concentration, drug diffusion, and

cell transit through several states ultimately leading to cell death, was developed to describe

previously reported data regarding tobramycin and colistin treatment of *Pseudomonas aeruginosa*

biofilms in a well-defined flow cell[9]. In the experiments, biofilm populations were established

for 48 hours under flow. Subsequently, the biofilms were provided continuous or transient

treatment of drug using the flow cell system, and data were collected continuously for up to 24

hours. The transit time of the drug within the tube was approximately 90 minutes, which was

accounted for in our model by subtracting 1.5 hours from the raw data. Propidium iodide (PI) dye

was included in the flow solution to stain the nonviable biomass, and the resulting fluorescence

was recorded by automated microscopy and normalized to the maximum fluorescence intensity

recorded. As a result, the experimentally reported quantity to which model predictions were

compared is the "Relative Biovolume," representing the normalized values of dead biovolume.

Pharmacodynamic model

In the proposed pharmacodynamic model (Figure 1), exposure to drug induces healthy biofilm

cells (B) to enter and progress through one or more transit states (D_1, D_2, …) in which the cell

membrane integrity is maintained (i.e., they do not stain with propidium iodide) but the cells are

no longer able to divide. Progression from the last transit state produces dead cells (X),

corresponding experimentally to nonviable biovolume. Mass balances were used to derive kinetic

equations describing the populations of healthy biofilm cells, the respective transit compartments,

and dead cells. For tobramycin administration, the number of transit compartments was determined

by optimization to be five, leading to the following set of balance equations. Colistin

administration followed the same model structure; however, there was only one transit

compartment as opposed to five.

$$\frac{dB^*}{dt} = B^* \cdot \left[\mu \cdot (1 - B^* - D_1^* - D_2^* - D_3^* - D_4^* - D_5^* - X^*) - k_s(\alpha, \beta)C_0^\gamma \right] \quad (1)$$

$$\frac{dD_1^*}{dt} = k_s(\alpha, \beta)C_0^\gamma \cdot B^* - k_t \cdot D_1^* \quad (2)$$

$$\frac{dD_2^*}{dt} = k_t \cdot (D_1^* - D_2^*) \quad (3)$$

$$\frac{dD_3^*}{dt} = k_t \cdot (D_2^* - D_3^*) \quad (4)$$

$$\frac{dD_4^*}{dt} = k_t \cdot (D_3^* - D_4^*) \quad (5)$$

$$\frac{dD_5^*}{dt} = k_t \cdot (D_4^* - D_5^*) \tag{6}$$

$$\frac{dX^*}{dt} = k_t \cdot D_5^* \tag{7}$$

Fig 1. Pharmacodynamic model structure for tobramycin. The pharmacodynamic model for response to tobramycin (subscript 't') tracks the transit of biofilm cells going from a viable (B) to nonviable state (X_t) following administration of tobramycin at bulk concentration C_{0t}. There is a flux, $J(D)$ of drug from the bulk to the biofilm cells, where the local concentration is C_t. For tobramycin, there are five transit compartments (D_{1t}, D_{2t}, D_{3t}, D_{4t} and D_{5t}) mediating the cellular response to drug. Growth is governed by a specific growth rate, μ, the coupled diffusion and pharmacodynamic response are subject to parameters α, β, and γ, and the transit rate to subsequent compartments is given by k_{tt}.

In the above equations, the values for each compartment were normalized, as indicated by the
asterisks, to the maximum biovolume observed in accordance with the experimental data [9]. It is
assumed that all of the cells start in the healthy biofilm state, from which they can proliferate with
a specific growth rate (μ) that is modified with a capacity constraint term (Equation 1). The rate
of healthy cell entry into the transit rates is given in terms of a rate constant, k_s, and the bulk
concentration, C_0, raised to a cooperativity factor, γ (Equation 2). The rate constant is proportional
to the diffusive flux (SI Appendix) and can be expressed in terms of two model parameters, α and
β, each of which is a grouping of physical constants, to give:

$$k_s = \alpha \left[1 + 2 \sum_{n=1}^{\infty} (-1)^n e^{-n^2 \beta t} \right] \tag{8}$$

Biofilm cells affected by tobramycin eventually progress through five compartmental transit states

(Equations 2-6) at a rate of k_t in which they become progressively less viable than the previous

state. In the final compartment (X^*), the biofilm cells are nonviable (dead). It is this quantity that

can be compared with the measured nonviable biovolume.

The coupled set of ordinary differential equations (1) – (7) was solved using an ode45 solver in

MATLAB where the initial relative density of the biofilm state was set to 0.81 for the tobramycin

treatment and 0.89 for the colistin treatment, and the rest of the compartments started with no

biovolume. For each drug (tobramycin and colistin), five adjustable model parameters (μ, f_c, k_t, α

and β) were fit to the composite experimental data [9] across varying respective concentrations

and time courses of 24 hours. An error function was first created to evaluate the squared difference

between the output of the model for a given set of parameters and the given data at a specific

timepoint. This function was then minimized using the MATLAB implementation of genetic

algorithm (ga), which produced the desired parameter values. The initial condition was essentially

an extra parameter within the model. To find these values for each respective drug, values ranging

between 0.70 and 0.95 were tested, and the error from the data and model output were compared.

The initial concentration producing the least error was then used.

Results

A detailed pharmacodynamic model was proposed to describe the dynamic response of

Pseudomonas aeruginosa to the antibiotics, tobramycin and colistin (Fig 1). The available dataset

used to validate this model consists of the amount of dead biofilm (relative biovolume) as a

function of time for several different biofilm concentrations for two different treatment protocols

(tobramycin and colistin). In the proposed model, the drug first must diffuse through a boundary

layer to get to the biofilm. When the drug reaches the biofilm, the biofilm cells go through a

progression of transit compartments, the number of which is specific to and reflects the mechanism

of action of the drug. Progression through the transit compartments is irreversible; consequently,

the cells ultimately die after passing through them. The model was fit to continuous-time data for

the killing of *P. aeruginosa* in a flow-cell chamber [9].

The model was first tested on data for *P. aeruginosa* treated with tobramycin (TOB). The

mechanism of action for tobramycin involves binding to the 30S ribosomal unit, thereby inhibiting

protein synthesis, which gradually incapacitates the bacterium and ultimately induces cell death

[12]. This is a prolonged process, which was modeled using five transit compartments, as described

in the Methods. At the TOB 20 µg/mL concentration, experimentally, there is a delay of

approximately 5.5 hours between drug exposure (with the dead volume of the system already taken

into account) and the emergence of nonviable biovolume, which subsequently increases rapidly

(Fig 2). This behavior is captured by the model following a continuous treatment of tobramycin

for 24 hours. The time required for the drug to diffuse to the biofilm is seen within the flat region

of the graph, and as the cells progress through the transit compartments, they are still viable until

death in the X* compartment. After the composite time for drug diffusion and cellular

compartment transit, there is a rapid increase in the number of dead cells observed experimentally

and predicted by the model. For the other studied concentrations of tobramycin, 5 and 50 µg/mL

(not shown), the proposed model shows the same pattern of delay, progression through the transit

compartments, and increase in the dead biovolume population, all of which are consistent with the

experimental results [9].

Fig 2. **Dynamics of *Pseudomonas aeruginosa* killing in response to tobramycin at 20 μg/mL.** The left panel shows the model comparison of the dead cells, shown in red, with the experimental data, shown in blue. The right panel shows the populations in each of the cellular compartments over the same time course.

A useful pharmacodynamic model should be able to capture not only the dynamics, but also the concentration (dose) dependence of response. To this end, the model's parameters were fit to the ensemble data of 5, 20, and 50 μg/mL tobramycin exposure to produce one set of fit parameters (Table 1). This single set of parameters successfully describes the dynamics of cell killing for the concentration range of 5 - 50 μg/mL (Fig 3). The dependence of the lag time before onset of dead biovolume is consistent with the drug diffusion aspect of the model. Delay due to diffusion is seen for all drug concentrations, and it is amplified for lower concentrations. Because the diffusive flux of drug to the biofilm cells is directly proportional to the concentration driving force, less delay and higher dead biovolume concentrations are observed at shorter times for greater drug

163 concentrations. Consequently, tobramycin concentrations of 20 and 50 µg/mL yield shorter lag
164 times as compared to 5 µg/mL (Fig 3).

Table 1. **Pharmacodynamic model parameters.**

Parameter	Description	Units	TOB Value	CST Value
μ_B	Growth rate of biofilm population	h^{-1}	0.0321	0.0001
α	Rate constant for drug effect on biofilm cells	$(\mu g/mL)^{\gamma-1}\ h^{-1}$	0.0002	0.0082
β	Normalized drug diffusivity	h^{-1}	0.2088	0.3986
γ	Cooperativity in drug effect on biofilm		3.5330	4.4313
k_t	Intercompartmental transit rate of drug	h^{-1}	0.5424	1.8924

Fig 3. Tobramycin model fits across a ten-fold concentration range. The PD model was used to simulate
the treatment of biofilms treated with tobramycin at drug concentrations of 5, 20, and 50 µg/mL. The model

was fit to the ensemble data of all three concentrations. The experimental data are shown in blue, and the model is shown in red.

A key application of a pharmacodynamic model is its use to predict the response to varying drug
administration protocols. Experimental data are available for the response of *P. aeruginosa* to a
transient exposure to tobramycin, where the drug is administered for the first four hours and then
turned off for the remaining twenty hours. The pharmacodynamic model predicts that drug effects
will continue to be observed after removal of drug from the bulk, due to continued flux of
remaining drug through the boundary layer and continued progression of cells through the transit
compartments. As a result, a sharp increase in dead cell biovolume is predicted by the model and
observed experimentally during the period from 5-20 hours after initial exposure, i.e., after the
drug was turned off (Fig 4). At the higher concentrations of 20 and 50 µg/mL, regrowth was
observed in the experimental model about 12 hours after the drug administration ceased. Only at
the lowest TOB concentration of 5 µg/mL was there a reduction in killing in the transient exposure
experiment as compared to continuous exposure. This behavior is explained by the model as being
due to an insufficient amount of drug having diffused into the boundary layer during the four hours
of drug exposure (Fig 4).

Fig 4. Transient exposure of biofilms to tobramycin. The same set of parameters for the continuous data was used to simulate the treatment of biofilms using tobramycin transiently for four hours at the same drug concentrations of 5, 20, and 50 µg/mL. The experimental data are shown in blue, and the model is shown in red.

Experimental data on the treatment of *P. aeruginosa* in the same flow system are available for colistin, whose mechanism of action provides a contrast to that of tobramycin. Colistin is a lipopeptide which binds to phospholipids found on the membrane of the cells and replaces cations [13]. This induces cell rupture and leakage of the inner contents of the cell, leading to death. Because this drug has a more rapid mode of killing than tobramycin, the pharmacodynamic model was modified to contain only one transit compartment (Fig 5A), such that the progression of cells from the exposure to the drug to cell death is more rapid than for tobramycin. Analogously to tobramycin, we fit data for multiple concentrations of colistin (CST) into one set of parameters

191 and used these to model various concentrations of colistin administered continuously over a period
192 of twelve hours.

Fig 5. **The pharmacodynamic model for colistin. (A)** Model structure with a single transit compartment. The symbols have the same meanings as in Figure 1, with subscripts changed from t (tobramycin) to c (colistin). **(B)** Data and model fit for colistin killing of *P. aeruginosa* biofilms at 10 µg/mL. The model fit uses parameters determined by fitting to the ensemble of colistin continuous response data.

As seen with tobramycin, a delay is observed in the response to colistin due to the time required
for diffusion through the boundary layer (Fig 5B). However, the delay is shorter due to the
existence of only one transit compartment and more rapid transit throughout. As with
tobramycin, a single set of parameters accurately describes the ensemble of data over the tested
concentration range (Fig 6). These same parameters were used when applying the model to a
transient exposure to colistin, where the drug was administered for the first four hours and shut
off for the remaining time (Fig 7). The same trend was seen as in the continuous treatment,
where at higher concentrations, the diffusive flux of colistin is greater, therefore resulting in less
delay and rapid onset of cell killing. Additionally, at these high concentrations of colistin, all of
the biofilm cells were observed to become nonviable at earlier time points in comparison to
tobramycin, again largely due to drug-treated cells spending less time in transit compartments.

Fig 6. **Colistin model fits across a ten-fold concentration range.** The PD model was used to simulate the treatment of biofilms using colistin concentrations of 2.5, 10, and 25 µg/mL. The data of all three concentrations were used to fit one set of parameters, which was used to model the various concentrations shown above. The experimental data are shown in blue, and the model is shown in red.

Fig 7. **Transient exposure of biofilms to colistin.** The same set of parameters for the continuous data was used to simulate the treatment of biofilms using colistin for four hours at concentrations of 2.5, 10, and 25 μg/mL. The experimental data are shown in blue, and the model is shown in red.

Since tobramycin and colistin have different mechanisms of action, they might produce additive
or synergistic effects when used in combination. If there are no strong synergies or antagonisms,
the original model may be able to predict outcomes of combined treatments using only the
parameters determined earlier for each respective drug. The model proposed for this mechanism
involves a combination of both treatments running in parallel (Fig 8A). It proved necessary to add
a path by which the biofilm cells could initially be affected by tobramycin or colistin, and the cells
in transit due to (slower acting) tobramycin exposure could be killed directly by (faster acting)
colistin. It is assumed that cells in a transit compartment due to tobramycin were equally likely as
naïve cells to be affected by colistin; thus, this path does not introduce any additional fitting
parameters into the model. Because of the more rapid killing mechanism of colistin, the response

to colistin dominates the experimentally observed and model behaviors (Figure 8B), where the

biofilm cell death occurs at earlier times, even with lower concentrations of colistin.

Fig 8. **Treatment with drug combinations. (A)** A schematic model for the combination treatment with two drugs. Transit compartments resulting from exposure to tobramycin and colistin are considered as parallel death pathways, with the possibility for crossover from the slow-acting tobramycin to fast-acting colistin pathway. The same parameters that were derived earlier for each respective drug were used to predict outcomes of combination treatments. **(B)** Combination treatments were tested using various dose combinations of tobramycin and colistin over the course of 24 hours. The mathematical model is shown in red, and the experimental data are shown in blue.

Discussion

Bacterial biofilms are a significant problem in human infections because they form communities
that both pose physical barriers to drug transport and allow metabolic adaptations that can alter the
pharmacology of antibiotic treatment [14, 15]. An improved understanding of the response of
biofilm-associated bacteria to antibiotic treatment is needed to optimize the administration route
and timing of existing drugs and to focus efforts on novel antibiotic development. Experimental
datasets wherein the response of a biofilm to treatment is monitored continuously over time
provide a signature of the pharmacologic response. The development of a mathematical model that
captures this response serves as a complementary tool that enables the interpretation of these data
in terms of physicochemical mechanisms.

Conventional pharmacologic expressions based on receptor theory are used to describe the
pharmacodynamics (PD) of bacterial response to antibiotics [16, 17]. These in turn are
incorporated into pharmacokinetic-pharmacodynamic (PK-PD) models, which are an important
tool in understanding the dose and time dependence of outcomes in preclinical studies and serve
as the basis for early phase clinical dose and administration scheduling [18, 19]. Traditionally, the
dynamics in PK-PD models of anti-infectives are dictated by the distribution of the drug, and the
concentration dependence is reflected in the pharmacodynamic expression. The simplest such
expression, which is commonly employed in practice, treats the encounter between drug and target
cell as a first-order reaction resulting in instantaneous cell killing [20]. This approach does not
capture important trends observed in preclinical and human infections including a delay between
drug exposure and drug effect and more complex dose response relationships.

More elaborate mathematical models have been proposed to describe the growth and treatment of
biofilms, taking into account physical effects such as diffusion of substrate for growth and of
antibiotic for killing within the biofilm, interfacial detachment of biofilm-associated cells,
advection, and chemotaxis [21-24]. Furthermore, additional cellular states that reflect
heterogeneity of response, e.g. persister states [25], or transitional states due to cell damage[26],
have been incorporated. These models promote our quantitative understanding of the role that
these physical and cell physiologic effects can play in the growth and antibiotic treatment of
biofilms. However, incorporation of mechanisms that depend on both space and time involves
partial differential equation-based continuous or agent-based simulation models that do not
incorporate readily into PK-PD models [27, 28]. We sought to develop a model with sufficient

mechanistic detail as to describe the dynamics of the biofilm response while still being tractable

for eventual incorporation into a PK-PD framework.

Recent experiments that monitor biofilm response dynamically demonstrate that there is a delay

between the onset of drug exposure and cell killing, and that the magnitude of the delay depends

on the particular drug being used [9, 20]. Thus, while diffusion can play a role in the temporal

response, cell physiology and the therapeutic mechanism of action are also evidently important.

Based on these observations, we developed a pharmacodynamic model whose response has two

critical aspects: diffusion of drug through a boundary layer to the cells, and a cell physiological

response in which a cascade of events is initiated whose number and rates can be modified

depending on the mechanism of action of the drug. Specifically, we introduced "transit

compartments" to account for cell states that are affected by drug: non-proliferative, but not yet

dead (Fig 1). Incorporating these elements, the model has five adjustable parameters with distinct

mechanistic interpretations.

The parameter μ_B represents the specific growth rate of the biofilm. As little cell growth is observed

during the time course of the experiments being modeled, its value is low, and no finer detail needs

to be incorporated. The inclusion of drug diffusion results in two lumped parameters, α and β

(Equation 6 and SI Appendix). The β parameter is the value of $\frac{\pi D}{H^2}$, where D is the diffusion

coefficient, and H is the thickness of the diffusion layer, which is a combination of the

hydrodynamic layer resulting from the experimental setup in a flow cell, as well as the physical

barrier imposed by the biofilm itself. The lumped parameter β results from the scaling of the

diffusion problem and is the inverse of the characteristic time for diffusion. Using the biofilm

thickness ~20 μm, the fit values of β would correspond to diffusion coefficients (1.3 - 5.2 x10^{-5}

mm^2/h). These values are several orders of magnitude lower than those for water [29], suggesting

that the diffusivity of the drugs is reduced in the biofilm and/or there is also a mass transfer

boundary layer [30]. For this reason, β was retained as a fit, rather than fixed, parameter. Since the

boundary layer thickness should be the same for both drugs, the slightly higher value fit for colistin

than for tobramycin can be interpreted as a higher effective diffusion coefficient for the former

compound. Although colistin has a higher molecular weight than tobramycin, it has biosurfactant

properties that may allow it to diffuse (penetrate) more rapidly in the biofilm barrier [31].

The α value is a lumping of $\frac{k_C D}{H}$, which includes the aforementioned parameters that describe

diffusive flux, as well as a rate constant, k_C , to denote the rate at which biofilm cells are affected

by the drug and enter into a transit compartment to begin its death cascade. The γ value is a purely

pharmacodynamic parameter representing cooperativity in terms of the drug binding to and

poisoning of the biofilm cells. The last parameter, k_t, corresponds to the intercompartmental transit

rate of the drug. This is not typically found in other PD models that are designed to capture data at

one time point; however, it is critical in capturing the overall dynamic behavior of the drug and its

effect on biofilm killing.

Fig 9. **Impact of model simplifications on performance.** The TOB continuous treatment was used to analyze the parameters within the model. Each TOB concentration was fit to the data without one indicated parameter. The 'Full Model' column shows the fit with the original parameter values found in Table 1.

In order to better understand the influence of model structure and parameter values on the model
output, several analyses were performed. First, each component of the model was removed in turn
and the model re-fit to the experimental data (Fig 9), with the exception of the cell killing
component (α parameter), which would give a trivial result. Removal of either the diffusion term
or the transit compartment terms significantly reduced the ability of the model to capture the
dynamics of the biofilm response across the three different concentration levels of tobramycin.
Likewise, removing cooperativity from the model (i.e., constraining the cooperativity parameter,
γ, to a value of one) prevented accurate reflection of the concentration dependence. Removal of
bacterial growth, μ, did not have a major effect on the model output. However, when neglecting
this parameter in the optimization program, the error between the model and data was increased
by 60% in comparison to incorporating growth rate (Table 2). As a result, the growth rate was
retained in the model.

Table 2. Impact of removing model components on total error

	Full model	Without μ_B	Without β	Without γ	Without k_t
μ_B	0.0321	-	0.5372	0.0289	0.0000
α	0.0002	0.0012	0.0072	0.0094	0.0351
β	0.2088	0.0815	-	0.2480	0.0200
γ	3.5330	3.2862	1.0705	-	3.4315
k_t	0.5424	0.4370	0.4228	0.8053	-
Error	0.0561	0.0971	0.4337	0.7883	1.6449

The effect of these five parameters on the model output can be further understood through a
parametric sensitivity analysis in which each parameter's value is varied while holding all others
constant (Fig 10). It is evident that variation in the value of k_t has the greatest impact on the model
overall. Conceptually, this is expected as the intercompartmental transit rate dictates progress
through the "death cascade" as well as the cellular response dynamics to the drug. The value of β
dictates the delay due to diffusion between drug administration and cellular effects. It couples with
k_t and has a strong influence on the output. The pharmacologic rate constant, α, has a modest effect
on the output, while the cooperativity, γ, exerts a stronger influence.

Fig 10. **Parametric sensitivity analysis.** The TOB 20 µg/mL continuous data were used as the base case. The time course of dead biovolume (X*) production was compared for values of each parameter, varied one at a time. The black line represents the output when the parameter is at its best fit value (Table 1), the red and pink lines show the effect of higher values, and the green and blue lines illustrate the effect of lower values of each parameter.

The number of transit compartments can be considered as an additional model parameter. We
varied the number systematically, refitting the model each time to determine the value most
consistent with the experimental data (Fig 11). For colistin treatment, it was found that one
compartment produces the most accurate model, whereas for tobramycin five compartments
provides the best fit. For either tobramycin or colistin treatment, as the number of compartments
increases, k_t increases in order to mimic the effect of one transit compartment (S1 Table).

Fig 11. **Optimal number of transit compartments.** The TOB 20 µg/mL and CST 10 µg/mL continuous data were used to determine the ideal number of compartments for the model. On the left graph, the model with three compartments gave the least error between the model and data for tobramycin. The graph on the right shows the model with a differing number of compartments for colistin, and as shown, only one compartment produced the least error between the model and data. The parameter values for each compartment can be found in the tables below.

The proposed model fit with a single set of parameters for each respective drug was able to

reproduce the response to drug concentrations that vary over an order of magnitude in both

continuous and transient combined exposure (Figs 3, 4, 6, and 7). It is challenging for a model to

capture both concentration (dose) and time effects. The ability of the present model to do so with

a single set of fit parameters is promising. Furthermore, the effect of combined treatment was

captured accurately using parameters for individual drug treatments (Fig 8B).

Areas where agreement was less robust point to limitations in the model and/or in the

experimental dataset. For instance, in the transient exposure experiments (Fig 4), a reduction in

biovolume (adherent, dead cells) occurred around 20 hours after treatment onset. This represents

detachment, which has been incorporated into some biophysical models of biofilm growth [32].

However, it was not considered in the present work, as the physical location of dead cells

(adherent versus detached) is not of great interest in pharmacologic applications. In addition, the

model somewhat underpredicts the extent of cell death throughout the transient experiment. This

underprediction is because the model parameters were fit based on the continuous treatment

experiments (Fig 3), and the rate of cell killing was more rapid during transient exposure, a result

that most likely reflects experimental variation rather than a physical or pharmacologic effect.

Another limitation is that, because experimental data were collected for only 24 hours,

pharmacologic effects occurring at longer times might not be captured accurately in the model.

This would include effects such as regrowth of the biofilm and development of drug resistance,

both of which tend to evolve over longer periods of time.

In summary, we have shown that a pharmacodynamic model which integrates diffusion of drug

from the bulk to the cells, drug-cell interactions, and a series of transit compartments for affected

cells is able to describe accurately the dynamics of *Pseudomonas aeruginosa* response to

tobramycin, colistin, and their combinations. The model was fit to an ensemble of data covering

multiple drug concentrations to obtain one set of fit parameters for each drug. Among these, the

specific growth rate proved inconsequential during the time course of the experiments studied, but

each of the other four parameters exerted a distinct influence on the model output and contributed

to its ability to capture experimentally observed dynamics. Overall, the model is robust enough to

show the general behaviors of tobramycin, colistin, and the combination of the two drugs at various

concentrations. This pharmacodynamic model can be paired with a pharmacokinetic description

in vivo to predict the drug's effect on an infection. This could be of potential interest in tissues

such as lung where both systemic and regional (e.g., pulmonary) delivery are possible [33]. Our

model can be useful in simulating effects of different strategies in drug administration and

scheduling to promote better eradication of challenging biofilm infections.

References

1.   Donlan RM. Biofilms: microbial life on surfaces. Emerg Infect Dis. 2002;8(9):881-90.

Epub 2002/08/27. doi: 10.3201/eid0809.020063. PubMed PMID: 12194761; PubMed Central

PMCID: PMCPMC2732559.

2.   Martinez-Melendez A, Morfin-Otero R, Villarreal-Trevino L, Baines SD, Camacho-Ortiz

A, Garza-Gonzalez E. Analysis of biofilm production and expression of adhesion structures of

circulating Clostridioides difficile strains from Mexico. Enferm Infecc Microbiol Clin (Engl Ed).

2021. Epub 2021/03/15. doi: 10.1016/j.eimc.2021.01.017. PubMed PMID: 33714640.

3.   Rabin N, Zheng Y, Opoku-Temeng C, Du Y, Bonsu E, Sintim HO. Biofilm formation

mechanisms and targets for developing antibiofilm agents. Future Med Chem. 2015;7(4):493-

512. Epub 2015/04/16. doi: 10.4155/fmc.15.6. PubMed PMID: 25875875.

4.   Ehre C, Ridley C, Thornton DJ. Cystic fibrosis: an inherited disease affecting mucin-

producing organs. Int J Biochem Cell Biol. 2014;52:136-45. Epub 2014/04/02. doi:

10.1016/j.biocel.2014.03.011. PubMed PMID: 24685676; PubMed Central PMCID:

PMCPMC4449140.

5.   Khatoon Z, McTiernan CD, Suuronen EJ, Mah TF, Alarcon EI. Bacterial biofilm

formation on implantable devices and approaches to its treatment and prevention. Heliyon.

2018;4(12):e01067. Epub 2019/01/09. doi: 10.1016/j.heliyon.2018.e01067. PubMed PMID:

30619958; PubMed Central PMCID: PMCPMC6312881.

6.   Krajewski J, Bode-Boger SM, Troger U, Martens-Lobenhoffer J, Mulrooney T,

Mittelstadt H, et al. Successful treatment of extensively drug-resistant Pseudomonas aeruginosa

osteomyelitis using a colistin- and tobramycin-impregnated PMMA spacer. Int J Antimicrob

Agents. 2014;44(4):363-6. Epub 2014/09/04. doi: 10.1016/j.ijantimicag.2014.05.023. PubMed

PMID: 25182711.

7.   Kaiser P, Wachter J, Windbergs M. Therapy of infected wounds: overcoming clinical

challenges by advanced drug delivery systems. Drug Deliv Transl Res. 2021. Epub 2021/02/22.

doi: 10.1007/s13346-021-00932-7. PubMed PMID: 33611768.

8.   Mouton JW, Jacobs N, Tiddens H, Horrevorts AM. Pharmacodynamics of tobramycin in

patients with cystic fibrosis. Diagn Microbiol Infect Dis. 2005;52(2):123-7. Epub 2005/06/21.

doi: 10.1016/j.diagmicrobio.2005.02.011. PubMed PMID: 15964500.

9.   Musken M, Pawar V, Schwebs T, Bahre H, Felgner S, Weiss S, et al. Breaking the

Vicious Cycle of Antibiotic Killing and Regrowth of Biofilm-Residing Pseudomonas aeruginosa.

Antimicrob Agents Chemother. 2018;62(12). Epub 2018/10/10. doi: 10.1128/AAC.01635-18.

PubMed PMID: 30297365; PubMed Central PMCID: PMCPMC6256772.

10.   Lawrence JR, Neu TR. Confocal laser scanning microscopy for analysis of microbial

biofilms. Methods Enzymol. 1999;310:131-44. Epub 1999/11/05. doi: 10.1016/s0076-

6879(99)10011-9. PubMed PMID: 10547787.

11.   Wood SR, Kirkham J, Marsh PD, Shore RC, Nattress B, Robinson C. Architecture of

intact natural human plaque biofilms studied by confocal laser scanning microscopy. J Dent Res.

2000;79(1):21-7. Epub 2000/02/26. doi: 10.1177/00220345000790010201. PubMed PMID:

10690656.

12.   Moustafa DA, Wu AW, Zamora D, Daly SM, Sturge CR, Pybus C, et al. Peptide-

Conjugated Phosphorodiamidate Morpholino Oligomers Retain Activity against Multidrug-

Resistant Pseudomonas aeruginosa In Vitro and In Vivo. mBio. 2021;12(1). Epub 2021/01/14.

doi: 10.1128/mBio.02411-20. PubMed PMID: 33436433; PubMed Central PMCID:

PMCPMC7844538.

13. Chamoun S, Welander J, Martis-Thiele MM, Ntzouni M, Claesson C, Vikstrom E, et al.

Colistin Dependence in Extensively Drug-Resistant Acinetobacter baumannii Strain Is

Associated with ISAjo2 and ISAba13 Insertions and Multiple Cellular Responses. Int J Mol Sci.

2021;22(2). Epub 2021/01/13. doi: 10.3390/ijms22020576. PubMed PMID: 33430070; PubMed

Central PMCID: PMCPMC7827689.

14. Angeles-Martinez L, Hatzimanikatis V. Spatio-temporal modeling of the crowding

conditions and metabolic variability in microbial communities. PLoS Comput Biol.

2021;17(7):e1009140. Epub 2021/07/23. doi: 10.1371/journal.pcbi.1009140. PubMed PMID:

34292935; PubMed Central PMCID: PMCPMC8297787.

15. Martin C, Low WL, Gupta A, Amin MC, Radecka I, Britland ST, et al. Strategies for

antimicrobial drug delivery to biofilm. Curr Pharm Des. 2015;21(1):43-66. Epub 2014/09/06.

doi: 10.2174/1381612820666140905123529. PubMed PMID: 25189862.

16. Jung WJ, Park JH, Goo S, Chae JW, Kim J, Shin S, et al. Dose Optimization of

Vancomycin Using a Mechanism-based Exposure-Response Model in Pediatric Infectious

Disease Patients. Clin Ther. 2021;43(1):185-94 e16. Epub 2020/12/29. doi:

10.1016/j.clinthera.2020.10.016. PubMed PMID: 33358258.

17. Wen X, Gehring R, Stallbaumer A, Riviere JE, Volkova VV. Limitations of MIC as sole

metric of pharmacodynamic response across the range of antimicrobial susceptibilities within a

single bacterial species. Sci Rep. 2016;6:37907. Epub 2016/12/03. doi: 10.1038/srep37907.

PubMed PMID: 27905408; PubMed Central PMCID: PMCPMC5131373.

18. Danhof M, de Lange EC, Della Pasqua OE, Ploeger BA, Voskuyl RA. Mechanism-based

pharmacokinetic-pharmacodynamic (PK-PD) modeling in translational drug research. Trends

Pharmacol Sci. 2008;29(4):186-91. Epub 2008/03/21. doi: 10.1016/j.tips.2008.01.007. PubMed

PMID: 18353445.

19.   Rajman I. PK/PD modelling and simulations: utility in drug development. Drug Discov

Today. 2008;13(7-8):341-6. Epub 2008/04/15. doi: 10.1016/j.drudis.2008.01.003. PubMed

PMID: 18405847.

20.   Haagensen J, Verotta D, Huang L, Engel J, Spormann AM, Yang K. Spatiotemporal

pharmacodynamics of meropenem- and tobramycin-treated Pseudomonas aeruginosa biofilms. J

Antimicrob Chemother. 2017;72(12):3357-65. Epub 2017/09/30. doi: 10.1093/jac/dkx288.

PubMed PMID: 28961810; PubMed Central PMCID: PMCPMC5890705.

21.   Stewart PS. Biofilm accumulation model that predicts antibiotic resistance of

Pseudomonas aeruginosa biofilms. Antimicrob Agents Chemother. 1994;38(5):1052-8. Epub

1994/05/01. doi: 10.1128/AAC.38.5.1052. PubMed PMID: 8067737; PubMed Central PMCID:

PMCPMC188149.

22.   Stewart PS, White B, Boegli L, Hamerly T, Williamson KS, Franklin MJ, et al.

Conceptual Model of Biofilm Antibiotic Tolerance That Integrates Phenomena of Diffusion,

Metabolism, Gene Expression, and Physiology. J Bacteriol. 2019;201(22). Epub 2019/09/11. doi:

10.1128/JB.00307-19. PubMed PMID: 31501280; PubMed Central PMCID: PMCPMC6805107.

23.   Stewart PS, Zhang T, Xu R, Pitts B, Walters MC, Roe F, et al. Reaction-diffusion theory

explains hypoxia and heterogeneous growth within microbial biofilms associated with chronic

infections. NPJ Biofilms Microbiomes. 2016;2:16012. Epub 2017/07/20. doi:

10.1038/npjbiofilms.2016.12. PubMed PMID: 28721248; PubMed Central PMCID:

PMCPMC5515263.

24.   Sheraton MV, Yam JKH, Tan CH, Oh HS, Mancini E, Yang L, et al. Mesoscopic Energy

Minimization Drives Pseudomonas aeruginosa Biofilm Morphologies and Consequent

Stratification of Antibiotic Activity Based on Cell Metabolism. Antimicrob Agents Chemother.

2018;62(5). Epub 2018/02/22. doi: 10.1128/AAC.02544-17. PubMed PMID: 29463543; PubMed

Central PMCID: PMCPMC5923133.

25.   Cogan NG, Szomolay B, Dindos M. Effect of periodic disinfection on persisters in a one-

dimensional biofilm model. Bull Math Biol. 2013;75(1):94-123. Epub 2013/01/09. doi:

10.1007/s11538-012-9796-z. PubMed PMID: 23296996.

26.   Roberts ME, Stewart PS. Modeling antibiotic tolerance in biofilms by accounting for

nutrient limitation. Antimicrob Agents Chemother. 2004;48(1):48-52. Epub 2003/12/25. doi:

10.1128/AAC.48.1.48-52.2004. PubMed PMID: 14693517; PubMed Central PMCID:

PMCPMC310152.

27.   Kindler O, Pulkkinen O, Cherstvy AG, Metzler R. Burst statistics in an early biofilm

quorum sensing model: the role of spatial colony-growth heterogeneity. Sci Rep.

2019;9(1):12077. Epub 2019/08/21. doi: 10.1038/s41598-019-48525-2. PubMed PMID:

31427659; PubMed Central PMCID: PMCPMC6700081.

28.   Mattei MR, Frunzo L, D'Acunto B, Pechaud Y, Pirozzi F, Esposito G. Continuum and

discrete approach in modeling biofilm development and structure: a review. J Math Biol.

2018;76(4):945-1003. Epub 2017/07/26. doi: 10.1007/s00285-017-1165-y. PubMed PMID:

28741178.

29.   Nichols WW, Dorrington SM, Slack MP, Walmsley HL. Inhibition of tobramycin

diffusion by binding to alginate. Antimicrob Agents Chemother. 1988;32(4):518-23. Epub

1988/04/01. doi: 10.1128/AAC.32.4.518. PubMed PMID: 3132093; PubMed Central PMCID:

PMCPMC172213.

30.   Stewart PS. Diffusion in biofilms. J Bacteriol. 2003;185(5):1485-91. Epub 2003/02/20.

doi: 10.1128/JB.185.5.1485-1491.2003. PubMed PMID: 12591863; PubMed Central PMCID:

PMCPMC148055.

31.   Banat IM, De Rienzo MA, Quinn GA. Microbial biofilms: biosurfactants as antibiofilm

agents. Appl Microbiol Biotechnol. 2014;98(24):9915-29. Epub 2014/11/02. doi:

10.1007/s00253-014-6169-6. PubMed PMID: 25359476.

32.   Stewart PS. A model of biofilm detachment. Biotechnol Bioeng. 1993;41(1):111-7. Epub

1993/01/05. doi: 10.1002/bit.260410115. PubMed PMID: 18601252.

33.   Sou T, Kukavica-Ibrulj I, Soukarieh F, Halliday N, Levesque RC, Williams P, et al.

Model-Based Drug Development in Pulmonary Delivery: Pharmacokinetic Analysis of Novel

Drug Candidates for Treatment of Pseudomonas aeruginosa Lung Infection. J Pharm Sci.

2019;108(1):630-40. Epub 2018/09/27. doi: 10.1016/j.xphs.2018.09.017. PubMed PMID:

30257195; PubMed Central PMCID: PMCPMC6307981.

Supplementary information

S1 Appendix. Diffusive flux to biofilm cells.

S1 Table: Effect of transit compartment number on error.