miR167-ARF8, an auxin-responsive module involved in the formation of root-knot nematode-induced galls in tomato

Yara Noureddine[1], Martine da Rocha[1], Jing An[2], Clémence Médina[1], Karine Mulet[1], Joffrey Mejias[1], Michael Quentin[1], Pierre Abad[1], Mohamed Zouine[2], Bruno Favery[1,3] and Stéphanie Jaubert-Possamai[1*].

[1] INRAE, Université Côte d'Azur, CNRS, ISA, F-06903 Sophia Antipolis, France

[2] Laboratoire de Recherche en Sciences Végétales, Université de Toulouse, CNRS, UPS, Toulouse INP, 31320 Auzeville-Tolosane, France.

[3] International Research Organization for Advanced Science and Technology, Kumamoto University, Kumamoto 860-8555, Japan

* Corresponding author: stephanie.jaubert@inrae.fr

Abstract

- Root-knot nematodes (RKN) from genus *Meloidogyne* induce the dedifferentiation of root vascular cells into giant multinucleate feeding cells. These feeding cells result from an extensive reprogramming of gene expression in targeted root cells, as shown by transcriptomic analyses of galls or giant cells from various plant species.

- Small non-coding RNAs, and messenger RNAs from tomato (*Solanum lycopersicum*) galls and uninfected roots were sequenced. *De novo* microRNA prediction in the tomato genome identified microRNAs expressed in galls and uninfected roots. Statistical analyses identified 174 miRNA genes differentially expressed in galls at 7 and/or 14 days post infection (dpi).

- Integrative analyses combining small non-coding RNA and transcriptome datasets with the specific sequencing of cleaved transcripts identified miRNA targets in tomato galls. Functional analyses of promoter-GUS fusions and CRISPR-Cas9 mutants highlighted the role of the miR167-regulated transcription factor AUXIN RESPONSE FACTOR 8 (ARF8) in giant cell formation.

Key words: root-knot nematodes, galls, microRNAs, tomato, auxin, ARF8

Introduction

Root-knot nematodes (RKNs) are major crop pests causing massive yield losses, estimated at millions of Euros annually, worldwide (Blok *et al.*, 2008; Abad & Williamson, 2010). These microscopic worms of genus *Meloidogyne* have a wide host spectrum encompassing more than 5,000 plant species, and a wide geographic distribution. After infecting the root, these obligatory parasites induce the *de novo* formation of a specialized feeding site that is crucial for nematode survival. The second-stage RKN juveniles (J2) penetrate the roots and migrate within them; they then inject a cocktail of molecules into five to seven root parenchyma cells (Favery et al., 2016). In response to RKN signals, targeted root parenchyma cells dedifferentiate into giant multinucleate hypermetabolic feeding cells. These "giant cells" form the feeding site supplying the nematode with the nutrients it requires for its development (Favery *et al.*, 2020). Dedifferentiation into giant cells involves an initial phase of successive mitoses without cytokinesis, followed a second phase of endoreduplication (de Almeida Engler & Gheysen, 2013). During feeding cell formation, the neighboring cells begin to divide. This whole process results in a swelling of the root to form a gall, the characteristic symptom of RKN infection. Feeding site formation involves several biological processes, including the cell cycle (de Almeida-Engler et al., 2011), metabolic reprogramming (Marella *et al.*, 2013), cytoskeleton organization (Caillaud *et al.*, 2008), and auxin signaling (Gheysen & Mitchum, 2019). Auxin (indole-3 acetic acid, IAA), is a major plant hormone that plays a key role in root development by regulating cell division and the establishment/maintenance of root primordia (De Smet *et al.*, 2007; Weijers & Wagner, 2016). The formation of RKN-induced feeding sites has been shown to involve a peak in auxin levels (Karczmarek *et al.*, 2004; Absmanner *et al.*, 2013) and gall transcriptome analyses have shown that auxin biosynthesis and auxin-responsive genes are upregulated in *A. thaliana* early galls, whereas the genes encoding repressors of auxin response genes are repressed (Barcala *et al.*, 2010).

Multiple transcriptome analyses have been performed on RKN-infected roots, galls or specifically on giant feeding cells, from various plant species, including tomato, initially by microarrays, and more recently by RNA sequencing (Bar-Or *et al.*, 2005; Portillo *et al.*, 2013; Shukla *et al.*, 2018). Four time points in feeding site formation have frequently been investigated in transcriptome analyses: the early phase of feeding site formation at 3 days post

infection (dpi), 7 dpi, a time point corresponding to multiple mitoses without cytokinesis, 14

dpi, corresponding to the endoreduplication phase and cell expansion, and, finally, 21 dpi

when the feeding cells are mature and fully functional. All these analyses revealed a massive

reprogramming of plant gene expression in response to nematode infection, with about 10%

of protein-coding genes displaying changes in expression levels in response to RKN infection

(Cabrera et al., 2014). However, it remains unclear how this reprogramming occurs and how

these genes are regulated.

Small non-coding microRNAs may act as the master regulators of this reprogramming of

gene expression (Jaubert-Possamai *et al.*, 2019). MicroRNAs are major repressors of gene

expression in eukaryotes. Within the plant genome, microRNAs are encoded by *MIR* genes,

often organized into multigene families and transcribed as a single-stranded RNA precursor

that folds into a typical hairpin structure. This hairpin precursor is processed to generate a

microRNA duplex consisting of two complementary strands of 20-22 nucleotides. One of the

two strands is then loaded into the ARGONAUTE-1 protein and guides the RNA silencing

complex (RISC) to target messenger RNAs through miRNA/mRNA sequence

complementarity. The targeting of an mRNA by a microRNA induces its degradation or the

inhibition of its translation, depending on the mRNA/miRNA sequence complementarity.

Several recent studies have identified microRNAs expressed in galls (Jaubert-Possamai *et al.*,

2019) induced by RKN in *Arabidopsis thaliana* (Cabrera *et al.*, 2016; Medina *et al.*, 2017), in

tomato (*Solanum lycopersicum;* Zhao et al., 2015; Kaur et al., 2017), in cotton (*Gossypium*

hirsutum; Pan *et al.*, 2019; Cai *et al.*, 2021) or in rice (*Oryza sativa;* Verstraeten *et al.*, 2021).

However, the roles of only four microRNAs have been validated by functional analyses: the

miR390/tasiRNA/*ARF3* module (Cabrera *et al.*, 2016), the miR159/*MYB33* pair (Medina et

al., 2017), the miR172/*TOE1/FT* module (Díaz-Manzano *et al.*, 2018) in *Arabidopsis* and the

miR319/*TCP4* pair in tomato (Zhao et al., 2015).

We investigated the gene regulation network involved in the plant response to RKN through a

combination of transcriptome, microRNome and degradome sequencing in uninfected roots

of tomato and tomato galls induced by the RKN *M. incognita* at two key time points in gall

development: 7 and 14 dpi. We identified 12 miRNA/targeted transcript pairs as robust

candidates for the regulation of gall formation. A key role in tomato gall formation was

demonstrated for the auxin-responsive miR167/*ARF8* transcript pair in functional analyses.

Materials and methods

Biological materials, growth conditions and nematode infection

For *in vitro* experiments, seeds of *Solanum lycopersicum* cv St Pierre or Micro-Tom (wild-type, WT or transgenic *pARF8A::GUS* and *pARF8B::G*US lines (Bouzroud *et al.*, 2018)) were surface-sterilized with chlorine solution (44% active chlorine) and washed three times with water. Ten to 15 sterile seeds were sown on a Gamborg B5 (Duchefa Biochemie) agar plates (1x Gamborg B5; pH = 6.4; 1% Sucrose; 0,7% Agar), placed at 24ºC for 48 hours for germination, and finally transferred in a growth chamber (8h light; 16h dark, 20ºC). *M. incognita* (Morelos strain) J2s were sterilized with HgCl2 (0.01%) and streptomycin (0.7%) as described before (Caillaud & Favery, 2016). One to two weeks after germination, roots were inoculated with 1,000 sterile J2s per petri dishes.

For *in soil* infection assay, *S. lycopersicum* (cv Micro-Tom) WT plants and CRISPR lines (*arf8a*, *arf8b* and *arf8ab*) were sown and individually transferred in pots filled with a mixture of sand and soil (1:1, vol:vol), kept at 4ºC for 48 hours, then transferred in a growth chamber (16h light and 8h dark, at 24ºC). Two weeks after germination, each plant was inoculated with 200 J2s. Infection rate was evaluated six weeks after inoculation. The root system of each plant was collected, rinsed with tap water, weighted and stained for 30 s. in eosin (Sigma) solution (0.5%). Galls and egg masses were counted for each root under the binocular magnifier MZFLIII (Leica). Mann–Whitney U tests ($\alpha = 2.5\%$) were performed to determine the significance of the differences in the numbers of egg masses and galls per root observed between mutants and WT.

BABB clearing

For giant cell area measurements, galls were collected 21 days post-infection (dpi), cleared in benzyl alcohol/benzyl benzoate (BABB) as previously described (Cabrera *et al.*, 2018; Mejias *et al.*, 2021) and examined under an inverted confocal microscope (model LSM 880; Zeiss). The mean areas of giant cells in each gall, for WT and CRISPR lines, for two biological replicates, were measured with Zeiss ZEN software. The impact of the mutation on the giant cell surface was analysed using a Mann & Whitney Test.

RNA extraction

Total RNAs, including small RNAs (< 200 nt), were isolated from *in vitro* tomato (St Pierre)

galls or uninfected root fragments at 7 and 14 dpi. Approximately 40 galls or uninfected roots

devoid meristems were independently frozen into powder by using a tissue lyser (Retsch;

MM301) at 30 Hertz frequency for 30 seconds with 4 mm tungsten balls (Retsch; MM301).

Total RNAs were extracted from these samples with the miRNeasy Mini Kit (Qiagen),

according to the manufacturer's instructions, with three additional washes in RPE buffer.

RNA sequencing

Small RNA libraries were generated by ligation, reverse transcription and amplification (11

cycles) from total RNAs (1 µg), with the reagents of the NEB Next Small RNA Library Prep

Set for Illumina. Libraries were then quantified with the Bioanalyzer High Sensitivity DNA

Kit (Agilent) and sequenced at the Nice-Sophia Antipolis functional genomics platform

(France Géenomique, IPMC, Sophia Antipolis, France). The full raw sequencing data were

submitted to the GEO database (http://www.ncbi.nlm.nih.gov/geo/), accession number

PRJNA799360.

PolyA-RNA libraries from St Pierre tomato were generated from 500 ng of total RNA using

Truseq Stranded mRNA kit (Illumina). Libraries were sequenced on a NextSeq 500 platform

(Illumina) with 2□×□75bp paired-end chemistry as described in (Mejias *et al.*, 2022). RNA-

seq data are available at SRA database accession number #PRJNA799360. PolyA sequencing

of galls from the *arf8* mutants and microtom wild type was performed by the Beiging

Genomics Institute (BGI) by using the DNBSeq technology. RNA-seq data are available at

SRA database accession number #XXX

miRNAome and transcriptome analysis

For each small RNA library, adapters were trimmed and reads matching ribosomal RNA,

mitochondrial RNA and repeat sequences were removed by performing Blast analyses with

the sequences listed in the Rfam database (Nawrocki et al., 2015). The STAR 2.5 aligner (: --

twopassMode Basic --alignEndsType EndToEnd) was then used to align the trimmed reads

(Dobin *et al.*, 2013) on a virtual concatenated genome generated from the *S. lycopersicum*

genome (V3.01, annotation V3.2) and the *M. incognita* genome (Blanc-Mathieu *et al.*, 2017).

Each read was attributed to the *S. lycopersicum* and/or *M. incognita* genome on the basis of

the best alignment obtained. Low-quality mapped reads were removed. The htseq-count

package version 0.9.1 (Anders *et al.*, 2015) was used to count reads mapping perfectly onto

the *S. lycopersicum* genome. The counts for protein coding genes from each replicate were

used for differential expression analysis with the R package EdgeR version 3.4.1 (Robinson *et*

al., 2009) and DSeq2 (Anders & Huber, 2010). Differentially expressed miRNAs, identified

with a false discovery rate of 5% (adjusted pvalue<0.05; Benjamini-Hochberg adjustment).

De novo microRNA encoding genes were predicted in tomato genome V3.0 by using three

algorithms MirCat (Paicu *et al.*, 2017), Shortstack (Axtell, 2013) and MirDeep plant (Yang &

Li, 2011) with default parameters. The sequence homology between newly predicted miRNA

mature sequences and mature miRNA sequences listed in miRBase 22.1 was analysed by

using SSearch algorithm (Kozomara *et al.*, 2019). The HTSEQCOUNT package (Anders *et*

al., 2015) was used to count reads mapping perfectly onto the predicted *S. lycopersicum*

mature microRNA 5P or 3P sequence. Reads mapping to multiple loci were counted for each

of the loci concerned. The counts for mature miRNAs (5P and 3P) from each replicate were

used for differential expression analysis by using DSeq2 statistical analysis (Anders and

Huber, 2010). Mature miRNAs with an adjusted p value below 0.05 were considered as

differentially expressed.

GO analyses of genes differentially expressed in galls were performed by using over-

representation test from PANTHER analysis tools (Mi *et al.* 2019) with a Fisher's exact test,

a FDR threshold of 0.05 and by selecting "Biological Process » as GO category.

Degradome analysis

Degradome libraries were constructed from total RNAs extracted from galls at 7 and 14 dpi

by Vertis Biotechnologie (Freising, Germany) using the parallel analysis of RNA ends

(PARE) protocol described by German *et al*. (2009). The PARE libraries were sequenced on

an Illumina High Sequencing 2000 platform. The full raw sequencing data were submitted to

the GEO database (http://www.ncbi.nlm.nih.gov/geo/), accession XXX. To identify miRNA

targets, degradome reads were analysed and classified by using the CleaveLand 4.0 (Addo-

Quaye *et al.*, 2009) algorithm with default parameters. All hits are classified into five

categories based on the abundance of the diagnostic cleavage tag relative to the overall profile

of degradome tags matching the targets.

GUS staining analysis

We localized the promoter activity in tomatoes transgenic lines expressing a reporter gene

GUS fused to the promoter of the two tomato genes *ARF8A* (*Solyc02g037530*) and *ARF8B*

(*Solyc03g031970*) (pARF8A:GUS and pARF8B:GUS lines) (Bouzroud *et al.*, 2018). We

inoculated 21-day-old seedlings *in vitro,* as described above. GUS staining was performed 7

and 14 dpi as previously described (Noureddine *et al*. 2022), and the roots were observed

under a Zeiss Axioplan 2 microscope. Stained galls were dissected, fixed by incubation in 1%

glutaraldehyde and 4% formaldehyde in 50 mM sodium phosphate buffer pH 7.2, dehydrated,

and embedded in Technovit 7100 (Heraeus Kulzer, Wehrheim, Germany), according to the

manufacturer's instructions. Sections were cut and mounted in DPX (VWR International Ltd,

Poole, UK), and observed under a Zeiss Axioplan 2 microscope (Zeiss, Jena, Germany).

Quantitative RT-PCR

Total RNA was extracted from galls and uninfected roots produced in soil with the miRNeasy

kit (QIAGEN) according to the manufacturer's instruction. 500ng of total RNA were

subjected to reverse transcription with the Superscript IV reverse transcriptase (Invitrogen).

qPCR analyses were performed as described by Nguyen *et al*. (2018). We performed qPCR

on triplicate samples of each cDNA from three independent biological replicates. *SlPSKR1*

(*Solyc01g008140*) and a gene coding for a Sucrose Synthase (*SuSy3*, *Solyc07g042550*) were

used for the normalization of qRT-PCR data. Quantifications and statistical analyses were

performed with SATqPCR (Rancurel *et al.*, 2019), and the results are expressed as

normalized relative quantities. Primers used to amplify the premature miRNAs and the

transcripts are listed in Table S1.

Generation of *ARF8* mutants by CRISPR/Cas9

For CRISPR/Cas9 construct, the sgRNA sequence (AAGCTTTCAACATCAGGAA)

commune to SlARF8a and SlARF8b was designed by using the CRISPR-P website tool

(http://cbi.hzau.edu.cn/crispr/). The sgRNA was cloned into pAGM4723 final vector by

golden gate ligation method. Construct was confirmed by sequencing before introduction into

the C58 *Agrobacterium tumefaciens* strain. Tomato seedlings were used for the next step

plant transformation according to Hao et al. (2015).

Results

Gall formation results from a massive reprogramming of gene expression in root cells

Two statistical methods, DSeq2 and EdgeR, were used to compare transcript levels between

tomato galls and uninfected roots. These two methods were applied to 19,918 genes, and

those found to be differentially expressed by both methods, with an adjusted *p*-value below

0.05, were identified as differentially expressed genes (DEG). We found 1,958 DEGs at 7 dpi

(Table S2) and 3,468 DEGs at 14 dpi **(Table S3 and Figure 1a)**. In total, 1,239 genes were

identified as DEGs at both 7 and 14 dpi, including 625 genes downregulated and 600 genes

upregulated at both time points and 14 DEGs with opposite patterns of change in expression

levels at 7 and 14 dpi. The 719 genes displaying differential expression in galls specifically at

7 dpi comprised 327 upregulated and 392 downregulated genes. The 2,229 genes displaying

differential expression in galls specifically at 14 dpi comprised 1,006 upregulated and 1,223

downregulated genes. The change in gene expression in galls detected by sequencing was

confirmed by RT-qPCR for six of these genes at 7 dpi and six at 14 dpi (**Figure S1 and**

Table S4). Gene ontology (GO) analysis DEGs in galls at 7 and/or 14 dpi revealed an

overrepresentation of genes associated with biological processes previously reported to be

involved in the formation of giant cells **(Table S5),** including i) "cell division", with multiple

categories linked to cytokinesis and cell wall biogenesis, ii) "response to auxin", iii)

"response to endogenous stimulus" (including response to hormone and to cytokinin) and iv)

"response to abiotic stress". This analysis confirms that a massive reprogramming of gene

expression in root cells underlies the formation of galls and feeding cells.

microRNAs regulate gene expression in galls

For the identification of regulators of gene expression in galls, we constructed libraries of

small non-coding RNAs from tomato galls and uninfected roots, from three independent

replicates at two points of gall development: 7 and 14 dpi. These libraries were sequenced,

generating a total of 333,949,327 raw reads **(Table S6)**. The reads were cleaned and mapped

to a virtual genome constructed from the *S. lycopersicum* genome (genome V3.0; ITAG3.3)

concatenated with the genome of *M. incognita* (genome V2.0; Blanc-Mathieu et al., 2017) to

reflect the dual composition of root galls. A *de novo* prediction of microRNAs was then

performed (**Table S7**), based on an integration of the results of three prediction algorithms:

MirCat, Shortstack and MirDeep plant.

Levels of microRNA expression were compared between galls and uninfected roots in DSeq2

statistical analyses (Anders & Huber, 2010). We identified 174 mature microRNAs (5P

and/or 3P) corresponding to 148 *MIR* genes as differentially expressed (DE) between

uninfected roots and galls at 7 and/or 14 dpi **(Table S8)**. We identified 129 of the 174 mature

microRNAs DE in galls as specifically DE at 7 dpi, 11 as specifically DE at 14 dpi and 34

mature microRNAs were found to be DE in galls at both 7 and 14 dpi (**Figure 1b**). These 148

MIR genes DE in galls comprised 65 known *MIR* genes listed in miRbase (Kozomara *et al.*,

2019) and 73 previously unknown *MIR* genes. The 65 known *MIR* genes DE in galls are

organized into 20 miRNA families, 14 of which are conserved between tomato and other

plants, the other six being specific to tomato.

Integration of data from the transcriptome, small RNAs and degradome sequencing to

construct a gene-microRNA regulation network in galls

Once the miRNAs expressed in galls had been identified, the transcripts cleaved by the

microRNAs in galls were identified by degradome sequencing (German *et al.*, 2009) on

mRNA extracted from galls at 7 (G7) and 14 dpi (G14). The CleaveLand pipeline (Addo-

Quaye *et al.*, 2009) was used to analyze degradome sequencing data and to predict the

mRNAs cleaved by miRNAs in galls. We restricted our analysis to the highest confidence

targets by selecting CleaveLand categories 0 and 1 with a degradome *p*-value below 0.05. In

total, 153 transcripts targeted by microRNAs in galls were identified **(Table S9),** including 58

targets common to both the G7 and G14 libraries, whereas 45 targets were identified

specifically in the G7 library and 50 were found only in the G14 library. We identified 111

targets of 135 known miRNAs from 39 known miRNA families. The 298 newly identified

miRNAs in galls included 46 that targeted 47 transcripts in galls at 7 and/or 14 dpi.

We integrated transcriptome, microRNA and degradome sequencing data to construct a a

gene-miRNA regulation network putatively involved in the gall formation. Transcriptome

analysis showed that 32 of the 153 transcripts identified as targeted by microRNAs expressed

in galls were DE in galls. Nineteen of the targeted genes were DE in galls at both 7 and 14

dpi; 11 of these genes were upregulated and eight were downregulated. Five targeted genes

were specifically DE at 7 dpi, including three transcripts that were upregulated and two that

were downregulated in galls. At 14 dpi, only eight transcripts identified as targets were DE,

three of which were upregulated, the other five being downregulated. Most plant miRNAs

silence gene expression by cleaving the targeted transcripts. An inverse correlation of

expression profiles between the microRNA and its target gene is, therefore, usually expected.

We identified 12 miRNA/mRNA pairs for which such an inverse correlation of expression

levels was observed **(Table 1).** These miRNA/mRNA pairs are the most robust candidates for

involvement in gall formation.

ARF8 auxin-related transcription factors are expressed in nematode-induced feeding sites

Among the 12 microRNA/mRNA pairs with opposite patterns of gene expression, two of the strongest genes candidates for regulation by a microRNA are the *AUXIN RESPONSE FACTORS 8A* (*Solyc02g037530*) and *8B* (*Solyc03g031970*), both of which are cleaved miR167. These two genes are ARF transcription factors, which relay auxin signaling at the transcriptional level by regulating the expression of auxin-responsive genes (Guilfoyle & Hagen, 2007). Transcriptomic analyses of galls showed that *ARF8B* was overexpressed in tomato galls at 7 and 14 dpi and *ARF8A* was overexpressed at 14 dpi **(Table 1)**.

Five *MIR167* genes were identified in the tomato genome, including the four described by Liu *et al*. (2014), all of which have the same mature sequence and are downregulated in galls, whereas both the *ARF8* genes were found to be upregulated. The fifth *MIR167* gene was annotated as SLYMIR167B in miRBase (Kozomara *et al.*, 2019) and was not DE in galls. However, it was expressed at a much lower level than the other four *MIR167* genes (**Figure S2**). *ARF8B* transcripts have been shown to be cleaved by miR167 in tomato (Liu *et al.*, 2014) and this regulation in conserved in *A. thaliana* (Wu *et al.*, 2006). The downregulation of miR167 and the upregulation of *ARF8A* and *ARF8B* observed in galls suggest that, by repressing *MIR167* expression, the RKN prevents the *ARF8* silencing by miR167 that occurs in uninfected roots.

We investigated the spatiotemporal expression of *ARF8A* and *ARF8B* in RKN-infected roots *in vivo* further, by analyzing the activity of both the *ARF8A* and *ARF8B* promoters in transgenic tomato lines expressing promoter-GUS fusions (Bouzroud *et al.*, 2018). Strong blue staining indicating GUS activity was observed 7 and 14 dpi in galls from two independent *pARF8B::GUS* and *pARF8A::GUS* lines and in root tips from uninfected roots **(Figure 2a-f)**. Histological sections of the galls showed strong GUS staining within the giant feeding cells and in neighboring cells (NC) at 7 and 14 dpi for both *ARF8A* and *ARF8B* lines **(Figure 3a-d)**. The strong activity of both the *ARF8A* and *ARF8B* promoters observed in galls *in vivo* confirms the upregulation in galls observed on transcriptomic analysis.

Generation of *SlARF8A* and *SlARF8B* mutants by the CRISPR/Cas9 gene-editing system.

We investigated the function of *SlARF8A* and *SlARF8B* during plant-nematode interaction, by

generating tomato Micro-Tom *slarf8* KO mutants with CRISPR/Cas9 gene-editing

technology. *SlARF8A* and *SlARF8B* single and double mutants were obtained with a sgRNA

complementary to a region identical in both genes (Figure 4a). The transformed lines were

screened and six independent R0 lines were generated and validated for the presence of the

construct in their genome. All mutants had mutations in the targeted region and the features

of the mutant plants were similar. In the R1 and R2 generations, the presence of mutation was

confirmed in more progeny lines. Three Cas9-free and homozygous mutant lines containing

single (*SlARF8A* or *SlARF8B)* or double mutation (*SlARF8A* and *SlARF8B)* were selected in

the following experiments. These three mutants types can be classified as *SlARF8A* single

mutation (*slarf8acr*), *SlARF8B* single mutation (*slarf8b-cr*) and *SlARF8A&B* double mutation

(*slarf8a&b-cr*) (Figure 4b). As shown in figure 4, the deletion/insertion mutations led to a

frame shift mutation followed by an early stop codon leading to the expression of truncated

SlARF8 proteins that do not contain the ARF family functional domains B3, III and IV. For

slarf8a-cr, a deletion of 2 nt was detected in the sgRNA1 targeted region, leading to a 13

amino acid (aa) protein rather than the 845 aa WT protein; for *slarf8b-cr*, an 11nt insertion

was observed in the sgRNA1 target region, leading to a 9 aa protein sequence rather than 843

aa in WT protein; and, for *slarf8a&b-cr*, there was a 2 nt deletion on *SlARF8A* and a 4 nt

deletion on *SlARF8B* resulting in a 13 aa SlARF8A protein and a 20 AA SlARF8B protein.

ARF8 auxin-related transcription factors are involved in tomato-RKN interactions

We investigated the role of both *ARF8A* and *ARF8B* in gall development, by analyzing the

effect of CRISPR deletions within *ARF8* coding sequences on *M. incognita* infection. The

arf8a[CR-2], *arf8b*[CR-11] and *arf8ab*[CR-2,4] double-mutant CRISPR lines had root phenotypes

identical to that of the WT **(Figure S3 and Table S10)**. The rate of infection of these

CRISPR lines after inoculation with *M. incognita* was determined by counting the galls on

infected roots and the egg masses produced by the adult females at the root surface at the end

of the RKN lifecycle. A significant large decrease, by approximately 50%, in the numbers of

galls and egg masses was observed for the *arf8a*[CR-2], *arf8b*[CR-11] and *arf8ab*[CR-2,4] lines relative

to WT plants **(Figure 5a and Table S10)**. Thus, *ARF8* disruption decreases susceptibility to

infection, thereby demonstrating that *ARF8A* and *ARF8B* play a key role in the plant-RKN

interaction. We investigated the reasons for the lower susceptibility of the *arf8a*[CR-2], *arf8b*[CR-11]

[11] and *arf8ab*[CR-2,4] lines, by measuring the area covered by giant cells directly with a confocal

microscope, after gall clearing with BABB (Cabrera *et al.* 2017). A comparison of the mean

surface areas covered by giant cells in each gall showed that giant cells from the CRISPR

lines were approximately 30% smaller than those from control plants (**Figure 5b-c**). These

results demonstrate that the expression of *ARF8A* and *ARF8B* is required for correct giant cell

development during the tomato-RKN interaction.

Identification of *ARF8A*- and *ARF8B*-regulated genes in galls

For the identification of genes regulated by ARF8A and ARF8B in galls, mRNA from 14 dpi

galls of *arf8a^{CR-2}, arf8b^{CR-11}* mutants and WT Micro-Tom tomatoes were sequenced.

Transcript levels in the galls from WT and mutant plants were compared in DESeq2 and

EdgeR statistical analyses. These two methods identified 189 and 66 genes, respectively, as

differentially expressed between galls from WT and *arf8a^{CR-2}* or *arf8ab$^{CR-2,4}$* mutants (**Table**

S11a and b). Several auxin-inducible genes have already been identified as candidate genes

downstream from ARF8s in tomato floral tissue (Liu *et al*. 2014): *SMALL AUXIN-*

UPREGULATED RNAs (SAUR; *Solyc07g042490.1.1*) and two *EXPANSINS*

(*Solyc08g077900.3.1* and *Solyc08g077910.3.1*) were found to be repressed in *arf8a^{CR-2}*tomato

galls. Only 16 DEGs were common to both mutants (**Table S11c**). These genes are located

directly or indirectly downstream from ARF8 in tomato galls. No defense marker genes, such

as orthologs of salicylic acid-mediated response marker *PATHOGENESIS-RELATED*

PROTEIN-1 (PR1), jasmonic acid-mediated defense maker *PROTEINASE INHIBITOR*

2 (*PIN2*) or *PLANT DEFENSINs,* were found to be induced in *arf8a^{CR-2}* or *arf8ab$^{CR-2,4}$* galls.

This absence of defense marker gene induction in the galls of *arf8a^{CR-2}* and *arf8ab$^{CR-2,4}$*

mutants indicates that the decrease in galls and egg masses observed is due to a loss of

susceptibility rather than the induction of a plant defense mechanism. Together with the

requirement of ARF8s for correct giant cell development, this findings supports a key role for

ARF8s in feeding site formation.

Discussion

Identification of miRNA/mRNA target pairs involved in gall formation

RKN of the genus *Meloidogyne* are highly polyphagous sedentary plant parasites that can

induce the formation of giant feeding cells in most crop species. The formation of feeding

cells by RKN has been shown to result from a massive reprogramming of plant gene

expression induced by the nematode (Jammes *et al.*, 2005; Fuller *et al.*, 2007; Ibrahim *et al.*,

2011; Damiani *et al.*, 2012; Yamaguchi *et al.*, 2017; Kaur *et al.*, 2017). In this study, we

identified 4187 protein-coding genes, corresponding to 12.3% of all annotated tomato genes,

as differentially expressed in tomato galls 7 and 14 dpi with *M. incognita* relative to

uninfected roots. This proportion is consistent with previous transcriptomic analyses in

Arabidopsis and tomato (Yamaguchi *et al.* 2017; Portillo *et al*. 2013). We investigated the

regulators of this reprogramming of gene expression, by analyzing the expression of

microRNAs and mRNAs in galls and uninfected tomato roots and using degradome

sequencing to identify transcripts cleaved under the guidance of microRNAs. Finally, a gene

regulation network for gall development was built by integrating all these -omics data.

Twelve of the 153 transcripts identified as targeted by microRNAs in tomato galls in

degradome analysis were considered the most robust candidates, based on their expression

profiles, which were inversely correlated with those of the corresponding microRNA. Some

of these 12 miRNA/mRNA pairs have already been reported to be involved in the plant-

nematode interaction in *Arabidopsis:* the miR408/UCCLACYANINE (blue copper protein)

and the miR398/copper superoxide dismutase (Noureddine *et al*. 2022), and the auxin-

regulated miR172/APETALA2 pair (Díaz-Manzano *et al*. 2018). Moreover, other pairs as

also appear to be interesting candidates based on the processes in which they are involved.

This is the case, for example for the miR164/NAC transcription factor involved in lateral root

development (Guo *et al*. 2005). In *Arabidopsis* root, the auxin-mediated induction of miR164

induces the silencing of *NO APICAL MERISTEM-1 (NAC1)* transcripts, thereby affecting

transmission of the auxin signal and regulating lateral root growth.

ARF8s are regulated at posttranscriptionally by miR167 in tomato roots

Among the 12 most robust miRNA/mRNA pairs identified as putatively involved in the

formation of galls, the *ARF8A* and *ARF8B* targets of miR167 were selected for further

functional analyses based on their role in auxin signaling, as auxins are a class of hormones

controlling root development and architecture (De Smet *et al.*, 2007; Quint & Gray, 2008;

Majda & Robert, 2018). ARF8 is an auxin-responsive factor (ARF). The transcription factors

of this family regulate the activation or repression of auxin-induced genes by binding to the

auxin response elements (AuxREs) in their promoters (reviewed in Guilfoyle and Hagen,

2007; Chandler, 2016; Li et al., 2016). The ARF gene family is conserved across the plant

kingdom and is well described in various plant species, including *A. thaliana* (23 genes)

(Hagen & Guilfoyle, 2002), *S. lycopersicum* (22 genes) (Zouine *et al.*, 2014), and *Oryza*

sativa (25 genes) (Wang *et al.*, 2007). The ARF family has been implicated in the regulation

of plant developmental processes, such as embryo morphogenesis (Rademacher *et al.*, 2011),

the formation of lateral roots in response to low levels of nitrogen (Gifford *et al.* 2008), the formation of adventitious roots (Lee *et al.*, 2019), leaf structure and senescence (Wilmoth *et al.*, 2005), flower development (Ellis *et al.*, 2005) and fruit initiation (Liu *et al.*, 2014). Like ARF5, 6, 7 and 19, ARF8 has been described as a transcriptional activator (reviewed in Guilfoyle and Hagen, 2007). In *Arabidopsis*, a partial redundancy between ARF8 and ARF6 has been reported, with both these activators silenced by miR167 (Reeves *et al.* 2012). The *Arabidopsis arf6arf8* double mutant has defective flower development, as the flower is entirely sterile (Nagpal, 2005), whereas the *arf8* mutant presents defects of pollination and fertilization (Tian *et al.*, 2004; Vernoux *et al.*, 2011). A role for ARF8 has been reported in the formation of lateral roots in *Arabidopsis* and soybean (Gifford *et al.*, 2008; Wang *et al.* 2015) and in the formation of adventitious roots (Gutierrez *et al.*, 2009). ARF6 and ARF8 were recently implicated in cambium establishment and maintenance (Ben-Targem *et al.*, 2021). The *arf6arf8* double mutant displays low levels of xylem occupancy and an absence of fiber accumulation until very late stages of plant growth.

In *A. thaliana* and tomato, *ARF8* genes are regulated posttranscriptionally by miR167 (Wu *et al.*, 2006; Liu *et al.*, 2014). Transcriptomic analyses of galls showed that *ARF8B* is overexpressed in tomato galls at 7 and 14 dpi, whereas *ARF8A* is overexpressed at 14 dpi. The infection of tomato lines expressing the GUS reporter gene under the control of the *ARF8A* or *ARF8B* promoter revealed high levels of activity for both *ARF8* promoters in giant cells and neighboring cells at 7 and 14 dpi, confirming the overexpression observed in the transcriptomic analyses. Gall degradome analysis identified *ARF8A* and *ARF8B* transcripts as cleaved by members of the miR167 family. Four tomato *MIR167* genes encode mature proteins with identical sequences and are downregulated in tomato galls at 7 and 14 dpi. This suggests that the *ARF8A* and *ARF8B* transcripts are cleaved by miR167 in uninfected tomato roots, as demonstrated in *A. thaliana* roots. We showed that RKN infection induces the inhibition of miR167 in galls, thereby decreasing the cleavage of *ARF8* transcripts by miR167, resulting in an overexpression of *ARF8A* and *ARF8B* in galls.

Auxin is a major factor regulating the formation of feeding cells in tomato
We used tomato lines with CRISPR deletions within the *ARF8A, ARF8B* and *ARF8AB* coding sequences to analyze the function of *ARF8* in plant-nematode interactions. The *arf8a*, *arf8b* and *arf8ab* lines displayed decreased susceptibility to nematode infection, with fewer gall and egg masses in mutants than in wild-type tomato plants. Moreover, the phenotyping of giant

feeding cells within cleared galls showed the giant cells from the three CRISPR lines to be
smaller than those from the wild type. These defects, associated with CRISPR mutations,
confirmed the involvement of *ARF8A* and *ARF8B* in the tomato response to RKN interaction
and that requirement for functional ARF8A and ARF8B for the correct development of
feeding cells.

MicroRNAs and transcription factors regulate *ARF8* expression, whereas auxin peaks
regulate ARF8 activity: when auxin levels are high, ARF8 (class II ARFs) activates the
transcription of auxin-responsive genes (Tiwari *et al.*, 2003). Auxin is known to be a major
factor regulating the formation of feeding cells in response to RKN signals (reviewed in
Gheysen and Mitchum, 2019; Oosterbeek *et al.*, 2021). Microarray analyses of *A. thaliana*
gall transcripts have revealed an early activation of genes responsible for auxin homeostasis
and auxin-responsive genes, and a downregulation of repressors of auxin responses (Hammes
et al., 2005; Jammes *et al.*, 2005; Barcala *et al.*, 2010). Studies of lines expressing reporter
genes under the control of the synthetic auxin-responsive DR5 promoter showed that this
promoter is activated in galls induced by RKN (Hutangura *et al.*, 1999; Karczmarek *et al.*,
2004; Absmanner *et al.*, 2013b). In *A. thaliana* galls, a strong signal was detected in both
giant cells and neighboring cells at 4 dpi for DR5:GUS lines (Cabrera *et al.*, 2014). This
increase in auxin levels in the galls may be controlled by either the plant or the nematode.
Auxin-mimicking compounds have been found in nematode secretions (De Meutter *et al.*,
2003, 2005).

ARF8 is involved in plant responses to biotic and abiotic stresses
ARF8 transcription factors have been implicated in plant responses to microorganisms. *ARF8*
is regulated in tomato leaves in response to biotic stresses, such as flagellin treatment or
infection with *Pseudomonas syringae* (Bouzroud *et al.*, 2018). A recent study in *Arabidopsis*
provided evidence for a suppression of miR167 expression, together with an induction of
ARF6 and *ARF8,* in response to infection with *P. syringae* pv. tomato DC3000 in *A. thaliana*
(Caruana *et al.*, 2020). The P35S:*MIR167* and *arf6 arf8* double mutants were found to be
more resistant to bacterial infection than the wild type. The authors suggested that ARF6 and
ARF8 modulate salicylic acid defenses response to *P. syringae* infection under miR167
regulation. Furthermore, soybean ARF8A and ARF8B have been shown to downregulate the
nodulation induced by miR167 (Wang et al., 2015). All these studies suggest that the auxin-
responsive pathway, including miR167/ARF8, is a key actor in the response to

microorganisms. Analysis of the transcriptomes of two tomato CRISPR *arf8* mutants showed

no induction of genes associated with plant defense in galls from *arf8* mutants. This finding

supports the notion that the lower levels of RKN infection observed in *arf8* mutants result

from a loss of susceptibility rather than an enhancement of plant defense.

ARF8, an environmental hub connecting development and stress responses

We have shown that *ARF8* genes are posttranscriptionally regulated by miR167 in galls, but

the transcriptional regulation of these genes has yet to be deciphered. *ARF8* was recently

shown to be regulated by a complex network of multiple activating and repressing

transcription factors in *A. thaliana* (Truskina *et al.*, 2021). Interestingly, some of these

transcription factors, such as WUSCHEL, Squamosa Promoter Binding Protein Like-13

(SPL13) and WRKY33, have also been implicated in plant development, and many *ARF8*

regulators are also associated with plant responses to biotic and abiotic stresses. Based on

these results, Truskina *et al*. suggested that ARF8 may act as an environmental hub

connecting development and stress responses mediating auxin responsiveness. The formation

of RKN-induced feeding sites interferes with plant developmental processes, including, in

particular, the development of lateral roots (Cabrera *et al.*, 2014, Olmo *et al.* 2020), which

suggests that the nematode may hijack this process. For example, the transcription factor

LBD16 and the microRNA miR390a, two key components of the auxin pathway and

transducers of lateral root development, are involved in gall formation in *Arabidopsis* and

tomato (Cabrera *et al.*, 2014, Olmo *et al*. 2020). *ARF8* has also been implicated in lateral root

formation in *Arabidopsis* (Gifford *et al*. 2008) and may, therefore, integrate biotic stress and

developmental processes during the formation of giant feeding cells. The common induction,

in tomato and *Arabidopsis* galls, of *ARF8* and of the transcription factor *LBD16* (Olmo *et al*.

2020) and miR390a (Diaz Manzano *et al*. 2016), suggests that there may be a conserved

auxin-mediated molecular pathway in galls. Early in the development of galls in *Arabidopsis*,

at 3 dpi, the silencing of *ARF3* by miR390 via tasiRNAs, and the induction of *ARF5* have

been shown to be required for parasitism (Cabrera *et al*. 2014a; Olmo *et al*. 2020). These

results suggest that there is a complex network involving ARFs and microRNAs responsible

for mediating auxin signaling during the development of galls induced by RKN.

Author contributions

Y.N., B.F and S.J.P. designed the study, performed the experimental work and wrote the manuscript. Y.N. and C.M. produced biological material for sequencing. M.dR. analyzed NGS data. M.Z. and JA designed, generated and characterized *arf8* CRISPR-Cas9 mutants. M.Q., J.M. and P.A. participated in the writing of the manuscript. All the authors analyzed and discussed the data.

Acknowledgments

The microscopy work was performed at the SPIBOC imaging facility of Institut Sophia Agrobiotech. We thank Dr Olivier Pierre and the entire team of the platform for assistance with microscopy. This work was funded by the INRAE SPE department and the French Government (National Research Agency, ANR) through the 'Investments for the Future' LabEx SIGNALIFE: program reference #ANR-11-LABX-0028-01 and IDEX UCAJedi ANR-15-IDEX-0, and by the French-Japanese bilateral collaboration programme PHC SAKURA 2019 #43006VJ. Y.N. was supported by a doctoral fellowship from Lebanon (Municipal Council of Aazzée).

References

Abad P, Williamson VM. **2010**. Plant Nematode Interaction: A Sophisticated Dialogue. *Advances in Botanical Research* **53**: 147–192.

Absmanner B, Stadler R, Hammes UZ. **2013a**. Phloem development in nematode-induced feeding sites: The implications of auxin and cytokinin. *Frontiers in Plant Science* **4**.

Absmanner B, Stadler R, Hammes UZ, Jammes F, Lecomte P, Almeida-Engler J De, Bitton F, Martin-Magniette M-L, Renou JP, Abad P, *et al.* 2013b. Parasitic nematodes modulate PIN-mediated auxin transport to facilitate infection (J Jones, G Gheysen, and C Fenoll, Eds.). *Mol Plant Microbe Interact* **26**: 107–109.

Addo-Quaye C, Miller W, Axtell MJ. **2009**. CleaveLand: A pipeline for using degradome data to find cleaved small RNA targets. *Bioinformatics* **25**: 130–131.

de Almeida Engler J, Gheysen G. **2013**. Nematode-induced endoreduplication in plant host cells: why and how? *Molecular Plant-Microbe Interactions* **26**: 17–24.

Anders S, Huber W. **2010**. Differential expression analysis for sequence count data. *Genome*

Biology **11**: R106.

Anders S, Pyl PT, Huber W. **2015**. HTSeq--a Python framework to work with high-throughput sequencing data. *Bioinformatics* **31**: 166–169.

Axtell MJ. **2013**. ShortStack: Comprehensive annotation and quantification of small RNA genes. *RNA* **19**: 740–751.

Bar-Or C, Kapulnik Y, Koltai H. **2005**. A broad characterization of the transcriptional profile of the compatible tomato response to the plant parasitic root knot nematode Meloidogyne javanica. *European Journal of Plant Pathology* **111**.

Barcala M, García A, Cabrera J, Casson S, Lindsey K, Favery B, García-Casado G, Solano R, Fenoll C, Escobar C. **2010**. Early transcriptomic events in microdissected Arabidopsis nematode-induced giant cells. *The Plant Journal* **61**: 698–712.

Ben-Targem M, Ripper D, Bayer M, Ragni L. **2021**. Auxin and gibberellin signaling cross-talk promotes hypocotyl xylem expansion and cambium homeostasis. *Journal of Experimental Botany* **72**: 3647–3660.

Blanc-Mathieu R, Perfus-Barbeoch L, Aury J-M, Da Rocha M, Gouzy J, Sallet E, Martin-Jimenez C, Bailly-Bechet M, Castagnone-Sereno P, Flot J-F, *et al.* **2017**. Hybridization and polyploidy enable genomic plasticity without sex in the most devastating plant-parasitic nematodes (T Gojobori, Ed.). *PLOS Genetics* **13**: e1006777.

Blok VC, Jones JT, Phillips MS, Trudgill DL. **2008**. Parasitism genes and host range disparities in biotrophic nematodes : the conundrum of polyphagy versus specialisation. *BioEssays* **30**: 249–259.

Bouzroud S, Gouiaa S, Hu N, Bernadac A, Mila I, Bendaou N, Smouni A, Bouzayen M, Zouine M. **2018**. Auxin Response Factors (ARFs) are potential mediators of auxin action in tomato response to biotic and abiotic stress (Solanum lycopersicum) (K Wu, Ed.). *PLOS ONE* **13**: e0193517.

Cabrera J, Barcala M, García A, Rio-Machín A, Medina C, Jaubert-Possamai S, Favery B, Maizel A, Ruiz-Ferrer V, Fenoll C, *et al.* **2016**. Differentially expressed small RNAs in Arabidopsis galls formed by Meloidogyne javanica : a functional role for miR390 and its TAS3-derived tasiRNAs. *New Phytologist* **209**: 1625–1640.

Cabrera J, Díaz-Manzano FE, Sanchez M, Rosso MN, Melillo T, Goh T, Fukaki H,

Cabello S, Hofmann J, Fenoll C, *et al*. 2014. A role for LATERAL ORGAN BOUNDARIES-DOMAIN 16 during the interaction Arabidopsis-Meloidogyne spp. provides a molecular link between lateral root and root-knot nematode feeding site development. *New Phytologist* **203**: 632–645.

Cabrera J, Olmo R, Ruiz-Ferrer V, Abreu I, Hermans C, Martinez-Argudo I, Fenoll C, Escobar C. **2018**. A Phenotyping Method of Giant Cells from Root-Knot Nematode Feeding Sites by Confocal Microscopy Highlights a Role for CHITINASE-LIKE 1 in Arabidopsis. *International Journal of Molecular Sciences* **19**: 429.

Cai C, Li C, Sun R, Zhang B, Nichols RL, Hake KD, Pan X. **2021**. Small RNA and degradome deep sequencing reveals important roles of microRNAs in cotton (Gossypium hirsutum L.) response to root-knot nematode Meloidogyne incognita infection. *Genomics* **113**: 1146–1156.

Caillaud M-C, Dubreuil G, Quentin M, Perfus-Barbeoch L, Lecomte P, de Almeida Engler J, Abad P, Rosso M-N, Favery B. **2008**. Root-knot nematodes manipulate plant cell functions during a compatible interaction. *Journal of Plant Physiology* **165**: 104–113.

Caillaud M-C, Favery B. **2016**. In Vivo Imaging of Microtubule Organization in Dividing Giant Cell. *Methods in molecular biology (Clifton, N.J.)* **1370**: 137–44.

Caruana JC, Dhar N, Raina R. **2020**. Overexpression of Arabidopsis microRNA167 induces salicylic acid□dependent defense against Pseudomonas syringae through the regulation of its targets ARF6 and ARF8. *Plant Direct* **4**.

Chandler JW. **2016**. Auxin response factors. *Plant Cell and Environment* **39**: 1014–1028.

Damiani I, Baldacci-Cresp F, Hopkins J, Andrio E, Balzergue S, Lecomte P, Puppo A, Abad P, Favery B, Hérouart D. **2012**. Plant genes involved in harbouring symbiotic rhizobia or pathogenic nematodes. *New Phytologist* **194**: 511–522.

De Meutter J, Tytgat T, Prinsen E, Gheysen G, Van Onckelen H, Gheysen G. **2005**. Production of auxin and related compounds by the plant parasitic nematodes Heterodera schachtii and Meloidogyne incognita. *Communications in agricultural and applied biological sciences* **70**: 51–60.

De Meutter J, Tytgat T, Witters E, Gheysen G, Van Onckelen H, Gheysen G. **2003**. Identification of cytokinins produced by the plant parasitic nematodes Heterodera schachtii

and Meloidogyne incognita. *Molecular Plant Pathology* **4**: 271–277.

Díaz-Manzano FE, Cabrera J, Ripoll J-JJ, Del Olmo I, Andrés MF, Silva AC, Barcala
M, Sánchez M, Ruíz-Ferrer V, de Almeida-Engler J, *et al.* 2018. A role for the gene
regulatory module microRNA172/TARGET OF EARLY ACTIVATION TAGGED
1/FLOWERING LOCUS T (miRNA172/TOE1/FT) in the feeding sites induced by
Meloidogyne javanica in Arabidopsis thaliana. *New Phytologist* **217**: 813–827.

Díaz-Manzano FE, Barcala M, Engler G, Fenoll C, de Almeida-Engler J, Escobar C.
2016. A Reliable Protocol for In situ microRNAs Detection in Feeding Sites Induced by
Root-Knot Nematodes. *Front Plant Sci.* 7: 966.

Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, Batut P, Chaisson M,
Gingeras TR. **2013**. STAR: ultrafast universal RNA-seq aligner. *Bioinformatics*: 1–7.

Ellis CM, Nagpal P, Young JC, Hagen G, Guilfoyle TJ, Reed JW. **2005**. AUXIN
RESPONSE FACTOR1 and AUXIN RESPONSE FACTOR2 regulate senescence and floral
organ abscission in Arabidopsis thaliana. *Development* **132**: 4563–4574.

Favery B, Dubreuil G, Chen MS, Giron D and Abad P. 2020. Gall-Inducing Parasites :
convergent and conserved strategies of plant manipulation by insect and nematodes. *Annu.*
Rev. Phytopathol. **58**(1) :1-22.

Favery B, Quentin M, Jaubert-Possamai S, Abad P. **2016**. Gall-forming root-knot
nematodes hijack key plant cellular functions to induce multinucleate and hypertrophied
feeding cells. *Journal of insect physiology* **84**: 60–69.

Fuller VL, Lilley CJ, Atkinson HJ, Urwin PE. **2007**. Differential gene expression in
Arabidopsis following infection by plant-parasitic nematodes Meloidogyne incognita and
Heterodera schachtii. *Molecular Plant Pathology* **8**: 595–609.

German MA, Luo S, Schroth G, Meyers BC, Green PJ. **2009**. Construction of parallel
analysis of rna ends (Pare) libraries for the study of cleaved mirna targets and the rna
degradome. *Nature Protocols* **4**: 356–362.

Gheysen G, Mitchum MG. **2019**. Phytoparasitic Nematode Control of Plant Hormone
Pathways. *Plant Physiology* **179**: 1212–1226.

Gifford ML, Dean A, Gutierrez RA, Coruzzi GM, Birnbaum KD. **2008**. Cell-specific
nitrogen responses mediate developmental plasticity. *Proceedings of the National Academy of*

Sciences of the United States of America **105**: 803–808.

Guilfoyle TJ, Hagen G. **2007**. Auxin response factors. *Current Opinion in Plant Biology* **10**: 453–460.

Gutierrez L, Bussell JD, Pacurar DI, Schwambach J, Pacurar M, Bellini C. **2009**. Phenotypic Plasticity of Adventitious Rooting in Arabidopsis Is Controlled by Complex Regulation of AUXIN RESPONSE FACTOR Transcripts and MicroRNA Abundance. *The Plant Cell* **21**: 3119–3132.

Hagen G, Guilfoyle T. **2002**. Auxin-responsive gene expression: Genes, promoters and regulatory factors. *Plant Molecular Biology* **49**: 373–385.

Hammes UZ, Schachtman DP, Berg RH, Nielsen E, Koch W, McIntyre LM, Taylor CG. **2005**. Nematode-induced changes of transporter gene expression in Arabidopsis roots. *Molecular Plant-Microbe Interactions* **18**: 1247–1257.

Hao Y, Hu G, Breitel D, Liu M, Mila I, Frasse P, Fu Y, Aharoni A, Bouzayen M, Zouine M. **2015**. Auxin Response Factor SlARF2 Is an Essential Component of the Regulatory Mechanism Controlling Fruit Ripening in Tomato. *PLoS Genetics* **11**.

Huang S-Y, Zhao G-H, Fu B-Q, Xu M-J, Wang C-R, Wu S-M, Zou F-C, Zhu X-Q. **2012**. Genomics and molecular genetics of Clonorchis sinensis: current status and perspectives. (J Jones, G Gheysen, and C Fenoll, Eds.). *Parasitology international* **61**: 71–6.

Hutangura P, Mathesius U, Jones MGK, Rolfe BG. **1999**. Auxin induction is a trigger for root gall formation caused by root-knot nematodes in white clover and is associated with the activation of the flavonoid pathway. *Functional Plant Biology* **26**: 221.

Ibrahim HMMM, Hosseini P, Alkharouf NW, Hussein EHA a, Gamal El-Din AEKY, Aly MAMM, Matthews BF. **2011**. Analysis of Gene expression in soybean (Glycine max) roots in response to the root knot nematode Meloidogyne incognita using microarrays and KEGG pathways. *BMC Genomics* **12**: 220.

Jammes F, Lecomte P, Almeida-Engler J, Bitton F, Martin-Magniette M-LL, Renou JP, Abad P, Favery B, De Almeida-Engler J, Bitton F, *et al.* 2005. Genome-wide expression profiling of the host response to root-knot nematode infection in Arabidopsisa. *The Plant Journal* **44**: 447–458.

Jaubert-Possamai S, Noureddine Y, Favery B. **2019**. MicroRNAs, New Players in the

Plant–Nematode Interaction. *Frontiers in Plant Science* **10**: 1–8.

Karczmarek A, Overmars H, Helder J, Goverse A. **2004**. Feeding cell development by
cyst and root-knot nematodes involves a similar early, local and transient activation of a
specific auxin-inducible promoter element. *Molecular Plant Pathology* **5**: 343–346.

Kaur P, Shukla N, Joshi G, VijayaKumar C, Jagannath A, Agarwal M, Goel S, Kumar
A. **2017**. Genome-wide identification and characterization of miRNAome from tomato
(Solanum lycopersicum) roots and root-knot nematode (Meloidogyne incognita) during
susceptible interaction (Q Zou, Ed.). *PLOS ONE* **12**: e0175178.

Kozomara A, Birgaoanu M, Griffiths-Jones S. **2019**. MiRBase: From microRNA
sequences to function. *Nucleic Acids Research* **47**: D155–D162.

Lee HW, Cho C, Pandey SK, Park Y, Kim M-J, Kim J. **2019**. LBD16 and LBD18 acting
downstream of ARF7 and ARF19 are involved in adventitious root formation in Arabidopsis.
BMC Plant Biology **19**: 46.

Li SB, Xie ZZ, Hu CG, Zhang JZ. **2016**. A review of auxin response factors (ARFs) in
plants. *Frontiers in Plant Science* **7**.

Liu N, Wu S, Houten J Van, Wang Y, Ding B, Fei Z, Clarke TH, Reed JW, Van Der
Knaap E. **2014**. Down-regulation of AUXIN RESPONSE FACTORS 6 and 8 by microRNA
167 leads to floral development defects and female sterility in tomato. *Journal of*
Experimental Botany **65**: 2507–2520.

Majda M, Robert S. **2018**. The Role of Auxin in Cell Wall Expansion. *International Journal*
of Molecular Sciences **19**: 951.

Marella HH, Nielsen E, Schachtman DP, Taylor CG. **2013**. The amino acid permeases
AAP3 and AAP6 are involved in root-knot nematode parasitism of arabidopsis. *Molecular*
Plant-Microbe Interactions **26**: 44–54.

Medina C, Rocha M, Magliano M, Ratpopoulo A, Revel B, Marteu N, Magnone V,
Lebrigand K, Cabrera J, Barcala M, *et al.* **2017**. Characterization of microRNAs from
Arabidopsis galls highlights a role for miR159 in the plant response to the root knot
nematode Meloidogyne incognita. *New Phytologist* **216**: 882–896.

Mi H, Muruganujan A, Elbert D, Huang X and Thomas PD. 2019. PANTHER version
14: more genomes, a new PANHER GO-slim and improvements in enrichment analysis tools.

Nucleic Acid Res. **47**: D419-D426.

Mejias J, Bazin J, Truong NM, Chen Y, Marteu N, Bouteiller N, Sawa S, Crespi MD,

Vaucheret H, Abad P, *et al.* **2021**. The root-knot nematode effector MiEFF18 interacts with

the plant core spliceosomal protein SmD1 required for giant cell formation. *New Phytologist*

229: 3408–3423.

Mejias J, Chen Y, Bazin J, Truong N-M, Mulet K, Noureddine Y, Jaubert-Possamai S,

Ranty-Roby S, Soulé S, Abad P, *et al.* **2022**. Silencing the conserved small nuclear

ribonucleoprotein SmD1 target gene alters susceptibility to root-knot nematodes in plants.

Plant Physiology **189**: 1741–1756.

Nagpal P. **2005**. Auxin response factors ARF6 and ARF8 promote jasmonic acid production

and flower maturation. *Development* **132**: 4107–4118.

Nguyen CN, Perfus-Barbeoch L, Quentin M, Zhao J, Magliano M, Marteu N, da Rocha

M, Nottet N, Abad P and Favery B. 2018. A root-knot nematode small glycine and

cysteine-rich effector, MiSGCR1, is involved in plant parasitism. *New Phytol.* **217**(2): 687-

699.

Nikovics K, Blein T, Peaucelle A, Ishida T, Morin H, Aida M, Laufs P. **2006**. The

Balance between the MIR164A and CUC2 Genes Controls Leaf Margin Serration in

Arabidopsis. *The Plant Cell* **18**: 2929–2945.

Noureddine Y, Mejias J, da Rocha M, Thomine S, Quentin M, Abad P, Favery B,

Jaubert-Possamai S. 2022. Copper microRNAs modulate the formation of giant feeding

cells induced by the root knot nematode Meloidogyne incognita in Arabidopsis thaliana.

New Phytol. **8**. In press.

Oosterbeek M, Lozano-Torres JL, Bakker J, Goverse A. **2021**. *Sedentary Plant-Parasitic*

Nematodes Alter Auxin Homeostasis via Multiple Strategies.

Paicu C, Mohorianu I, Stocks M, Xu P, Coince A, Billmeier M, Dalmay T, Moulton V,

Moxon S. **2017**. MiRCat2: Accurate prediction of plant and animal microRNAs from next-

generation sequencing datasets. *Bioinformatics* **33**: 2446–2454.

Palatnik JF, Wollmann H, Schommer C, Schwab R, Boisbouvier J, Rodriguez R,

Warthmann N, Allen E, Dezulian T, Huson D, *et al.* **2007**. Sequence and Expression

Differences Underlie Functional Specialization of Arabidopsis MicroRNAs miR159 and

miR319. *Developmental Cell* **13**: 115–125.

Pan X, Nichols RL, Li C, Zhang B. **2019**. MicroRNA-target gene responses to root knot
nematode (Meloidogyne incognita) infection in cotton (Gossypium hirsutum L.). *Genomics*
111: 383–390.

Portillo M, Cabrera J, Lindsey K, Topping J, Andr MF, Emiliozzi M, Oliveros JC, Garc
G, Solano R, Koltai H, *et al*. **2013**. Distinct and conserved transcriptomic changes during
nematode-induced giant cell development in tomato compared with Arabidopsis: A functional
role for gene repression. *New Phytologist* **197**: 1276–1290.

Quint M, Gray WM. **2008**. Auxin signaling Marcel. *Curr. Opin. Plant Biol.* **9**: 448–453.

Rademacher EH, Möller B, Lokerse AS, Llavata-Peris CI, Van Den Berg W, Weijers D.
2011. A cellular expression map of the Arabidopsis AUXIN RESPONSE FACTOR gene
family. *Plant Journal* **68**: 597–606.

Rancurel C, van Tran T, Elie C, Hilliou F. **2019**. SATQPCR: Website for statistical
analysis of real-time quantitative PCR data. *Molecular and Cellular Probes* **46**: 101418.

Reeves PH, Ellis CM, Ploense SE, Wu MF, Yadav V, Tholl D, Chételat A, Haupt I,
Kennerley BJ, Hodgens C, *et al*. **2012**. A regulatory network for coordinated flower
maturation. *PLoS Genetics* **8**.

Robinson MD, McCarthy DJ, Smyth GK. **2009**. edgeR: A Bioconductor package for
differential expression analysis of digital gene expression data. *Bioinformatics* **26**: 139–140.

Shukla N, Yadav R, Kaur P, Rasmussen S, Goel S, Agarwal M, Jagannath A, Gupta R,
Kumar A. **2018**. Transcriptome analysis of root-knot nematode (Meloidogyne incognita)-
infected tomato (Solanum lycopersicum) roots reveals complex gene expression profiles and
metabolic networks of both host and nematode during susceptible and resistance responses.
Molecular Plant Pathology **19**: 615–633.

De Smet I, Tetsumura T, De Rybel B, Frey NF dit, Laplaze L, Casimiro I, Swarup R,
Naudts M, Vanneste S, Audenaert D, *et al*. **2007**. Auxin-dependent regulation of lateral
root positioning in the basal meristem of Arabidopsis. *Development* **134**: 681–690.

Tian CE, Muto H, Higuchi K, Matamura T, Tatematsu K, Koshiba T, Yamamoto KT.
2004. Disruption and overexpression of auxin response factor 8 gene of Arabidopsis affect
hypocotyl elongation and root growth habit, indicating its possible involvement in auxin

homeostasis in light condition. *Plant Journal* **40**: 333–343.

Tiwari SB, Hagen G, Guilfoyle T. **2003**. The roles of auxin response factor domains in

auxin-responsive transcription. *Plant Cell* **15**: 533–543.

Truskina J, Han J, Chrysanthou E, Galvan-Ampudia CS, Lainé S, Brunoud G, Macé J,

Bellows S, Legrand J, Bågman AM, *et al*. 2021. A network of transcriptional repressors

modulates auxin responses. *Nature* **589**: 116–119.

Van Ha C, Le DT, Nishiyama R, Watanabe Y, Sulieman S, Tran UT, Mochida K, Van

Dong N, Yamaguchi-Shinozaki K, Shinozaki K, *et al*. 2013. The auxin response factor

transcription factor family in soybean: Genome-wide identification and expression analyses

during development and water stress. *DNA Research* **20**: 511–524.

Vernoux T, Brunoud G, Farcot E, Morin V, Van den Daele H, Legrand J, Oliva M, Das

P, Larrieu A, Wells D, *et al*. 2011. The auxin signalling network translates dynamic input

into robust patterning at the shoot apex. *Molecular Systems Biology* **7**: 508.

Verstraeten B, Atighi MR, Ruiz-Ferrer V, Escobar C, De Meyer T, Kyndt T. **2021**. Non-

coding RNAs in the interaction between rice and Meloidogyne graminicola. *BMC Genomics*

22: 560.

Wang Y, Li P, Cao X, Wang X, Zhang A, Li X. **2009**. Identification and expression

analysis of miRNAs from nitrogen-fixing soybean nodules. *Biochemical and Biophysical*

Research Communications **378**: 799–803.

Wang Y, Li K, Chen L, Zou Y, Liu H, Tian Y, Li D, Wang R, Zhao F, Ferguson BJ, *et*

***al*. 2015**. MicroRNA167-Directed Regulation of the Auxin Response Factors *GmARF8a*and

*GmARF8b*Is Required for Soybean Nodulation and Lateral Root Development. *Plant*

Physiology **168**: 984–999.

Wang D, Pei K, Fu Y, Sun Z, Li S, Liu H, Tang K, Han B, Tao Y. **2007**. Genome-wide

analysis of the auxin response factors (ARF) gene family in rice (Oryza sativa). *Gene* **394**:

13–24.

Weijers D, Wagner D. **2016**. Transcriptional Responses to the Auxin Hormone. *Annual*

Review of Plant Biology **67**: 539–574.

Wilmoth JC, Wang S, Tiwari SB, Joshi AD, Hagen G, Guilfoyle TJ, Alonso JM, Ecker

JR, Reed JW. **2005**. NPH4/ARF7 and ARF19 promote leaf expansion and auxin-induced

lateral root formation. *Plant Journal* **43**: 118–130.

Wu G, Park MY, Conway SR, Wang JW, Weigel D, Poethig RS. **2009**. The Sequential

Action of miR156 and miR172 Regulates Developmental Timing in Arabidopsis. *Cell* **138**:

750–759.

Wu M-F, Tian Q, Reed JW. **2006**. Arabidopsis microRNA167 controls patterns of ARF6

and ARF8 expression, and regulates both female and male reproduction. *Development* **133**:

4211–4218.

Yamaguchi YL, Suzuki R, Cabrera J, Nakagami S, Sagara T, Ejima C, Sano R, Aoki Y,

Olmo R, Kurata T, *et al.* **2017**. Root-Knot and Cyst Nematodes Activate Procambium-

Associated Genes in Arabidopsis Roots. *Frontiers in Plant Science* **8**: 1–13.

Yang X, Li L. **2011**. miRDeep-P: A computational tool for analyzing the microRNA

transcriptome in plants. *Bioinformatics* **27**: 2614–2615.

Zhao W, Li Z, Fan J, Hu C, Yang R, Qi X, Chen H, Zhao F, Wang S. **2015**. Identification

of jasmonic acid-associated microRNAs and characterization of the regulatory roles of the

miR319/TCP4 module under root-knot nematode stress in tomato. *Journal of Experimental*

Botany **66**: 4653–4667.

Zouine M, Fu Y, Chateigner-Boutin A-L, Mila I, Frasse P, Wang H, Audran C, Roustan

J-P, Bouzayen M. **2014**. Characterization of the Tomato ARF Gene Family Uncovers a

Multi-Levels Post-Transcriptional Regulation Including Alternative Splicing (S Maas, Ed.).

PLoS ONE **9**: e84203.

miR	miRNA family	padj 14dpi	logFC 14dpi	padj 7 dpi	logFC 7dpi	Degradome Category
SL3...	miR164	1.03541e-06	-2.05	0.27	-0.45	0
SL3...	miR164	0.27	-0.63	0.04	-0.54	0
SL3...	miR167	0.001	0.84	0.18	0.32	0
SL3...	miR167	0.001	0.9	0.0001	0.82	0
SL3...	miR397	0.005	-1.26	0.07	-0.32	0
SL3...	miR398	0.0001	-1.51	0.04	-0.83	0
SL3...	miR403	0.001	1.16	0.18	0.46	0
SL3...	miR408	0.16	-0.692	0.03	-0.79	0
SL3...	miR7981			0.002	1.1	0
SL3...	miR7981	0.004	2.09	0.07	0.64	0
SL3...	miR172	0.0001	1.26			0
SL3...	miR172	4.45e-05	1.53	0.04	0.99	0

Supporting Information

Fig. S1. Quantification of tomato transcripts by RT-qPCR in galls relative to uninfected

tomato roots, at 7 and 14 days post infection

Fig. S2. Expression of the Sly-miR167 family in galls and uninfected tomato roots.

Fig S3. Root phenotype of *crispr arf8* mutants

Table S1. Primers used for RT-qPCR.

Table S2. Protein-coding genes DE in galls at 7 dpi

Table S3. Protein-coding genes DE in galls at 14 dpi

Table S4. Quantification of tomato transcripts in galls by RT-qPCR (G) relative to uninfected tomato roots (R), at 7 and 14 days post inoculation (dpi).

Table S5. Gene ontology (GO) analysis of the genes DE in galls at 7 and/or 14 dpi.

Table S6. The number of raw reads from the three libraries obtained from galls and uninfected roots of *S. lycopersicum* at 7 dpi and 14 dpi.

Table S7. *De novo* prediction of *Solanum lycopersicum MIR* genes.

Table S8. MicroRNAs differentially expressed in galls at 7 and/or 14 dpi.

Table S9. 153 transcripts targeted by miRNAs in galls at 7 and/or 14 dpi identified by degradome sequencing and CleaveLand analysiss.

Table S10. Infection assays in *arf8 CRISPR* lines infected with *M. incognita* and comparison to WT.

Table S11. Genes differentially expressed between *crisprarf8* mutant galls and wild-type (WT) galls.

Figure legends

Figures and Tables

Table 1. The 12 miRNA/mRNA pairs with inversely regulated expression profiles. Degradome analysis identified 12 genes targeted by miRNAs in galls. *Solanum lycopersicum* (Solyc) gene expression levels were compared between galls and uninfected roots, by two statistical methods (DSeq2 and EdgeR), and the expression of mature miRNAs was compared by DSeq2. Gall/root fold-change differences in expression (LogFC) at 7 and 14 days post infection (dpi) and the adjusted *p*-value obtained by the Benjamini-Hochberg method (adj *p*-value) are indicated for genes and miRNAs. The genes and mature microRNAs upregulated in galls are shown in red, and those downregulated in galls are shown in green.

Figure 1. Tomato protein-coding genes and miRNAs differentially expressed in *M. incognita*-induced galls relative to the corresponding uninfected roots, at 7 and/or 14 dpi. The numbers of genes differentially expressed at 7 days post inoculation (dpi) (pink)

and/or 14 dpi (yellow) between galls and the corresponding uninfected roots are indicated in

Venn diagrams.

Figure 2. *ARF8A* and *ARF8B* are strongly transcribed in tomato galls induced by *M.*

***incognita*.** The activity of the *ARF8A* and *ARF8B* promoters (*pARF8A* and *pARF8B*) was

studied in galls induced by *M. incognita* in *S. lycopersicum* lines expressing the

pARF8A::GUS or the *pARF8B::GUS* construct, at 7 and 14 days post inoculation (dpi). Blue

staining indicating GUS activity under the control of *pARF8A* was observed in (a) uninfected

root tips and (b-c) galls induced by *M. incognita* at 7 dpi (b) and 14 dpi (c). GUS activity

under the control of *pARF8B* was observed in (d) uninfected root tips and in (e-f) galls at 7

dpi (e) and 14 dpi (f). Bars: 500 µm.

Figure 3. *ARF8A* and *ARF8B* are strongly transcribed in 14 dpi nematode-induced

feeding sites. The activity of the *ARF8A* and *ARF8B* promoters (*pARF8A* and *pARF8B*) was

studied in galls induced by *M. incognita* in *S. lycopersicum* expressing the *pARF8A::GUS* (a-

b) or the *pARF8B::GUS* (c-d) construct, at 7 and 14 days post inoculation (dpi). Gall sections

(5 µm) were cut after GUS staining and observed by dark-field microscopy. Red staining,

reflecting GUS activity, was observed in giant cells and neighboring cells in *pARF8A::GUS*

galls (a) 7 dpi and (b) 14 dpi. Strong GUS activity was observed in giant cells and

neighboring cells in the galls of *pARF8B::GUS* plants at (c) 7 dpi and (d) 14 dpi. *, giant

cells; nc: neighboring cells; Bars: (a,b) 100 µm (c,d) 50 µm.

Figure 4. Tomato *slarf8-KO* lines. Generation of slarf8-KO mutant lines by CRISPR/Cas9.

Guide RNAs (sgRNA, red bar) anchored next to the Zinc Finger Motif (ZFM) were designed

for CRISPR/Cas9 strategy. Protospacer adjacent motif (PAM) are indicated in blue. Mutations

within *slarf8* coding sequences corresponding to nucleotide deletions are shown in green.

Three types of mutants predicted to produce heavily truncated proteins were chosen for

further phenotypic characterization. These mutants are annotated as *arf8a-cr* (*arf8a* single

mutant), *arf8a-cr* (*arf8b* single mutant), *arf8ab-cr* (*arf8a* and *arf8b* double mutant). The

predicted mutated proteins are schematically illustrated (right panel).

Figure 5. The single mutants *arf8a*[CR-2] and *arf8b*[CR-11] and the double mutant *arf8ab*[CR-2,4]
were significantly less susceptible to *M. incognita* than the wild type. (a), The
susceptibility of the single and double CRISPR-Cas9 mutant lines and wild-type (WT)
MicroTom plants to *M. incognita* was evaluated by counting the numbers of galls and egg
masses per plant (G/plant and EM/plant, respectively) in two independent infection assays in
soil. (b) The effect of deleting *ARF8A* and/or *ARF8B* on the development of giant feeding
cells was further evaluated by measuring the size of the feeding site produced in each mutant
line and comparing it to that in the WT. Galls were collected 21 days post infection (dpi) *in*
vitro to measure the area (mm^2) covered by the giant cells by the BABB clearing method
(Cabrera et al., 2018). The impact of plant genotype was assessed in Mann-Whitney tests. *,
$P < 0.05$. Boxes indicate the interquartile range (25th to 75th percentile). The central lines
within the boxes represent the medians. Whiskers indicate the minimum and maximum usual
values present in the dataset. The circle outside the box represents an outlier. n, the number of
plants analyzed in each assay. Bars 50 µm.

(a)

Test1 μtom n=11 ; *arf8a* n=9; *arf8b* n=10; *arf8ab* n=7

Test2 μtom n=16 ; *arf8a* n=13; *arf8b* n=16; *arf8ab* n=14

Test3 μtom n=8 ; *arf8a* n=17; *arf8b* n=11; *arf8ab* n=6

Galls/plant   Egg Masses/plant

(b)

Test 1 : μtom n=X ; *arf8a* n=X; *arf8b* n=X; *arf8ab* n=X
Test 2 : μtom n=14 ; *arf8a* n=16; *arf8b* n=8; *arf8ab* n=10

Nematode Feeding site area (mm2)

Test1   Test2

(c)