Proton and alpha radiation-induced mutational profiles in human cells

Tiffany M. Delhomme[1], Manuela Buonanno[2], Veljko Grilj[2], Josep Biayna[1], Fran Supek[1,3]*

[1] Genome Data Science, Institute for Research in Biomedicine (IRB Barcelona), Barcelona, Spain

[2] Radiological Research Accelerator Facility (RARAF), Columbia University, New York, USA.

[3] Catalan Institution for Research and Advanced Studies (ICREA), Barcelona, Spain.

*correspondence to: fran.supek@irbbarcelona.org

Abstract

Ionizing radiation (IR) is known to be DNA damaging and mutagenic, however less is known about which mutational footprints result from exposures of human cells to different types of IR. We were interested in the mutagenic effects of particle radiation exposures on genomes of various human cell types, in order to gauge the genotoxic risks of space travel, and of certain types of tumor radiotherapy. To this end, we exposed cultured cell lines from the blood, breast and lung to intermittent proton and alpha particle (helium nuclei) beams at doses sufficient to affect cell survival. Whole-genome sequencing revealed that mutation rates were not overall markedly increased upon proton and alpha exposures. However, there were changes in mutation spectra and distributions, such as the increases in clustered mutations and of certain types of indels and structural variants. The spectrum of mutagenic effects of particle beams may often be cell-type and/or genetic background specific. Overall, the mutational effects of recurrent exposures to proton and alpha radiation on human cells appear subtle, however further work is warranted to understand effects of chronic, long-term exposures on various human tissues.

Introduction

Humans have a keen interest in travelling in space, from deep space exploration, to extraterrestrial colonization. In addition, the space environment was also considered for biomedical research purposes, for example the "Tumors in space" experiment aims to study the effects of microgravity and exposure to space radiation on tumor organoids, with implications to cancer risk and treatment[1]. Spaceflight, i.e. flight beyond the Earth's atmosphere[2], is usually defined as crossing the Kármán separation line i.e. 80-100 kilometres altitude above the sea level[2–4]. Above this boundary, the environment dramatically changes, notably in terms of microgravity, temperature and radiation[5]. Studies revealed the pathological consequences on human physiology from travelling in space and reported cardiovascular, musculoskeletal and immune changes[6]. However, a better understanding of space travel impact on the human body at the cellular and sub-cellular levels is needed, for instance how the genome integrity is affected. Important effects of space travel are exerted by the various types of radiation to which tissues are exposed outside the Earth's atmosphere and magnetosphere. Those include the solar radiation, the radioactive environment of the planet, and galactic cosmic radiation (or rays;

GCR).

GCR originates mainly from the Sun or outside of the Solar system but within the Milky
Way galaxy (there exist also the extra-galactic cosmic radiations, which are less relevant as a
space travel hazard due to their low flux and extremely high energies). The GCR encompasses
photon-based radiation as well as particle radiations, including neutrons and charged particles;
among the latter, protons and helium nuclei are the most abundant. Cosmic radiation has
emerged as an issue of concern, since exposure to radiation may increase cancer risk in
astronauts[7,8]. A better understanding of the mutagenic mechanisms of various cosmic radiation
types upon human DNA would be helpful to understand the basis of the increased risk of cancer
or of reproductive harm resulting from space travel.

Photons are the better-studied component of the GCR, since they consist of gamma
rays and X-rays, whose biological impact was studied in other contexts apart from GCR. For
instance, gamma rays were released during the Chernobyl nuclear power plant disaster,
prompting studies of how this type of radiation impacts human genome stability. The genomic
profiles of thyroid carcinomas in irradiated children post-Chernobyl contained similar driver
mutations and gene fusions as in non-radiation-associated thyroid tumors[9]. However there was
a radiation dose-dependant increase in the fusion driver events, as well as a general radiation-
associated increase in certain types of structural variants (rearrangements)[10]. Consistent with a
mutational impact that can generate somatic driver mutations, ionizing radiation-generating
incidents (e.g. the Chernobyl disaster, and also the Hiroshima and Nagasaki A-bomb attacks)
also increased the risk of developing cancer, proportionally to the dose of exposure[11,12].

In addition to nuclear incidents, radiation exposures can result from medical uses.
Commonly this involves X-rays, used for both diagnostic purposes, and also in applications to
tumor radiotherapy. The impact of therapeutic X-ray exposure on human DNA was examined
via genomic analyses of radiation-associated second malignancies and of post-treatment
metastatic tumors, reporting an increase of small and large deletions, and also certain
mutational signatures[13,14]. For instance, this entails a particular signature of indels (the PCAWG
Signature ID8) characterized by larger deletions ($\geq$ 5-bp) without flanking micro-homology; this
was previously linked to double-strand break repair[15]. Consistently with their mutagenic effect,
medical use of X-rays has been reported to increase cancer risk[16] .

Regardless of the source of the radiation – either the cosmos, nuclear accidents or
medical uses – it is not known if ionizing radiation consisting of charged-particle radiations, such
as protons or alpha rays, would have similar mutagenic effects to more commonly studied
photons (X-rays and gamma-rays). Generally, all types of ionizing radiations, including gamma-
rays, X-rays, energetic charged particles (protons, alpha particles [helium nuclei] and heavier
ions) and neutrons have been classified as human carcinogens by the World Health
Organization (WHO)[17]. This suggests mutagenic impacts of various radiation types, however
research on the specific mutational footprints of charged particles on the integrity of the human
genome has been limited.

These are of interest because during transit beyond low Earth orbit, every cell nucleus within an astronaut's body is traversed, on average, by an energetic proton every few days, and by a helium nucleus once every few weeks. Mutational effect of proton radiation is of interest also for reasons other than space travel. This type of radiation is increasingly adopted in the clinic to treat cancer[18] , and appears promising for toxicity profiles[19–22]. Proton therapy may reduce the health risks (compared to X-rays) of secondary malignancies[23], and of severe radiation-induced lymphopenia[24,25]. The above provides motivation for studying the mutagenic effects of proton, and of helium nuclei (alpha) radiation exposures, which have the potential to result in long term health effects in exposed individuals.

The activity of mutational processes in DNA, which may arise from both endogenous factors such as DNA repair deficiencies and exogenous factors such as tobacco smoking, can be captured by mutational signatures[26], mathematical constructs that describe the differential frequencies of mutation types across many genomes. Such signatures are commonly based on trinucleotide spectra of single nucleotide variants (SNVs), however they can also be based on indels, structural variation (rearrangements), or mutation clusters[15,27,28]. SNV and indel mutational signatures in human cancer are well characterized and organized in comprehensive catalogues[15,29], including also signatures in tumors pre-treated with ionizing radiation[13,14,30] (here referring to photons).

In addition, some genomic studies have reported mutational signatures on experimental models, including experimentally-induced mutational signatures of (non-ionizing) UV radiation in human cell lines[31], and also in cultured cells from healthy human tissues[32,33]. Additionally, signatures of ionizing radiations were reported in exposed mice[34,35] and in the worm *C. elegans*[36]; both reported SNV signatures enriched in C>T transitions[15].

In order to increase our understanding of the potential impact of particle radiation encountered during space travel on human tissues, it is important to systematically, in a genome-wide manner examine the DNA mutations arising from the exposure in different human tissues. To this end, we analyse genomes three human cancer cell lines, A549 (lung adenocarcinoma, epithelial cells), HAP1 (chronic myelogenous leukaemia, blood cells) and MCF7 (breast carcinoma, epithelial cells). Those cell lines were irradiated with two types of particle radiations – proton and helium fluxes – in order to identify the mutational footprints induced and their potential cell type-specificity.

Results

Determining proton and helium ion fraction, and irradiating the cell lines

The dosage of radiation was determined experimentally in order to achieve between 40% and 50% lethality (corresponding to 50%-60% clonogenic survival), independently across the two types of cosmic rays (Figure 1A-B).

A

B

C

Figure 1. Overview of the experimental design. A-B. Experimental identification of radiation dose in
order to achieve ~50% of cell lethality (i.e. ~50% of cell survival) independently across the three cell lines
and across the two types of radiations. SF, survival fraction. C: Schematic overview of the experimental
design of the study. A549, MCF7 and HAP-1 cell lines were exposed to GCR (protons or helium ions).
GCR: Galactic Cosmic Radiation. Single cells from treated and untreated conditions were separated by
FACS and clonally expanded. DNA from single-cell derived populations was extracted and subjected to
whole genome sequencing.

This led to a dose of 0.5 Gy for helium fluxes and a dose of 1 Gy for proton fluxes. After

an expansion of the three cell lines collected (A549, HAP1 and MCF7), the cells were irradiated

using the 5.5 MV Singletron accelerator at the Radiological Research Accelerator Facility

(RARAF; see Methods for details about the procedure, with the corresponding dosage). We

exposed each cell line four times every 4 days to allow cell recovery between each exposure to

facilitate mutation accumulation.

After the final latency period of 4 days, the cells were collected, sorted in a 96-well plate,

and a clonal expansion was performed to obtain colonies with a near-identical genome to

facilitate identifying mutations (Figure 1C). Finally, the DNA was extracted from those colonies

and whole-genome sequenced (WGS) on an Illumina NovaSeq 6000 machine. Details of the

following bioinformatics analysis are described in Methods.

Overall burden of various mutation types generated by cosmic radiations

After the GCR treatments, the cell line genomes contained newly-acquired point mutations

(single nucleotide variants, SNVs), short insertions and deletions (indels), and structural variants

(SVs) in all three cell lines assayed, HAP1, A549 and MCF7, and regardless of the radiation

type, protons or helium ions. As a control, we considered sham-irradiated (i.e. untreated) cells

for HAP1 and A549 cell lines, which also accumulated certain numbers of SNVs, small indels

and SVs, consistent with them undergoing cell divisions and also potentially being exposed to

various stresses during handling.

The global number of observed mutation events (SNVs, indels and SVs) does not

markedly differ between the conditions, suggesting that proton and alpha radiation exposure is

not highly mutagenic to cultured human cells using the fractionated dosing regimen applied

herein. Nevertheless, we detected roughly double the amount of SNVs in the MCF7 cell line

than in the two other cell lines after exposure to either particle (with the exception of one helium

clone, Figure 2A).

Figure 2. General description of point mutations detected in the clones. A. Distribution of the number of SNVs, small insertions and small deletions per clone in each of the three analysed cell lines. Proton radiation generated the highest number of point mutations and small deletions. In HAP1 cell line, this treatment generated also more insertions compared to the other conditions. B. Distribution of the variant allelic fraction (VAF) of detected point mutations for each clone of each cell lines. A VAF peak around 50% or 100% would correspond to a selection of precisely one cell from the cell culture, whereas a peak close de 25% would correspond to the isolation of two cells.

This may indicate a mutator phenotype in our MCF7 cell line, possibly due to DNA mismatch repair (MMR) failure (see below). However this is difficult to ascertain since the absolute number of mutations between cell lines and also between conditions cannot be compared precisely, due to a potentially variable number of cell cycles the cells have undergone.

In all irradiated cell lines, the proton treatment generated both the highest number of SNVs (around 4.8 thousands) and the highest number of small deletions (around 500), compared to the other conditions (Figure 2A). An enrichment of deletions, as we observed in the proton-treated clones, has recently been highlighted in ionizing radiation-associated tumors[13,14] which are typically treated with X-rays. The globally most-mutated clone is a single proton-irradiated sample of the HAP1 cell line, which contains 1,986 more SNVs (Figure 2A) than the most-mutated control (non-irradiated) HAP1 sample. Proton flux generated also the highest number of insertions in the HAP1 cell line (~500 in both replicates), whereas, for this type of alterations, there was not a clear impact of the proton treatment in the A549 and MCF7 cell lines

196 (Figure 2A). This suggests that mutagenic impact of proton irradiation on SNVs, insertions and
197 deletions may be specific to cell-type and/or the genetic background.
198
Variant allelic fractions (VAFs), a metric corresponding to the proportion of sequencing
reads that contain a particular variant, is a proxy for the proportion of cells in the sequenced
population that harbour the mutation. In our experiments, we aimed to select a single treated
cell and expand it into a clone, thus it is expected that the VAFs observed should be centered at
0.5 for heterozygous diploid genome segments or at 1 to homozygous mutations e.g. in LOH
segments (or also genome-wide in the haploid cell line HAP1; note that this cell line can grow as
haploid or diploid and may switch spontaneously). In practice, three different types of VAF
peaks were observed, one at 0.25 which suggests that two cells were collected at the time of
bottlenecking (or alternatively, that the genome segment may be tetraploid), one peak at 0.5
which suggests that one cell was collected, and one peak at 1 for the HAP1 cell line, as
expected due to the possible haploid state of that cell line (Figure 2B).
210

Identification of SNV trinucleotide mutational signatures

212
Upon classifying SNVs into the 6 different categories, tallying DNA strand-symmetrically, i.e.
C>A, C>G, C>T, T>A, T>C, T>G, the most prevalent class of SNVs in all cell lines/clones was
C>A. One important cause of C>A transversion mutations in cancer genomes is oxidative
damage to guanine (Signatures SBS18 and SBS36 as reported previously[26]). Moreover, an
abundance of C>A mutations was observed in recent mutation-accumulation experiments on
human cell lines similar to ours[31,37], plausibly due to exposure to atmospheric oxygen during cell
culture.
220
221   Additionally, we also observed an enrichment of the C>T and T>C transition mutation
222 types (Figure 3A), which is consistent with commonly observed mutational signatures in dividing
223 cells, namely the 'clock-like' mutational signatures SBS1 and SBS5/SBS40[15].

224
Figure 3. Mutational signatures of SNVs in irradiated human cell lines. A. Break-down of point
mutations into the 6 main type of mutations (*i.e.* C>A, C>G, C>T, T>A, T>C, T>G), per clone and per cell
line. Values are shown log$_{10}$ transformed. All genomes were enriched for C>A (possibly due to oxidative
damage to DNA), and C>T and T>C transitions (likely due to 'clock-like' mutational processes). B.
Number of point mutations attributed to each of the 7 different extracted NMF mutational signatures, for
each sample. A signature was considered to be active in a sample if the proportion of total number of
mutations attributed to this particular signature is at least 10%. The SBS96A, SBS96B and SBS96C (all

likely related to experimental procedures) were found in the majority of the samples. SBS96D (consistent
with an MMR deficiency) was found in many MCF7 clones, but also in the control samples of the HAP1
cells. One particular signature, the SBS96E, a signature enriched in T>A transversions, was found in
MCF7 but was not active in non-radiated samples (of other cell line; MCF7 did not have a corresponding
unirradiated sample available). C. Activity matrix of extracted mutational signatures in various clones
herein. D. Cosine similarities between PCAWG mutational signatures and our 7 extracted SBS
signatures. The SigsPack R package was used to obtain the mutational profiles of PCAWG signatures.
239
240   To further classify SNV mutation types, as is usually performed, the mutations were
241 classified into 96 categories, taking into account both the observed class of mutation and the
242 trinucleotide context (A_A, A_C, ..., T_T), in order to infer the 96-component spectrum of SNVs
243 for each cell line and each clone.
244
We extracted *de* novo mutational signatures from our irradiated and control clones via
the SigProfiler tool[15]. To better identify similarities with known mutational processes from other
controlled experiments on human cell line models, our mutation data from proton and alpha
particle irradiated clones was pooled with previous genomic data: (i) 54 cell line genomes
treated with various mutagens, largely chemicals, in which a specific SNV signature was able to
be identified[31], and (ii) additional 173 cell line genomes generated from CRISPR-Cas9 gene
knockouts of 43 DNA repair genes[37]. By a joint analysis of our and previous cell line mutation
accumulation data, we identified 7 independent mutational signatures (see Methods) based on
SNVs (here called SBS [single base substitution]) (Supp. Figure 2). In order to attribute a
potential aetiology to each signature, we computed the similarity with the catalogue of known
somatic signatures, referred as PCAWG signatures[15] (Figure 3C; of note, these signatures are
often referred to in the literature as "COSMIC", by the name of the database, however this is
unrelated with "cosmic radiation" and this previous catalogue does not contain known signatures
of cosmic radiation). We further computed the similarity to the signatures of deficient DNA repair
or of exposure to mutagenic chemicals from two previous studies[31,37].
260
The SBS96A was assigned to (Figure 3D) both PCAWG SBS4 (a tobacco smoking-
associated signature), SBS45 (C>A-rich signature, likely an experimental artefact due to 8-oxoG
generated during preparation of DNA for sequencing[38]), and also to experimental exposure to
PAHs (polycyclic aromatic hydrocarbons, a component of tobacco smoke) in cell lines[31]. The
SBS96B was assigned to the PCAWG signature SBS18, which was reported to result from a
damage to guanines by reactive oxygen species[15,39,40]. This signature was also found to be
significantly active in a *OGG1* knockout cell line genome (Supp. Figure 1A), consistent with the
known function of base excision repair protein OGG1 in mending nucleobase oxidative
damage[41]. The SBS96C was not assigned to a known reported signature in PCAWG catalog
(Figure 3D), but 45% of mutations in the previous *EXO1-/-* cell line WGS[37] was attributed to this
mutational signature. EXO1 is a nuclease with various roles in the MMR, translesion synthesis,
and homologous recombination repair pathway (reviewed in ref. [42]), suggesting that a defect in
one or more of the DNA repair pathways generates the signature SBS96C. The SBS96D was
assigned to a set of four PCAWG signatures: SBS46, SBS37, SBS12, SBS26. Of those, SBS26
is known to be associated with defective DNA mismatch repair (MMR), and is dominated by
C>T transitions as the three other signatures, suggesting that SBS96D may be signature of

MMR deficiency. Consistently, SBS96D was the most active signature in the four *PMS2*-/- cell
lines that were available[37], suggesting that a particular type of MMR deficiency may be
generating this mutational footprint. The SBS96E was assigned to the PCAWG signature
SBS22, resulting from an exposure to aristolochic acid, an agent generating bulky nucleotide
adducts and consisting of T>A transversions, and SBS25, associated with a chemotherapy
treatment[43,44], again rich in T>A changes. The SBS96F was assigned to the PCAWG signature
SBS44 and to the *MLH1* gene knockout[37], indicating another signature of a defective DNA
mismatch repair, thus mechanistically related to but distinct from SBS96D mentioned above.
Finally, the SBS96G was assigned both to the known mutational signature associated with
ultraviolet light exposure (SBS7b), and also to a simulated solar (i.e. non-ionizing, UV-
containing) radiation treatment of a cell line[31].
288
289 SNV mutational signatures predominantly active in certain samples
290
To identify the mutational processes operative in our cell line samples, we considered a
signature to be active in a particular sample if the proportion of mutations attributed to that
signature is at least 10% of the total amount of mutations. The signatures SBS96A, SBS96B
and SBS96C were found in all the three cell lines and in the majority of the samples of each line
and were also abundant in the non-irradiated control samples (Figure 3C). This means that
signatures of adducts at C:G pairs (SBS96A), of oxygen reactive damage (SBS96B), and
possibly deficient various repair pathways (see above for SBS96C) are not a result of the effects
of radiation, because there were detected also in the non-irradiated clones at similar
proportions. The signature of DNA oxidation reaction, SBS96B, has a higher activity in the A549
lung cancer cell line (Figure 3C), suggesting either a more rapid formation of reactive oxygen
species in that cell line, and/or a higher rate of resulting DNA damage, and/or a lower rate (or
accuracy) of repair of oxidized DNA in that cell line.
303
The signature associated with MMR deficiency via the *PMS2* gene and/or SBS26
(SBS96D) was found in almost all samples of the MCF7 cell line (7/8 samples), and also in the
two non-irradiated clones from the HAP1 cell line. This suggests that a form of MMR deficiency
was probably acquired during the cell growth of HAP1, and is present in all our MCF7 clones,
and so this signature did not result from radiation (we note there is no available MCF7 non-
irradiated WGS data to compare). The second, "canonical" MMR-associated signature that
associated to the *MLH1* gene loss, the SBS96F, was not active in our samples; instead, it was
seen only in the external data sets we integrated. Thus GCR exposure does not commonly
select for cells with a deficient MMR system, as is the case for some chemical exposures that
generate mismatch-like DNA lesions, such as drugs TMZ or 6-TG; reviewed in ref. [45]. Overall,
signatures of deficient MMR in our data appear unrelated with alpha or proton radiation
exposure.
316
317   The mutational signature of the non-ionizing UV radiation SBS96G, was, expectedly, not
318 detected at notable levels in our clones. The minor fraction (around ~5%) of observed mutations
319 attributed to this process was similarly active in the non-irradiated conditions, and so is not
320 linked with ionizing radiation, but is more likely to result from imprecisions in the mutational

321   signature inference methodology.
322
One particular signature was associated with ionizing radiation exposure, specifically,
the T>A transversion-rich signature SBS96E (Figure 3B-C). This was detected in MCF7
irradiated samples, both in helium-treated and proton-treated clones, without being active in the
non-irradiated samples of various cell lines (note that non-irradiated MCF7 WGS was not
available). This observation suggests two novel insights. Firstly, the SNV mutational
consequences of particle radiations are likely to be tissue-specific and/or genetic background
specific, since they were observed consistently across clones in one of the three cell lines
(breast), but not in the other two (lung or blood). Secondly, the proton/helium radiation effect
can mimic the mutational footprint of mutagens generating bulky adducts at A:T nucleotide
pairs, similar to aristolochic acid or certain chemotherapies. Speculatively, such a bulky DNA
lesion might be generated by cross-linking nucleotides, or by cross-linking DNA with proteins.
Since this mutational signature of point mutations was found in both types of radiation we
tested, it does appear broadly relevant to various particle radiation types present in GCR.
Moreover it is expected to result from other sources of alpha or proton radiation (e.g. in cancer
radiotherapy) in the susceptible cell types and/or genetic backgrounds.
338
339   **Mutational signatures of indels across cell types and conditions**
340
Small insertions and deletions (indels) can be classified into 83 types (the ID83 categorization,
based on PCAWG indel signatures[15]). In brief, the indels are separated into 1-bp or longer
indels, insertions or deletions, at repeats or not, and finally deletions with micro-homology are
counted separately; these classes can further be divided into subclasses based on the
gained/lost nucleotide, or based on the length of the alteration. Each observed indel is assigned
to one category, thus providing the mutational spectrum of indels for each cell line and each
condition (Figure 4A).
348

349
Figure 4. Mutational signatures of small indels in irradiated human cell lines. A. Mutational profiles
of the 4 indel signatures extracted from pooled data (from our data and from the two other cell line studies
that reported DNA repair deficient background and mutagenic chemical exposed cells[31,37]) using the
SigProfiler algorithm. Indels were classified into 83 distinct categories following the procedure used in
Alexandrov *et al.*[15]. B. Number of indels attributed to each of the 7 different extracted signatures, for each
sample. ID83A and ID83B, characterized by DNA replication errors are found in all the clones. ID83C and
ID83D are unique to radiated samples. C. Ratio of number of deletions over number of insertions.

In order to estimate the potential indel signatures as footprints of the helium and alpha radiation
treatments, we ran the SigProfiler extractor tool on the generated indel count matrices. The
matrices were from the pooled dataset, consisting as above of genomes from our irradiated
MCF7, A549 and HAP1 cell lines, as well as genomes of mutagen-treated and DNA repair
deficient cell lines from previous work[31,37] (see Methods for details on samples).

Four indel signatures were identified: ID83A, ID83B, ID83C and ID83D (Figure 4A), as
well as the absolute activity of each in our experimental cell line samples (Figure 4B). To learn
about the potential underlying mechanisms, we also estimated the signatures' activity in WGS of
previous DNA repair knockouts and the mutagen-treated samples[31,37], as for SNV signatures
above. ID83A was linked to DNA mismatch repair deficiency (*MSH2*, *MLH1* and *MSH6*). ID83B
was found in DNA polymerase knockout samples (*POLB*, *POLM* and *POLE4* [37], see Supp.
Figure 1B). The ID83C may be linked to various mechanisms (Supp. Figure 1B), and its
spectrum resembles the PCAWG ID1, suggested to be related to slippage during DNA
replication[15].

Finally, the ID83D signature, consisting mostly of 1-bp deletions and ≥5bp deletions at
micro-homology regions, was found to be active in a cell line knocked out in *XRCC4*[37], a core

factor of the non-homologous end joining (NHEJ) pathway for DSB repair (Supp. Figure 1B). A
loss of this gene (whose full name is "X-Ray Repair Cross Complementing 4") is well-
appreciated to increase sensitivity to ionizing radiation[46–48]. Moreover, this indel signature is
similar to the indel mutational signature provoked by ionizing photon radiation, reported in a
recent preprint, where clonal organoids from mouse and human cells were irradiated[49].
382
383   ID83A and ID83B were found in majority of our cell lines/conditions including controls,
384 suggesting they may represent errors intrinsic to DNA replication. These errors may be
385 enhanced in case of MMR failures, and thus observed more abundantly in MMR mutants (see
386 above).
387
The signatures ID83C and ID83D were found in only a subset of our samples and more
interestingly they were absent in the controls. ID83C was active in HAP1/proton, A549/helium
and MCF7/helium conditions (Figure 4B), suggesting that radiation may cause a mutational
footprint resembling that of strand slippage during DNA replication. In our set of cell lines, this
mutational effect does not appear not-tissue specific nor genetic background-specific,
suggesting an universal indel mutational footprint of particle-based radiation. However, we do
not rule out that different cell types might be differentially affected by different radiation types
(protons *versus* helium) generating different indel signature burdens.
396
397   ID83D was found only in the MCF7 cell line and in both irradiated conditions, suggesting
398 a possible tissue-specific mutagenic effect (it was not seen in blood or lung cells); this leaves
399 similar footprints to those reported elsewhere for photon-based ionizing radiation[49].
400
401 **Mutational signatures of structural variation**
402
403   Large structural variants (SVs, also referred to as rearrangements) were grouped into four
404 master classes: large deletion (DEL), large insertions (INS), tandem duplications (DUP) and
405 inversions (INV). No particular SV category has been found to be the most prevalent across the
406 cell lines and across the cell lines/conditions (Figure 5A).
407

Figure 5. Structural variation in irradiated human cell lines. A. Distribution of the number of structural variants (SVs), decomposed into large deletions, large insertions, duplications and inversions, per clone inside each of the three analysed cell lines. Large insertion was the less-represented class of the SVs detected in the clones. B. Ratio of the number of large deletions over number of large insertions.

The rarest event type – insertions – is similarly observed across the cell lines and clones; their low frequency may however stem from technical difficulties in calling insertion variants from short-read WGS, resulting in many undetected variants. More generally, in our experiments, we observed globally a low number of SVs of any type, suggesting that the impact of the proton and alpha ray treatment on rearrangement rates in human cells may not be remarkable, at least in the treatment regimen applied here. Irradiated clones are not enriched in SVs when compared to control cells, except in the case of HAP1 cell line/alpha radiation, where we observed approximately two times more large deletions (23 in the most mutated clone) compared to the controls (10 in the most mutated clone).

Globally, helium (alpha) treatment generated the higher total number of SVs in both HAP1 and MCF7 cell lines, compared to proton treatment and to untreated (for HAP1). Also in the A549 lung cell line, results were generally consistent with this trend of propensity of alpha radiation to generate SVs, however less pronounced since both the non-treated clone and the helium-treated clone exhibited a high number of SVs.

We also investigated another quantitative measure of the identified variations: the
distributions of the SV sizes, as commonly done in cancer genome analysis[27,29]. Again, we
asked if there is a difference between two types of radiation treatments, also comparing them
with the non-irradiated controls. Because insertions were rarely detected, here we considered
only deletions, duplications and inversions. While the distribution of SV lengths was not
significantly different across the conditions, nevertheless, we observed a trend for deletions that
is consistent across all the cell lines. In particular, proton radiation generated the smallest
deletion events, followed by alpha, and finally longest deletions were observed in untreated
controls (Supp. Figure 3). Inversions behaved similarly: we also observed that treated clones,
whatever the radiation type, contains inversions smaller than those in non-treated samples.
Moreover, there is also a trend toward longer inversions resulting from helium compared to
protons (Supp. Figure 3). Overall, proton radiation generated SVs of shorter lengths than alpha
rays or than the default mutagenesis in untreated cells, suggesting a different mode of DNA
damage and/or repair thereof following proton exposure in various human cell types.

Finally, we investigated the enrichment of deletion SVs compared to insertion and
duplication SV (the latter two were considered to be similar and merged for this analysis). As
previously reported for ionizing radiation-associated second malignancies[14], the ratio of
deletions to insertions increases in our particle-irradiated human cell lines. We did not find this
bias towards deletions to be tissue-specific (Figure 5B). Overall, different types of ionizing
radiation, either X-rays or gamma as commonly employed for cancer therapy[13,14], or protons
and helium/alpha particles as tested here, exhibit an overall similar spectrum of increase in
deletion SVs relative to insertion/duplication SVs.

Regional mutation rate with respect to replication time and gene activity

DNA replication timing (RT) is correlated with regional variability in somatic mutation rates in
human tissues. There is an increased mutation rate in late-replicating heterochromatic regions
of the genome, because DNA repair mechanisms preferentially protect the early-replicating
euchromatic, gene-rich regions[50,51]. We were interested if there is an enrichment of radiation-
associated mutation rates in regions with a specific RT. Although the correlation between the
SNV density and the RT bins are not significantly different between untreated cells and
irradiated cells (and between conditions), we did observe a trend for radiation-associated SVs
(Supp. Figure 4). In particular, alpha radiation, compared to untreated clones, generated more
SV mutations in early-replicating regions, compared to late-replicating regions; this was
observed in A549 and MCF7 cell lines. In proton radiation this trend is observed more strongly
in the HAP1 cell line (Supp Figure 4B). Of note, the mutation enrichment in early-replicating
DNA was not observed for point mutations (Supp Figure 4A), suggesting that this global
"redistribution" of mutation rates[52] upon irradiation is specific to SV mutagenesis. Early-
replicating regions have a higher gene density and are enriched with active genes. Therefore
the increased rate of SV formation in early-replicating DNA upon particle radiation may have
higher potential for functional impact, such as tumor suppressor gene inactivation.

Additionally, we considered mutation rate association in relation to presence of DNA

methylation, which can be mutagenic e.g. it is thought to underlie generation of the very

common mutational signature SBS1[53,54]. Also, gene body DNA methylation tends to be

positively associated with gene expression levels, thus DNA methylation-associated mutational

processes are expected to be more impactful[55]. We did not observe a clear association signal

of mutation rates with DNA methylation for all the three cell lines (Supp Figure 4C). However,

there is a trend in HAP1 blood cells for both SNVs and SVs, with fewer alterations in lower-

methylated regions in irradiated cells compared to untreated cells (Supp Figure 4D). Prompted

by this, we also investigated the distribution of mutations across regions harbouring various

levels of gene expression. We observed that HAP1 proton condition generated more SV in

highly expressed genes (Supp. Figure 4F) compared to the control non-radiated cell lines. This

is consistent with the above RT analysis, suggesting that the exposures to particle radiation may

increase SV mutation rates preferentially in active-chromatin regions, and thus would be

expected to result in impactful alterations. However this effect may be cell type specific and/or

genetic background specific.

Clustered mutation processes due to irradiation

Next, we considered patterns of mutation clusters, which can be particularly informative about

various mutational processes in human cells[28], most prominently with regard to APOBEC

enzyme DNA damaging activity[53,55,56] and also error-prone DNA polymerase usage[28].

Operationally, we defined a mutation cluster as a set of point mutations at a pairwise

intermutational distance lower than 1Kb; given the overall modest mutation burden in our

genomes, this threshold is unlikely to result in many artefactual (false-positive) mutation

clusters. In the A549 cell line, 61 mutations were clustered in control clones, 109 in helium-

irradiated clones and 93 in proton-irradiated samples. In the HAP1 cell lines, a total of 57

mutations were clustered in control clones, 262 in helium-irradiated clones and 181 in proton-

irradiated samples. Finally, in the MCF7 cell line, we detected 344 clustered mutations in the

helium-irradiated clones and 373 in the proton-irradiated clones (Figure 6A); note that untreated

control DNA WGS is not available for MCF7.

505
506 **Figure 6. Clustered point mutations associated with irradiation.** A. Distribution of the genomic
507 distance between point mutations that were identified as being clustered together (see methods for the
508 definition of a mutational cluster). *P*-values shown were computed using the *ggpubr* R package (t-test
509 mean comparison). B. Distribution of the size of each individual cluster (number of mutations), defined as
510 a graph component (see methods for details).

511
512

We detected more clustered mutations in radiation conditions compared to non-radiation
conditions for both cell lines that could be analyzed, indicating that alpha and proton radiation
generates mutation clusters in human cells. This pattern of mutagenesis is consistent with many
reports of ionizing radiation damaging DNA in a clustered manner[57–59]. With respect to types of
radiation, in the HAP1 and A549 cell lines (but not MCF7 cells) more clustered mutations are
found in helium clones compared to proton clones.

519
520   We next asked if the clustered mutations, depending on the cell line and the radiation type,
521 were found at closer distances. In HAP1 cells, clustered mutations were spaced closer together
522 for both helium and proton radiations compared to the control clones. In that cell line, helium
523 radiation generated more distant clustered mutations compared to proton radiation, but this was
524 found to be the opposite in the MCF7 cell line (Figure 6A).

525
We were then interested in the repartition of the clustered mutations, *i.e.* do they form a lot

of small clusters or do they form a few large (multi-mutation) clusters, the latter case
corresponding to a kataegis-like phenomenon[53] and the former to an omikli-like phenomenon[55].
We applied a graph-theory approach to quantify this (Methods). Considering large clusters as
connected components of size higher than 5, we identified one large cluster in A549/helium,
four in HAP1/helium, one in MCF7/helium and one in HAP1/proton. However there were no
large clusters in the untreated cells, suggesting mutation showers can be generated by ionizing
radiation (Figure 6B). Of note, the majority of clustered mutations, both in the irradiated and in
the untreated conditions, were in smaller, *omikli*-like clusters..
Overall, this suggests that the clustering of mutations often results from exposure to particle
radiation, that it may be tissue and/or genetic background specific, and that it may occasionally
generate multi event, kataegis-like mutation clusters,.
538

Discussion

540
In this study, we analysed genomes of three human cancer cell lines irradiated with two different
types of GCR – proton and helium fluxes – in order to characterize the spectra of DNA
alterations that GCR would produce in different human tissues. Overall burden of SNV
mutations, indels and SV events was not markedly different between treated and untreated
cells, which suggests that GCR exposure – at least in the regime applied in our experiments – is
not grossly mutagenic (see below for further considerations).
547
However, there were multiple differences in mutation spectra and distributions between
GCR-treated and untreated cells. For instance, in agreement with a recent study analysing
ionizing radiation-associated tumors[14], we observed an enrichment of deletions compared to
insertions, and this was even more the case for large (SV) events (Figure 4C, Figure 5B). This
phenomenon was also observed in a recent study that reported a 3.6-fold increase of small
deletion burden when comparing radiotherapy treated glioma patients to untreated glioma
patients[13]. We note that tumor radiotherapy usually employs X rays or gamma rays, rather than
the protons and alpha particles tested here, and so this enrichment of deletion mutations
appears broadly independent of the type of IR applied to cells.
557
We found that GCR generates point mutations that are more often clustered than those
found in non-irradiated samples, moreover the distance between mutations in clusters and size
of the clusters may vary between tissues and conditions. This is in line with recent work
suggesting clustered mutations to be more specific indicators of mutagenic processes[28,55,56]
than the general, genome-wide mutation signatures. The types of clusters were also variable:
proton treatment generated clustered mutations with shorter inter-mutational distances
compared to helium treatment in some cell lines. Helium treatment generates larger, multi-
mutational clusters compared to proton treatment. Those two observations, overall, suggest that
the clustered mutational distributions generated by helium and proton radiation exposure may
differ, and potentially exhibit tissue-specificity and/or genetic background-specificity; the latter
point would be clarified with additional experiments in future work.
569
We were also interested in the SNV trinucleotide mutational signatures that helium and

proton GCR can generate, and whether known biological mechanisms can be linked to those
footprints. While most observed SNV signatures were consistent across all treated and
untreated samples, e.g. reactive oxygen species mutagenesis was noted universally, some
samples exhibited signatures consistent with MMR deficiency (this was however unlikely to
have resulted from, or have been selected by irradiation, because we detected it also in a
control sample). Interestingly, we observed a SNV signature in both irradiation conditions of the
MCF7 cell line, but not in the controls we analysed (note these controls were in other cell lines
apart from MCF7, see Methods): an T>A rich SBS signature, resembling previous signatures of
aristolochic acid and some chemotherapy treatments[53,54], suggesting, speculatively, that particle
radiation may generate DNA damage resembling bulky adducts in some cell types.
581
Similar to SNVs, some indel signatures were radiation-associated: ID83C and ID83D.
ID83C was found in the three cell lines, in proton and helium exposed samples, thus suggesting
a likely general effect over tissues for this particular mechanism. The second GCR-associated
indel signature in our experiment, ID83D, was also highly active in *XRCC4*-knockout sample
from an earlier study[37], suggesting that ID83D represents the mutational signature of a backup
DSB repair pathway employed in absence of XRCC4 mediated non-homologous end joining
repair. While ionizing radiation is expected to generate DSBs in all cell types similarly, the
ID83D signature was observed only in MCF7 cells, suggesting this signature might derive from
radiation applied to cells with a particular genetic background, potentially a DNA repair deficient
one.
592
593 The solar radiation SNV signatures identified previously[15,31] (presumably UV generated)
594 were observed at low intensities in our irradiated samples, likely stemming from imprecisions
595 during statistical analysis. This is consistent with prior knowledge that ionizing cosmic radiation
596 affects the DNA differently than the non-ionizing UV radiation[60].
597
Even though overall mutation rates were not increased in irradiated cells, we did note a
relative increase of some mutation types, prominently SVs, in the active regions of the genome,
which are early-replicating and contain highly expressed genes. This type of redistribution of
mutation rates across the genome can increase the likelihood of impactful mutations occurring
even in the absence of overall increased mutation burdens[52,55]. Such phenomena were
previously linked with failures in DNA repair[50] and with activity of the APOBEC3A cytosine
deaminase enzyme[55], and here we show that to a certain extent this happens also with ionizing
radiation. Therefore, mutational impact on active genes may be increased by ionizing radiation
exposures via directing SVs towards active chromatin.
607
608 Our study also has several limitations. For instance, some components of GCR were not
609 considered in this study, in particular the heavy atom nuclei (often referred to as "HZE ions"),
610 highly energetic particles likely originating from supernova explosions. Despite their small
611 proportion in cosmic rays, their biological impact might be large, and their mutagenic effects on
612 genome stability remain to be studied.
613
Next, we considered only a single dose of radiation that resulted in a ~40% to 50%

reduction of cell viability assessed by the colony formation assay, and a fractionation regime of
four fractions (given every four days) exposures of this dose. The dose studied here had rather
strong biological effects on cell viability, and such dosages would probably be well sufficient to
observe the mutagenic effects of GCR. Since a gross increase in mutation rates was not
observed, we infer that GCR exposures are not highly mutagenic. However, we cannot rule out
the (less parsimonious) scenario where lower doses of GCR than those employed here might
be more mutagenic e.g. by failing to trigger cell cycle checkpoints, thus 'slipping under the radar'
of the mechanisms protecting genome integrity and introducing mutations.
623
624 Moreover, given that for several mutation patterns we observed tissue-specific and/or
625 genetic background specific responses (these scenarios cannot be distinguished), we
626 emphasize more experiments would be needed on different cell lines or other experimental
627 models to ascertain tissue-specific responses to radiation types.
628
Finally, our data overall suggests that components of GCR are not overtly mutagenic to
human cells, with some effects on SV distributions, and on indel and clustered mutation spectra
However, the intermittent exposure regime that we employed for reasons of feasibility is
different from longer-term, chronic exposures to GCR. Even modest increases in mutation rates
under chronic GCR exposures e.g. during space flight might have detrimental effects on cancer
rates, and possibly neurodegeneration and other aging phenotypes. Importantly, germline cells
might be affected, with potential effects on reproductive health and on genetic disease incidence
in the progeny. Therefore we highlight the necessity of further experimental work on cell and
animal models, using longer-term exposures to components of galactic cosmic rays to ascertain
their effects on the stability of the human genome.
639
640
641 **Methods**
642
643 **Cell lines**
644
We used three different cell lines: A549, HAP1 and MCF7. The lung adenocarcinoma cell line
A549 (CCL-185) and the breast adenocarcinoma cell line MCF7 (HTB-22) were purchased from
ATCC (the American Type Culture Collection, Manassas, VA). The near-haploid cell line derived
from the KBM-7 cell line HAP1 (C859) was purchased from Horizon (Carle Place, NY). All cell
lines were cultured and maintained according to recommended protocols. All the cell lines were
authenticated by STR profiling by the corresponding repositories.
All the cells were grown at 37°C in a 5% CO_2 humidified incubator; the A549 cells were grown in
Eagle's Minimum Essential Medium (MEM), MCF7 in Dulbecco modified MEM, and HAP-1 cells
in Iscove's modified Dulbecco's medium (IMDM). All the media were supplemented with 10%
heat-inactivated (56ºC, 30 min) fetal bovine serum (FBS), and 100 U/ml penicillin and 100
mg/ml streptomycin (all from Sigma–Aldrich Corp).
656
657 **Cell irradiation, culture and clones isolation**
658
A clonogenic assay[61] pilot experiment was conducted to assess the dose that resulted in 40-

50% lethality (i.e., 50-60% survival) to 11 MeV helium ions (LET = 65 keV/µm) or 5.4 MeV
protons (LET = 10 keV/µm), which was 0.5 and 1 Gy, respectively (Figure 1A,B). Cells for the
experiment were then exposed to four fractions delivered every four days. Cells were collected
four days after the last fraction. For each sample, half of the cells were frozen at -80 C while the
other half was re-plated until confluency. The cells were then collected, separated by vigorously
passing through a 21 G syringe and 3 ul/ml of DAPI was added to the solution for cell sorting at
the BD Influx cell sorter. For each cell line, single cells were sorted in two 96-well plates and
incubated in 250 ul of medium supplemented with 20% FBS. Once colonies were visible, we
picked ~30 colonies and re-plated each one in an individual well of a 12-well plate in the case of
A549 and MCF7 and in 48-well plate in the case of HAP1 cells up to confluence. The cells were
then collected and half frozen at -80C while the other half was replated in 6-well plates and then
in 100-mm Petri dish until confluence. DNA was extracted with PureLink Genomic DNA kit
(Invitrogen), frozen and shipped to the sequencing centre.
673
674   **Cell irradiations at the track segment irradiation platform**
675
The 5.5 MV Singletron accelerator at the Radiological Research Accelerator Facility (RARAF)
served as source of energetic protons and alpha particles. The accelerator was operated at the
maximum terminal voltage generating beams with nominal energies of 5.5 MeV and 11 MeV for
protons and alphas, respectively. Cells were irradiated at the so called "track segment"
irradiation platform whose name indicates the traversal of thin samples (typically cell
monolayers) by a short segment of the ion's trajectory resulting in a small variation of the linear
energy transfer (LET) throughout the sample. It is valid therefore to assume that radiation doses
at the track segment platform are delivered by mono-LET beams. The LET values of the applied
beams were 10 keV/µm for protons and 65 keV/µm for alphas. Detailed description of the track
segment irradiation platform and its operational principles can be found elsewhere[1,2,3,4]. We will
state here only the main features of the irradiation protocol. The protocol can be generally
divided in two parts. First, beam characterization and dosimetry were performed with the use of
different detectors mounted on the metal wheel that rotates over the beam exit aperture. The
aims of this step were to verify the energy and the LET of the beam, to check the uniformity of
the irradiation field covering the 6 mm x 20 mm beam exit aperture (2.9 µm thick havar foil) and,
finally, to calibrate the online beam monitor in terms of the absorbed dose delivered to the
sample. In the second step, the dosimetry wheel was replaced with the sample-carrying wheel
that can accommodate up to 20 custom made dishes. The dishes were manufactured by gluing
6 µm thick mylar foil over metal rings having a diameter of 5 cm. The mylar foil serves as the
bottom of the dish on which the cells are attached, allowing thus the ions to penetrate through
and reach the cells without losing much of their energy. The in-house developed software drives
the stepper motor and controls the movement of sample dishes over the beam. A desired dose
was delivered to the sample by exposing it to the beam until the appropriate number of monitor
counts was reached according to dosimetry measurements performed during the first part of the
irradiation protocol.
701
702   **Irradiations of the cells for mutational analysis and clone isolations**
703

The obtained survival curves were fitted with the linear quadratic (LQ) model. Doses of 1 Gy
and 0.5 Gy were selected for investigating the mutational signatures of protons and alpha
particles, respectively. According to the LQ fits, the selected doses resulted in preservation of
between 50% and 60% clonogenic capacity for all cell lines. Four fractions of the same dose
were delivered to the cells every 96 hours to amplify the number of mutations in surviving cells.
Between the fractions, the cells were not removed from the mylar dishes. To prevent the cells to
overgrow the dish size before the end of the irradiations, initial number of plated cells was <
1000.
712
713 **Whole-genome sequencing**
714
Total genomic DNA was extracted from pelleted cells using PureLink™ Genomic DNA Mini Kit
(K182001, Thermo Fisher Scientific). DNA was sequenced on NovaSeq 6000, 150 nt paired-
end mode. Reads were aligned to the reference human genome Hg38 (GRCh38.d1.vd1) using
BWA version 0.7.17[62]. GATK Base Quality Score Recalibration was applied to adjust quality
scores assigned during the sequencing process[63]. The average coverage of the various
samples was 18.5X to 44.8X.
721
722 **Variant detection**
723
Strelka2 variant calling algorithm was used to detect point mutations and small insertions and
deletions[64]. Strelka2 was launched in the joint genotyping mode independently across each
sample and each cell line, *i.e.* using all the samples of one particular cell line as the input of the
variant calling process, in one batch. Mutations were annotated using annovar tool[65] and the
gnomAD genome database v2.1.1[66]. To call structural variants, i.e. large insertions and
deletions, duplications and inversions, we used the Manta tool in the joint calling mode[67], in the
same manner we used Strelka2. For inversions, Manta returns a list of breakpoints that
described the inversions it detected. To transform those breakpoints into inversions, we applied
the convertInversion.py python script provided by the Manta developers
(https://github.com/Illumina/manta). In order to remove common germline variants, we removed
all the mutations reported in the gnomAD database at a frequency higher than 0.1%. In order to
remove the rest of germline variations that are shared mutations across the clones of the same
cell lines, we removed all mutations present in more than one clone. From further analyses, we
excluded the data from the MCF7 control (unirradiated) sample, that harboured an implausibly
high number of mutations (116,971) possibly due to the low quality of the sequenced DNA, or
other artefacts.
740
741 **Mutation clustering**
742
Point mutations were considered to be clustered together if they appear at positions with a
genomic distance lower than 1kb; such mutations would be considered to fall into the omikli
("mutation fog") category[55.] To consider large, multi-mutation clusters resembling a kataegis
(mutation shower) event[53,68], we used an approach based on graph theory. Roughly, we
considered each mutation to be a node of the graph, and edges connect two nodes if the

distance between corresponding mutations is lower than 1kb. Then, we computed the

components of the graph, that corresponds to a connected sub-graph, and subsequently the

size of the components, *i.e.* how many nodes or mutations are present in the component.

Extraction of mutational signatures

We pooled our samples with two already published datasets in order to help NMF to converge

and in order to compare our datasets to signatures of DNA damage and signatures of various

mutagens. We first pooled our data with the mutations reported in Zou *et al.* study[37], which

analysed 173 clones generated by CRISPR-Cas9 knockouts of 73 DNA repair/replicative

pathway genes in human induced pluripotent stem cells. We then pooled the data with the

mutations reported in Kucab *et al.* study[31], which reported 54 samples containing a mutational

signature (note there were additional samples in the study without a prominent mutation pattern,

which were not included here), together with our cosmic-radiation exposed samples of three cell

lines. This pooled dataset of a total of 247 individual samples. In order to extract SNV

signatures generated by our radiation treatments, we applied SigProfiler Extractor algorithm[15]

with default settings on the set of detected mutations, classified into the 96 SNV categories. The

suggested solution contained 12 extracted signatures. Nevertheless, SigProfiler also provided

the summary of all tester set of signatures, and a set of 7 signatures was also found to be

potentially plausible by SigProfiler (based on low reconstruction error and high reproducibility in

mutational signatures, corresponding to an high signature stability, see refs.[54,69] for details). We

observed that, when considering the set of 12 mutational signatures, some mechanisms were

spread across multiple signatures suggesting that some could be merged. We thus considered

the set of 7 mutational signatures for the rest of the analyses. Extracted signature were

assigned to a known PCAWG "SBS" signature[15], if the cosine similarity between the two

mutational profiles was at least 0.85. The SigsPack R package[70] was used to obtain the

mutational profiles of PCAWG signatures.

For indel signatures, we used the SigProfiler Matrix Generator tool to correctly classify the

observed indels into one of the 83 categories defined in Alexandrov *et al.*[15]. Then, the

SigProfiler Extractor tool was launched in the a total set of 321 samples from Kucab *et al.*, Zou

et al., and our experiments, using those generated matrices. We considered the optimal solution

given by the tool, resulting in a total set of 4 extracted indel signatures.

Regional enrichment analysis

We performed a regional enrichment analysis in order to estimate if the mutations detected in

the irradiated clones were located in regions with a specific epigenetic pattern compared to the

mutations detected in the non-treated clones. We tested three genomic features: the replication

timing, the DNA repair histone mark H3K36me3 and the gene expression levels based on RNA-

seq data. Replication timing bins, genomic bins extracted from chromatin mark H3K36me3, and

regions based on variable gene expression were computed as in Supek and Lehner[28]. To detect

significant association of mutations in specific genomic regions, we fitted a negative binomial

regression using the glm.nb function from the MASS R package, on the counts of mutations per

bins, controlling for the 3-nucleotide context and type of mutation in the case of SNV, as in

Supek and Lehner[28]. The regional enrichment analysis source code is implemented using

Nextflow[71] and is available on the GitHub platform

(https://github.com/tdelhomme/RegionalEnrichment-nf/).

Statistical analyses

All the statistical analyses were performed using R software version 3.6.0. VariantAnnotation[72]

R package version 1.32 was used to read the VCF files. The R packages

BSgenome.Hsapiens.UCSC.hg38 version 1.4.1 and BSgenome.Hsapiens.UCSC.hg19 version

1.4.0 were used to load the human genome inside R. MutationalPatterns[73] R package version

1.12 was used to generate the trinucleotide contexts. Ggraph version 2.0.5 and Igraph version

1.2.6 R packages was used to identify the cluster components.

References

1. Larose, T. L. *Tumors in Space: Preparation for Spaceflight*. *Preparation of Space Experiments* (IntechOpen, 2020). doi:10.5772/intechopen.93465.
2. Jastrow, R. Definition of Air Space. in *First Colloquium on the Law of Outer Space* (eds. Haley, A. G. & Heinrich, W.) 82–82 (Springer, 1959). doi:10.1007/978-3-7091-4414-5_16.
3. 100km Altitude Boundary for Astronautics. https://www.fai.org/page/icare-boundary (2017).
4. McDowell, J. C. The edge of space: Revisiting the Karman Line. *Acta Astronaut.* **151**, 668–677 (2018).
5. Thirsk, R., Kuipers, A., Mukai, C. & Williams, D. The space-flight environment: the International Space Station and beyond. *CMAJ Can. Med. Assoc. J.* **180**, 1216–1220 (2009).
6. Panesar, S. S. & Ashkan, K. Surgery in space. *BJS Br. J. Surg.* **105**, 1234–1243 (2018).
7. Hellweg, C. E. & Baumstark-Khan, C. Getting ready for the manned mission to Mars: the astronauts' risk from space radiation. *Naturwissenschaften* **94**, 517–526 (2007).
8. Cucinotta, F. A. Space Radiation Risks for Astronauts on Multiple International Space Station Missions. *PLoS ONE* **9**, e96099 (2014).
9. Nikiforov, Y. E. Thyroid carcinoma: molecular pathways and therapeutic targets. *Mod. Pathol. Off. J. U. S. Can. Acad. Pathol. Inc* **21 Suppl 2**, S37-43 (2008).
10. Morton, L. M. *et al.* Radiation-related genomic profile of papillary thyroid carcinoma after the Chernobyl accident. *Science* **372**, eabg2538 (2021).
11. Brenner, A. V. *et al.* I-131 dose response for incident thyroid cancers in Ukraine related to the Chornobyl accident. *Environ. Health Perspect.* **119**, 933–939 (2011).
12. Ozasa, K. Epidemiological research on radiation-induced cancer in atomic bomb survivors. *J. Radiat. Res. (Tokyo)* **57**, i112–i117 (2016).
13. Kocakavuk, E. *et al.* Radiotherapy is associated with a deletion signature that contributes to poor outcomes in patients with cancer. *Nat. Genet.* **53**, 1088–1096 (2021).
14. Behjati, S. *et al.* Mutational signatures of ionizing radiation in second malignancies. *Nat. Commun.* **7**, 12605 (2016).
15. Alexandrov, L. B. *et al.* The repertoire of mutational signatures in human cancer. *Nature* **578**, 94–101 (2020).
16. Berrington de González, A. & Darby, S. Risk of cancer from diagnostic X-rays: estimates for the UK and 14 other countries. *Lancet Lond. Engl.* **363**, 345–351 (2004).
17. IARC. *Ionizing Radiation, Part 1: X- and Gamma (γ)-Radiation, and Neutrons*.
18. Hughes, J. R. & Parsons, J. L. FLASH Radiotherapy: Current Knowledge and Future Insights Using Proton-Beam Therapy. *Int. J. Mol. Sci.* **21**, E6492 (2020).
19. Yuan, T.-Z., Zhan, Z.-J. & Qian, C.-N. New frontiers in proton therapy: applications in cancers. *Cancer Commun.* **39**, 61 (2019).
20. Li, M. *et al.* Clinical Efficacy and Safety of Proton and Carbon Ion Radiotherapy for Prostate Cancer: A Systematic Review and Meta-Analysis. *Front. Oncol.* **11**, (2021).
21. Hu, M., Jiang, L., Cui, X., Zhang, J. & Yu, J. Proton beam therapy for cancer in the era of precision medicine. *J. Hematol. Oncol.J Hematol Oncol* **11**, 136 (2018).
22. Vogel, J. *et al.* Proton therapy for pediatric head and neck malignancies. *Pediatr. Blood Cancer* **65**, e26858 (2018).
23. Jain, V. *et al.* Predicted Secondary Malignancies following Proton versus Photon Radiation for Oropharyngeal Cancers. *Int. J. Part. Ther.* **6**, 1–10 (2020).
24. Huang, J. & Mehta, M. Can proton therapy reduce radiation-related lymphopenia in glioblastoma? *Neuro-Oncol.* **23**, 179–181 (2021).
25. Mohan, R. *et al.* Proton therapy reduces the likelihood of high-grade radiation-induced lymphopenia in glioblastoma patients: phase II randomized study of protons vs photons. *Neuro-Oncol.* **23**, 284–294 (2021).
26. Alexandrov, L. B., Nik-Zainal, S., Wedge, D. C., Campbell, P. J. & Stratton, M. R. Deciphering signatures of mutational processes operative in human cancer. *Cell Rep.* **3**, 246–259 (2013).
27. Li, Y. *et al.* Patterns of somatic structural variation in human cancer genomes. *Nature* **578**, 112–121 (2020).
28. Supek, F. & Lehner, B. Clustered Mutation Signatures Reveal that Error-Prone DNA Repair Targets Mutations to Active Genes. *Cell* **170**, 534-547.e23 (2017).
29. Degasperi, A. *et al.* Substitution mutational signatures in whole-genome–sequenced cancers in

the UK population. *Science* **376**, abl9283 (2022).

30. Pleasance, E. *et al.* Pan-cancer analysis of advanced patient tumors reveals interactions between therapy and genomic landscapes. *Nat. Cancer* **1**, 452–468 (2020).

31. Kucab, J. E. *et al.* A Compendium of Mutational Signatures of Environmental Agents. *Cell* **177**, 821-836.e16 (2019).

32. Saini, N. *et al.* The Impact of Environmental and Endogenous Damage on Somatic Mutation Load in Human Skin Fibroblasts. *PLOS Genet.* **12**, e1006385 (2016).

33. Franco, I. *et al.* Whole genome DNA sequencing provides an atlas of somatic mutagenesis in healthy human cells and identifies a tumor-prone cell type. *Genome Biol.* **20**, 285 (2019).

34. Davidson, P. R., Sherborne, A. L., Taylor, B., Nakamura, A. O. & Nakamura, J. L. A pooled mutational analysis identifies ionizing radiation-associated mutational signatures conserved between mouse and human malignancies. *Sci. Rep.* **7**, 7645 (2017).

35. Sherborne, A. L. *et al.* Mutational Analysis of Ionizing Radiation Induced Neoplasms. *Cell Rep.* **12**, 1915–1926 (2015).

36. Volkova, N. V. *et al.* Mutational signatures are jointly shaped by DNA damage and repair. *Nat. Commun.* **11**, 2169 (2020).

37. Zou, X. *et al.* A systematic CRISPR screen defines mutational mechanisms underpinning signatures caused by replication errors and endogenous DNA damage. *Nat. Cancer* **2**, 643–657 (2021).

38. Costello, M. *et al.* Discovery and characterization of artifactual mutations in deep coverage targeted capture sequencing data due to oxidative DNA damage during sample preparation. *Nucleic Acids Res.* **41**, e67 (2013).

39. Sekiguchi, M. & Tsuzuki, T. Oxidative nucleotide damage: consequences and prevention. *Oncogene* **21**, 8895–8904 (2002).

40. Kawanishi, S., Hiraku, Y. & Oikawa, S. Mechanism of guanine-specific DNA damage by oxidative stress and its role in carcinogenesis and aging. *Mutat. Res.* **488**, 65–76 (2001).

41. Wang, R. *et al.* OGG1-initiated base excision repair exacerbates oxidative stress-induced parthanatos. *Cell Death Dis.* **9**, 628 (2018).

42. Keijzers, G. *et al.* Human Exonuclease 1 (EXO1) Regulatory Functions in DNA Replication with Putative Roles in Cancer. *Int. J. Mol. Sci.* **20**, 74 (2018).

43. Nik-Zainal, S. *et al.* Landscape of somatic mutations in 560 breast cancer whole-genome sequences. *Nature* **534**, 47–54 (2016).

44. Alexandrov, L. B. *et al.* Clock-like mutational processes in human somatic cells. *Nat. Genet.* **47**, 1402–1407 (2015).

45. Mas-Ponte, D., McCullough, M. & Supek, F. Spectrum of DNA mismatch repair failures viewed through the lens of cancer genomics and implications for therapy. *Clin. Sci. Lond. Engl. 1979* **136**, 383–404 (2022).

46. Bassing, C. H. *et al.* Increased ionizing radiation sensitivity and genomic instability in the absence of histone H2AX. *Proc. Natl. Acad. Sci. U. S. A.* **99**, 8173–8178 (2002).

47. Delacôte, F., Han, M., Stamato, T. D., Jasin, M. & Lopez, B. S. An xrcc4 defect or Wortmannin stimulates homologous recombination specifically induced by double-strand breaks in mammalian cells. *Nucleic Acids Res.* **30**, 3454–3463 (2002).

48. Wen, Y., Dai, G., Wang, L., Fu, K. & Zuo, S. Silencing of XRCC4 increases radiosensitivity of triple-negative breast cancer cells. *Biosci. Rep.* **39**, BSR20180893 (2019).

49. Youk, J. *et al.* Mutational impact and signature of ionizing radiation. 2021.01.12.426324 (2021) doi:10.1101/2021.01.12.426324.

50. Supek, F. & Lehner, B. Differential DNA mismatch repair underlies mutation rate variation across the human genome. *Nature* **521**, 81–84 (2015).

51. Zheng, C. L. *et al.* Transcription restores DNA repair to heterochromatin, determining regional mutation rates in cancer genomes. *Cell Rep.* **9**, 1228–1234 (2014).

52. Supek, F. & Lehner, B. Scales and mechanisms of somatic mutation rate variation across the human genome. *DNA Repair* **81**, 102647 (2019).

53. Nik-Zainal, S. *et al.* The life history of 21 breast cancers. *Cell* **149**, 994–1007 (2012).

54. Alexandrov, L. B. *et al.* Signatures of mutational processes in human cancer. *Nature* **500**, 415–421 (2013).

55. Mas-Ponte, D. & Supek, F. DNA mismatch repair promotes APOBEC3-mediated diffuse

hypermutation in human cancers. *Nat. Genet.* **52**, 958–968 (2020).

56. Bergstrom, E. N. *et al.* Mapping clustered mutations in cancer reveals APOBEC3 mutagenesis of ecDNA. *Nature* **602**, 510–517 (2022).

57. Sage, E. & Shikazono, N. Radiation-induced clustered DNA lesions: Repair and mutagenesis. *Free Radic. Biol. Med.* **107**, 125–135 (2017).

58. Sutherland, B. M., Bennett, P. V., Weinert, E., Sidorkina, O. & Laval, J. Frequencies and relative levels of clustered damages in DNA exposed to gamma rays in radioquenching vs. nonradioquenching conditions. *Environ. Mol. Mutagen.* **38**, 159–165 (2001).

59. Hada, M. & Georgakilas, A. G. Formation of clustered DNA damage after high-LET irradiation: a review. *J. Radiat. Res. (Tokyo)* **49**, 203–210 (2008).

60. Ravanat, J.-L. & Douki, T. UV and ionizing radiations induced DNA damage, differences and similarities. *Radiat. Phys. Chem.* **128**, 92–102 (2016).

61. Puck, T. T. & Marcus, P. I. Action of x-rays on mammalian cells. *J. Exp. Med.* **103**, 653–666 (1956).

62. Li, H. & Durbin, R. Fast and accurate short read alignment with Burrows-Wheeler transform. *Bioinformatics* **25**, 1754–1760 (2009).

63. McKenna, A. *et al.* The Genome Analysis Toolkit: a MapReduce framework for analyzing next-generation DNA sequencing data. *Genome Res.* **20**, 1297–1303 (2010).

64. Kim, S. *et al.* Strelka2: fast and accurate calling of germline and somatic variants. *Nat. Methods* **15**, 591–594 (2018).

65. Wang, K., Li, M. & Hakonarson, H. ANNOVAR: functional annotation of genetic variants from high-throughput sequencing data. *Nucleic Acids Res.* **38**, e164 (2010).

66. Karczewski, K. J. *et al.* The mutational constraint spectrum quantified from variation in 141,456 humans. *Nature* **581**, 434–443 (2020).

67. Chen, X. *et al.* Manta: rapid detection of structural variants and indels for germline and cancer sequencing applications. *Bioinforma. Oxf. Engl.* **32**, 1220–1222 (2016).

68. Roberts, S. A. *et al.* Clustered Mutations in Yeast and in Human Cancers Can Arise from Damaged Long Single-Strand DNA Regions. *Mol. Cell* **46**, 424–435 (2012).

69. Islam, S. M. A. *et al.* Uncovering novel mutational signatures by de novo extraction with SigProfilerExtractor. 2020.12.13.422570 (2022) doi:10.1101/2020.12.13.422570.

70. Schumann, F. *et al.* SigsPack, a package for cancer mutational signatures. *BMC Bioinformatics* **20**, 450 (2019).

71. Di Tommaso, P. *et al.* Nextflow enables reproducible computational workflows. *Nat. Biotechnol.* **35**, 316–319 (2017).

72. Obenchain, V. *et al.* VariantAnnotation: a Bioconductor package for exploration and annotation of genetic variants. *Bioinforma. Oxf. Engl.* **30**, 2076–2078 (2014).

73. Blokzijl, F., Janssen, R., van Boxtel, R. & Cuppen, E. MutationalPatterns: comprehensive genome-wide analysis of mutational processes. *Genome Med.* **10**, 33 (2018).

Supplementary figures

810
811

Supp. Figure 1. Sample attribution to a mutational signature. A signature was considered to
be generated by a condition (basically a mutagen or a gene knock-out) if at least 80% of the
total number of mutations detected in the condition sample are attributed to the signature. Note
that this is different than the activity of a signature that can be detected in one particular sample.
This was computed independently for point mutations (A) and for indels (B). Samples were
coloured according to the corresponding study: blue are Zou *et al.* knock-out samples[37], orange
are Kucab *et al.* mutagenesis samples[31] and yellow are the samples from this study.
819

Supp. Figure 2. Extracted SNV spectra. Mutational spectrum of the 7 SNV signatures
identified in the pooled-data, from the SigProfiler analysis. Each signature corresponds to the
distribution of point mutations into one of the 96 possible substitution categories taking a
trinucleotide context: 6 possible point mutations (taking into account base complementarity, *i.e.*
A=T; G=C) at the target position and 4 at each surrounding genomic location = 6*4*4=96
combinations.

Supp. Figure 3. Structural variation lengths. Distribution of the length of structural variations,
classified into the 4 main categories, *i.e.* large deletions, large insertions, duplications and
inversions. Length values are shown in the $\log_{10}$ scale. Rows represented the three different cell
lines, A549, HAP1 and MCF7. *P*-values shown in the plots were computed using the ggpubr R
package (t-test mean comparison). We observed a particular trend: proton radiation provokes
the smallest deletion and inversion events, followed by helium, and finally controls.
836

Supp. Figure 4. Regional mutation rates analysis, comparing negative binomial regression coefficient distribution. A negative binomial regression was computed for each cell line/condition/clone, in order to detect association between number of mutations (mutation rate) and genomic region, corrected by trinucleotide context in the case of SNVs. A similar approach than previously developed[55] was applied. The regression was applied to 3 different types of genomic components: replication time (A: SNVs, B: SVs), methylation (C: SNVs, D: SVs), and gene expression (E: SNVs, F: SVs). See methods for details on regional bins computations.

848
849

Supp. Figure 5. Correlation between SNV signatures and indel signature activities. For each sample we computed, for each SNV signature and each indel signature, the proportion of mutations attributed to that particular signature. The plot shows the correlation between each pair of indel signature activities and SNV signature activities. In other words, a high value would mean that the corresponding indel and SNV signatures are both highly active in the same samples, whereas a low value would mean they are both slightly active in the same samples.