Coronaviruses using different strategies to antagonize antiviral responses and pyroptosis

Xinyu Fu[1,7], Yang Yang[2,7], Weilv Xu[1,3,4,7], Danyue Li[1,3,4,7], Xinyue Li[1,3,4], Nan Chen[1,3,4], Qian Lv[1,3,4], Yuhua Shi[1,3,4], Xiaoliang Li [1,4,5,*] Fushan Shi,[1,3,4,6,*]

[1] Department of Veterinary Medicine, College of Animal Sciences, Zhejiang University, Hangzhou 310058, Zhejiang, China

[2] Key Laboratory of Applied Technology on Green-Eco-Healthy Animal Husbandry of Zhejiang Province, Zhejiang Provincial Engineering Research Center for Animal Health Diagnostics & Advanced Technology, Zhejiang International Science and Technology Cooperation Base for Veterinary Medicine and Health Management, China-Australia Joint Laboratory for Animal Health Big Data Analytics, College of Animal Science and Technology & College of Veterinary Medicine of Zhejiang A&F University, Hangzhou 310058, Zhejiang, China

[3] MOA Key Laboratory of Animal Virology, Center for Veterinary Sciences, Zhejiang University, Hangzhou 310058, Zhejiang, China

[4] Institute of Preventive Veterinary Medicine, Zhejiang Provincial Key Laboratory of Preventive Veterinary Medicine, Zhejiang University, Hangzhou 310058, Zhejiang, China

[5] Hainan Institute of Zhejiang University, Sanya 572025, Hainan, China

[6] Veterinary Teaching Hospital, Center for Veterinary Sciences, Zhejiang University, Hangzhou 310058, Zhejiang, China

[7] These authors contributed equally to this work

[*] Corresponding author: Xiaoliang Li, E-mail: xlli@zju.edu.cn; Fushan Shi, E-mail: sfs@zju.edu.cn

Abstract

Viral infection triggers inflammasome-mediated caspase-1 activation. However, less is known about how viruses use the active caspase-1 to evade host immune response. Here, we use porcine epidemic diarrhea virus (PEDV) as a model of coronaviruses (CoVs) to illustrate the sophisticated regulation of CoVs to counteract IFN-I signaling and pyroptosis. We show that PEDV infection stabilizes caspase-1 expression via papain-like protease PLP2's deubiquitinase activity and the enhanced stabilization of caspase-1 disrupts IFN-I signaling by cleaving RIG-I at D189 residue. Meanwhile, PLP2 can degrade GSDMD-p30 by removing its K27-linked ubiquitin chain at K275 to restrain pyroptosis. Papain-like proteases from other genera of CoVs (PDCoV and SARS-CoV-2) have the similar activity to degrade GSDMD-p30. We further demonstrate that SARS-CoV-2 N protein induced NLRP3 inflammasome activation also uses the active caspase-1 to counter IFN-I signaling by cleaving RIG-I. Therefore, our work unravels a novel antagonistic mechanism employed by CoVs to evade host antiviral response.

Keywords: Coronavirus; Papain-like protease; Antiviral immunity; GSDMD; Type I interferon signaling

Introduction

Coronaviruses which belong to order Nidovirales, family Coronaviridae, and subfamily Orthocoronavirinae, are a group of viruses packaged in envelop covered with spikes and contain a linear, single and positive-stranded RNA[1]. Coronaviruses

are subdivided into four genera, including α, β, γ, and δ. As a member of α genus,

PEDV infection was characterized by vomiting, severe diarrhea, dehydration and high

mortality in suckling piglets[2]. The viral genome of PEDV contains a full-length

genome of 28 kb and encodes a total of seven open reading frames (ORFs)[3]. Like

other CoVs, PEDV utilizes its own proteases, including papain-like protease (PLpro)

and main protease to cleave the polyprotein. The papain-like protease of PEDV is

encoded by the largest nonstructural protein Nsp3, which contains two papain-like

protease domains (PLP1 and PLP2 or PLpro)[4]. In general, α coronaviruses and clade

A of β coronaviruses harbor PLP1 and PLP2, and other coronaviruses have one

functional PLpro only[5]. Studies have shown that CoVs PLP2 (PLpro) act as a viral

deubiquitinase to negatively regulate type I IFN signaling pathway[4,6]. For example,

mouse hepatitis virus (MHV) PLP2 can bind to and deubiquitinate IRF3 to prevent its

nuclear translocation[7]. Transmissible gastroenteritis virus (TGEV) PLP2 is able to

inhibit the degradation of IκBα by decreasing its ubiquitination, resulting in the

suppression of NF-κB signaling[8]. As to PEDV PLP2, it has been reported that PLP2

deubiquitinates RIG-I and STING to block the IFN signaling pathway[4]. The

catalytically dead mutants of PLP2 (C1729A, H1888A and D1901A) can abrogate

their deubiquitinase activity and fail to inhibit PEDV-induced IFN-β production.

Caspase-1 is a cysteine protein hydrolase, which participates in cell death and

inflammatory response[9]. Caspase-1 exists in the form of proteasome (pro-caspase-1)

at static state and acts as an enzyme after activation, which relies on the activation of

different inflammasomes[10]. Interestingly, high expression of pro-caspase-1 can also

result in its auto-activation in the absence of a ligand[11-13]. Pyroptosis is an inflammatory caspases-dependent, pro-inflammatory programmed cell death characterized by losing of cell membrane integrity, pore forming and swelling and rupture of the cells[14]. The canonical pathway of pyroptosis is mediated by caspase-1[15,16]. When host cells' pattern-recognition receptors (PRRs) are stimulated by pathogen-associated molecular patterns (PAMPs) or damage-associated molecular patterns (DAMPs), inflammasomes will assemble automatically and activate themselves, leading to the activation of caspase-1. Researches have demonstrated that NLRP3 inflammasome can be activated by ZIKA[17], SARS-CoV-2[18] and MERS-CoV[19,20], and AIM2 inflammasome can be activated by SARS-CoV-2[21] while NLRP1 inflammasome can be activated by human rhinovirus (HRV) and picornaviruses[22]. Activated caspase-1 cleaves GSDMD to produce N-terminal GSDMD-p30 fragments with perforating activity on cell membrane, which leads to pyroptosis, and it also cleaves pro-IL-1β to produce IL-1β to improve antiviral capacity[15,16].

What is clear is that, inflammasomes-mediated caspase-1 activation and pyroptosis induced by viruses will lead to enhancement of cell antiviral capability and destruction of cell integrity, and these changes are detrimental to virus replication, especially at the early stage of viral infection. So what is the purpose of viral activation of caspase-1 as well as pyroptosis? Here, we report that PEDV infection increases the stabilization of caspase-1 by removing its K11-linked ubiquitin chains at K134 via its papain-like protease PLP2. Consequently, caspase-1 targets to RIG-I for

cleavage at D189 residue which leads to decreased IFN-I signaling and enhanced

PEDV replication. Meanwhile, to assure the integrity of host cells which might be

broken by GSDMD-p30 cleaved by caspase-1, PLP2 can degrade GSDMD-p30 by

removing its K27-linked ubiquitin chain at K275 residue to restrain pyroptosis. We

further demonstrate that papain-like proteases from other genera of CoVs, such as

PDCoV and SARS-CoV-2, have the similar activity to degrade GSDMD-p30.

Importantly, we confirm that SARS-CoV-2 N protein induced NLRP3 inflammasome

activation also uses the active caspase-1 to counter IFN-I signaling. Therefore, our

findings uncover a distinctive feature of papain-like protease in antagonizing antiviral

responses that might serve as a target for CoVs treatment in the future.

Results

PEDV infection elevates the expression of caspase-1

In order to investigate the effect of PEDV on porcine caspase-1, endogenous

caspase-1 expression was analyzed by immunoblotting in intestinal tissues of

mock-infected and PEDV naturally-infected piglets. The results showed that

expression of caspase-1 was increased in the PEDV-infected piglets (Fig. 1A).

qRT-PCR showed that *caspase-1* mRNA abundance was not altered in IPEC-J2 cells

upon PEDV infection (Fig. 1B). Immunoblotting also confirmed that PEDV infection

promoted the amount of caspase-1 in IPEC-J2 cells and CHX treatment further

elevated caspase-1 expression (Fig. 1C), suggesting that the increased caspase-1

expression was not caused by the transcription of mRNA and synthesis of new

proteins. In addition, Vero cells were transfected with plasmids encoding caspase-1

and then infected with PEDV. We found that the increased caspase-1 by PEDV

infection was both MOI and time dependent (Fig. 1D and E). Since the high

expression of caspsae-1 causes auto-activation and produces active caspase-1 p20[11-13],

which can lead to the cleavage of GSDMD, we investigated the relationship between

PEDV infection and pyroptosis. Vero cells were co-transfected with GSDMD-FL (full

length) and caspase-1 and then infected with PEDV. Caspase-1 expression and

cleavage of GSDMD increased progressively in both MOI and time dependent

manner during infection (Fig. 1F and G). Interestingly, PEDV infection resulted in

declines in LDH release compared to mock-infected group, despite the increased

GSDMD cleavage (Fig. 1H and I). These results indicate that PEDV infection

promotes the amount of caspase-1.

The papain-like protease 2 of PEDV inhibits the proteasomal degradation of

caspase-1

The increased expression of caspase-1 induces GSDMD cleavage but inhibits

pyroptosis. Therefore, we speculated that some kind of viral proteins affected this

process. Vero cells transfected with GSDMD-FL were infected with PEDV, following

performed with coimmunoprecipitation (Co-IP) and mass spectrometry (MS) analysis.

The database search revealed that the papain-like protease 2 (PLP2) of PEDV binds to

GSDMD-FL during infection (Supplementary Fig. 1A). The Co-IP assay also

confirmed that PEDV PLP2 interacted with GSDMD-FL (Supplementary Fig. 1B).

Since it is known that GSDMD-FL and caspase-1 are interacted[23], we further

examined whether PLP2 interacts with caspase-1. HEK293T cells were co-transfected

with FLAG-tagged caspase-1 and MYC-tagged PLP2, following treated with VX765,

an inhibitor of caspase-1, to curb caspase-1 activation. Co-IP assay showed that PLP2

interacted with caspase-1 (Supplementary Fig. 1C). Indirect immunofluorescence

further demonstrated that GSDMD, caspase-1 and PEDV PLP2 were colocalized in

the cytoplasm (Supplementary Fig. 1D). Thus, PLP2 interacts with both caspase-1 and

GSDMD.

Given that high expression of caspase-1 leads to auto-activation, caspase-1-C285A

(caspase-1 mutant, which lacks its protease activity) and PLP2 were co-transfected

into HEK293T cells. As shown in Fig. 2A, we observed an increased protein amount

of caspase-1-C285A in the presence of PLP2. In addition, co-transfection of PLP2,

GSDMD-FL and caspase-1 in HEK293T also revealed that PLP2 caused the

accumulation of caspase-1 but not GSDMD-FL (Supplementary Fig. 2A). We next

measured caspase-1 degradation by a cycloheximide (CHX) chase assay to determine

whether PLP2 could stabilize caspase-1. The results demonstrated that PEDV PLP2

could stabilize caspase-1 but not GSDMD-FL by delaying its degradation (Fig. 2B

and Supplementary Fig. 2B). Ubiquitination is a key signal for proteasomal

degradation, and PLP2 has the deubiquitinase activity. To identify whether the

deubiquitinase activity of PLP2 is required for the stabilization of caspase-1, we

generated the catalytic mutants (C113A, H272A, D285A) of PLP2 and co-transfected

them with caspase-1-C285A. As expected, all of the three mutants lost the ability to

stabilize caspase-1 (Fig. 2C). Furthermore, we co-expressed the wild type PLP2 or its mutants with HA-tagged ubiquitin (HA-Ub) and FLAG-tagged caspase-1 in HEK293T cells. Compared to the empty vector, wild type PLP2 substantially inhibited ubiquitination of caspase-1 while the three mutants had no effect on it (Fig. 2D), suggesting that the deubiquitinase activity of PLP2 is indispensable for the stabilization of caspase-1.

It has been reported that the K11-linked ubiquitination of human caspase-1 at Lys-134 is a way to induce its degradation[24]. To determine whether K134 residue of porcine-derived caspase-1 is also critical for its ubiquitination, we constructed porcine-derived caspase-1-K134R mutant in which the lysine (K) residue was replaced with arginine (R). As shown in Fig. 2E, wild type caspase-1 was degraded by K11 in a dose-dependent manner, while caspase-1-K134R could not be degraded by K11. Moreover, we found that caspase-1-K134R mutant exhibited a dramatic decrease in K11-linked ubiquitination (Supplementary Fig. 2C). To further confirm the role of PLP2 in this process, we co-expressed PLP2 with caspase-1 and K11 ubiquitin, the results demonstrated that the degradation effect of K11 on caspase-1 was abrogated by PLP2 (Fig. 2F). Immunoblotting also showed that PLP2 stabilized caspase-1 but not caspase-1-K134R (Fig. 2G). Additionally, PEDV infection decreased K11-linked poly-ubiquitination of caspase-1, further fortifying the above findings (Fig. 2H). Taken together, these results indicate that PLP2 of PEDV stabilizes porcine caspase-1 by removing K11-linked ubiquitin chains at Lys134.

Caspase-1 attenuates IFN-I signaling during PEDV infection

It has been reported that caspase-1 can negatively regulate IFN-I signaling by cleaving cGAS during DNA virus infection[25]. As an RNA virus, PEDV activates IFN-I signaling through the RLRs pathway. Since IFN-I plays a vital role in antiviral process, we wondered whether the stabilized caspase-1 can affect it. We initially found that overexpression of caspase-1 in IPEC-J2 cells attenuated the *IFN-α*, *IFN-β*, *ISG15* and *OAS1* mRNA levels induced by poly (I:C) treatment (Fig. 3A-D). In addition, caspase-1 inhibitor VX765 reversed the attenuated IFN-I signaling during poly (I:C) treatment (Fig. 3E and F). Moreover, knockout of caspase-1 in THP-1 cells further promoted the expression of *IFN-β* and *ISG15* mRNA induced by poly (I:C) (Fig. 3G and H). Same results were observed for PLP2 in IPEC-J2 cells (Fig. 3I and J). Thus, we determined that both caspase-1 and PLP2 were antagonists of IFN-I. Next, the above-mentioned findings were further verified in PEDV-infected IPEC-J2 cells, which were consistent with our observations (Fig. 3 K-P). These results indicate that PEDV-induced IFN-I signaling was counteracted by the stabilized caspase-1.

The role of caspase-1 and PLP2 in viral replication was further analyzed considering the vital antiviral effect of IFN-I. PLP2 was transfected into IPEC-J2 cells and then infected with PEDV. Consistent with its antagonistic effect on IFN-I, the results showed that PLP2 could promote PEDV replication, which was also confirmed by immunoblotting (Fig. 3Q). Since the high expression of caspase-1 leads to autoprocessing, we transfected caspase-1 with different doses and then treated with PEDV infection. As shown in Fig. 3R, all doses of caspase-1 could promote PEDV

replication, while the effect of high concentration of caspase-1 was attenuated, which

might relate to pyroptosis caused by caspase-1 activation. Moreover, the use of

VX765 inhibited viral replication co-treated with necrosulfonamide (NSA), a specific

GSDMD inhibitor (Fig. 3S). Together, these data suggest that the enhanced caspase-1

stabilized by PEDV infection attenuates type I IFN signaling and thus benefits to viral

replication.

Cleavage of RIG-I by caspase-1 ensures PEDV replication

As caspase-1 is vital to suppress type I IFN during PEDV infection, we hypothesized

that caspase-1 exerted its effect on RIG-I. Therefore, we analyzed whether caspase-1

could directly interact with RIG-I. HEK293T cells were then transfected with

MYC-tagged RIG-I and FLAG-tagged caspase-1, Co-IP assay showed that RIG-I

interacted with caspase-1 (Fig. 4A). Interestingly, we found that RIG-I was

diminished after co-transfected with caspase-1 (Fig. 4B). Thus, we co-transfected

caspase-1 or its mutant with RIG-I at different doses in HEK293T, followed with

immunoblotting. As shown in Fig. 4C, a clear band with a molecular weight around

25-35 kDa was observed, while the caspase-1-C285A (caspase-1 mutant) impaired the

cleavage. Because RIG-I was N-terminal MYC-tagged, we speculated that caspase-1

cut RIG-I at N-terminus. We therefore generated mutants of RIG-I at different sites to

identify the exact cleavage site of RIG-I by caspase-1. RIG-I or its mutants (D163A,

D189A, D194A, D199/202/203A and D209A) were co-transfected with caspase-1 in

HEK293T cells. As shown in Fig. 4D and Supplementary Fig. 3A, wild-type RIG-I,

D163A, D194A, D199/202/203A and D209A could still be cleaved by caspase-1

successfully, while RIG-I-D189A could not be cleaved. These results imply that RIG-I

was a cleaved target of caspase-1 at D189. Moreover, we investigated the relationship

between human derived caspase-1 and RIG-I. As shown in Supplementary Fig. 3B, a

band with a molecular weight around 25-35 kDa was observed (lane 2). Based on the

multiple-sequence alignment of RIG-Is (Supplementary Fig. 3C), different mutants,

including single or double mutants of human RIG-I (h-RIG-I) were generated and

co-expressed with empty vector or h-caspase-1. The results showed that the cleaved

fragments from all of the mutants were reduced compared to the wild type RIG-I

(Supplementary Fig. 3B). We next generated a triple mutant of RIG-I

(h-RIG-I-D163/194/234A) and transfected it with caspase-1 in HEK293T cells.

Supplementary Fig. 3D revealed that the cleaved band between 25-35 kDa was further

weakened, suggesting that the D163, D194 and D234 residues of RIG-I might be the

cleavage sites of human derived caspase-1.

To further investigate the effects of RIG-I cleavage on viral replication and type I

IFN signaling, Vero cells were transfected with empty vector (EV), RIG-I or

co-transfected with caspase-1 and RIG-I, followed by PEDV infection. As shown in

Fig. 4E, RIG-I significantly attenuated viral replication, while the co-transfected

group reversed this effect. We then generated a RIG-I mutant, RIG-I-D189A, and

transfected into IPEC-J2 cells and then infected with PEDV. As expected, wild type

RIG-I facilitated *IFN-β* and *ISG15* production and inhibited PEDV replication, and

the RIG-I-D189A mutant further enhanced IFN-I signaling and impaired viral

replication (Fig. 4F-H). Next, we evaluated the function of RIG-I-1-189aa and

RIG-I-190-943aa in activating the innate immune response as well as in viral

replication. We found that wild-type RIG-I promoted IFN induction and inhibited

PEDV replication, while RIG-I-1-189aa or RIG-I-190-943aa potently reversed IFN-I

production and restored viral reproduction (Fig. 4I-K). Taken together, these results

collectively demonstrate that caspase-1 targets the D189 residue of porcine RIG-I for

cleavage and the cleaved fragments lose their function to inhibit PEDV replication.

PLP2 degrades GSDMD-p30 by removing the K27-linked ubiquitin chains at

K275 residue to inhibit pyroptosis

The activation of caspase-1 can lead to GSDMD cleavage and thus cause pyroptotic

cell death. However, according to the above results, PEDV infection did not cause

substantial pyroptosis at early stage (Fig. 1H and I). To determine whether PLP2 also

targets GSDMD and regulates pyroptosis, HEK293T cells were initially

co-transfected with GSDMD-FL and caspase-1 for 6 h, and then PLP2 was transfected

in a dose-dependent manner. The immunoblotting results showed that the cleaved

fragment of GSDMD (GSDMD-p30) was decreased in the presence of PLP2 (Fig.

5A). Meanwhile, we repeatedly observed a reduced protein level of GSDMD-p30

after overexpression of PLP2, but had no variation on GSDMD-FL (Fig. 5B and

Supplementary Fig. 4A). The LDH release assay and the propidium iodide (PI)

staining also confirmed that PLP2 had an inhibitory effect on pyroptotic cell death

induced by GSDMD-p30 (Fig. 5C and Supplementary Fig. 4B). As shown in Fig. 5D,

PLP2 had a stronger interaction with GSDMD-p30 than GSDMD-FL, which further demonstrated the complex relationship between GSDMD-p30 and PLP2. In order to evaluate the effect of PLP2 on GSDMD-p30 more directly, we co-transfected GFP-tagged PLP2 and FLAG-tagged GSDMD-p30 in HEK293T cells. GSDMD-p30 was found to localize mainly in the cytoplasm, formed specks and ultimately resulted in rupture of the membrane (Fig. 5E top row). In the presence of PLP2, GSDMD-p30 was evenly dispersed in cytoplasm, failed to form pyroptotic specks (Fig. 5E bottom row). Taken together, we reveal that PLP2 degrades GSDMD-p30 and thus blocks pyroptosis.

To determine which degradation system is responsible for the degradation of GSDMD-p30, we examined the effect of the proteasome inhibitor MG132, the autophagy inhibitor 3MA and CQ on GSDMD-p30 degradation in the presence of PLP2. The results showed that PLP2-mediated degradation of GSDMD-p30 was rescued by MG132, which meant that GSDMD-p30 degradation was corresponded with proteasome pathway (Fig. 6A). We further revealed that all the catalytic mutants of PLP2 (C113A, H272A, D285A) lost the ability to degrade GSDMD-p30 which was also confirmed by the LDH release assay (Fig.6B and C), suggesting that the deubiquitinase activity of PLP2 is essential for the GSDMD-p30 degradation. To determine the lysine residues of GSDMD-p30 to which ubiquitin is attached, we used the UbPred program (http://www.ubpred.org), to predict the potential ubiquitination sites of porcine GSDMD-p30. This analysis revealed three putative lysine residues: K103, K177 and K275. We then generated GSDMD-p30-K103R, K177R and K275R

mutants in which each of these lysine residues was replaced with arginine (R). We

co-transfected the wild type porcine GSDMD-p30 and three of its mutants with

HA-tagged ubiquitin, followed with immunoblotting. As shown in Fig. 6D,

co-transfection of HA-Ub failed to promote GSDMD-p30-K275R mutant

ubiquitination, suggesting that Lys275 residue might be the ubiquitination site of

GSDMD-p30. Next, we co-transfected HA-tagged ubiquitin mutants (with only one of

the seven lysine residues retained as lysine, while the other six replaced with arginine)

with GSDMD-p30 in HEK293T cells. The LDH release assay revealed that

K27-linked ubiquitin mutant markedly elevated the LDH level (Fig. 6E), which was

consistent with our previous study[26]. Ubiquitination assay further confirmed that

K27-linked ubiquitin promoted the ubiquitination of GSDMD-p30 (Fig. 6F, lane 2

and 4) and K275R mutant displayed a significant reduction of K27-linked

ubiquitination (Fig. 6F, lane 4 and 5). Moreover, we co-expressed the wild type PLP2

or its mutants with HA-tagged K27-linked ubiquitin and FLAG-tagged GSDMD-p30

and then performed Co-IP assay. As expected, wild type PLP2 dramatically reduced

K27-linked ubiquitination of GSDMD-p30, while neither of its mutants performed the

same effect (Fig. 6G). To further confirm the role of K275 residue of GSDMD, we

designed three siRNAs targeting porcine GSDMD. The efficiency of GSDMD

knockdown in IPEC-J2 cells was shown in Supplementary Fig. 4C, and we selected

si-GSDMD#1 for further studies. IPEC-J2 cells were transfected with GSDMD-FL or

GSDMD-FL-K275R upon siRNA-mediated knockdown of GSDMD, followed by

PEDV infection. As shown in Supplementary Fig. 4D, GSDMD-FL-K275R facilitated

the viral replication compared to wild type GSDMD-FL, the LDH assay further confirmed this (Supplementary Fig. 4E). Taken together, our results demonstrate that PLP2 is able to degrade GSDMD-p30 by removing the K27-linked ubiquitin chain at K275 residue of GSDMD-p30 and thus abrogate pyroptosis.

CoVs using different strategies to disrupt antiviral response and inhibit pyroptosis

Next, we tested whether papain-like proteases in other genus of CoVs have the same effects on GSDMD-p30 and caspase-1. PEDV-PLP2, SARS-CoV-2-PLpro and PDCoV-PLpro were generated and then transfected into HEK293T cells with GSDMD-p30. As shown in Fig. 7A and B, all these papain-like proteases could inhibit LDH release and degrade GSDMD-p30, which revealed their ability in abrogating pyroptosis. Furthermore, we examined the role of these viral proteins in stabilizing caspase-1 and found that only PEDV PLP2 could promote caspase-1 expression (Fig. 7C). It is possible that PDCoV and SARS-CoV-2 use different ways to activate caspase-1. To evaluate the effect of these viral proteins on PEDV replication, IPEC-J2 cells were transfected with empty vector or these papain-like proteases, and then infected with PEDV. All these papain-like proteases were able to promote viral replication (Fig. 7D). Collectively, GSDMD-p30 was attenuated by papain-like proteases of CoVs, which led to higher viral replication.

It has been reported that the nucleocapsid protein (N) of SARS-CoV-2 activates the NLRP3 inflammasome, thus we hypothesized that NLRP3 inflammasome-mediated

caspase-1 activation could also antagonize IFN production. In that case, we generated THP-1 cell lines stably expressing FLAG-EV-Lentivirus (negative control) or FLAG-N-Lentivirus (Lentivirus carrying the SARS-CoV-2 N gene) (Fig. 7E). Consistent with previous studies[18], FLAG-N-Lentivirus elevated the secretion of IL-1β, suggesting that SARS-CoV-2 N can activate NLRP3 inflammasome (Fig. 7F). To evaluate the role of SARS-CoV-2 N in IFN production, the differentiated macrophages were then stimulated with poly (I:C) and we found that the endogenous RIG-I was significantly reduced in the presence of SARS-CoV-2 N (Fig. 7G). As shown in Fig. 7H and I, the production of IL-1β was elevated while the IFN-β was dramatically reduced by SARS-CoV-2 N. Moreover, qRT-PCR analyses revealed that the expression of *IFN-β* and ISG15 mRNA level were notably repressed by SARS-CoV-2 N (Fig. 7J and K), indicating that the activated caspase-1 targeted RIG-I to abrogate IFN-I signaling. LDH release assay also confirmed the inhibitory effect of SARS-CoV-2 N on GSDMD-mediated pyroptosis (Fig. 7L). Taken together, we unravel a novel antagonistic mechanism employed by different viral proteins from CoVs and thus evade host IFN signaling and pyroptosis (Fig.8).

Discussion

A variety of viruses have been reported to activate inflammasomes. The inflammasome operates as a platform for caspase-1 activation, resulting in caspase-1-dependent IL-1β release and subsequent pyroptosis. This, in turn, increases the expression of antiviral genes to control invading pathogens[27]. As a countermeasure,

viruses have evolved strategies to antagonize host immune response. Previous study

has demonstrated that DNA virus-induced inflammasome-mediated active caspase-1

cleaves human cGAS at D140 and D157 to dampen IFN production[25]. In the present

study, we demonstrated the sophisticated regulation of RNA virus to counteract IFN-I

signaling and pyroptosis using active caspase-1 and papain-like protease. We found

that CoVs-induced active caspase-1 disrupts IFN-I signaling by cleaving RIG-I to

produce inactive fragments. Meanwhile, papain-like protease of CoVs can degrade

GSDMD-p30 by removing its K27-linked ubiquitin chain at K275 residue to restrain

pyroptosis. Therefore, our work unravels a novel antagonistic mechanism employed

by CoVs to evade host antiviral response.

To date, most viruses induced caspase-1 activation is mediated by inflammasomes.

EV71 3D and ZIKV NS5 activate the NLRP3 inflammasome by interacting with

NACHT and the LRR domain of NLRP3[17,28,29]. NS1 of ZIKA also enhances NLRP3

inflammasome activation to facilitate its infection[24]. The influenza A virus (IAV)

non-structural protein PB1-F2 contributes to severe pathophysiology through

triggering NLRP3-dependent caspase-1 activation[30]. Except for NLRP3, viruses can

also induce other inflammasomes activation. Human rhinovirus (HRV) 3C protease

directly cleaves human NLRP1 between the Glu130 and Gly131 junction, which

liberates the activating C-terminal fragment and subsequently promoting NLRP1

inflammasome activation[31]. NLRP9b recognizes short double-stranded RNA stretches

via RNA helicase Dhx9 and forms inflammasome complexes with ASC and

pro-caspase-1 to promote caspase-1-dependent IL-1β release and pyroptosis[32]. Herpes

simplex virus 1 (HSV1) can induce AIM2-, pyrin- and ZBP1-mediated caspase-1

activation, cytokine release and cell death[33]. For coronaviruses, studies have shown

that MERS-CoV and at least three different proteins of SARS-CoV-2 can induce

NLRP3 inflammasome activation and caspase-1-mediated subsequent

pyroptosis[18,20,34,35]. Evidence also demonstrated that circulating monocytes from

COVID-19 patients show signs of AIM2 inflammasome activation and

caspase-1-meidated GSDMD cleavage and pyroptosis[21]. Unlike the reported manner

of caspase-1 activation, our results suggest that PEDV induces porcine caspase-1

activation via its papain-like protease PLP2 through elevating pro-caspase-1

expression by removing its K11-linked ubiquitin chains at K134 residue. Studies have

demonstrated that accumulated pro-caspase-1 lead to autoprocessing within the

catalytic domain[11-13]. Consequently, active caspase-1 targets to porcine RIG-I for

cleavage at D189 residue leading to decreased IFN-I signaling and enhanced PEDV

replication. Interestingly, PLPro from SARS-CoV-2 and PDCoV did not use this way

to activate caspase-1, as they can trigger NLRP3- or AIM2-mediated caspase-1

activation. Similarly, inflammasome-mediated caspase-1 activation also cleaved

human RIG-I to abrogate IFN-I signaling pathway. Therefore, our study suggests that

the main purpose of virus-induced inflammasome-mediated caspase-1 activation is to

antagonize IFN-I signaling. A newly published study further indicated that

SARS-CoV-2 infection induced high expression of pro-caspase-4/11, which may

result in its auto-activation in the absence of a ligand[36]. Thus, our results uncover a

new and different mechanism of CoVs inducing caspase-1 activation.

Activation of inflammasome is often vital in the host antiviral immune

response[35,37-39]. It is obvious that triggering of pyroptosis and inflammasomes

activation is detrimental to the replication and survival of viruses. Thus, viruses have

also developed strategies to antagonize extensive inflammasome activation and

pyroptosis, especially at the early stage of viral infection. The human papillomavirus

(HPV) oncoprotein E7 promotes the TRIM21-mediated K33-linked ubiquitination of

the IFI16 inflammasome for degradation to inhibit pyroptosis. African swine fever

virus (ASFV) pS273R cleaves GSDMD at G107-A108 to generate a shorter

N-terminal of GSDMD (GSDMD-N$_{1-107}$) which is not capable of triggering pyroptosis.

The accessory protein (PB1-F2) of H5N1 and H3N2 influenza A viruses (IAV) can

bind to the pyrin and LRR domains of NLRP3 and prevent NEK7 binding to stabilize

the auto-repressed and closed conformation of NLRP3. 2A and 3C proteases of EV71

can cleave NLRP3 directly to inhibit the activation of inflammasome, while the EV71

3C-like protease can also cleave GSDMD to block the pyroptosis pathway. Our recent

research also proved that 3C-like proteases Nsp5 of coronaviruses (PEDV, PDCoV,

SARS-CoV-2 and MERS-CoV) can cleave GSDMD at Q193 to produce two

fragments that are unable to trigger pyroptosis. SARS-CoV-2 N induces

pro-inflammatory cytokines through promoting the assembly and activation of the

NLRP3 inflammasome[18]. However, SARS-CoV-2 N protein is also capable of

binding to GSDMD directly to inhibit the GSDMD cleavage from caspase-1 to avoid

pyroptosis[40]. Our results demonstrated that SARS-CoV-2 N-induced NLRP3

inflammasome activation can antagonize IFN production through cleaving RIG-I by

activated caspase-1. In the present study, we further demonstrated that CoVs (PEDV,

PDCoV and SARS-CoV-2) can abrogate pyroptosis by degradation of GSDMD-p30

via their papain-like proteases. Therefore, we discovered a novel mechanism for CoVs

to antagonize pyroptosis.

In conclusion, we used PEDV as a model of CoVs to illustrate the sophisticated

regulation of CoVs to counteract IFN-I signaling and pyroptosis. For the first time, we

demonstrate the role of inflammasome dependent or independent caspase-1 activation

in antagonizing IFN-I signaling. Furthermore, we show that papain-like protease of

CoVs can inhibit pyroptosis by degradation of GSDMD-p30 through proteasome

pathway. Thus, our present study unveils a novel antagonistic mechanism by which

CoVs manipulates the cross-talk of IFN signaling and pyroptosis, which might

indicate a framework for design of anti-CoVs therapies.

Methods

Reagents and antibodies

Anti-HA (3724) and anti-Caspase-1 (2225) antibodies were purchased from Cell

Signaling Technology. Anti-FLAG antibody (F1804), anti-MYC antibody (C3956)

and anti-FLAG magnetic beads (10004D) were obtained from Sigma. Anti-FLAG

antibody (rabbit source), Goat pAb to MS IgG (Chromeo 546, ab60316), Dnk pAb to

Rb IgG (Alexa Fluor 647, ab150075) fluorescent secondary antibodies were acquired

from Abcam. Anti-GAPDH antibody, HRP-labeled goat anti-rabbit IgG and goat

anti-mouse IgG were from Hangzhou Fudebio. Anti-GSDMDC1 antibody (sc-393581)

was purchased from Santa Cruz. Anti-RIG-I antibody (20566) was obtained from

proteintech. The anti-PEDV N monoclonal antibody was prepared in our laboratory as

previously described[41]. Efficient eukaryotic transfection reagent VigoFect was

obtained from Vigorous Biotechnology (Beijing), Lipo8000TM transfection reagent

was obtained from Beyotime Biotechnology while Lipofectamine 2000 transfection

reagent was obtained from Invitrogen.

Plasmids

Eukaryotic expression vectors used in this subject were saved in our laboratory. The

PLP2 fragment sequence was amplified from cDNA of PEDV strain ZJ15XS0101

(GenBank accession no. KX55SO0281) which was isolated by our collaborating

laboratories for generational preservation, while amplification of the porcine GSDMD

gene and the porcine caspase-1 gene used cDNA of IPEC-J2 as a templet. The PLP2

gene was cloned to PRK-MYC vector, and the porcine GSDMD gene was cloned onto

p3×FLAG-CMV vector. The porcine caspase-1 gene was cloned to PCMV-HA and

p3×FLAG-CMV vector, respectively. Porcine GSDMD-p30 fragment was amplified

from the p3×FLAG-N-GSDMD-FL plasmid and cloned onto p3×FLAG-CMV. PLP2,

caspase-1 and GSDMD-p30 enzyme active site mutants were constructed based on

the eukaryotic expression plasmids PRK-MYC-PEDV-PLP2, HA-PCMV-Caspase-1

and p3×FLAG-GSDMD-p30. The plasmids encoding SARS-CoV-2 PLPro, PDCoV

PLPro and SARS-CoV-2-N were synthesized by Tsingke Biotech. Primers used in the

plasmids construction are listed in table 1 in supplementary materials.

Cell culture and transfection

IPEC-J2 cells, Vero cells and HEK293T cells were cultured in Dulbecco's modified

Eagle's medium (DMEM) with added 10% FBS and 1% penicillin-streptomycin. Wild

type THP-1 and caspase-1 deficient THP-1 were cultured in PRIM 1640 medium

containing 10% FBS and 1% penicillin-streptomycin solution. The PEDV strain

ZJ15XS0101 (GenBank accession no. KX55SO0281) was isolated and stored in our

laboratory[42]. When HEK293T cells seeded in plates grow to about a density of

60%~80%, indicated plasmids were transfected to the cells by VigoFect according to

the manufacturer's operation guide. When IPEC-J2 and Vero seeded in plates grow to

approximately 70%~80%, they were transfected with corresponding plasmids using

Lipo8000 transfection reagent according to the manufacturer's introduction. Lipo2000

was applied in the transfection of poly(I:C) (Merck) to IPEC-J2 during the second

transfection period.

Viral infection

Viral infection was performed when the IPEC-J2 or Vero cells grow to the density

around 80%~90% or after the transfection of other indicated plasmids.

Lentivirus infection

SARS-COV-2-N over-expressing THP-1 cells were generated using lentiviral infection

technique. After cloning *SARS-COV-2-N* coding sequence to the

pLVX-IRES-Puro-3×Flag lentiviral vector plasmid, HEK293T cells were

co-transfected with pLVX-IRES-Puro-3×Flag-SARS-COV-2-N, pMD2G and

PSPAX2. Supernatant was collected after 48 h incubation for further lentiviral

concentration.

Inhibitors treatment

The caspase-1 specific inhibitor VX765 diluted to 20 μM was used after plasmid transfection, during viral infection or before poly(I:C) transfection. Protein synthesis inhibitor cycloheximide (CHX) diluted to 25 μg/ml was added at different time points before receiving samples. Fresh culture medium containing proteasome inhibitor MG132, autophagy initiation inhibitor 3MA or autophagy lysosome inhibitor CQ of appropriate concentration were added to different wells of cell-culturing plates 6 h before receiving samples.

Cytotoxicity assay

Cells transfected with corresponding plasmids for 24 h were applied to cytotoxicity test using CytoTox 96® Reagent (Promega) according to the manufacturer's manual. OD values were read at 492 nm on an enzyme marker (Thermo Scientific).

Immunoblotting

Total proteins were extracted from cells lysed by RIPA lysis buffer (Beyotime Biotechnology) supplemented with 1% Phenylmethanesulfonyl fluoride (PMSF) (Beyotime Biotechnology). After being separated on the 10% SDS-PAGE gel (Verde Biotechnology), protein stripes were transferred onto the polyvinylidene difluoride (PVDF) membranes (Bio-rad). Membranes were blocked in the blocking buffer (Beyotime Biotechnology), followed by incubation with primary antibodies and positioning with secondary antibodies. Chemiluminescent signals were captured by an ECL chemiluminescence imaging analysis system (Clinx Science Instruments).

Co-immunoprecipitation

Cells were lysed using IP lysis buffer (Beyotime Biotechnology) for 15 min at 4°C,

and then centrifuged at speed 10000 rpm at 4°C for 15 min. After being incubated

with the supernatant of the cell lysis solution in a flip mixer (Kylin-Bell) overnight,

the magnetic beads coated with FLAG-antibodies (sigma) were washed with

coIP-buffer for 5 times. Soaking in 1×loading buffer, magnetic beads absorbed with

target proteins were denatured for 10 min in a boiling water bath. After discarding

magnetic beads, the remaining protein samples were applied to subsequent

immunoblotting assay.

Confocal immunofluorescence assay

HEK293T cells were seeded on glass slides placed in 24-well plates. After being

transfected for 24 h, cells were fixed with Immunol Staining Fix Solution (Beyotime),

permeabilized with Immunostaining Permeabilization Solution with Saponin

(Beyotime), blocked with QuickBlock Blocking Buffer for Immunol Staining

(Beyotime), and then incubated with indicated antibodies. The cells were observed

under a laser confocal microscope (Olympus).

ELISA

Supernatants collected from transfected cells were applied to the detection of IL-1β

and IFN-β according to the manufacturer's instructions. Each trial group was

conducted independently for three times.

Total RNA extraction and reverse transcription

Add RNA-easy Isolation Reagent (Vazyme Biotechnology) to cells to gather lysis

solution containing cells. Follow the manufacturer's introduction to extract the RNA.

After measuring the product RNA concentration, use the HiScript III RT SuperMix for

qPCR (+gDNA wiper) (Vazyme Biotechnology) to reverse transcribe RNA to cDNA.

Quantitative real-time polymerase chain reaction (qRT-PCR)

qRT-PCR was performed with ChamQ Universal SYBR qPCR Master Mix (Vazyme

Biotechnology) according to the manufacturer's requirements. Primers demanded in

this analysis were listed in table 2 in supplementary materials.

RNA interference

SiRNAs (Genepharma) specific for porcine GSDMD were transfected into IPEC-J2

cells using the Lipofectamine 8000 transfection reagent according to the

manufacturer's instructions. The sequences of GSDMD siRNAs were listed in table 3

in supplementary materials.

Propidium iodide staining

Cells were incubated with Propidium iodide (PI) (BD Bioscience) for 15 min under

light-proof conditions after transfection with indicated plasmids for 24 h. Dyeing

condition was observed under fluorescence microscope (Nikon).

Statistical analysis

All experiments were performed independently at least three times. Data were

presented as the mean ± standard deviation (SD), analyzed and used for statistical

graphing by GraphPad Prism 8, the significance of differences was determined by

One-way ANOVA or Student's *t*-test. The significance of differences ranked as:

****stands for $P<0.0001$, ***stands for $P<0.001$, ** stands for $P<0.01$, * stands for

$P<0.05$ and ns stands for non-significant difference.

Data availability

The authors declare that all data supporting the findings of this article are available

within the paper and the supplementary information files or are available from the

authors upon request.

References

1. Zhang, Q. & Yoo, D. Immune evasion of porcine enteric coronaviruses and viral modulation of antiviral innate signaling. *Virus Res* **226**, 128-141 (2016).

2. Stevenson, G.W. *et al.* Emergence of Porcine epidemic diarrhea virus in the United States: clinical signs, lesions, and viral genomic sequences. *J Vet Diagn Invest* **25**, 649-654 (2013).

3. Jung, K. & Saif, L.J. Porcine epidemic diarrhea virus infection: Etiology, epidemiology, pathogenesis and immunoprophylaxis. *Vet J* **204**, 134-143 (2015).

4. Xing, Y. *et al.* The papain-like protease of porcine epidemic diarrhea virus negatively regulates type I interferon pathway by acting as a viral deubiquitinase. *J Gen Virol* **94**, 1554-1567 (2013).

5. Chu, H.F. *et al.* Structural and Biochemical Characterization of Porcine Epidemic Diarrhea Virus Papain-Like Protease 2. *J Virol* **96**, e0137221 (2022).

6. Clementz, M.A. *et al.* Deubiquitinating and interferon antagonism activities of coronavirus papain-like proteases. *J Virol* **84**, 4619-4629 (2010).

7. Zheng, D., Chen, G., Guo, B., Cheng, G. & Tang, H. PLP2, a potent deubiquitinase from murine hepatitis virus, strongly inhibits cellular type I interferon production. *Cell Res* **18**, 1105-1113 (2008).

8. Wang, Y. *et al.* Porcine transmissible gastroenteritis virus inhibits NF-kappaB activity via nonstructural protein 3 to evade host immune system. *Virol J* **16**, 97 (2019).

9. Man, S.M. & Kanneganti, T.D. Converging roles of caspases in inflammasome activation, cell death and innate immunity. *Nat Rev Immunol* **16**, 7-21 (2016).

10. Ball, D.P. *et al.* Caspase-1 interdomain linker cleavage is required for pyroptosis. *Life Sci Alliance* **3** (2020).

11. Boucher, D. *et al.* Caspase-1 self-cleavage is an intrinsic mechanism to terminate inflammasome activity. *J Exp Med* **215**, 827-840 (2018).

12. Ross, C. *et al.* Inflammatory Caspases: Toward a Unified Model for Caspase Activation by Inflammasomes. *Annu Rev Immunol* **40**, 249-269 (2022).

13. Bambouskova, M. *et al.* Itaconate confers tolerance to late NLRP3 inflammasome activation. *Cell Rep* **34**, 108756 (2021).

14. Jorgensen, I. & Miao, E.A. Pyroptotic cell death defends against intracellular pathogens. *Immunol Rev* **265**, 130-142 (2015).

591   15.   Liu, X. *et al.* Inflammasome-activated gasdermin D causes pyroptosis by forming
592   membrane pores. *Nature* **535**, 153-158 (2016).

593   16.   Sborgi, L. *et al.* GSDMD membrane pore formation constitutes the mechanism of
594   pyroptotic cell death. *EMBO J* **35**, 1766-1778 (2016).

595   17.   Wang, W. *et al.* Zika virus infection induces host inflammatory responses by facilitating
596   NLRP3 inflammasome assembly and interleukin-1beta secretion. *Nat Commun* **9**, 106
597   (2018).

598   18.   Pan, P. *et al.* SARS-CoV-2 N protein promotes NLRP3 inflammasome activation to
599   induce hyperinflammation. *Nat Commun* **12**, 4664 (2021).

600   19.   Ahn, M. *et al.* Dampened NLRP3-mediated inflammation in bats and implications for a
601   special viral reservoir host. *Nat Microbiol* **4**, 789-799 (2019).

602   20.   Jiang, Y. *et al.* Complement Receptor C5aR1 Inhibition Reduces Pyroptosis in
603   hDPP4-Transgenic Mice Infected with MERS-CoV. *Viruses* **11** (2019).

604   21.   Junqueira, C. *et al.* SARS-CoV-2 infects blood monocytes to activate NLRP3 and AIM2
605   inflammasomes, pyroptosis and cytokine release. *Res Sq* (2021).

606   22.   Tsu, B.V. *et al.* Diverse viral proteases activate the NLRP1 inflammasome. *Elife* **10** (2021).

607   23.   Shi, F. *et al.* Coronaviruses Nsp5 Antagonizes Porcine Gasdermin D-Mediated Pyroptosis
608   by Cleaving Pore-Forming p30 Fragment. *mBio*, e0273921 (2022).

609   24.   Zheng, Y. *et al.* Zika virus elicits inflammation to evade antiviral response by cleaving
610   cGAS via NS1-caspase-1 axis. *EMBO J* **37** (2018).

611   25.   Wang, Y. *et al.* Inflammasome Activation Triggers Caspase-1-Mediated Cleavage of
612   cGAS to Regulate Responses to DNA Virus Infection. *Immunity* **46**, 393-404 (2017).

613   26.   Shi, Y. *et al.* E3 ubiquitin ligase SYVN1 is a key positive regulator for GSDMD-mediated
614   pyroptosis. *Cell Death Dis* **13**, 106 (2022).

615   27.   Shrivastava, G., Leon-Juarez, M., Garcia-Cordero, J., Meza-Sanchez, D.E. &
616   Cedillo-Barron, L. Inflammasomes and its importance in viral infections. *Immunol Res* **64**,
617   1101-1117 (2016).

618   28.   Wang, W. *et al.* EV71 3D Protein Binds with NLRP3 and Enhances the Assembly of
619   Inflammasome Complex. *PLoS Pathog* **13**, e1006123 (2017).

620   29.   He, Z. *et al.* NLRP3 Inflammasome Activation Mediates Zika Virus-Associated
621   Inflammation. *J Infect Dis* **217**, 1942-1951 (2018).

622   30.   McAuley, J.L. *et al.* Activation of the NLRP3 inflammasome by IAV virulence protein
623   PB1-F2 contributes to severe pathophysiology and disease. *PLoS Pathog* **9**, e1003392
624   (2013).

625   31.   Robinson, K.S. *et al.* Enteroviral 3C protease activates the human NLRP1 inflammasome
626   in airway epithelia. *Science* **370** (2020).

627   32.   Zhu, S. *et al.* Nlrp9b inflammasome restricts rotavirus infection in intestinal epithelial cells.
628   *Nature* **546**, 667-670 (2017).

629   33.   Lee, S. *et al.* AIM2 forms a complex with pyrin and ZBP1 to drive PANoptosis and host
630   defence. *Nature* **597**, 415-419 (2021).

631   34.   Sun, X. *et al.* SARS-CoV-2 non-structural protein 6 triggers NLRP3-dependent
632   pyroptosis by targeting ATP6AP1. *Cell Death Differ* **29**, 1240-1254 (2022).

633   35.   Lee, S., Channappanavar, R. & Kanneganti, T.D. Coronaviruses: Innate Immunity,
634   Inflammasome Activation, Inflammatory Cell Death, and Cytokines. *Trends Immunol* **41**,

1083-1099 (2020).

36.　Eltobgy, M.M. *et al.* Caspase-4/11 exacerbates disease severity in SARS-CoV-2 infection
by promoting inflammation and immunothrombosis. *Proc Natl Acad Sci U S A* **119**,
e2202012119 (2022).

37.　Kanneganti, T.D. *et al.* Bacterial RNA and small antiviral compounds activate caspase-1
through cryopyrin/Nalp3. *Nature* **440**, 233-236 (2006).

38.　Kanneganti, T.D. *et al.* Critical role for Cryopyrin/Nalp3 in activation of caspase-1 in
response to viral infection and double-stranded RNA. *J Biol Chem* **281**, 36560-36568
(2006).

39.　Thomas, P.G. *et al.* The intracellular sensor NLRP3 mediates key innate and healing
responses to influenza A virus via the regulation of caspase-1. *Immunity* **30**, 566-575
(2009).

40.　Ma, J. *et al.* SARS-CoV-2 nucleocapsid suppresses host pyroptosis by blocking
Gasdermin D cleavage. *EMBO J* **40**, e108249 (2021).

41.　Shan, Y. *et al.* Nucleocapsid protein from porcine epidemic diarrhea virus isolates can
antagonize interferon-lambda production by blocking the nuclear factor-kappaB nuclear
translocation. *J Zhejiang Univ Sci B* **19**, 570-580 (2018).

42.　Zhou, Y. *et al.* Effect of route of inoculation on innate and adaptive immune responses to
porcine epidemic diarrhea virus infection in suckling pigs. *Vet Microbiol* **228**, 83-92
(2019).

Acknowledgements

This work was financially supported by the National Natural Science Foundation of

China (32072817), the Zhejiang Provincial Key R&D Program of China

(2021C02049), the Scientific Research Fund of Zhejiang Provincial Education

Department (Y202045613, Y202148324), the Zhejiang Provincial Natural Science

Foundation of China (LY18C180001, LY21C180001), and the Intergovernmental

Collaborative Project in S & T Innovation under the National Key R & D Program

(Grant No. 2018YFE0111700).

Author contributions

666　X.F., Y.Y., W.X. and F.S. contributed to the design of experiments and performed most

667 of the experiments. D.L, X.L., N.C., Q.L. and Y.S. performed some experiments. X.F,

668 D.L., W.X., X.L. and F.S. contributed to data analysis. X.F., D.L., Y.Y. and F.S.

669 contributed to writing the manuscript.

Competing interests

672 The authors declare no competing interests.

Figure legends

Figure 1. PEDV infection elevates the expression of caspase-1. A The intestinal tissue lysates were extracted from healthy or PEDV naturally-infected piglets for immunoblotting (IB). **B, C** IPEC-J2 cells were mock infected or infected with PEDV at an MOI of 0.5, followed by cycloheximide (CHX) treatment (25 μg/ml) for 6 h. The indicated gene mRNA was quantified by qRT-PCR (**B**). The indicated proteins were analyzed by immunoblotting (**C**). **D** Vero cells were transfected with plasmid encoding porcine caspase-1. At 24 h after transfection, the cells were mock infected or infected with different doses of PEDV for another 24 h, the cell lysates were then processed for immunoblotting. **E** Vero cells were transfected with plasmid encoding porcine caspase-1. At 24 h after transfection, the cells were mock infected or infected with PEDV at an MOI of 0.1. At indicated time points after infection, the cell lysates were analyzed by immunoblotting. **F, H** Vero cells were co-transfected with plasmids encoding GSDMD-FL and caspase-1. At 12 h after transfection, the cells were mock infected or infected with different doses of PEDV for another 24 h, the cell lysates

were processed for immunoblotting (**F**), the supernatants were collected for LDH

release assay (**H**). **G, I** Vero cells were co-transfected with plasmids encoding

GSDMD-FL and caspase-1. At 12 h after transfection, the cells were mock infected or

infected with PEDV at an MOI of 0.05. At indicated time points after infection, the

cell lysates were analyzed by immunoblotting (**G**), the supernatants were collected for

LDH release assay (**I**). All results shown are representative of at least three

independent experiments. ****stands for P<0.0001, ***stands for P<0.001, ** stands

for P<0.01, * stands for P<0.05 and ns stands for non-significant difference.

Figure 2. PLP2 inhibits the proteasomal degradation of caspase-1 by removing

K11-linked ubiquitin chains at Lys134. A HEK293T cells were co-transfected with

HA-caspase-1-C285A and increasing amount of MYC-PEDV-PLP2. Cell lysates were

analyzed by immunoblotting. **B** HEK293T cells were co-transfected with

HA-caspase-1-C285A together with empty vector or MYC-PEDV-PLP2 then treated

with cycloheximide (CHX) (25 μg/ml) for the indicated time points. Cell lysates were

analyzed by immunoblotting. **C** HEK293T cells were co-transfected with

HA-caspase-1-C285A together with wild-type PEDV PLP2 or its protease-defective

mutants (C113A, H272A and D285A). Cell lysates were analyzed by immunoblotting.

D Immunoprecipitation analysis of HEK293T cells expressing FLAG-caspase-1 and

HA-UB together with MYC-tagged empty vector, wild-type PEDV-PLP2 or its

mutants as indicated. Anti-FLAG immunoprecipitates were analyzed by

immunoblotting (IP). The expression levels of the transfected proteins were analyzed

by immunoblotting (INPUT). **E** HEK293T cells were co-transfected with FLAG-caspase-1 or its mutant FLAG-caspase-1-K134R together with increasing amount of HA-tagged K11-linked ubiquitin (HA-K11). Cell lysates were analyzed by immunoblotting. **F** HEK293T cells were transfected with FLAG-caspase-1 together with HA-K11 and MYC-PEDV-PLP2 as indicated. Cell lysates were analyzed by immunoblotting. **G** HEK293T cells were transfected with HA-caspase-1-C285A or its mutant HA-caspase-1-C285A-K134R together with MYC-tagged empty vector or PEDV PLP2. Cell lysates were analyzed by immunoblotting. **H** Vero cells were transfected with FLAG-caspase-1 together with HA-K11 or empty vector. At 24 h after transfection, the cells were mock infected or infected with PEDV at an MOI of 0.05 for another 24 h. Immunoprecipitation was performed with anti-FLAG binding beads and analyzed by immunoblotting. All results shown are representative of at least three independent experiments.

Figure 3. Caspase-1 attenuates IFN-I signaling during PEDV infection. A-D Relative qRT-PCR analysis of *IFN-α*, *IFN-β*, *ISG15* and *OAS1* mRNA levels in IPEC-J2 cells transfected with empty vector or caspase-1 for 24 h, and then stimulated with poly (I:C) (200 ng/ml) for another 12 h. **E, F** Relative qRT-PCR analysis of *IFN-β* and *ISG15* mRNA levels in IPEC-J2 cells treated with VX765 (20 μM) or DMSO, followed by poly (I:C) (200 ng/ml) transfection for 12 h. **G, H** Relative qRT-PCR analysis of *IFN-β* and *ISG15* mRNA levels in wild type (WT) or caspase-1 knockout (KO) THP-1 cells transfected with poly (I:C) (200 ng/ml) for 12 h. **I, J**

Relative qRT-PCR analysis of *IFN-β* and *ISG15* mRNA levels in IPEC-J2 cells

transfected with empty vector or PEDV PLP2 for 24 h, followed with poly (I:C) (200

ng/ml) transfection for another 12 h. **K, L** Relative qRT-PCR analysis of *IFN-β* and

ISG15 mRNA levels in IPEC-J2 cells transfected with empty vector or caspase-1 for

24 h, followed with PEDV infection (MOI=0.5) for another 24 h. **M, N** Relative

qRT-PCR analysis of *IFN-β* and *ISG15* mRNA levels in IPEC-J2 cells treated with

VX765 (20 μM) or DMSO, followed with PEDV infection (MOI=0.5) for 24 h. **O, P**

Relative qRT-PCR analysis of *IFN-β* and *ISG15* mRNA levels in IPEC-J2 cells

transfected with empty vector or PEDV PLP2 for 24 h, followed with PEDV infection

(MOI=0.5) for another 24 h. **Q** IPEC-J2 cells were transfected with empty vector or

PEDV PLP2 for 24 h, followed with PEDV infection at an MOI of 0.5 for another 24

h. The indicated gene mRNA levels were quantified by qRT-PCR and the indicated

proteins were analyzed by immunoblotting. **R** Relative qRT-PCR analysis of *PEDV S*

mRNA levels in IPEC-J2 cells transfected with empty vector or increasing amount of

caspase-1 for 12 h, followed with PEDV infection (MOI=0.1) for another 24 h. **S**

Relative qRT-PCR analysis of *PEDV S* mRNA levels in IPEC-J2 cells transfected

with empty vector and treated with or without VX765 (20 μM), co-treated with

necrosulfonamide (NSA 8 μM), followed with PEDV infection (MOI=0.1) for another

24 h. All results shown are representative of at least three independent experiments.

****stands for $P<0.0001$, ***stands for $P<0.001$, ** stands for $P<0.01$, * stands for

$P<0.05$ and ns stands for non-significant difference.

Figure 4. Cleavage of RIG-I by caspase-1 ensures PEDV replication. A HEK293T cells co-transfected with FLAG-caspase-1 and S-MYC-RIG-I (porcine derived) were lysed and immunoprecipitated with anti-FLAG beads and analyzed by immunoblotting. **B, C** HEK293T cells were co-transfected with S-MYC-RIG-I and increasing amount of HA-caspase-1 or HA-caspase-1-C285A. Cell lysates were analyzed by immunoblotting. **D** HEK293T cells were co-transfected with wild-type (WT) S-MYC-RIG-I or its mutants (D163A, D189A, D194A, D199/202/203A, D209A) together with empty vector or HA-caspase-1. Cell lysates were analyzed by immunoblotting. **E** Vero cells were transfected with empty vector, RIG-I together with or without caspase-1. At 12 h after transfection, the cells were mock infected or infected with PEDV at an MOI of 0.05. The indicated gene mRNA levels were quantified by qRT-PCR. **F-H** Relative qRT-PCR analysis of *PEDV S, IFN-β* and *ISG15* mRNA levels in IPEC-J2 cells transfected with empty vector, RIG-I or RIG-I-D189A for 24 h, followed by PEDV infection (MOI=0.5) for another 24 h. **I-K** Relative qRT-PCR analysis of *PEDV S, IFN-β* and *ISG15* mRNA levels in IPEC-J2 cells transfected with empty vector, RIG-I, RIG-I-1-189aa or RIG-I-190-943aa for 24 h, followed by PEDV infection (MOI=0.5) for another 24 h. All results shown are representative of at least three independent experiments. ****stands for P<0.0001, ***stands for P<0.001, ** stands for P<0.01, * stands for P<0.05 and ns stands for non-significant difference.

Figure 5. PLP2 degrades GSDMD-p30 to inhibit pyroptosis. A HEK293T cells

 were initially co-transfected with FLAG-GSDMD-FL and HA-caspase-1. At 6 h after

transfection, the cells were transfected with increasing amount of MYC-PEDV-PLP2.

Cell lysates were analyzed by immunoblotting. **B, C** HEK293T cells were

co-transfected with FLAG-GSDMD-p30 and increasing amount of

MYC-PEDV-PLP2. Cell lysates were analyzed by immunoblotting (**B**). The

supernatants were collected for LDH release assay (**C**). **D** HEK293T cells

co-transfected with MYC-PEDV-PLP2 together with empty vector,

FLAG-GSDMD-FL or FLAG-GSDMD-p30 were lysed and immunoprecipitated with

anti-FLAG beads and analyzed by immunoblotting. **E** HEK293T cells were

transfected with FLAG-GSDMD-p30 and GFP-PLP2 respectively or together for 24 h,

and then FLAG-GSDMD-p30 cells were labeled with indicated antibodies.

Subcellular localization of FLAG-GSDMD-p30 (red), GFP-PLP2 (green), and DAPI

(blue, nucleus marker) were visualized with confocal microscopy. All results shown

are representative of at least three independent experiments. ****stands for P<0.0001,

***stands for P<0.001, ** stands for P<0.01, * stands for P<0.05 and ns stands for

non-significant difference.

Figure 6. PLP2 removes the K27-linked ubiquitin chain at K275 residue of

GSDMD-p30. A HEK293T cells were transfected with FLAG-GSDMD-p30 together

with empty vector or MYC-PEDV-PLP2 then treated with DMSO, MG132 (10 μM),

3MA (1 mM) or CQ (40 μM) for 6 h. Cell lysates were analyzed by immunoblotting.

B, C HEK293T cells were co-transfected with FLAG-GSDMD-p30 together with

empty vector, wild type PLP2 or its protease-defective mutants (C113A, H272A,

D285A). The supernatants were collected for LDH release assay (**B**). Cell lysates

were analyzed by immunoblotting (**C**). **D** HEK293T cells were co-expressed with

FLAG-GSDMD-p30 or its mutants (K103R, K177R, K275R) together with empty

vector or HA-UB. Cell lysates were analyzed by immunoblotting. **E** HEK293T cells

were co-transfected with FLAG-GSDMD-p30 and HA-tagged WT ubiquitin or its

mutants (K6, K11, K27, K29, K33, K48, K63, KO). The supernatants were collected

for LDH release assay. **F** Co-immunoprecipitation and immunoblot analysis of

extracts of HEK293T cells transfected with FLAG-GSDMD-p30 or

FLAG-GSDMD-p30-K275R together with HA-UB or HA-K27 ubiquitin. **G**

Co-immunoprecipitation and immunoblot analysis of extracts of HEK293T cells

transfected with FLAG-GSDMD-p30 and HA-K27 ubiquitin together with empty

vector, wild type PLP2 or its mutants (C113A, H272A, D285A). All results shown are

representative of at least three independent experiments. ****stands for P<0.0001,

***stands for P<0.001, ** stands for P<0.01, * stands for P<0.05 and ns stands for

non-significant difference.

Figure 7. Papain-like proteases in other CoVs can also abrogate pyroptosis by

degradation of GSDMD-p30. A, B HEK293T cells were transfected with

FLAG-GSDMD-p30 together with empty vector, PEDV-PLP2, SARS-CoV-2-PLPro

or PDCoV-PLPro. The supernatants were collected for LDH release assay (**A**). Cell

lysates were analyzed by immunoblotting (**B**). **C** HEK293T cells were transfected

with HA-caspase-1-C285A together with empty vector, PEDV-PLP2, SARS-CoV-2-PLPro or PDCoV-PLPro. Cell lysates were analyzed by immunoblotting. **D** IPEC-J2 cells were transfected with empty vector, PEDV-PLP2, SARS-CoV-2-PLPro or PDCoV-PLPro. At 24 h after transfection, the cells were infected with PEDV at an MOI of 0.1 for another 24 h. The indicated gene mRNA levels were quantified by qRT-PCR (top). Cell lysates were analyzed by immunoblotting (bottom). **E** Immunoblot analysis of extracts of THP-1 cells stably infected with FLAG-EV-Lentivirus or FLAG-N-Lentivirus. **F** PMA-differentiated THP-1 macrophages were stably infected with FLAG-EV-Lentivirus or FLAG-N-Lentivirus. The supernatants were collected and analyzed by ELISA for IL-1β. **G-L** PMA-differentiated THP-1 macrophages were stably infected with FLAG-EV-Lentivirus or FLAG-N-Lentivirus, followed with poly (I:C) (200 ng/ml) transfection for 6 h. Cell lysates were analyzed by immunoblotting (**G**). The supernatants were collected and analyzed by ELISA for IL-1β (**H**), IFN-β (**I**) and for LDH release assay (**L**). The mRNA levels of *IFN-β* (**J**) and *ISG15* (**K**) were quantified by qRT-PCR. All results shown are representative of at least three independent experiments. ****stands for $P<0.0001$, ***stands for $P<0.001$, ** stands for $P<0.01$, * stands for $P<0.05$ and ns stands for non-significant difference.

Figure 8. Mechanistic diagram of CoVs antagonizing antiviral responses and pyroptosis.

Supplementary Fig. 1. PEDV PLP2 interacts with both caspase-1 and GSDMD. A

Vero cells were transfected with FLAG-GSDMD-FL and then infected with PEDV at

an MOI of 0.05. Immunoprecipation was performed with anti-FLAG antibody. The

potential GSDMD-binding viral proteins were evaluated using MS analysis. **B**

Co-immunoprecipitation and immunoblot analysis of extracts of HEK293T cells

transfected with MYC-PEDV-PLP2 together with FLAG-GSDMD-FL or

FLAG-tagged empty vector. **C** Co-immunoprecipitation and immunoblot analysis of

extracts of HEK293T cells transfected with MYC-PEDV-PLP2 together with

FLAG-caspase-1 or FLAG-tagged empty vector, then treated with VX765 (20 μM). **D**

HEK293T cells were transfected with FLAG-GSDMD-FL, GFP-PLP2 and

HA-caspase-1 respectively or together for 24 h, and then the cells were labeled with

indicated antibodies. Subcellular localization of FLAG-GSDMD-FL (cyan),

GFP-PLP2 (green), HA-caspase-1 (red) and DAPI (blue, nucleus marker) were

visualized with confocal microscopy. All results shown are representative of at least

three independent experiments.

Supplementary Fig. 2. The papain-like protease 2 of PEDV inhibits proteasomal

degradation of caspase-1. A HEK293T cells were transfected with

FLAG-GSDMD-FL and HA-caspase-1 together with increasing amount of

MYC-PEDV-PLP2. Cell lysates were analyzed by immunoblotting. **B** HEK293T cells

were co-transfected with FLAG-GSDMD-FL together with empty vector or

MYC-PEDV-PLP2 then treated with cycloheximide (CHX) (25 μg/ml) for the

indicated time points. Cell lysates were analyzed by immunoblotting. **C**

Co-immunoprecipitation and immunoblot analysis of extracts of HEK293T cells

transfected with FLAG-caspase-1 or FLAG-caspase-1-K134R together with HA-K11

ubiquitin. All results shown are representative of at least three independent

experiments.

Supplementary Fig. 3. RIG-I was cleaved by caspase-1. A (long exposure of Fig.

4D) HEK293T cells were co-transfected with wild-type (WT) S-MYC-RIG-I or its

mutants (D163A, D189A, D194A, D199/202/203A, D209A) together with empty

vector or HA-caspase-1. Cell lysates were analyzed by immunoblotting. **B** HEK293T

cells were co-transfected with wild-type (WT) h-FLAG-RIG-I or its single, double

mutants (D163A, D194A, D234A, D163/194A, D163/234A, D194/234A) together

with empty vector or H-HA-caspase-1. Cell lysates were analyzed by immunoblotting.

C Alignment of the amino acid sequence of RIG-Is from different species. **D**

HEK293T cells were co-transfected with wild-type (WT) h-FLAG-RIG-I or its triple

mutant (D163/194/234A) together with empty vector or h-HA-caspase-1. Cell lysates

were analyzed by immunoblotting.

Supplementary Fig. 4. PLP2 degrades GSDMD-p30 by removing the K27-linked

ubiquitin chains at K275 residue to inhibit pyroptosis. A HEK293T cells were

transfected with FLAG-GSDMD-FL together with increasing amount of

MYC-PEDV-PLP2. Cell lysates were analyzed by immunoblotting. **B** HEK293T cells

were co-transfected with FLAG-GSDMD-p30 and MYC-PEDV-PLP2, then the cells

were performed with propidium iodide (PI) staining analysis. **C** Immunoblot analysis

of extracts of IPEC-J2 cells transfected with negative control (NC) siRNA or GSDMD

siRNA (si-GSDMD#1, si-GSDMD#2, si-GSDMD#3). Cell lysates were analyzed by

immunoblotting. **D, E** IPEC-J2 cells were transfected with GSDMD siRNA

(si-GSDMD#1) for 24 h and then expressed with GSDMD-FL or GSDMD-FL-K275R

for another 24 h. The cells were then infected with PEDV at an MOI of 0.1 for 24 h.

The indicated gene mRNA levels were quantified by qRT-PCR (**D**). The supernatants

were collected for LDH release assay (**E**). All results shown are representative of at

least three independent experiments. ****stands for $P<0.0001$, ***stands for $P<0.001$,

** stands for $P<0.01$, * stands for $P<0.05$ and ns stands for non-significant difference.