Mapping the Macrostructure and Microstructure of the in vivo Human Hippocampus using Diffusion MRI

Bradley G. Karat[1,2], Jordan DeKraker[1,3], Uzair Hussain[4], Stefan Köhler[5], Ali R. Khan[1,6]

[1]Robarts Research Institute, Schulich School of Medicine and Dentistry, University of Western Ontario, Canada, [2]Neuroscience Graduate Program, University of Western Ontario, Canada, [3]Montreal Neurological Institute, McGill University, Montreal, Quebec, Canada, [4]University Health Network, Toronto, Ontario, Canada, [5]Department of Psychology, University of Western Ontario, Canada, [6]Western Institute for Neuroscience, University of Western Ontario, Canada,

Data and code availability

The *HippUnfold* software is openly available and can be found at https://github.com/khanlab/hippunfold. Code used for the study can be found at https://github.com/Bradley-Karat/Hippo_Macro_Micro.
Data were provided [in part] by the Human Connectome Project, WU-Minn Consortium (Principal Investigators: David Van Essen and Kamil Ugurbil; 1U54MH091657) funded by the 16 NIH Institutes and Centers that support the NIH Blueprint for Neuroscience Research; and by the McDonnell Center for Systems Neuroscience at Washington University; https://www.humanconnectome.org/study/hcp-young-adult/document/1200-subjects-data-release

Abstract

The hippocampus is classically divided into mesoscopic subfields which contain varying microstructure that contribute to their unique functional roles. It has been challenging to characterize this microstructure with current MR based neuroimaging techniques. In this work, we used diffusion MRI and a novel surface-based approach in the hippocampus which revealed distinct microstructural distributions of neurite density and dispersion, T1w/T2w ratio as a proxy for myelin content, fractional anisotropy, and mean diffusivity. We used the Neurite Orientation Dispersion and Density Imaging (NODDI) model optimized for gray matter diffusivity to characterize neurite density and dispersion. We found that neurite dispersion was highest in the Cornu Ammonis (CA) 1 and subiculum subfields which likely captures the large heterogeneity of tangential and radial fibers, such as the Schaffer collaterals, perforant path, and pyramidal neurons. Neurite density and T1w/T2w were highest in the subiculum and CA3 and lowest in CA1, which may reflect known myeloarchitecture differences between these subfields. Using a simple logistic regression model, we showed that neurite density, dispersion, and T1w/T2w measures provided good separability across the subfields, suggesting that they may be sensitive to the known variability in subfield cyto- and myeloarchitecture. We report macrostructural measures of gyrification, thickness, and curvature that were in line with ex vivo descriptions of hippocampal anatomy. We employed a multivariate orthogonal projective non-negative matrix factorization (OPNNMF) approach to capture co-varying regions of macro- and microstructure across the hippocampus. The clusters were highly variable along the medial-lateral (proximal-distal) direction, likely reflecting known differences in morphology, cytoarchitectonic profiles, and connectivity. Finally, we show that by examining the main direction of diffusion relative to canonical hippocampal axes, we could identify regions with stereotyped microstructural orientations that may map onto specific fiber pathways, such as the Schaffer collaterals, perforant path, fimbria, and alveus. These results highlight the value of combining in vivo diffusion MRI with computational approaches for capturing hippocampal microstructure, which may provide useful features for understanding cognition and for diagnosis of disease states.

Keywords: Hippocampus; Subfields; Microstructure; Neurite Density; Neurite Dispersion; Diffusion MRI

1. Introduction

The hippocampus is classically divided into structurally distinct mesoscopic subfields according

to differences in cyto-, myelo-, and chemoarchitecture (Duvernoy et al., 2013; Ding & Van

Hoesen, 2015). The neuronal circuitry that compose the microstructure of the hippocampus exist

within and across these subfields. For example, the pyramidal neurons that exist within the

Cornu Ammonis (CA) and subiculum subfields have apical and basal dendrites which project

across the laminae, while their axons project to the alveus, a white matter bundle adjoining the

hippocampus. The trisynaptic pathway is the major circuitry component which connects the

subfields of the hippocampus. The entorhinal cortex connects to the dentate gyrus (DG) and

other subfields through the myelinated perforant path. The DG then projects to the pyramidal

neurons of CA3 through the mossy fibers, which then project to CA1 through the largely

unmyelinated Schaffer collaterals. Finally, CA1 projects to the subiculum and back to the

entorhinal cortex as a large hippocampal efferent. Hippocampal microstructure is key in

producing unique cognitive functions such as memory formation and storage and spatial

navigation among others (Voss et al., 2017; Goodroe et al., 2018; Horner et al., 2015).

Furthermore, the hippocampus is typically one of the earliest aberrant structures in many disease

states, where specific microstructural properties are differentially afflicted or spared (Moodley &

Chan, 2014; Dhikav & Anand, 2012; Small et al., 2011). While much work has addressed

volumetric characterization of the hippocampus and its subfields, understanding hippocampal

microstructure can provide key insights into its complex cognitive functions as well as its early

deterioration in disease.

Diffusion magnetic resonance imaging (dMRI) is a particular technique which holds

promise in probing the hippocampal circuitry by sensitizing the measured MRI signal to the

movement of water molecules, which diffuse more readily parallel to microstructure. Several

models have been proposed that attribute measures of the dMRI signal to compartments which

have varying diffusivity environments (Assaf et al., 2008; Assaf & Basser, 2005; Zhang et al.,

2012). One of the earliest and most widely used models proposed by Basser et al. (1994) is

diffusion tensor imaging (DTI). DTI estimates quantitative parameters such as fractional

anisotropy (FA - a measure of the anisotropy of diffusion), mean diffusivity (MD – magnitude of

diffusion), and the ellipsoidal orientation of the diffusion process. However, DTI has some

notable limitations. At increasing b-values (approximately greater than 1000-1500 s/mm^2) there

is contribution from multiple compartments with varying diffusivities (such as restricted intra-

cellular water), which goes beyond the monoexponential signal modelled by DTI (Assaf &

Cohen. 2000). As well, regions of crossing fibers result in planar DTI ellipsoids with understated

FA values (Campbell et al., 2005). Furthermore, DTI measures are sensitive to multiple

microstructural properties at the same time, decreasing their specificity (Pierpaoli et al., 1996).

Other models aim to utilize increasingly complex diffusion acquisitions to relate the diffusion

signal attenuation to varying sets of biophysically motivated compartments.

One popular compartment model is Neurite Orientation Dispersion and

Density Imaging (NODDI), which aims to provide a biophysical interpretation of the diffusion

signal (Zhang et al., 2012). NODDI assumes that three microstructural environments consisting

of an intra-cellular, extra-cellular, and cerebrospinal fluid (CSF) compartment contribute to the

diffusion signal. The intra-cellular compartment is modeled as a set of infinitely anisotropic

sticks (diffusion can only be parallel to the main orientation of the stick), while the extra-cellular

compartment is modeled as a zeppelin (or cylindrically symmetric tensor) with hindered

diffusion perpendicular to its main axis. The CSF compartment is modeled as a sphere with

gaussian isotropic diffusion. Diffusion is assumed to be contained separately within each

compartment, where the resulting signal is the sum from all compartments. NODDI aims to

overcome the limitations of DTI by providing microstructural scalars such as the neurite density

index (NDI) and orientation dispersion index (ODI) which are sensitive to fiber crossings and are

biophysically grounded (Zhang et al., 2012). NDI is meant to represent the volume fraction of

the intra-cellular compartment, which is believed to correspond to the density of dendrites,

axons, and other "stick" like processes in a voxel. ODI is meant to capture the essence of the

orientation distribution of the diffusion signal, where more complex or disperse microstructural

configurations correspond to a higher ODI value.

Contemporary work has attempted to examine hippocampal microstructure with both DTI

and NODDI. Some such studies have found age-related changes of hippocampal microstructure

by averaging NODDI measures within subfields (Radhakrishnan et al., 2020) while others have

shown regionally specific changes using DTI (Yassa et al., 2010). Recent work has also begun to

use measures derived from structural MRI to interrogate microstructural characteristics, such as

the ratio of T1-weighted over T2-weighted signal as a correlate of myelin (Glasser & Van Essen,

2011; Glasser et al., 2014). These measures may be useful to capture the myelinated intra-

hippocampal circuitry like the perforant path. A recent study investigated the variation of DTI

and intra-cortical myelin through the ratio of T1w/T2w images across the hippocampus using

non-negative matrix factorization, however, they did not make quantitative comparisons of

microstructure within and across the subfields (Patel et al., 2020). As well, they note the limited

specificity nature of the DTI and T1w/T2w metrics investigated. A recent study examined the

distribution of NODDI metrics and cortical thickness across the entire cerebral cortex including

the hippocampus by averaging metrics across all subjects within each cortical parcel (Fukutomi

et al., 2018). Thus, they only examined coarse-grained averages across the entire hippocampal

volume. The spatial distributions of NODDI and DTI measures have not been extensively

investigated within the hippocampal subfields and across its longitudinal axis.

The orientation and trajectory of the hippocampal circuitry including the trisynaptic

circuit has been probed previously using tractography and polarized light imaging (PLI). Ex vivo

work has indeed resolved major parts of the hippocampal circuitry using dMRI tractography

(Beaujoin et al., 2018) and PLI (Zeineh et al., 2017) in a small number of samples. While these

studies serve as a close to ground-truth reference for the orientation of hippocampal circuitry, a

difficult step has been recapitulation of this circuitry in vivo. Some in vivo work has attempted to

use DTI to capture parts of the trisynaptic circuit such as the perforant path (Yassa et al., 2010)

or the whole hippocampal circuitry (Zeineh et al., 2012). However, it is unclear whether the

found trajectories are anatomically valid. Furthermore, at lower resolutions, tracts can be

spurious requiring complex acquisition and correction schemes, and since acquisitions can vary

across studies, tractography practically always requires separate optimization of its parameters

(Zeineh et al., 2012). Thus, understanding hippocampal microstructure in vivo may benefit from

a simpler characterization of the orientation of hippocampal circuitry with reference to metrics

derived from common models like NODDI within and across the subfields.

In the current study, we examined the spatial distribution of NODDI and DTI metrics,

T1w/T2w ratio, and macrostructural features of thickness, gyrification, and curvature across the

hippocampus using high-resolution in vivo human connectome project (HCP) data (Van Essen et

al., 2013), something that has not been extensively investigated. We aimed to examine if the

microstructural metrics systematically vary across the cytoarchitectonic defined subfields.

Furthermore, we used Orthogonal Projective Non-Negative Matrix Factorization (OPNNMF) as

a multivariate approach to capture regions of the hippocampus where these metrics co-vary. We

aimed to consider the current OPNNMF representation of disparate macro- and microstructural

metrics under the framework of previous research which has suggested modes of hippocampal

organization along its medial-lateral (across subfields) and anterior-posterior (longitudinal) axes

(Genon et al., 2021; Robinson et al., 2015; Zhong et al., 2019; Cheng et al., 2020; Plachti et al.,

2019; Plachti et al., 2020; Patel et al., 2020, DeKraker et al., 2020). While most hippocampus

representations use voxel-based approaches, we utilized a novel surface-based approach called

HippUnfold (DeKraker et al., 2018; DeKraker et al., 2022). Much like in the neocortex, surface-

based methods are better suited to account for interindividual differences in tissue curvature and

digitation than voxel-based approaches (DeKraker et al., 2022; DeKraker et al., 2021). Aligning

hippocampi on a 2D surface preserves topology and the known contiguity of subfields, allowing

for improved anatomical detail to be captured. Finally, hippocampal gray matter shows a laminar

distribution similar to that of other cortical areas with large radial and tangential neurite

components, although the highly curved structure of the hippocampus is reflected in the

complexity of its neurite orientations. Importantly, these neurite orientations tend to be highly

aligned along one of the axes of the hippocampus that span the anterior-posterior (AP

longitudinal), proximal-distal (PD - across subfields), or inner-outer (across laminae) directions

(Figure 1A and B) (Zeineh et al., 2017; Nieuwenhuys et al., 2008; Duvernoy et al., 2013). Thus,

we also aimed to determine if the known stereotyped orientations of microstructure can be

elucidated by analyzing the primary direction of diffusion along each of the axes in vivo, as

depicted in Figure 1B.

Figure 1. Depiction of hippocampal structural axes, the stereotyped organization of

microstructure, and diffusion vectors of the hippocampus. (A) A coronal slice depicting the

structural axes of the hippocampus defined as anterior-posterior (AP), proximal-distal (PD), and

inner-outer (IO) provided by *HippUnfold.* The inner gray area corresponds to the SRLM, while

the rest of the uncoloured regions correspond to the stratum pyramidale and stratum oriens

layers. The white arrow in the top right inset depicts the orientation of the anterior-posterior axis.

The colour of the hippocampal surface is the anterior-posterior Laplace coordinates. Finally, the

black arrow depicts the intended level of the coronal slice for the cartoon depiction. (B)

Simplified cartoon depiction of known microstructural circuits within the hippocampus and their

approximate main orientation relative to the hippocampus, defined by the colour coded legend

on the right (Zeineh et al., 2017; Nieuwenhuys et al., 2008; Duvernoy et al., 2013). (C) Primary

diffusion directions for one subject (μ of the watson distribution from NODDI) overlaid on a

coronal slice approximately through the hippocampal body. Coloured borders represent the

hippocampal subfields provided by *HippUnfold.* (D) Pictorial example representing the NODDI

and hippocampal axis vectors in a single voxel defined in (C). Cosine similarities are represented

as the angle between the NODDI vector and each hippocampal vector, providing a measure of

orientation coherence along each cardinal axis (see section *2.5)*. Sub - Subiculum, CA - Cornu

Ammonis, DG - Dentate Gyrus, SRLM - Stratum Radiatum Lacunosum Moleculare, ETC -

Entorhinal Cortex.

2. Methods

2.1 Overview

A subset of 100 unrelated subjects from the publicly available Human Connectome Project

(HCP) 1200 dataset were used for this study (Van Essen et al., 2013). All 100 subjects were run

through *HippUnfold* (DeKraker et al., 2022), a new automated tool for surface-based subfield

segmentation and hippocampal unfolding (see section *2.3*; DeKraker et al., 2018). The Laplacian

coordinates generated from *HippUnfold* within each subject were used to calculate gradient

vector fields along each axis of the hippocampus (see section *2.5 &* Figure 1A*). NODDI and DTI

metrics were calculated in each subject's native space using whole-brain diffusion images (see

2.2 and *2.4*). Cosine similarities between the NODDI orientational vector (defined as μ of the

Watson distribution; see section *2.4*) and the vectors along each of the 3 axes (AP, PD, and IO)

were calculated at each voxel. Furthermore, the T1w/T2w ratio was calculated as a proxy for

myelin content. Macrostructural measures of curvature, gyrification, and thickness were

calculated along the midthickness surface (middle of the hippocampal gray matter) of the

hippocampus across all subjects. NODDI measures of ODI and NDI, DTI measures of FA and

MD, and the cosine similarities were all sampled along the midthickness surface within each

subject and averaged in unfolded space (DeKraker et al., 2018; DeKraker et al., 2022). Plots of

NODDI and DTI metrics, cosine similarities, and macrostructure metrics across the midthickness

surface were visualized as folded and unfolded surfaces. Logistic regression was performed at

the level of subfield averages using the NODDI and T1w/T2w metrics to elucidate their

variability across the subfields. Logistic regression was chosen due to its simplicity and

interpretability. Finally, Orthogonal Projective Non-Negative Matrix Factorization (OPNNMF)

was used to capture co-varying regions of the hippocampus and to examine the dimensions of

macro- and microstructure hippocampal organization.

2.2 Data acquisition and preprocessing

We used the publicly available HCP young adult dataset (ages 22-35), which consisted of

structural and diffusion MRI data for 1200 subjects (Van Essen et al., 2013). To avoid any biases

caused by family structures, we chose the 100 unrelated subjects subset for analysis (mean age:

27.52 years +/- 3.47 years; F/M: 54/46). Data included T1-weighted (T1w) and T2-weighted

(T2w) structural images at 0.7 mm^3 isotropic resolution and diffusion-weighted data at 1.25 mm^3

isotropic resolution. Structural images were obtained using a 3D MPRAGE sequence (TR-

2400ms, TE–2.14ms, TI-1000ms, FOV-224x224 mm). Diffusion images were obtained using a

spin-echo echo-planar imgaging sequence (b=0 (18 acquisitions), 1000, 2000, 3000s/mm^2, 90

diffusion-encoding directions, TR-5520ms, TE-89.5ms, FOV-210x180mm). Data used in the

preparation of this work were obtained from the Human Connectome Project (HCP) database

(Van Essen et al., 2013). In this work we utilized the preprocessed structural and diffusion

images for the HCP dataset. Preprocessing of structural images included: gradient distortion

correction, coregistration and averaging of repeated T1w and T2w runs using 6-DOF rigid

transformation, initial brain extractions for T1w and T2w, field map distortion correction and

registration of T2w with T1w images, bias field correction, and atlas registration. Preprocessing

of diffusion images included: intensity normalization across runs, EPI distortion correction, eddy

current and motion correction, gradient nonlinearity correction, and registration of the mean b0

image to T1w native space. The full pre-processing pipeline for structural and diffusion images

were published elsewhere (Andersson et al., 2015; Glasser et al., 2013; Jenkinson et al., 2002;

Sotiropoulos et al., 2013) and can be found at the HCP website

(https://www.humanconnectome.org/study/hcp-young-adult). To derive a correlate of myelin, we

divided the T1w image intensity by the T2w image intensity and corrected for the bias field

(Glasser & Van Essen, 2011; Glasser et al., 2014), which is referred to as T1w/T2w for the rest

of the paper.

2.3 HippUnfold - Hippocampal unfolding and surface-based segmentation

The newly developed *HippUnfold* tool was used in the current study for surface-based

segmentation and unfolding of the hippocampus (DeKraker et al., 2022). *HippUnfold* is

predicated on the idea that the hippocampal subfields are topologically constrained as they

differentiate from a flat cortical mantle (Duvernoy et al., 2013). Using a 'U-net' deep

convolutional neural network (Isensee et al., 2021), *HippUnfold* provides a detailed subject-

specific tissue segmentation of the hippocampal gray matter and its topological boundaries (for

example, the Hippocampal Amygdala Transition Area and the Indusium Griseum as the most

anterior and posterior topological boundaries, respectively) required for unfolding (DeKraker et

al., 2022). Segmentation is done for each individual hippocampi preserving its topologically

consistent structure, which is critical for inter-individual alignment across variably shaped

hippocampi. See Dekraker et al. (2022) for more detailed information on *HippUnfold.*

The outputs of *HippUnfold* utilized in this study included the provided subfield

segmentation, Laplacian coordinates for gradient field calculation (see section 2.5),

macrostructure measures of curvature, gyrification, and thickness, and the midthickness surface

representation for sampling volumetric space metrics onto the surface (see section 2.4). Due to

the small size of the Dentate Gyrus (DG) and CA4, we combined them into a single DG/CA4

subfield label. As well, portions of the DG were excluded in our surface representation. All

subfield segmentations for both hemispheres were reviewed for gross errors by BK. The

midthickness surface used in this study was composed of 2004 vertices with a spacing of roughly

1mm. Averaging of each metric across subjects was performed at each vertex in unfolded space.

2.4 Characterization of microstructure with NODDI & DTI

NODDI models the diffusion signal as a combination from 3 microstructural environments:

intra-cellular, extra-cellular, and cerebrospinal fluid (CSF) (Zhang et al., 2012). The intra-

cellular compartment is considered the space that is bounded by neurites, which is modelled as a

set of sticks. The stick geometry captures the restricted diffusion of water perpendicular to

neurites, and the relatively unhindered diffusion along them. Furthermore, sticks can capture the

wide range of neurite orientations, from highly coherent to highly dispersed tissue. The extra-

cellular compartment is the space around the neurites, which consists of glial cells and in gray

matter, the somas. In extra-cellular space, the signal is modeled as Gaussian anisotropic diffusion

to represent the hindered but not restricted movement of water. Finally, the CSF compartment is

modelled as Gaussian isotropic diffusion, representing the free movement of water. NODDI does

not draw any *a priori* assumptions about whether a voxel is gray matter, white matter or CSF,

and thus it treats each voxel as a possible combination of different compartments (Zhang et al.,

2012). Thus, the normalized dMRI signal can be written as:

$$E_{NODDI} = f_{iso}E_{iso}(d_{iso}) + W(\mu, \kappa)[f_{ec}E_{ec}(d_{perp}, d_{par}) + f_{ic}E_{ic}(d_{par})] \quad (1)$$

Where $f_{iso}E_{iso}, f_{ec}E_{ec}$ and $f_{ic}E_{iC}$ are the volume and signal fractions of the CSF, extra-cellular,

and intra-cellular (NDI) compartments, respectively. The extra-cellular and intra-cellular

compartments are linked orientationally by the Watson distribution $W(\mu, \kappa)$, where κ is the

concentration parameter (ODI $= \frac{2}{\pi}\arctan(\frac{1}{\kappa})$) and μ is the mean orientation of the Watson

distribution (herein referred to as the NODDI microstructural vector or primary diffusion

direction). The hindered perpendicular diffusion of the extra-cellular compartment d_{perp} is set via

a tortuosity model. The original NODDI model which was developed mainly for white matter sets

the parallel diffusivity value d_{par} equal to $1.7 \times 10^{-3} \frac{mm^2}{s}$ and the isotropic or CSF compartment

diffusion to $3.0 \times 10^{-3} \frac{mm^2}{s}$. Previous studies in the gray matter have sought to optimize d_{par}, and

have consistently found that the lowest mean squared error is achieved with d_{par} equal to

$1.1 \times 10^{-3} \frac{mm^2}{s}$ (Guerrero et al., 2019; Fukutomi et al., 2018). Thus, in the current study we

used the gray matter optimized d_{par} value of $1.1 \times 10^{-3} \frac{mm^2}{s}$ for fitting the NODDI model. The

Microstructure Diffusion Toolbox (MDT; Harms et al., 2017) was utilized to fit the NODDI model

using whole-brain diffusion images aligned to their respective T1w space with all b-values (b=0,

1000, 2000, 3000 s/mm^2). The validity of the assumptions of the NODDI model are discussed in

section 4.6.

We also used the MDT (Harms et al., 2017) to calculate metrics of FA and MD using DTI.

DTI was performed using only the $b = 1000 \frac{s}{mm^2}$ volumes to align with typical DTI experiments

(Behrens & Johansen-Berg, 2014). Both the NODDI and DTI metrics were mapped onto the

hippocampal midthickness surface using the process described below.

We used Connectome Workbench (https://github.com/Washington-

University/workbench) to sample values at each surface vertex from volume data. In this study

we used 2004 vertices defined along the midthickness surface of the hippocampus. Sampling

along the midthickness surface helps reduce partial volume effects. To sample voxel data along

the midthickness surface we used a ribbon-constrained mapping algorithm which also requires

the inner and outer surfaces also generated by *HippUnfold.* The ribbon method constructs a

polyhedron from the vertex's neighbor on each surface defined, and then estimates the volume of

the polyhedron that falls inside any nearby voxels to use as weights. We further reduced the

weight of any voxel based on its distance from the midthickness surface, where the scaling value

was calculated using a Gaussian with a standard deviation determined by the laminar thickness at

each vertex. This had the effect of more aggressively down-weighting voxels further from the

midthickness surface where the hippocampus is thinner. We then averaged each metric at each

vertex across all subjects to generate the average maps which were plotted in folded and

unfolded space.

2.5 Examining the primary direction of diffusion relative to hippocampal axes

Water molecules diffuse more readily parallel to microstructure, which in the hippocampus tends

to be aligned along the AP, PD, and IO axes (Zeineh et al., 2017; Nieuwenhuys et al., 2008;

Duvernoy et al., 2013). Thus, analyses were performed to examine how the primary direction of

diffusion was oriented relative to these axes, with the goal of elucidating the stereotyped

orientation of hippocampal microstructure (Figure 1). We obtained gradient vector fields along

the AP, PD, and IO axes by taking the first derivative of the respective Laplacian coordinates

provided by *HippUnfold* (Figure 1A), such that the vectors only pointed along one of the axes.

That is, we computed the partial derivative of the Laplacian coordinate function along the x, y,

and z spatial dimension:

338
$$\overrightarrow{G_{AP}} = \left[\frac{\partial \psi_{AP}}{\partial x}, \frac{\partial \psi_{AP}}{\partial y}, \frac{\partial \psi_{AP}}{\partial z}\right]$$

339
$$\overrightarrow{G_{PD}} = \left[\frac{\partial \psi_{PD}}{\partial x}, \frac{\partial \psi_{PD}}{\partial y}, \frac{\partial \psi_{PD}}{\partial z}\right] \quad (2)$$

340
$$\overrightarrow{G_{IO}} = \left[\frac{\partial \psi_{IO}}{\partial x}, \frac{\partial \psi_{IO}}{\partial y}, \frac{\partial \psi_{IO}}{\partial z}\right]$$

Where the function ψ_* represents the spatial Laplacian coordinates along a particular

hippocampal axis (AP, PD, or IO), which were calculated by solving Laplaces equation along

each axis ($\nabla^2(\psi) = 0$) (DeKraker et al., 2022). The result was 3 distinct vector images within a

hemisphere for each subject and axis. With the aligned NODDI microstructural vectors

(representing the primary diffusion direction), we calculated cosine similarities between the

generated vectors along the AP, PD, and IO axes and the NODDI microstructural vector at each

voxel (Figure 1C & D). All vectors were normalized before calculating cosine similarities. The

cosine similarity was defined as the inner product between vectors:

349
$$\text{abs}\left(\frac{\bar{a} \cdot \bar{u}}{|\bar{a}|\,|\bar{u}|}\right) \quad (3)$$

Where $\bar{a}$ was a hippocampal axis vector in a single voxel (AP, PD, or IO), and $\bar{u}$ was the

NODDI microstructural vector (μ of the Watson distribution) at the same voxel. A higher cosine

similarity meant that diffusion is increasingly oriented along that hippocampal axis (cosine

similarities of 0 and 1 correspond to angles of 90 degrees and 0 degrees, respectively). The

cosine similarities were then put into context of the known spatial arrangement of hippocampal

microstructure and their stereotyped orientation (Figure 1A and B) under the assumption that the

primary direction of diffusion modelled the main microstructural orientation at each voxel. There

were a total of 3 cosine similarity images (one for AP, PD, and IO similarity values) within each

hemisphere for each subject. Each scalar cosine similarity image was sampled along the

midthickness surface and averaged as described above.

2.6 Correlations between all metrics

Spearman correlations were performed at both the subject-level and at the level of the vertex

averaged maps (Figure 2C-J average maps). However, it is difficult to test the significance of the

spatial correspondence between maps. One proposed solution that has been implemented using

the spherical representation of the cortical surface has been to perform spin testing, where the

cortical sphere is randomly permuted to derive a null distribution of association which preserves

the spatial autocorrelation in the data (Alexander-Bloch et al., 2018). Critically, the geometry of

our unfolded hippocampus cannot be well represented by a sphere. We thus developed a similar

method on the unfolded hippocampus. Our developed method uses periodic boundary conditions

on the unfolded plane, which can be thought of as performing spin testing on a torus geometry.

This allows us to perform rotations and translations and thus derive permuted null distributions.

We then used this hippocampal spin test to analyze the significance of the observed associations

between the averaged maps. We performed 2500 permutations between any two averaged maps,

and we then performed false-discovery rate correction on the derived p-values using the

Benjamini-Hochberg method. Technical and practical background on the developed

hippocampal spin test can be found in Supplementary Document 1, and the spin test code can be

found at Karat (2023).

2.7 Orthogonal Projective NNMF (OPNNMF)

Orthogonal Projective NNMF (OPNNMF) was used in this work to attempt to identify co-

varying regions in the hippocampus using the metrics described above (Sotiras et al., 2015; Yang

& Oja, 2010). OPNNMF decomposes an input matrix X of dimensions a x b into a component

matrix C (a x k) and a weight matrix W (k x b). The number of components (k) is defined *a*

priori. The component and weight matrices are derived such that their multiplication best

reconstructs the input data (X ~ C x W). OPNNMF solves the following minimization problem

to estimate C (Sotiras et al., 2015):

$$\|X - CC^TX\|^2 \text{ subject to } C \geq 0, C^TC = I, \text{ and } W = C^TX \tag{4}$$

Where $\| \|^2$ represents the squared Frobenius norm and I denotes the identity matrix which

enforces orthogonality among C. C is first initialized using a non-negative double singular value

decomposition (Boutsidis & Gallopoulos, 2008). Then, C is updated through an iterative process

until it converges on an optimal solution. The iterative multiplicative update rule is as reported by

Yang and Oja (2010):

$$\hat{C}_{ij} = C_{ij} \frac{(XX^TC)_{ij}}{(CC^TXX^TC)_{ij}} \tag{5}$$

Where i represents the number of vertices and j represents the number of components. The

component matrix C represents the latent structure in the data and allows for an examination of

the underlying covariance in multivariate data. As done in Patel et al. (2020) the sparse and non-

overlapping component matrix allows for each vertex to be assigned an output component using a

winner take all method which improves the interpretability of the spatial output components. The

weight matrix W represents the subject-metric coefficients, allowing for an examination of subject-

specific and metric-specific contributions to each component.

2.8 Implementing OPNNMF

A total of 11 metrics were included in the OPNNMF implementation (ODI, NDI, T1w/T2w, FA,

MD, gyrification, thickness, curvature, AP cosine similarity, PD cosine similarity, IO cosine

similarity) with subsets of these metrics used for more specific analyses (i.e. NODDI only, DTI

only, macrostructure only, and cosine similarity only). The input matrix X was built using all

2004 vertices of the midthickness surface in unfolded space for all 11 metrics across all 100

subjects per hemisphere. That is, each subject contributed 11 unfolded space maps to the input

matrix. Thus, the input matrix had 1100 columns (100 subjects x 11 metrics - defined as subject-

metrics) and 2004 rows (2004 vertices) for a single hemisphere. Normalization was required

since the metrics exist on different scales. First, each metric was z-scored within each

hemisphere. Then, each z-scored metric distribution was shifted by the minimum value from all

the z-scored metrics to ensure all metrics were on the same scale and there were no negative

values. All distributions were manually inspected to ensure the minimum value used was not an

outlier.

OPNNMF was implemented using publicly available and open MATLAB code at

https://github.com/asotiras/brainparts (Sotiras et al., 2015; Yang & Oja, 2010; Boutsidis &

Gallopoulos, 2008; Halko et al., 2011). OPNNMF was run with a max number of iterations =

10000, tolerance = 0.00001, and non-negative double singular value decomposition initialization.

2.9 Stability & Reconstruction Error

The quantification of OPNNMF decomposition stability followed that of Patel et al. (2020).

Stability was assessed by examining the similarity of the spatial component matrix C across

varying splits of data. All 100 subjects were randomly split into two groups of equal sizes.

OPNNMF was then performed on each split independently. A within-split similarity matrix was

then derived by multiplying a particular splits component matrix by the transpose of itself (i.e. a

cosine similarity). The result is a 2004x2004 (number of vertices x number of vertices) matrix

where each row contains the cosine similarity of component scores between a vertex and all

other vertices. Finally, a Pearson's correlation coefficient was calculated across the rows of the

cosine similarity matrix between splits to quantify if the decomposition maintained the

relationships between vertices. The above process was repeated 6 times, each with a new random

split of the data. The mean and standard deviation of the correlation coefficient was taken across

all vertices and splits for a given component solution. A correlation of 1 represented perfect

stability (i.e. each split of the data had perfect correspondence between vertex relationships),

whereas -1 represented instability. The above process was then repeated for different component

decompositions, from k=2 to k=12. Reconstruction error was calculated through 3-steps. First,

the component matrix C and the weight matrix W were estimated and then multiplied together to

return the reconstructed input matrix. The original and reconstructed input matrix were then

subtracted to obtain a reconstruction error matrix. The Frobenius norm of the reconstruction

error matrix was then taken to get the reconstruction error. The gradient in the reconstruction

error was taken across solutions with varying component numbers to assess the magnitude of the

improvement in reconstruction error when adding more components.

2.10 Interpreting OPNNMF

The output component matrix C contains a component value for each vertex while the weight

matrix W describes how each subject-metric is projected onto each component. A large value in

the component matrix can be interpreted as a particular vertex being identified as a part of the

variance pattern. The weight matrix can be used to elucidate which metrics contributed to each

component as well as inter-subject variance within metrics. In the current study these 2 matrices

were used to explore spatial patterns and the contributions of particular metrics to each

component. A winner-take-all method was used where a vertex was assigned the integer of the

component with the highest component weighting value from the matrix C. The matrix W was z-

scored within each component row and then plotted to examine metric-specific trends.

3. Results

The results begin with qualitative descriptions of average macro- and microstructural measures

and their correlations on the midthickness hippocampal surface (middle of the hippocampal gray

matter). We also show the significance of these correlations using the developed hippocampal

spin test. We then present results from a logistic regression model used to examine the

separability of the subfields using microstructural features. We then show the cosine similarities

between NODDI microstructural vectors and the hippocampal axis vectors. Finally, we present

the OPNNMF results including a stability analysis and a 6-component solution.

3.1 Distributions of Hippocampal Metrics

Figure 2 presents Spearman's rho correlations, mean macro- and microstructural metrics, along

with subfield segmentations shown on an averaged hippocampal midthickness surface in folded

and unfolded space. The standard deviation of these metrics is shown in supplementary Figure 1.

The orientation dispersion (ODI - Figure 2C) is highest in the anterior and body of CA1 and the

distal parts of the subiculum, while dispersion is lowest in the body and posterior of DG/CA4,

CA3 and CA2 and at the most proximal edge of the subiculum. Neurite density (NDI - Figure

2D) is highest in the body and tail of the subiculum, while there is lower neurite density in CA1

and the DG/CA4. The T1w/T2w ratio has a strikingly similar distribution to that of NDI, as

found in previous cortical studies (Figure 2E; Fukutomi et al., 2018). T1w/T2w, ODI, and NDI

maps appear to vary across the subfields.

Macrostructure features of thickness, gyrification, and curvature are shown in Figure 2H-

J. Gyrification is largest in anterior CA1 and the DG/CA4. The thickest regions are the anterior

and posterior of the subiculum and CA1, as well as throughout the DG/CA4, while CA3 and

CA2 are thin. Curvature tends to be highest in the anterior part of the subiculum, along the spine

of the hippocampus (red arrows in Figure 2J), and in CA3. These findings are largely in line with

previous work (DeKraker et al., 2020).

Spearman's rho correlations between averaged maps of these metrics can be seen in the

lower triangle of Figure 2A while the false-discovery rate (FDR) corrected p-values derived from

the hippocampal spin test can be seen in the upper triangle. Similar correlations between all maps

at the subject level can be seen in supplementary Figure 2. The null distributions given by the

spin test using the averaged maps can be seen in supplementary Figure 3.

3.2 Correlations between NODDI metrics & T1w/T2w

In Figure 2D and E, strong qualitative similarities can be seen between NDI and T1w/T2w. Both

are high in the subiculum and CA3/CA2 regions, while hypointensities are noted in CA1. A

significant spatial overlap is seen between the positively correlated NDI and T1w/T2w maps (ρ

= 0.86, FDR corrected p-value < 0.05). ODI (Figure 2C) and T1w/T2w (Figure 2D) are also

significantly overlapping and are negatively correlated (ρ = -0.61, FDR corrected p-value <

0.05). A similar correlation has also been noted across the entire cortex (Fukutomi et al., 2018).

3.3 Correlations between NODDI & DTI metrics

Qualitatively, the map of FA (Figure 2F) resembles the NDI map (Figure 2D) and the inverse of

the ODI map (Figure 2C). Particularly, the distinct pattern of high dispersion equates to low FA,

while high neurite density equates to high FA. Furthermore, the region of low neurite density

(Figure 2D) in the body of CA1 and CA2 corresponds to a region of high MD (Figure 2G). ODI

and FA have a significant spatial overlap and are negatively correlated (ρ = -0.88, FDR corrected

p-value < 0.05). Furthermore, NDI and FA are significantly overlapping and are positively

correlated (ρ = 0.75, FDR corrected p-value < 0.05) and NDI and MD are significantly

overlapping and negatively correlated (ρ = -0.46, FDR corrected p-value < 0.05).

A disentangling of FA as determined by ODI and NDI has been reported previously in

cortical gray matter (Zhang et al., 2012). Two voxels with different neurite densities can have the

same FA if the one with the larger neurite density also has the larger dispersion (Zhang et al.,

2012). That is, anisotropy as measured with FA conflates neurite density and orientation

dispersion. We report a similar level of disentangling within the hippocampus in supplementary

Figure 4, where within any bin of FA there is a range of potential ODI and NDI values.

3.4 Correlations between macrostructure and NODDI metrics

The only significant overlap between a macrostructural and microstructural map is gyrification

and T1w/T2w, which are negatively correlated (ρ = -0.46, FDR corrected p-value < 0.05).

Regions of high gyrification in CA1 spatially correspond to regions of low T1w/T2w content.

While there are some other notable correlations, such as a moderate positive correlation between

gyrification and ODI (ρ = 0.45) and a moderate negative correlation between gyrification and

FA (ρ = -0.49), none were significantly overlapping. Correlations between NODDI and

macrostructural measures at the subfield-averaged level can be seen in supplementary Figure 5.

Figure 2. Correlations and plots of mean macro- and microstructure metrics on averaged

hippocampal midthickness surfaces in folded and unfolded space for left and right hemispheres.

(A) Lower triangle of the heatmap shows the Spearman's rho correlations of the average maps

after combining both left and right hemispheres. The upper triangle represents the false-

discovery rate corrected p-values derived from the hippocampus spin testing using 2500

permutations. Note that the colour bar is inverted such that any brighter component of the

heatmap corresponds to a significant p-value. (B) Left and right hippocampal subfields from a

manual segmentation of a histological reference (Ammunts et al., 2013; DeKraker et al., 2020).

Unfolded space is shown in the same orientation for left and right hemispheres. DG - Dentate

Gyrus, CA - Cornu Ammonis. (C,D) Orientation Dispersion Index (ODI) and Neurite Density

Index (NDI) from NODDI. White lines represent subfield borders shown in (B). (E) T1w/T2w

ratio. (F,G) Diffusion Tensor Imaging metrics of Fractional Anisotropy (FA) and Mean

Diffusivity (MD - m^2/s). (H-J) Macrostructure measures of thickness, gyrification, and

curvature. (J) Red arrows highlight the highly curved "spine" of the hippocampus.

3.5 Evaluating the variability of the microstructural metrics across the subfields

To examine if the microstructural metrics used here could provide differentiability to the

subfields, we trained a simple logistic regression classifier with L2 regularization on subfield

averaged microstructural measures using the scikit-learn (version 1.1.2) package in python

(Pedregosa et al., 2011). 70 subjects were included in the training set, while 30 subjects were

included in the testing set (both hemispheres were combined). ODI, NDI, and T1w/T2w

averages across the midthickness surface within each of the 5 subfields per hemisphere were

obtained within each subject in both the training and testing set. A logistic regression model was

then trained to predict the subfield label given the subfield averaged microstructural metrics.

This model was then tested on the unseen data from the testing set to quantify if the

microstructural metrics are differentiable across the subfields. Examining all the data points for

ODI, NDI, and T1w/T2w, colour coded by subfield class, it appears that the subfields present as

relatively separable clusters apart from CA2 which appears to have lots of microstructural

overlap with the other subfields (Figure 3A). The same clustering can be seen in the test set

(Figure 3B) left out to evaluate the performance of the logistic regression classifier (Figure 3C).

Using a confusion matrix and F1-scores defined as (2*precision*recall) / (precision+recall), it

can be seen that the simple logistic regression classifier was able to perform well in delineating

the subiculum (F1-score = 0.87), CA1 (F1-score = 0.85), CA3 (F1-score = 0.69), and the

DG/CA4 (F1-score = 0.83) using microstructural measures of ODI, NDI, and T1w/T2w (Figure

3D). However, CA2 was not as differentiable from the rest of the subfields using these metrics,

with an F1-score of 0.53. Furthermore, we found that ODI and NDI generally had higher F1-

scores and greater separability across the subfields then the DTI metrics of FA and MD

(supplementary Figure 6).

Figure 3. The performance of a logistic regression classifier to capture microstructural

variability across the subfields. (A) Subfield averaged ODI, NDI, and T1w/T2w from all subjects

and hemispheres combined coloured by subfield label. (B) Test set data from 30 subjects

coloured by subfield. (C) Logistic regression classified labels on the test set seen in (B). (D)

Confusion matrix and subfield-specific F1-scores to evaluate logistic regression performance on

classifying the subfields using the microstructural measures of the test set.

3.6 Examination of the primary direction of diffusion relative to hippocampal axes

This section qualitatively analyzes the mean of the cosine similarities (Figure 1D; Figure 4)

between the hippocampal vectors along the anterior-posterior (AP), proximal-distal (PD), and

inner-outer (IO) (Figure 1A) axes and the NODDI microstructural vectors along the

midthickness surface (Figure 1C).

High AP alignment can be seen in the body of the DG to CA3, as well as in the

subiculum. Furthermore, there is relatively little AP oriented diffusion in CA1.

High PD oriented diffusion can be seen in the head of CA3, CA2, CA1, and in the body

of CA1. There was little PD oriented diffusion in the body of DG/CA4, CA3, CA2, and the

subiculum.

High IO alignment can be seen in the tail of the subiculum and CA1, as well as

throughout the body of CA1, while low IO oriented diffusion can be seen in the body of the

DG/CA4, CA3, CA2, and the subiculum.

The cosine similarities varied the greatest in CA1 and the subiculum across AP, PD, and

IO directions (supplementary Figure 7). Noticeable differences in the orientation of diffusion

across the subfields can be seen in supplementary Figure 8.

Figure 4. Mean of the cosine similarities between hippocampal axis vectors and NODDI

microstructural vectors in the left and right hemisphere along the midthickness surface. High

cosine similarities correspond to a high alignment of the NODDI microstructural vector along

that particular hippocampal axis. (A) Distribution of cosine similarities along the anterior-

posterior direction. (B) Distribution of cosine similarities along the proximal-distal (tangential)

direction. (C) Distribution of cosine similarities along the inner-outer (laminar) direction.

3.7 Stability Analysis

The results of the stability analysis can be seen in Figure 5. Figure 5A presents the stability and

the gradient in the reconstruction error using all the metrics combined that are shown in Figure

5B-E. The goal of the stability analysis was to elucidate the largest component value that was

still stable and provided a relative gain in reconstruction error. In Figure 5A it can be seen that a

component solution of k=6 has good stability with relatively low standard deviation.

Comparatively, decomposing into a larger number of components decreases the stability of the

OPNNMF solution. As well, k=6 does provide a relative gain in reconstruction error, although

the largest gain in reconstruction error occurs when moving from a k=2 to a k=3 solution. The

stability analysis suggests that k=6 is the highest component value that is largely stable, thus we

use this for the decomposition results using all metrics.

Figure 5. Stability coefficient and the gradient in reconstruction error based on the number of

components used for the OPNNMF solution. Filled in circles plus solid lines are the left

hemisphere and filled in squares plus dotted lines are the right hemisphere. Error lines show +/-

1 SD. (A) Stability coefficient (blue) and the gradient in reconstruction error (red) as a function

of the number of components using all metrics for NMF in B-E. (B) Stability coefficient for

NODDI metrics (ODI and NDI) plus T1w/T2w. (C) Stability coefficient for DTI metrics (FA

and MD) plus T1w/T2w. (D) Stability coefficient for macrostructure metrics (gyrification,

thickness, curvature). (E) Stability coefficient for cosine similarities (AP, PD, and IO). Points

between hemispheres are slightly offset along the x-axis so that error bars are visible.

Another goal of the stability analysis was to compare the stability of the decomposition

using all metrics versus using smaller groupings of metrics, such as NODDI (ODI and NDI) plus

T1w/T2w. Comparing Figure 5A with Figure 5B-E, it can be seen that for almost all component

values the all metric solution tends to be more stable than any of the smaller metric groupings.

This is especially true for the larger component values above k=6. These results suggest that the

use of multiple metrics results in more stable parcellations, as found in Patel et al. (2020).

The 6-component solution using all metrics is presented in Figure 6 for both the left and

right hippocampus. 4, 5, 6, and 7-component solutions using all the metrics can be found in

supplementary Figure 9. 4-component solutions for all smaller metric combinations shown in

Figure 5B-E can be found in supplementary Figure 10.

3.8 Description of the 6-component Solution

Figure 6A depicts the winner-take-all method applied at each vertex in folded and unfolded

space for 6-components. Figure 6B shows the z-scored subject-metric weight matrices. In the

following paragraph we describe the first 3 components including their location relative to the

subfields (proximal-distal/medial-lateral axes) as well as along the anterior-posterior

(longitudinal) axis. We then briefly describe components 4-6. We also describe the features

which contribute to each component. The left and right hippocampus do have similar covariance

patterns although the component ordering differs. For example, the vertices of left component 1

correspond to the vertices of right component 2. To simplify our descriptions, we take the

ordering of the left hippocampus for the rest of the paper, and we have made this adjustment

such that the difference in ordering does not need to be considered for Figure 6 (Patel et al.,

2020).

Component 1 is characterized by a cluster of vertices through the body and tail of the

most proximal edge of the subiculum and through the body and tail of CA3. This component

spans around the bottom two-thirds of the hippocampus across its anterior-posterior axis.

Component 1 is characterized by high NDI, T1w/T2w, FA, and AP cosine similarity, with lower

ODI, PD and IO cosine similarity. This may reflect the large anisotropic AP oriented fiber

bundles that are myelinated such as the cingulum bundle for the proximal edge of the subiculum

and the fimbria for CA3.

Component 2 is characterized by vertices that are present only in the body or middle one-

third along the anterior-posterior axis of CA1. This component is characterized by high ODI,

gyrification, and PD and IO cosine similarities. This likely reflects a high heterogeneity in fiber

orientation in CA1.

Component 3 is characterized by vertices that cross the subiculum and CA1 in a

proximal-distal fashion in the head of the hippocampus, as well as vertices that span the anterior-

posterior body of the hippocampus at the border between the subiculum and CA1. This

component is characterized by high ODI, thickness, and IO cosine similarity.

Component 4 corresponds to a large posterior cluster that stretches into the body of CA1

characterized by a moderate range of metrics with the largest being the PD cosine similarity and

FA. Component 5 corresponds to a cluster of vertices in the body of CA2 and CA1 and is

represented by a large MD, curvature, and AP cosine similarity and low NDI, T1w/T2w, FA, and

thickness. Interestingly, component 6 is characterized by a thin cluster of vertices roughly

corresponding to the anterior and body of the DG/CA4 region. This was represented by high

macrostructural measures of thickness, gyrification, and curvature and low weights for all other

metrics.

A general pattern is noticed when examining the whole 6-components rather than looking

at its parts. Generally, more parcellations exist along the proximal-distal direction than do in the

anterior-posterior direction.

Figure 6. 6-component NMF solution on the midthickness surface for left and right hippocampi. (A) Winner-take all output at each vertex shown in folded and unfolded space. White lines denote subfield borders. (B) Z-scored subject-metric weight matrices across each of the 6 components, denoting the z-scored contribution of each metric to each component. AP, PD, and IO represent the 3 cosine similarity metrics.

4. Discussion

In the current study we examined the microstructure of the hippocampus using the in vivo HCP

dMRI and structural data, along with a novel surface-based method for subfield segmentation. We

found that ODI was highest in the CA1 subfield, likely capturing the large heterogeneity of

tangential and radial fibers. NDI and T1w/T2w were found to be strongly correlated and were

highest in the subiculum and lowest in CA1 and the DG/CA4, suggesting that NODDI is likely

sensitive to the myelin content of the hippocampus. Using these microstructural measures, we

found that the cytoarchitectonic defined subfields were largely separable using a simple logistic

regression model. OPNNMF components appeared to capture unique co-varying clusters within

the hippocampus, with high medial-lateral variability. Finally, we showed distinct regions of

similar microstructural orientations by examining the main direction of diffusion relative to the

three hippocampal axes, which may correspond to specific microstructural properties.

4.1 Dispersion of neurites in the hippocampus may reflect heterogeneous radial and tangential

neurite components

The Orientation Dispersion Index (ODI) is meant to characterize the variation in diffusion

orientations around a single dominant direction at every voxel. A previous study using ODI and

patch-wise circular variance measured using histology (measures variability in neurite

orientations) has shown that both measures have lower dispersion in demyelinated lesions in

patients with multiple sclerosis, where there is reduced geometrical complexity of neurites

(Grussu et al., 2017). The hippocampal gray matter has a general distribution of microstructure

that is similar to the neocortex, with tangential (proximal-distal) and radial (inner-outer)

components that follow the curvature of the hippocampus. In the current study we showed that

CA1 had the largest ODI, suggesting that it had the largest heterogeneity in neurite orientations.

CA1 has large tangential neural processes, like the Schaffer collaterals and perforant path, as

well as a large (yet dispersed) radial pyramidal cell layer (Duvernoy et al., 2013). By measuring

the orientation of the main direction of diffusion relative to the three hippocampal axes in CA1

(Figure 4), we found either high tangential or radial diffusion, supporting the idea that ODI

reflected the heterogeneity of these components. Conversely, ODI was lower in DG/CA4, CA3,

and at the most proximal edge of the subiculum. In these regions the primary diffusion direction

was minimally tangential or radial, and was mainly anterior-posterior or oblique with large

macroscopic diffusion anisotropy (i.e. large NDI and FA). The apparent reduction in

orientational heterogeneity (less radial/tangential components) and a resulting increase in

anisotropy may potentially explain the low ODI in these regions. In the DG/CA4 and CA3

region this could be a result of partial voluming with the nearby fimbria, and in the subiculum it

could be due to partial voluming with the nearby cingulum bundle or the perforant path at its

most proximal edge (supplementary Figure 11). As hypothesized in the rest of the cortex

(Fukutomi et al., 2018), it is likely that ODI in the hippocampal gray matter is largely driven by

the heterogeneity of radial and tangential neurite components.

4.2 Hippocampal neurite density is highly correlated with T1w/T2w

The distribution of the NDI and T1w/T2w across hippocampal gray matter was similar, as seen

in Figure 2D and E and as shown by their strong positive correlation and significant overlap.

While the diffusion signal is generally agnostic to water within myelin, previous work has shown

that myelinated axons restrict diffusion to a greater degree than unmyelinated axons (Behrens &

Johansen-Berg, 2014). This increase in restriction due to myelin would result in an increase in

NDI since there is more "stick" like diffusion occurring (i.e., a monoexponentially decaying

signal with a slope defined by the parallel diffusivity). Furthermore, due to the short exchange

time of water within dendrites and glia to the extra-cellular space, it is likely that any restricted

diffusion would be reflective of myelinated axons (Jespersen et al., 2010). This suggests that

NDI reflected the myelin content of the hippocampus. The T1w/T2w content and NDI were

largest in the body and tail of the subiculum. High myelin content in the subiculum has been

noted previously with histology (Ding & Van Hoesen, 2015). Furthermore, it is likely that the

white matter of the cingulum bundle or perforant path contribute to the large T1w/T2w content

seen in the subiculum. Conversely, T1w/T2w and NDI were lower in CA1, which is likely a

result of a relatively sparse layer of pyramidal cells along the midthickness surface or the largely

unmyelinated Schaffer collaterals (Jürgen et al., 2011; Szirmai et al., 2012). Overall, the

distribution of T1w/T2w found here agreed with previous studies (DeKraker et al., 2018;

Ábrahám et al., 2012). A strong positive correlation between NDI and T1w/T2w was found

previously across the cortex. However, the hippocampus was found to have high values in NDI

but low values of T1w/T2w when compared to the rest of the cortical areas (Fukutomi et al.,

2018). Here we showed that a strong correlation between NDI and T1w/T2w still exists in the

hippocampus when comparing them at a finer spatial scale. This correlation is further

corroborated by another cortical study at high ex vivo resolutions in the rodent brain, in which

cortical NDI was strongly correlated with staining intensity of myelinated axons (Jespersen et al.,

2010). Histological work has found similar correlations in white matter, where myelin content

was found to be strongly correlated with axon count (Schmierer et al., 2007). However, a recent

study utilizing a multicomponent relaxometry method for imaging myelin water fraction found

no significant correlation between myelin and NDI measured using NODDI in most white matter

structures (Qian et al., 2020). While NDI and T1w/T2w as a proxy of myelin do appear to be

correlated in gray matter including the hippocampus, further work is needed to examine this

correlation in other white matter structures, including white matter surrounding the hippocampus

such as the fimbria, fornix, and alveus.

4.3 Microstructure metrics systematically vary across the subfields

Microstructural metrics such as intra-cortical myelin and macrostructural cortical thickness have

been shown to be useful in parcellating the neocortex into subregions (Nieuwenhuys, 2013;

Glasser et al., 2014; Glasser & Van Essen, 2011). Furthermore, using non-negative matrix

factorization of T1w/T2w, MD, and FA it was found that a 4-component solution qualitatively

resembled hippocampal subfield borders (Patel et al., 2020), suggesting that T1w/T2w and

microstructure may provide sufficient separability to parcellate hippocampal subfields. In the

current study, by examining their averaged maps, we found that T1w/T2w, NDI, and ODI

appeared to be separable qualitatively across the subfields (Figure 2). Quantitatively, we showed

that a simple logistic regression model performed well in predicting the subfield label based on

the subfield-averaged metrics of T1w/T2w, NDI, and ODI. This result suggests that these

microstructural measures are potentially sensitive to the known microstructural differences

across the subfields as defined via cytoarchitectonics. Critically, it appeared that NDI and ODI

provided more separability across the subfields then FA and MD, suggesting that NODDI may

be more useful than DTI in capturing known microstructural differences across subfields

(supplementary Figure 6). CA2 appeared to be the exception, with a distinct pattern of MD when

compared to the other subfields, while this separability of CA2 was not captured with NODDI

metrics (supplementary Figure 6). Myelin content has been demonstrated previously to closely

correspond to averaged subfield borders (DeKraker et al., 2018). To a lesser extent,

macrostructure appeared to also follow the subfield borders, which has been noted previously for

thickness (DeKraker et al., 2018). While thickness is consistently low in CA3 and CA2, and

gyrification is consistently high in CA1, these measures alone may not differentiate all subfield

boundaries. Thus, a combination of NODDI and macrostructural measures may provide

complimentary information needed for subject-specific subfield delineation.

4.4 The orientation of diffusion relative to the hippocampus may be useful in identifying

hippocampal microstructure

We quantified the main direction of diffusion relative to the 3 main hippocampal axes which

microstructure tends to align closely with. Here we provide descriptions of microstructure

(Figure 1B) that likely contribute to the orientation results. The following three paragraphs

describe the anterior-posterior (AP), proximal-distal (PD), and inner-outer (IO) alignment,

respectively.

The high AP alignment in the body of the DG to CA3 was likely driven by the

neighbouring fimbria, the largest bundle in that region oriented AP. High AP alignment in the

subiculum was likely caused by the cingulum, a large fiber bundle that traverses the

parahippocampal gyrus. Some partial voluming from the outer (where the cingulum exists) to the

midthickness surface was expected, which may drive this alignment.

High PD alignment in the head of CA3 was expected to be either Schaffer collaterals

which curve immediately PD off of the apical dendrites of the pyramidal cells or from perforant

projections coming from the entorhinal cortex and entering CA3 which are also oriented PD.

High PD alignment in CA1 was likely a result of the Schaffer collaterals. The Schaffer

collaterals make synaptic contact at the apical and basal dendrites of CA1 in a PD fashion

(Nieuwenhuys et al., 2008; Swanson et al., 1978). However, the perforant path could have also

contributed to a higher PD alignment as it moves from the entorhinal cortex to the DG synapsing

on CA1 along the way (Nieuwenhuys et al., 2008). Furthermore, a high PD cosine similarity in

the CA3 region could have been a result of partial voluming with the alveus, a highly PD

oriented bundle that sits atop the hippocampal gray matter.

High IO alignment seen in CA1 was likely a result of the pyramidal neurons. The

pyramidal somas exist in the stratum pyramidale layer of the midthickness surface, and are

generally scattered in CA1 (Nieuwenhuys et al., 2008). Their axons and basal dendrites move IO

towards the alveus/outer surface and their large apical dendrites move IO towards the stratum

radiatum/inner surface. All IO alignment seen in CA1 was expected to be caused by the

pyramidal neurons or other afferent CA1 paths such as the Schaffer collaterals which may curve

IO before making contact with the apical dendrites of the pyramidal neurons. High IO alignment

in the subiculum was also likely caused by pyramidal neurons as in CA1.

The cosine similarities across subjects varied the greatest in CA1 and the subiculum across AP,

PD, and IO directions (supplementary Figure 7).

Typical hippocampal microstructural analyses average scalar diffusion metrics (such as

FA, MD, NDI, etc.) either across whole hippocampi (van Uden et al., 2015; Salmenpera et al.,

2006) or whole subfields (Radhakrishnan et al., 2020), which are inherently non-specific towards

microstructure that exists within and across subfields and the hippocampal long-axis. However,

tractography analyses which aim to capture the continuous intra-hippocampal circuitry are

difficult to perform, as at lower resolutions, tracts can be spurious requiring complex acquisition

and correction schemes (Zeineh et al., 2012). The orientational analyses described here have the

potential to increase specificity at in vivo resolutions by leveraging the known anatomical

orientations of hippocampal microstructure. Furthermore, capturing the essence of hippocampal

microstructural orientations with vertex-wise scalar values can make qualitative observations and

statistical analyses more tenable than with the complex 3D orientations provided by

tractography. Future studies could relate the hippocampal axis vectors to other dedicated

methods of diffusion orientation representation that have the ability to faithfully represent more

complex fiber configurations.

Applications of the proposed orientational methods may be useful to identify

microstructure deterioration in disease states, where affected microstructure may be less

prominent, and may appear as smaller cosine similarities along a particular axis. For example,

perforant path lesions in rats caused rapid memory loss which was akin to early-stage

Alzheimer's disease (Kirkby & Higgins, 2001). A 2010 study found deterioration of the

perforant path in aged humans using diffusion tensor imaging (Yassa et al., 2010). Perforant path

degradation should result in less attenuation of the diffusion signal along its length, which may

potentially show up as smaller PD cosine similarities specifically in the subiculum, CA3, and

CA1, as there should be less PD oriented diffusion. This may be possible for other neurological

diseases where specific microstructure is affected, such as pyramidal neuron degradation which

should result in smaller IO cosine similarities. However, to draw such conclusions, further ex

vivo validation with the ability to measure more ground-truth microstructural orientations will be

essential to evaluate the usefulness of this method. As well, future studies will have to evaluate

the efficacy of this method at clinical resolutions.

4.5 6-component OPNNMF solution displays distinct co-varying regions of macro- and

microstructure

The hippocampus is believed to have two main interacting dimensions of organization along its

medial-lateral/proximal-distal or subfield axis, and across its long or anterior-posterior axis (see

Genon et al., 2021 for review). In the current study with a 6-component OPNNMF solution we

found a gradient in the clusters along the proximal-distal direction, suggesting that regions along

this axis had disparate macro- and microstructural properties. That is, the macro- and

microstructural "axes" of the hippocampus as defined by the metrics used here tended to vary

along the same axis that the subfields were defined. Variability along this axis was expected, as

the subfields show differences in morphology, cytoarchitectonic profiles (Duvernoy et al., 2013;

Ding & Van Hoesen, 2015), and connectivity (Andersen et al., 1971), which likely manifest as

changes in macro- and microstructure. However, it is notable that a wide array of disparate

metrics at in vivo resolutions show such patterns. Recently there has been interest in the long-

axis organization of the hippocampus, with evidence coming from anatomical and physiological

recordings in rodents (Chase et al., 2015). In the current study there appeared to be a smaller

gradient in the clusters along the anterior-posterior axis, suggesting that the metrics used here

varied less along the long-axis when compared to the proximal-distal axis. While functional

studies have found stark variability across the long axis, it is unclear to the extent which the

hippocampal microstructure varies across this axis (Genon et al., 2021, Plachti et al., 2019).

Future work could look to examine the variability of dMRI measures across the long axis at

higher resolutions.

4.6 Limitations

There are some limitations of the current research that largely pertain to the metrics used here.

As noted in the introduction, DTI aims to capture the macroscopic diffusion anisotropy at each

voxel, assuming that the diffusion process can be well characterized by a Gaussian distribution.

Due to its popularity in both clinical and research settings, we sought to examine the distribution

of two common DTI measures (FA and MD) in the hippocampus. However, in most gray matter

regions there tends to be a complex arrangement of fiber orientations that cannot be well

modelled via a single tensor. This can result in drastically understated FA values with generally

spherical or planar DTI ellipsoids which contain minimal information on diffusion orientations

(Campbell et al., 2005). While DTI has been used extensively in the hippocampus, it has

generally been purported to be sensitive but not specific to its microstructure (Coras et al., 2014).

In ex vivo tissue at microscopic resolutions DTI metrics of FA and MD have shown good

contrast to the hippocampal laminae, demonstrating their sensitivity (Coras et al., 2014;

Shepherd et al., 2007; Stolp et al., 2018). However, at lower in vivo resolutions used here a

plethora of microstructure partial voluming within each voxel is inevitable, which leads to a

diffusion signal that may show little macroscopic diffusion anisotropy and thus cannot be

captured by a single tensor. Indeed, here we found that even the largest FA values of the

hippocampus in regions of purportedly high T1w/T2w tended to range around 0.28, a relatively

low amount of macroscopic anisotropy. Furthermore, it is unclear the extent to which extra-

hippocampal white matter such as the fimbria and angular bundle might influence our measures

sampled along the middle of the hippocampal gray matter.

The NODDI model has also been used extensively in both healthy and diseased states to

provide metrics that are biologically grounded. Such work involved explicitly modelling the

diffusion signal as a sum of compartments representing varying tissue geometries (Zhang et al.,

2012). However, some assumptions of the NODDI model are likely not valid in practice. The

tortuosity assumption that links the extra-cellular perpendicular diffusivity to the axon volume

fraction does not hold for tight axon packings (Jelescu et al., 2016). The NODDI model also sets

the intra-axonal parallel diffusivity equal to the extra-axonal parallel diffusivity. However, the

intra-axonal parallel diffusivity has been shown to be much higher than the extra-axonal parallel

diffusivity (Jelescu et al., 2016; Novikov et al., 2018). While we altered the fixed NODDI

diffusivity values to reflect what has been found in gray matter, the modelling degeneracies

mentioned here could influence the NODDI metrics, such that they may not correspond to the

biophysical reality of the tissue. To this end, there has been recent work which has sought to re-

structure the basic microstructural building blocks used for white matter modelling to improve

characterization of the gray matter. Two primary examples of this include a model which

accounts specifically for the presence of the soma (Palombo et al., 2020) and a model which

considers the water exchange across the cell membranes (i.e., from neurites to extra-cellular

space) (Jelescu et al., 2022). Future work should look to apply gray matter specific models in the

hippocampus.

Finally, the T1w/T2w ratio as a proxy for myelin content was used in the current study

based on previous work that demonstrated good correspondence with cortical patterns of myelin

distribution (Glasser & Van Essen 2011). However, recently it has been suggested that the

T1w/T2w ratio is a suboptimal proxy of myelin in the subcortex (Arshad et al., 2017). At

present, it is unclear whether the T1w/T2w ratio is also a suboptimal marker of myelin for

archicortical areas such as the hippocampus. Future work should look to investigate the use of

other techniques derived from quantitative MRI including magnetization transfer and myelin

water imaging to get an improved measure of myelin content within the hippocampus (Tardif et

al., 2016).

5. Conclusion

In the current study we show distinct in vivo microstructural distributions and orientations within and across the hippocampal subfields, something that has not been investigated with comparable granularity up to this point. Furthermore, we provide context for the use of surface-based approaches to investigate hippocampal microstructure.

Our findings have several important implications for future work. The hippocampus is particularly vulnerable to certain neurological diseases such as Alzheimer 's disease and epilepsy, in which it is often one of the earliest aberrant structures (Dhikav et al., 2012). Examining the microstructure of the hippocampus at fine spatial resolutions in the simplified unfolded space, as done in this study, may provide potentially useful markers of hippocampal integrity. Furthermore, we noticed relatively large radial and tangential components of diffusion mainly in CA1 and the subiculum. Future work could attempt to tease apart these two orientationally distinct populations, providing estimates which may be useful to examine microstructurally specific deterioration. Furthermore, using the same orientation methods in this study, future work should focus on capturing multiple microstructure orientations as the hippocampus contains a complex configuration of fiber orientations. Future work could also relate all the identified OPNNMF components to demographic and cognitive variables to identify if there is a relationship between variability in cognitive performance and variability in the metrics used in this study. Finally, the microstructural metrics observed in this study appear to show good separability across hippocampal subfields, suggesting they may be sensitive to the underlying cyto- and myeloarchitectonic differences.

Acknowledgements

This work was supported in part by funding provided by Brain Canada, in partnership with Health Canada, for the Canadian Open Neuroscience Platform initiative. BK is supported by a post-graduate scholarship from the Natural Sciences and Engineering Research Council of

Canada (NSERC). JD is supported by a postdoctoral fellowship NSERC grant. ARK was

supported by the Canada Research Chairs program (#950-231964), NSERC Discovery Grant

(#6639), and Canada Foundation for Innovation (CFI) John R. Evans Leaders Fund project

(#37427), the Canada First Research Excellence Fund, and Brain Canada. ARK and SK were

supported by a Canadian Institute for Health Research grant (CIHR Project grant #366062). SK

is supported by a NSERC discovery grant (#05770). This research was enabled in part by the

support provided by the Digital Research Alliance of Canada. Data were provided [in part] by

the Human Connectome Project, WU-Minn Consortium (Principal Investigators: David Van

Essen and Kamil Ugurbil; 1U54MH091657) funded by the 16 NIH Institutes and Centers that

support the NIH Blueprint for Neuroscience Research; and by the McDonnell Center for Systems

Neuroscience at Washington University.

References

Ábrahám, H., Vincze, A., Veszprémi, B., Kravják, A., Gömöri, É., Kovács, G. G., & Seress, L. (2012). Impaired myelination of the human hippocampal formation in down syndrome. *International Journal of Developmental Neuroscience*, *30*(2), 147–158. https://doi.org/10.1016/j.ijdevneu.2011.11.005

Alexander-Bloch, A. F., Shou, H., Liu, S., Satterthwaite, T. D., Glahn, D. C., Shinohara, R. T., Vandekar, S. N., & Raznahan, A. (2018). On testing for spatial correspondence between maps of human brain structure and function. *NeuroImage*, *178*, 540–551. https://doi.org/10.1016/j.neuroimage.2018.05.070

Amunts, K., Lepage, C., Borgeat, L., Mohlberg, H., Dickscheid, T., Rousseau, M.-E., Bludau, S., Bazin, P.-L., Lewis, L. B., Oros-Peusquens, A.-M., Shah, N. J., Lippert, T., Zilles, K., & Evans, A. C. (2013). Bigbrain: An ultrahigh-resolution 3d human brain model. *Science*, *340*(6139), 1472–1475. https://doi.org/10.1126/science.1235381

Andersen, P., Bliss, T. V. P., & Skrede, K. K. (1971). Lamellar organization of hippocampal excitatory pathways. *Experimental Brain Research*, *13*(2). https://doi.org/10.1007/bf00234087

Andersson, J. L. R., & Sotiropoulos, S. N. (2015). Non-parametric representation and prediction of single- and multi-shell diffusion-weighted MRI data using Gaussian processes. *NeuroImage*, *122*, 166–176. https://doi.org/10.1016/j.neuroimage.2015.07.067

Arshad, M., Stanley, J. A., & Raz, N. (2017). Test-retest reliability and concurrent validity of in vivo myelin content indices: Myelin water fraction and calibrated t1w/t2W image ratio. *Human Brain Mapping*, *38*(4), 1780–1790. https://doi.org/10.1002/hbm.23481

Assaf, Y., & Basser, P. J. (2005). Composite hindered and restricted model of diffusion (charmed) MR imaging of the human brain. *NeuroImage*, *27*(1), 48–58.

10.1016/j.neuroimage.2005.03.042

Assaf, Y., Blumenfeld-Katzir, T., Yovel, Y., & Basser, P. J. (2008). Axcaliber: A method for

measuring Axon diameter distribution from diffusion MRI. *Magnetic Resonance in*

Medicine, *59*(6), 1347–1354. 10.1002/mrm.21577

Assaf, Y., & Cohen, Y. (2000). Assignment of the water slow-diffusing component in the central

nervous system using Q-Space Diffusion Mrs: Implications for Fiber Tract Imaging.

Magnetic   Resonance   in   Medicine,   *43*(2),   191–199.

https://doi.org/10.1002/(sici)1522-2594(200002)43:2<191::aid-mrm5>3.0.co;2-b

Basser, P. J., Mattiello, J., & LeBihan, D. (1994). Mr diffusion tensor spectroscopy and

imaging. *Biophysical Journal*, *66*(1), 259–267.

https://doi.org/10.1016/s0006-3495(94)80775-1

Beaujoin, J., Palomero-Gallagher, N., Boumezbeur, F., Axer, M., Bernard, J., Poupon, F., . .

. Poupon, C. (2018). Post-mortem inference of the human Hippocampal connectivity

and microstructure using ultra-high field diffusion MRI at 11.7 T. *Brain Structure*

and Function, 223(5), 2157-2179.

Behrens, T., & Johansen-Berg, H. (2014). *Diffusion Mri: From quantitative measurement to*

in vivo neuroanatomy. Elsevier.

Bendel, O., Bueters, T., Euler, M. V., Ögren, S. O., Sandin, J., & Euler, G. V. (2005).

Reappearance of Hippocampal CA1 Neurons after Ischemia is Associated with

Recovery of Learning and Memory. *Journal of Cerebral Blood Flow & Metabolism,*

25(12), 1586-1595. 10.1038/sj.jcbfm.9600153

Blümcke, I., Coras, R., Miyata, H., & Özkara, C. (2012). Defining clinico-neuropathological

subtypes of Mesial Temporal Lobe Epilepsy with   hippocampal sclerosis. *Brain*

Pathology, *22*(3), 402–411. https://doi.org/10.1111/j.1750-3639.2012.00583.x

Boutsidis, C., & Gallopoulos, E. (2008). SVD based initialization: A head start for nonnegative

matrix factorization. *Pattern Recognition*, *41*(4), 1350–1362.

https://doi.org/10.1016/j.patcog.2007.09.010

Burggren, A. C., Zeineh, M. M., Ekstrom, A. D., Braskie, M. N., Thompson, P. M., Small, G.

W., & Bookheimer, S. Y. (2008). Reduced cortical thickness in hippocampal subregions

among cognitively normal apolipoprotein e E4 carriers. *NeuroImage*, *41*(4), 1177–1183.

https://doi.org/10.1016/j.neuroimage.2008.03.039

Campbell, J. S. W., Siddiqi, K., Rymar, V. V., Sadikot, A. F., & Pike, G. B. (2005). Flow-based

fiber tracking with diffusion tensor and Q-ball data: Validation and comparison to

principal diffusion direction techniques. *NeuroImage*, *27*(4), 725–736.

https://doi.org/10.1016/j.neuroimage.2005.05.014

Chase, H. W., Clos, M., Dibble, S., Fox, P., Grace, A. A., Phillips, M. L., & Eickhoff, S. B.

(2015). Evidence for an anterior–posterior differentiation in the human hippocampal

formation revealed by meta-analytic parcellation of fmri coordinate maps: Focus on the

Subiculum. *NeuroImage*, *113*, 44–60. https://doi.org/10.1016/j.neuroimage.2015.02.069

Cheng, H., Zhu, H., Zheng, Q., Liu, J., & He, G. (2020). Functional parcellation of the

hippocampus by semi-supervised clustering of resting state fmri data. *Scientific Reports*,

10(1). https://doi.org/10.1038/s41598-020-73328-1

Cherubini, E., & Miles, R. (2015). The CA3 region of the hippocampus: How is it? What

is it for? How does it do it? *Frontiers in Cellular Neuroscience, 9*.

10.3389/fncel.2015.00019

Coras, R., Milesi, G., Zucca, I., Mastropietro, A., Scotti, A., Figini, M., Mühlebner, A.,

Hess, A., Graf, W., Tringali, G., Blümcke, I., Villani, F., Didato, G., Frassoni, C.,

Spreafico, R., & Garbelli, R. (2014). 7T MRI features in control Human hippocampus

and hippocampal sclerosis: An ex vivo study with histologic correlations. *Epilepsia*,

55(12), 2003–2016. https://doi.org/10.1111/epi.12828

DeKraker, J., Ferko, K., Lau, J., Köhler, S., & Khan, A. (2018). Unfolding the hippocampus: An

intrinsic coordinate system for subfield segmentations and quantitative mapping.

NeuroImage, 167, 408418. 10.1016/j.neuroimage.2017.11.054

DeKraker, J., Haast, R. A. M., Yousif, M. D., Karat, B., Lau, J. C., Köhler, S., & Khan, A. R.

(2022). Automated hippocampal unfolding for morphometry and subfield segmentation

with Hippunfold. *ELife*, *11*. https://doi.org/10.7554/elife.77945

DeKraker, J., Köhler, S., & Khan, A. R. (2021). Surface-based hippocampal subfield

segmentation. *Trends in Neurosciences*. https://doi.org/10.1016/j.tins.2021.06.005

DeKraker, J., Lau, J. C., Ferko, K. M., Khan, A. R., & Köhler, S. (2020). Hippocampal

subfields revealed through unfolding and unsupervised clustering of laminar and

morphological features in 3D BigBrain. *NeuroImage*, *206*, 116328.

https://doi.org/10.1016/j.neuroimage.2019.116328

Dhikav, V., & Anand, K. S. (2012). Hippocampus in health and disease: An overview. *Annals of*

Indian Academy of Neurology, *15*(4), 239. https://doi.org/10.4103/0972-2327.104323

Ding, S.-L., & Van Hoesen, G. W. (2015). Organization and detailed parcellation of human

hippocampal head and body regions based on a combined analysis of cyto- and

chemoarchitecture. *Journal of Comparative Neurology*, *523*(15), 2233–2253.

https://doi.org/10.1002/cne.23786

Dinkelacker, V., Valabregue, R., Thivard, L., Lehéricy, S., Baulac, M., Samson, S., &

Dupont, S. (2015). Hippocampal-thalamic wiring in medial temporal lobe epilepsy:

Enhanced connectivity per hippocampal voxel. *Epilepsia*, *56*(8), 1217–1226.

https://doi.org/10.1111/epi.13051

Duvernoy, H. M., Cattin, F., & Risold, P.-Y. (2013). *The human hippocampus functional*

anatomy, vascularization and serial sections with MRI. Springer.

Fukutomi, H., Glasser, M. F., Murata, K., Akasaka, T., Fujimoto, K., Yamamoto, T., Autio, J.

A., Okada, T., Togashi, K., Zhang, H., Van Essen, D. C., & Hayashi, T. (2019).

Diffusion tensor model links to neurite orientation dispersion and density imaging at

high B-value in cerebral cortical gray matter. Scientific Reports, 9(1).

https://doi.org/10.1038/s41598-019-48671-7

Fukutomi, H., Glasser, M. F., Zhang, H., Autio, J. A., Coalson, T. S., Okada, T., Togashi,

K., Van Essen, D. C., & Hayashi, T. (2018). Neurite imaging reveals

microstructural variations in human cerebral cortical gray matter. *NeuroImage*, *182*,

488–499.10.1016/j.neuroimage.2018.02.017

Garcia, K. E., Kroenke, C. D., & Bayly, P. V. (2018). Mechanics of cortical folding: Stress,

growth and stability. *Philosophical Transactions of the Royal Society B: Biological*

Sciences, *373*(1759), 20170321. https://doi.org/10.1098/rstb.2017.0321

Garyfallidis, E., Brett, M., Amirbekian, B., Rokem, A., van der Walt, S., Descoteaux, M., &

Nimmo-Smith, I. (2014). Dipy, a library for the analysis of diffusion MRI data.

Frontiers in Neuroinformatics, *8*. https://doi.org/10.3389/fninf.2014.00008

Genon, S., Bernhardt, B. C., La Joie, R., Amunts, K., & Eickhoff, S. B. (2021). The many

dimensions of human hippocampal organization and (dys)function. *Trends in*

Neurosciences, *44*(12), 977–989. https://doi.org/10.1016/j.tins.2021.10.003

Glasser, M. F., & Van Essen, D. C. (2011). Mapping human cortical areas in vivo based on

myelin content as revealed by T1- and T2-weighted MRI. *Journal of Neuroscience*,

31(32), 11597–11616. https://doi.org/10.1523/jneurosci.2180-11.2011

Glasser, M. F., Sotiropoulos, S. N., Wilson, J. A., Coalson, T. S., Fischl, B., Andersson, J. L.,

Xu, J., Jbabdi, S., Webster, M., Polimeni, J. R., Van Essen, D. C., & Jenkinson, M.

(2013).

The minimal preprocessing pipelines for the Human Connectome Project. *NeuroImage*,

80, 105–124. https://doi.org/10.1016/j.neuroimage.2013.04.127

Glasser, M. F., Goyal, M. S., Preuss, T. M., Raichle, M. E., & Van Essen, D. C. (2014). Trends

and properties of human cerebral cortex: Correlations with cortical myelin content.

NeuroImage, *93*, 165–175. https://doi.org/10.1016/j.neuroimage.2013.03.060

Goodroe, S. C., Starnes, J., & Brown, T. I. (2018). The Complex Nature of

Hippocampal-Striatal Interactions in Spatial Navigation. *Frontiers in Human*

Neuroscience, 12. 10.3389/fnhum.2018.00250

Grussu, F., Schneider, T., Tur, C., Yates, R. L., Tachrount, M., Ianuş, A., Yiannakas, M. C.,

Newcombe, J., Zhang, H., Alexander, D. C., DeLuca, G. C., & Gandini Wheeler-

Kingshott, C. A. (2017). ? *Annals of Clinical and Translational Neurology*, *4*(9), 663–

679. https://doi.org/10.1002/acn3.445

Guerrero, J. M., Adluru, N., Bendlin, B. B., Goldsmith, H. H., Schaefer, S. M., Davidson, R. J.,

Kecskemeti, S. R., Zhang, H., & Alexander, A. L. (2019). Optimizing the fitting initial

condition for the parallel intrinsic diffusivity in NODDI: An extensive empirical

evaluation. https://doi.org/10.1101/630541

Halko, N., Martinsson, P. G., & Tropp, J. A. (2011). Finding structure with randomness:

Probabilistic algorithms for constructing approximate matrix decompositions. *SIAM*

Review, *53*(2), 217–288. https://doi.org/10.1137/090771806

Harms, R. L., Fritz, F. J., Tobisch, A., Goebel, R., & Roebroeck, A. (2017). Robust and fast

nonlinear optimization of diffusion mri microstructure models. *NeuroImage*, *155*, 82–96.

https://doi.org/10.1016/j.neuroimage.2017.04.064

Horner, A. J., Bisby, J. A., Bush, D., Lin, W., & Burgess, N. (2015). Evidence for holistic

episodic recollection via hippocampal pattern completion. *Nature Communications, 6*(1).

10.1038/ncomms8462

Isensee, F., Jaeger, P. F., Kohl, S. A., Petersen, J., & Maier-Hein, K. H. (2020). NNU-net: A

self-configuring method for deep learning-based biomedical image segmentation. *Nature*

Methods, *18*(2), 203–211. https://doi.org/10.1038/s41592-020-01008-z

Jelescu, I. O., Veraart, J., Adisetiyo, V., Milla, S. S., Novikov, D. S., & Fieremans, E. (2015).

One diffusion acquisition and different white matter models: How does microstructure

change in human early development based on WMTI and Noddi? *NeuroImage*, *107*,

242–256. https://doi.org/10.1016/j.neuroimage.2014.12.009

Jelescu, I. O., Veraart, J., Fieremans, E., & Novikov, D. S. (2016). Degeneracy in model

parameter estimation for multi-compartmental diffusion in neuronal tissue. *NMR in*

Biomedicine, *29*(1), 33–47. https://doi.org/10.1002/nbm.3450

Jelescu, I. O., de Skowronski, A., Geffroy, F., Palombo, M., & Novikov, D. S. (2022). Neurite

Exchange Imaging (NEXI): A minimal model of diffusion in gray matter with inter-

compartment water exchange. *NeuroImage*, *256*, 119277.

https://doi.org/10.1016/j.neuroimage.2022.119277

Jenkinson, M., Bannister, P., Brady, M., & Smith, S. (2002). Improved optimization for the

robust and accurate linear registration and motion correction of brain images.

NeuroImage, *17*(2), 825–841. https://doi.org/10.1006/nimg.2002.1132

Jespersen, S. N., Bjarkam, C. R., Nyengaard, J. R., Chakravarty, M. M., Hansen, B., Vosegaard,

T., Østergaard, L., Yablonskiy, D., Nielsen, N. C., & Vestergaard-Poulsen, P. (2010).

Neurite density from magnetic resonance diffusion measurements at ultrahigh field:

Comparison with light microscopy and electron microscopy. *NeuroImage*, *49*(1),

205–216. https://doi.org/10.1016/j.neuroimage.2009.08.053

Jürgen M. K., Paxinos, G., Insausti, R., & Amaral, D. G. (2011). Hippocampal Formation. In *The*

Human Nervous System (3rd ed., pp. 871–914). Elsevier Academic Press.

Karat., B (2023). Bradley-Karat/Hippo_Spin_Testing; 0.1.0 (v0.1.0). Zenodo.

https://doi.org/10.5281/zenodo.7837990

Kirkby, D. L., & Higgins, G. A. (2001). Characterization of perforant path lesions in rodent

models of memory and attention. *European Journal of Neuroscience*, *10*(3), 823–838.

https://doi.org/10.1046/j.1460-9568.1998.00087.x

Knierim, J. J. (2015). The hippocampus. *Current Biology, 25*(3), R1116-R1121.

La Joie, R., Perrotin, A., de La Sayette, V., Egret, S., Doeuvre, L., Belliard, S., Eustache, F.,

Desgranges, B., & Chételat, G. (2013). Hippocampal subfield volumetry in mild

cognitive impairment, Alzheimer's disease and semantic dementia. *NeuroImage:*

Clinical,

3, 155–162. https://doi.org/10.1016/j.nicl.2013.08.007

Liu, C., Ye, F. Q., Newman, J. D., Szczupak, D., Tian, X., Yen, C. C.-C., Majka, P., Glen, D.,

Rosa, M. G., Leopold, D. A., & Silva, A. C. (2020). A resource for the detailed 3D

mapping of white matter pathways in the marmoset brain. *Nature Neuroscience*, *23*(2),

271–280. https://doi.org/10.1038/s41593-019-0575-0

Maillot, J., Yahia, H., & Verroust, A. (1993). Interactive texture mapping. *Proceedings of the*

20th Annual Conference on Computer Graphics and Interactive Techniques -

SIGGRAPH '93. https://doi.org/10.1145/166117.166120

Moodley, K. K., & Chan, D. (2014). The hippocampus in neurodegenerative disease. *Frontiers*

of Neurology and Neuroscience, 95–108. https://doi.org/10.1159/000356430

Nieuwenhuys, R., Huijzen, C. van, & Voogd, J. (2008). *The human central nervous system*.

Springer.

Nieuwenhuys, R. (2013). The myeloarchitectonic studies on the human cerebral cortex of the

Vogt-vogt school, and their significance for the interpretation of functional neuroimaging

data. *Microstructural Parcellation of the Human Cerebral Cortex*, 55–125.

https://doi.org/10.1007/978-3-662-45766-5_3

Novikov, D. S., Veraart, J., Jelescu, I. O., & Fieremans, E. (2018). Rotationally-invariant

mapping of scalar and orientational metrics of neuronal microstructure with diffusion

MRI. *NeuroImage*, *174*, 518–538. https://doi.org/10.1016/j.neuroimage.2018.03.006

Ozarslan, E., Koay, C.G., Shepherd, T., Blackband, S., & Basser, P. (2009). Simple harmonic

oscillator based reconstruction and estimation for three-dimensional q-space MRI.

Proceedings of the International Society for Magnetic Resonance in Medicine, 16.

Palombo, M., Ianus, A., Guerreri, M., Nunes, D., Alexander, D. C., Shemesh, N., & Zhang, H.

(2020). SANDI: A compartment-based model for non-invasive apparent Soma and neurite

imaging by Diffusion MRI. *NeuroImage*, *215*, 116835.

https://doi.org/10.1016/j.neuroimage.2020.116835

Patel, R., Steele, C. J., Chen, A. G. X., Patel, S., Devenyi, G. A., Germann, J., Tardif, C. L., &

Chakravarty, M. M. (2020). Investigating microstructural variation in the human

hippocampus using non-negative matrix factorization. *NeuroImage*, *207*, 116348.

https://doi.org/10.1016/j.neuroimage.2019.116348

Pedregosa, F., Varoquaux, G., Gramfort, A., Michel, V., Thirion, B., Grisel, O., Blondel, M.,

Prettenhofer, P., Weiss, R., Dubourg, V., Vanderplas, J., Passos, A., Cournapeau, D.,

Brucher, M., Perrot, M., & Duchesnay, É. (2011). Scikit-learn: Machine Learning in

Python. *Journal of Machine Learning Research*, *12*(85), 2825–2830.

Pierpaoli, C., Jezzard, P., Basser, P. J., Barnett, A., & Di Chiro, G. (1996). Diffusion tensor MR

imaging of the human brain. *Radiology*, *201*(3), 637–648.

https://doi.org/10.1148/radiology.201.3.8939209

Plachti, A., Eickhoff, S. B., Hoffstaedter, F., Patil, K. R., Laird, A. R., Fox, P. T., Amunts, K., &

Genon, S. (2019). Multimodal parcellations and extensive behavioral profiling tackling

the hippocampus gradient. *Cerebral Cortex*, *29*(11), 4595–4612.

https://doi.org/10.1093/cercor/bhy336

Plachti, A., Kharabian, S., Eickhoff, S. B., Maleki Balajoo, S., Hoffstaedter, F., Varikuti, D. P.,

Jockwitz, C., Caspers, S., Amunts, K., & Genon, S. (2020). Hippocampus co-atrophy

pattern in dementia deviates from covariance patterns across the lifespan. *Brain*, *143*(9),

2788–2802. https://doi.org/10.1093/brain/awaa222

Poppenk, J., Evensmoen, H. R., Moscovitch, M., & Nadel, L. (2013). Long-axis specialization

of the human hippocampus. *Trends in Cognitive Sciences*, *17*(5), 230–240.

https://doi.org/10.1016/j.tics.2013.03.005

Qian, W., Khattar, N., Cortina, L. E., Spencer, R. G., & Bouhrara, M. (2020). Nonlinear

associations of neurite density and myelin content with age revealed using

multicomponent diffusion and relaxometry magnetic resonance imaging. *NeuroImage*,

223, 117369. https://doi.org/10.1016/j.neuroimage.2020.117369

Quezada, S., van de Looij, Y., Hale, N., Rana, S., Sizonenko, S. V., Gilchrist, C.,

Castillo-Melendez, M., Tolcos, M., & Walker, D. W. (2020). Genetic and

microstructural differences in the cortical plate of gyri and sulci during gyrification in

fetal sheep. *Cerebral Cortex*, *30*(12), 6169–6190. https://doi.org/10.1093/cercor/bhaa171

Radhakrishnan, H., Stark, S. M., & Stark, C. E. (2020). Microstructural alterations in

Hippocampal Subfields Mediate Age-related Memory decline in humans. *Frontiers in*

Aging Neuroscience, *12*. https://doi.org/10.3389/fnagi.2020.00094

Raffelt, D. A., Tournier, J.-D., Smith, R. E., Vaughan, D. N., Jackson, G., Ridgway, G. R., &

Connelly, A. (2017). Investigating white matter fibre density and morphology using

fixel-based analysis. *NeuroImage*, *144*, 58–73.

https://doi.org/10.1016/j.neuroimage.2016.09.029

Robinson, J. L., Barron, D. S., Kirby, L. A., Bottenhorn, K. L., Hill, A. C., Murphy, J. E., Katz,

J.S., Salibi, N., Eickhoff, S. B., & Fox, P. T. (2015). Neurofunctional topography of the

human hippocampus. *Human Brain Mapping*, *36*(12), 5018–5037.

https://doi.org/10.1002/hbm.22987

Salmenpera, T. M., Simister, R. J., Bartlett, P., Symms, M. R., Boulby, P. A., Free, S. L.,

Barker, G. J., & Duncan, J. S. (2006). High-resolution diffusion tensor imaging of the

hippocampus in temporal lobe epilepsy. *Epilepsy Research*, *71*(2-3), 102–106.

https://doi.org/10.1016/j.eplepsyres.2006.05.020

Schmierer, K., Wheeler-Kingshott, C. A. M., Boulby, P. A., Scaravilli, F., Altmann, D. R.,

Barker, G. J., Tofts, P. S., & Miller, D. H. (2007). Diffusion tensor imaging of Post

Mortem Multiple Sclerosis Brain. *NeuroImage*, *35*(2), 467–477.

https://doi.org/10.1016/j.neuroimage.2006.12.010

Shepherd, T. M., Özarslan, E., Yachnis, A. T., King, M. A., & Blackband, S. J. (2007).

Diffusion Tensor Microscopy Indicates the Cytoarchitectural Basis for Diffusion

Anisotropy in the Human Hippocampus. *American Journal of Neuroradiology*, *28*(5),

958–964.

Small, S. A., Schobel, S. A., Buxton, R. B., Witter, M. P., & Barnes, C. A. (2011). A

pathophysiological framework of hippocampal dysfunction in ageing and disease. *Nature*

Reviews Neuroscience, *12*(10), 585–601. https://doi.org/10.1038/nrn3085

Sotiras, A., Resnick, S. M., & Davatzikos, C. (2015). Finding imaging patterns of structural

covariance via non-negative matrix factorization. *NeuroImage*, *108*, 1–16.

https://doi.org/10.1016/j.neuroimage.2014.11.045

Sotiropoulos, S. N., Moeller, S., Jbabdi, S., Xu, J., Andersson, J. L., Auerbach, E. J., Yacoub,

E., Feinberg, D., Setsompop, K., Wald, L. L., Behrens, T. E., Ugurbil, K., & Lenglet, C.

(2013). Effects of image reconstruction on fiber orientation mapping from multichannel

diffusion mri: Reducing the noise floor using sense. *Magnetic Resonance in Medicine*,

70(6), 1682–1689. https://doi.org/10.1002/mrm.24623

Stolp, H. B., Ball, G., So, P.-W., Tournier, J.-D., Jones, M., Thornton, C., & Edwards, A. D.]

(2018). Voxel-wise comparisons of cellular microstructure and diffusion-MRI in mouse

hippocampus using 3D bridging of optically-clear histology with neuroimaging data (3D-

bond). *Scientific Reports*, *8*(1). https://doi.org/10.1038/s41598-018-22295-9

Strang, B. A., Witter, M. P., Lein, E. S., & Moser, E. I. (2014). Functional organization of the

Hippocampal Longitudinal Axis. *Nature Reviews Neuroscience*, *15*(10), 655–669.

https://doi.org/10.1038/nrn3785

Swanson, L. W., Wyss, J. M., & Cowan, W. M. (1978). An autoradiographic study of the

organization of intrahippocampal association pathways in the rat. *The Journal of*

Comparative Neurology, *181*(4), 681–715. https://doi.org/10.1002/cne.901810402

Szirmai, I., Buzsáki, G., & Kamondi, A. (2012). 120 years of hippocampal Schaffer Collaterals.

Hippocampus, *22*(7), 1508–1516. https://doi.org/10.1002/hipo.22001

Tardif, C. L., Gauthier, C. J., Steele, C. J., Bazin, P.-L., Schäfer, A., Schaefer, A., Turner, R., &

Villringer, A. (2016). Advanced MRI techniques to improve our understanding of

experience-induced neuroplasticity. *NeuroImage*, *131*, 55–72.

https://doi.org/10.1016/j.neuroimage.2015.08.047

Van Essen, D., Smith, S., Barch, D., Behrens, T., Yacoub, E., & Ugurbil, K. (2013). The

WU-Minn Human Connectome Project: An overview. *NeuroImage, 80*(62), 62-79.

van Uden, I. W. M., Tuladhar, A. M., van der Holst, H. M., van Leijsen, E. M. C., van

    Norden, A. G. W., de Laat, K. F., Rutten-Jacobs, L. C. A., Norris, D. G., Claassen, J. A.

    H. R., van Dijk, E. J., Kessels, R. P. C., & de Leeuw, F.-E. (2015). Diffusion tensor

    imaging of the HIPPOCAMPUS predicts the risk of dementia; the RUN DMC study.

    Human Brain Mapping, *37*(1), 327–337. https://doi.org/10.1002/hbm.23029

Voss, J. L., Bridge, D. J., Cohen, N. J., & Walker, J. A. (2017). A Closer Look at the

    Hippocampus and Memory. *Trends in Cognitive Sciences, 21*(8), 577-588.

    10.1016/j.tics.2017.05.008

Yang, & Oja, E. (2010). Linear and nonlinear projective nonnegative matrix factorization. *IEEE*

    Transactions on Neural Networks, *21*(5), 734–749.

    https://doi.org/10.1109/tnn.2010.2041361

Yassa, M. A., Muftuler, L. T., & Stark, C. E. (2010). IC-P-056: Microstructural diffusion tensor

    Imaging reveals perforant path degradation in aged humans in vivo. *Alzheimer's &*

    Dementia, 6.

Yushkevich, P. A., Wang, H., Pluta, J., Das, S. R., Craige, C., Avants, B. B., Weiner, M. W., &

    Mueller, S. (2011). Nearly automatic segmentation of hippocampal subfields in in vivo

    focal T2-weighted MRI. *NeuroImage*, *53*(4), 1208–1224.

    https://doi.org/10.1016/j.neuroimage.2010.06.040

Zeineh, M. M., Holdsworth, S., Skare, S., Atlas, S. W., & Bammer, R. (2012). Ultra-high

    resolution diffusion tensor imaging of the microscopic pathways of the medial temporal

    lobe. *NeuroImage, 62*(3), 2065-2082.

Zeineh, M. M., Palomero-Gallagher, N., Axer, M., Gräßel, D., Goubran, M., Wree, A., Woods,

    R., Amunts, K., & Zilles, K. (2017). Direct visualization and mapping of the spatial course

    of fiber tracts at microscopic resolution in the Human Hippocampus. *Cerebral Cortex*.

    https://doi.org/10.1093/cercor/bhw010

Zhang, H., Schneider, T., Wheeler-Kingshott, C. A., & Alexander, D. C. (2012). NODDI:

Practical in vivo neurite orientation dispersion and density imaging of the human brain.

NeuroImage, 61(4), 1000-1016. 10.1016/j.neuroimage.2012.03.072

Zhong, Q., Xu, H., Qin, J., Zeng, L.-L., Hu, D., & Shen, H. (2019). Functional parcellation of the

hippocampus from resting-state dynamic functional connectivity. *Brain Research*, *1715*,

165–175. https://doi.org/10.1016/j.brainres.2019.03.023

Zhou, Y., Dougherty, J. H., Hubner, K. F., Bai, B., Cannon, R. L., & Hutson, R. K. (2008).

Abnormal connectivity in the posterior cingulate and hippocampus in early alzheimer's

disease and mild cognitive impairment. *Alzheimer's & Dementia*, *4*(4), 265–270.

https://doi.org/10.1016/j.jalz.2008.04.006