Sex-biased infections scale to population impacts for an emerging wildlife disease

Macy J. Kailing[1], Joseph R. Hoyt[1], J. Paul White[2], Heather M. Kaarakka[2], Jennifer A. Redell[2],

Ariel E. Leon[3], Tonie E. Rocke[3], John E. DePue[4], William H. Scullon[5], Katy L. Parise[6], Jeffrey

T. Foster[6], A. Marm Kilpatrick[7], and Kate E. Langwig[1]

1 Department of Biological Sciences, Virginia Polytechnic Institute, Blacksburg, VA, USA

2 Wisconsin Department of Natural Resources, Madison, WI, USA

3 US Geological Survey, National Wildlife Health Center, Madison, WI, USA

4 Michigan Department of Natural Resources, Baraga, MI, USA

5 Michigan Department of Natural Resources, Norway, MI, USA

6 Pathogen and Microbiome Institute, Northern Arizona University, Flagstaff, AZ, USA

7 Department of Ecology and Evolutionary Biology, University of California, Santa Cruz, CA,

USA

* Corresponding author: Macy Kailing, macykailing@gmail.com

Keywords: Sex biased infection, emerging infectious disease, white-nose syndrome, bats, Allee

effects, fungal disease

ABSTRACT:

Demographic factors are fundamental in shaping infectious disease dynamics. Aspects of populations that create structure, like age and sex, can affect patterns of transmission, infection intensity and population outcomes. However, studies rarely link these processes from individual to population-scale effects. Moreover, the mechanisms underlying demographic differences in disease are frequently unclear. Here, we explore sex-biased infections for a multi-host fungal disease of bats, white-nose syndrome, and link disease-associated mortality between sexes, the distortion of sex ratios, and the potential mechanisms underlying sex differences in infection. We collected data on host traits, infection intensity, and survival of five bat species at 42 sites across seven years. We found females were more infected than males for all five species. Females also had lower apparent survival over winter and accounted for a smaller proportion of populations over time. Notably, female-biased infections were evident by early hibernation and likely driven by sex-based differences in autumn mating behavior. Male bats were more active during autumn which likely reduced replication of the cool-growing fungus. Higher disease impacts in female bats may have cascading effects on bat populations beyond the hibernation season by limiting recruitment and increasing the risk of Allee effects.

1. INTRODUCTION

Emerging infectious diseases are a serious threat to wildlife health (1, 2). Population structure can shape epidemics by influencing spatial spread, outbreak size, and host impacts (2-4). Elements of populations that create structure such as classes of individuals of specific ages, sexes, or breeding stages, can have profound effects on disease dynamics (3, 5-9). Sex is an especially important factor because sex-biases in infections can contribute to differential transmission through populations due to behavior (10-15), amplify outbreaks due to seasonal changes in susceptibility (16-19), and modify population impacts through disproportionate mortality (20-22). Differences in infection and mortality can also modulate virulence evolution through sex-specific immune responses that affect pathogen replication and growth (23, 24). As such, determining patterns and mechanisms of sex-biases in infections will improve efforts to minimize outbreaks and manage impacts.

Behavioral and physiological traits are two mechanisms that can produce sex-biases in transmission, susceptibility, infection intensity, or disease-induced mortality (25-27). In most systems, males have an elevated risk of disease compared to females due to territory defense and

promiscuous mating, which increases contacts with conspecifics or increase pathogen

susceptibility due to physiological stress (9, 25, 28). Sex hormones can also have strong effects

on host immunity (29), such that testosterone suppresses immune responses while estrogens

enhance it, resulting in weaker immune responses in males, which can increase susceptibility to

pathogen infection (25, 30, 31). As such, the majority of empirical studies find that infections are

typically male-biased (28, 32, 33), although this generalization is sometimes reversed (34-40) or

weak (41) and may be linked to seasonal reproductive stress associated with pregnancy,

parturition, or parental care (16, 17). Thus, it is likely that the effects of sex-biased infections

may be highly pronounced in disease systems where host reproductive strategies differ

seasonally among sexes (19).

White-nose syndrome (WNS) is a highly seasonal fungal disease of bats that has caused

population collapse, with declines exceeding 95% in many populations of multiple species (42-

46). WNS is caused by the fungal pathogen, *Pseudogymnoascus destructans,* which invades the

epidermal tissue of bats during hibernation, and disrupts bat homeostasis, causing water and

electrolyte imbalances that increase arousals and deplete stored fat (47-54). Transmission of *P.*

destructans occurs through host-to-host contacts and contact between hosts and contaminated

environmental reservoirs inside winter hibernation sites (55-59). *Pseudogymnoascus destructans*

has a thermal growth range of 0-20ºC and optimal growth between 12-16 ºC (60). This limits on-

host growth to the periods when bats are in torpor and they lower their body temperature to the

ambient temperatures of their hibernation sites (60-62). The restricted growth of *P. destructans*

above 20ºC drives the seasonal patterns of WNS, with infection only occurring during winter

hibernation and mortality peaking during late winter (56, 59, 63).

Temperate bat species have seasonal sex-biased differences in behaviors that may influence

their exposure, susceptibility, and mortality from WNS (64-69). Bats mate in autumn swarms,

and females store sperm over winter, delaying ovulation, and giving birth at maternity colonies

in spring (70). Male reproductive energy expenditures are highest during autumn when they

aggregate at hibernacula (subterranean sites where bats spend the winter) and mate

indiscriminately with females (71). Since autumn swarm coincides with the seasonal

transmission of *P. destructans* (63), breeding stress among males as well as exposure to the

environmental reservoirs (56, 58) could increase their susceptibility to infection. Female energy

expenditures are greatest during pregnancy and lactation in spring and summer when hosts

typically clear infection (72, 73) and transmission of *P. destructans* is low (63), suggesting

females may be the less infected sex. However, female bats may spend more time torpid during

winter than males to conserve energy for spring reproduction (66) which could influence the

growth of *P. destructans* (60). Due to the differences in reproductive investment and seasonal

energy expenditure, sex-biased traits have the potential to affect exposure (e.g., contacting *P.*

destructans from other hosts or in the environment) and pathogen growth (e.g., time spent at

torpid temperatures that permit fungal growth). Thus, we predict sex-specific behaviors, and the

highly seasonal pattern of WNS may drive intersex differences in infection, mortality, and

population impacts. Here, we examined differences in *P. destructans* infection between females

and males across five bat species in 42 sites across seven years. We then assessed the

consequences of intersex differences in infection at both the individual and population-level.

Lastly, we explored autumn activity patterns between sexes as a potential mechanism of sex-

biased infections.

2. MATERIALS AND METHODS

a. Study sites and sampling design

We sampled bats at hibernacula in Illinois, Massachusetts, Michigan, New York, Vermont,

Virginia, and Wisconsin between 2011 and 2021 twice during a seasonal epizootic. Generally,

bats were sampled at two time points: early winter (between November and December) and

again in late winter (between March and April). During site visits, we walked each section of the

hibernacula and counted all the bats of each species and resighted any bats that were previously

banded. We sampled up to 20 bats of each species during each visit when possible and stratified

sample collection throughout hibernacula to obtain a sample reflective of the spatial distribution

of bats at each site. Bat species included little brown (*Myotis lucifugus*), northern long-eared

(*Myotis septentrionalis*), tricolored (*Perimyotis subflavus*), big brown (*Eptesicus fuscus*), and

Indiana (*Myotis sodalis*) bats. The animal sampling protocols were approved by the Virginia

Tech IACUC protocol 17-180. We followed the field decontamination procedures outlined by

the United States Fish and Wildlife Service Decontamination Guidelines as well as the

recommendations provided by individual states.

b. *P. destructans* sample collection and quantification

For every bat we sampled, we determined species and biological sex based on external

morphology and attached an aluminum-lipped band (2.4, 2.9, and 4.2mm; Porzana Ltd.,

Icklesham, E. Sussex, U.K.) to the wing for individual identification and resighting. We

collected an epidermal swab from individuals to quantify infection prevalence (the fraction of

individuals positive for *P. destructans*) and severity (the quantity of *P. destructans* on infected

bats) using a standardized swabbing technique (45, 63). We placed swabs in RNAlater® for

storage before testing. We extracted DNA and tested for *P. destructans* presence and quantity by

qPCR using protocols developed specifically for this fungal pathogen that included a standard

growth curve (63, 74).

133   **c.   Activity data collection**

At three sites in Wisconsin in 2020, we installed radio frequency identification (RFID)

systems, consisting of a passive antenna and an automated data logger (IS10001; Biomark,

Boise, ID) at the entrances of each hibernaculum (2-4 per site). The systems at each entrance

were equipped with a solar or direct power source to run continuously. During autumn swarm,

we captured little brown bats swarming near hibernacula and injected 12.5mm PIT tags

(Biomark APT12; Biomark, Boise, ID). Each RFID system was programmed to record each time

a unique individual passed through an entrance with a one-minute delay to avoid duplicate

detections of tags. We scored a detection as an active bat and used it to characterize activity of

females and males.

143   **d.   Statistical analyses**

144   ***i. Infection***

We used generalized (GLMM) and linear (LMM) mixed effect models (75) to compare

differences in *P. destructans* prevalence and infection severity between males and females of

each of the five species. We defined prevalence as the fraction of bats testing positive for *P.*

destructans on qPCR (0|1), and infection severity as the $\log_{10}$-transformed quantity of fungal

DNA (ng) on positive bats ($\log_{10}$ fungal loads) which is a well-established metric of infection

severity (43, 76, 77). We estimated differences in prevalence between sexes using a GLMM with

a binomial distribution and sex and species as fixed effects (Model 1, Appendix 1.1) and

compared additive and interactive versions of the model using Akaike Information Criterion

(AIC; Appendix 1.2). For prevalence models, we only used data from early hibernation as every

bat was infected at the end of hibernation.

We used two datasets to examine changes in fungal loads over winter (between early and late

hibernation) between sexes. First, to estimate population-level differences in fungal loads, we

used a LMM with fixed additive effects of species and sex and date as a fixed effect interacting

with each term (Model 2, Appendix 2.1). Fungal loads are well-correlated with mortality (43, 76,

78), and bats that begin hibernation with higher fungal loads are more likely to die of their

infections before the end of winter (79), potentially resulting in survival biases in population-

level analyses (43) .Therefore, in a second analysis, we also explored whether growth rates of *P.*

destructans on individual female and male bats that were recaptured over winter clearly differed

(43). For example, if one sex had higher fungal loads than the other and subsequently died of

their infections, these highly infected bats will be omitted from the late hibernation population-

level sample (77). By comparing the difference between late and early hibernation fungal loads

on individual recaptured bats, we can directly assess whether female and male bats have different

growth rates and better identify the timing of transmission differences that may underlie sex-

biases in infection. Therefore, we compared the over winter increase in fungal loads on

recaptured individuals (the $\log_{10}$ difference between late hibernation and early hibernation loads)

using a LMM with sex as a fixed effect and site as a random effect (Model S1A, Appendix

5.1.1). Lastly, to explore differences in sex-based ecology that could lead to differences in

infection between sexes, we also examined models including *a priori* hypothesized effects of

roosting temperature and body mass on early hibernation prevalence and fungal loads but found

no clear improvements over models with sex and species using AIC (Appendix 5.2 and 5.3). We

used site as a random effect for all infection analyses and confirmed the performance of our

infection models using 5-fold cross validation (see Supplemental Material for description).

177   *ii.   Individual survival*

We examined differences between sexes in over winter survival using data from individual

little brown bats, which were abundant enough to obtain reasonable sample sizes. We compared

the probability of observing bats in late hibernation (March) that we observed in early

hibernation (November). Bats affected by WNS often emerge prematurely (mid-winter) and die

on the landscape (80), enabling the use of recapture as an estimate of apparent survival between

sampling dates (79, 81). We used a mixed effects model with site as a random effect with a

binomial distribution and a logit link to quantify how sex affected the probability of a bat being

resighted over winter (Model 3A; Appendix 3.1).

186   *iii.*   *Population-level impact of disease*

We estimated the proportion of female little brown bats sampled at each visit from a core set

of sites (N=14) that were sampled in consecutive years to evaluate how female proportions were

changing over time. We estimated the probability of sampling a female vs a male (1|0) over time

using generalized linear mixed model with a binomial distribution and a logit link with

continuous years since *P. destructans* invasion as a fixed effect, and site as a random effect

(Model 3B; Appendix 3.2).

193   *iv.*   **RFID activity**

To explore differences in autumn activity of female and male bats, we used a generalized

linear mixed model with a binomial distribution and a logit link with sex and site as fixed effects

(we did not include site as a random effect due to the small number of sites) and individual bat

identification as a random effect (Model 4; Appendix 4.1). For our response variable, we treated

each bat as a series of binomial trials where it could be detected (=1) or not detected (=0) on a

given night throughout autumn swarm that any RFID system was operational at an entrance, as

determined by the detection of a programmed test tag. We determined the period of autumn

swarm to be between the dates in which more than 95% of all individual detections occurred

(beginning of swarm; August 19) and all individual detections concluded (end of swarm;

October 01). Results were qualitatively similar if we included all dates in which any bat was

detected from early August to November. Since bats are nocturnal and daily activity patterns

span two calendar days, we treated detections that occurred between hours 0:00-07:00 as a

detection during the previous night for consistency with bat ecology. We used 5-fold cross-

validation to check the performance of our model (see Supplemental Material for description).

3. RESULTS

We examined the effects of sex on *P. destructans* infections of five bat species during winter

(Supplemental Table 1). We sampled a total of 665 females and 1071 males across all species,

sites, years, and seasons (Supplemental Table 2A-D). On average, females had higher *P.*

destructans prevalence than male bats in early winter (GLMM with site and species as random

effects; female coef +/- SE: 0.630 +/- 0.221, P = 0.0043) and suffered from higher fungal loads

(LMM with site and species as random effects; female coef +/- SE: 0.487 +/- 0.086, P < 0.0001).

The model predicting early winter *P. destructans* prevalence, including species and an additive

effect of sex, was better supported over the interactive version (ΔAIC = -3.11; Appendix 1.2-

1.3), suggesting female infections are generally higher across all bat species in our study

(GLMM with site as random effect; female coef +/- SE: 0.637 +/- 0.223, P = 0.0044; Figure 1;

Appendix 1.1).

The differences in infection prevalence during early winter aligned with higher fungal loads

on female bats during the hibernation season. Across species, females began hibernation with

higher fungal loads than males (LMM with site as random effect; female coef +/- SE: 0.490 +/-

0.093, P < 0.0001; Figure 2; Appendix 2.1). Fungal loads appeared to become more similar

between females and males by the end of winter (smaller date slope for females (=0.405) vs

males (=0.501); Figure 2; Appendix 2.4). However, this may be an artefact of higher female

mortality (Figure 3; Appendix 3.1) such that individuals with higher fungal loads in early

hibernation are less likely to survive to late winter, resulting in only bats with low early

hibernation infections surviving to be sampled in late hibernation (77, 79). To assess whether

mortality bias might be responsible for the apparent similarity of infections between sexes in late

hibernation, we used data on differences in fungal loads of recaptured individual little brown bats

sampled in both early and late winter. We found that the change in fungal loads over winter did

not differ between sexes in the absence of selective mortality (LMM with site as a random effect;

female coef +/- SE: 0.090 +/- 0.118, P = 0.448; Supplemental Figure1A; Appendix 5.1.1).

We did not find support that hibernation temperatures or body mass significantly improved

models of prevalence and fungal loads that already included sex (ΔAIC < 2; Appendix 5.2 and

5.3) and hibernation temperatures of male and female bats broadly overlapped (Supplemental

Figure 2; Appendix 5.2).

Figure 1. Early winter *P. destructans* prevalence by sex for five bat species. The colored points

show the proportion of individuals that were infected at a given site and year. The size of colored

points is scaled to the number of bats sampled during the site visit. The black circles show model

predicted prevalence with standard error bars. Sample sizes are provided in Supplemental Table

2A.

Figure 2. (a-e) Change in *P. destructans* infection severity (fungal loads) throughout winter by sex for each of the five host species. A point represents an individual bat, and females and males are shown in blue and orange, respectively. Lines represent model predicted fungal loads over time. Sample sizes by species are provided in Supplemental Table 2A. **(f)** Fungal loads in early winter by sex for each host species. Larger circles outlined in black show estimated fungal loads on November 1 extracted from model predictions used in a-e to visualize the differences in early winter infections across species.

At an individual scale, using data from little brown bats that were banded in early hibernation

(Supplemental Table 2B), we found clear support that female little brown bats were less likely to

be recaptured in late winter than males (GLMM with site as random effect; female coef +/- SE: -

0.855 +/- 0.380, P = 0.0240; Figure 3A; Appendix 3.1). At a population level, the proportion of

females sampled in the same populations at the end of winter (Supplemental Table 2C) decreased

with time since WNS invasion (GLMM with site as random effect; years since invasion coef +/-

SE: -0.363 +/- 0.173, P = 0.0365; Figure 3B; Appendix 3.2). Collectively, these data suggest that

females have higher, more severe infections (Figure 1, 2), and correspondingly experience higher

mortality during winter (Figure 3).

Females begin hibernation with more severe infections than male bats and differential

activity between sexes may influence infections. We found that the probability of a female being

active on a given night was four-fold lower than males during autumn. (GLMM with individual

bat as random effect; female coef +/- SE: -1.401 +/- 0.171, P < 0.0001; Figure 4; Supplemental

Table 2D; Appendix 4.1).

Figure 3. **(A)** Recapture probability of little brown bats by sex. Blue and orange points show the

observed probability of recapture from early (November) and late (March) winter from a given

site and year for females and males, respectively. Black points and vertical bars represent the

estimated mean probability of recapture and standard errors. Sample sizes are provided in

Supplemental Table 2B. **(B)** Proportional change in females with time since *P. destructans*

invasion. Points show the proportion of all sampled individuals that were female at the same sites

sampled over time since the invasion of *P. destructans*. The solid black line shows the model

predicted proportion of females and dashed lines show the standard error around the model

estimates. The size of points in both panels is scaled to the number of bats sampled during the

site visit. Sample sizes are provided in Supplemental Table 2C.

Figure 4. Differences in autumn activity of female and male little brown bats at three

hibernacula. Each data point represents the proportion of nights an individual bat was active at a

site. The proportion of nights active was calculated as the number of nights an individual was

detected at least once on the RFID systems divided by the total number of nights the RFID

systems at each site were operating throughout autumn swarm. Points at 0 represent tagged bats

that were never detected on a reader. Black points show model predicted activity by sex with

standard errors denoted with vertical bars. Sample sizes are provided in Supplemental Table 2D.

4. DISCUSSION

Our results demonstrate that sex differences in infection have an important role in disease

dynamics. Across all species, females experienced higher infection prevalence (Figure 1) and

fungal loads by the beginning of winter hibernation (Figure 2). In little brown bats, females were

also less likely to be recaptured over winter (Figure 3A), as we would expect if more severe

infections resulted in higher mortality among females (43, 56, 76, 77, 79). Finally, female-biased

mortality associated with time since disease arrival likely contributed to population-level

reductions in females over time (Figure 3B). Higher impacts to female bats may mask even more

severe long-term impacts from WNS (46, 82) as a disproportionate loss of recruiting females

may be expected to have more negative population-level consequences than male-biased

declines.

We consistently observed higher pathogen prevalence and intensity in females than males

across bat species (Figure 1, Figure 2). In other vertebrates, males maximize fitness at the cost of

greater exposure and susceptibility to pathogens (33, 83-85), in part through higher testosterone

concentrations which often correlate negatively with immune defense (86-88). Given male bats

undergo their most substantial investment in reproduction during autumn swarm compared to

females that invest relatively little towards reproduction throughout autumn and over winter (70),

we expected infection to be higher in males. However, we found the opposite pattern with

females having more severe disease. Drivers of higher female infections likely include sex-based

physiological differences (25). Temperate female bats use fat stores acquired in autumn more

slowly than males suggesting a more substantial constraint on energy expenditure compared to

males throughout the autumn to spring period (66, 67). Therefore, different physiological

strategies (e.g., torpor patterns) between female and male bats could be shaping sex-biased

infections.

Differences in sex-based physiology and behavior were supported by our data showing that

the proportion of nights males were active during autumn was substantially higher than females

(Figure 4). Differences in autumn activity patterns are likely motivated by differences in

reproductive energy allocation between sexes. For males, their fitness is enhanced by remaining

active so they can maximize mating opportunities with females. However, female bats, which

store sperm and delay ovulation until spring, may prefer to conserve energetic resources during

fall and use torpor more extensively (67). As bats arrive to contaminated hibernacula (56, 58) for

autumn swarm and are exposed to the pathogen, torpor use in autumn may permit pathogen

growth as bats reduce their body temperatures to be within the thermal range for *P. destructans*

replication. The relationship between sex-based activity and fungal intensity may arise through

two potential pathways associated with torpor expression. First, less active females may provide

favorable conditions for pathogen growth for a longer period prior to hibernation compared to

males, thus resulting in the more severe female infections. Second, more active males may be

able to actively inhibit pathogen growth through euthermia compared to females, as euthermic

mammals mount more robust immune defenses than torpid or hibernating mammals (89, 90).

Our activity estimates, which are consistent with other studies (65), suggest that vast differences

between male and female bats' energy use strategies during the mating period likely contribute to

differences in infection.

We were less likely to recapture female little brown bats over winter than males, suggesting

greater mortality in females (Figure 3A). Further, the sex ratio shifted to be more male-biased as

WNS progressed (Figure 3B), suggesting that fungal infection may be driving female-specific

mortality. Generally, infections increase on bats over winter before reaching a threshold at which

bats die from their infections (43), and this bias in survival of the least infected individuals in

early hibernation (Supplemental Figure 1B; Appendix 5.1.3) could explain why male and female

infections become more similar at the end of hibernation (Figure 2). On average, female bats

began hibernation with higher fungal loads than males (Figure 1) but have similar fungal growth

rates (Supplemental Figure 1A; Appendix 5.1) suggesting that females reach high infection

levels that result in mid-winter mortality, thus, are removed from the sample population by late

winter. This relationship between high early hibernation fungal loads and mortality is further

supported by previous studies that directly link early hibernation fungal loads with mortality and

impacts (43, 79). Our findings are supported by previous work demonstrating that WNS severity

is positively correlated with mortality, which is mechanistically linked by more frequent arousals

(43, 47, 53, 77, 81). Several lines of evidence also suggest that females did not leave hibernation

sites early and survive elsewhere. In our study region, emergence from hibernation in healthy

populations does not begin until four to six weeks after our sampling (91). Given that the

proportion of females declined with WNS progression (Figure 3B), our results suggest that

disease associated mortality may be contributing to the reduction in overwinter recaptures of

females. Previous studies on intersex differences in survival associated with WNS have found

contrasting results. One study observed lower female survival in naturally infected bats (68)

whereas another reported higher female survival in experimental infections where each sex was

inoculated with the same dose of *P. destructans* at identical times (92). Our results suggest that

differences in disease outcomes between females and males may originate from differences in

pathogen growth during autumn. Therefore, inoculating both sexes with the same dose

simultaneously in the experimental infection may have eliminated the underlying difference in

early winter infections between female and males.

We provide the first spatiotemporally broad evidence from the field that sex-based survival

of bats affected by WNS likely scales to population-level impacts. In the first year after *P.*

destructans invasion, the mean proportion of females in late hibernation was 45%, but this

percentage declined to 21% after four years of WNS (Figure 3B). The reduction in the percent of

females with time since pathogen invasion differs from previous studies of healthy bat

populations that show relatively consistent interannual proportions of females (93-96). This

suggests that a comparable decline in females is not typical in non-diseased populations. Losing

females at higher rates than males will likely affect how bat populations respond to WNS. First,

reducing the number of females could limit the recruitment potential of bat populations and is

likely to be especially critical for imperiled populations that remain small and vulnerable to Allee

effects. Many temperate bat species, including little brown bats, form maternity colonies during

summer when they cooperatively rear offspring and larger colony sizes afford females lower

energy expenditure (97). If birth and offspring survival rates decline with density, as shown in

other temperate bat species (98), the disproportionate loss of females could further impact

recruitment. Second, distorted sex-ratios may negatively affect reproductive success during

autumn. In other taxa, a significant increase in male to female sex-ratios resulted in increased

stress to females from exacerbated pressure for males to mate with limited females, negatively

affecting female fecundity (99, 100).

Our results include impacts to female bats up to four years after the arrival of WNS, which

includes the epidemic phase of the disease, when the majority of mortality occurs (46). Females

are likely under strong selection pressure to evolve mechanisms of survival given their increased

mortality and will need to adapt for populations to rebound (101). Future work focusing on the

effects of female infection and mortality biases on bat population persistence and recovery could

benefit conservation efforts, especially as the negative effects are likely to compound over time

if sex ratios become increasingly distorted.

We find a novel example in which female-biased infections may contribute to population-

level impacts of an emerging disease. Our results provide a clear, yet less frequently observed,

instance of an emerging pathogen that consistently impacts females more than males regardless

of host species. We describe a new mechanistic explanation to female-biased infections that links

temperature-dependent fungal growth to sex-specific seasonal physiology. Ultimately, disparate

infections among demographic classes of hosts are fundamental for understanding and managing

emerging infectious diseases, and cross-scale analyses can provide insights into the important

consequences of demographic biases on disease systems.

ACKNOWLEDGEMENTS

We thank Steffany Yamada for data curation support, Skylar Hopkins for analysis support, Rick

Reynolds and Carl Herzog for logistical support, and the many landowners for site access.

FUNDING

The research was funded by NSF grants DEB-1115895 to A.M.K. & J.T.F., DEB-1911853 to

K.E.L., J.R.H., A.M.K. & J.T.F., the USFWS (F17AP00591) to K.E.L., and by Virginia Tech

Institute for Critical Technology and Applied Science to M.J.K.

DISCLAIMER

Any use of trade, firm, or product names is for descriptive purposes only and does not imply

endorsement by the U.S. Government.

AUTHOR CONTRIBUTIONS

M.J.K.: conceptualization, investigation, methodology, visualization, data curation, writing-original draft, writing-review and editing, formal analysis; J.R.H.: conceptualization, investigation, methodology, visualization, funding acquisition, resources, project administration, data curation, writing-review and editing; J.P.W.: investigation, writing-review and editing; H.M.K.: investigation, writing-review and editing; J.A.R.: investigation; A.E.L.: resources, data curation, writing-review and editing; T.E.R.: resources, writing-review and editing; J.E.D.: investigation, writing-review and editing; W.H.S.: investigation; K.L.P.: investigation, writing-review and editing; J.T.F.: investigation, funding acquisition, writing-review and editing; A.M.K.: investigation, funding acquisition, writing-review and editing; K.E.L.: conceptualization, investigation, visualization, methodology, funding acquisition, resources, project administration, data curation, writing-original draft, writing-review and editing, supervision

All authors gave final approval for publication and agreed to be held accountable for the work performed therein.

DATA, CODE, AND MATERIALS
The datasets and code generated in this study have been uploaded as electronic supporting material for review and will be deposited in Dryad Digital Repository upon final submission. Exact site locations are not disclosed to protect endangered species and landowners.

CONFLICT OF INTEREST DECLARATION
The authors declare no competing interests.

ETHICS
All bat handling procedures were reviewed and approved by Virginia Tech Institute for Animal Care and Use Committee protocol 17-180.

LITERATURE CITED

1.   Daszak P, Cunningham AA, Hyatt AD. Emerging infectious diseases of wildlife - Threats to biodiversity and human health. Science. 2000;287(5452):443-9.

2.   De Castro F, Bolker B. Mechanisms of disease-induced extinction. Ecology Letters.
2004;8(1):117-26.
3.   Silk MJ, Hodgson DJ, Rozins C, Croft DP, Delahay RJ, Boots M, McDonald RA.
Integrating social behaviour, demography and disease dynamics in network models: Applications
to disease management in declining wildlife populations. Philosophical Transactions of the
Royal Society B: Biological Sciences. 2019;374(1781).
4.   Lloyd-Smith JO, Schreiber SJ, Kopp PE, Getz WM. Superspreading and the effect of
individual variation on disease emergence. Nature. 2005;438(7066):355-9.
5.   VanderWaal KL, Ezenwa VO. Heterogeneity in pathogen transmission: mechanisms and
methodology. Functional Ecology. 2016;30(10):1606-22.
6.   Rogers W, Brandell EE, Cross PC. Epidemiological differences between sexes affect
management efficacy in simulated chronic wasting disease systems. Journal of Applied Ecology.
2022;59(4):1122-33.
7.   Silk MJ, Croft DP, Delahay RJ, Hodgson DJ, Boots M, Weber N, McDonald RA. Using
Social Network Measures in Wildlife Disease Ecology, Epidemiology, and Management.
Bioscience. 2017;67(3):245-57.
8.   Gorsich EE, Ezenwa VO, Cross PC, Bengis RG, Jolles AE. Context-dependent survival,
fecundity and predicted population-level consequences of brucellosis in African buffalo. Journal
of Animal Ecology. 2015;84(4):999-1009.
9.   Perkins SE, Ferrari MF, Hudson PJ. The effects of social structure and sex-biased
transmission on macroparasite infection. Parasitology. 2008;135(13):1561-9.
10.   Rogers LM, Delahay R, Cheeseman CL, Langton S, Smith GC, Clifton-Hadley RS.
Movement of badgers (Meles meles) in a high–density population: individual, population and
disease effects. Proceedings of the Royal Society of London Series B: Biological Sciences.
1998;265(1403):1269-76.
11.   Graham J, Smith GC, Delahay RJ, Bailey T, McDonald RA, Hodgson D. Multi-state
modelling reveals sex-dependent transmission, progression and severity of tuberculosis in wild
badgers. Epidemiology and Infection. 2013;141(7):1429-36.
12.   Ferrari N, Cattadori IM, Nespereira J, Rizzoli A, Hudson PJ. The role of host sex in
parasite dynamics: field experiments on the yellow-necked mouse Apodemus flavicollis.
Ecology Letters. 2003;7(2):88-94.
13.   Grear DA, Luong LT, Hudson PJ. Sex-biased transmission of a complex life-cycle
parasite: Why males matter. Oikos. 2012;121(9):1446-53.
14.   Stephenson JF. Parasite-induced plasticity in host social behaviour depends on sex and
susceptibility. Biology Letters. 2019;15(11).
15.   Ezenwa VO, Snider MH. Reciprocal relationships between behaviour and parasites
suggest that negative feedback may drive flexibility in male reproductive behaviour. Proceedings
of the Royal Society B: Biological Sciences. 2016;283(1831):20160423.
16.   Sweeny AR, Corripio-Miyar Y, Bal X, Hayward AD, Pilkington JG, McNeilly TN,
Nussey DH, Kenyon F. Longitudinal dynamics of co-infecting gastrointestinal parasites in a wild
sheep population. Parasitology. 2022;149(5):593-604.
17.   Cattadori IM, Boag B, Bjørnstad ON, Cornell SJ, Hudson PJ. Peak shift and
epidemiology in a seasonal host-nematode system. Proceedings of the Royal Society B:
Biological Sciences. 2005;272(1568):1163-9.

18.   Altizer S, Davis AK, Cook KC, Cherry JJ. Age, sex, and season affect the risk of
mycoplasmal conjunctivitis in a southeastern house finch population. Canadian Journal of
Zoology. 2004.
19.   Altizer S, Dobson A, Hosseini P, Hudson P, Pascual M, Rohani P. Seasonality and the
dynamics of infectious diseases. Ecology Letters. 2006;9(4):467-84.
20.   Miller MR, White A, Wilson K, Boots M. The Population Dynamical Implications of
Male-Biased Parasitism in Different Mating Systems. PLOS ONE. 2007;2(7):e624.
21.   Rosa GM, Bosch J, Martel A, Pasmans F, Rebelo R, Griffiths RA, Garner TWJ.
Sex biased disease dynamics increase extinction risk by impairing population recovery. Animal
Conservation. 2019;22(6):579-88.
22.   Härkönen T, Harding K, Rasmussen TD, Teilmann J, Dietz R. Age- and Sex-Specific
Mortality Patterns in an Emerging Wildlife Epidemic: The Phocine Distemper in European
Harbour Seals. PLOS ONE. 2007;2(9):e887.
23.   Úbeda F, Jansen VAA. The evolution of sex-specific virulence in infectious diseases.
Nature Communications. 2016;7.
24.   Hall MD, Mideo N. Linking sex differences to the evolution of infectious disease life-
histories. Philosophical Transactions of the Royal Society B: Biological Sciences.
2018;373(1757).
25.   Guerra-Silveira F, Abad-Franch F. Sex Bias in Infectious Disease Epidemiology: Patterns
and Processes. PLOS ONE. 2013;8(4).
26.   McDonald JL, Smith GC, McDonald RA, Delahay RJ, Hodgson D. Mortality trajectory
analysis reveals the drivers of sex-specific epidemiology in natural wildlife–disease interactions.
Proceedings of the Royal Society B: Biological Sciences. 2014;281(1790).
27.   Zuk M. The Sicker Sex. PLoS Pathogens. 2009;5(1):e1000267-e.
28.   Zuk M, McKean KA. Sex Differences in Parasite Infections: Patterns and Processes.
International Journal for Parasitology. 1996;26(10):1009-24.
29.   Ezenwa VO, Stefan Ekernas L, Creel S. Unravelling complex associations between
testosterone and parasite infection in the wild. Functional Ecology. 2012;26(1):123-33.
30.   Foo YZ, Nakagawa S, Rhodes G, Simmons LW. The effects of sex hormones on immune
function: a meta-analysis. Biological Reviews. 2017;92(1):551-71.
31.   Klein SL. Hormonal and immunological mechanisms mediating sex differences in
parasite infection. Parasite Immunology. 2004;26(6-7):247-64.
32.   Klein SL. The effects of hormones on sex differences in infection: From genes to
behavior. Neuroscience & Biobehavioral Reviews. 2000;24(6):627-38.
33.   Klein SL. Hormones and mating system affect sex and species differences in immune
function among vertebrates. Behavioural Processes. 2000;51(1-3):149-66.
34.   Morales-Montor J, Chavarria A, León MAD, Castillo LID, Escobedo EG, Sánchez EN,
Vargas JA, Hernández-Flores M, Romo-González T, Larralde C. Host gender in parasitic
infections of mammals: an evaluation of the female host supremacy paradigm. Journal of
Parasitology. 2004;90(3):531-46.
35.   Christe P, Glaizot O, Evanno G, Bruyndonckx N, Devevey G, Yannic G, Patthey P,
Maeder A, Vogel P, Arlettaz R. Host sex and ectoparasites choice: preference for, and higher
survival on female hosts. Journal of Animal Ecology. 2007;76(4):703-10.
36.   Sanchez A, Devevey G, Bize P. Female-biased infection and transmission of the
gastrointestinal nematode Trichuris arvicolae infecting the common vole, Microtus arvalis.
International Journal for Parasitology. 2011;41(13-14):1397-402.

37.   Kiffner C, Stanko M, Morand S, Khokhlova IS, Shenbrot GI, Laudisoit A, Leirs H,
Hawlena H, Krasnov BR. Sex-biased parasitism is not universal: Evidence from rodent-flea
associations from three biomes. Oecologia. 2013;173(3):1009-22.
38.   Hillegass MA, Waterman JM, Roth JD. The influence of sex and sociality on parasite
loads in an African ground squirrel. Behavioral Ecology. 2008;19(5):1006-11.
39.   McCurdy DG, Shutler D, Mullie A, Forbes MR. Sex-Biased Parasitism of Avian Hosts:
Relations to Blood Parasite Taxon and Mating System. Oikos. 1998;82(2):303-12.
40.   Duneau D, Luijckx P, Ruder LF, Ebert D. Sex-specific effects of a parasite evolving in a
female-biased host population. BMC Biology. 2012;10(1):104.
41.   Kelly CD, Stoehr AM, Nunn C, Smyth KN, Prokop ZM. Sexual dimorphism in immunity
across animals: a meta-analysis. Ecology Letters. 2018;21(12):1885-94.
42.   Frick WF, Pollock JF, Hicks AC, Langwig KE, Reynolds DS, Turner GG, Butchkoski
CM, Kunz TH. An Emerging Disease Causes Regional Population Collapse of a Common North
American Bat Species. Science. 2010;329(5992):679-82.
43.   Langwig KE, Frick WF, Hoyt JR, Parise KL, Drees KP, Kunz TH, Foster JT, Kilpatrick
AM. Drivers of variation in species impacts for a multi-host fungal disease of bats. Philosophical
Transactions of the Royal Society B: Biological Sciences. 2016;371(1709).
44.   Langwig KE, Frick WF, Bried JT, Hicks AC, Kunz TH, Marm Kilpatrick A. Sociality,
density-dependence and microclimates determine the persistence of populations suffering from a
novel fungal disease, white-nose syndrome. Ecology Letters. 2012;15(9):1050-7.
45.   Frick WF, Puechmaille SJ, Hoyt JR, Nickel BA, Langwig KE, Foster JT, Barlow KE,
Bartonička T, Feller D, Haarsma A-J, et al. Disease alters macroecological patterns of North
American bats. Global Ecology & Biogeography. 2015;24(7):741-9.
46.   Hoyt JR, Kilpatrick AM, Langwig KE. Ecology and impacts of white-nose syndrome on
bats. Nature Reviews Microbiology. 2021;19(3):196-210.
47.   Warnecke L, Turner JM, Bollinger TK, Lorch JM, Misra V, Cryan PM, Wibbelt G,
Blehert DS, Willis CKR. Inoculation of bats with European Geomyces destructans supports the
novel pathogen hypothesis for the origin of white-nose syndrome. Proceedings of the National
Academy of Sciences of the United States of America. 2012;109(18):6999-7003.
48.   Warnecke L, Turner JM, Bollinger TK, Misra V, Cryan PM, Blehert DS, Wibbelt G,
Willis CKR. Pathophysiology of white-nose syndrome in bats: a mechanistic model linking wing
damage to mortality. Biology Letters. 2013;9(4).
49.   Verant ML, Meteyer CU, Speakman JR, Cryan PM, Lorch JM, Blehert DS. White-nose
syndrome initiates a cascade of physiologic disturbances in the hibernating bat host. 2014.
50.   Cryan PM, Meteyer CU, Boyles JG, Blehert DS. Wing pathology of white-nose
syndrome in bats suggests life-threatening disruption of physiology. BMC Biology. 2010;8:135.
51.   McGuire LP, Mayberry HW, Willis CKR. White-nose syndrome increases torpid
metabolic rate and evaporative water loss in hibernating bats. American Journal of Physiology -
Regulatory Integrative and Comparative Physiology. 2017;313(6):R680-R6.
52.   Wilcox A, Warnecke L, Turner JM, McGuire LP, Jameson JW, Misra V, Bollinger TC,
Willis CKR. Behaviour of hibernating little brown bats experimentally inoculated with the
pathogen that causes white-nose syndrome. Animal Behaviour. 2014;88:157-64.
53.   Reeder DM, Frank CL, Turner GG, Meteyer CU, Kurta A. Frequent Arousal from
Hibernation Linked to Severity of Infection and Mortality in Bats with White-Nose Syndrome.
PLoS ONE. 2012;7(6):38920.

54.   Lilley TM, Johnson JS, Ruokolainen L, Rogers EJ, Wilson CA, Schell SM, Field KA,
Reeder DAM. White-nose syndrome survivors do not exhibit frequent arousals associated with
Pseudogymnoascus destructans infection. Frontiers in Zoology. 2016;13(1).
55.   Lorch JM, Meteyer CU, Behr MJ, Boyles JG, Cryan PM, Hicks AC, Ballmann AE,
Coleman JTH, Redell DN, Reeder DM, et al. Experimental infection of bats with Geomyces
destructans causes white-nose syndrome. Nature. 2011;480(7377):376-8.
56.   Hoyt JR, Langwig KE, Sun K, Parise KL, Li A, Wang Y, Huang X, Worledge L, Miller
H, White JP, et al. Environmental reservoir dynamics predict global infection patterns and
population impacts for the fungal disease white-nose syndrome. Proceedings of the National
Academy of Sciences. 2020:201914794.
57.   Hoyt JR, Langwig KE, White JP, Kaarakka HM, Redell JA, Kurta A, Depue JE, Scullon
WH, Parise KL, Foster JT, et al. Cryptic connections illuminate pathogen transmission within
community networks. Nature. 2018;563:710-3.
58.   Laggan NA, Parise KL, White JP, Kaarakka HM, Redell JA, DePue JE, Scullon WH,
Kath J, Foster JT, Kilpatrick AM, et al. Host abundance and heterogeneity in infectiousness
determine extent of the environmental reservoir. bioRxiv. 2022:2022.09.20.508714.
59.   Langwig KE, White JP, Parise KL, Kaarakka HM, Redell JA, DePue JE, Scullon WH,
Foster JT, Kilpatrick AM, Hoyt JR. Mobility and infectiousness in the spatial spread of an
emerging fungal pathogen. Journal of Animal Ecology. 2021;90(5):1134-41.
60.   Verant ML, Boyles JG, Waldrep W, Wibbelt G, Blehert DS. Temperature-Dependent
Growth of Geomyces destructans, the Fungus That Causes Bat White-Nose Syndrome. PLoS
ONE. 2012;7(9):46280.
61.   Perry RW. A review of factors affecting cave climates for hibernating bats in temperate
North America. Environmental Reviews 21: 28-39. 2013;21:28-39.
62.   Webb PI, Speakman JR, Racey PA. How hot is a hibernaculum? A review of the
temperatures at which bats hibernate. Canadian Journal of Zoology. 1996;74(4):761-5.
63.   Langwig KE, Frick WF, Reynolds R, Parise KL, Drees KP, Hoyt JR, Cheng TL, Kunz
TH, Foster JT, Marm Kilpatrick A. Host and pathogen ecology drive the seasonal dynamics of a
fungal disease, white-nose syndrome. Proceedings of the Royal Society B: Biological Sciences.
2015;282(1799).
64.   Burns LE. Dynamics of Autumn Swarming and Population Structure of Little Brown and
North Myotis Bats (Myotis lucifugus and M. septentrionalis): Dalhousie University; 2014.
65.   Burns LE, Broders HG. Maximizing mating opportunities: higher autumn swarming
activity in male versus female Myotis bats. Journal of Mammalogy. 2015;96(6):1326-36.
66.   Czenze ZJ, Jonasson KA, Willis CKR. Thrifty females, frisky males: Winter energetics
of hibernating bats from a cold climate. Physiological and Biochemical Zoology.
2017;90(4):502-11.
67.   Jonasson KA, Willis CKR. Changes in Body Condition of Hibernating Bats Support the
Thrifty Female Hypothesis and Predict Consequences for Populations with White-Nose
Syndrome. PLoS ONE. 2011;6(6).
68.   Grieneisen LE, Brownlee-Bouboulis SA, Johnson JS, Reeder DM. Sex and hibernaculum
temperature predict survivorship in white-nose syndrome affected little brown myotis (Myotis
lucifugus). Royal Society Open Science. 2015;2.
69.   Jonasson KA, Willis CKR. Hibernation energetics of free-ranging little brown bats.
Journal of Experimental Biology. 2012;215(12):2141-9.

70.   Racey P, Entwistle A, Crichton E, Krutzsch P. Reproductive biology of bats. Academic
Press, London; 2000.
71.   Thomas DW, Brock Fenton M, Barclay RMR. Social behavior of the little brown bat,
Myotis lucifugus. Behavioral Ecology and Sociobiology. 1979;6(2):129-36.
72.   Kurta A, Kunz TH. Size of bats at birth and maternal investment during pregnancy.
Symposia of the Zoological Society of London. 1987;57:79-106.
73.   Wimsatt WA. Notes on Breeding Behavior, Pregnancy, and Parturition in Some
Vespertilionid Bats of the Eastern United States. Journal of Mammalogy. 1945;26(1):23.
74.   Muller LK, Lorch JM, Lindner DL, O'Connor M, Gargas A, Blehert DS. Bat white-nose
syndrome: a real-time TaqMan polymerase chain reaction test targeting the intergenic spacer
region of Geomyces destructans. Mycologia. 2013;105(2):253-9.
75.   Bates D, Mächler M, Bolker B, Walker S. Fitting Linear Mixed-Effects Models Using
lme4. Journal of Statistical Software. 2015;67(1):1 - 48.
76.   McGuire LP, Turner JM, Warnecke L, McGregor G, Bollinger TK, Misra V, Foster JT,
Frick WF, Kilpatrick AM, Willis CKR. White-nose syndrome disease severity and a comparison
of diagnostic methods. EcoHealth. 2016;13(1):60-71.
77.   Langwig KE, Hoyt JR, Parise KL, Frick WF, Foster JT, Kilpatrick AM. Resistance in
persisting bat populations after white-nose syndrome invasion. Philosophical Transactions of the
Royal Society B: Biological Sciences. 2017;372(1712):20160044.
78.   Hopkins SR, Fleming Davies AE, Belden LK, Wojdak JM. Systematic review of
modeling assumptions and empirical evidence: does parasite transmission increase nonlinearly
with host density? Methods in Ecology and Evolution. 2020.
79.   Hopkins SR, Hoyt JR, White JP, Kaarakka HM, Redell JA, DePue JE, Scullon WH,
Kilpatrick AM, Langwig KE. Continued preference for suboptimal habitat reduces bat survival
with white-nose syndrome. Nature Communications. 2021;12(1):166.
80.   Frick WF, Puechmaille SJ, Willis CKR. White-Nose Syndrome in Bats. In: Voigt CC,
Kingston T, editors. Bats in the Anthropocene: Conservation of Bats in a Changing World.
Cham: Springer International Publishing; 2016. p. 245-62.
81.   Hoyt JR, Langwig KE, White JP, Kaarakka HM, Redell JA, Parise KL, Frick WF, Foster
JT, Kilpatrick AM. Field trial of a probiotic bacteria to protect bats from white-nose syndrome.
Scientific Reports. 2019;9(1):9158.
82.   Maslo B, Valent M, Gumbs JF, Frick WF. Conservation implications of ameliorating
survival of little brown bats with white-nose syndrome. Ecological Applications.
2015;25(7):1832-40.
83.   Luong LT, Perkins SE, Grear DA, Rizzoli A, Hudson PJ. The relative importance of host
characteristics and co-infection in generating variation in Heligmosomoides polygyrus fecundity.
Parasitology. 2010;137(6):1003-12.
84.   Jenkins HE, Cox DR, Delahay RJ. Direction of Association between Bite Wounds and
Mycobacterium bovis Infection in Badgers: Implications for Transmission. PLoS ONE.
2012;7(9).
85.   Delahay RJ, Walker NJ, Forrester GJ, Harmsen B, Riordan P, Macdonald DW, Newman
C, Cheeseman CL. Demographic correlates of bite wounding in Eurasian badgers, Meles meles
L., in stable and perturbed populations. Animal Behaviour. 2006;71(5):1047-55.
86.   Folstad I, Karter AJ. Parasites, bright males, and the immunocompetence handicap.
American Naturalist. 1992;139(3):603-22.

87.   Edler R, Goymann W, Schwabl I, Friedl TWP. Experimentally elevated testosterone
levels enhance courtship behaviour and territoriality but depress acquired immune response in
Red Bishops Euplectes orix. Ibis. 2011;153(1):46-58.
88.   Roberts M, Peters A. Is testosterone immunosuppressive in a condition-dependent
manner? An experimental test in blue tits. Journal of Experimental Biology. 2009;212(12):1811-
8.
89.   Prendergast BJ, Freeman DA, Zucker I, Nelson RJ. Periodic arousal from hibernation is
necessary for initiation of immune responses in ground squirrels. American Journal of
Physiology - Regulatory Integrative and Comparative Physiology. 2002;282(4 51-4).
90.   Bouma HR, Carey HV, Kroese FGM. Hibernation: the immune system at rest? Journal of
Leukocyte Biology. 2010;88(4):619-24.
91.   Meyer GA, Senulis JA, Reinartz JA. Effects of temperature and availability of insect prey
on bat emergence from hibernation in spring. Journal of Mammalogy. 2016;97(6):1623-33.
92.   Johnson JS, Reeder DM, McMichael JW, Meierhofer MB, Stern DWF, Lumadue SS,
Sigler LE, Winters HD, Vodzak ME, Kurta A, et al. Host, pathogen, and environmental
characteristics predict white-nose syndrome mortality in captive little brown myotis (Myotis
lucifugus). PLoS ONE. 2014;9(11).
93.   Davis WH, Hitchcock HB. Biology and migration of the bat, Myotis lucifugus, in New
England. Journal of Mammalogy. 1965;46(2):296-313.
94.   Mohr CE. Sex ratios of bats in Pennsylvania. Proceedings of the Pennsylvania Academy
of Science. 1945;19:65-9.
95.   Tinkle DW, Milstead WW. Sex Ratios and Population Density in Hibernating Myotis.
The American Midland Naturalist. 1960;63(2):327-34.
96.   Lynda MSL, William C. Sex ratio variation of Myotis velifer (Chiroptera:
Vespertilionidae) in Oklahoma The Southwestern Naturalist. 2007;52(1):67-74.
97.   Willis CKR, Brigham RM. Social thermoregulation exerts more influence than
microclimate on forest roost preferences by a cavity-dwelling bat. Behavioral Ecology and
Sociobiology. 2007;62(1):97-108.
98.   Linton DM, Macdonald DW. Spring weather conditions influence breeding phenology
and reproductive success in sympatric bat populations. Journal of Animal Ecology.
2018;87(4):1080-90.
99.   Le Galliard J-F, Fitze Patrick S, Ferrière R, Clobert J. Sex ratio bias, male aggression,
and population collapse in lizards. Proceedings of the National Academy of Sciences.
2005;102(50):18231-6.
100.   Johanos TC, Becker BL, Baker JD, Ragen TJ, Gilmartin WG, Gerrodette T. Impacts of
sex ratio reduction on male aggression in the Critically Endangered Hawaiian monk seal
Monachus schauinslandi. Endangered Species Research. 2010;11(2):123-32.
101.   Grimaudo AT, Hoyt JR, Yamada SA, Herzog CJ, Bennett AB, Langwig KE. Host traits
and environment interact to determine persistence of bat populations impacted by white-nose
syndrome. Ecology Letters. 2022;25(2):483-97.
701