Geobacter sulfurreducens' unique metabolism results in cells with a high iron and lipid content

Ethan Howley[1,3], Dongwon Ki[1,4], Rosa Krajmalnik-Brown[1,3], César I. Torres[1,2]

[1] Biodesign Swette Center for Environmental Biotechnology, Arizona State University, USA

[2] School for Engineering of Matter Transport and Energy, Arizona State University, USA

[3] School for Sustainable Engineering and the Built Environment, Arizona State University, USA

[4] Division of Living and the Built Environment Research, Seoul Institute of Technology, South Korea

Abstract

Geobacter sulfurreducens is a ubiquitous iron reducing bacterium in soils, and in engineered systems it can respire an electrode to produce measurable electric current. Its unique metabolism, heavily dependent on an extensive network of cytochromes, requires a unique cell composition. In this work we used metallomics, cell fraction and elemental analyses, and transcriptomics to study and analyze the cell composition of *G. sulfurreducens*. Elemental composition studies (C,H,O,N, ash content) showed a high C:O and H:O ratios of approximately 1.7:1 and 0.25:1, indicative of more reduced cell composition that is consistent with a high lipid content. Our study shows that *G. sulfurreducens* cells have a large amount of iron (2 ± 0.2 µg/gdw) and lipids ($32 \pm 0.5\%$ dw/dw) and that this composition does not change whether the cells are grown with a soluble or an insoluble electron acceptor. The high iron concentration, higher than similar microorganisms, is attributed to the production of cytochromes that are abundant in

transcriptomic analyses in both solid and soluble electron acceptor growth. The unique cell

composition of *G. sulfurreducens* must be considered when growing this microorganism for lab

studies and commercial applications.

Importance

Geobacter sulfurreducens is an electroactive microorganism. In nature, it grows on metallic

minerals by transferring electrons to them, effectively 'breathing' metals. In a manmade system,

it respires an electrode to produce an electric current. It has become a model organism for the

study of electroactive organisms. There are potential biotechnological applications of an

organism that can bridge the gap between biology and electrical signal, and as a ubiquitous iron

reducer in soils around the world, *G. sulfurreducens* and its relatives impact the global iron cycle.

We measured the concentrations of metals, macromolecules, and basic elements in *G.*

sulfurreducens to define this organism's composition. We also used gene expression data to

discuss which proteins those metals could be associated with. We found that *G. sulfurreducens*

has a large amount of lipid and iron compared to other bacteria — these observations are

important for future microbiologists and biotechnologists working with the organism.

Introduction

Anode-respiring electroactive bacteria , such as *Geobacter sulfurreducens*, have been

studied for almost two decades for their capability to produce electrical current from metabolic

respiration of organic compounds while in multi-layered biofilms (1–3) . A unique feature of

these biofilms is the extracellular matrix that allows the transport of electrons over tens of

micrometers (4–6). As part of this extracellular matrix, several components have been proposed

to be crucial in achieving extracellular electron transport (EET). The transport of electrons starts

at the inner membrane, and travels across the periplasm and outer membrane before it reaches

the extracellular environment. Cytochromes at these locations are known to play an important

role in delivering electrons outside the cell (7–10). Microbial nanowires, now also identified as

cytochrome polymers (4, 6), are the main path by which electrons are thought to be conducted in

the extracellular environment, reaching a solid electron acceptor. Extracellular polymeric

substances (EPS) have also been proposed to play a role in EET in *G. sulfurreducens* as well (3,

11). On the other hand, *Shewanella oneidensis* MR-1 has been shown to produce outer

membrane and periplasmic extensions, which are lipid bilayers and contain extracellular

cytochromes (12, 13). In both cases, it is clear that the EET mechanism creates an extra

metabolic burden to electroactive organisms and that it can alter their cell composition and

nutrient requirements when compared against microorganisms performing respiration of soluble

electron acceptors. These nutrient requirements, however, have not been assessed in a systematic

way.

Transcriptomic and proteomic studies in *G. sulfurreducens* have highlighted the

importance of respiratory and EET proteins for their growth on anodes and metal oxides.

Extracellular and outer membrane proteins, including pili, outer membrane channels, and c-type

membrane cytochromes, are essential to the metabolism of *G. sulfurreducens* (9, 14–26). The

high abundance of these proteins and other possible extracellular components may result in a

unique cellular composition. For example, each cytochrome contains one or more iron-

containing heme complexes which can influence the iron content of the cell. An analysis of cell

composition can provide insights into the composition of the extracellular matrix and EET-

related components.

Several studies have provided insights into the cellular composition of *G. sulfurreducens*.

For example, the lipid fraction of *Geobacter sulfurreducens* was reported 15% wt/wt by

Mahadevan et al., 2006, where an additional 4% lipopolysaccharides fraction was assumed (27).

In comparison, the lipid in *Escherichia coli* has been reported to be 9.1% wt (28) while

cyanobacterium *Synechocystis* sp. PCC6803, known to produce thylakoid membranes, has been

reported to be as high as 14% wt(29) and the lipid-rich microalgae *Schizochytrium* sp. can

contain up to 30% lipid (30). The high lipid content of *G. sulfurreducens* has potential

implications for biotechnology applications.

To our knowledge, few studies have performed a metallomic analysis in *G.*

sulfurreducens, and the existing literature has conflicting results. Previous research showed that

G. sulfurreducens, when grown on fumarate as electron acceptor, has similar metal content to

that of *E. coli*. On the other hand, the closely related organism *Geobacter metallireducens* grown

on iron citrate showed an order of magnitude higher iron content; but the possible formation of

inorganic precipitates was reported to be a possible hindrance to the measurement (31). Another

study found that *G. sulfurreducens* had a per cell iron content an order of magnitude higher than

E. coli, and that limiting growth medium iron content inhibited EET in *G. sulfurreducens* (15).

In this study, we hypothesized that *G. sulfurreducens* has a significantly different cellular

composition compared to other cells performing soluble respiratory metabolisms. The

differences that stem out of the EET requirements can help explain how EET develops.

Understanding these characteristics can lead to a better growth and maintenance of this

microorganisms in laboratory and applied systems. We performed a metallomic analysis on *G.*

sulfurreducens grown on an anode versus fumarate as electron acceptors and compared it to *E.*

coli K12. The use of an anode allows us to study EET and eliminates the interference of possible

iron oxides reported by Budhraja et al. (31). We also performed an elemental analysis (C, H, O,

N, ash content) and a fraction analysis (protein, carbohydrates, and proteins) to obtain a

comprehensive cell composition and determine if the previously observed lipid fractions

measured in fumarate samples are also observed during anodic respiration. The results are

complemented with a transcriptomic analysis of *G. sulfurreducens* grown under similar

conditions.

Materials and Methods

Bacterial strain and culture media

We subcultured *Geobacter sulfurreducens* PCA (ATCC 51573) and *Escherichia coli* K-

12 from commercially available stocks. Medium compositions are listed in detail in Table S1 for

four different cases: 1. *G. sulfurreducens* grown in microbial electrochemical cell (electrode), 2.

G. sulfurreducens grown in sodium fumarate-containing serum bottle (fumarate), 3. *E. coli*

grown in *Geobacter* medium, and 4. *E. coli* grown in M9 medium. In brief, *Geobacter* medium

contained sodium acetate (50 mM), $NaHCO_3$ (30 mM), NH_4Cl (20 mM), NaH_2PO_4 (4 mM), KCl

(1 mM), vitamin mix (10 mL), and trace minerals (10 mL). Trace minerals contained

Nitrilotriacetic acid, trisodium salt (5.5 mM), $MgSO_4·7H_2O$ (12 mM), $MnSO_4·H_2O$ (2.9 mM),

NaCl (17 mM), $FeSO_4·7H_2O$ (0.36 mM), $CaCl_2·2H_2O$ (0.68 mM), $CoCl_2·6H_2O$ (0.42 mM),

$ZnCl_2$ (0.95 mM), $CuSO_4·5H_2O$ (0.4 mM), $AlK(SO_4)_2·12H_2O$ (0.2 mM), H_3BO_4 (0.16 mM),

$Na_2MoO_4·H_2O$ (0.01 mM), $NiCl_2·6H_2O$ (0.01 mM), and $Na_2WO_4·2H_2O$ (8.5 µM). We provided

higher concentration of acetate (50 mM) than the previous studies[32–34] (10 mM), as we used a

higher electrode surface area requiring more reduced electron donor. For *G. sulfurreducens*

grown in serum bottles, sodium fumarate (100 mM) was added in the medium. We bubbled the

media with N_2/CO_2 (80:20 v/v) to remove oxygen before autoclaving. After autoclaving,

$FeCl_2·4H_2O$ (20 µM), $Na_2S·9H_2O$ (54 µM), sodium bicarbonate, and vitamins were added in

anaerobic glove box. *E. coli* medium (ATCC Medium 2511 - M9 Minimal Broth) contained

glucose (44 mM), Na_2HPO_4 (180 mM), KH_2PO_4 (44 mM), NaCl (17 mM), NH_4Cl (37 mM),

$MgSO_4·7H_2O$ (0.2 mM), CaCl2 (10 µM), and thiamin (24 µM). Media for *E. coli* (M9 and

Geobacter medium) were autoclaved without any gas sparging. *Geobacter* medium for *E. coli*

contained the same ingredients of *G. sulfurreducens* medium for microbial electrochemical cells

except for the electron donor; the same concentration of glucose as M9 (44 mM) was used

instead of acetate.

Electrochemical setup and operation

Single-chamber microbial electrochemical cells were constructed in 500 mL bottles with

rubber stoppers located on top having PTFE tubing for gas inflow and outflow, carbon anode as

working electrode, nickel wire cathode as counter electrode, and reference electrode. We used

two square graphite electrodes to grow biofilms of *G. sulfurreducens* with a surface area of

~20.9 cm^2, and an Ag/AgCl reference electrode (BASi, West Lafayette, IN). We mixed the

chambers with magnetic stirrer bars at 180 rpm and flushed humidified N_2/CO_2 gas (80:20 v/v)

continuously. Before filling up the media in the anaerobic glove box, electrochemical cells were

autoclaved for sterilization. We set -0.3 V vs Ag/AgCl (-0.03 V vs. SHE) as fixed anode

potential using a VMP3 digital potentiostat (Bio-Logic USA, Knoxville, TN). This is at the

lower end of the range of electrode potentials that *G. sulfurreducens* can respire. Fumarate-

grown *G. sulfurreducens* reactors were set up in 250 mL serum bottles. Both electrochemical

cells and serum bottles were in a temperature-controlled room at 30 °C. Also, the media filling

along with inoculation was performed in the anaerobic glove box. We used 250 mL flasks for *E.*

coli cultivation with M9 and *Geobacter* media. *Geobacter* medium (G medium) for *E. coli* was

used to compare the cell composition with *G. sulfurreducens*. After filling the media and

inoculation for *E. coli*, we placed the flasks in an incubator with shaking at 180 rpm and

temperature at 37 °C.

Sample preparation

For the determination of carbohydrate, protein, lipid, metal, and element per dried cell,

we collected the *G. sulfurreducens* grown on anodes in the microbial electrochemical cells and in

serum bottle reactors and the *E. coli* grown in flasks. *G. sulfurreducens* biofilms grown on the

anodes were scraped off with a needle in an anaerobic glove box. Grown cells of *G.*

sulfurreducens and *E. coli* from the serum bottles and flasks were separated by centrifugation

(Eppendorf Centrifuge 5810 R, USA) at 4000 rpm in microcentrifuge tubes. Cells were washed

once with a Ringer's solution (25% strength) and centrifuged again (Table S1). The cells dried

overnight at 105 °C in plastic tubes and the cool pellets were then broken up with a sterile

stainless-steel spatula.

Analytical methods

We measured proteins by bicinchoninic acid (BCA) protein assay (35). In brief, dried cell

biomass (2-3 mg) treated with 0.1 N NaOH at 90 °C for 30 min, re-suspended and centrifuged

the lysate, and used 0.1 mL of supernatant for the assay. Carbohydrates were measured by a

colorimetric method (36). In brief, dried cell biomass (2-3 mg) was acidified in sulfuric acid with

sonication for 2 hours, and dissolved samples (0.5 mL) added in the test tubes with distilled

water (0.5 mL), phenol (50 μL), and sulfuric acid (5 mL) for overnight reaction. Concentrations

of proteins and carbohydrates were determined using calibration curve with bovine serum

albumin and glucose with the absorbance at wavelengths of 485 and 562 nm, respectively. Crude

lipids were extracted from the dried cell biomass using the Folch method (37). The dried biomass

(~15 mg) was sonicated for 1 hour and vortexed for 1 hour with Folch solution (chloroform-

methanol, 2:1, v/v) at room temperature. Solvent extracts were obtained after removing the

biomass by centrifugation at 4000 rpm. The crude lipid weight was determined by evaporating in

the water bath (60 °C) and weighing the tube before and after the evaporation of lipid. For metal

extraction, we added dried cell biomass (3-5 mg) to glass vials along with hydrochloric acid (12

M) and sonicated at 60 °C for 2 hours. The dissolved metals in acid were analyzed by

inductively coupled plasma - optical emission spectrometer (ICP-OES, Thermo iCAP6300).

Carbon, hydrogen, and nitrogen in the dried cell biomass (~2 mg) were measured using CHN

Elemental Analyzer (PE2400). Instrumentation for ICP-OES and CHN Elemental Analyzer was

done in Goldwater Environmental Laboratory at Arizona State University. For oxygen estimation,

we measured ash content of the dried cell biomass (~10 mg) following the previous method (38),

burning the biomass at 600 °C in an alumina crucible. Then we subtracted the fraction of C, H, N,

and ash content from the dried cell biomass (100%) for O estimation. Based on the elemental

analysis data (%C, %H, %O, and %N), we obtained the empirical biomass formula followed by

Equation 1 below (39),

$C_N H_a O_b N_c + \left(\frac{2n+0.5a-1.5c-b}{2}\right) O_2 \longrightarrow nCO_2 + cNH_3 + \frac{a-3c}{2} H_2O$ (Eq. 1)

where, n $= \frac{\%C}{12T}$, a $= \frac{\%H}{T}$, b $= \frac{\%O}{16T}$, and c $= \frac{\%N}{14T}$,

and, $T = \frac{\%C}{12T} + \frac{\%H}{T} + \frac{\%O}{16T} + \frac{\%N}{14T}$.

Transcriptomic analysis

We sequenced reverse transcribed RNA from *G. sulfurreducens* grown in a single chamber

microbial electrochemical cell as an anode biofilm with an anode poised at -0.28 V vs. Ag/AgCl

as the electron acceptor, and in anaerobic test tubes with fumarate as the electron acceptor. Both

conditions were collected in biological triplicate. We extracted RNA with the Qiagen/MOBIO

PowerMicrobiome RNA extraction kit and used the ThermoFisher MicrobExpress bacterial

mRNA enrichment kit to reduce the fraction of rRNA following manufacturer's

recommendations. RNA library preparation and sequencing were performed by the genomics

core facility at ASU. We mapped the reads to the RefSeq assembly for *G. sulfurreducens PCA*

and used DESeq2 in R for differential expression analysis. The cutoff determination for

differential expression was set as a $\log_2$ fold change of at least 1.5 and a multiple comparison

adjusted *p*-value of less than 0.05. The sequencing data we used here is a subset of data analyzed

in a previous publication, and raw sequence data is available from NCBI under accession number

GSE200066 (40).

Results and Discussion

Elemental analysis

We analyzed the trace metals and macronutrients of dried *G. sulfurreducens* and *E. coli*

grown in the lab with different environments (Table S3, Table 3, Figure 1). We found significant

differences between the two species in the mass fraction of several trace metals (Figure 1, Table

S3, Figure S1). Media compositions were different in each growth condition, matching each

organism's requirements (Table S2). There were few differences between anode-grown and

fumarate grown *G. sulfurreducens*. Significant differences in Mn and Fe content were observed

in *E. coli* when growing in M9 medium versus *Geobacter* medium. Given the low abundance of

nutrients in M9's minimal medium, significant increases in Cu, Fe, Mn, and Se were observed

when *E. coli* was grown in *Geobacter* medium. We use this latter condition as point of reference

when comparing *G. sulfurreducens* and *E. coli*.

Several metals were present at higher concentrations in anode- and fumarate-grown *G.*

sulfurreducens when compared to *E. coli*, but only a subset of them had statistically significant

differences between the organisms (Table S3, Figure 1).

Figure 1. Relevant differences in metal concentrations between *E. coli*-GS (Geobacter medium),

E. coli-M9 (M9 medium), *G. sulf*-A (biofilm grown on an electrode), and *G. sulf*-F (planktonic

cells using fumarate as the electron acceptor). See Figure S1 for more comparisons. *($p<0.05$),

pairwise t-test with multiple comparison correction performed with the Benjamini-Hochberg

method. We chose to omit alkali metals from this figure, but Lithium did have significant

differences as well (Table S1).

Metals used as cofactors

Fe is a required trace metal for cellular respiration in many organisms as the cofactor in

cytochromes. In both growth conditions for *G. sulfurreducens*, Fe was much higher than in *E.*

coli, which suggest a much higher abundance of Fe-containing metalloproteins and other iron-

containing biomolecules (Figure 1).

In previous studies, listed in Table 1, anaerobic bacteria are reported to have more iron

per cell biomass than aerobic bacteria. Phototrophic bacteria (*Rhodospirillum*,

Rhodopseudomonas, *Chromatium*) and facultative bacteria (*Escherichia*, *Enterobacter*) have

150-500 μg/gdw Fe. *Desulfovibrio vulgaris*, an anaerobic Deltaproteobacterium like *G.*

sulfurreducens, has over 900 μg/gdw. Not included in our analysis are iron oxidizing bacteria,

whose Fe precipitates can lead to Fe concentrations over 2% by dry mass (41). To our

knowledge, *G. sulfurreducens* has the highest Fe content among bacteria studied, with almost

twice the content of *D. vulgaris*. This is consistent with their production of heme-containing

cytochromes in much higher abundance than other microorganisms, leading to not only

cytoplasmic and membrane metalloproteins, but also an extensive abundance of extracellular

cytochromes. Growing *G. sulfurreducens* on the anode versus fumarate did not change the total

Fe amount, suggesting similar abundance of Fe metalloproteins. OmcS nanowires have been

isolated from fumarate culture (6), indicating that *G. sulfurreducens* does not necessarily

downregulate its EET metabolism when it is not needed. Our gene expression data also shows a

high expression of cytochromes associated with EET in fumarate and anode biofilm culture

(Figure 2).

Table 1. Amount of iron in different prokaryotic species

Bacterial species	Iron (ug Fe/gdw)	References
Rhodospirillum rubrum	Light - 202 (± 13)	
	Dark - 198 (± 10)	
Rhodopseudomonas	Light - 163 (± 21)	Kassner & Kamen, 1968*
spheroides	Dark - 230 (± 24)	(42)
Chromatium	Light - 456 (± 76)	
Desulfovibrio vulgaris	952	Lancaster et al., 2014**
Enterobacter cloacae	154	(43)
Escherichia coli	223	Hartmann and Braun, 1981 (44)
	280	Abdul-Tehrani et al., 1999

		(45)
	300	
Micrococcus roseus	200	Rouf, 1964
Bacillus cercus	300~400	(41)
Psudomonas aeruginosa	0.1	Ma et al.,1999 (46)
Shewanella oneidensis	147$^{\square}$	Daly et al., 2004 (47)
Escherichia coli	G medium - 430 (± 90) M9 medium - 130 (± 50)	This study
Geobacter sulfurreducens	Electrode - 1970 (± 226) Fumarate - 1960 (± 229)	

* Photosynthetic bacteria were grown in different growth conditions with and without light

exposure. ** Iron in *D. vulgaris* and *E. cloacae* were estimated with the measured iron per total

protein of the cells; we used a conversion factor of 0.55 to convert from protein to volatile solids

(28). $^{\square}$ *S. oneidensis* iron content was converted from nmol Fe/ mg protein to µg Fe/gdw using

52.8% protein content as measured previously (48).

Fe may be a limiting trace mineral in commonly used *G. sulfurreducens* media. Estevez-

Canales et al. found that a medium concentration of 2 µM Fe limits biomass culture in a

chemostat growing *G. sulfurreducens* led to a culture with 1.9 x 10^{-6} ng of Fe per cell (15). If we

assume the average cell dry weight of *G. sulfurreducens* is between 0.1 and 1 picograms, as has

been shown in *E. coli* (49), our results would give an iron content of 2 x 10^{-7} ng to 2 x 10^{-6} ng Fe

per cell in *G. sulfurreducens*.

Nickel, cobalt, and chrome content were significantly higher in both *G. sulfurreducens*

conditions relative to *E. coli* (Table S3, Figure 1). Nickel is a cofactor in Ni-Fe hydrogenases,

and the genome of *G. sulfurreducens* encodes for several (50, 51). *G. sulfurreducens* is able to

assimilate cobalt through its cobamide-synthesis pathways (52), but it may also be precipitated

on the cell surface as a defense mechanism against cobalt toxicity (53).

Precipitating metals

There are some metals that may be overrepresented in our *G. sulfurreducens* samples due to

precipitation. *G. sulfurreducens* requires at least two multicopper proteins, OmpC and OmpB, to

respire Fe (III) oxide (54), and while these and other metalloproteins are a likely reservoir of Cu

in our samples, *G. sulfurreducens* is capable of reducing Cu(II) to Cu_2S nanoparticles that

associate with cells (55). This phenomenon makes it difficult to estimate how much copper was

required for metalloproteins, and how much may have been trapped as inorganic precipitates. *G.*

sulfurreducens can also immobilize copper through dissimilatory reduction (56). In our data, the

G. sulfurreducens samples were enriched in Cr compared to *E. coli* (Figure 1), while differences

in Cu were not statistically significant. Manganese was significantly lower in the *E. coli* grown

with M9 medium compared to all other conditions including *E. coli* grown with the *G.*

sulfurreducens medium recipe because M9 medium does not contain manganese (Figure 1, Table

S2).

Based on the metal content of the *G. sulfurreducens* cells collected, we can estimate a

maximum cell density from the available mineral content in the common Geobacter medium

(ATCC 1957). Table S4 shows the estimated growth cell assuming cells require the observed

metal concentrations and only have the medium as a source. As expected, Fe is the most limiting

metal in the medium, allowing for only 0.10 g cells/L. Cu and Zn are also close to this limitation

and could lead to a multi-nutrient limitation when growing *G. sulfurreducens* at ~0.1 g/L. This

nutrient limitation can either limit cell density in cell suspensions or limit current generation in

microbial electrochemical technologies when operated in batch mode. Assuming a current

production of ~0.28 A/g protein (57) or 0.6 A/g cell (based on Table 2), one liter of *Geobacter*

medium can support enough *G. sulfurreducens* cells to produce 60 mA, an amount of current that

is enough for most experimental setups but might be limiting in electrochemical cells with a high

specific surface area.

Cell composition of G. sulfurreducens is different to an average bacterium

We also studied the cell composition and elemental analysis (C, H, O, N, Ash) of *G.*

sulfurreducens. Interestingly, the *G. sulfurreducens* cell showed a high abundance of lipids in

both growth conditions (Table 3). The values of ~32% lipid content were much higher than

previously reported and similar to lipid-accumulating algal cultures (58–60). We do not know the

reason why *G. sulfurreducens* requires such a high lipid content. Their smaller diameter (~ 0.5

µm) and distinct morphology (61) compared to other rod-shape bacteria certainly plays a role in

the increased lipid content. *Shewanella oneidensis*, another electrogenic organism, is known to

produce outer membrane extensions for electron transfer (13). While it is likely that our analysis

captured some extracellular polymeric substances, the extracellular matrix of *G. sulfurreducens*

has not been found to have a significant lipid component (62). Most microorganisms exhibiting

this high lipid fraction have either lipid accumulation, as in the case of certain algal species (58–

60) or have internal lipid structures that increase its relative fraction as in the case of thylakoid

membranes and intracytoplasmic membranes (63–65). Our method for lipid quantification would

likely collect other nonpolar compounds like polyhydroxyalkanoates if they were present in the

biofilm, and while *G. sulfurreducens* has not been shown to produce this type of carbon storage

polymer, biofilms enriched in other *Geobacter* species do use some uncharacterized carbon

storage mechanism (66). We anticipate that future studies may add context to the data we present.

The lipid fraction in *E. coli* was lower than *G. sulfurreducens* and higher than has been

previously reported at $24.3 \pm 3.6\%$. However, there is a high variance in reported *E. coli* lipid

content with values ranging from 9 to 19% by dry weight (67, 68).

Because of the higher lipid content, *G. sulfurreducens* cells show a significantly lower

protein content when compared to other microorganisms (~22-26%, Table 3). Fumarate-grown

cells had a larger protein fraction than in anode-grown cells. On the other hand, total

carbohydrates were ~2 times higher in anode-grown cell; exopolysaccharide (EPS) excreted from

G. sulfurreducens to form a biofilm on the electrode probably increases the carbohydrate content

in this growth condition.

Our elemental analysis of *G. sulfurreducens* cells is consistent with the low protein, high

lipid content measured. Following Eq. 1, empirical cell biomass formulas of *G. sulfurreducens*,

normalized to N, were calculated as $C_{5.77}H_{10.61}O_{2.43}N$ for electrode-grown and $C_{6.58}H_{12.48}O_{3.02}N$

for fumarate-grown cells. Compared with general formulas for bacterial biomass, such as

$C_5H_7O_2N$ (39), *G. sulfurreducens* has a higher C:N ratio typical of a low protein content. It also

has a higher hydrogen content, due to the higher lipid content that has approximately a 1:2 for

fumarate-grown cells.

Table 2 Cell compositions of *G. sulfurreducens* and *E. coli* in different growth conditions. Error

is the sample standard deviation.

Growth condition	*G. sulf.* electrode	*G. sulf.* fumarate	*E. coli* M9 medium
Proteins (mg BSA/gdw)	215 ($\pm$ 7)	262 ($\pm$ 56)	284 ($\pm$ 15)

Crude Lipids (mg/gdw)		323 (± 45)	321 (± 35)	243 (± 36)
Carbohydrates (mg glucose/gdw)		193 (± 11)	87 (± 24)	68 (± 19)
Elements	C (%)	47.0 (± 1.1)	46.8 (± 1.8)	46.9 (± 0.7)
	H (%)	7.2 (± 0.6)	7.4 (± 0.1)	7.4 (± 1.0)
	N (%)	9.5 (± 0.2)	8.3 (± 0.3)	12.5 (± 0.0)
	O (%)	26.4 (± 3.3)	28.6 (± 3.9)	24.8 (± 2.0)
	Ash (%)	9.9 (± 3.0)	9.0 (± 3.3)	8.4 (± 1.3)

We compared our cellular composition of *G. sulfurreducens* to that reported in Mahadevan et al.

2006 (27). The main differences between the fraction distributions reported here and those

reported in Mahadevan et al. is the higher lipid content at the expense of a lower protein content.

We do not know the reason for the discrepancy, but in both cases the lipid content is significantly

higher than *E. coli* and other bacterial cells.

Iron-containing genes are highly expressed

Our analysis identified 434 genes that were differentially expressed between the anode biofilm

samples and the planktonic fumarate cells out of 3434 annotated genes detected at quantifiable

levels. 205 genes were more highly expressed in the anode biofilm, and 229 genes were more

highly expressed in the planktonic samples. In Figure 2, MA plots visualize the differential

expression and highlight several types of iron-containing protein-coding genes. While a greater

number of cytochromes were significantly upregulated in the anode biofilm than the number

upregulated in the planktonic samples, most cytochromes were not differentially expressed.

Ferritin domain containing protein- and nonheme Fe-S domain protein-coding genes were also

342 present among the differentially expressed genes. Our data show that iron-containing protein-

343 coding genes are expressed in both planktonic fumarate cultures and anode biofilms, but that

344 there are specific iron-related genes whose expression depends on growth conditions. The

345 abundance of expression of Fe-containing proteins is consistent with the high Fe abundance in

346 both conditions. Our gene expression data is similar in overall patterns to the data in Otero et al.

347 2018, with multiheme cytochromes like the PpcA, OmcZ, and CbcL among the most expressed

348 genes overall, and only a subset of all cytochromes differentially expressed between electrode

349 and fumarate conditions (7). We did, however, observe a greater number of cytochrome genes

350 differentially expressed between electrode and fumarate conditions than Otero et al. 2018, likely

351 due to the lower electrode potential that we used. We also investigated the expression of lipid

352 synthesis pathway genes and found that many of them were present in the transcriptome but not

353 differentially expressed, although a few lipid-associated genes showed significant differences

354 (Figure S2).

Figure 2 MA plots of mRNA gene expression data comparing planktonic cells and anode

biofilms. Positive log2 fold change indicates higher expression in the anode biofilm condition.

Solid points indicate a log2 fold change greater than 1.5 and an adjusted *p*-value below 0.05.

Conclusions

G. sulfurreducens is a bacterium with a complex system of electroactive proteins, and those

electroactive proteins largely require iron. This may be a factor in the high iron concentration we

measured in *G. sulfurreducens* relative to non-electroactive Gram-negative *E. coli* and values

reported in literature for other bacteria. Our analysis complements previous work showing that

restricting iron limits EET in *G. sulfurreducens* (15). This study estimates what the nutrient

limitations might be for *G. sulfurreducens*, and this information is valuable for biotechnologists

developing applications using this and similar organisms. The nearly identical composition

between anode-grown and fumarate-grown cells supports the hypothesis that *G. sulfurreducens*

is not adapted to efficiently grow on fumarate – it makes electron carriers for EET regardless of

the electron acceptor if the nutrients are available. The lipid content measured in *G.*

sulfurreducens was higher than what has been reported before, and relatively high for a

bacterium without lipidic storage. While all samples were taken from active biofilms or

suspended cultures, we did not have a mechanism to separate dead cells from active cells, and it

is probable that the composition of an individual cell may differ from the composition of the bulk

samples analyzed. When compared to similar studies on other bacteria and the *E. coli* in our

study, we have shown that *G. sulfurreducens* has a unique composition to support its complex

metabolism.

381
Acknowledgements

The funding for this work was provided by Office of Naval Research (ONR awards N0014-15-1-

2702 and N0014-20-1-2269). We also thank Roy Erickson and Adam Smith for assistance of

ICP-OES and elemental analysis (Goldwater Environmental Laboratory at ASU) and the

Genomics Core at ASU for sequencing.

References

1.   Torres CI. 2014. On the importance of identifying, characterizing, and predicting

fundamental phenomena towards microbial electrochemistry applications. Curr Opin

Biotechnol 27:107–114.

2.   Reguera G, Nevin KP, Nicoll JS, Covalla SF, Woodard TL, Lovley DR. 2006. Biofilm

and nanowire production leads to increased current in Geobacter sulfurreducens fuel cells.

Appl Environ Microbiol 72:7345–7348.

3.   Bond DR, Strycharz-Glaven SM, Tender LM, Torres ICI. 2012. On Electron Transport

through Geobacter Biofilms. ChemSusChem 5:1099–1105.

4.   Lovley DR, Walker DJFF. 2019. Geobacter Protein NanowiresFrontiers in Microbiology.

Frontiers Media S.A.

5.   Chadwick GL, Jiménez F, Gralnick JA, Bond DR, Orphan VJ, Otero FJ, Gralnick JA,

Bond DR, Orphan VJ, Jiménez F, Gralnick JA, Bond DR, Orphan VJ. 2019. NanoSIMS

imaging reveals metabolic stratification within current-producing biofilms. Proc Natl

Acad Sci 116:20716–20724.

6.   Wang F, Coureuil M, Osinski T, Orlova A, Altindal T, Gesbert G, Nassif X, Egelman EH,

Craig L. 2017. Cryoelectron Microscopy Reconstructions of the Pseudomonas aeruginosa

and Neisseria gonorrhoeae Type IV Pili at Sub-nanometer Resolution. Structure 25:1423-

1435.e4.

7.   Otero FJ, Chi Ho Chan DRB, Jiménez Otero F, Chan CH, Bond DR. 2018. Identification

of Different Putative Outer Membrane Electron Conduits Necessary for Fe(III) Citrate,

Fe(III) Oxide, Mn(IV) Oxide, or Electrode Reduction by Geobacter sulfurreducens. J

Bacteriol 200:1–20.

8.   Levar CE, Hoffman CL, Dunshee AJ, Toner BM, Bond DR. 2017. Redox potential as a

master variable controlling pathways of metal reduction by Geobacter sulfurreducens

11:741–752.

9.   Zacharoff L, Chan CH, Bond DR. 2016. Reduction of low potential electron acceptors

requires the CbcL inner membrane cytochrome of Geobacter sulfurreducens.

Bioelectrochemistry 107:7–13.

10.   Pessanha M, Morgado L, Louro RO, Londer YY, Pokkuluri PR, Schiffer M, Salgueiro CA.

2006. Thermodynamic Characterization of Triheme Cytochrome PpcA from Geobacter

sulfurreducens□: Evidence for a Role Played in e - / H + Energy Transduction.

Biochemistry 45:13910–13917.

11.   Rollefson JB, Stephen CS, Tien M, Bond DR. 2011. Identification of an extracellular

polysaccharide network essential for cytochrome anchoring and biofilm formation in

Geobacter sulfurreducens. J Bacteriol 193:1023–1033.

12.   Subramanian P, Pirbadian S, El-Naggar MY, Jensen GJ. 2018. Ultrastructure of

Shewanella oneidensis MR-1 nanowires revealed by electron cryotomography. Proc Natl

Acad Sci 201718810.

13.   Pirbadian S, Barchinger SE, Leung KM, Byun HS, Jangir Y, Bouhenni RA, Reed SB,

Romine MF, Saffarini DA, Shi L, Gorby YA, Golbeck JH, El-Naggar MY. 2014.

Shewanella oneidensis MR-1 nanowires are outer membrane and periplasmic extensions

of the extracellular electron transport components. Proc Natl Acad Sci U S A 111:12883–

8.

14.   Kim BC, Leang C, Ding YHR, Glaven RH, Coppi M V., Lovley DR. 2005. OmcF , a

Putative c -Type Monoheme Outer Membrane Cytochrome Required for the Expression of

Other Outer Membrane Cytochromes in Geobacter sulfurreducens. J Bacteriol 187:4505–

4513.

15.   Estevez-Canales M, Kuzume A, Borjas Z, Füeg M, Lovley D, Wandlowski T, Esteve-

Núñez A. 2015. A severe reduction in the cytochrome C content of Geobacter

438   sulfurreducens eliminates its capacity for extracellular electron transfer. Environ

439   Microbiol Rep 7:219–226.

440 16. Peng L, Zhang Y. 2017. Cytochrome OmcZ is essential for the current generation by

441   Geobacter sulfurreducens under low electrode potential. Electrochim Acta 228:447–452.

442 17. Jiménez Otero F, Chadwick GL, Yates MD, Mickol RL, Saunders SH, Glaven SM,

443   Gralnick JA, Newman DK, Tender LM, Orphan VJ, Bond DR. 2021. Evidence of a

444   Streamlined Extracellular Electron Transfer Pathway from Biofilm Structure, Metabolic

445   Stratification, and Long-Range Electron Transfer Parameters. Appl Environ Microbiol 87.

446 18. Joshi K, Chan CH, Bond DR. 2021. Geobacter sulfurreducens inner membrane

447   cytochrome CbcBA controls electron transfer and growth yield near the energetic limit of

448   respiration. Mol Microbiol 116:1124–1139.

449 19. Leang C, Adams LA, Chin KJ, Nevin KP, Methé BA, Webster J, Sharma ML, Lovley DR.

450   2005. Adaptation to disruption of the electron transfer pathway for Fe(III) reduction in

451   Geobacter sulfurreducens. J Bacteriol 187:5918–5926.

452 20. Lloyd JR, Leang C, Hodges Myerson AL, Coppi M V., Cuifo S, Methe B, Sandler SJ,

453   Lovley DR. 2003. Biochemical and genetic characterization of PpcA, a periplasmic c-type

454   cytochrome in Geobacter sulfurreducens. Biochem J 369:153–161.

455 21. Mehta T, Coppi M V., Childers SE, Lovley DR. 2005. Outer Membrane c-Type

456   Cytochromes Required for Fe(III) and Mn(IV) Oxide Reduction in Geobacter

457   sulfurreducens. Microbiology 71:8634–8641.

458 22. Shelobolina ES, Coppi M V., Korenevsky AA, DiDonato LN, Sullivan SA, Konishi H, Xu

459   H, Leang C, Butler JE, Kim BC, Lovley DR. 2007. Importance of c-type cytochromes for

460   U(VI) reduction by Geobacter sulfurreducens. BMC Microbiol 7:1–15.

23. Nevin KP, Kim BC, Glaven RH, Johnson JP, Woodward TL, Methé BA, Didonato RJ,

Covalla SF, Franks AE, Liu A, Lovley DR. 2009. Anode biofilm transcriptomics reveals

outer surface components essential for high density current production in Geobacter

sulfurreducens fuel cells. PLoS One 4.

24. Aklujkar M, Coppi M V., Leang C, Kim BC, Chavan MA, Perpetua LA, Giloteaux L, Liu

A, Holmes DE. 2013. Proteins involved in electron transfer to Fe(III) and Mn(IV) oxides

by Geobacter sulfurreducens and Geobacter uraniireducens. Microbiol (United Kingdom)

159:515–535.

25. Levar CE, Chan CH, Mehta-Kolte MG, Bond DR. 2014. An inner membrane cytochrome

required only for reduction of high redox potential extracellular electron acceptors. MBio

5:1–9.

26. Liu Y, Fredrickson JK, Zachara JM, Shi L. 2015. Direct involvement of ombB, omaB, and

omcB genes in extracellular reduction of Fe(III) by Geobacter sulfurreducens PCA. Front

Microbiol 6:1–8.

27. Mahadevan R, Bond DR, Butler JE, Coppi M V. 2006. Characterization of Metabolism in

the Fe (III) -Reducing Organism Geobacter sulfurreducens by Constraint-Based

Modeling. Appl Environ Microbiol 72:1558–1568.

28. Neidhart FC, Ingraham JL, Schaechter M. 1990. Physiology of the Bacterial Cell: A

Molecular Approach. Sinauer Associates, Sunderland, Mass.

29. Cuellar-Bermudez SP, Romero-Ogawa MA, Vannela R, Lai YJS, Rittmann BE, Parra-

Saldivar R. 2015. Effects of light intensity and carbon dioxide on lipids and fatty acids

produced by Synechocystis sp. PCC6803 during continuous flow. Algal Res 12:10–16.

30. Byreddy AR, Barrow CJ, Puri M. 2016. Bead milling for lipid recovery from

thraustochytrid cells and selective hydrolysis of Schizochytrium DT3 oil using lipase.

Bioresour Technol 200:464–469.

31. Budhraja R, Ding C, Walter P, Wagner S, Reemtsma T, Gary Sawers R, Adrian L. 2019.

The impact of species, respiration type, growth phase and genetic inventory on absolute

metal content of intact bacterial cells. Metallomics 11:925–935.

32. Snider RM, Strycharz-Glaven SM, Tsoi SD, Erickson JS, Tender LM. 2012. Long-range

electron transport in Geobacter sulfurreducens biofilms is redox gradient-driven. Proc Natl

Acad Sci 109:15467–15472.

33. Strycharz-Glaven SM, Snider RM, Guiseppi-Elie A, Tender LM. 2011. On the electrical

conductivity of microbial nanowires and biofilms. Energy Environ Sci 4:4366–4379.

34. Yates MD, Strycharz-Glaven SM, Golden JP, Roy J, Tsoi S, Erickson JS, El-Naggar MY,

Barton SC, Tender LM. 2016. Measuring conductivity of living geobacter sulfurreducens

biofilms. Nat Nanotechnol 11:910–913.

35. Brown RE, Jarvis KL, Hyland KJ. 1989. Protein measurement using bicinchoninic acid:

elimination of interfering substances. Anal Biochem 180:136–139.

36. DuBois M, Gilles KA, Hamilton JK, Rebers PA, Smith F. 1956. Colorimetric Method for

Determination of Sugars and Related Substances. Anal Chem 28:350–356.

37. Folch J, Lees M, Sloane Stanley GH. 1957. A simple method for the isolation and

purification of total lipides from animal tissues. J Biol Chem 226:497–509.

38. Sluiter A, Hames B, Ruiz R, Scarlata C, Sluiter J, Templeton D, Crocker D. 2008.

Determination of structural carbohydrates and lignin in Biomass. Natl Renew Energy Lab

17.

39. McCarty PL, Rittmann BE. 2001. Environmental biotechnology: principles and

applications. McGraw-Hill.

40. Howley E, Krajmalnik-brown R, Torres CI. 2022. Cytochrome expression shifts in

Geobacter sulfurreducens to maximize energy conservation in response to changes in

redox conditions 1–34.

41. Rouf MA. 1964. Spectrochemical Analysis of Inorganic Elements in Bacteria. J Bacteriol

88:1545–1549.

42. Kassner RJ, Kamen MD. 1968. Trace Metal Composition of Photosynthetic Bacteria.

Biochim Biophys Acta 153:270–278.

43. Lancaster WA, Menon AL, Scott I, Poole FL, Vaccaro BJ, Thorgersen MP, Geller J,

Hazen TC, Hurt RA, Brown SD, Elias DA, Adams MWW. 2014. Metallomics of two

microorganisms relevant to heavy metal bioremediation reveal fundamental differences in

metal assimilation and utilization. Metallomics 6:1004–1013.

44. Hartmann A, Braun V. 1981. Iron uptake and iron limited growth of Escherichia coli K-12.

Arch Microbiol 130:353–356.

45. Abdul-Tehrani H, Hudson AJ, Chang YS, Timms AR, Hawkins C, Williams JM, Harrison

PM, Guest JR, Andrews SC. 1999. Ferritin mutants of Escherichia coli are iron deficient

and growth impaired, and fur mutants are iron deficient. J Bacteriol 181:1415–1428.

46. Ma JF, Ochsner UA, Klotz MG, Nanayakkara VK, Howell ML, Johnson Z, Posey JE,

Vasil ML, Monaco JJ, Hassett DJ. 1999. Bacterioferritin A modulates catalase A (KatA)

activity and resistance to hydrogen peroxide in Pseudomonas aeruginosa. J Bacteriol

181:3730–3742.

47. Daly MJ, Gaidamakova EK, Matrosova VY, Vasilenko A, Zhai M, Venkateswaran A,

Hess M, Omelchenko M V., Kostandarithes HM, Makarova KS, Wackett LP, Fredrickson

JK, Ghosal D. 2004. Accumulation of Mn(II) in Deinococcus radiodurans Facilitates

Gamma-Radiation Resistance. Science (80-) 306:1025–1028.

48.   Pinchuk GE, Hill EA, Geydebrekht O V., de Ingeniis J, Zhang X, Osterman A, Scott JH,

Reed SB, Romine MF, Konopka AE, Beliaev AS, Fredrickson JK, Reed JL. 2010.

Constraint-based model of Shewanella oneidensis MR-1 metabolism: A tool for data

analysis and hypothesis generation. PLoS Comput Biol 6:1–8.

49.   Loferer-Kro☐ßbacher M, Klima J, Psenner R. 1998. Determination of Bacterial Cell Dry

Mass by Transmission Electron Microscopy and Densitometric Image Analysis. Appl

Environ Microbiol 64:688–694.

50.   Coppi M V. 2005. The hydrogenases of Geobacter sulfurreducens: A comparative

genomic perspective. Microbiology 151:1239–1254.

51.   Tremblay PL, Lovley DR. 2012. Role of the NiFe hydrogenase hya in oxidative stress

defense in Geobacter sulfurreducens. J Bacteriol 194:2248–2253.

52.   Yan J, Ritalahti KM, Wagner DD, Löffler FE. 2012. Unexpected specificity of

interspecies cobamide transfer from geobacter spp. to organohalide-respiring

dehalococcoides mccartyi strains. Appl Environ Microbiol 78:6630–6636.

53.   Dulay H, Tabares M, Kashefi K, Reguera G. 2020. Cobalt Resistance via Detoxification

and Mineralization in the Iron-Reducing Bacterium Geobacter sulfurreducens. Front

Microbiol 11:1–17.

54.   Holmes DE, Mester T, O'Neil RA, Perpetua LA, Larrahondo MJ, Glaven R, Sharma ML,

Ward JE, Nevin KP, Lovley DR. 2008. Genes for two multicopper proteins required for

Fe(III) oxide reduction in Geobacter sulfurreducens have different expression patterns

both in the subsurface and on energy-harvesting electrodes. Microbiology 154:1422–1435.

55.   Nanoparticles CS, Kimber RL, Bagshaw H, Smith K, Buchanan DM, Coker VS, Cavet JS,

Lloyd JR. 2020. Biomineralization of Cu2S Nanoparticles by Geobacter sulfurreducens.

Appl Environ Microbiol 86:1–11.

56.   Gong Y, Werth CJ, He Y, Su Y, Zhang Y, Zhou X. 2018. Intracellular versus extracellular

accumulation of Hexavalent chromium reduction products by Geobacter sulfurreducens

PCA. Environ Pollut 240:485–492.

57.   Otero FJ, Chadwick GL, Yates MD, Mickol RL, Saunders SH, Glaven SM, Gralnick JA,

Newman DK, Tender LM, Orphan VJ, Bond DR, Jiménez Otero F, Chadwick GL, Yates

MD, Mickol RL, Saunders SH, Glaven SM, Gralnick JA, Newman DK, Tender LM,

Orphan VJ, Bond DR. 2021. Evidence of a Streamlined Extracellular Electron Transfer

Pathway from Biofilm Structure, Metabolic Stratification, and Long-Range Electron

Transfer Parameters. Appl Environ Microbiol 87:1–16.

58.   Ma C, Wen H, Xing D, Pei X, Zhu J, Ren N, Liu B. 2017. Molasses wastewater treatment

and lipid production at low temperature conditions by a microalgal mutant Scenedesmus

sp. Z-4. Biotechnol Biofuels 10:1–13.

59.   Wahidin S, Idris A, Shaleh SRM. 2013. The influence of light intensity and photoperiod

on the growth and lipid content of microalgae Nannochloropsis sp. Bioresour Technol

129:7–11.

60.   Piligaev A V., Sorokina KN, Shashkov M V., Parmon VN. 2018. Screening and

comparative metabolic profiling of high lipid content microalgae strains for application in

wastewater treatment. Bioresour Technol 250:538–547.

61.   Caccavo F, Lonergan DJ, Lovley DR, Davis M, Stolz JF, McInerney MJ. 1994. Geobacter

sulfurreducens sp. nov., a Hydrogen- and Acetate-Oxidizing Dissimilatory Metal-

Reducing Microorganism. Appl Environ Microbiol 60:3752–3759.

62.   Stöckl M, Teubner NC, Holtmann D, Mangold KM, Sand W. 2019. Extracellular

Polymeric Substances from Geobacter sulfurreducens Biofilms in Microbial Fuel Cells.

ACS Appl Mater Interfaces 11:8961–8968.

63.   Davies SL, Whittenbury R. 1970. Fine structure of methane and other hydrocarbon-

utilizing bacteria. J Gen Microbiol 61:227–232.

64.   Murray RG, Watson SW. 1965. Structure of Nitrosocystis Oceanus and Comparison With

Nitrosomonas and Nitrobacter. J Bacteriol 89:1594–1609.

65.   Greening C, Lithgow T. 2020. Formation and function of bacterial organelles. Nat Rev

Microbiol 18:677–689.

66.   Kubannek F, Schröder U, Krewer U. 2018. Revealing metabolic storage processes in

electrode respiring bacteria by differential electrochemical mass spectrometry.

Bioelectrochemistry 121:160–168.

67.   Damoglou AP, Dawes EA. 1968. Studies on the lipid content and phosphate requirement

of glucose-and acetate-grown Escherichia coli. Biochem J 110:775–781.

68.   Lee S, Jeon E, Yun HS, Lee J. 2011. Improvement of fatty acid biosynthesis by

engineered recombinant Escherichia coli. Biotechnol Bioprocess Eng 16:706–713.