Acinetobacter baumannii represses type VI secretion system through an Mn^{2+}-dependent sRNA-mediated regulation

Somok Bhowmik[1], Avik Pathak[1], Kuldip Devnath[1], Abhiroop Sett[1], Shivam Pandey[1], Nishant Jyoti[1], Timsy Bhando[1,†], Jawed Akhter[1], Saurabh Chugh[2], Ramandeep Singh[2], Tarun Kumar Sharma[2,††] & Ranjana Pathania[1,3,*]

[1]Department of Biosciences and Bioengineering, Indian Institute of Technology Roorkee, Uttarakhand-247667, India

[2]Translational Health Science and Technology Institute, Faridabad, Haryana-121001, India

[3]Centre of Excellence in Disaster Mitigation and Management, Indian Institute of Technology Roorkee, Uttarakhand-247667, India

[†]Present address: Michael G. DeGroote Institute of Infectious Disease Research, McMaster University, Hamilton, Ontario, L8N, Canada

[††]Present address: Department of Medical Biotechnology, Gujarat Biotechnology University, Gujarat-382355, India

[*]Corresponding author. Tel: 01332-285324; E-mail: ranjana.pathania@bt.iitr.ac.in

ABSTRACT

Type VI secretion system (T6SS) is utilized by many Gram negative bacteria for

eliminating competing bacterial species and manipulating host cells. *Acinetobacter*

baumannii ATCC 17978 utilizes T6SS at the expense of losing pAB3 plasmid to induce

contact-dependent killing of competitor microbes, resulting in the loss of antibiotic

resistance carried by pAB3. However, the regulatory network associated with T6SS in *A.*

baumannii remains poorly understood. Here, we identified an Mn^{2+}-dependent post-

transcriptional regulation of T6SS mediated by a bonafide small RNA, AbsR28. *A.*

baumannii utilizes MumT (Mn^{2+}-uptake inner membrane transporter) for the uptake of

extracellular Mn^{2+} during oxidative stress. We demonstrate that the abundance of

intracellular Mn^{2+} enables complementary base-pairing of AbsR28-*tssM* mRNA (that

translated to TssM, one of the vital inner membrane components of T6SS), inducing RNase

E-mediated degradation of *tssM* mRNA and resulting in T6SS repression. Thus, AbsR28

mediates a crosstalk between MumT and T6SS in *A. baumannii*.

IMPORTANCE

Small RNAs (sRNAs) are identified as critical components within the bacterial regulatory

networks involved in fine regulation of virulence-associated factors. The sRNA-mediated

regulation of *Acinetobacter baumannii*'s T6SS was unchartered. Our findings reveal a

novel underlying mechanism of an Mn^{2+}-dependent sRNA-mediated regulation of T6SS in

A. baumannii. We show that binding of Mn^{2+} to AbsR28 aids in the complementary base-

pairing of AbsR28-*tssM* mRNA, resulting in RNase E-mediated processing of *tssM* and

T6SS repression. The findings also shed light on *A. baumannii*'s preference for antibiotic

resistance over contact-dependent killing during infection.

Keywords: Mn^{2+}; oxidative stress; post-transcriptional regulation; small RNA; T6SS

INTRODUCTION

Acinetobacter baumannii is a Gram-negative pathogen and has gained importance due to

its ability to cause wound and burn infections, sepsis, meningitis, urinary tract infections,

bloodstream infections, and ventilator-associated pneumonia (1-5). Treating hospital-

acquired *A. baumannii* infections is a major hassle, as most *A. baumannii* clinical strains

are multidrug-resistant (MDR) (6). Due to this severe global impact on public health, it is

essential to identify the molecular machinery adapted by *A. baumannii* to escape from host-

mediated immune response and establish pathogenesis.

One of the vital host-mediated immune responses encountered by pathogens is the

phagocytic cell-mediated free metal ion limitation, termed as "host-mediated nutritional

immunity" and generation of reactive oxygen species (ROS) at the site of infection (7, 8).

Mn^{2+} sequestration by neutrophils at the infection site reduces the activity of bacterial

metalloenzymes such as superoxide dismutase (SOD) and catalase, which bacteria use to

break down ROS (7, 9, 10). To acquire Mn^{2+} during oxidative stress and host-mediated

Mn^{2+} restriction, *A. baumannii* utilizes a high-affinity Mn^{2+} acquisition system MumT

(<u>m</u>anganese and <u>u</u>rea <u>m</u>etabolism <u>t</u>ransporter) (11, 12).

A synergy between the type VI secretion system (T6SS) and Mn^{2+}-transporter has been

observed in *Burkholderia thailandensis* (13). *B. thailandensis* secretes an effector protein

TseM through T6SS that scavenges Mn^{2+} from the extracellular milieu and delivers Mn^{2+}

through an outer membrane transporter (MnoT) to compensate for the scarcity of

intracellular Mn^{2+} under oxidative stress (13). T6SS is one of the secretion systems

frequently utilized by Gram-negative bacteria to promote contact-dependent killing of

competitors by injecting harmful toxins into it (14-17). Although Mn^{2+}-uptake systems

have been reported as one of the major virulence factors in pathogenic bacteria (18, 19),

crosstalk between Mn^{2+}-transporter and T6SS remains unknown in *A. baumannii*.

The biogenesis of T6SS assembly is an enormous and energetically expensive process

for bacteria (20). As a contact-dependent system with specific cellular targets (21), the

expression of T6SS might be precisely regulated transcriptionally and post-

transcriptionally (22, 21). Transcriptional repression of T6SS by *tetR*-like regulators

carrying by pAB3 plasmid or its derivative is observed in *A. baumannii* (23), whereas post-

transcriptional regulation of T6SS has not been explored yet. Small regulatory RNAs

(sRNAs) are identified as critical components within the bacterial regulatory networks

involved in post-transcriptional regulation of virulence-associated factors. The sRNAs are

mostly 50-500 nucleotides long, function as global regulators of numerous bacterial

physiological processes, and play a crucial role in regulating several virulence factors (24,

25). The sRNAs base pair with their cognate mRNA and regulate the translational activity

and/or the stability of that particular mRNA often with the assistance of the RNA

chaperone, Hfq (26). The sRNA-mediated regulation of T6SS in *A. baumannii* remains

unexplored.

In this work, we show that the breakdown in Mn^{2+}-uptake by deleting *mumT* increases

T6SS expression during oxidative stress. Moreover, the elevated level of intracellular Mn^{2+}

by overexpressing MumT reduces T6SS expression. These observations indicate the

existence of an Mn^{2+}-dependent T6SS regulation in *A. baumannii*. We explored the

crosstalk between MumT and T6SS with the aim of understanding the underlying

mechanism of this crosstalk. Intriguingly, we identify an sRNA, AbsR28 (<u>*A*</u>. <u>*baumannii*</u>

small RNA 28), that mediates the crosstalk between MumT and T6SS. Collectively, we

show that the uptake of Mn^{2+} by MumT is necessary for AbsR28-mediated post-

transcriptional regulation of *tssM* mRNA, leading to T6SS repression in *A. baumannii*.

RESULTS

***A. baumannii* cells that switch to T6+ are unable to withstand oxidative stress**

The T6SS in bacteria is often found to be essential for bacterial pathogenesis as it

modulates the host immune response during infection (27-30). In contrast, certain T6SSs

seem to function as antivirulence mechanisms, as evidenced by increased pathogenicity in

T6SS deficient mutants relative to wild-type bacteria (31, 32). Such involvement of *A.*

baumannii's T6SS during infection remains enigmatic. Oxidative burst by phagocytic cells

serves as a primary mechanism for microbial eradication in the host during infection. To

assess whether *A. baumannii* cells expressing T6SS (T6+) would have an advantage in

survival during phagocytic cell-mediated killing compared to *A. baumannii* cells with silent

T6SS (T6-), we co-incubated phagocytic cells either with wild-type (WT) T6- or WT cells

that switched to T6+ cells and checked their survival. The WT T6- and WT cells that

switched to T6+ cells were isolated from *A. baumannii* ATCC 17978 based on the secretion

of T6SS-associated protein Hcp (hemolysin-coregulated protein, a positive marker for

expression of T6SS) using several rounds of Hcp-ELISA followed by western blot to detect

secreted Hcp. The cells that secreted Hcp were considered as T6+ (WT T6+ cells after

losing pAB3 plasmid which carries *tetR*-repressors for T6SS are referred as WT T6+

throughout the study) and the cells that showed no detectable Hcp in the cell-free

supernatant were considered as T6- (WT T6- cells carrying pAB3 plasmid which carries

tetR-repressors for T6SS are referred as WT T6- throughout the study) (Fig S1A). To our

surprise, WT T6- cells exhibited ~60-70% survival, whereas WT T6+ cells exhibited only

~20-30% survival against phagocytic cell-mediated killing compared to untreated controls

(Fig 1A and S1B). This observation suggests that WT T6+ cells cannot withstand

phagocytic cell-mediated killing. As neutrophils induce oxidative stress by producing

deleterious superoxide radicals during infection, the WT T6- and WT T6+ cells were grown

in an LB medium supplemented with methyl viologen (MV) to induce superoxide radicals

(33-35), and bacterial growth was measured at the indicated time points. The strains

displayed almost equal growth in LB, however, upon addition of MV, WT T6+ cells

exhibited a remarkable growth defect (Fig 1B). This reinforced our previous observation

that WT T6+ cells display enhanced sensitivity to oxidative stress compared to WT T6-

cells. Interestingly, There are two mixed variants of *A. baumannii* ATCC 17978, one

variant has a 44 kb island in the genome (referred as +AbaAL44), and the other one is

devoid of that 44 kb island (referred as -AbaAL44) (36). It was demonstrated that the

presence of *katX* in AbaAL44 island confers resistance to *A. baumannii* strain carrying

AbaAL44 island against H_2O_2-mediated oxidative stress (36). It was surprising that WT

T6+ strain is sensitive to oxidative stress despite possessing AbaAL44 island, while WT

T6- strain exhibits more resistance to oxidative stress despite lacking the AbaAL44 island

suggesting that the growth defect is not dependent on AbaAL44 status (Fig S1C).

Intracellular ROS generation via Fenton chemistry is a hallmark of oxidative stress

response (37, 38). Thus, we measured intracellular ROS level in WT T6- and WT T6+ cells

by growing them in either LB broth or LB broth supplemented with MV by measuring the

fluorescence of 2',7'-dichlorofluorescein (DCF). A significant increase in ROS level was

observed in WT T6+ cells compared to WT T6- cells treated with MV (Fig 1C), suggesting

that WT T6+ cells are more sensitive to oxidative stress due to higher intracellular ROS

accumulation.

***A. baumannii* T6+ cells cannot withstand oxidative stress due to inadequate**

intracellular Mn^{2+}

Bacteria utilize metal ion-dependent SODs (Mn/Fe SOD and Cu/Zn SOD) and catalases

for breaking down intracellular ROS. Therefore, metal ion homeostasis governs the cellular

ROS level. *A. baumannii* employs specialized metal uptake systems to import metal ions

for the activity of these metalloenzymes during oxidative stress and to survive against host-

mediated metal limitation (12, 39-41). Due to the compromised survival of WT T6+ cells

in phagocytic cell-mediated killing, we hypothesized that WT T6+ cells may be defective

in the uptake of free metal ions, which serve as cofactors for SODs and catalases required

to break down ROS. To test this, we co-incubated neutrophils with either WT T6- or WT

T6+ cells and checked the expression of *mumT*, *znuB*, and *tonB* genes involved in Mn^{2+}-

uptake, Zn^{2+}-uptake, and Fe$^{2+/3+}$-uptake, respectively, by quantitative reverse transcription

PCR (qRT-PCR). There was no significant difference in the expression of *znuB* and *tonB*

genes, but in the case of *mumT*, a ~3.5-log$_2$ fold reduction was observed in WT T6+ cells

as compared to WT T6- cells (Fig 1D). Similarly, there was a significant fold reduction of

mumT transcript levels in WT T6+ cells compared to WT T6- cells during oxidative stress

when the bacterial cells were grown LB-medium supplemented with MV (Fig S1D). Due

to the fold reduction of *mumT* expression in WT T6+ cells, we hypothesized that this could

lead to intracellular Mn^{2+} deficiency in WT T6+ cells under oxidative stress. To examine

this, we measured the concentration of intracellular Mn^{2+} using inductively coupled plasma

mass spectrometry (ICP-MS). To determine the intracellular concentration of a specific

metal ion, minimal media was preferred over complex media in this assay. The WT T6+

cells had ~3-fold lower Mn^{2+} levels in cell pellets than WT T6- cells in the presence of

$MnCl_2$+MV (Fig 1E). In contrast, there were no significant differences in Zn^{2+} and $Fe^{2+/3+}$

levels in cell pellets of WT T6+ cells as compared to the WT T6- cells in the presence of

$ZnSO_4$+MV and $FeCl_3$+MV, respectively (Fig 1F). Next, we hypothesized that the addition

of sufficient $MnCl_2$ might alleviate the oxidative stress toxicity in WT T6+ cells. The WT

T6- strain grew almost equally in all conditions; supplementation of $ZnSO_4$ and $FeCl_3$

reverted the growth of WT T6+ cells to some extent, but supplementation of $MnCl_2$ did not

have much effect under oxidative stress (Fig S1E). Taken together, these data suggest that

WT T6+ cells display impaired growth under oxidative stress due to inadequate

intracellular Mn^{2+}, which is crucial for *A. baumannii*'s SOD and catalase activity, and this

defect in activity could not be restored even after supplementation of sufficient Mn^{2+} (Fig

S1F, G). We isolated T6+ cells of two clinical strains of *Acinetobacter baumannii* (RPTC2

and RPTC3) and compared the growth of WT and T6+ cells of these isolates under

oxidative stress. We observed that the T6+ strains of RPTC2 and RPTC3 showed a growth

defect under oxidative stress compared to their respective WT strains (Fig S2A-C). Like

A. baumannii ATCC 17978, RPTC2 and RPTC3 T6+ strains are more susceptible to

neutrophil-mediated killing (Fig S2D).

pAB3 plasmid has no role in mitigating oxidative stress

In view of the fact that WT T6- cells of *A. baumannii* ATCC 17978 switch to WT T6+

cells upon losing pAB3 plasmid (23) and the WT T6+ cells are sensitive to oxidative stress

due to inadequate intracellular Mn^{2+}, we speculated that the sensitivity might be due to the

cells losing pAB3 plasmid which encodes several transcription factor regulators. To

investigate this, we transformed pAB3 into the WT T6+ competent cells (which were

devoid of pAB3), confirmed the transformants (Fig S3A, B), and assessed the growth of

WT T6-, WT T6+, and WT T6+ cells transformed with pAB3 (referred as WT T6+/pAB3

throughout the study) strains in LB supplemented with MV. We observed that the presence

of pAB3 does not confer resistance to oxidative stress as there is no growth difference

between WT T6+ and WT T6+/pAB3 cells (Fig S3C). Next, we sought to determine

whether *mumT* expression is reduced by T6SS functionality, rendering the T6+ cells

defective in Mn^{2+} uptake (scenario 1), or if the opposite is true and a breakdown in Mn^{2+}

uptake results in the T6+ phenotype (scenario 2) (Fig S3D). Consistent with the previous

observation, we also observed that T6SS expression is repressed in WT T6+ strain upon

transformation of pAB3 which contains *tetR*-repressors (Fig S3E) whereas the

downregulation of *mumT* is unaffected in comparison to WT T6+ cells by the

transformation of pAB3 into the WT T6+ cells (WT T6+/pAB3) (Fig S3F). This led us to

infer that alteration of T6SS status does not impact *mumT* expression and ruling out the

possibility of scenario 1.

Breakdown in Mn^{2+}-uptake results in an increase in T6SS expression

Mn^{2+} is an essential micronutrient for bacterial physiology, virulence, and survival against

oxidative stress (18). *A. baumannii* utilizes an inner membrane transporter, MumT, for

Mn^{2+} uptake during host-mediated metal restriction (12). To assess whether intracellular

Mn^{2+} has any impact on T6SS regulation (scenario 2; Fig S3D), we first checked the effect

of *mumT* on T6SS expression. To evaluate this, we created a deletion mutant of *mumT* in

A. baumannii containing pAB3 plasmid (referred as Δ*mumT* throughout the study), grew

both the WT T6- and Δ*mumT* cells under oxidative stress supplemented with $MnCl_2$, and

checked the expression of *hcp* by qRT-PCR. Hcp is a structural component of T6SS (Fig

2A). Interestingly, we observed a ~6-$\log_2$ fold increase in *hcp* expression in Δ*mumT* cells

with respect to WT T6- cells (Fig 2B). The data were further validated by Hcp-Western

blot, where Hcp expression was highly induced in Δ*mumT* cells compared to WT T6- cells

in both cell lysate (CL) and culture supernatant (S) (Fig 2C). We used a prey-predator assay

to assess the T6SS-mediated killing of prey cells by WT T6- and Δ*mumT* cells, and the

killing of prey cells confirmed the expression of T6SS. As expected, Δ*mumT* cells

exhibited an efficient increase in the killing of prey cells compared to WT T6- cells (Fig

2D, S4A). We overexpressed *mumT* into WT T6+ and Δ*mumT* strains. We observed that

the elevated intracellular Mn^{2+} (Fig S4B) due to an increase in MumT expression (Fig S4C)

remarkably reduced Hcp secretion (Fig S4D) and T6SS activity compared to their

respective control (Fig S4F). The percentage of Hcp-secretion positive cells increased to

~50% of the entire population in Δ*mumT* strain, while only ~5% of WT T6- cells were

positive under oxidative stress supplemented with $MnCl_2$ (Fig 2E and S4G). Taken

together, these data suggest that deletion of MumT results in an increase in T6SS

expression and thus MumT seems to have a negative impact on T6SS regulation in *A.*

baumannii under oxidative stress (scenario 2).

AbsR28 mediates the crosstalk between *mumT* and T6SS

We wondered how a metal transporter could impact the bacterial secretion system. Our

observation that Δ*mumT* cells are predominant in T6+ despite having *tetR1* and *tetR2* (Fig

S5A), which are transcriptional repressors of T6SS, suggests the presence of a MumT-

dependent regulation of T6SS in *A. baumannii*. As the deletion of *mumT* in *A. baumannii*

mimics an Mn^{2+}-depleted stress condition (11), we speculated that the deletion of *mumT*

might disrupt some transcriptional or post-transcriptional regulation of T6SS that causes

T6SS upregulation in the Δ*mumT* cells. To begin with this hypothesis, we first checked the

expression of several transcriptional regulators (i.e., *oxyR*, *zur*, and *fur*), which are known

to regulate T6SS in other bacteria (13, 42, 43) at the transcriptional level in WT T6-, WT

T6+, and Δ*mumT* cells grown under oxidative stress supplemented with $MnCl_2$ using qRT-

PCR. We also checked the transcript level of *mumR*, a transcriptional regulator of *mumT*

in *A. baumannii* (12). None of the tested transcriptional regulators displayed any significant

changes in expression (Fig S5B). sRNAs are known to play a vital role in post-

transcriptional gene regulation in bacteria. The sRNA-mediated post-transcriptional

regulation of T6SS is observed in *P. aeruginosa* (44-46). In light of this, we performed a

CopraRNA search for all thirty-one putative sRNAs in *A. baumannii* that were predicted

earlier using bioinformatic analysis by our lab (47), and eighteen of these sRNAs showed

T6SS as a putative target (Table S1). To test the hypothesis further, we first picked four

sRNAs (AbsR1, AbsR11, AbsR25, and AbsR28) out of those eighteen whose presence was

already confirmed in *A. baumannii* and whose sequences were RACE mapped in our earlier

studies (47). We checked the expression of four validated sRNAs (i.e., AbsR1, AbsR11,

AbsR25, and AbsR28), and AbsR29 (which was not validated earlier to characterize it for

its function), and *hfq* at the transcriptional level in WT T6-, WT T6+, and Δ*mumT* cells

grown under oxidative stress supplemented with $MnCl_2$ using qRT-PCR. Intriguingly, the

expression of one sRNA, AbsR28, showed a notable fold reduction in the WT T6+ and

Δ*mumT* cells compared with WT T6- cells (Fig 3A and S5C). Due to the maximum fold

increase at the transcription level in the presence of $MnCl_2$ under oxidative stress (Fig S5D)

and significant fold reduction in WT T6+ cells and Δ*mumT* cells which are defective in

Mn^{2+} utilization, we focused our study on evaluating the role of AbsR28 in *A. baumannii*'s

T6SS regulation. We created a deletion mutant of AbsR28 in *A. baumannii* containing

pAB3 plasmid (referred as ΔAbsR28 throughout the study) and confirmed that the deletion

had no polar effect on transcript abundance (Fig S5E, F). To identify the genes regulated

by AbsR28, we performed RNA-seq analysis comparing the relative abundance of total

mRNA transcripts of WT T6- and ΔAbsR28 cells (upon exposure to oxidative stress and

supplemented with $MnCl_2$). In a comparison of WT T6- and ΔAbsR28 strains, the

expression of seven structural genes of the T6SS [*tslA* (A1S_1292); *hcp* (A1S_1296); *tssF1*

(A1S_1298); *tssF2* (A1S_1299); *tssG* (A1S_1300); *tsmK* (1301) and *tsgF* (A1S_1304)]

and six genes encoding VgrGs/T6SS effector molecules (A1S_0086, A1S_0550,

A1S_0551, A1S_1290, and A1S_3363) displayed a significant fold-induction in ΔAbsR28

strain (Fig 3B). All the genes encoding T6SS displayed a significant fold-increase in

ΔAbsR28 cells compared to WT T6- cells grown under oxidative stress supplemented with

$MnCl_2$, indicating that AbsR28 negatively regulates T6SS (Fig. 3C). Like Δ*mumT* cells,

ΔAbsR28 cells are also predominant in T6+ despite having *tetR1* and *tetR2* (Fig S5G).

Next, we investigated whether AbsR28 regulates T6SS in *A. baumannii* and if there is a

role of Mn^{2+} in this regulation. To evaluate this, we cloned AbsR28 under the control of an

arabinose inducible promoter in the pWBAD30 vector and complemented the ΔAbsR28

strain. We observed that the elevated level of AbsR28 due to arabinose pulse, decreased

the expression of Hcp, and the expression was further reduced when grown in the presence

of $MnCl_2$ in ΔAbsR28-pWBAD30AbsR28 strain (Fig 3D). The data was further validated

by Hcp-western blot to detect secreted Hcp and a prey-predator assay, where increased

expression of AbsR28 resulted in a notable reduction in Hcp secretion and increase in prey

cell survival, respectively, (Fig 3E, F). These data indicate that AbsR28 efficiently

represses T6SS expression in *A. baumannii* under oxidative stress in the presence of Mn^{2+}.

Binding of Mn^{2+} to AbsR28 alters its native structure

An sRNA's ability to target a certain mRNA is contingent on its ability to form a base-

pairing interaction with that mRNA at a specific location, known as the seed region (48).

In AbsR28, the region that forms the most frequent base pairs with the predicted targets is

positioned ~1-50 nt from the start site, making it inaccessible for interaction due to the

stem formation (Fig. 4A). Due to this, we hypothesized that Mn^{2+} might bind to AbsR28

and alter its native structure, which might be essential for base pairing and its regulatory

function. The dissociation constant (K_D) between Mn^{2+} and AbsR28 was 5.95×10^{-6} ±

1.21×10^{-6} M, as measured by isothermal titration calorimetry (ITC) (Fig 4B). The affinity

to Mn^{2+} is more efficient than the controls where the binding of another small RNA (i.e.,

AbsR25) with Mn^{2+} and AbsR28 with Mg^{2+} were assessed (Fig S6A, B). Mutations in

AbsR28 (U_{89}, $A_{91}U_{94}A_{95}$, and $U_{84}U_{85}$) reduced its affinity to Mn^{2+} except one mutant

($A_{76}A_{77}$) (Fig S6C). To determine whether the binding of Mn^{2+} affects the native structure

of AbsR28, we performed *in vitro* structural probing with 5'-end labeled $[\gamma\text{-}^{32}P]ATP$

AbsR28 using lead(II) acetate (PbAc). The Mn^{2+} concentration-dependent changes in the

cleavage of AbsR28 were observed in a denaturing-PAGE (Fig 4C), which were not very

evident in the presence of Mg^{2+}, used as a control (Fig 4D). These data confirm that Mn^{2+}

binds to AbsR28 and alters its native structure.

Mn^{2+} is required for AbsR28 to interact with *tssM* mRNA

sRNAs regulate gene expression by directly base pairing with target mRNA (49). Since

T6SS in *A. baumannii* is a large gene cluster (50), we used the CopraRNA bioinformatic

tool (51) to predict the targets of AbsR28 in *A. baumannii*. A CopraRNA search using the

whole-genome sequence of *A. baumannii* ATCC 17978 and full-length AbsR28 (RACE

mapped sequence) as inputs predicted base pairing between *tssM* mRNA position 258-291

and position 27-55 of AbsR28 with a predicted hybridization energy value of -14.10

kcal/mol (Fig 5A). To validate the direct interaction between AbsR28 and *tssM* mRNA, an

in vitro RNA-RNA interaction was assessed. The addition of full-length AbsR28 at

increasing concentrations to *tssM in vitro* transcripts resulted in the retardation of *tssM*

transcripts in a native PAGE in the presence of Mn^{2+} (Fig 5B); in contrast, the retardation

of *tssM* transcripts was very weak in the presence of Mg^{2+}, used as a control (Fig 5C and

5D). The addition of full-length AbsR28 at increasing concentrations to an unrelated

mRNA *pntB in vitro* transcripts (used as a negative control) did not show any retardation

(Fig. S7A). For further validation of AbsR28-*tssM* binding, AbsR28 was mutated at the

seed region (AbsR28-M9; $A_{27}G_{28}G_{29}A_{30}$ was mutated to $U_{27}C_{28}C_{29}U_{30}$) (Fig S6B). The

retardation of *tssM* transcripts was much weaker even with higher concentrations of

AbsR28-M9 compared to native AbsR28 (Fig S7C and S7D). Furthermore, we show that

the compensatory mutation in *tssM* mRNA (*tssM**) restored the binding of AbsR28-M9

(S7E). Collectively, these findings indicate that the direct interaction of AbsR28 and *tssM*

transcripts is Mn^{2+}-dependent.

AbsR28 potentiates RNase E-mediated *tssM* mRNA degradation

To study the molecular mechanism of AbsR28-meditated T6SS repression in *A.*

baumannii, we checked the effect of AbsR28-*tssM* interaction on the stability of target

mRNA (i.e., *tssM*) via *in vivo* pulse expression. AbsR28 markedly reduced the stability of

tssM mRNA *in vivo* in the presence of Mn^{2+} (Fig 5E). We observed that the elevated level

of native AbsR28 reduced the level of *tssM* transcripts more significantly compared to the

AbsR28-M9 via *in vivo* pulse expression assay (Fig S7F). This result confirmed that *tssM*

processing depends on the direct interaction between AbsR28 and *tssM*. sRNAs induce the

decay of target mRNA by RNase E-mediated mRNA degradation in an Hfq-dependent

manner and repress its translation (52-58). We hypothesized that this might also happen in

the case of AbsR28. To examine this, we performed an *in vitro* RNase E-mediated

degradation assay where the 5'-end of the *in vitro* transcribed *tssM* was labeled with [γ-

^{32}P]ATP and incubated with purified RNase E in the presence or absence of unlabeled

AbsR28 with *A. baumannii* Hfq72 or *E. coli* Hfq protein. RNase E alone cleaves *tssM*

transcripts, but the cleavage is more prominent in the presence of AbsR28. As expected,

there is a distinct RNase E-mediated degradation of *tssM* transcripts in the presence of

AbsR28 and Hfq (Fig. 5F). These data demonstrate the molecular mechanism behind

AbsR28-mediated T6SS regulation where the degradation of *tssM* mRNA by RNase E is

strongly enhanced by base pairing to AbsR28 in the presence of Mn^{2+} and reduces the

abundance of *tssM* transcripts for T6SS assembly.

AbsR28 contributes to the fitness of *A. baumannii* during infection

In addition to T6SS regulation, we observed that the deletion of AbsR28 primarily affects

genes involved in bacterial metabolism (carbohydrate, phenylalanine, cysteine, and

methionine metabolism) and stress response (two-component system and ABC

transporters) in RNAseq data (Table S2). Since AbsR28 remains uncharacterized in *A.*

baumannii pathophysiology; therefore, we aimed to explore the importance of AbsR28-

mediated gene regulation in the pathogenesis of *A. baumannii*. We co-incubated human

blood-derived neutrophils with either WT T6-, WT T6+, Δ*mumT*, or ΔAbsR28 cells and

checked their survival. WT T6- cells exhibited ~85% survival, whereas ΔAbsR28 cells

showed only ~40% survival against neutrophil-mediated killing compared to untreated

cells (Fig 6A). We next assessed the pathogenesis of WT T6-, WT T6+, ΔmumT, and

ΔAbsR28 strains in a mouse model of *A. baumannii*-induced pneumonia where the strains

were intranasally administered to BALB/c mice. In consistency with the previous

observation that the deletion of *mumT* in *A. baumannii* results in compromised survival in

mice due to a defect in Mn^{2+}-uptake (12), we also observed that Δ*mumT* strain had a lower

organ burden compared to the WT T6- strain. Additionally, WT T6+ and ΔAbsR28 strains

had significantly reduced burdens in the lungs and liver when compared with the WT T6-

strain (Fig 6B). Lungs isolated from mice infected with the WT T6- strain exhibited a

remarkable tissue infiltration as indicated by necrosis and alveolar inflammation, whereas

the mice lungs infected with either WT T6+, Δ*mumT*, or ΔAbsR28 strains did not show

any notable damage (Fig 6C). Together, the data suggest that AbsR28 contributes to *A.*

baumannii fitness during infection.

DISCUSSION

During bacterial infection, the host recruits phagocytic cells at the site of infection, which

creates oxidative stress and limits the availability of free metal ions (12, 59, 60). For certain

bacteria, the T6SS sequesters extracellular free metal ions and helps them cope with

oxidative stress and host-mediated free metal restriction, so it is advantageous for them to

express T6SS during oxidative stress (13, 44, 61). However, such role of T6SS in *A.*

baumannii's ability to survive under phagocytic cell-mediated oxidative stress has not yet

been explored. Unlike *B. thailandensis*, where a synergistic relationship exists between the

Mn^{2+}-uptake system and T6SS expression (13), we found that *A. baumannii* cells that

switch to T6+ cells are sensitive to oxidative stress due to inadequate intracellular Mn^{2+},

which is required as a cofactor for SOD and catalase to break down intracellular ROS

generated during oxidative stress (Fig 1 and S1). This distinguishes *A. baumannii* from

other pathogens with distinctive gene regulatory mechanisms and emphasizes the need for

studying *A. baumannii* pathophysiology in depth. To check the impact of intracellular Mn^{2+}

on T6SS expression, we observed that the deletion of *mumT* promotes a significant increase

in T6SS expression under oxidative stress (Fig 2). The T6SS expression is reduced due to

adequate intracellular Mn^{2+} in the *mumT*-overexpressed strains (Fig S4), indicating the

existence of a *mumT*-dependent T6SS regulation. Next, we focused on determining the

molecular mechanism behind this regulation.

As deletion of *mumT* resulted in an increase in T6SS expression in *A. baumannii*

(Fig 2) and there were no significant changes in the expression of known transcriptional

regulators, we hypothesized that this might be due to Mn^{2+}-dependent post-transcriptional

repression, which is affected by the deletion of *mumT*. We observed that AbsR28 mediates

the crosstalk between MumT and T6SS in *A. baumannii* during oxidative stress (Fig 3).

Next, we focused on elucidating the mechanistic details of AbsR28-mediated T6SS

repression. We show that the binding of Mn^{2+} to AbsR28 alters its native structure and

results in a complementary base pairing with *tssM* transcripts (Fig. 4B, 4C and 5B). The

exact Mn^{2+}-binding region of AbsR28 remains to be determined. TssM is an indispensable

structural component of the T6SS membrane complex which is essential for T6SS

assembly (62-65). Hence, the deletion of *tssM* disrupts the functionality of T6SS in *A.*

baumannii (66). We observed that the stability of *tssM* transcripts was significantly reduced

when AbsR28 was pulse expressed in the presence of Mn^{2+} (Fig 5E). In

Gammaproteobacteria, RNase E is the primary catalyst of mRNA decay (52). Since we

observed AbsR28 expression-dependent decay of *tssM* transcripts *in vivo*, we hypothesized

that AbsR28 triggers RNase E-mediated processing of *tssM* transcripts. Our in-line probing

data demonstrate that AbsR28 potentiates RNase E-mediated degradation of *tssM*

transcripts with the assistance of RNA chaperone Hfq (Fig 5F).

The T6SS is often not expressed in most of the clinical isolates of *A. baumannii*

(67-69). The reason behind the inactive T6SS in *A. baumannii* is not fully understood. The

following factors may account for potentially silent T6SS in *A. baumannii* clinical isolates:

(i) high energy is required for T6SS expression, so to utilize the energy for other cellular

processes, T6SS is silent; (ii) T6SS components are immunogenic (70, 71), so it may be

advantageous for *A. baumannii* T6SS inactive strains to evade the host-mediated immune

response (72); (iii) *A. baumannii* cells switch from T6- to T6+ at the expense of losing

antibiotic resistance encoded by the plasmid (23); (iv) the T6+ cells of *A. baumannii*

urinary tract infection (UTI) isolate have been previously reported to exhibit decreased

virulence due to loss of pAB5 (a pAB3-family member), which regulates the expression of

several chromosomally-encoded virulence factors (73); (v) firing the T6SS arrow involves

bacteria rupturing the outer membrane, which could potentially disrupt the membrane

integrity and render *A. baumannii* T6+ cells more sensitive to oxidative stress. Thus, to

maintain the advantage of antibiotic resistance and virulence factors over the contact-

dependent killing mechanism, it might be preferable for *A. baumannii* cells to inactivate

the T6SS by other mechanisms independent of plasmid-mediated repression. A

comparative study on the *A. baumannii* clinical strains and lab strains about the regulatory

machinery of T6SS regulation during oxidative stress can be conducted in future for a
better understanding of T6SS regulation.

In summary, we explored the molecular mechanism of AbsR28-mediated T6SS
repression in *A. baumannii* ATCC 17978 (Fig 7). The sequential events of this proposed
pathway are as follows: (i) during oxidative stress, *A. baumannii* cells upregulate *mumT* to
increase Mn^{2+} uptake; (ii) intracellular Mn^{2+} binds to AbsR28 and alters its native structure;
(iii) the altered structure of AbsR28 aids in the complementary base pairing with *tssM*
mRNA transcripts and triggers RNase E processing; (iv) degradation by RNase E reduces
the abundance of *tssM* transcripts and thereby represses the T6SS expression. *A. baumannii*
cells that lose this controlled regulation result in T6+ phenotype. Our results unravel a
novel mechanism of an Mn^{2+}-dependent sRNA-mediated T6SS regulation in *A. baumannii*.
Collectively, the findings presented here widen our knowledge about sRNA-mediated gene
regulation and its role in shaping *A. baumannii*'s pathophysiology.

MATERIALS AND METHODS

Bacterial strains and growth conditions

Bacterial strains and plasmids used in this study are listed in Tables S3 and S4. *A. baumannii* strains were grown at 37 ^{0}C in LB broth or LB agar. *E. coli* J53 strains were grown in LB broth or LB agar, supplemented with 100 µg/mL sodium azide when necessary. *E. coli*-pNYL GFP strains were grown in LB broth or LB agar, supplemented with 50 µg/mL kanamycin when necessary. *P. aeruginosa* PAO1 strains were grown in LB broth or LB agar. *A. baumannii* strains that contain the pWBAD30 vector were grown in LB broth or LB agar, supplemented with 50 µg/mL kanamycin.

Estimation of cell survival from phagocytosis

All experiments using human blood-derived neutrophils under protocol BT/IHEC-612020/7865 were reviewed and approved by the Institute Human Ethics Committee (HEC) of the Indian Institute of Technology Roorkee. The estimation of bacterial survival from phagocytic cell-mediated killing was performed as described previously (74). The detailed procedure is provided in the Supplemental material.

Growth assay under oxidative stress

From the mid-log phase cultures of the indicated strains, 0.1% inoculum was inoculated in a 200 µL of fresh LB medium containing methyl viologen (MV) at 250 µM (final conc.). All the growth assays were performed in a sterile 96-well plate (Genaxy) at 37 ^{0}C with shaking linearly at 180 CPM (6 mm), and the OD_{600} as a measurement of growth was measured at every 30 min interval for the indicated total time in the Synergy microplate reader (BioTek). Only media without any culture served as a negative control for this assay. The represented data is after background correction.

ROS quantification

Bacterial cells were grown at 37 ^{0}C with shaking to an OD$_{600}$ of 0.6 (mid-log phase) in LB-medium. Bacterial cells were harvested by centrifugation and washed in sterile 1X PBS. The bacterial cell pellets were resuspended in 1X PBS, and 2',7'-dichlorofluorescein diacetate (Thermo Fisher Scientific, D399) was added at a final concentration of 100 μM. After incubation for 30 min at 37 ^{0}C, the cells were washed with 1X PBS to remove excess dye and transferred to 100 μL/well of a 96-well transparent bottom black well plate (BRAND). MV (250 μM final conc.) was added to the wells containing bacterial cells and incubated at 37 ^{0}C with shaking linearly at 180 CPM (6 mm). OD$_{600}$ and fluorescence (excitation/emission at 485/535 nm) were measured every 10 min interval for the indicated total time in the Synergy microplate reader (BioTek). The represented data is after background correction and OD$_{600}$ normalization.

Quantitative RT-PCR analysis

Human blood-derived neutrophils were co-incubated with the bacterial strains as described in the above section. After 4 h of incubation, the tissue culture plate was centrifuged at 400g for 5 min to settle down the neutrophils. Sample supernatants containing the bacterial cells were collected and harvested the bacterial cells by centrifugation at maximum speed. After washing the cell pellet with 1X PBS, RNA was extracted from the bacterial cells by the classic phenol-chloroform method. The cDNA synthesis was performed using PrimeScript 1st strand cDNA Synthesis Kit (TakaRa, 28704) according to the manufacturer's instructions. Amplifications were achieved using a 3-step program on a QuantStudio 5 system (Thermo Fisher Scientific). Transcript abundance was calculated using the $\Delta\Delta C_T$ method (75) and normalized by the 16s gene.

Quantification of intracellular metal content

The concentration of metal content in bacterial cells was determined as described previously (12) with some modifications. The detailed procedure is provided in the Supplemental material.

Bacterial survival assay upon uptake of metal ions under oxidative stress

Bacterial cells were grown in 5 mL M9-medium (supplemented with 1% casamino acids) with or without $MnCl_2$, $ZnSO_4$, or $FeCl_3$ at a final concentration of 250 μM and grown at 37 ^{0}C with shaking to an OD_{600} of 0.6 (mid-log phase). The cultures were then treated with MV at a final concentration of 250 μM to induce oxidative stress and grow for another 2 h at 37 ^{0}C with shaking. Cells were harvested from a 1 mL culture, washed with 1X PBS, and dissolved into 50 μL of 1X PBS. For spot assay, after a serial dilution in 1X PBS, 5 μL from each dilution was spotted onto LB agar containing MV at a final concentration of 100 μM, incubated at 37 ^{0}C for O/N, and images were taken by a gel documentation system (Bio-Rad).

Western blot analysis for Hcp

From the mid-log phase cultures of the indicated strains, 0.1% inoculum from the O/N cultures was then subcultured into fresh 5 mL of LB-medium containing $MnCl_2$ at a final concentration of 250 μM and grown at 37 ^{0}C with shaking to an OD_{600} of 0.6 (mid-log phase) for each strain. MV was added to the culture at a final concentration of 250 μM and incubated for another 2 h. The Hcp secretion phenotype of the strains was performed by Hcp-Western blot as described in Supplemental material. Primary anti-Hfq-antibody raised in rabbit at a dilution of 1:1000 was used as a loading control for cell lysate (CL).

Bacterial killing assay

Bacterial killing assays were performed as described previously (23). *A. baumannii* T6-
and Δ*mumT* strains were used as killers/predators, while *E. coli* J53 or *E. coli*-pNYL GFP,
or *P. aeruginosa* were used as prey for this assay. The detailed procedure is provided in
the Supplemental material.

ELISA assay for HCP

From the mid-log phase cultures of the indicated strains, 0.1% inoculum from the O/N
cultures was then subcultured into fresh 5 mL of LB-medium containing $MnCl_2$ at a final
concentration of 250 µM and grown at 37 ^{0}C with shaking to an OD_{600} of 0.6 (mid-log
phase) for each strain. MV was added to the culture at a final concentration of 250 µM and
incubated for another 2 h. The rest of the protocol for Hcp-ELISA was performed as
mentioned in Supplemental material.

RNA-sequencing data analysis

RNA was isolated from WT T6- and ΔAbsR28 cells grown in LB supplemented with
$MnCl_2$ (250 µM) to an OD_{600} of 0.6 (mid-log phase), treated with MV (250 µM) for 2 h,
and purified as described above. RNA sequencing was performed by Biokart India Pvt.
Ltd. (India) using the Illumina HiSeq 4000 platform (Illumina). Analysis was performed
by Rockhopper v 2.0.3. Comparative and statistical analyses were performed using iGeak
v1.0a using the reference *A. baumannii* ATCC 17978 genome (NCBI: CP000521.1).

Pulse expression studies

An arabinose inducible vector, pWBAD30, was modified from the pBAD30 backbone for
this study. The pWBAD30-AbsR28 plasmid was transformed into ΔAbsR28
electrocompetent cells and the transformants were selected on LB-agar plates containing

495   kanamycin (50 µg/mL). ΔAbsR28 transformed with pWBAD30 served as a vector control.

496   The bacterial survival assay was performed as described Supplemental material in the

497   presence of arabinose (0.2% w/v final concentration).

Isothermal calorimetry

ITC titrations were performed using the ITC-200 (GE Healthcare). After degassing, 600

µM $MnCl_2$ was placed into the syringe and 30 µM *in vitro* transcribed AbsR28 was placed

in the reaction cell. ITC was performed over 20 injections, each 1.8 µl of $MnCl_2$ with an

interval of 120 s to allow for equilibration of the mixture between injections at a constant

stirring of 500 rpm. All reactions were performed in a buffer containing 10 mM Tris, 100

mM KCl, and pH 8.0 at 25 ^{0}C. A control experiment was performed at the same condition

in the absence of sRNA. Analysis after subtracting the control data was performed using

the Origin version 7.0 software provided with the system and data was fitted as an

independent binding model.

In vitro RNA transcription and 5'-end labeling

509   *In vitro* transcription (IVT) was performed as described earlier (76). The template for IVT

510   was obtained using *A. baumannii* ATCC 17978 genomic DNA as a PCR template and

511   forward primers containing the T7 promoter listed in Table S5. The detailed procedure is

512   provided in the Supplemental material.

In vitro structural probing

In vitro structural probing was performed as described earlier (77) with some modifications

using *in vitro* transcribed AbsR28 5'-labelled with [^{32}P]-γ-ATP. The detailed procedure is

provided in the Supplemental material.

Gel retardation assay

The gel retardation assay was performed as described earlier (78) with some modifications

using unlabeled *tssM in vitro* transcripts (250 nt upstream and 250 nt downstream from

ATG) and full-length AbsR28 *in vitro* transcripts. The detailed procedure is provided in

the Supplemental material.

RNase E-mediated degradation assay

Rnase E-mediated degradation assay was performed using the *in vitro* transcript *tssM* 5'-

labeled with $[^{32}P]$-γ-ATP. The detailed procedure is provided in the Supplemental material.

Mice infection model for *A. baumannii* pneumonia

All animal experiments under protocol BT/IAEC/2018/07 were reviewed and approved by

the Institute Animal Ethics Committee of the Indian Institute of Technology Roorkee.

Adult (6–8 week old) age-matched male BALB/c mice (n = 8 for each group, determined

using G*Power analysis) were infected intranasally with *A. baumannii* and CFU was

enumerated as previously described (79). Briefly, the mice were anesthetized and infected

intranasally with 20 µL of inoculum containing 4×10^4 CFU of the indicated strains. Mice

were euthanized at 36 h of infection, and the lungs and livers were harvested, immediately

transferred on ice, and washed with ice-cold 1X sterile PBS. The harvested organs were

chopped into pieces to enumerate the bacterial burden, histopathology, and Western blot

analysis.

Quantification and statistical analysis

Statistical analyses were performed using GraphPad Prism 8. Quantifying bands from gels

was performed using ImageJ and analyzed using GraphPad Prism 8. The details about the

number of repeated experiments, statistical tests used, significance values, and group sizes

are indicated in each figure legend.

ACKNOWLEDGEMENTS

We thank all the members of the Pathania Lab for their suggestions and fruitful discussions

regarding this manuscript. We are grateful to Prof. S.P. Mukherjee for the training required

to handle [γ-^{32}P]ATP at IIT Roorkee. We extend heartfelt thanks to Prof. N.K. Navani (IIT

Roorkee) for his scientific inputs during experiments and critical views on the manuscript.

We are thankful to Dr. Arpita Dey for helping with the ITC assay and data analysis. We

thank Dr. Amit Gaurav for helping with RNA sequencing data deposition on the NCBI

SRA. We thank Prof. P. Roy (IIT Roorkee) and members of his laboratory for animal cell

lines. We also thank Dr. Siva for the clonings required for antibody generation and

construction of TssM knockout in ATCC 17978. We are grateful to the Institute

Instrumentation Centre (IIC) in IIT Roorkee for providing facilities like ICP-MS and ITC.

This work was supported by the DBT/Wellcome Trust India Alliance Fellowship (grant

number IA/S/21/1/505588) awarded to R.P. S.B. was supported by a Research Fellowship

(Sr. No. 2061530396; Ref. No. 21/06/2015(i)EU-V) from University Grants Commission

(UGC). A.P. was supported by a fellowship (Ref. No. 09/143(1019)/2020-EMR-I) from

CSIR. K.D. was supported by a fellowship (Ref. No. DST/INSPIRE/03/2017/000008)

from DST. A.S. was supported by a fellowship from MHRD. S.P. was supported by a

fellowship (Ref. No. DBTHRDPMU/JRF/BET-22/I/2022-23/196) from DBT. The funders

had no role in study design, data collection and interpretation, or the decision to submit the

work for publication.

AUTHOR CONTRIBUTIONS

S.B. and R.P. designed the study, conceptualized the experiments and interpreted the data. S.B. performed most of the experiments, analyzed the data, and prepared figures. A.P., K.D., A.S., S.P., and N.J. helped during the experimentation. T.B. generated preliminary data for AbsR28 that implied its role in oxidative stress. J.A. helped during animal experiments. R.S. S.C. and T.K.S. provided the radioactive facility for lead acetate probing at THSTI. R.P. engaged in the acquisition of funds. S.B. and R.P. wrote the manuscript, R.P. edited the manuscript and all the authors commented on it.

CONFLICT OF INTEREST

The authors declare no competing interests.

DATA AVAILABILITY

Raw RNA-sequence data have been deposited at NCBI Sequence Read Archive (NCBI SRA) with accession number PRJNA865908 and are publicly available as of the date of publication. Target prediction of AbsR28 was performed using CopraRNA (https://rna.informatik.uni-freiburg.de/CopraRNA [accessed 2nd October 2020]) using the RACE-mapped sequence of AbsR28 (45) as input for *Acinetobacter baumannii* ATCC 17978 (NCBI Reference Sequence: NZ_CP018664, NZ_CP049363, NZ_CP039028, and NZ_CP039023) as an organism of interest. Any additional information required to analyze the data reported in this paper is available upon request from the lead contact.

REFERENCES

1. Bergogne-Bérézin E, Towner KJ. 1996. *Acinetobacter* spp. as nosocomial pathogens: Microbiological, clinical, and epidemiological features. Clin Microbiol Rev 9: 148–165.
2. Dexter C, Murray GL, Paulsen IT, Peleg AY. 2015. Community-acquired *Acinetobacter baumannii*: clinical characteristics, epidemiology and pathogenesis. Expert Rev. Anti Infect Ther 13: 567–573.
3. Falagas ME, Bliziotis IA, Siempos II. 2006. Attributable mortality of *Acinetobacter baumannii* infections in critically ill patients: A systematic review of matched cohort and case-control studies. Crit Care 10: 1–8.
4. Munoz-Price LS, Zembower T, Penugonda S, Schreckenberger P, Lavin MA, Welbel S, Vais D, Baig M, Mohapatra S, Quinn, JP, et al. 2010. Clinical outcomes of carbapenem-resistant *Acinetobacter baumannii* bloodstream infections: Study of a 2-State Monoclonal Outbreak. Infect Control Hosp Epidemiol 31; 1057–1062.
5. Peleg AY, Seifert H, Paterson DL. 2008. *Acinetobacter baumannii*: Emergence of a successful pathogen. Clin Microbiol Rev 21; 538–582.
6. Kaye KS, Pogue JM. 2015. Infections Caused by resistant Gram-negative bacteria: Epidemiology and Management. Pharmacotherapy 35; 949–962.
7. Hood MI, Skaar EP. 2012. Nutritional immunity: Transition metals at the pathogen-host interface. Nat Rev Microbiol 10; 525–537.
8. Porcheron G, Garénaux A, Proulx J, Sabri M, Dozois, CM. 2013. Iron, copper, zinc, and manganese transport and regulation in pathogenic Enterobacteria: Correlations between strains, site of infection and the relative importance of the different metal transport systems for virulence. Front Cell Infect Microbiol 3; 1–24.
9. Diaz-Ochoa VE, Lam D, Lee CS, Klaus S, Behnsen J, Liu JZ, Chim N, Nuccio SP, Rathi SG, Mastroianni JR, et al. 2016. *Salmonella* mitigates oxidative stress and thrives in the Inflamed gut by evading calprotectin-mediated manganese sequestration. Cell Host & Microbe 19; 814–825.
10. Kehl-Fie TE, Chitayat S, Hood MI, Damo S, Restrepo N, Garcia C, Munro KA, Chazin WJ, Skaar EP. 2011. Nutrient metal sequestration by calprotectin inhibits bacterial superoxide defense, enhancing neutrophil killing of *Staphylococcus aureus*. Cell Host & Microbe 10; 158–164.
11. Green ER, Juttukonda LJ, Skaar EP. 2020. The manganese-responsive transcriptional regulator MumR protects *Acinetobacter baumannii* from oxidative stress. Infect Immun 88; e00762–19.
12. Juttukonda LJ, Chazin WJ, Skaar EP. 2016. *Acinetobacter baumannii* coordinates urea metabolism with metal import to resist host-mediated metal limitation. mBio 7; e01475–16.
13. Si M, Zhao C, Burkinshaw B, Zhang B, Wei D, Wang Y, Dong TG, Shen X. 2017. Manganese scavenging and oxidative stress response mediated by type VI secretion system in *Burkholderia thailandensis*. Proc Natl Acad Sci U S A 114; 2233–2242.
14. Cao Z, Casabona MG, Kneuper H, Chalmers JD, Palmer T. 2016. The type VII secretion system of *Staphylococcus aureus* secretes a nuclease toxin that targets competitor bacteria. Nat Microbiol 2; 1-11.
15. García-Bayona L, Guo MS, Laub MT. 2017. Contact-dependent killing by *Caulobacter crescentus* via cell surface-associated, glycine zipper proteins. Elife 6; 1–26.
16. Souza DP, Oka GU, Alvarez-Martinez CE, Bisson-Filho AW, Dunger G, Hobeika L, Cavalcante NS, Alegria MC, Barbosa LR, Salinas RK, et al. 2015. Bacterial killing via a type IV secretion system. Nat Commun 6; 1–9.
17. Vassallo CN, Cao P, Conklin A, Finkelstein H, Hayes CS, Wall D. 2017. Infectious polymorphic toxins delivered by outer membrane exchange discriminate kin in myxobacteria. Elife 6; 1–24.
18. Juttukonda LJ, Skaar EP. 2015. Manganese homeostasis and utilization in pathogenic bacteria. Mol Microbiol 97; 216–228.
19. Kehres DG, Maguire ME. 2003. Emerging themes in manganese transport, biochemistry and pathogenesis in bacteria. FEMS Microbiol Rev 27; 263–290.
20. Lin, L., Capozzoli, R., Ferrand, A., Plum, M., Vettiger, A., Basler, M., 2022. Subcellular localization of Type VI secretion system assembly in response to cell–cell contact. The EMBO Journal 41(13), p.e108595.
21. Silverman JM, Brunet YR, Cascales E, Mougous JD. 2012. Structure and regulation of the type VI secretion system. Annu Rev Microbiol 66; 453–472.

22. Bernard CS, Brunet YR, Gueguen E, Cascales E. 2010. Nooks and crannies in type VI secretion regulation. J. Bacteriol 192; 3850–3860.
23. Weber BS, Ly PM, Irwin JN, Pukatzki S, Feldman MF. 2015. A multidrug resistance plasmid contains the molecular switch for type VI secretion in *Acinetobacter baumannii*. Proc Natl Acad Sci U S A 112; 9442–9447.
24. Gripenland J, Netterling S, Loh E, Tiensuu T, Toledo-Arana A, Johansson J. 2010. RNAs: Regulators of bacterial virulence. Nat Rev Microbiol 8; 857–866.
25. Romby P, Vandenesch F, Wagner EGH. 2006. The role of RNAs in the regulation of virulence-gene expression. Curr Opin Microbiol 9; 229–236.
26. Schuppli D, Georgijevic J, Weber H. 2000. Synergism of mutations in bacteriophage Qβ RNA affecting host factor dependence of Qβ replicase. J Mol Biol 295; 149–154.
27. Aubert DF, Xu H, Yang J, Shi X, Gao W, Li L, Bisaro F, Chen S, Valvano MA, Shao F. 2016. A *Burkholderia* type VI effector deamidates Rho GTPases to activate the pyrin inflammasome and trigger inflammation. Cell Host & Microbe 19(5); 664–674.
28. Hachani A, Wood TE, Filloux A. 2016. Type VI secretion and anti-host effectors. Curr Opin Microbiol 29; 81–93.
29. Jiang F, Wang X, Wang B, Chen L, Zhao Z, Waterfield NR, Yang G, Jin Q. 2016. The *Pseudomonas aeruginosa* type VI secretion PGAP1-like effector induces host autophagy by activating endoplasmic reticulum stress. Cell Rep 16(6); 1502–1509.
30. Pukatzki S, Ma AT, Revel AT, Sturtevant D, Mekalanos JJ. 2007. Type VI secretion system translocates a phage tail spike-like protein into target cells where it cross-links actin. Proc Natl Acad Sci U S A 104(39); 15508–15513.
31. Ahn S, Jung J, Jang IA, Madsen EL, Park W. 2016. Role of glyoxylate shunt in oxidative stress response. J Biol Chem 291; 11928–11938.
32. Coady A, Xu M, Phung Q, Cheung TK, Bakalarski C, Alexander MK, Lehar SM, Kim J, Park S, Tan MW, et al. 2015. The *Staphylococcus aureus* ABC-type manganese transporter MntABC is critical for reinitiation of bacterial replication following exposure to phagocytic oxidative burst. PLoS One 10; 1–24.
33. Ma AT, Mekalanos JJ. 2010. In vivo actin cross-linking induced by *Vibrio cholerae* type VI secretion system is associated with intestinal inflammation. Proc Natl Acad Sci U S A 107: 4365–4370.
34. Parsons DA, Heffron F. 2005. *sciS*, an *icmF* homolog in *Salmonella enterica* serovar Typhimurium, limits intracellular replication and decreases virulence. Infect Immun 73: 4338–4345.
35. Hassan H, Fridovich I. 1977. Regulation of the synthesis of superoxide dismutase in *Escherichia coli*. Induction by methyl viologen. J Biol Chem 252; 7667–7672.
36. Wijers, C.D., Pham, L., Menon, S., Boyd, K.L., Noel, H.R., Skaar, E.P., Gaddy, J.A., Palmer, L.D., Noto, M.J. 2021. Identification of two variants of *Acinetobacter baumannii* strain ATCC 17978 with distinct genotypes and phenotypes. Infection and Immunity 89(12).
37. Hong Y, Zeng J, Wang X, Drlica K, Zhao X. 2019. Post-stress bacterial cell death mediated by reactive oxygen species. Proc Natl Acad Sci U S A 116; 10064–10071.
38. Schieber M, Chandel NS. (2014). ROS function in redox signaling. Curr Biol 24; 453–462.
39. Hood MI, Mortensen BL, Moore JL, Zhang Y, Kehl-Fie TE, Sugitani N, Chazin WJ, Caprioli RM, Skaar EP. 2012. Identification of an *Acinetobacter baumannii* zinc acquisition system that facilitates resistance to calprotectin-mediated zinc sequestration. PLoS Pathog 8; 20–24.
40. Mortensen BL, Skaar EP. 2013. The contribution of nutrient metal acquisition and metabolism to *Acinetobacter baumannii* survival within the host. Front Cell Infect Microbiol 3; 1–10.
41. Zimbler DL, Arivett BA, Beckett AC, Menke SM, Actis LA. 2013. Functional features of TonB energy transduction systems of *Acinetobacter baumannii*. Infect Immun 81; 3382–3394.
42. Si M, Wang Y, Zhang B, Zhao C, Kang Y, Bai H, Wei D, Zhu L, Zhang L, Dong TG, et al. 2017. The type VI secretion system engages a redox-regulated dual-functional heme transporter for zinc acquisition. Cell Rep 20; 949–959.
43. Zuo Y, Li C, Yu D, Wang K, Liu Y, Wei Z, Yang Y, Wang Y, Shen X, Zhu L. 2023. A Fur-regulated type VI secretion system contributes to oxidative stress resistance and virulence in *Yersinia pseudotuberculosis*. Stress Biology 3(1); 2.
44. Janssen KH, Diaz MR, Golden M, Graham JW, Sanders W, Wolfgang MC, Yahr TL. 2018. Functional analyses of the RsmY and RsmZ small noncoding regulatory RNAs in *Pseudomonas aeruginosa*. J. Bacteriol 200; 10–1128.

45. Marden JN, Diaz MR, Walton WG, Gode CJ, Betts L, Urbanowski ML, Redinbo MR, Yahr TL, Wolfgang MC. 2013. An unusual CsrA family member operates in series with RsmA to amplify post-transcriptional responses in *Pseudomonas aeruginosa*. Proc Natl Acad Sci U S A 110; 15055–15060.

46. Romero M, Silistre H, Lovelock L, Wright VJ, Chan KG, Hong KW, Williams P, Cámara M, Heeb S. 2018. Genome-wide mapping of the RNA targets of the *Pseudomonas aeruginosa* riboregulatory protein RsmN. Nucleic Acids Res 46; 6823–6840.

47. Sharma R, Arya S, Patil SD, Sharma A, Jain PK, Navani NK, Pathania R. 2014. Identification of novel regulatory small RNAs in *Acinetobacter baumannii*. PLoS One 9; e93833.

48. Bandyra KJ, Said N, Pfeiffer V, Górna MW, Vogel J, Luisi BF. 2012. The seed region of a small RNA drives the controlled destruction of the target mRNA by the endoribonuclease RNase E. Mol Cell 47; 943-953.

49. Balasubramanian D, Vanderpool CK. 2013. New developments in post-transcriptional regulation for operons by small RNAs. RNA Biol 10; 337–341.

50. Cianfanelli FR, Monlezun L, Coulthurst SJ. 2016. Aim, Load, Fire: The type VI secretion system, a bacterial nanoweapon. Trends Microbiol 24; 51–62.

51. Wright PR, Richter AS, Papenfort K, Mann M, Vogel J, Hess WR, Backofen R, Georg J. 2013. Comparative genomics boosts target prediction for bacterial small RNAs. Proc Natl Acad Sci U S A 110; 3487-3496.

52. Afonyushkin T, Večerek B, Moll I, Bläsi U, Kaberdin VR. 2005. Both RNase E and RNase III control the stability of sodB mRNA upon translational inhibition by the small regulatory RNA RyhB. Nucleic Acids Res 33; 1678–1689.

53. Caron MP, Lafontaine DA. Massé E. 2010. Small RNA-mediated regulation at the level of transcript stability. RNA Biol 7; 140–144.

54. Guillier M, Gottesman S. 2008. The 5′ end of two redundant sRNAs is involved in the regulation of multiple targets, including their own regulator. Nucleic Acids Res 36; 6781–6794.

55. Morita T, Maki K, Aiba H. 2006. RNase E-based ribonucleoprotein complexes: Mechanical basis of mRNA destabilization mediated by bacterial noncoding RNAs. Genes Dev 19; 2176–2186.

56. Prévost K, Desnoyers G, Jacques JF, Lavoie F, Massé E. 2011. Small RNA-induced mRNA degradation achieved through both translation block and activated cleavage. Genes Dev 25; 385–396.

57. Urban JH, Vogel J. 2007. Translational control and target recognition by *Escherichia coli* small RNAs in vivo. Nucleic Acids Res 35; 1018–1037.

58. Vogel J, Luisi BF. 2011. Hfq and its constellation of RNA. Nat Rev Microbiol 9; 578–589.

59. Monteith AJ, Skaar EP. 2021. The impact of metal availability on immune function during infection. Trends Endocrinol Metab 32; 916–928.

60. Murdoch CC, Skaar EP. 2022. Nutritional immunity: the battle for nutrient metals at the host–pathogen interface. Nat Rev Microbiol 20; 657–670.

61. Lin J, Zhang W, Cheng J, Yang X, Zhu K, Wang Y, Wei G, Qian PY, Luo ZQ, Shen X. 2017. A *Pseudomonas* T6SS effector recruits PQS-containing outer membrane vesicles for iron acquisition. Nat Commun 8; 14888.

62. Brunet YR, Zoued A, Boyer F, Douzi B, Cascales E. 2015. The type VI secretion TssEFGK-VgrG phage-like baseplate is recruited to the TssJLM membrane complex via multiple contacts and serves as assembly platform for tail tube/sheath polymerization. PLoS Genet 11; 1–21.

63. Felisberto-Rodrigues C, Durand E, Aschtgen MS, Blangy S, Ortiz-Lombardia M, Douzi B, Cambillau C, Cascales E. 2011. Towards a structural comprehension of bacterial type VI secretion systems: Characterization of the TssJ-TssM complex of an *Escherichia coli* pathovar. PLoS Pathog 7; 1–11.

64. Ma LS, Narberhaus F, Lai EM. 2012. IcmF family protein TssM exhibits ATPase activity and energizes type VI secretion. J Biol Chem 287; 15610–15621.

65. Stietz MS, Liang X, Li H, Zhang X, Dong TG. 2020. TssA–TssM–TagA interaction modulates type VI secretion system sheath-tube assembly in *Vibrio cholerae*. Nat Commun 11; 1–11.

66. Weber BS, Miyata ST, Iwashkiw JA, Mortensen BL, Skaar EP, Pukatzki S, Feldman MF. 2013. Genomic and functional analysis of the type VI secretion system in *Acinetobacter*. PLoS One 8; e55142.

67. Meumann EM, Anstey NM, Currie BJ, Piera KA, Kenyon JJ, Hall RM, Davis JS, Sarovich DS. 2019. Genomic epidemiology of severe community-onset *Acinetobacter baumannii* infection. Microb Genomics 5; 1–13.

68. Traglia G, Chiem K, Quinn B, Fernandez JS, Montaña S, Almuzara M, Mussi MA, Tolmasky ME, Iriarte A, Centrón D, et al. 2018. Genome sequence analysis of an extensively drug-resistant

Acinetobacter baumannii indigo-pigmented strain depicts evidence of increase genome plasticity. Sci Rep 8; 1–15.

69. Wright MS, Haft DH, Harkins DM, Perez F, Hujer KM, Bajaksouzian S, Benard MF, Jacobs MR, Bonomo RA, Adams MD. 2014 New insights into dissemination and variation of the health care-associated pathogen *Acinetobacter baumannii* from genomic analysis. mBio 5; e00963-13.

70. Mougous JD, Cuff ME, Raunser S, Shen A, Zhou M, Gifford CA, Goodman AL, Joachimiak G, Ordoñez CL, Lory S, et al. 2006. A virulence locus of *Pseudomonas aeruginosa* encodes a protein secretion apparatus. Science 312; 1526–1530.

71. Zheng J, Shin OS, Cameron DE, Mekalanos JJ. 2010. Quorum sensing and a global regulator TsrA control expression of type VI secretion and virulence in *Vibrio cholerae*. Proc Natl Acad Sci U S A 107; 21128-21133.

72. Lopez J, Ly PM, Feldman MF. 2020 The tip of the VgrG spike is essential to functional type VI secretion system assembly in *Acinetobacter baumannii*. mBio 11; e02761–19.

73. Di Venanzio G, Flores-Mireles AL, Calix JJ, Haurat MF, Scott NE, Palmer LD, Potter RF, Hibbing ME, Friedman L, Wang B, Dantas G. 2019. Urinary tract colonization is enhanced by a plasmid that regulates uropathogenic *Acinetobacter baumannii* chromosomal genes. Nat Commun 10(1); p.2763.

74. Knippel RJ, Wexler AG, Miller JM, Beavers WN, Weiss A, de Crécy-Lagard V, Edmonds KA, Giedroc DP, Skaar EP. 2020. *Clostridioides difficile* senses and hijacks host heme for incorporation into an oxidative stress defense system. Cell Host & Microbe 28; 411-421.

75. Livak KJ, Schmittgen TD. 2001. Analysis of relative gene expression data using real-time quantitative PCR and the 2-ΔΔCT method. Methods 25; 402–408.

76. Desnoyers G, Massé E. 2012. Noncanonical repression of translation initiation through small RNA recruitment of the RNA chaperone Hfq. Genes Dev 26; 726–739.

77. Dambach M, Sandoval M, Updegrove TB, Anantharaman V, Aravind L, Waters LS, Storz G. 2015. The Ubiquitous yybP-ykoY riboswitch is a manganese-responsive regulatory element. Mol Cell 57; 1099–1109.

78. Matera G, Altuvia Y, Gerovac M, El Mouali Y, Margalit H, Vogel J. 2022. Global RNA interactome of *Salmonella* discovers a 5′ UTR sponge for the MicF small RNA that connects membrane permeability to transport capacity. Mol Cell 82; 629–644.

79. Palmer LD, Green ER, Sheldon JR, Skaar EP. 2019. Assessing *Acinetobacter baumannii* virulence and persistence in a murine model of lung infection. Methods Mol Biol 1946; 289–305.

Figure 1. *A. baumannii* **T6+ cells are defective in Mn^{2+}-uptake and unable to cope with**

oxidative stress. (**A**) Bacterial cells were co-incubated with the phagocytic cells (human

blood-derived neutrophils and macrophage RAW 264.7 cell line) for 4 h at an MOI of 1:1.

The percentage of bacterial survival was enumerated by accounting for the respective

untreated control (without phagocytic cells) as 100%. The data represent the mean of four

independent experiments, each in biological triplicates ± SEM. (**B**) Growth of the indicated

strains in the presence or absence of MV (250 μM) in LB broth. The data represents the

mean of four biological replicates, each in technical triplicates ± SD. (**C**) Bacterial

intracellular ROS generation was determined by measuring the fluorescence of DCF. The

data represents the mean of four biological replicates ± SD. (**D**) Neutrophils were co-

incubated with either WT T6- or WT T6+ cells, and the expression of *mumT*, *znuB*, and

tonB in the bacterial cells was determined by qRT-PCR. The data represent the mean of

three independent experiments, each in technical triplicates ± SEM. (**E and F**) Intracellular

^{55}Mn, ^{66}Zn, and ^{56}Fe in cell pellet (OD$_{600}$ normalized volume) were quantified by ICP-MS.

The data represents the mean of three biological replicates ± SD. Statistical significance

was determined using a multiple comparison two-way ANOVA test with the Sidak

correction for multiple comparisons comparing the means of each group to one another (A,

E, F) and Student's t-test (B, C, D). ** denotes p-value <0.01, *** denotes p-value <0.001,

**** denotes p-value <0.0001, ns denotes not significant.

Figure 2. Deletion of *mumT* exhibited a significant increase in *hcp* expression under oxidative stress. (**A**) A schematic representation of T6SS (upper panel) and a schematic layout of the T6SS gene cluster in *A. baumannii* ATCC 17978 (lower panel). (**B**) Transcription of *hcp* was determined by qRT-PCR. The data represent the mean of three independent experiments, each in technical triplicates ± SEM. Statistical significance was determined using the Student's t-test. * denotes p-value <0.05. (**C**) The Hcp secretion profile of the indicated strains was confirmed by a Western blot of cell lysate (CL) and culture supernatants (S). WT T6+ cell and purified His$_6$-Hcp were used as a positive control

for this assay. The Hfq antibody was used as a loading control for CL. The data represents

three independent experiments where the samples were pooled and run on one gel. (**D**) In

the T6SS competition assay, prey cells were subjected to killing by incubation with WT

T6-, WT T6+, Δ*mumT*, and Δ*tssM* cells. The survival percentage of prey cells was

enumerated by accounting for the respective untreated control (without predator/killer

cells) as 100%. WT T6+ and Δ*tssM* (the deletion mutant of TssM in WT-17978, which

does not have functional T6SS) cells were used as T6+ and T6-, respectively, control for

this assay. The data represents the mean of three biological replicates ± SD. Statistical

significance was determined using two-way ANOVA with Tukey's multiple comparison

test for each sample with the same prey killed by the killer strains. **** denotes p-value

<0.0001. (**E**) Detection of Hcp secretion from individual colonies of WT T6-, and Δ*mumT*

strains by Hcp-ELISA. * Well indicates purified His_6-Hcp, used as a positive control for

Hcp-ELISA.

Figure 3. AbsR28 effectively represses T6SS in the presence of Mn²⁺. (**A**) The expression of several sRNAs and *hfq* was examined by qRT-PCR. The data represent the mean of three independent experiments, each in technical triplicates ± SEM. (**B**) RNA-seq analysis comparing RNA from MV and MnCl₂ (both at 250 μM final conc.) treated

ΔAbsR28 to a WT T6- control in duplicates. A dotted black line denotes $p < 0.05$. (**C**) The

transcription of each T6SS structural gene in WT T6- and ΔAbsR28 cells was determined

by qRT-PCR. The data represent the mean of three independent experiments, each in

technical triplicates ± SEM. (**D**) Transcription of *hcp* was determined by qRT-PCR. Un

and In denote uninduced (without arabinose) and induced (with arabinose), respectively.

The data represent the mean of three independent experiments, each in technical triplicates

± SEM. (**E**) The Hcp secretion profile of the bacterial strains (in biological duplicates) was

tested by Western blot of cell lysate (CL) and cell-free supernatant (S). The Hfq antibody

was used as a loading control for CL and to confirm that the supernatants were free from

the bacterial cell. Un and In denote uninduced (without arabinose) and induced (with

arabinose), respectively. (**F**) In the T6SS competition assay, *E. coli* J53 prey cells were

subjected to killing by ΔAbsR28-pWBAD30AbsR28 and ΔAbsR28-pWBAD30 strains.

The data represents the mean of three biological replicates ± SD.

Figure 4. Mn²⁺ binds to AbsR28 and alters its structure. (**A**) The secondary structure of

A. baumannii AbsR28 sRNA was predicted by Rfam software. (**B**) Isothermal titration

calorimetry (ITC) of Mn²⁺ to AbsR28. (**C and D**) Lead acetate probing of the 5'end-labeled

[γ-^{32}P]ATP AbsR28 in an increasing concentration of either $MnCl_2$ or $MgCl_2$ provided in

the structure buffer. Lanes indicated as T1 and OH ladders were obtained from the same

labeled AbsR28 after incubation with RNase T1 and hydroxyl anions, respectively. RNase

T1 digestion was performed in duplicates at 0.1 and 1.0 U concentrations. The position of

cleaved G residues is marked at the left of the gel. Statistical significance was determined

using the multiple comparison two-way ANOVA test with the Sidak correction for multiple

comparisons comparing the means of each group to one another (A, D, F) and Student's t-

test (C). * denotes p-value <0.05, ** denotes p-value <0.01, *** denotes p-value <0.001,

**** denotes <0.0001, ns denotes not significant.

Figure 5. Mn²⁺ is required for AbsR28 to base pair with *tssM* mRNA and represses T6SS by potentiating RNase E-mediated degradation of *tssM* mRNA. (**A**) Predicted base pairing between *tssM* mRNA and AbsR28 by the CopraRNA bioinformatics tool. (**B**

and C) Gel retardation assay of unlabeled *tssM in vitro* transcripts and unlabeled full-

length AbsR28 *in vitro* transcripts in structure buffer containing either 10 mM $MnCl_2$ or

$MgCl_2$. (**D**) Using ImageJ software, unbound *tssM* transcripts were quantified from Figures

B and C (n = 2 independent experiments). Nonlinear regression was used to fit the curve.

(**E**) The stability of *tssM* transcripts was determined after adding a RNA synthesis inhibitor

Rifampicin (400 µg/mL), by qRT-PCR and plotted as the fold change relative to 0 min for

each sample. The data represents the mean of technical triplicates ± SD. (**F**) The cleavage

of the 5'end-labeled [γ-^{32}P]ATP *tssM in vitro* transcripts was assessed by incubating with

RNase E in structure buffer containing $MnCl_2$ for 0 and 3.5 h in the presence of either

unlabeled AbsR28 (lanes 5-6) or *A. baumannii* Hfq72 (lanes 7-8) or unlabeled AbsR28 +

A. baumannii Hfq72 (lanes 9-10) or *E. coli* Hfq (lanes 11-12) or unlabeled AbsR28 + *E.*

coli Hfq (lanes 13-14). Addition of Hfq results in a slower migration which is evident from

the retardation of bands (7-14). The lower panel represents the quantification of the

complete 5'-labeled *tssM* transcripts that remain after RNase E-mediated cleavage

(denoted by a filled triangle from lanes 1-6 and an empty triangle from lanes 7-14) using

ImageJ software (n = 3 independent experiments). Statistical significance was determined

using the one-way ANOVA test with Dunnett's multiple comparisons (E) and Student's t-

test (F). *** denotes p-value <0.001, **** denotes p-value <0.0001, ns denotes not

significant.

Figure 6. Mn^{2+}-dependent AbsR28-mediated T6SS repression is required for *A.
***baumannii* pathogenesis.** (**A**) Bacterial strains were co-incubated with neutrophils for 4 h

at an MOI of 1:1. The percentage of bacterial survival was enumerated, accounting for the

respective untreated control (without neutrophils) as 100%. The data represent the mean of

three independent experiments, each in biological triplicates ± SEM. (**B**) Enumeration of

bacterial burden recovered from mice lungs and liver (n = 8) infected with WT T6-, WT

T6+, Δ*mumT,* or ΔAbsR28 strains at 36 hpi. Data represents mean ± SD. (**C**)

Histopathology of the mice lungs (H&E stained) infected with WT T6-, WT T6+, Δ*mumT*,

or ΔAbsR28 strains at 36 hpi. Tissue infiltration is indicated by necrosis (arrowheads) and

alveolar inflammation (asterisks). The scale bar is 100 μM. Statistical significance was

688 determined using the one-way ANOVA test with Tukey's multiple comparisons (A, B). *

689 denotes p-value <0.05, *** denotes p-value <0.001, ns denotes not significant.

Figure 7. A proposed model of AbsR28-mediated post-transcriptional repression of T6SS in *A. baumannii* during oxidative stress. In the presence of Mn^{2+}, AbsR28 base pairs with *tssM* mRNA and induces RNase E-mediated processing of *tssM*, which results in the repression of T6SS in *A. baumannii*.