Evolution to increased positive charge on the viral spike protein may be part of

the adaptation of SARS-CoV-2 to human transmission.

Matthew Cotten*, My V.T. Phan

Author affiliations: Medical Research Council–University of Glasgow Centre for Virus

Research, Glasgow, Scotland, UK (MC), UK Medical Research Council–Uganda

Virus Research Institute and London School of Hygiene and Tropical Medicine

Uganda Research Unit, Entebbe, Uganda (MC, MVTP); UK

*Correspondence to Matthew Cotten, email: matthew.cotten@lshtm.ac.uk

Abstract

The severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2), the causative agent of

the coronavirus disease 2019 (COVID-19) pandemic, continues to evolve and infect individuals. The

exterior surface of the SARS-CoV-2 virion is dominated by the spike protein and the current work

examined spike protein biochemical features that have changed during the 2 years that SARS-CoV-2

has infected humans. These biochemical properties may influence virion survival and promote

movement through the environment and within the human airway to reach target cells to bind, enter

and establish the next round of infection. In addition to selective pressure to avoid immune

recognition of viral proteins, we hypothesised that SARS-CoV-2 emerged from an animal reservoir

capable of human infection and transmission but in a sub-optimum state and a second level of

selective pressure is acting on these biochemical features. Our analysis identified a striking change in

spike protein charge, from -8.3 in the original Lineage A and B viruses to -1.26 in the current Omicron

viruses. In summary, we conclude that in addition to immune selection pressure, the evolution of

SARS-CoV-2 has also altered viral spike protein biochemical properties. Future vaccine and

therapeutic development should also exploit and target these biochemical properties.

Introduction

The severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2), the causative agent of

the coronavirus disease 2019 (COVID-19) epidemic, continues to evolve and infect individuals.

Similar to other viruses, the SARS-CoV-2 virion biochemical properties play an important role in

controlling virus transmission. After replication in an infected individual and release from an infected

cell, onward transmission requires survival of the virion to reach susceptible cells in a new host

individual initiating the next round of infection. The physical properties of the surface proteins of the

virus such as charge, size, hydrophobicity and folding may influence movement of the virion through

the environment, promoting or limiting binding of the virion to the external surfaces. Once reaching a susceptible individual, virion physical properties may influence movement within the human airway and determine the ability of an infecting virion to reach target cells to bind, enter and replicate (Adamczyk et al. 2021). The exterior surface of the SARS-CoV-2 virion is dominated by the spike protein and the current work examines simple spike protein features that have changed during the 2 years of the SARS-CoV-2 pandemic. In addition to selective pressure to avoid immune recognition of viral proteins, we hypothesise that SARS-CoV-2 emerged from an animal reservoir capable of human infection and transmission but in a sub-optimum state. Additionally, there is a second level of selective pressure to adjust to the physical transmission between humans. Evidence for this adaptation can be found in changes in the SARS-CoV-2 spike protein over recent evolution. With over 11 million SARS-CoV-2 genomic sequences generated globally from across the pandemic, many of these sequences have intact spike gene sequences that can be used to monitor change across the 2 years of human host evolution of this virus.

Much of the observed spike protein substitutions may be in response to the developing immune response to this new pathogen, which is reflected in substitutions occurring in the immune-exposed S1 domain of the spike protein and there is ample evidence that many of these spike protein changes allow escape from host immunity (Tzou et al. 2022)(Greaney, Loes, et al. 2021)(Greaney, Starr, et al. 2021)(Greaney et al. 2022) (Cao et al. 2022) (Dejnirattisai et al. 2022) (DeGrace et al. 2022). There may also be evolutionary selection for protein changes that improve host interactions apart from immune evasion. These include altering spike/receptor binding kinetics, protease cleavage events, tertiary structure (S1/S2 interactions after cleavage) or the physical properties of the virion (charge, hydrophobicity, and protein folding or secondary structure) in ways that might improve transmission. To explore the role of the biochemical features of the spike protein in human transmission, we monitored changes in spike biochemical features over the two years that SARS-CoV-2 has been evolving in humans and report an increase in spike protein positively charge especially among the virus lineages that were highly prevalent.

Results.

The SARS-CoV-2 spike protein physical features were calculated from spike protein sequences from across 2 years of the COVID-19 epidemic. Features that could be quantitated from protein sequence were used (see Methods), including charge at pH 7.4, Kyle and Doolittle GRAVY score (Kyte & Doolittle 1982) (which is a measure of hydrophobicity), an instability index derived from dipeptide content (Guruprasad et al. 1990), properties influencing protein folding (percent helix, fold or sheet as predicted from amino acid content), individual amino acid total fraction and di-amino acid total fraction.

A dominant pattern of SARS-CoV-2 evolution during the two years of human adaptation has been the regular appearance and the subsequent regional and then global dominance of lineages. These lineages typically encode a small set of amino acid changes from earlier lineages, many of

which are likely to provide temporary or long-term advantage for the viral lineage. An analysis was

performed to identify spike physical features most strongly linked with SARS-CoV-2 lineages (Figure

1). The first 300 reported genomes from each major lineage were collected, spike protein sequences

were extracted and the physical features of each protein were collected into a matrix. The top features

distinguishing SARS-CoV-2 lineages were identified with charge as the most important feature (Figure

1A). A principal component analysis using the top 8 features (charge, gravy, fraction T, fraction R,

instability, fraction G, fraction D and fraction K), provided clustering of spike sequences by lineage

(Figure 1B). These results support the idea that spike protein charge (among other features) is an

important determinant of the lineages that have evolved during the first two years of the COVID-19

epidemic.

85
Figure 1. Identification of spike protein charge association with SARS-CoV-2 lineage. **Panel A:**

A set of 300 spikes sequences extracted from the first 300 SARS-CoV-2 genomes per lineage (by

date of collection) was analyzed, features for each sequence were collected (see Methods). SKLearn

feature selection (Pedregosa, F. and Varoquaux, G. and Gramfort, A. and Michel, V. et al. 2011) was

used to identify features that most accurately identified the sequence lineage. The importance of

features were ranked in order. **Panel B:** The top 8 features (charge, gravy, fraction T, fraction R,

instability, fraction G, fraction D, fraction K) were further used in a principal component analysis to

cluster the same set of SARS-CoV-2 spike sequences. Each node represents a single spike

sequence, nodes were coloured by Pangolin lineage assigned to the genome from which the spike

sequence was obtained. Lineage colouring is explained in the figure legend to the right**.**

96
97
Changes in charge of spike protein across the epidemic were investigated. Plotting total spike

charge for all genomes per month of the epidemic showed a clear pattern of increase in charge over

two years of evolution (Figure 2, panel A). Median spike charge was -8.3 in the original SARS-CoV-2

viruses reported in late 2019 to early 2020, by March 2020, an increase in positive charge to -7.28

was observed. Subsequently, an additional increase in positive charge occurred in mid-2021 to -3.28,

and most recently a charge increase occurred in late 2020/early 2021 to -1.26.

A.

B.

105
Figure 2.**Panel A. Total SARS-CoV-2 spike charge per epidemic month**. All available SARS-CoV-

2 genomes up to 24 September 2022 were retrieved from GISAID (GISAID 2020) and the spike

protein sequence was extracted (if intact). Total charge at pH 7.4 was calculated and values were

plotted using a violin plot by month of sample collection. For each epidemic month the violin plot

depicts the distributions of calculated spike charge for all available SARS-CoV-2 genomes. **Panel B.**

Spike charge in major SARS-CoV-2 lineages. For each lineage, all available spike sequences were

collected (up to 24 September 2022), total charge was measured and violin plots prepared to show

the charge distribution by lineage. Lineages (indicated at bottom of chart) were ordered by their

appearance in the epidemic.

These spike protein charge increases can be attributed to the major successful lineages

reported over time (Figure 2B). The B.1, B.1.1 and B.1.1.7 (Alpha) lineages that dominated the first

year of the epidemic encoded spike proteins with charges between -8 and -6 while the B.1.351 (Beta)

and B.1.525 (Eta) lineages showed a further increase in charge to around -4.5. The B.1.617.2 (Delta)

lineage and sublineages (AY.x) displayed further increase in charge. Most recently, the Omicron

variants (including BA.1, BA.1.1, BA.2, BA.2.12,BA.3, BA.4 and BA.5) show further spike charge

increases with the majority of Omicron encoded spike proteins showing charge at -1.26 (Figure 2,

panel B).

Some indication of functional consequences of the observed changes in spike charge can be

obtained from the location on the charged amino acid substitutions in the spike protein. Sets of spike

sequences (extracted from the first 300 reported genomes per select lineage) were processed to

illustrate the changes to more negative charge (blue) or more positive charge (orange/red) in the

protein relative to the initial Lineage B genome sequences (Supplementary Figure 1). The initial

change in charge was a substitution of an aspartic acid residue (D, with a calculated charge of -1) by

a glycine (G, neutral). In some early lineages (e.g. A.23.1), proline (P) at position 681was substituted

with the positively charged arginine (R),or Q680 was substituted with a partially charged histidine H

residue. The P681R positive substitution promotes furin cleavage and activation of the spike protein

for cell fusion (Lubinski et al. 2022)(Liu et al. 2022). The Delta lineage spike proteins encoded

additional positive charge in the ACE2 binding region, as well as in the far amino terminal region and

near the heptad repeat (HR1) which may also enhance membrane fusion activity. More recently, a

number of positive substitutions have occurred in the Omicron lineage virus spike proteins with

predominance of positively charged changes in the receptor binding domain (Supplementary Figure

1), suggesting a role of increased charge in spike/receptor interactions.

It is probable that the spike protein has an upper limit to the amino acid charge that it can

allow for proper folding, assembly and function. This upper charge value will be determined by the

acquisition of optimum transmission properties in balance with immune selection. After the regular

increase of spike protein charge observed up to the appearance of the Omicron lineages, an

indication of a stasis in positively-charged amino acid accumulation is now displayed by SARS-CoV-2

Omicron lineages. The majority of Omicron sub-lineages remain at spike charge -1.26 (Figure 3A)

although a few specific Omicron sub-lineages show changes toward more positive or negative charge

(e.g. BA.2.75.1 more positive, BA.4.6 more negative, as illustrated in Figure 2B) with the additional

changes often associated with immune selection. To monitor the current trends of spike protein

changes, we calculated the fraction of reported genomes with spike charge greater than or less than

the Omicron mean charge of -1.26 and documented how these fractions had changed over the last 4

months of the pandemic (Figure 3B). The majority of encoded spike proteins are almost exclusively

from Omicron lineage viruses and show a charge of -1.26. However, a small fraction of genomes

encode spikes proteins with slightly more or less charge (Figure 3B) with the greater trend (20% of all

reported genomes in September 2022) showing more negative charge (Figure 3B).

Figure 3. Recent changes in spike protein charge. **Panel A**: All available spike proteins from genomes

with sample collection dates of June-Sept 2022 were analyzed for total spike charge. A histogram of

the calculate total spike charges for the entire set is shown in panel A with the **kernel density**

estimation (**KDE**) line in red. A major peak at -1.26 is observed wit small outlier peaks of genomes

with more negative and more positive spike proteins **Panel B**: For each month (over the period June 1

to Sept 26 2022) the fraction of reported genomes for that month with charge greater than or less than

the majority value of -1.26 was calculated.

Lastly, we investigated if a similar pattern of spike charge evolution could be observed in
other coronaviruses that have made a transition to human transmission. In recent history, several
coronaviruses (in addition to SARS-CoV-2) have been observed to jump hosts. For example,
coronavirus 229E is commonly detected in humans and very close coronaviruses have been identified
in bats (Victor Max Corman et al. 2015) (Tao et al. 2017) and camels (Victor M. Corman et al. 2016)
(Sabir et al. 2016) suggesting movement of the virus between hosts. All available coronavirus 229E
full genomes sequences were retrieved from GenBank, the spike coding region was extracted from
the genomes, translated and total charge was calculated. A difference from -26 to -8, or almost 18
charge units is seen comparing 229E-like viruses from bats to 229E from humans (Figure 4A) and
almost 9 charge unit difference was observed in spike median charge comparing 229E viruses from
camel vs. human infections (Figure 4A).

Infection with coronavirus OC43 is common in humans and closely related viruses are found
in cattle (bovine coronavirus, BCoV) (Vijgen et al. 2006) and pigs (porcine hemagglutinating
encephalomyelitis virus, PHEV) (Vijgen et al. 2005) (Vijgen et al. 2006). Comparing the three OC43-
type virus groups, the human virus OC43 has an increased charge of ca. 5 units compared to PHEV
and ca.7 units compared to BCoV (Figure 4B).

The commonly known host for the Middle East Respiratory Syndrome coronavirus (MERS-
CoV) is dromedary camels; however, zoonosis and serious human infections occur frequently
(Cotten et al. 2013) (Cotten et al. 2014) (Memish et al. 2014) (Zhou et al. 2021) (So et al. 2019) as
reviewed in (Peiris & Perlman 2022). From 698 full MERS-CoV genomes available in GenBank, there
was no strong difference in the encoded spike charge of virus sequences derived from human versus
camels infections (Figure 4C).

Considering the location of the charge differences in the spike proteins, for coronavirus 229E,
the charge increases occurred throughout the protein, although there is a slightly higher number of
positive changes in the receptor binding region of the human infection derived viruses (Figure 4D).
For OC43, the porcine and human viruses also show increases in positive charge throughout the
spike protein, the porcine PHEV also showed a slight enrichment in positive charge in the receptor
binding region (Figure 4E).

Figure 4. Spike charges from select groups of coronaviruses that have moved into humans

195 **(Panels A-C)**. All available full genomes for the indicated coronaviruses were retrieved from

GenBank, the spike coding region was identified and translated into protein and total charge at ph 7.4 was calculated. Violin plots indicate the charges of each collection of spike proteins, median values are indicated by the open square. **Panel A**. Coronavirus 229E from bat, camel or human infections, **Panel B**. BCoV (from bovine infections) PHEV (from porcine infections) and OC43 (from human infection), **Panel C.** MERS-CoV from camel or human infection. **Panel D-F: Consensus spike protein sequences were generated from the indicated virus groups and charged amino acid changes were determined.** Charge changes were colored from dark blue (change from positively to negatively charged amino acid (AA)), blue change from neutral to negatively charged AA), orange (change from neutral to positively charged AA) and red (change from negative to positively charged AA). **Panel D:** 229E spike from human infections compared to 229E spike from camel infections, **Panel E**: Human OC43 spike compared to BCoV spike, **Panel F**: Human OC43 spike compared to PHEV spike. Key spike protein features of each group's spike protein are shown in the upper portion of each panel.

Discussion

After more than two years of the COVID-19 pandemic and with the availability of >11 million SARS-CoV-2 genome sequences, a trend of SARS-CoV-2 spike protein charge can be observed, with successive lineages showing an increase in positive charge over earlier lineages. Over the course of the pandemic, the SARS-CoV-2 spike protein has evolved from a protein with a total charge of -8.28 in the original Lineage A and B viruses to a protein with a total charge of -1.26 in the majority of the currently circulating Omicon lineage viruses. This pattern has been noted previously (Pawłowski 2021) (Nie et al. 2022). We expand on these observations, and document lineage patterns and sites of change in the spike protein and explore similar phenomena of evolution to more positive charge in two other coronaviruses (OC43 and 229E) that have moved between animals and humans.

This study does not identify a mechanistic basis for the increased spike charge although there are several possible transmission steps that might be promoted by increasing charge. Exposed, positively charged spike amino acids should promote interactions with negatively charged cellular structures. Interactions with negatively-charged heparin have been reported with SARS-CoV-2 spike (Kim et al. 2020) and negatively-charged sialylated glycans are reported to promote entry of SARS-CoV-2 (Nguyen et al. 2022). The upper respiratory tract is coated by and protected by mucins, frequently modified with sialic acid or phosphorylated, high mannose N-glycans (Byrd-Leotis et al. 2021) which present a negatively charged matrix that could either promote or protect against viral transmission. The SARS-CoV-2, OC43, and BCoV virions display binding to negatively charged carbohydrate structures found in the airway (Byrd-Leotis et al. 2021) and the ionic environment of the human upper respiratory tract may favour binding and transmission of viruses with increased positive charge. Perhaps it is not surprising that both OC43 and 229E coronaviruses exhibited increases in spike positive charge after moving from animal hosts (cow, pig and camel) to human hosts (Figure 5).

A similar change in MERS-CoV was not observed, however MERS-CoV currently shows only limited

human to human transmission with most known transmission chains ending after 2 to 3 human to

human transmission events as shown in (Assiri et al. 2013) and (Cotten et al. 2013). MERS-CoV

might not have experienced sufficient number of human replication cycles or have undergone the

same level of selection for human transmission that OC43, 229E and SARS-CoV-2 have experienced.

For both OC43 and 229E coronaviruses moving to humans, the broad location of the positive

changes across the spike protein sequence suggested that positive charge may be promoting several

functions including receptor binding, furin cleavage, cell fusion as well as antigenic changes or less

specific changes to avoid or promote ionic interactions during transmission.

There is likely a limit to the accumulation of positively charged residues in the SARS-CoV-2

spike protein. Functional constraints exist, there may also be penalties associated with non-specific

binding due to excess positive charge, and there are certainly charge influences on protein folding

and higher order protein interactions (Creighton 2002). Our prediction is that the SARS-CoV-2 protein

will reach some upper limit of charge defined by these constraints. Indeed, we observe that the

majority of Omicron lineages encode spike proteins with charge -1.26, after more than 6 months of

evolution (Figures 2A and 2B). A small fraction of genomes with more positive charge or less positive

charge have appeared, but the global tendency across all reported genomes from June to September

2022 is a modest decline in the positive charge (Figure 3b) which suggest the upper limit to charge

has been reached.

Could these changes in spike charge have occurred by chance and not be a response to

selective pressure? Of the 20 standard amino acids (AA), only 2 AA have negatively charged side

chains, 2 AA have positively charged side chains while the remaining 16 AA are neutral at pH 7.4..

Assuming equal probability of any AA change, there is an 18/20 chance of a negative AA being

substituted by a neutral or positively charged AA and the majority of change opportunities would result

in loss of negative charge. However, natural selection is more complex, because the genetic code

uses 3 adjacent nucleotides to encode an AA, there are multiple encoding possibilities for each AA,

the codon redundancy is not identical for each AA and the number of nucleotide changes required to

produce any particular AA change can be 1, 2 or 3. This has resulted in an evolved protein stability in

the genetic code (Chan et al. 2020) with AA changes that maintain rather than change physical

properties (negative, positive, polar, non-polar, aromatic) more likely based on the codon array

(Livingstone & Barton 1993) and the nucleotide changes required for an AA change. For example, the

probability of a negative AA to negative AA change is 0.333 while the probabilities of change of a

negative AA to a non-polar, aromatic, polar or positive AA are 0.051, 0.044, 0.028 and 0.044

respectively, with changes away from a negative charged AA nearly 10-fold less likely to occur than

conserving the negative charge at that position (Livingstone & Barton 1993). For these reasons, it

appears that the accumulation of positive charge on spike protein has not occurred by chance and is

likely providing some selective advantage for the virus. It should also be noted that the observed

charge changes in exposed virion proteins seem to be limited to spike. Two additional SARS-CoV-2

proteins are externally exposed, the E protein (ORF4) and the M protein (ORF5), showing no consistent change in the charge of either of these proteins across the 2 years of the epidemic (results not shown).

Obermeyer et al. documented AA substitutions associated with SARS-CoV-2 fitness (Obermeyer et al. 2022). Consistent with the idea that the increase in positive charges is not by chance, of the top 20 substitutions increasing SARS-CoV-2 fitness, 14 substitutions were in the spike protein, among which 4 were changes that increased positive charge while only 1 of 14 introduced a negative charge in spike (Obermeyer et al. 2022).

Natural selection could be acting on multiple features of the spike protein. The necessity to avoid host immune responses is likely to be the major selective force acting on the virus. This results in the amino acid changes, which in turn are determined by epitopes. The selection for increased charge in the spike protein is probably occurring in the background, not as a major shift needed to bypass immune responses. However, the increase in charge may improving survival and transmission in humans in subtle ways, and this advantage, when multiplied over the millions of infections can provide some of the growth and infection advantages seen by new SARS-CoV-2 variants. It is proposed that the N764K, N856K and N969K substitutions (all increasing spike positive charge) may enhance S1/S2 subunit interactions after proteolytic processing of the spike protein, resulting in reduced S1 shedding and improving transmission (Martin et al. 2022) Increased charge may also alter receptor interactions. In the Omicron (BA.1) spike protein, the Q493R and Q498R substitutions are predicted to allow two additional salt bridges with ACE2 receptor position 35Glu and 38Glu (McCallum et al. 2022). Indeed, looking at the timing of charge shifts in each major lineage, the changes to more positive charge accumulate later than the changes that first allow a lineage to emerge and dominate global infections. In this model, the primary spike changes are driven by immune selection and allow a new lineage to bypass existing immune responses. Once a successful new variant emerges, the large number of new infections allow selection for the accumulation of beneficial positive charge changes. The similar pattern of increased positivity of spike protein in other coronaviruses that have moved between animals and humans (OC43, 229E, Figure 4) suggest that the change in surface protein charge may be a more general phenomenon with coronaviruses and might be a useful parameter to examine when monitoring zoonosis. This study provides a framework to monitor viral evolution through changes in biochemical properties, which can be easily applied to other viruses important to public and global health. An important note, our analyses on viral spike protein biochemical properties to monitor virus evolution are not meant to replace traditional phylogenetic analyses. The observed pattern of biochemical properties changes should completement phylogenetic signals. However, in situations where there are limited sequences available to produce reliable phylogenetic signals (e.g. the 229E and OC43 viruses examined in Figure 4), this kind of analysis using virus biochemical properties from different host species would certainly help provide important information on the virus evolution, zoonosis as well as aiding the prediction of patterns of viral changes.

In conclusion, our study provides an novel analytical framework to monitor viral evolution

through changes in biochemical properties, which can be easily applied to other viruses important to

public and global health. We also showed that natural virus evolution is more complicated and may

involve multiple factors including immune selection, as well as spike protein biochemical properties.

The observation of increase of SARS-CoV-2 spike protein charge over time provides useful

information for future vaccine and therapeutic development.

Methods

Full alignments of SARS-CoV-2 genomes were obtained from GISAID (GISAID 2020) with

collection dates to 15 June 2022. All spaces in fasta IDs were removed using sed (sed -i -e 's/ /_/g'

msa_xxxx.fasta), the alignment was dealigned ("-" characters removed) and genomes were classified

using Pangolin (Áine O'Toole et al. 2020) with the most recent database updates (pangolin v4.1.1,

pangolin-data v1.11

constellations v0.1.10 and scorpio v0.3.17). The spike coding region from each genome (if present

and intact (no Ns)) was translated into protein. Features of the protein that could be quantitated from

the spike protein sequence were determined using the ProteinAnalysis functions from BioPython

(Cock et al. 2009). These features included charge at pH 7.4, Kyle and Doolittle GRAVY score

(Kyte & Doolittle 1982) (a measure of hydrophobicity), an instability index derived from dipeptide

content (Guruprasad et al. 1990), the total percent helix, fold or sheet properties of the protein and the

total fractions of individual amino acids and fractions of di-amino acids. A matrix of all spike protein

features plus collection date, and lineage was prepared and used for analysis. Similar analyses were

performed for other coronaviruses such as 229E, OC43 and MERS-CoV by retrieving all complete

genomes available from GenBank (15 June 2022). The spike protein was also extracted using the

same method as aforementioned. Additional details are provided in the figure legends. The python

code used for the analyses is available here: https://github.com/mlcotten13/SARS-CoV-

2_spike_charge .

Acknowledgements

We thank all global SARS-CoV-2 sequencing groups for the open sharing of sequence data

and to the GISAID platform and team for making these data available. We are grateful to Andrew

Rambaut, Áine O'Toole and the Pangolin team for the Pangolin typing tool and resources.

Funding

This work was supported by the UK Medical Research Council (MRC/UK Research and

Innovation) and the UK Department for International Development (DFID) under the MRC/DFID

Concordat agreement (grant agreement no. MC_PC_20010) and Wellcome Trust, UK FCDO-

Wellcome Epidemic Preparedness-Coronavirus (grant agreement no. 220977/Z/20/Z).

References

Adamczyk, Z., Batys, P., & Barbasz, J. (2021). 'SARS-CoV-2 virion physicochemical characteristics
pertinent to abiotic substrate attachment', *Current Opinion in Colloid & Interface Science*, 55:
101466. DOI: 10.1016/j.cocis.2021.101466

Áine O'Toole et al. (2020). 'Phylogenetic Assignment of Named Global Outbreak LINeages
(PANGOLIN)',.

Assiri, A., McGeer, A., Perl, T. M., Price, C. S., Al Rabeeah, A. A., Cummings, D. A. T., Alabdullatif, Z.
N., et al. (2013). 'Hospital Outbreak of Middle East Respiratory Syndrome Coronavirus', *New*
England Journal of Medicine, 369/5: 407–16. DOI: 10.1056/NEJMoa1306742

Byrd-Leotis, L., Lasanajak, Y., Bowen, T., Baker, K., Song, X., Suthar, M. S., Cummings, R. D., et al.
(2021). 'SARS-CoV-2 and other coronaviruses bind to phosphorylated glycans from the
human lung', *Virology*, 562: 142–8. DOI: 10.1016/j.virol.2021.07.012

Cao, Y., Wang, J., Jian, F., Xiao, T., Song, W., Yisimayi, A., Huang, W., et al. (2022). 'Omicron
escapes the majority of existing SARS-CoV-2 neutralizing antibodies', *Nature*, 602/7898:
657–63. DOI: 10.1038/s41586-021-04385-3

Chan, K.-F., Koukouravas, S., Yeo, J. Y., Koh, D. W.-S., & Gan, S. K.-E. (2020). 'Probability of
change in life: Amino acid changes in single nucleotide substitutions', *Biosystems*, 193–194:
104135. DOI: 10.1016/j.biosystems.2020.104135

Cock, P. J. A., Antao, T., Chang, J. T., Chapman, B. A., Cox, C. J., Dalke, A., Friedberg, I., et al.
(2009). 'Biopython: freely available Python tools for computational molecular biology and
bioinformatics', *Bioinformatics*, 25/11: 1422–3. DOI: 10.1093/bioinformatics/btp163

Corman, Victor M., Eckerle, I., Memish, Z. A., Liljander, A. M., Dijkman, R., Jonsdottir, H., Juma
Ngeiywa, K. J. Z., et al. (2016). 'Link of a ubiquitous human coronavirus to dromedary
camels', *Proceedings of the National Academy of Sciences*, 113/35: 9864–9. DOI:
10.1073/pnas.1604472113

Corman, Victor Max, Baldwin, H. J., Tateno, A. F., Zerbinati, R. M., Annan, A., Owusu, M., Nkrumah,
E. E., et al. (2015). 'Evidence for an Ancestral Association of Human Coronavirus 229E with
Bats', (S. Schultz-Cherry, Ed.)*Journal of Virology*, 89/23: 11858–70. DOI: 10.1128/JVI.01755-
15

Cotten, M., Watson, S. J., Kellam, P., Al-Rabeeah, A. A., Makhdoom, H. Q., Assiri, A., Al-Tawfiq, J.
A., et al. (2013). 'Transmission and evolution of the Middle East respiratory syndrome
coronavirus in Saudi Arabia: a descriptive genomic study', *Lancet (London, England)*,
382/9909: 1993–2002. DOI: 10.1016/S0140-6736(13)61887-5

Cotten, M., Watson, S. J., Zumla, A. I., Makhdoom, H. Q., Palser, A. L., Ong, S. H., Al Rabeeah, A.
A., et al. (2014). 'Spread, circulation, and evolution of the Middle East respiratory syndrome
coronavirus', *mBio*, 5/1: e01062-13. DOI: 10.1128/mBio.01062-13

Creighton, T. E. (2002). *Proteins: structures and molecular properties.*, 2. ed., [Nachdr.]. New York:
Freeman.

DeGrace, M. M., Ghedin, E., Frieman, M. B., Krammer, F., Grifoni, A., Alisoltani, A., Alter, G., et al.
(2022). 'Defining the risk of SARS-CoV-2 variants on immune protection', *Nature*, 605/7911:
640–52. DOI: 10.1038/s41586-022-04690-5

Dejnirattisai, W., Huo, J., Zhou, D., Zahradník, J., Supasa, P., Liu, C., Duyvesteyn, H. M. E., et al.
(2022). 'SARS-CoV-2 Omicron-B.1.1.529 leads to widespread escape from neutralizing
antibody responses', *Cell*, 185/3: 467-484.e15. DOI: 10.1016/j.cell.2021.12.046

GISAID. (2020). 'The GISAID Initiative',.

Greaney, A. J., Loes, A. N., Crawford, K. H. D., Starr, T. N., Malone, K. D., Chu, H. Y., & Bloom, J. D.
(2021). 'Comprehensive mapping of mutations in the SARS-CoV-2 receptor-binding domain
that affect recognition by polyclonal human plasma antibodies', *Cell Host & Microbe*, 29/3:
463-476.e6. DOI: 10.1016/j.chom.2021.02.003

Greaney, A. J., Starr, T. N., Barnes, C. O., Weisblum, Y., Schmidt, F., Caskey, M., Gaebler, C., et al.
(2021). 'Mapping mutations to the SARS-CoV-2 RBD that escape binding by different classes
of antibodies', *Nature Communications*, 12/1: 4196. DOI: 10.1038/s41467-021-24435-8

Greaney, A. J., Starr, T. N., & Bloom, J. D. (2022). 'An antibody-escape estimator for mutations to the
SARS-CoV-2 receptor-binding domain', *Virus Evolution*, 8/1: veac021. DOI:
10.1093/ve/veac021
Guruprasad, K., Reddy, B. V. B., & Pandit, M. W. (1990). 'Correlation between stability of a protein
and its dipeptide composition: a novel approach for predicting *in vivo* stability of a protein from
its primary sequence', *Protein Engineering, Design and Selection*, 4/2: 155–61. DOI:
10.1093/protein/4.2.155
Kim, S. Y., Jin, W., Sood, A., Montgomery, D. W., Grant, O. C., Fuster, M. M., Fu, L., et al. (2020).
'Characterization of heparin and severe acute respiratory syndrome-related coronavirus 2
(SARS-CoV-2) spike glycoprotein binding interactions', *Antiviral Research*, 181: 104873. DOI:
10.1016/j.antiviral.2020.104873
Kyte, J., & Doolittle, R. F. (1982). 'A simple method for displaying the hydropathic character of a
protein', *Journal of Molecular Biology*, 157/1: 105–32. DOI: 10.1016/0022-2836(82)90515-0
Liu, Y., Liu, J., Johnson, B. A., Xia, H., Ku, Z., Schindewolf, C., Widen, S. G., et al. (2022). 'Delta
spike P681R mutation enhances SARS-CoV-2 fitness over Alpha variant', *Cell Reports*, 39/7:
110829. DOI: 10.1016/j.celrep.2022.110829
Livingstone, C. D., & Barton, G. J. (1993). 'Protein sequence alignments: a strategy for the
hierarchical analysis of residue conservation', *Bioinformatics*, 9/6: 745–56. DOI:
10.1093/bioinformatics/9.6.745
Lubinski, B., Frazier, L. E., Phan, M. V. T., Bugembe, D. L., Cunningham, J. L., Tang, T., Daniel, S.,
et al. (2022). 'Spike Protein Cleavage-Activation in the Context of the SARS-CoV-2 P681R
Mutation: an Analysis from Its First Appearance in Lineage A.23.1 Identified in Uganda',
Microbiology Spectrum, e0151422. DOI: 10.1128/spectrum.01514-22
Martin, D. P., Lytras, S., Lucaci, A. G., Maier, W., Grüning, B., Shank, S. D., Weaver, S., et al. (2022).
'Selection analysis identifies clusters of unusual mutational changes in Omicron lineage BA.1
that likely impact Spike function', (K. Crandall, Ed.)*Molecular Biology and Evolution*, msac061.
DOI: 10.1093/molbev/msac061
McCallum, M., Czudnochowski, N., Rosen, L. E., Zepeda, S. K., Bowen, J. E., Walls, A. C., Hauser,
K., et al. (2022). 'Structural basis of SARS-CoV-2 Omicron immune evasion and receptor
engagement', *Science*, 375/6583: 864–8. DOI: 10.1126/science.abn8652
Memish, Z. A., Cotten, M., Meyer, B., Watson, S. J., Alsahafi, A. J., Al Rabeeah, A. A., Corman, V. M.,
et al. (2014). 'Human infection with MERS coronavirus after exposure to infected camels,
Saudi Arabia, 2013', *Emerging Infectious Diseases*, 20/6: 1012–5. DOI:
10.3201/eid2006.140402
Nguyen, L., McCord, K. A., Bui, D. T., Bouwman, K. M., Kitova, E. N., Elaish, M., Kumawat, D., et al.
(2022). 'Sialic acid-containing glycolipids mediate binding and viral entry of SARS-CoV-2',
Nature Chemical Biology, 18/1: 81–90. DOI: 10.1038/s41589-021-00924-1
Nie, C., Sahoo, A. K., Netz, R. R., Herrmann, A., Ballauff, M., & Haag, R. (2022). 'Charge Matters:
Mutations in Omicron Variant Favor Binding to Cells', *ChemBioChem*. DOI:
10.1002/cbic.202100681
Obermeyer, F., Jankowiak, M., Barkas, N., Schaffner, S. F., Pyle, J. D., Yurkovetskiy, L., Bosso, M.,
et al. (2022). 'Analysis of 6.4 million SARS-CoV-2 genomes identifies mutations associated
with fitness', *medRxiv: The Preprint Server for Health Sciences*, 2021.09.07.21263228. DOI:
10.1101/2021.09.07.21263228
Pawłowski, P. (2021). 'SARS-CoV-2 variant Omicron (B.1.1.529) is in a rising trend of mutations
increasing the positive electric charge in crucial regions of spike protein S.', *Acta Biochimica
Polonica*. DOI: 10.18388/abp.2020_6072
Pedregosa, F. and Varoquaux, G. and Gramfort, A. and Michel, V., and Thirion, B. and Grisel, O. and
Blondel, M. and Prettenhofer, P., and Weiss, R. and Dubourg, V. and Vanderplas, J. and
Passos, A. and, & Cournapeau, D. and Brucher, M. and Perrot, M. and Duchesnay, E. (2011).
'Scikit-learn: Machine Learning in Python', *Journal of Machine Learning Research*, 12: 2825–
30.
Peiris, M., & Perlman, S. (2022). 'Unresolved questions in the zoonotic transmission of MERS',
Current Opinion in Virology, 52: 258–64. DOI: 10.1016/j.coviro.2021.12.013

Sabir, J. S. M., Lam, T. T.-Y., Ahmed, M. M. M., Li, L., Shen, Y., E. M. Abo-Aba, S., Qureshi, M. I., et
al. (2016). 'Co-circulation of three camel coronavirus species and recombination of MERS-
CoVs in Saudi Arabia', *Science*, 351/6268: 81–4. DOI: 10.1126/science.aac8608
So, R. T. Y., Chu, D. K. W., Miguel, E., Perera, R. A. P. M., Oladipo, J. O., Fassi-Fihri, O., Aylet, G., et
al. (2019). 'Diversity of Dromedary Camel Coronavirus HKU23 in African Camels Revealed
Multiple Recombination Events among Closely Related Betacoronaviruses of the Subgenus
Embecovirus', *Journal of Virology*, 93/23: e01236-19. DOI: 10.1128/JVI.01236-19
Tao, Y., Shi, M., Chommanard, C., Queen, K., Zhang, J., Markotter, W., Kuzmin, I. V., et al. (2017).
'Surveillance of Bat Coronaviruses in Kenya Identifies Relatives of Human Coronaviruses
NL63 and 229E and Their Recombination History', (S. Perlman, Ed.)*Journal of Virology*, 91/5:
e01953-16. DOI: 10.1128/JVI.01953-16
Tzou, P. L., Tao, K., Pond, S. L. K., & Shafer, R. W. (2022). 'Coronavirus Resistance Database (CoV-
RDB): SARS-CoV-2 susceptibility to monoclonal antibodies, convalescent plasma, and
plasma from vaccinated persons', (J. Bhattacharya, Ed.)*PLOS ONE*, 17/3: e0261045. DOI:
10.1371/journal.pone.0261045
Vijgen, L., Keyaerts, E., Lemey, P., Maes, P., Van Reeth, K., Nauwynck, H., Pensaert, M., et al.
(2006). 'Evolutionary history of the closely related group 2 coronaviruses: porcine
hemagglutinating encephalomyelitis virus, bovine coronavirus, and human coronavirus
OC43', *Journal of Virology*, 80/14: 7270–4. DOI: 10.1128/JVI.02675-05
Vijgen, L., Keyaerts, E., Moës, E., Thoelen, I., Wollants, E., Lemey, P., Vandamme, A.-M., et al.
(2005). 'Complete genomic sequence of human coronavirus OC43: molecular clock analysis
suggests a relatively recent zoonotic coronavirus transmission event', *Journal of Virology*,
79/3: 1595–604. DOI: 10.1128/JVI.79.3.1595-1604.2005
Zhou, Z., Hui, K. P. Y., So, R. T. Y., Lv, H., Perera, R. A. P. M., Chu, D. K. W., Gelaye, E., et al.
(2021). 'Phenotypic and genetic characterization of MERS coronaviruses from Africa to
understand their zoonotic potential', *Proceedings of the National Academy of Sciences of the
United States of America*, 118/25: e2103984118. DOI: 10.1073/pnas.2103984118
480
481

Supplementary Figure 1. Location of charged amino acid changes in the spike protein. The spike protein sequences encoded by the first 300 reported genomes for the indicated SARS-CoV-2 lineages were collected, and charged amino acid changes from the original B lineage spike sequence were plotted. Charge changes were colored from dark blue (change from positive to negative charged amino acid (AA)), blue change from neutral to negative charged AA), orange (change from neutral to positive charged AA) and red (change from negative to positive charged AA). Substitutions are indicated by original AA/position in reference sequence spike/novel AA. The GenBank NC_045512 genome was used as reference. Key spike protein features of the SARS-CoV-2 spike protein are shown in the upper panel of the figure.