Modern human expansion from (southern?) Africa

Zarus Cenac

Department of Psychology, City, University of London, Rhind Building, St John Street, London, United Kingdom, EC1R 0JD
Email: zarus.cenac@city.ac.uk

Abstract

Previous research favours the idea that modern humans spread worldwide from Africa. For instance, autosomal diversity falls as geographical distance from Africa rises, and, through autosomal diversity, an area of origin for this worldwide expansion is indicated to be within Africa entirely. Declines are also known to happen for mitochondrial, X-chromosomal, and Y-chromosomal diversities. What was not known is if those diversities signal an area of origin which is totally in Africa. Additionally, there is disagreement regarding where in Africa the origin is. Considered in the present research was whether the expansion is suggested through various genetic diversities (including mitochondrial, X-chromosomal, and Y-chromosomal) and cranial traits (e.g., cranial size dimorphism), and where in Africa the expansion started. For each genetic or cranial variable, it was seen if the estimated area of origin was solely in Africa. To generally estimate where the expansion began in Africa, a centroid was calculated from potential origins that were obtained through indicators of the expansion. The area of origin was completely within Africa for each variable except one – for Y-chromosomal diversity, the area was possibly in Asia only. The centroid of the potential origins was in southern Africa. However, the support for a southern African origin (over elsewhere in Africa) could potentially reflect variation in non-African ancestry within Africa rather than the expansion. Nevertheless, a southern African origin was pointed to via the autosomal diversity of sub-Saharan African populations which had been adjusted for non-African ancestry. Southern Africa possible is the origin of the expansion.

Introduction

Findings in the literature align with modern humans having journeyed across the world via an expansion that originated in Africa (Manica et al., 2007; Ramachandran et al., 2005; see Henn, Cavalli-Sforza, & Feldman, 2012, for a review). Research agrees with population bottlenecks having been encountered during the expansion (more and more bottlenecks with greater and greater distance from Africa) thereby leading to a reduction in autosomal microsatellite diversity (expected heterozygosity) as the expansion reached greater distances (Prugnolle et al., 2005). Not only does the diversity of autosomal microsatellites decrease as geographical distance from Africa widens (Prugnolle et al., 2005), but declines are also present in other aspects of biological diversity, such as

diversity in autosomal single nucleotide polymorphisms (SNPs) (Balloux et al., 2009) and cranial shape (von Cramon-Taubadel & Lycett, 2008).

Through measuring the fit between geographical distance (the distance from geographical locations to populations) and a biological variable (e.g., the autosomal microsatellite diversity of populations), it is possible to estimate a geographical area out of which the expansion started (Manica et al., 2007). This area contains the location which gives the best fit, and locations which are similar in fit to the best (e.g., Manica et al., 2007) – the area can be called a *geographical area of best fit* (see Figure 1A for an example). More than one geographical area of best fit may be found with a biological variable (as in the present research with mitochondrial diversity, and also with cranial shape diversity for female crania). For instance, one geographical area of best fit may have correlation coefficients (for relationships between distance and diversity) that are negative, whereas another geographical area of best fit has positive coefficients (see *Results and discussion*).[1] Results in prior research have been in alignment with types of biological diversity declining as distance increases from the origin of the expansion (e.g., Ramachandran et al., 2005; von Cramon-Taubadel & Lycett, 2008), so the geographical area of best fit which has negative correlation coefficients (rather than the one with positive coefficients) would be favoured for being the area where the expansion has its origin.

Figure 1

Autosomal SNP Diversity Suggests the Worldwide Expansion Originated in Africa

Note. A map of Africa is presented in Figure 1A. Whilst autosomal SNP haplotype diversity has been found to decline as distance from Africa increases (Balloux et al., 2009), an origin area was not known to have been estimated when using this type of diversity. For Figure 1A, the area was estimated (see *Method* for details) and is the darker grey area. The lowest Bayesian information criterion (BIC) regarding the origins used was in Niger (15°N, 5°E).

Skeletal

Variables which appear to mark the expansion from Africa i) relate to distance from Africa (e.g., Prugnolle et al., 2005; von Cramon-Taubadel & Lycett, 2008). Moreover, like autosomal microsatellite diversity (Manica et al., 2007), variables which indicate the expansion should ii) lead to

an estimated origin area entirely within Africa. Ideally, an indicator of the expansion should meet both of these points (i.e., association with distance from Africa, and the estimated area being in Africa alone). Of the three studies on skeletal diversity which have indicated origin areas (Betti et al., 2009, 2013; Manica et al., 2007), only cranial form diversity amongst males (e.g., using 105 populations [n = 4,666]) has fulfilled both points (Betti et al., 2009; Manica et al., 2007) – the area is fully within Africa when adjusting the cranial diversity of males for climate (Manica et al., 2007) and without amending for climate (Betti et al., 2009) (see Manica et al., 2007, and Betti et al., 2009, for methodological differences between the two studies). Female cranial form diversity (Betti et al., 2009) and hip bone shape diversity for males and females have met point one but not point two (Betti et al., 2013). With female cranial diversity in form (39 populations [n = 1,579]), the origin area is not in Africa alone (Betti et al., 2009). Indeed, for females, the origin area is wider than it is for males, although the sample size for females was not large in that study (Betti et al., 2009). Regarding hip bone shape diversity, the origin area is not exclusive to Africa for males (27 populations [n = 891]) or females (20 populations [n = 552]) (Betti et al., 2013). Other studies on whether distance from Africa is related to types of skeletal diversity (e.g., Betti et al., 2012), such as cranial shape diversity (von Cramon-Taubadel & Lycett, 2008), have not found origin areas.

In unpublished research, cranial sexual size dimorphism (adjusted for absolute latitude) was found to attain point one, i.e., it correlated with distance from Africa (Cenac, 2022). Research was not known to have tested if it meets the second point, and therefore finding the origin area through this adjusted dimorphism is something which could be explored (Cenac, 2022).

Genetic diversity

As geographical distance from Africa grows, the diversity (heterozygosity) of autosomal microsatellites decreases (Prugnolle et al., 2005). Using autosomal microsatellite heterozygosity, the area of origin is estimated to solely be within Africa (Manica et al., 2007). Aside from autosomal diversity, does any other type of genetic diversity i) decline with rising distance from Africa (Balloux et al., 2009; Shi et al., 2010) and ii) result in Africa completely having the estimated origin area? Balloux et al. (2009) found declines (as distance from Africa increases) in X-chromosomal microsatellite diversity, Y-chromosomal microsatellite diversity, and mitochondrial diversity. Figure 3E in Balloux et al. (2009) presents Y-chromosomal microsatellite diversities for populations at distances from Africa. That figure may indicate that this Y-chromosomal diversity is not at its highest in Africa. Indeed, when diversities are grouped by continent and averaged, the numerically highest diversity is not found to be with Africa, but with Asia (Table 1). When research on diversity or/and expansion (Manica et al., 2007; Ramachandran et al., 2005; Romero et al., 2009) is considered, one would expect Y-chromosomal diversity to numerically be greater in Africa than anywhere else if Y-chromosomal diversity is a reflection of the expansion from Africa. Therefore, Table 1 seems at odds with Y-chromosomal diversity actually indicating the *worldwide expansion*.[2] Given Table 1, it would

not be surprising if Y-chromosomal microsatellite diversity leads to an origin area which is outside of Africa. Elsewhere, in Chiaroni et al. (2009), there was a negative gradient regarding the Y-chromosomal haplotype diversity of populations and distance from Africa. Yet Figure S6 of that study (i.e., Chiaroni et al., 2009) may suggest that this diversity is not at its greatest in Africa.

Table 1

Average Y-Chromosomal Microsatellite Diversity

Continent	Y-Chromosomal Diversity
Africa	0.42
Asia	0.46
Europe	0.44
North America	0.37
Oceania	0.39
South America	0.19

Note. Using diversities within each of 51 populations in Balloux et al. (2009), averages (means) were calculated by continent. Also, see estimates of Y-chromosomal diversity elsewhere (e.g., Hammer et al., 2001; Seielstad et al., 1999), for instance, in Jorde et al. (2000), microsatellite diversity was numerically higher in Africa than in Asia or Europe (other continents were not featured).

Balloux et al. (2009) did not explore where origin areas are located regarding genetic diversities; they did not determine if mitochondrial, and X- and Y-chromosomal diversities suggest an origin area entirely within Africa. Other studies have occurred concerning distance from Africa and the diversity of the X chromosome (Li et al., 2008) and Y chromosome (Chiaroni et al., 2009; Shi et al., 2010), regarding X-chromosomal haplotypes (Li et al., 2008), Y-chromosomal haplotypes (Chiaroni et al., 2009), and Y-chromosomal haplogroups (Shi et al., 2010). However, these studies also did not estimate origin areas from X- and Y-chromosomal diversities. Consequently, it was of interest to estimate origin areas through X-chromosomal, Y-chromosomal, and mitochondrial diversities.

Expansion origin

Where in Africa did the expansion originate? Inferring from previous research/reasoning (e.g., Henn, Cavalli-Sforza, & Feldman, 2012; Manica et al., 2007; Ray et al., 2005), the answer is not clear. For instance, eastern Africa has been suggested based on autosomal microsatellites (Ray et al., 2005), whereas southern Africa was favoured in a review of the literature (Henn, Cavalli-Sforza, & Feldman, 2012).[3] Figure 1 of Batini and Jobling (2011) summarised some findings in research concerning human origins onto two maps of mainland Africa. This led to an idea, in the current analyses, of gathering coordinates at which relationships between geographical distance and

indicators of the expansion from Africa are at their strongest in Africa (such coordinates are called *peak points* in the present research), and showing their locations through a representation of mainland Africa. It could be seen if peak points (for different types of indicators of the expansion) generally cluster in any particular region(s); a clustering in a region would support the expansion having its origin in that region. Moreover, a centroid could be calculated from the peak points (found with indicators) as a general approximation of where the expansion started. However, the centroid may not necessarily be useful. For instance, if the peak points were not clustered in a region, but dispersed across Africa, a centroid might not be indicative of which region(s) the expansion originated in. Therefore, clustering and the centroid would both be of interest.

Analyses

In the present research, something was defined as being an indicator of the expansion if it met the two points raised earlier, i.e., i) it is related to distance from Africa, and ii) it produces an origin area only in Africa. The present analyses estimated where in Africa the worldwide expansion began. It was established if origin areas are completely in Africa for diversity in cranial shape, mitochondrial DNA, X-chromosomal microsatellites, and Y-chromosomal microsatellites. Given that these cranial/genetic diversities do have negative correlations with distance from Africa (Balloux et al., 2009; von Cramon-Taubadel & Lycett, 2008), the expansion from Africa would be signalled by these diversities if the estimated origin areas are only within Africa. Additionally, more of an insight was sought into whether cranial form diversity is indicative of the expansion from Africa. Following on from previous research (Cenac, 2022), analyses found whether cranial size dimorphism (whether adjusted for absolute latitude or not) indicates an origin area which is in Africa only. After finding whether certain cranial or genetic variables are indicative of the expansion from Africa, it was ascertained whether peak points (found in the current analyses and previous studies) were clustered in any region of Africa. Furthermore, a centroid (i.e., as an overall estimate of the origin) was determined from peak points. Later, in *Results and discussion*, some potential caveats are considered. For instance, extrapolating from previous research regarding admixture (e.g., Haber et al., 2016; Pickrell & Reich, 2014) and phonemic diversity (Van Tuyl & Pereltsvaig, 2012), the non-African ancestry in Africa could have affected where origin areas and peak points suggest the worldwide expansion originated. As a step to addressing this, additional analyses occurred. The autosomal microsatellite heterozygosity of 106 sub-Saharan African populations (Pemberton et al., 2013) was adjusted for non-African ancestry – it was seen where in Africa this adjusted heterozygosity suggests the origin area and peak point are, along with whether geographical distance from the peak point correlates with adjusted heterozygosity.

Method

Analyses of data, and calculations more generally, were through R Versions 4.0.3 (R Core Team, 2020) and 4.0.5 (R Core Team, 2021), and Microsoft Excel. A summary of some types of outcomes in the present research and previous studies is presented in Table 2.

Table 2

Summary

Cranial/genetic variable	Association with distance from Africa?	Peak point coordinates stated	Origin area in Africa alone?
Cranial form diversity	Betti et al. (2009), present research		Betti et al. (2009), Manica et al. (2007), present research
Cranial shape diversity	von Cramon-Taubadel and Lycett (2008), present research	Present research	Present research
Cranial size dimorphism	Cenac (2022), present research	Present research	Present research
Autosomal microsatellite heterozygosity	Balloux et al. (2009), Prugnolle et al. (2005), present research	Ramachandran et al. (2005), present research	Manica et al. (2007)
Autosomal microsatellite repeat length diversity	Tishkoff et al. (2009)	Tishkoff et al. (2009)	Tishkoff et al. (2009)
Autosomal SNP heterozygosity	Balloux et al. (2009)	Present research	Present research
Mitochondrial diversity	Balloux et al. (2009)	Present research	Present research
X-chromosomal microsatellite diversity	Balloux et al. (2009)	Present research	Present research
Y-chromosomal microsatellite diversity	Balloux et al. (2009)		Present research

Note. The table is filled in without respect to whether or not cranial/genetic variables were adjusted for another variable (see main text and references, e.g., Manica et al., 2007, adjusted cranial form diversity for climate) or if populations were across the globe or only in Africa. The second column lists research which used *p*-values.

Cranial

The cranial size dimorphism of Holocene modern humans (and dimorphism within cranial dimensions), which had been calculated from measurements in the Howells (1973, 1989, 1995, 1996) dataset (Cenac, 2022), was used in the present analyses. Measurements (56 dimensions) were from 26

populations ($n_{\text{males}} = 1{,}256$; $n_{\text{females}} = 1{,}156$) (Howells, 1989, 1996), and the calculation of size dimorphism (from geometric means) had utilised a natural log ratio in Smith (1999), i.e., ln[(male size)/(female size)] in the prior study (Cenac, 2022). The cranial size dimorphism (calculated using the Howells data) had previously been adjusted for absolute latitude (Cenac, 2022), and that adjusted dimorphism was employed in the present research.

Analyses of cranial diversity also featured the Howells (1973, 1989, 1995, 1996) data (so did Cenac, 2022). Specifically, male crania from 28 populations ($n = 1{,}348$) and female crania from the 26 populations (Howells, 1989, 1996) were used and the cranial dimensions were the same 56 used in the dimorphism analysis of Cenac (2022). Cranial form diversity can be quantified by i) standardising each dimension by way of a z-score transformation, ii) calculating the variance of each of the standardised dimensions for each population, and iii) taking a mean from the variances of all those standardised dimensions (i.e., a mean variance) per population (e.g., Betti et al., 2009). In the present analyses, this method was used on female crania from the Howells dataset. Through that method, the mean form diversity of male crania and the mean shape diversity of female crania had been calculated beforehand from the Howells data (Cenac, 2022) – these diversities were used in the current analyses. In a previous study, cranial shape diversity was determined amongst male crania in the Howells data (28 populations) by calculating the geometric mean of each cranium, dividing the cranial measurements within a cranium by the geometric mean, and then calculating the mean variances after standardising dimensions (von Cramon-Taubadel & Lycett, 2008). For the current analyses, this technique was used to calculate cranial shape diversity with the same data as that previous study (i.e., male crania in the Howells data). From preceding research, the coefficient (r) for the correlation between distance from Africa and the cranial shape diversity of females in the Howells data (Cenac, 2022) was used. Cenac (2022) used the Howells dataset obtained through http://web.utk.edu/~auerbach/HOWL.htm, and so did the present research.[4]

Genetic

The mitochondrial diversity of 51 populations, featured in the supplementary materials of Balloux et al. (2009), was utilised. Balloux et al. (2009) sourced genetic data from GenBank and (citing Ingman & Gyllensten, 2006) the Human Mitochondrial Genome Database (mtDB); for details on either GenBank or the mtDB, see Benson et al. (2013) and Ingman and Gyllensten (2006) respectively. The mitochondrial diversity was from the complete mitochondrial genome minus hypervariable segments I and II (Balloux et al., 2009). Additionally, Balloux et al. (2009) did present mitochondrial diversity from hypervariable segment I (of 109 populations in HVRBase++). In the present analyses, for representativeness of mitochondrial diversity in general, the diversity from the complete genome (minus the two segments) was used rather than diversity from that hypervariable segment. Diversity (expected heterozygosity) was found in the supplementary material of Balloux et al. (2009) for X- and Y-chromosomal microsatellites, and autosomal SNP haplotypes of 51

populations, and these diversities were used in the present research. Balloux et al. (2009) calculated X- and Y-chromosomal microsatellite heterozygosities using the HGDP-CEPH Human Genome Diversity Cell Line Panel. Balloux et al. (2009) found SNP haplotype heterozygosity in Li et al. (2008); Li et al. (2008) calculated mean heterozygosity from the panel. Information on the panel can be found, for instance, in Cann et al. (2002). From summing population sample sizes in Balloux et al. (2009), N is 633 for mitochondrial and 963 for autosomal and regarding the allosomes.

Geographical distances

As for geographical distances, ones calculated beforehand (Cenac, 2022) were used in the present research. Also used were additional distances which were calculated through a means featured by Williams (2011) on http://edwilliams.org/avform.htm. Coordinates for locations in Africa (Betti et al., 2013), coordinates offered for populations regarding the genetic data (Balloux et al., 2009), and coordinates presented elsewhere, including waypoints (Cenac, 2022; von Cramon-Taubadel & Lycett, 2008), were used in the geographical distance calculations. Previous research has used waypoints when calculating geographical distances (e.g., von Cramon-Taubadel & Lycett, 2008) – waypoints were used to calculate distances in the present research. Regarding the waypoints, four of five waypoints listed in a previous study (von Cramon-Taubadel & Lycett, 2008) were featured in calculations like (with the finalised distances) in Cenac (2022); an alternative waypoint, as in Cenac (2022), was made use of rather than the Cairo waypoint which von Cramon-Taubadel and Lycett (2008) used. This alternative waypoint shares the latitude of that Cairo waypoint, but is further east (Cenac, 2022). A distance of 0 km was used for the distance between a population and itself. Geographical maps (Bartholomew illustrated reference atlas of the world, 1985), including one with representations of locations for populations in the HGDP-CEPH data (López Herráez et al., 2009), were used regarding the calculation of geographical distances. Maps (Bartholomew illustrated reference atlas of the world, 1985) were also utilised for determining which country/region peak points or centroids are located in.

Origins

A number of *origin* coordinates inside/outside of Africa (Betti et al., 2013; von Cramon-Taubadel & Lycett, 2008) were used in the present research. Coordinate pairs were used, in previous research, as *origins* in Africa – 99 in Betti et al. (2013) and three in von Cramon-Taubadel and Lycett (2008) for instance; the present research utilised those 99 coordinate pairs in Betti et al. (2013) and one of the three (Botswana) used by von Cramon-Taubadel and Lycett (2008) as origins too. In von Cramon-Taubadel and Lycett (2008), three origins outside of Africa were used, and these three featured as origins in the present research. Coordinates applied to populations in the Howells data (von Cramon-Taubadel & Lycett, 2008) were also employed in the present research as origins like in Cenac (2022). A number of origin coordinates (Betti et al., 2013) are shown/stated in the present research in figures/text (e.g., the note of Figure 1, Figure 8).

Correlation

Correlation tests in the present analyses were two-tailed. When seeing if a variable is associated with geographical distance, previous studies (e.g., Atkinson, 2011; Betti et al., 2013) have employed different methods for choosing which geographical location the population distances are from. For instance, Betti et al. (2013) used the centroid of the origin area. Atkinson (2011) utilised the lowest BIC; in a number of analyses in the present research, the lowest BIC was used. Indeed, if the lowest BIC was found to be at a location outside of Africa for a diversity (and the location gave a negative *r* correlation coefficient), a correlation test (through the cor.test R function) was to be run between diversity and the distance from the origin which gave that lowest BIC. Like previously (Cenac, 2022), Botswana functioned as the origin in Africa for correlation tests regarding the cranium.

Origin area

After prior research (e.g., Betti et al., 2009), the BIC was the metric for finding which geographical origin gives the peak relationship between variables, and for deciding on the other geographical origins which are just as good as the peak. BICs regarding adjusted cranial size dimorphism were found using a formula in Masson (2011). For each variable, the origin giving the lowest (and therefore strongest) BIC was found, and locations within four BICs of this value were defined as being equally as good a fit (e.g., Manica et al., 2007) for the relationship of geographical distance and the variables (diversities and unadjusted/adjusted dimorphism). However, as alluded to in the *Introduction*, if there were competing geographical areas of best fit for a type of diversity or unadjusted/adjusted dimorphism, correlation coefficients were considered when defining the origin area.

Betti et al. (2013) used 99 coordinate pairs alongside others to present two maps of Africa and beyond. Each showed coordinates included and not included in the origin area and peak points regarding the hip bone shape diversity of males (one graph) and females (the other graph). Following Betti et al. (2013) (who used origins within and outside of Africa), 99 coordinate pairs (found in Betti et al., 2013) used as origins in Africa were employed to display maps of Africa, with it being indicated whether origins were, or were not, in the estimated origin area (Figures 1A, 2A, 2C, 2E, 2G, 3B, 4C, 5A, 5C, 8A, 8B, S3A, S5B, and S5D). Moreover, in the present research, those 99 coordinate pairs were used to make maps of Africa that had peak points (Figures 6, 7, 8A, 8B, S5B, and S5D). Regarding the 99 *origin* coordinate pairs (in Africa) from Betti et al. (2013) which were used in the present research – Betti et al. (2013) utilised coordinates at 5° intervals, e.g., (30°S, 30°E), (25°S, 30°E) etc.

BICs were calculated through R, except regarding Figures 8B, S3, S5B, and S5D, and Text S1; when R was not used, BICs were arrived at through Microsoft Excel using a formula in Masson (2011) which allows for BIC to be calculated through R^2. Yet, when utilising the formula, R^2 was not used in the present research (except for some analyses in Text S1). For instance, when BICs were

calculated from adjusted cranial size dimorphism, squared semi-partial Pearson correlation coefficients were used instead of R^2.

Graphs

The symbols used for the continents of populations (e.g., Figure 1B) continued, or were adapted, from such symbols used in Cenac (2022). Indeed, graphs have extents of similarity to ones in Cenac (2022) in terms of design/axes/labelling, for instance, BIC graphs (e.g., Figures 1B, 2B) and correlation coefficient graphs (e.g., Figures 4B, S2B). Hunley et al. (2012) presented correlation coefficients for diversity and distance (*y*-axis) against geographical regions (*x*-axis). Geographical regions consisted of locations used as origins (but with one *origin* location being used in the Australia category) (Hunley et al., 2012). Similarly, Cenac (2022) presented coefficients (*y*-axis), but against the geographical distance between origins and some location (*x*-axis). For certain graphs (Figures 1B, 2B, 2D, 2F, 2H, 3C, 4A, 4B, 5B, 5D, S2A-F, S3B, S4A-B), following Cenac (2022), associations between distance and a variable were plotted on a graph (or, unlike Cenac, 2022, BICs were represented) when each one of 32 geographical coordinate pairs was utilised as an origin. The same *origin* locations (32) used in Cenac (2022) were featured in these graphs. Those 32 coordinate pairs used in Cenac (2022) were from von Cramon-Taubadel and Lycett (2008) for Beijing, Delhi, Tel Aviv, 28 populations in the Howells dataset, and Botswana. The Botswana coordinates are in the north of Botswana (Cenac, 2022). Cenac (2022) constructed graphs using *r*-, *sr*-, and sr_s- values; the present research used *r*-values, but also BICs. Indeed, the present research graphed correlation coefficients against the distances of origins (Figures 4B, S2, S4), and presented BICs against distances of origins (Figures 1B, 2B, 2D, 2F, 2H, 3C, 4A, 5B, 5D, S3B).

Positive spatial autocorrelation

Regarding spatial autocorrelation, the three-step method can be used in a Microsoft Excel spreadsheet to find a spatial Durbin-Watson (spatial *DW*) (Chen, 2016) upon which the Durbin-Watson bounds (Savin & White, 1977) can be utilised (Chen, 2016). This method, through the spreadsheet, and bounds were used to check for positive spatial autocorrelation in the correlation analyses regarding the cranium and the Y chromosome. Bounds at the 5% significance level (Savin & White, 1977) were looked to. Within the Chen (2016) spreadsheet is a graph. In any instance of positive spatial autocorrelation, or uncertainty with regard to if such autocorrelation occurred, the graph (i.e., with data in the current analyses put into the spreadsheet) was examined to see which datapoints may be the cause.

Origin centroid

Locations which lead to the strongest relationships numerically (out of locations in Africa) between geographical distance and diversity/dimorphism (using populations within and outside of Africa), whether calculated through *r*-values (Ramachandran et al., 2005) or BICs, were found from previous research or calculated in the present analyses. Also, a location found through bootstrapping

(Tishkoff et al., 2009) was included. Locations were with respect to relationships between distance and i) cranial shape diversity (males, females), ii) cranial size dimorphism, iii) mitochondrial diversity (the present research), iv) autosomal diversity of microsatellites regarding heterozygosity (Ramachandran et al., 2005), repeat lengths (Tishkoff et al., 2009), and SNP heterozygosity, and v) X-chromosomal diversity (present research). These types of diversity/dimorphism each suggest an estimated origin area in Africa alone (Figures 1-5; Manica et al., 2007; Tishkoff et al., 2009). The ability of cranial form diversity to indicate the expansion from Africa is suggested to arise more from diversity in cranial shape rather than cranial size diversity (*Results and discussion*). Therefore, to reduce redundancy between origin locations, the locations with respect to cranial form diversity were not included when determining the centroid.

Using the geosphere (Version 1.5-10) package (Hijmans, 2019) in R, a midpoint and centroids were calculated: i) a midpoint was taken between the coordinates from cranial shape diversity (as there was one set of coordinates for males, and one for females), ii) a geographical centroid was calculated from the three locations from the autosomal diversities, iii) the midpoint (cranial shape diversity), the centroid (autosomal diversity), and the three other locations (cranial size dimorphism, mitochondrial diversity, and X-chromosomal diversity) were used when calculating the expansion origin centroid.

Sub-Saharan Africa and admixture

Autosomal microsatellite (mean) expected heterozygosity for each of 106 populations in Tishkoff et al. (2009) (Pemberton et al., 2013) was used. From population sample sizes in Pemberton et al. (2013), $N = 2340$. Following some previous research concerning geographical distance and genetic diversity (Pemberton et al., 2013), populations that do not have a population sample size of five or more were not featured in the present analyses. To avoid overlap with Ramachandran et al. (2005), who used autosomal microsatellite heterozygosity of HGDP-CEPH data (the origin centroid of Figure 6 was contributed towards by the peak location in Ramachandran et al., 2005), heterozygosity for the HGDP-CEPH populations was not used despite being available in Pemberton et al. (2013). Estimates of different types of ancestry (East Asian, European / Middle Eastern, Indian, Native American, and Oceanian), as proportions, (Tishkoff et al., 2009) were summed together to represent non-African ancestry.

Parametric and nonparametric analyses were conducted. BICs and correlation coefficients (r, sr, r_s, or sr_s) were found regarding geographical distances (between locations in Africa and populations) and the heterozygosity of the populations which either had or had not been adjusted for non-African ancestry (in nonparametric analysis, a rank-transformation had been applied to heterozygosity, distances, and non-African ancestry). BICs were determined in R regarding the relationship between autosomal microsatellite diversity and distance in the parametric analysis. Otherwise (e.g., when autosomal microsatellite diversity was adjusted for non-African ancestry), BICs

were calculated in Excel using a formula for BIC calculation which was presented in Masson (2011). The formula found in Masson (2011) for calculating BIC uses R^2. Instead of using R^2s, i) for the parametric analysis between adjusted heterozygosity and distance, the semi-partial Pearson correlation coefficient was squared, and ii) for the nonparametric analyses, BICs were calculated through squared Spearman correlation coefficients and squared semi-partial Spearman coefficients.

Geographical distances were found as described above, i.e., through Williams (2011). Locations in Africa were from Betti et al. (2013). Coordinates for populations were found in Tishkoff et al. (2009). From the location which had the lowest BIC (alongside a negative correlation coefficient), it was tested whether distance to populations is correlated with the unadjusted/adjusted heterozygosity.

For zero-order correlation tests, cor.test in R was used. ppcor (Kim, 2015) was employed in R regarding the tests for semi-partial correlations. It was seen whether autosomal heterozygosity (whether adjusted for non-African ancestry or not) correlates with distance from southern Africa, and if non-African ancestry (adjusted for distance from southern Africa or not) correlates with autosomal heterozygosity. As detailed above, positive spatial autocorrelation was looked for (in parametric analyses) via Chen (2016) and the Savin and White (1977) bounds. To conduct the autocorrelation analyses, interpopulation distances were established through Williams (2011), and 0 km was used i) as the distance when populations had the same coordinates and, ii) for the distance a population was from itself.

Multiple testing

Statistical significance tests were initially from two sets of analyses which were for two manuscripts; one on the cranium and mitochondrial DNA, and the other on the allosomes and the origin centroid. As a result, the Holm-Bonferroni correction was given to p-values (Holm, 1979) for each set separately. Regarding each set, a spreadsheet (Gaetano, 2013) was used to apply the correction. Any p-value calculated after the combination of manuscripts (i.e., for the correlation tests involving autosomal microsatellite heterozygosity) was added to the second set. The corrections were not used on the spatial autocorrelation tests.

<div align="center">

Results and discussion

</div>

Cranial form and shape diversities

Aligning with previous research (Betti et al., 2009), there was a negative correlation between distance from Africa and cranial form diversity for males, $r(26) = -.52$, $p = .023$, spatial $DW = 2.50$ (Figure S1A), and females $r(24) = -.49$, $p = .044$, spatial $DW = 2.10$ (Figure S1B). Using the Howells data, prior research had already found that the shape diversity of male crania negatively correlates with distance from Africa (von Cramon-Taubadel & Lycett, 2008).[5] Amongst female crania, distance from Africa correlated negatively with cranial shape diversity, $r(24) = -.59$, $p = .009$, spatial $DW = $

2.26 (Figure S1C). For cranial form diversity, the geographical area of best fit was only in Africa (Figure 2A-2D). Regarding cranial form diversity, in preceding research the area had been entirely in Africa for males (Betti et al., 2009; Manica et al., 2007), but not for females (Betti et al., 2009).

Figure 2

Cranial Diversity Suggests African Origin of Worldwide Expansion

Note. The fit of geographical distance to types of diversity: male cranial form (2A and 2B), female cranial form (2C and 2D), male cranial shape (2E and 2F), and female cranial shape (2G and 2H). In Figure 2A, 2C, 2E, and 2G, darker grey indicates an area for where the expansion originated. In Figure 2B, 2D, 2F, and 2H, a lower BIC signals a better fit to the relationship of distance and diversity. Distance from Botswana was used on the *y*-axes in right panels (also in Figures 3C, S2, S3B, and S4) rather than the location which gives the lowest BIC because correlation tests regarding the cranium were run using distances from Botswana. Moreover, correlation coefficients are given within various cranial dimensions in Figures S2 and S4. It is likely that peak points would vary between dimensions. As in Cenac (2022), distances of origins were from Botswana so that the distributions of correlation coefficients could be compared generally (dimensional vs. *overall*).

As for cranial shape diversity, the area was just in Africa for male crania (Figure 2E and 2F). In prior research with male crania, R^2s representing the extent of association between shape diversity and geographical distance were numerically greater when distance was from Africa (Botswana, Democratic Republic of the Congo, or Ethiopia) rather than from Asia (China, India, or Israel) (von Cramon-Taubadel & Lycett, 2008) which is congruent with the present research which found that the area of best fit is in Africa. However, for the cranial shape diversity of females, there was more than one geographical area of best fit, with one being in Africa and at least one other being outside of Africa (Figure 2G and 2H). This finding arose from some locations in Oceania having low BICs. For female crania, correlation coefficients were negative for locations in Africa but positive for locations in Oceania (Figure S2C). Because research agrees with a decline in types of diversity with distance from the expansion origin (Ramachandran et al., 2005; von Cramon-Taubadel & Lycett, 2008), the area with negative coefficients is the estimated area of origin (as discussed in the *Introduction*). Therefore, results with female cranial shape diversity are in step with the expansion being from Africa and not elsewhere. As a result, in the present research, cranial diversity in form and shape lead to the estimations of origin areas which are only in Africa.

Cranial shape diversity appears to indicate a smaller origin area than cranial form diversity does (Figure 2). Hence, cranial shape diversity may be better at indicating the origin compared to cranial form diversity. Such an outcome should not be unexpected. This is because shape, along with size, is part of form (Betti et al., 2010) and cranial size diversity (controlled for absolute latitude) has no correlation with distance from Africa (Cenac, 2022).

Cranial size dimorphism

Regarding a test of whether cranial size dimorphism correlates with distance from Africa (Botswana), positive spatial autocorrelation presented when using 26 populations, spatial $DW = 1.17$. A graph (produced through the Chen, 2016, spreadsheet) indicated that this was caused by two populations: Ainu and North Japan. The positive spatial autocorrelation was resolved by the absence of those populations, spatial $DW = 1.87$. With the remaining 24 populations, it was found that cranial size dimorphism and distance from Africa have a positive correlation, $r(22) = .66$, $p = .003$ (Figure 3A).

Using cranial size dimorphism, an origin area was suggested within Africa alone (Figure 3B and 3C). This seems congruent with cranial size dimorphism having a link to the expansion from Africa, and cranial size dimorphism being a useful indicator of this expansion. Whilst prior research had explored whether cranial size dimorphism, when adjusted for absolute latitude, does have a correlation with distance from Africa (Cenac, 2022), such an adjustment is likely to be without value (Text S1). Nevertheless, with adjusted cranial size dimorphism, an origin area was also found to fully be in Africa (Figure S3). And so, given that adjusted cranial size dimorphism does correlate with

distance from Africa (Cenac, 2022; Text S1), cranial size dimorphism appears to remain as an indicator of the expansion after adjusting it for absolute latitude.

The level of correlation between adjusted cranial size dimorphism and geographical distance does not appear to be a good yardstick of coefficients for correlations between the adjusted dimorphism of individual dimensions and geographical distance (Cenac, 2022). This is apparent with unadjusted dimorphism (i.e., zero-order correlation coefficients) (Figure S4A and S4C). Such an occurrence is present regarding relationships between geographical distance and both cranial form diversity (Cenac, 2022; Figure S2A and S2B) and cranial shape diversity (Figure S2C-S2F).

Figure 3

Cranial Size Dimorphism Indicates the Origin of the Expansion

Note. In Figure 3B, a darker grey indicates the origin area, unlike the lighter grey. The lowest BIC (-170.71) was when using a location in South Africa (30°S, 20°E) as the origin.

Mitochondrial diversity

As for mitochondrial diversity, when using 32 worldwide origins, the lowest BIC was in North America, specifically at the Arikara location which had a BIC of 382.75 (Figure 4A). Origins with BICs within four of Arikara were in North America and Africa (Figure 4A). As with cranial shape diversity, one can note the direction of *r*-values. Such values were negative for Africa and positive for North America (Figure 4B). Hence, the geographical area of best fit in Africa can be considered as the estimated origin area for the same reason as with cranial shape diversity amongst females. Using the four-BIC route (Manica et al., 2007), when therefore not considering origin locations in North America, the geographical area of best fit was only within Africa (Figure 4C).

Figure 4

Mitochondrial Diversity Suggests that Africa has the Expansion Origin

Note. A) BICs are for the relationship between geographical distance and mitochondrial diversity, with distances being from 32 locations (origins) worldwide. B) Correlation coefficients are displayed, each being the level to which distance from Africa and mitochondrial diversity are correlated. The distances are from the 32 locations. C) The darker grey is the origin area suggested.

Diversity of allosomes

Results indicated that the geographical area of best fit for the relationship of X-chromosomal microsatellite heterozygosity and geographical distance was exclusively in Africa (Figure 5A and 5B). Therefore, as this type of diversity does correlate negatively with geographical distance from Africa (Balloux et al., 2009), X-chromosomal microsatellite heterozygosity does indeed seem to have an indication of the expansion from Africa.

With Y-chromosomal microsatellite heterozygosity, the best BIC was found to be within India (Figure 5D). Distance from India did correlate negatively with Y-chromosomal diversity, $r(49) = -.55$, $p < .001$, spatial $DW = 1.83$. All origins in Africa were more than four BICs away from the lowest BIC. Therefore, Africa did not have any of the geographical area of best fit (Figure 5C). No origin (of those used) outside of Asia was within four BICs of the lowest, and so the geographical area of best fit for Y-chromosomal microsatellite heterozygosity was possibly within Asia fully.

However, what if more than one origin was indicated? Previous research had considered this with respect to autosomal heterozygosity in expansion (Manica et al., 2007) and the origin of language (Atkinson, 2011). For instance, Atkinson (2011) constructed models for predicting phonemic diversity from types of geographical distance (distance from the peak point in Africa,

distance from elsewhere) and population size. Atkinson (2011) did not find that phonemic diversity (in the model) had a negative correlation with distance from elsewhere. A similar sort of analysis was used in the present research; when adjusting Y-chromosomal diversity for distance from India, the numerically lowest correlation coefficient (*sr*) for the relationship between distance and adjusted diversity when using an African origin was found – distance from this origin was not correlated with the adjusted diversity, *sr*(48) = -.10, *p* = .97, spatial *DW* = 1.83 (the correlation test was run using the ppcor package of Kim, 2015). As a result, although there is a negative correlation between Y-chromosomal microsatellite heterozygosity and distance from Africa (Balloux et al., 2009), this relationship would not appear to actually be reflective of the expansion from Africa.

Figure 5

X-Chromosomal, but not Y-Chromosomal, Microsatellite Diversity Signals African Origin for Expansion

Note. Figure 5A and 5B are with respect to X-chromosomal heterozygosity, whilst 5C and 5D are for Y-chromosomal. Regarding X-chromosomal diversity (5A), of the locations in Africa used as origins, the origin leading to the best BIC was at the coordinates (15°S, 15°E), i.e., in Angola. Origins (locations) in the area shaded with a darker grey are inside the geographical area of best fit – origins in the lighter grey are outside of the area. For 5B and 5D, however, calculation of BICs used 32 origins from across the globe. In 5B, origins outside of Africa were not within the geographical area of best fit. As for Y-chromosomal diversity, of the origins used (whether within Africa or outside), India had the best BIC (5D). Origins in Africa were not inside four BICs of the best, and so Africa was not within the geographical area of best fit regarding Y-chromosomal diversity (5C).

When using a combined African population and a number of non-African populations, a phylogenetic tree (arising from Y-chromosomal microsatellites) has its root located between Africans

and non-Africans (Seielstad et al., 1999), hence Table 1 and Figure 5C and 5D may be surprising. Nevertheless, studies are congruent with there being migration back into Africa (e.g., Henn, Botigué et al., 2012; Hodgson et al., 2014). Indeed, whilst research with the Y chromosome does suggest expansion from Africa into Asia, migration from Asia to Africa is also indicated (e.g., Hammer et al., 2001). Furthermore, if Y-chromosomal haplogroup E (which has been described as being the most common of Y haplogroups in Africa) was from an expansion from Asia (Wang & Li, 2013), maybe it should not be that surprising if extents of Y-chromosomal diversity do not reflect the worldwide expansion from Africa. Hence, the signal of expansion from Asia in Y-chromosomal microsatellite diversity (Figure 5D) could very well represent migration from Asia to other continents, including back into Africa.

Nevertheless, in Balloux et al. (2009), Y-chromosomal diversity was calculated from six microsatellites, whilst X-chromosomal and autosomal microsatellite diversities were determined from 36 and 783 microsatellites respectively. Ten Y-chromosomal microsatellites is few (Seielstad et al., 1999). A tally of six microsatellites could be described similarly. Therefore, the apparent signalling of expansion from Asia (rather than from Africa) in Y-chromosomal diversity (Figure 5C and 5D) was found through few microsatellites. It would be unwise to discount Y-chromosomal microsatellite diversity from reflecting the expansion from Africa until far more microsatellites are used.

Finding genetic diversity to not be greatest in Africa can be indicative of ascertainment bias (e.g., Conrad et al., 2006, with SNP heterozygosity). Previous research has considered if there are indications of ascertainment bias regarding microsatellites in the HGDP-CEPH data (e.g., Ray et al., 2005, who utilised microsatellite data used in Rosenberg et al., 2002, and Rosenberg et al., 2002, employed autosomal microsatellites) – perhaps future research could address whether ascertainment bias is present in those six Y-chromosomal microsatellites.

Expansion origin centroid

Because Y-chromosomal diversity was not found to signal the expansion from Africa, coordinates regarding Y-chromosomal diversity were not included when calculating the origin centroid. Given the lack of support for using an adjustment for absolute latitude on cranial size dimorphism (Text S1), the centroid (Figure 6) was not calculated using the adjusted dimorphism.

The shape used when estimating the expansion origin centroid was formed of lines connecting coordinates (with coordinates being vertices) – from the autosomal diversity coordinate to mitochondrial diversity to cranial size dimorphism to cranial shape diversity to X-chromosomal diversity to autosomal diversity. Regarding indicators of the expansion from Africa, most of the types of peak points are in southern Africa (four out of five), which generally supports southern Africa being the origin (Figure 6, with two of these *peak points* themselves being a centroid or an average). Indeed, a centroid of the peak points suggests that southern Africa is the most probable origin of the expansion (Figure 6), in the context of the peak points generally clustering in the southern African

region (like in Choudhury et al. 2021, Angola, Malawi, and Zambia were regarded as part of southern Africa in the present research).

Figure 6

The Expansion Originated in Southern Africa?

Note. Smaller grey circles (peak points) represent the coordinates regarding cranial size dimorphism (30°S, 20°E), mitochondrial diversity (25°S, 15°E), autosomal diversity (.57°N, 10.10°E), cranial shape diversity (25.02°S, 17.50°E), and X-chromosomal diversity (15°S, 15°E). The peak point for autosomal diversity is itself a centroid, and the peak for cranial shape is an average. The larger grey circle stands for the centroid (coordinates) of (the shape resulting from) those other five coordinates – a location (17.86°S, 14.70°E) which is close to the Angola/Namibia border, similar to the location found in Tishkoff et al. (2009) regarding autosomal microsatellite diversity. The grey background was made using 99 coordinate pairs in Africa from Betti et al. (2013).

Number of populations

From the number of populations (Balloux et al., 2009; Howells, 1989, 1996; Ramachandran et al., 2005; Tishkoff et al., 2009) used in analyses, few populations in Africa were involved when finding peak points via diversity/dimorphism except for autosomal repeat length diversity. Given previous research on phonemic diversity (Van Tuyl & Pereltsvaig, 2012), in the context of calculating origins, a concern may be those instances in which there were only a small number of populations in Africa. Language has featured in the discourse regarding the expansion from Africa, with reference being made to Atkinson (2011) (e.g., Atkinson, 2012; Henn, Cavalli-Sforza, & Feldman, 2012). In Atkinson (2011), it was found that phonemic diversity falls as distance from Africa rises, and that the

geographical area of best fit for the negative association between distance and phonemic diversity is only within Africa (for discussion regarding Atkinson, 2011, see Atkinson, 2012, or Van Tuyl & Pereltsvaig, 2012, for example). The location leading to the lowest negative association between geographical distance and phonemic diversity in Atkinson (2011) was in Africa (Van Tuyl & Pereltsvaig, 2012). However, if populations are grouped by geographical region (e.g., North America, South America, etc.), it can be seen that when adjusting the *origin* in Africa there is a shift in the correlation of distance and phonemic diversity for populations in Africa, but no shift for populations in regions outside of Africa (Van Tuyl & Pereltsvaig, 2012). Indeed, the location producing the strongest negative correlation is only a bit different when using worldwide populations compared to populations in Africa alone (Van Tuyl & Pereltsvaig, 2012); populations outside Africa are immaterial to defining which particular location in Africa has the lowest negative association between distance and phonemic diversity – only populations within Africa define the location (Van Tuyl & Pereltsvaig, 2012), or so it would appear.

Figure 7

Peak Points with Populations Globally, and in Africa

Notes. Peak points were found (through the *r* correlation coefficient) using populations worldwide (larger circles) and populations in Africa alone (smaller circles) with A) cranial shape diversity of males, B) cranial shape diversity of females, C) cranial size dimorphism, D) autosomal SNP heterozygosity, E) mitochondrial diversity, F) X-chromosomal microsatellite diversity. Data described in the *Method* were used. Like in Figure 6, the coordinates in Betti et al. (2013) were utilised to form the grey backgrounds.

Indeed, when considering the relationships between geographical distance and variables more generally, it seems that there is variety in how close peak points in Africa are when the peaks are found with worldwide populations compared to Africa alone (Figure 7). Yet, there indeed appears to

generally be agreement between peak points regardless of whether they are calculated from populations worldwide or in Africa only (Figure 7). And so, peak points in Africa (when using populations across the globe) are not defined exclusively by African populations, but may only be particularly *driven* by such populations.

If the distribution of biological diversity/dimorphism in Africa drives where the peak point in Africa is found to be, there could indeed be a concern over the number of populations in Africa that were featured when calculating the origin centroid in Figure 6. However, the origin centroid did arise from different types of diversity/dimorphism variables (and some different samples), which should counter the issue of small numbers of populations to an extent.

Within Africa

Tishkoff et al. (2009) used worldwide populations, including 121 in Africa. The peak location in their study for the relationship between geographical distance and diversity in autosomal microsatellite repeat lengths is close to the origin centroid of Figure 6 (in the present research, that peak was used to calculate the centroid for autosomal diversity, which was used to calculate the origin centroid).

Figure 8

Origin of the Expansion from the Autosomal Microsatellite Diversity of Populations in Sub-Saharan Africa

Note. Diversity was not adjusted for non-African ancestry in 8A, but it was adjusted in 8B and 8C. In 8A and 8B, the darker grey background signifies locations which, when used as origins, were within four BICs of the lowest and led to negative correlation coefficients, and the location with that lowest BIC is the grey circle. Nonparametric analysis indicated a more southerly peak point (Figure S5).

Using populations across the world, autosomal microsatellite heterozygosity i) is found to negatively correlate with distance from Africa (Prugnolle et al., 2005), and ii) results in an origin area that is fully in Africa (Manica et al., 2007). If the location of the origin centroid *within* Africa is greatly due to populations in Africa, one would assume that the signal of the expansion is clear

amongst populations in Africa (although correlations between distance and autosomal diversity inside Africa were not present in Hunley & Cabana, 2016, but see proceeding paragraphs). In the present research, the magnitude of relationships between autosomal microsatellite heterozygosity (106 populations) in sub-Saharan Africa and geographical distance (within Africa) was strongest when distance was from Angola (15°S, 20°E), with there being a negative correlation between heterozygosity and distance from that location, $r(104) = -.31$, $p = .006$, spatial $DW = 2.01$, and an origin area in southern Africa was indicated (Figure 8A).[6] (This is by no means the first analysis to explore whether distance and a variable are associated within Africa, e.g., Henn, Gignoux et al., 2011, did explore this.) Therefore, autosomal microsatellite heterozygosity within sub-Saharan Africa seems to indicate a southern African origin of an expansion within Africa. Assuming that this expansion is part of the worldwide expansion, then the origin of the worldwide expansion appears to be indicated. Hence, this origin indicated with populations in Africa (Figure 8A) broadly agrees with the origin centroid.

However, a graph placing distance (from the peak) against microsatellite heterozygosity suggested that heteroscedasticity (i.e., non-constant variability) was present. Therefore, the confintr package (Mayer, 2022) was used in R to bootstrap a 99% confidence interval regarding the correlation coefficient (r) for the relationship between distance from the peak and heterozygosity. The default method of bias-corrected accelerated (Mayer, 2022) was chosen. In bootstrapping, a total of 1,000 samples was used – to match the number of bootstrap samples employed when comparing correlation coefficients to each other in the analysis presented in Text S1. The confidence interval [-.51, -.01] did not include 0 and it consisted entirely of negative values, thereby favouring there being a negative correlation. Furthermore, nonparametric analysis supported the presence of a negative correlation, albeit with a different peak point which was (25°S, 15°E), $r_s(104) = -.51$, $p < .001$ (Figure S5A), with the origin area still being indicated to fully be in southern Africa (Figure S5B).

Nevertheless, Hunley and Cabana (2016) did not find a correlation between distance from eastern Africa and autosomal microsatellite diversity within any of three categories of populations in Africa. Yet an absence of correlations should not be surprising because, in the present research, eastern Africa was not in the origin area for the relationship between distance and heterozygosity in sub-Saharan Africa (Figure 8A), with southern Africa looking to be a more plausible origin for the expansion than eastern Africa (Figures 6 and 8A).

Non-African ancestry

Keeping in mind previous research with respect to admixture (e.g., Haber et al., 2016; Pickrell et al., 2014; Pickrell & Reich, 2014), non-African ancestry within Africa should be considered. The pattern of heterozygosity falling as geographical distance from a population increases occurs in a scenario in which there are a small number of bottlenecks (two) and admixture (with the population not being used as the source of a worldwide expansion) (Pickrell & Reich, 2014). Hence, the support

for a southern African origin (over other regions in the continent) for a worldwide expansion (Figures 6 and 8A) could have occurred even in the absence of bottlenecks within sub-Saharan Africa, and instead this support could have resulted from admixture. Indeed, previous research is congruent with heterozygosity decreasing with greater Eurasian ancestry amongst Africans who are admixed (Haber et al., 2016). Therefore, it can be supposed that, more generally, the heterozygosity of Africans would decrease with increasing non-African ancestry (given the fall in autosomal heterozygosity with distance from Africa in Prugnolle et al., 2005), and perhaps there would be relationships with non-African ancestry beyond heterozygosity, e.g., with cranial shape diversity and cranial size dimorphism. So, non-African ancestry in Africa could plausibly affect origin areas, no matter whether those areas are found via genetic or cranial data. Moreover, given Figure 7 alongside Van Tuyl and Pereltsvaig (2012), it seems that peak points within Africa for cranial/genetic variables would likely be driven by populations in Africa, therefore non-African ancestry within Africa could greatly be determining the peak points.

Such an assertion would depend on there being variation in non-African ancestry in Africa. Pickrell et al. (2014) presented extents of West Eurasian ancestry for a number of populations. An observation of theirs (Pickrell et al., 2014, p. 2636) certainly can be interpreted as referring to West Eurasian ancestry in populations largely being lower in southern Africa than in eastern Africa. Such a (numerical) difference between regions can be backed up by Figures 2B of Gallego Llorente et al. (2015) which presents extents of West Eurasian ancestry on a map. Therefore, perhaps West Eurasian (and more generally non-African) ancestry contributed towards the apparent support in the present research for the origin of a worldwide expansion from southern Africa.

In the present research, the data in Tishkoff et al. (2009) was used to examine whether the expansion origin is indicated to be in southern Africa when autosomal heterozygosity in sub-Saharan Africa is adjusted for non-African ancestry (distance was not adjusted).[7] With the adjusted heterozygosity, an origin area in southern Africa was found (Figure 8B). The location at which the BIC was lowest was the same as with unadjusted heterozygosity (parametric analysis). Distance from (15°S, 20°E) was negatively correlated with adjusted heterozygosity, $sr(103) = -.32$, $p = .008$, spatial $DW = 2.00$ (Figure 8C). However, heteroscedasticity was indicated to occur (Figure 8C). Therefore, ppcor (Kim, 2015) was used to run a semi-partial Spearman correlation between distance from the peak of (25°S, 15°E) and autosomal diversity when the diversity was adjusted for non-African ancestry; there was a correlation, $sr_s(103) = -.54$, $p < .001$ (Figure S5C). Moreover, southern Africa was suggested to have the expansion origin area (Figure S5D). Therefore, the signalling of a southern African origin through autosomal heterozygosity would not appear to be attributable to non-African ancestry.[8]

Results suggest that autosomal microsatellite heterozygosity within sub-Saharan Africa reflects part of the worldwide expansion. This finding is favourable towards the origin centroid (Figure 6)

actually approximating where the worldwide expansion started. However, it remains unestablished if adjusting for non-African ancestry would have a minimal effect (or no effect) on the origin areas and peak points found with other variables (e.g., cranial size dimorphism). Given populations and their locations in Tishkoff et al. (2009), the present research used relatively few populations inside southern Africa (four populations) in the analyses with sub-Saharan African heterozygosity. Indeed, not many populations were close to the location of the lowest BIC (e.g., Figure 8C). This does undermine some confidence in whether adjusted/unadjusted heterozygosity actually declines linearly in sub-Saharan Africa as distance from southern Africa increases. Critically, it is not clear if an expansion signal in autosomal microsatellite heterozygosity is present in regions outside of Africa – if the signal is not in other regions (Text S2), it seems questionable whether the signal in Africa actually reflects the expansion.

Additional support?

Might findings with phonemic diversity (Atkinson, 2011; Van Tuyl & Pereltsvaig, 2012), nucleotide diversity (Luca et al., 2011), and linkage disequilibrium (Henn, Gignoux et al., 2011) bolster the origin centroid? The peak point concerning the negative relationship between phonemic diversity and geographical distance (Van Tuyl & Pereltsvaig, 2012) is in the vicinity of the overall peak point found with autosomal diversity (Figure 6). However, whilst Atkinson (2011) found that Africa exclusively holds the area that is within four BICs of the lowest BIC regarding phonemic diversity (phoneme inventory size), Africa was not included in such an area in Creanza et al. (2015). Indeed, Creanza et al. (2015) found the lowest BIC (or Akaike information criterion) was not even within Africa, but in Europe, with there being a negative association between phoneme inventory size and distance from Europe (see Creanza et al., 2015, for their thoughts on how the negative association came about). Generally, given previous research/discussion (e.g., Atkinson, 2011; Creanza et al., 2015), it seems uncertain whether phonemic diversity is useful for indicating the origin within Africa of the worldwide expansion. Consequently, it is not clear if findings with phonemic diversity (Atkinson, 2011; Van Tuyl & Pereltsvaig, 2012) lend support to the origin centroid.

In Luca et al. (2011), correlation coefficients were calculated regarding nucleotide diversity and distance from locations in Africa. The (numerically) largest correlation coefficient regarding nucleotide diversity and geographical distance was when distances were from southern Africa (and this coefficient was negative) (Luca et al., 2011).

Using only populations in Africa, in Henn, Gignoux et al. (2011), there was a positive correlation between linkage disequilibrium (5 kilobases, autosomal SNPs) and distance from southwestern Africa. Furthermore, out of locations used as origins in Africa, it was found that the strongest positive correlation was within the southern-western area of Africa (Henn, Gignoux et al., 2011). However, it has been noted by López et al. (2015) that, in Schlebusch et al. (2012), linkage disequilibrium in South Africa was roughly as low as it is in places within Africa that are outside of

South Africa. Given Schlebusch et al. (2012) and wording in López et al. (2015, p. 59), presumably López et al. (2015) were referring to southern Africa rather than the Republic of South Africa in particular.

More caveats, and future research

Origin area estimation problem?

As discussed above, building on Van Tuyl and Pereltsvaig, (2012), when genetic/cranial data consist of populations across the globe, the peak point (within Africa) appears to disproportionately be due to populations within Africa. A disproportionate weighting may imply a problem when estimating the origin area. From examining a formula for calculating the BIC (Masson, 2011), the number of populations (no matter whether they are inside Africa or outside) would affect the size of the origin area – a larger number of populations would reduce the area in general.[9] From previous research (e.g., Betti et al., 2009; Manica et al., 2007), it is clear that, when estimating origin areas, it is usual that relatively few populations are in Africa and most populations are outside of Africa. Therefore, the size of the estimated origin area would clearly be affected by the number of populations outside of Africa (and within). It could be bothersome that, when using populations worldwide, the peak point appears to be determined disproportionately (not fully) by populations in Africa (Figure 7) whereas the size of the origin area would be notably contributed to by populations outside of the continent (and not largely by populations within Africa). Surely the inside Africa vs. outside Africa influence really *ought* to be similar for both the peak point and the size of the origin area. So, it could be argued that the size of the origin area may be smaller than it ought to be. This may not only be an issue regarding the BIC. Origin areas estimated through the AIC (e.g., Betti et al., 2013) may also be affected. Consequently, estimated origin areas in prior research (e.g., Betti et al., 2013; Manica et al., 2007), and several in the present research, could be smaller than they should be. Nevertheless, this possible problem with origin areas was avoided in the analyses which used populations in sub-Saharan Africa alone (Figure 8).

For worldwide samples, a remedy to the possible estimation problem can perhaps be found in Tishkoff et al. (2009). Through a bootstrapping method, they represented on a map the frequency with which a location was the strongest (numerically) for the relationship between geographical distance and autosomal microsatellite diversity in repeat lengths (Tishkoff et al., 2009). The resulting area (Tishkoff et al., 2009) would likely not be subject to the underestimation which may affect BIC-based and AIC-based methods.

Defining which variables signal the expansion

In the present research, for a variable to be noted as an indicator of the expansion from Africa and be included when calculating the origin centroid, certain criteria were applied; the variable had to correlate with distance from Africa, and the variable had to signal an origin area that is only within

Africa. These criteria may be biased in favour of variables that have peak points which are more southerly/westerly within Africa.

Why may there be a bias? The answer to this question is in regard to the size of the origin area. An origin area can encompass quite a lot of continental Africa (e.g., Figure 2). Therefore, given support for a northeastern exit from Africa in the worldwide expansion (Tishkoff et al., 2009), a variable with a peak point (in Africa) closer to this exit is perhaps more likely to have an origin area which reaches outside of Africa. Consequently, with respect to peak points in Africa, a variable that has its peak point nearer to the exit may have a higher likelihood of being discarded as an indicator of the expansion. And so, a bias may have occurred when applying criteria used in the present research for defining an indicator of the expansion. Therefore, the origin centroid (Figure 6) could have been biased against peak points which are nearer to the exit. This bias could promote finding a southerly/westerly centroid in Africa. Indeed, the centroid is in southwestern Africa (Figure 6). More relaxed criteria might reduce the bias. The relaxed criteria could be: i) an association with distance, ii) a peak point in Africa, and iii) an origin area at least largely in Africa. Hip bone shape diversity falls with the increase in geographical distance from Africa, it has a peak point in Africa, and whilst it leads to an origin area not fully within Africa (Betti et al., 2013), Figure 4 of Betti et al. (2013) shows that the area is mostly in Africa. With hip bone shape diversity, the peak point is in southern Africa for males and for females (Betti et al., 2013). Hence, even with a bias in calculating the origin centroid, this bias might not necessarily have greatly affected which region of Africa the centroid is located in.

Mitochondrial diversity and climate

Whereas previous research has found an origin area and peak point using cranial form diversity after adjusting this type of cranial diversity for climate (Manica et al., 2007), research does not support cranial form diversity (overall) being associated with climate (Betti et al., 2009) (although climate is associated with the diversity of some cranial dimensions, Manica et al., 2007). There could be good reason to adjust mitochondrial diversity for climate. Mitochondrial diversity is associated with minimum temperature (Balloux et al., 2009). Furthermore, mitochondrial diversity (with distance from Africa being adjusted for) increases with minimum temperature (Balloux et al., 2009). Hence, the origin area (Figure 4) and peak point found with mitochondrial diversity (Figure 6) could perhaps be refined by adjusting mitochondrial diversity for minimum temperature.

However, the distribution of population locations in Africa may need to be kept in mind. The peak point found with (unadjusted) mitochondrial diversity is in southern Africa when using populations from across the globe (Figure 6). However, when analysing mitochondrial diversity in the present research, going by Balloux et al. (2009), no populations in southern Africa were featured. With this absence of southern African representation, there is reason to be hesitant about both the origin area and peak point for mitochondrial diversity.

Admixture and diversity

Previous research indicates that SNP heterozygosity in Africans decreases as Eurasian ancestry greatens (Haber et al., 2016). It could be worthwhile considering if this (to some extent) could be explained by bottlenecks within Africa in the worldwide expansion given that the present research suggests i) that the worldwide expansion began in southern Africa and ii) that the signal of the expansion is present in Africa. Bearing in mind research regarding Native Americans (Hunley & Healy, 2011), it would not be surprising if support for the association between Eurasian ancestry and genetic diversity in Africa (Haber et al., 2016) would still be indicated despite the expansion signal in Africa. In that previous research with Native Americans, there was a negative correlation between gene identity (of autosomal microsatellites) and European ancestry, and a negative correlation was still present when adjusting for the distance between Beringia and populations (Hunley & Healy, 2011).

Type of origin

The geographical area which is indicated to be the one from which modern humans expanded (i.e., expansion origin) can be interpreted as being the suggested area for where modern humans *originated* (i.e., where they first lived – modern human origin) (Manica et al., 2007); the origin centroid could be described as signaling the geographical location at which modern humans did originate. Regarding populations in Africa, it is possible that their ancestors were outside of southern Africa (but still within Africa) prior to going to the south and expanding (Henn, Bustamante et al., 2011). Therefore, origin areas for the worldwide expansion in the present analyses (e.g., Figure 1A; Figure 2A) need not imply that modern humans originated within those specific areas, and the expansion origin centroid (Figure 6) is not necessarily suggestive of modern humans having originated in southern Africa. Furthermore, the present research is not intended to suggest that all the ancestry of modern humans descends from the geographical origin of the worldwide expansion (for instance, see the literature on Neanderthal ancestry, e.g., Green et al., 2010).

Conclusion

The present analyses explored which cranial and genetic variables may signal the worldwide expansion of modern humans, and where in Africa this expansion commenced from. Support was found for the signal being present in various types of diversity (e.g., X-chromosomal microsatellites, cranial shape) and cranial size dimorphism. Nonetheless, despite Y-chromosomal microsatellite heterozygosity being related to geographical distance from Africa (Balloux et al., 2009), that type of diversity was not indicated to signal the expansion. Yet that diversity was from six Y-chromosomal microsatellites (Balloux et al., 2009), i.e., not many (given Seielstad et al., 1999), so it really is not clear if the signal is present in the heterozygosity of Y-chromosomal microsatellites. As for where in Africa the expansion originated, the location of a geographical centroid (determined through a number of expansion indicators) and locations of peak points (using populations worldwide) adds weight to

southern Africa being the origin (Figure 6). In the context of research with phonemic diversity (Van Tuyl & Pereltsvaig, 2012), the locations of these peak points and this centroid may greatly arise from biological diversity/dimorphism within Africa (Figure 7). This points towards the signal of the expansion being found amongst populations in the continent. Indeed, analyses with autosomal microsatellite diversity in sub-Saharan African populations suggested the presence of the expansion signal in Africa, with the expansion seeming more likely to have originated in southern Africa than in other regions of Africa. Nevertheless, a number of caveats (as described above) do remain. It is far from a certainty that southern Africa is the region from which the worldwide expansion originated.

Footnotes

[1] Indeed, depending on which location is utilised as the origin, correlation coefficients on the association between diversity and geographical distance can vary between negative and positive (e.g., Hunley et al., 2012; Ramachandran et al., 2005). An example of this can be found with autosomal microsatellite diversity – the correlation coefficient is negative when using Africa as an origin, whilst the coefficient is positive when South America is used (Ramachandran et al., 2005).

[2] Figure 1 of Henn, Cavalli-Sforza, and Feldman (2012) represents expansion of modern humans from inside Africa to Eurasia, and from Eurasia to Oceania and the Americas.

[3] After referring to previous research (specifically Prugnolle et al., 2005, and Ramachandran et al., 2005) on autosomal microsatellite heterozygosity falling as distance from eastern Africa increases, Deshpande et al. (2009) appear to have noted that the decline is strongest from East Africa. However, Prugnolle et al. (2005) did not describe examining whether a decline in autosomal microsatellite diversity is greater when distance is from East Africa rather than anywhere else. Additionally, Ramachandran et al. (2005) did not find that the peak decline occurs when using distance from East Africa. Moreover, Manica et al. (2007) estimated an origin area using autosomal microsatellite heterozygosity; that origin area did include East Africa, but also other regions of Africa (Manica et al., 2007).

[4] The names given for populations in the Howells data (Howells, 1989, 1996) are used in the present research.

[5] In the present analyses, distance from Africa (Botswana) was found to have a negative correlation with cranial shape diversity amongst males, $r(26) = -.67$, $p = .001$, spatial $DW = 2.39$. Because von Cramon-Taubadel and Lycett (2008) had already found a correlation coefficient and p-value regarding distance from Botswana and cranial shape diversity using the male crania (28 populations in the Howells dataset), it was not necessary for this correlation test to have been run in the current analyses. However, von Cramon-Taubadel and Lycett (2008) did not state whether they examined residuals for positive spatial autocorrelation.

[6] Research/discussion has occurred regarding genetic diversity in sub-Saharan Africa (e.g., Bergström et al., 2021; Schlebusch et al., 2012). Present analyses with autosomal microsatellite heterozygosity in Africa only used populations in sub-Saharan Africa. This is because of research in line with i) migration back into the continent of Africa (Henn, Botigué et al., 2012; Hodgson et al., 2014), ii) the presence of non-African ancestry

in the continent (Tishkoff et al., 2009), and iii) for Africans with admixture, heterozygosity falling as Eurasian ancestry increases (Haber et al., 2016). Therefore, solely populations in sub-Saharan Africa were used in order to lower any impact on results of migration back into Africa. That said, non-African ancestry in sub-Saharan Africa (e.g., Tishkoff et al., 2009) is acknowledged.

[7] Ancestry has been adjusted for (Hunley & Cabana, 2016; Hunley et al., 2016) previously. For instance, research on expansion has adjusted distance from Africa and the autosomal diversity of Native Americans for European ancestry (Hunley & Cabana, 2016).

[8] Previous research explored correlations between gene identity and European ancestry amongst Native Americans when adjusting both variables for how far from Beringia populations are, and when not adjusting variables for distance (with gene identity being the number one minus gene diversity) (Hunley & Healy, 2011); in the present research, also given the research aligning with a relationship between Eurasian ancestry and the heterozygosity of Africans (Haber et al., 2016), the Tishkoff et al. (2009) data was utilised to find if non-African ancestry correlates with the autosomal heterozygosity of populations in sub-Saharan Africa. The autosomal heterozygosity was adjusted for distance from the peak point in some correlation tests. Interestingly, there was no correlation between the non-African ancestry and heterozygosity, whether heterozygosity was, $sr(103) = .13$, $p = .72$, spatial $DW = 2.00$ [$sr_s(103) = .17$, $p = .44$] or was not adjusted for distance from the peak point, $r(104) = .09$, $p = .97$, spatial $DW = 1.89$ [$r_s(104) = .10$, $p = .97$]. There appears to have been little variability in the non-African ancestry (Figure S6), and this lack of variation could explain the absence of correlations.

[9] In a previous study, BIC was used for finding the origin area through cranial form diversity (adjusted for climate) (Manica et al., 2007). Regarding the crania of males (105 populations; 4,666 crania) and females (39 populations; 1,579 crania), amongst females, the dataset was too small to sift between possible origins (Manica et al., 2007) which suggests that the sample size influences the size of the origin area estimated through the BIC.

References

Atkinson, Q. D. (2011). Phonemic diversity supports a serial founder effect model of language expansion from Africa. *Science*, *332*(6027), 346-349. https://doi.org/10.1126/science.1199295

Atkinson, Q. D. (2012). Response to comments on "Phonemic Diversity Supports a Serial Founder Effect Model of Expansion from Africa". *Science*, *335*(6069), Article 657-e. https://doi.org/10.1126/science.1210005

Balloux, F., Lawson Handley, L.-J., Jombart, T., Liu, H., & Manica, A. (2009). Climate shaped the worldwide distribution of human mitochondrial DNA sequence variation. *Proceedings of the Royal Society B: Biological Sciences*, *276*(1672), 3447-3455. https://doi.org/10.1098/rspb.2009.0752

Bartholomew illustrated reference atlas of the world (1985). John Bartholomew & Son Limited.

Batini, C., & Jobling, M. A. (2011). The jigsaw puzzle of our African ancestry: Unsolved, or unsolvable? *Genome Biology*, *12*(6), Article 118. https://doi.org/10.1186/gb-2011-12-6-118

Benson, D. A., Cavanaugh, M., Clark, K., Karsch-Mizrachi, I., Lipman, D. J., Ostell, J., & Sayers, E. W. (2013). GenBank. *Nucleic Acids Research*, *41*(Database issue), D36-D42. https://doi.org/10.1093/nar/gks1195

Bergström, A., Stringer, C., Hajdinjak, M., Scerri, E. M. L., & Skoglund, P. (2021). Origins of modern human ancestry. *Nature*, *590*(7845), 229-237. https://doi.org/10.1038/s41586-021-03244-5

Betti, L., Balloux, F., Amos, W., Hanihara, T., & Manica, A. (2009). Distance from Africa, not climate, explains within-population phenotypic diversity in humans. *Proceedings of the Royal Society B: Biological Sciences*, *276*(1658), 809-814. https://doi.org/10.1098/rspb.2008.1563

Betti, L., von Cramon-Taubadel, N., & Lycett, S. J. (2012). Human pelvis and long bones reveal differential preservation of ancient population history and migration out of Africa. *Human Biology*, *84*(2), 139-152. https://doi.org/10.3378/027.084.0203

Betti, L., von Cramon-Taubadel, N., Manica, A., & Lycett, S. J. (2013). Global geometric morphometric analyses of the human pelvis reveal substantial neutral population history effects, even across sexes. *PLoS ONE*, *8*(2), Article e55909. https://doi.org/10.1371/journal.pone.0055909

Cann, H. M., de Toma, C., Cazes, L., Legrand, M.-F., Morel, V., Piouffre, L., Bodmer, J., Bodmer, W. F., Bonne-Tamir, B., Cambon-Thomsen, A., Chen, Z., Chu, J., Carcassi, C., Contu, L., Du, R., Excoffier, L., Ferrara, G. B., Friedlaender, J. S., Groot, H., … Cavalli-Sforza, L. L. (2002). A Human Genome Diversity Cell Line Panel. *Science*, *296*(5566), 261-262. https://doi.org/10.1126/science.296.5566.261b

Cenac, Z. (2022). *Indicators of migration from Africa in the human cranium: Dimorphism and diversity* [Unpublished manuscript]. https://www.researchgate.net/publication/340935838_Indicators_of_migration_from_Africa_in_the_human_cranium_Dimorphism_and_diversity

Chen, Y. (2016). Spatial autocorrelation approaches to testing residuals from least squares regression. *PLoS ONE*, *11*(1), Article e0146865. https://doi.org/10.1371/journal.pone.0146865

Chiaroni, J., Underhill, P. A., & Cavalli-Sforza, L. L. (2009). Y chromosome diversity, human expansion, drift, and cultural evolution. *Proceedings of the National Academy of Sciences of the United States of America*, *106*(48), 20174-20179. https://doi.org/10.1073/pnas.0910803106

Choudhury, A., Sengupta, D., Ramsay, M., & Schlebusch, C. (2021). Bantu-speaker migration and admixture in southern Africa. *Human Molecular Genetics*, *30*(2), R56-R63. https://doi.org/10.1093/hmg/ddaa274

Conrad, D. F., Jakobsson, M., Coop, G., Wen, X., Wall, J. D., Rosenberg, N. A., & Pritchard, J. K. (2006). A worldwide survey of haplotype variation and linkage disequilibrium in the human genome. *Nature Genetics*, *38*(11), 1251-1260. https://doi.org/10.1038/ng1911

Creanza, N., Ruhlen, M., Pemberton, T. J., Rosenberg, N. A., Feldman, M. W., & Ramachandran, S. (2015). A comparison of worldwide phonemic and genetic variation in human populations. *Proceedings of the National Academy of Sciences of the United States of America*, *112*(5), 1265-1272. https://doi.org/10.1073/pnas.1424033112

Deshpande, O., Batzoglou, S., Feldman, M. W., & Cavalli-Sforza, L. L. (2009). A serial founder effect model for human settlement out of Africa. *Proceedings of the Royal Society B: Biological Sciences*, *276*(1655), 291-300. https://doi.org/10.1098/rspb.2008.0750

Gaetano, J. (2013). *Holm-Bonferroni sequential correction: An EXCEL calculator*. https://www.researchgate.net/publication/236969037_Holm-Bonferroni_Sequential_Correction_An_EXCEL_Calculator

Gallego Llorente, M., Jones, E. R., Eriksson, A., Siska, V., Arthur, K. W., Arthur, J. W., Curtis, M. C., Stock, J. T., Coltorti, M., Pieruccini, P., Stretton, S., Brock, F., Higham, T., Park, Y., Hofreiter, M., Bradley, D. G., Bhak, J., Pinhasi, R., & Manica, A. (2015). Ancient Ethiopian genome reveals extensive Eurasian admixture in Eastern Africa. *Science*, *350*(6262), 820-822. https://doi.org/10.1126/science.aad2879

Green, R. E., Krause, J., Briggs, A. W., Maricic, T., Stenzel, U., Kircher, M., Patterson, N., Li, H., Zhai, W., Fritz, M., H.-Y., Hansen, N. F., Durand, E. Y., Malaspinas, A.-S., Jensen, J. D., Marques-Bonet, T., Alkan, C., Prüfer, K., Meyer, M., Burbano, H. A., … Pääbo, S. (2010). A draft sequence of the Neandertal genome. *Science*, *328*(5979), 710-722. https://doi.org/10.1126/science.1188021

Haber, M., Mezzavilla, M., Bergström, A., Prado-Martinez, J., Hallast, P., Saif-Ali, R., Al-Habori, M., Dedoussis, G., Zeggini, E., Blue-Smith, J., Wells, R. S., Xue, Y., Zalloua, P. A., & Tyler-Smith, C. (2016). Chad genetic diversity reveals an African history marked by multiple Holocene Eurasian migrations. *The American Journal of Human Genetics*, *99*(6), 1316-1324. https://doi.org/10.1016/j.ajhg.2016.10.012

Hammer, M. F., Karafet, T. M., Redd, A. J., Jarjanazi, H., Santachiara-Benerecetti, S., Soodyall, H., & Zegura, S. L. (2001). Hierarchical patterns of global human Y-chromosome diversity. *Molecular Biology and Evolution*, *18*(7), 1189-1203. https://doi.org/10.1093/oxfordjournals.molbev.a003906

Henn, B. M., Botigué, L. R., Gravel, S., Wang, W., Brisbin, A., Byrnes, J. K., Fadhlaoui-Zid, K., Zalloua, P. A., Moreno-Estrada, A., Bertranpetit, J., Bustamante, C. D., & Comas, D., (2012). Genomic ancestry of North Africans supports back-to-Africa migrations. *PLoS Genetics*, *8*(1), Article e1002397. https://doi.org/10.1371/journal.pgen.1002397

Henn, B. M., Bustamante, C. D., Mountain, J. L., & Feldman, M. W. (2011). Reply to Hublin and Klein: Locating a geographic point of dispersion in Africa for contemporary humans.

Proceedings of the National Academy of Sciences of the United States of America, *108*(28), Article E278. https://doi.org/10.1073/pnas.1107300108

Henn, B. M., Cavalli-Sforza, L. L., & Feldman, M. W. (2012). The great human expansion. *Proceedings of the National Academy of Sciences of the United States of America*, *109*(44), 17758-17764. https://doi.org/10.1073/pnas.1212380109

Henn, B. M., Gignoux, C. R., Jobin, M., Granka, J. M., Macpherson, J. M., Kidd, J. M., Rodríguez-Botigué, L., Ramachandran, S., Hon, L., Brisbin, A., Lin, A. A., Underhill, P. A., Comas, D., Kidd, K. K., Norman, P. J., Parham, P., Bustamante, C. D., Mountain, J. L., & Feldman, M. W. (2011). Hunter-gatherer genomic diversity suggests a southern African origin for modern humans. *Proceedings of the National Academy of Sciences of the United States of America*, *108*(13), 5154-5162. https://doi.org/10.1073/pnas.1017511108

Hijmans, R. J. (2019). *geosphere: Spherical trigonometry* (Version 1.5-10). https://CRAN.R-project.org/package=geosphere

Hodgson, J. A., Mulligan, C. J., Al-Meeri, A., & Raaum, R. L. (2014). Early back-to-Africa migration into the Horn of Africa. *PLoS Genetics*, *10*(6), Article e1004393. https://doi.org/10.1371/journal.pgen.1004393

Holm, S. (1979). A simple sequentially rejective multiple test procedure. *Scandinavian Journal of Statistics*, *6*(2), 65-70.

Howells, W. W. (1973). *Cranial variation in man: A study by multivariate analysis of patterns of difference among recent human populations. Papers of the Peabody Museum of Archaeology and Ethnology Volume 67.* Harvard University.

Howells, W. W. (1989). *Skull shapes and the map: Craniometric analyses in the dispersion of Modern Homo. Papers of the Peabody Museum of Archaeology and Ethnology Volume 79.* Harvard University.

Howells, W. W. (1995). *Who's who in skulls: Ethnic identification of crania from measurements. Papers of the Peabody Museum of Archaeology and Ethnology Volume 82.* Harvard University.

Howells, W. W. (1996). Howells' craniometric data on the internet. *American Journal of Physical Anthropology*, *101*(3), 441-442. https://doi.org/10.1002/ajpa.1331010302

Hunley, K., Bowern, C., & Healy, M. (2012). Rejection of a serial founder effects model of genetic and linguistic coevolution. *Proceedings of the Royal Society B: Biological Sciences*, *279*(1736), 2281-2288. https://doi.org/10.1098/rspb.2011.2296

Hunley, K. L., & Cabana, G. S. (2016). Beyond serial founder effects: The impact of admixture and localized gene flow on patterns of regional genetic diversity. *Human Biology*, *88*(3), 219-231. https://doi.org/10.13110/humanbiology.88.3.0219

Hunley, K., Gwin, K., & Liberman, B. (2016). A reassessment of the impact of European contact on the structure of Native American genetic diversity. *PLoS ONE*, *11*(8), Article e0161018. https://doi.org/10.1371/journal.pone.0161018

Hunley, K., & Healy, M. (2011). The impact of founder effects, gene flow, and European admixture on Native American genetic diversity. *American Journal of Physical Anthropology*, *146*(4), 530-538. https://doi.org/10.1002/ajpa.21506

Ingman, M., & Gyllensten, U. (2006). mtDB: Human Mitochondrial Genome Database, a resource for population genetics and medical sciences. *Nucleic Acids Research*, *34*(Database issue), D749-D751. https://doi.org/10.1093/nar/gkj010

Jakobsson, M., Scholz, S. W., Scheet, P., Gibbs, J. R., VanLiere, J. M., Fung, H.-C., Szpiech, Z. A., Degnan, J. H., Wang, K., Guerreiro, R., Bras, J. M., Schymick, J. C., Hernandez, D. G., Traynor, B. J., Simon-Sanchez, J., Matarin, M., Britton, A., van de Leemput, J., Rafferty, I., ... Singleton, A. B. (2008). Genotype, haplotype and copy-number variation in worldwide human populations. *Nature*, *451*(7181), 998-1003. https://doi.org/10.1038/nature06742

Jorde, L. B., Watkins, W. S., Bamshad, M. J., Dixon, M. E., Ricker, C. E., Seielstad, M. T., & Batzer, M. A. (2000). The distribution of human genetic diversity: A comparison of mitochondrial, autosomal, and Y-chromosome data. *American Journal of Human Genetics*, *66*(3), 979-988. https://doi.org/10.1086/302825

Kim, S. (2015). ppcor: An R package for a fast calculation to semi-partial correlation coefficients. *Communications for Statistical Applications and Methods*, *22*(6), 665-674. https://doi.org/10.5351/CSAM.2015.22.6.665

Li, J. Z., Absher, D. M., Tang, H., Southwick, A. M., Casto, A. M., Ramachandran, S., Cann, H. M., Barsh, G. S., Feldman, M., Cavalli-Sforza, L. L., & Myers, R. M. (2008). Worldwide human relationships inferred from genome-wide patterns of variation. *Science*, *319*(5866), 1100-1104. https://doi.org/10.1126/science.1153717

López, S., van Dorp, L., & Hellenthal, G. (2015). Human dispersal out of Africa: A lasting debate. *Evolutionary Bioinformatics Online*, *11*(S2), 57-68. https://doi.org/10.4137/EBO.S33489

López Herráez, D., Bauchet, M., Tang, K., Theunert, C., Pugach, I., Li, J., Nandineni, M. R., Gross, A., Scholz, M., & Stoneking, M. (2009). Genetic variation and recent positive selection in worldwide human populations: Evidence from nearly 1 million SNPs. *PLoS ONE*, *4*(11), Article e7888. https://doi.org/10.1371/journal.pone.0007888

Luca, F., Hudson, R. R., Witonsky, D. B., & Di Rienzo, A. (2011). A reduced representation approach to population genetic analyses and applications to human evolution. *Genome Research*, *21*(7), 1087-1098. https://doi.org/10.1101/gr.119792.110

Manica, A., Amos, W., Balloux, F., & Hanihara, T. (2007). The effect of ancient population bottlenecks on human phenotypic variation. *Nature*, *448*(7151), 346-348. https://doi.org/10.1038/nature05951

Masson, M. E. J. (2011). A tutorial on a practical Bayesian alternative to null-hypothesis significance testing. *Behavior Research Methods*, *43*, 679-690. https://doi.org/10.3758/s13428-010-0049-5

Mayer, M. (2022). *confintr: Confidence Intervals* (Version 0.1.2). https://CRAN.R-project.org/package=confintr

Pemberton, T. J., DeGiorgio, M., & Rosenberg, N. A. (2013). Population structure in a comprehensive genomic data set on human microsatellite variation. *G3: Genes, Genomes, Genetics*, *3*(5), 891-907. https://doi.org/10.1534/g3.113.005728

Pickrell, J. K., Patterson, N., Loh, P.-R., Lipson, M., Berger, B., Stoneking, M., Pakendorf, B., & Reich, D. (2014). Ancient west Eurasian ancestry in southern and eastern Africa. *Proceedings of the National Academy of Sciences of the United States of America*, *111*(7), 2632-2637. https://doi.org/10.1073/pnas.1313787111

Pickrell, J. K., & Reich, D. (2014). Toward a new history and geography of human genes informed by ancient DNA. *Trends in Genetics*, *30*(9), 377-389. https://doi.org/10.1016/j.tig.2014.07.007

Prugnolle, F., Manica, A., & Balloux, F. (2005). Geography predicts neutral genetic diversity of human populations. *Current Biology*, *15*(5), R159-R160. https://doi.org/10.1016/j.cub.2005.02.038

R Core Team (2020). *R: A Language and Environment for Statistical Computing* (Version 4.0.3) [Computer software]. https://www.R-project.org

R Core Team (2021). *R: A Language and Environment for Statistical Computing* (Version 4.0.5) [Computer software]. https://www.R-project.org

Ramachandran, S., Deshpande, O., Roseman, C. C., Rosenberg, N. A., Feldman, M. W., & Cavalli-Sforza, L. L. (2005). Support from the relationship of genetic and geographic distance in human populations for a serial founder effect originating in Africa. *Proceedings of the National Academy of Sciences of the United States of America*, *102*(44), 15942-15947. https://doi.org/10.1073/pnas.0507611102

Ray, N., Currat, M., Berthier, P., & Excoffier, L. (2005). Recovering the geographic origin of early modern humans by realistic and spatially explicit simulations. *Genome Research*, *15*(8), 1161-1167. https://doi.org/10.1101/gr.3708505

Romero, I. G., Manica, A., Goudet, J., Handley, L. L., & Balloux, F. (2009). How accurate is the current picture of human genetic variation? *Heredity*, *102*, 120-126. https://doi.org/10.1038/hdy.2008.89

Rosenberg, N. A., Pritchard, J. K., Weber, J. L., Cann, H. M., Kidd, K. K., Zhivotovsky, L. A., & Feldman, M. W. (2002). Genetic structure of human populations. *Science*, *298*(5602), 2381-2385. https://doi.org/10.1126/science.1078311

Savin, N. E., & White, K. J. (1977). The Durbin-Watson test for serial correlation with extreme sample sizes or many regressors. *Econometrica*, *45*(8), 1989-1996. https://doi.org/10.2307/1914122

Schlebusch, C. M., Skoglund, P., Sjödin, P., Gattepaille, L. M., Hernandez, D., Jay, F., Li, S., De Jongh, M., Singleton, A., Blum, M. G. B., Soodyall, H., & Jakobsson, M. (2012). Genomic variation in seven Khoe-San groups reveals adaptation and complex African history. *Science*, *338*(6105), 374-379. https://doi.org/10.1126/science.1227721

Seielstad, M., Bekele, E., Ibrahim, M., Touré, A., & Traoré, M. (1999). A view of modern human origins from Y chromosome microsatellite variation. *Genome Research*, *9*(6), 558-567.

Shi, W., Ayub, Q., Vermeulen, M., Shao, R.-g., Zuniga, S., van der Gaag, K., de Knijff, P., Kayser, M., Xue, Y., & Tyler-Smith, C. (2010). A worldwide survey of human male demographic history based on Y-SNP and Y-STR data from the HGDP–CEPH populations. *Molecular Biology and Evolution*, *27*(2), 385-393. https://doi.org/10.1093/molbev/msp243

Smith, R. J. (1999). Statistics of sexual size dimorphism. *Journal of Human Evolution*, *36*(4), 423-459. https://doi.org/10.1006/jhev.1998.0281

Tishkoff, S. A., Reed, F. A., Friedlaender, F. R., Ehret, C., Ranciaro, A., Froment, A., Hirbo, J. B., Awomoyi, A. A., Bodo, J.-M., Doumbo, O., Ibrahim, M., Juma, A. T., Kotze, M. J., Lema, G., Moore, J. H., Mortensen, H., Nyambo, T. B., Omar, S. A., Powell, K., … Williams, S. M. (2009). The genetic structure and history of Africans and African Americans. *Science*, *324*(5930), 1035-1044. https://doi.org/10.1126/science.1172257

Van Tuyl, R., & Pereltsvaig, A. (2012). Comment on "Phonemic Diversity Supports a Serial Founder Effect Model of Language Expansion from Africa". *Science*, *335*(6069), 657-d. https://doi.org/10.1126/science.1209176

von Cramon-Taubadel, N., & Lycett, S. J. (2008). Brief communication: Human cranial variation fits iterative founder effect model with African origin. *American Journal of Physical Anthropology*, *136*(1), 108-113. https://doi.org/10.1002/ajpa.20775

Wang, C.-C., & Li, H. (2013). Inferring human history in East Asia from Y chromosomes. *Investigative Genetics*, *4*, Article 11. https://doi.org/10.1186/2041-2223-4-11

Williams, E. (2011). *Aviation formulary V1.46*. http://edwilliams.org/avform.htm