***Streptococcus suis* infection on European farms is associated with an altered tonsil**
microbiome and resistome

Simen Fredriksen[1*], Carlos Neila-Ibáñez[2,3,4], Isabel Hennig-Pauka[5], Xiaonan Guan[1,6], Jenelle
Dunkelberger[7], Isabela Fernandes de Oliveira[1], Maria Laura Ferrando[1], Florencia Correa-
Fiz[2,3,4], Virginia Aragon[2,3,4], Jos Boekhorst[1], Peter van Baarlen[1], Jerry M. Wells[1*]

1. Host-Microbe Interactomics Group, Animal Sciences Department, Wageningen University,
Wageningen, The Netherlands
2. Unitat mixta d'Investigació IRTA-UAB en Sanitat Animal, Centre de Recerca en Sanitat
Animal (CReSA), Campus de la Universitat Autònoma de Barcelona (UAB), Bellaterra,
08193, Catalonia, Spain
3. IRTA. Programa de Sanitat Animal. Centre de Recerca en Sanitat Animal (CReSA).
Campus de la Universitat Autònoma de Barcelona (UAB), Bellaterra, 08193, Catalonia,
Spain
4. OIE Collaborating Centre for the Research and Control of Emerging and Re-emerging
Swine Diseases in Europe (IRTA-CReSA), Bellaterra, 08193 Barcelona, Spain
5. Field Station for Epidemiology, University of Veterinary Medicine Hannover, 49456
Bakum, Germany
6. Schothorst Feed Research B.V., Lelystad, The Netherlands
7. Topigs Norsvin USA, Burnsville, MN, United States

*simen.fredriksen@gmail.com, jerry.wells@wur.nl

23
24

Abstract

Introduction

Streptoccocus suis is a Gram-positive opportunistic pathogen causing systemic disease in

piglets around weaning age. Outbreaks of *S. suis* disease are controlled by metaphylactic use

of antibiotics, leading to high levels of antimicrobial resistance in *S. suis* isolates. This is an

issue for both animal and human health due to the zoonotic disease potential of *S. suis*. The

mechanisms facilitating invasive disease are not known but may involve host and

environmental factors. The palatine tonsils are considered a portal of entry for pathogenic

strains to cause systemic disease. We hypothesised that tonsil colonization by pathogenic and

commensal bacteria may impact on disease risk via colonization resistance and co-infections.

We conducted a case-control study on 9 European farms, comparing the tonsil microbiome of

piglets with *S. suis* systemic disease with asymptomatic controls. We also compared these to

piglets on control farms and piglets reared naturally in a forest.

Results

We found a small but significant difference in the tonsil microbiota composition of case and

control piglets. Case-control associations varied between amplicon sequence variants (ASVs)

and metagenome assembled genomes (MAGs) within the same species. Variants of putatively

commensal taxa including *Rothia nasimurium* were reduced in abundance in case piglets

compared to asymptomatic controls. Case piglets had higher relative abundance of

Fusobacterium gastrosuis, *Bacteroides heparinolyticus*, and uncultured *Prevotella* and

Alloprevotella species. There was, however, no higher abundance of *S. suis* itself at the

species-level or of clinical strain marker genes in case piglets. Piglets sampled prospectively

weeks prior to developing clinical signs had reduced microbiota alpha diversity. Despite

case-control pairs receiving equal antimicrobial treatment, case piglets had higher abundance

of antimicrobial resistance genes (ARGs) conferring resistance to antimicrobial classes used

to treat *S. suis*.

Conclusions

The tonsillar microbiota of *S. suis* case piglets had increased abundance of taxa not

previously linked to *S. suis* disease. This coincided with increased ARG abundance in case

piglets, possibly due to adaptation of the disease-associated microbiota to frequent

antimicrobial treatment.

Introduction

Streptococcus suis is a Gram-positive bacterium colonizing the upper respiratory tract of pigs. It is one of the major bacterial causes of disease in pigs and a zoonotic pathogen causing sepsis and meningitis in humans [1–4]. Swine infections are prevented by metaphylactic use of antimicrobials. This has led to increased antimicrobial resistance (AMR) in *S. suis* isolates [5], with macrolide and tetracycline resistance genes *erm(B)* and *tet(O)* being the most common [6]. The spread of antimicrobial resistance genes (ARGs) in zoonotic *S. suis* and to other streptococci is of concern to veterinary and human medicine [7].

The palatine tonsils have been suggested as a main habitat for *S. suis* colonization and a putative site of entry into the host bloodstream [8–15]. A recent study identified differences in taxonomic composition of the tonsillar microbiota in piglets diagnosed with *S. suis* disease [16]. Microbiota associations with *S. suis* infectious disease are of great interest to understand co-infection dynamics and to identify probiotic candidate species providing colonization resistance. It is thought that co-infection by bacterial and viral pathogens part of the porcine respiratory disease complex (PRDC) may predispose to *S. suis* invasive disease [17–19].

Tonsillar colonization by different *S. suis* strains may also impact on invasive disease risk [13, 20]. While *S. suis* has been described as an opportunistic pathogen, different strains have varying virulent potential and can be grouped into commensal and pathogenic clades [21–23]. Strains from pathogenic *S. suis* clades have reduced genome sizes [22] and putatively reduced ability to persist as colonizers in competition with commensals. Strains from pathogenic clades can also be isolated from the tonsils of healthy pigs, but it is not known whether *S. suis* invasive disease is preceded by outgrowth of disease-associated strains.

This study was aimed at identifying the tonsillar microbiota associated with *S. suis* systemic disease in weaning age piglets. We utilized a case-control study design and next generation sequencing to characterise the composition of the tonsillar microbiota of piglets with *S. suis* systemic infection at 9 European farms and compare these to piglets from 4 farms without *S. suis* disease. We used 16S rRNA gene amplicon- and metagenomic shotgun sequencing to quantify the tonsillar microbiota and resistome and analysed clinical and non-clinical *S. suis* strains from the farms by whole genome sequencing. This study design allowed us to assess microbiome composition and predict functionality, as well as to assess the abundance of marker genes prevalent in clinical or non-clinical *S. suis* strains.

Results

Sequencing

We sampled 45 case-control pairs of piglets with *S. suis* systemic clinical signs and asymptomatic control piglets from the same pen. Case-control pairs were equivalent in age, genetic background, and antimicrobial treatment. Additional control piglets from case farms and from control farms without *S. suis* outbreaks were sampled to assess farm differences. We also sampled one US farm and piglets raised organically in a forest for reference (see supplementary file 1 for information about the farms). Amplicon sequencing of the full set of 295 samples yielded an average of 76757 reads per sample after processing with DADA2 [24]. Metagenomic DNA from all case-control pairs and a subset of piglets from control farms was shotgun sequenced. A total of 109 metagenomic samples yielded on average 7.5 million reads per sample after removal of host DNA and plant DNA from feed.

The *Streptococcus suis* disease-associated microbiota

Case and control piglets on outbreak farms had significantly different tonsillar microbiota composition, but the effect size was small ($R^2 = 0.01$ and $p = 0.02$, PERMANOVA on Bray-Curtis dissimilarity comparing case and control piglets within outbreak farms). Analysis per country found varying results for Spain ($R^2 = 0.030$, $p = 0.01$) and Germany ($R^2 = 0.017$, $p = 0.93$), which had fewer samples. The largest difference in tonsil microbiota composition was found on the Dutch farm NL1 ($R^2 = 0.26$, $p < 0.01$), but this might be due to all 3 case piglets from this farm being sampled shortly after death allowing opportunistic pathogens to bloom. We did, however, not see a general association between the microbiota and clinical sign severity or (future) death within the other farms. In general, there was no consistent difference in alpha diversity between case and control piglets (Figure S1).

We identified amplicon sequence variants (ASVs) significantly associated with piglet case-control status (Figure 1A). SIAMCAT [25] identified 3 ASVs with significantly ($p < 0.05$) higher abundance in case piglets: ASV 29 (1.8% vs 0.7% mean abundance, *Prevotella* sp.), ASV 3 (3.6% vs 2.2%, *Alloprevotella* sp.), and ASV 21 (3.6% vs 1.5%, *Fusobacterium gastrosuis*). These were all part of the core microbiota in the European farms (minimum 99% prevalence in control piglets). The case-associations were stronger in the Dutch and Spanish farms, where the outbreaks were larger and/or symptoms more severe than in the German farms. These 3 ASVs also had the strongest association with case piglets using Wilcoxon Rank Sum Test ($p < 0.005$), but no ASVs were significantly different when applying multiple

testing correction (FDR > 0.05). No genera were significantly different between case and

control piglets (FDR > 0.05). Table S1 lists case-control associations for all ASVs and

genera.

While the 3 case-associated ASVs had the strongest case-control associations, few other

ASVs trended towards case-associations. A larger number of different ASVs had weaker,

non-significant (FDR > 0.05), control-associations. Among core microbiota members, ASV

48 (0.38% vs 0.74%, *Clostridium disporicum*), ASV 36 (0.47% vs 0.91%, *Rothia*

nasimurium), and ASV 6 (1.98% vs 2.62%, *S. suis*) had the strongest control associations.

ASV 11 (0.27% vs 0.63%, *Actinobacillus minor*, not present in farms ES4 and NL1) and

ASV 81 (0.22% vs 0.40%, *Terrisporobacter mayombei*) were the non-core ASVs with the

strongest control-associations. In terms of total relative abundance, ASV 1 (3.5% vs 8.1%,

Moraxella porci/pluranimalium) showed the largest difference between case and control

piglets. ASV 1 had strong control associations in farm ES2 (0.9% vs 5.6%, being completely

absent in several case piglets) and ES3 (15% vs 24%) but was absent from farm ES4 and

NL1 and equally abundant in German case and control piglets.

Porcine respiratory disease complex (PRDC) and its range of bacteria thought to co-infect

with *S. suis* is mainly associated with respiratory disease in older pigs, while this study

investigates weaning age systemic disease. Still, PRDC-associated bacteria were of interest in

possible co-infections. *Streptococcus suis* itself and *Glaesserella parasuis* were among the

most abundant tonsil microbiota members. Relative abundance of *S. suis* ASVs trended to be

higher in healthy controls, while *G. parasuis* ASVs had mixed associations. ASV 174 (100%

identical to both *G. parasuis* and *Actinobacillus indolicus*) trended towards case-association

(0.27% vs 0.16%). *Pasteurella multocida* was relatively prevalent (50%) but at low

abundance (mean 0.06%), although two outlier case piglets had over 1% abundance.

Actinobacillus pleuropneumoniae had 34% prevalence and 0.08% mean abundance but was

most abundant in control piglets. *Bordetella bronchiseptica* had lower prevalence (17%) but

higher abundance (0.10%) due to some control piglets having up to 12% abundance.

Mycoplasma hyopneumoniae was virtually absent from the dataset, with only 6 reads from a

single control piglet.

Figure 1: Microbiota case-control associations. **A.** ASVs significantly associated with case-control status on the outbreak farms, sorted by strongest case-control association with Wilcoxon Rank Sum Test. Top row: control-associated ASVs, bottom rom: case-associated ASVs. **B.** Shannon diversity (amplicon sequencing) of prospective samples collected pre-weaning from piglets that developed *S. suis* clinical signs 2-4 weeks later. Samples from future case piglets had lower Shannon diversity than control piglets which remained asymptomatic. Piglets from control farm ES1, which was managed by the same company and used sows from ES2, also had high mean diversity.

Early life microbiota diversity predicted clinical sign appearance weeks later

To assess the value of tonsillar microbiota composition to predict future invasive *S. suis* disease, we collected prospective samples on farm ES2, which had a history of post-weaning *S. suis* outbreaks, and farm ES1, a control farm without recorded *S. suis* problems despite use of sows produced on farm ES2. A cohort of 40 piglets were sampled on farm ES2 1 week before weaning, prior to visible clinical signs. Eight of the 40 piglets developed *S. suis* clinical signs 1-3 weeks post-weaning, and tonsil samples collected from these piglets post-weaning were included in the 45 case-control pairs.

Comparison of prospective samples collected from case and control piglets 1 week before weaning, i.e., 2-4 weeks before disease onset, revealed that the tonsil microbiota of case piglets had significantly lower Shannon diversity compared to control piglets that remained asymptomatic ($p = 0.005$; Figure 1B). Case piglets trended towards having lower diversity also post-weaning, during the outbreak, but the effect was smaller and less significant ($p = 0.12$). Asymptomatic siblings of case piglets also had a lower diversity than control piglets from litters without case piglets and piglets from control farm ES1 (Figure 1B). The difference in composition was, however, smaller 1 week pre-weaning ($R2 = 0.06$, $p = 0.15$,

PERMANOVA on Bray-Curtis dissimilarity) than during the outbreak (R2 = 0.13, p < 0.01).

The Spearman correlation between ASV ratio of case-control abundance pre- and post-

weaning on the farm was low (R = 0.09, p = 0.4), showing that separate ASVs were

differentially abundant and driving case-control compositional differences pre- and post-

weaning. The strongest pre-weaning case-associations were ASV 88 (*Pasteurellaceae*, 1.4%

vs 0.55%) and ASV 186 (*Actinobacillus indolicus/minor*, 1.1% vs 0.21%), while ASV 77

(Leptotrichia, 0.11% vs 0.33%) and ASV 22 (Actinobacillus minor, 0.06% vs 0.20%) had the

strongest control associations.

Metagenome-assembled genomes (MAG) analysis

We shotgun sequenced all case-control pairs and a subset of piglets from control farms. Co-

assembly per farm produced 802 metagenome-assembled genomes (MAGs) (>70%

completeness and <10% contamination). Read mapping to the MAGs yielded an overall case-

control compositional difference similar to that found by amplicon sequencing analysis

(Figure 2, R = 0.03, p < 0.01, Bray-Curtis dissimilarity PERMANOVA). The case-control

associated MAGs largely corresponded to the ASVs identified by amplicon-based analysis,

but some taxa, including case-associated *Bacteroides heparinolyticus*, were identified only

using MAGs. The MAG approach resulted in stronger case-control associations (FDR <

0.0001) than the ASV approach, suggesting that the case-associated ASVs represented

several strains with variable case-control associations. The two ASVs with strongest disease-

association putatively corresponded to a collection of different *Bacteroidales* MAGs. The

MAGs with the strongest disease-associations were classified as *Bacteroidales* UBA1309 and

Alloprevotella F0040, clades poorly described and with few or no publicly available related

isolate genomes. Case-control association of all MAGs are listed in Table S2.

Figure 2: Redundancy analysis (RDA) on the abundance of metagenome-assembled genomes (MAGs) constrained by case-control status. The points represent individual samples; samples in the direction of arrows have more of that taxon. The eclipse represents 75% confidence level.

The tonsillar microbiome of piglets is similar between farms but diverged in free-range forest piglets

The tonsillar microbiota composition was more similar between piglets on the same farm than to that of piglets from other farms (Figure 3A-B). Compared to tonsillar microbiota compositions of farm piglets, Tamworth free-range piglets living outdoors in a forest in The Netherlands had a strikingly different microbiota and resistome (figure 3C-D). While European farm piglets shared a large core microbiota with 83 ASVs being present in 80% of piglets or more, 44 of these were not found in the free-range piglets. Vice versa of 157 ASVs found in all 5 forest piglets, 117 were not found in any farm piglet. Free-range piglets had low abundance of the genera most abundant in farm piglets, in particular *Moraxella* (0.6% vs 13%) and *Streptococcus* (0.9% vs 12%), and higher abundance of Acinetobacter (10% vs

2.3%), "*Rikenellaceae* RC9 gut group" (9.9% vs 0.1%), and *Treponema pedis* (3.6% vs

0.04%). The sample with the lowest *S. suis* abundance in the study, 0.3% abundance of a

single ASV (ASV 1050), was from a free-range piglet. This ASV did not have 100% identity

to the 16S rRNA gene V3-V4 region of any *S. suis* strain publicly available in SILVA or

NCBI assembly databases. The other free-range piglets were colonised by *S. suis* ASVs

shared with farm piglets.

Figure 3: Microbiome differences between farms. **A.** PCA on microbiota composition (ASV

abundance) of all farm samples and countries, pre- and post-weaning (transformed with

log(1000*abundance+1)). Samples clustered broadly by country, but samples from farm DE6

clustered with NL1, and samples from ES4 clustered away from the other Spanish farms. **B.**

Mean pairwise Bray-Curtis dissimilarity between samples from the different farms. **C.** PCA

on ARG abundance (transformed with log(1000*abundance+1)). The pre- and post-weaning

Spanish samples separated in 2 clusters. **D.** Total abundance of all ARGs per farm.

Increased tetracycline ARG abundance in case piglets

We quantified the abundance of antibiotic resistance genes (ARGs), collectively called the resistome, by mapping metagenomic reads to the Resfinder database [26] and normalizing abundance by fragments per kilobase reference per million fragments (FPKM). Piglets on most farms received antibiotics via feed or water, and these were included for resistome analysis. Twenty-two piglets received intramuscular injections, and these were excluded from the main analysis. All case-control pairs included in the main analysis had received equal antimicrobial treatment. Farms varied in total ARG abundance (Figure 3D), but shared high abundances of the most abundant ARGs, such as *blaROB-1* conferring resistance to penicillin/amoxicillin/ampicillin (411 FPKM mean abundance), *sul2* conferring sulfamethoxazole resistance (338 FPKM), and streptomycin resistance genes *aph(3'')-Ib* (312 FPKM) and *aph(6)-Id* (220 FPKM). Table S3 lists all detected ARGs per farm. The ARGs most common in *S. suis*, *erm(B)* and *tet(O)* [6], were less abundant (66 and 111 FPKM, respectively). When comparing the overall ARG composition by PCA, samples clustered by country, except for Spanish pre- and post-weaning samples clustering separately by age and not by farm (Figure 3C).

ARG abundance may, in addition to by taxonomic composition, be influenced by both historical antimicrobial usage on the farm and by direct antimicrobial treatment of the sampled piglets. This study could not disentangle these two effects since most antimicrobial treatments were given equally to all sampled piglets at each farm. Farm NL1, a high health status research farm where the sampled piglets were not treated, and where piglets are rarely treated with antimicrobials, had the lowest ARG abundance (except for the free-range forest piglets). In Germany, high health status farms DE1 and DE2 had low ARG abundance, but so did DE6 despite a history of severe *S. suis* disease. Farms DE3, ES3, and ES4 were assessed to have the lowest health status and highest historical antimicrobial usage by veterinarians (supplementary file 1), and piglets on these farms were also administered antimicrobials before sampling. These three farms had the highest ARG abundance (Figure 3D). There was no consistent link between the antimicrobials administered and the abundance of ARGs conferring resistance to these. On farm DE1, piglets that had received tetracycline had lower tetracycline ARG abundance than untreated piglets (810 vs 391 FPKM), and on DE3, where all piglets had received tetracycline, tetracycline ARG abundance was lower than on DE2 and DE5 where piglets had received no antimicrobial treatment (804 vs 805 and 926 FPKM, respectively).

The total abundance of ARGs was 15% higher in case piglets compared to controls (3473 vs

3026 mean FPKM), but this overall effect was not statistically significant (p = 0.36). ARGs

conferring resistance to tetracycline (class level) were strongly case-associated (788 vs 620

FPKM, p = 0.01), especially within farms ES2 and ES3. Specifically, tetracycline and

doxycycline resistance, conferred largely by the same ARGs, were the most abundant

Resfinder ARG phenotype categories and had the strongest case-associations (Figure 4).

Tet(Q) was the individual gene with the strongest case-association (134 vs 75 FPKM, FDR <

0.01). ARGs of classes aminoglycoside, beta-lactam, and macrolide, which like tetracyclines

are commonly used to treat *S. suis* disease, also trended towards higher abundance in case

piglets (Table S4).

We investigated the presence of case-associated ARGs in MAGs to determine if the higher

ARG abundance was linked to the case-associated taxa. *Tet(Q)* was only found in case-

associated *Prevotella* MAGs, and case-associated ARGs *ant(6)-Ib* and *tet(44)* were only

found in case-associated *Fusobacteriales* MAGs. ARGs found in control-associated *Rothia*

and *Clostridium* MAGs, *lnu(P)*, *erm(Q)*, *tetA(P)*, and *erm(X)*, were not case-associated.

Figure 4: The abundance of 12 most abundant Resfinder ARG phenotype categories in case

and control piglets. Note that most genes confer resistance to several antimicrobials, making

the sum of phenotype abundances greater than the total ARG abundance. For instance,

tetracycline and doxycycline resistance is conferred by the same genes.

Comparison of the tonsil and nasal microbiota

To evaluate whether differences in microbiota composition between case and control piglets

were specific to the tonsillar microbiota or a general trend in the oropharyngeal cavity, we

collected 41 nasal swabs from three Spanish farms, ES1-3. Nasal and tonsillar swab

microbiota composition was significantly different, but more variation was explained by the

farm (R2 = 0.08, p < 0.001, Bray-Curtis PERMANOVA) than nasal/tonsil sample location

(R2 = 0.06, p < 0.001). The nasal microbiota was characterised by higher abundance of

genera *Moraxella* (32% vs 16%) and *Bergeyella* (5% vs 1%) and lower abundance of most

other genera. Case and control piglets had a significantly different nasal microbiota (R2 =

0.07, p = 0.04, Bray-Curtis PERMANOVA). This effect size was comparable to the

difference found in the tonsillar microbiota samples for the same subset of piglets (R2 = 0.07,

p = 0.09). However, the ASVs that were differentially abundant in the nasal microbiota of

case and control piglets were not the same ASVs that were differentially present in tonsillar

microbiota. Some ASVs had inverse case-control associations: ASV 21 (*F. gastrosuis*), ASV

29 (*Prevotella* sp.), and ASV 3 (*Alloprevotella* sp.) were disease-associated in the tonsillar

microbiota, both overall and within ES2 and ES3, but health-associated in the nasal

microbiota.

S. suis diversity in the tonsillar microbiota

Amplicon sequencing data showed higher *S. suis* abundance in control piglets compared to

case piglets on farms with pre-weaning outbreaks (farms ES3 and ES4, 3.9% vs 6.4%, p =

0.02). This effect was not significant on post-weaning outbreak farms (DE1, DE3-6, ES2,

NL1, 4.4% vs 5.1%, p = 0.27).

In total 89 ASVs were classified as *S. suis*, and 52 of these were present in 2 or more tonsil

samples, suggesting they are unlikely to be sequencing artefacts. Farm piglets were on

average colonised by 7 *S. suis* ASVs, with a range between 2 and 16. *Streptococcus suis*

relative abundance was similar in outbreak and control farms (Figure 5A). Comparison of the

89 *S. suis* ASVs to the 16S rRNA gene V3-V4 region of 2463 *S. suis* assemblies available on

NCBI assembly revealed that many ASVs were poorly or not at all represented by sequenced

genomes. This likely relates to the fact that many clinical but few non-clinical strains have

been sequenced. ASVs are, however, not good markers for assessing *S. suis* strain diversity as

the 16S rRNA gene V3-V4 region correlates poorly with whole genome phylogeny, and as

both clinical and non-clinical strains share the same ASVs. 82% of all *S. suis* assemblies had

a 16S rRNA gene V3-V4 region amplicon identical to ASV 6, which comprised 30% of *S.*

suis in the tonsillar microbiota. ASV 17, the second most abundant *S. suis* ASV, comprised

21% of all *S. suis* in the microbiota but was only found in 0.08% of the assemblies. In

addition to strain DE512T1 collected in the present study, ASV 17 was only found in 2

isolates recently sampled from diseased pigs in China (GCF_019793915.1 and

GCF_019794525.1).

Further, we used metagenomic data to assess the relative proportion of commensal and

pathogenic *S. suis* in the tonsillar microbiota of each piglet. We created a *S. suis* pangenome

by clustering protein coding genes from a previously published genome collection [21] at

80% identity, and mapped metagenomic reads to the representative sequences of each cluster

to assess their abundance in each sample. We calculated the ratio of prevalence of each

cluster in clinical and non-clinical genomes to assess their putative association with

pathogenicity. We found that in the tonsillar microbiota, genes predominantly found in non-

clinical strains occurred at higher abundance than genes common in clinical isolates (figure

5B). There was a small positive correlation between gene clinical/non-clinical genome

presence ratio and case-/control sample abundance ratio (Figure 5C, Spearman R = 0.05, p <

0.01), due to higher abundance of commensal *S. suis* in control samples.

One often used *S. suis* marker gene for strain virulence is the gene encoding suilysin (*sly*),

and this gene was found in at least 1 clinical strain from all farms where we sequenced

clinical strains. *Sly* is well suited for use in metagenomics analysis due to high sequence

conservation across *S. suis* strains. *Sly* abundance was low in the tonsillar metagenomes, and

similarly abundant in case and control piglets (Figure 5D). On some farms we did not detect

sly in any piglets, but this may be due to small sample size and insufficient sequencing depth.

Figure 5: *S. suis* abundance and diversity. **A)** Total *S. suis* relative abundance per farm, amplicon sequencing data. **B)** Within the *S. suis* pangenome, genes associated with non-clinical strains (x-axis) were more abundant (y-axis) in metagenomic samples from the tonsillar microbiota. Genes with ratio 0 are equally common in clinical and non-clinical strains, genes with a ratio of 1 are twice as common in clinical strains. **C)** The correlation between gene association with clinical/non-clinical genomes (x-axis) and association with abundance in case-control metagenomic samples. **D)** The abundance of suilysin encoding gene *sly*, a gene highly conserved in clinical strains, in case and control samples per farm.

Discussion

In this study, we found the tonsil microbiota composition of case piglets with *S. suis* clinical signs to differ significantly from the microbiota of asymptomatic controls. *Streptococcus suis* disease may occur as a part of polymicrobial infections collectively known as porcine respiratory disease complex (PRDC), which includes porcine reproductive and respiratory syndrome (PRRS) virus, swine influenza A and potentially bacterial primary and secondary (opportunistic) pathogens [18, 27]. We did not find disease-associations with the tonsil microbiota abundance of any species linked to PRDC, such as *G. parasuis* or *S. suis* itself.

This study included only piglets with weaning age systemic *S. suis* disease, and PRDC

associated taxa may be more relevant to respiratory disease in older finisher pigs.

We identified novel disease-associations with *Fusobacterium gastrosuis*, *Bacteroides*

heparinolyticus, and uncultured *Prevotella* and *Alloprevotella* species. *Fusobacterium*

gastrosuis has previously been linked to *Helicobacter suis* infection in the gastric microbiota

and shown to have genes involved in adhesion, invasion and induction of cell death as well as

in immune evasion in other *Fusobacterium* species [28]. The uncultured case-associated

Alloprevotella/*Prevotella*/*Bacteroides* species lack isolate genomes and are unknown in

relation to *S. suis* disease, possibly due to being unculturable. Case-associated taxa may also

interact with the host to facilitate *S. suis* to cross epithelial barriers without themselves

entering the bloodstream, thus remaining undetected by necropsy. Alternatively, disease-

associated taxa may increase in abundance due to host immune status and dysbiosis, as

suggested for oral *Prevotella*/*Alloprevotella* species in humans [29].

Case piglets had lower tonsillar *S. suis* abundance than control piglets. We assessed that this

was due to a reduced abundance of strains from commensal clades, most of which are poorly

represented among sequenced *S. suis* strains. While *S. suis* genes predominantly found in

non-clinical isolates were more abundant in control piglets, genes predominantly found in

clinical strains, such as *sly*, were low in abundance and more equally distributed between

cases and controls. This shows that the majority of *S. suis* colonising tonsils are commensal,

lacking genes required for invading the host, but also confirms that strains carrying genes

most prevalent in clinical strains are also colonising asymptomatic piglets at low abundance.

Based on these results, we conclude that tonsillar colonisation by *S. suis* itself cannot be used

to reliably predict or even confirm ongoing *S. suis* invasive disease.

Associations between *S. suis* disease and the tonsillar microbiota have been investigated

using amplicon sequencing in a previous study by Niazy et al. [16]. *Bacteroides* and

Lachnospiraceae were found to be more abundant in controls, while case piglets had higher

abundance of *Campylobacter* and *Porphyromonas*, among others. This may be due to

European and North American piglets having fundamentally different microbiomes.

Differences between studies may largely be due to methodological differences, but in the

present study we included one US farm and found the sampled piglets to have low diversity,

and that while microbiota members were shared at the ASV level, composition was

dominated by high *Actinobacillus* and *Streptococcus* abundance. The different findings in

Niazy et al. might also be due to their sampling of whole tonsillar tissue while we used
swabs. Furthermore, piglets suffering from other pathologies such as rectal prolapse and
hernia were sampled as controls, so the case piglets were not compared with healthy controls
as in the present study. While their study may include control-associated taxa associated with
other disease, our results may not be specific only to *S. suis* disease but include microbiome
traits associated with low health status in general. Niazy et al. also sampled uneven numbers
of case and control piglets per farm, and from some farms no controls. This confoundment of
case-control and farm comparison may have affected the results.

We found case piglets to have higher abundance of antimicrobial resistance genes than
controls, despite case-control pairs being treated with the same antimicrobials. ARGs
conferring resistance to doxycycline and tetracycline had the strongest case-association, and
ARGs conferring resistance to other antimicrobials used against *S. suis* also trended towards
being more abundant in case piglets. ARGs found in MAGs of disease-associated taxa
Fusobacterium, *Prevotella*, and *Alloprevotella* had strong disease-associations compared to
those found in health-associated *Rothia* and *Clostridium* MAGs. It is possible that frequent
antimicrobial usage has caused selection pressure on disease-associated taxa, leading to
accumulation of ARGs.

While antimicrobial treatment may have caused strong selective pressure over time, we found
limited increases in abundance of ARGs conferring resistance to the antimicrobials used to
treat the sampled piglets. It was unexpected to find that antimicrobials administered by water
and feed did not appear to have led to increases in ARG abundance. Antimicrobial usage
starting in the first few days after birth has previously been shown to influence piglet nasal
microbiota composition and diversity [30], but in the present study most antimicrobial
treatment started later in life, and only a few days before sampling. Another plausible
explanation for the limited effect of antimicrobial treatment on the tonsillar microbiota in our
study is intrinsic resistance of biofilms to antibiotics [31]. The route of antimicrobial
administration is known to determine the impact on gut microbiota [32, 33], but it is not
known to what extent antimicrobials provided in water, feed, or by injection are able to
penetrate oral biofilm.

Farms had significant differences in microbiota composition, but clustered by country. This
may be due to both sampler bias (a single person collected all samples in each country) and
farm environment, practices, and regulations that vary by country. The piglet microbiota may

for instance be influenced by factors such as cleaning, feed composition, temperature, and

antimicrobial treatment [34–41]. However, the microbiota composition was not more similar

on farms with comparable management practices. The genetics and source of the sows, and

their vertically transferred microbiota may be a more important determinant of the piglet

microbiota than farm conditions. Among the Spanish farms, ES1 and ES2, which were

operated by the same company and had frequent exchange of animals, had the most similar

microbiota composition to each other. However, unrelated German farms shared a more

similar microbiota composition than any of the Spanish farms. In both Spain and Germany,

farms were spread over a large geographic area and managed by different companies.

We sampled the tonsillar microbiota of 5 piglets living outdoors in a forest in The

Netherlands. These piglets have no recorded problems with diseases commonly affecting pigs

on intensive farms, including *S. suis* associated disease. We found the tonsillar microbiota of

the forest piglets to be fundamentally different from farm piglets, and to have lower ARG

abundance. Core ASVs in farm piglets were completely absent in the free-range piglets, and

core ASVs from the forest piglets were not found in farm piglets. The ecological farm DE4,

where piglets had straw bedding and outside access, did not have a more similar microbiota

to the forest piglets than other farms. Compared to DE4 and other farms, living outdoors may

shape the tonsillar microbiota via exposure to environmental microbes from soil and diverse

natural feed sources, and via high air quality and lower exposure to bacterial transfer from

other piglets. Previous studies have shown differences in the faecal [42–44] and nasal [45]

microbiota of wild pigs, but we are not aware of studies on specific factors that may drive

differences between the natural and domestic microbiota or between farms with varying

disease problems. Understanding these factors may be key to preventing disease by

opportunistic pathogens in pig farming.

We conducted a longitudinal sampling on farm ES2, collecting prospective samples from pre-

weaning, prior to clinical sign development. Future case piglets had reduced alpha diversity

compared to asymptomatic controls and piglets from control farm ES1. Asymptomatic

siblings of case piglets had alpha diversity intermediate between symptomatic siblings and

control piglets from litters without any *S. suis* cases. *Streptococcus suis* disease cases were

concentrated in a limited number of litters, despite most of the herd remaining unaffected.

This suggests that maternal effects involving early life immunity- or microbiota may be

important in predisposing to *S. suis* disease and potential dysbiosis. This may be due to

vertical transmission of a disease-prone microbiota, but also differences in maternal

immunity, with antibodies depleting prematurely [46]. *Streptococcus suis* disease most commonly occurs around the time colostral antibodies to *S. suis* start becoming depleted. Various studies have found other colostral antibodies to have similar half-lives as well as high variation in antibody abundance and half-life between individual piglets and litters of different sows [47–51]. Thus, it is possible that during *S. suis* outbreaks, some piglets have sufficiently high levels of maternal antibodies to opsonise invading *S. suis*, whereas other piglets in some litters lack a sufficient level of maternal immunity against *S. suis*.

In conclusion, there are small but significant differences between the tonsillar microbiota of *S. suis* case piglets and asymptomatic controls. We discovered novel taxa associated with case piglets, while *S. suis* abundance was higher in controls. The microbiota differences may originate from dysbiosis starting during early life prior to disease outbreak, but further research is needed to assert this. It is also not conclusively known whether *S. suis* invades through the tonsils or other locations. Case piglets had higher abundance of ARGs conferring resistance against classes commonly used to treat *S. suis* disease. This may be linked to high ARG prevalence in case-associated taxa driven by more frequent exposure antimicrobial treatment than control-associated taxa.

Materials and methods

Animals

We utilized a case-control study design to assess the association between tonsillar microbiota composition and incidence of *S. suis* invasive disease. To search for consistent correlations between microbiota composition and *S. suis* disease incidence we included farms from different countries with different livestock management systems (supplementary text 1). Tonsil swabs were obtained from 3- to 10-week-old piglets at 13 farms, of which 9 had ongoing *S. suis* disease outbreaks. Three sampled farms had no history of *S. suis* disease. We compared our European farms to samples obtained at one US farm with a history of *S. suis* disease but no cases at the time of sampling. Lastly, we sampled 5 piglets free-living in a forest in the Netherlands at approximately 1 month after separation from the sows.

We selected 45 pairs of case-control piglets for metagenomic sequencing (table 1). The pairs were selected to be as equal as possible, coming from the same pen, room, and/or sow, and having received equivalent antimicrobial treatment. The cross-sectional case-control study design was extended with longitudinal sampling at two farms, ES1 and ES2, to determine whether microbiota differences in early life may predict future *S. suis* disease occurrence.

ES1 is a production farm free of *S. suis* disease despite all sows and their carried microbiota

and *S. suis* strains originating from ES2, which had severe *S. suis* disease problems.

Clinical signs consistent with *S. suis* infection were recorded at each farm visit. Cases fell

into two categories: arthritis (typically presenting as lameness) and meningitis (including

otitis and sepsis, typically presenting as loss of balance, paralysis, paddling, shaking, and

convulsing). Confirmation of *S. suis* infection was not carried out on the individual piglets

due to welfare reasons. Most sampled piglets recovered after antimicrobial treatment.

Streptococcus suis disease was confirmed by necropsy in all 3 case piglets on farm NL1.

This study and all animal procedures were approved by the appropriate ethical committees.

Sampling of the forest piglets was conducted according to the restrictions of the Animal and

Human Welfare Codes in The Netherlands (2019.W-0026.001). On other farms, sampling

was carried out for diagnostic purposes (covered by EU Directive 2010/63/EU).

Table 1: Number of case and control samples collected from each farm. See detailed
information in supplementary text 1.

Farm	Country	Age	Amplicon		Metagenomics	
			Case	Control	Case	Control
DE1	Germany	Postweaning	2	6	2	2
DE2	Germany	Postweaning		6		4
DE3	Germany	Postweaning	2	5	2	2
DE4	Germany	Postweaning	3	4	3	3
DE5	Germany	Postweaning	2	4	2	2
DE6	Germany	Postweaning	3	4	2	2
ES1	Spain	Postweaning		15		3
ES1	Spain	Preweaning		15		2
ES2	Spain	Postweaning	11	16	11	11
ES2	Spain	Preweaning	8 prospective	16		
ES3	Spain	Preweaning	10	22	10	10
ES4	Spain	Preweaning	10	10	10	10
FST	Forest	Postweaning		5		5
NL1	Netherlands	Postweaning	3	9	3	3
US1	USA	Postweaning		30		5

Whole genome sequencing of bacterial isolates

We selected 9 clinical (isolated from lesions observed at necropsy on the farms, but not

necessarily from the same piglets sampled for tonsillar microbiota) and 7 non-clinical (from

tonsillar swabs) *S. suis* strains for whole-genome sequencing. The isolates were grown in

Todd-Hewitt broth with yeast extract overnight, and DNA was isolated with PowerSoil DNA

Isolation Kit (Qiagen) with 0.1mm silica bead beating. Isolated DNA was paired-end

Illumina sequenced. Reads were trimmed with trimmomatic v0.39 [52], assembled with

Spades v3.14.1 [53], and annotated with Prokka v1.14.5 [54]. Strain metadata and assembly

statistics is shown in Table S5.

Sample collection

The palatine tonsil microbiota of piglets was sampled by gently scraping the tonsillar surface

with HydraFlock swabs (Puritan, ME, USA) for 10 seconds. Swabs were immediately put in

vials containing Powerbead solution (Qiagen, The Netherlands) and transported at -20 □C

before being stored at -80 □C. DNA was isolated using the PowerSoil DNA Isolation Kit

with 0.7mm garnet bead beating according to the manufacturer's recommended protocol.

Amplicon sequencing

The V3-V4 region of the 16S rRNA gene was amplified with primers 341F (5′-

CCTAYGGGRBGCASCAG-3′) and 806R (5′-GGACTACNNGGGTATCTAAT-3′) and

paired-end 250 bp sequenced using either Illumina HiSeq 2500 or Novaseq 6000. Reads were

trimmed with cutadapt 2.3 [55] using default settings before being processed in DADA2 [24]

following the v1.4 workflow for paired-end big data. Different sequencing batches were run

separately before being merged to account for differences in the learnErrors step. Taxonomy

was assigned with SILVA database v138 [56]. Genus level taxonomy was assigned by the

DADA2 pipeline using the RDP Naive Bayesian Classifier algorithm, and we used mmseqs2

easy-search with default settings to detect the species in SILVA with the highest identity

alignment to each ASV [57]. The highest identity species alignment above 98.5% was

assigned. When several species had equally high identity all were assigned separated by a

slash. Amplicon sequence variants (ASVs) with taxonomic assignment as eukaryote,

mitochondria, or chloroplast were discarded. Alpha and beta diversity were calculated on

rarefied data (24325 reads) using R packages Phyloseq [58] and vegan [59], and the

vegan::adonis function was used to perform PERMANOVA to determine the overall

compositional differences between groups. Vegan function RDA was used for principal

component analysis (PCA) and redundancy analysis (RDA).

Shotgun sequencing

Metagenomic libraries were prepared with NEB Next Ultra DNA Library Prep Kit (New

England Biolabs, ME, USA) following the manufacturer's instructions. DNA was fragmented

to 350 bp, purified with AMPure XP (Beckman Coulter, CA, USA) and sequenced with 150

bp paired-end sequencing on an Illumina NovaSeq 6000 machine.

Analysis of metagenomic data

Pig and plant (feed) reads were removed with kneaddata

(https://github.com/biobakery/kneaddata) using the genomes of pig (GCF_000003025.6),

wheat (GCF_002162155.1), and maize (GCF_902167145.1). Since some samples still had a

large proportion on host or plant reads left after this, we further normalized read counts by

the proportion of plant and pig reads found by kraken analysis [60]. This prevented samples

with small proportions of bacterial reads from being outliers when mapping metagenomic

reads to marker genes and metagenome assembled genomes (MAGs). We created MAGs

using MetaWRAP v1.3.2 [61] with SPAdes v3.14.1 [53], and identified MAG taxonomy with

GTDBtk v1.3.0 [62] and ARGs with resfinder software v4.1 [26]. We also used the resfinder

database for quantifying ARG abundance directly from metagenomics reads. Before mapping

we clustered the genes in the database to 90% identity using mmseqs2 [57] easy-cluster with

settings "--min-seq-id 0.9 --cov-mode 0". Reads were mapped to the representative sequences

of the clusters using mmseqs2 easy-search with default settings, and reads aligning with

minimum 100 bp and at 95% identity were accepted.

To assess the *S. suis* population in the tonsillar microbiota we created a *S. suis* pangenome by

clustering all protein coding genes and mapping metagenomic reads to representative

sequences. We annotated the genomes using Prokka v1.14.5 and clustered all protein coding

genes at 80% residue identity using mmseqs easy-cluster (–min-seq-id 0.8 –cluster-mode 2 –

cov-mode 1). We used previously published metadata covering 1703 assemblies [21]. We

determined the association of each cluster with presence in clinical and non-clinical strains by

calculating the ratio of percent presence in each group. The ratio of clusters more present in

clinical strains were calculated by (% presence clinical/% presence non-clinical) - 1, and

clusters more present in non-clinical strains by ((% presence non-clinical/% presence clinical)

* -1) + 1, so that clusters equally prevalent in clinical and non-clinical strains had a ratio of 0.

We mapped metagenomic reads to the representative sequence of each cluster as described

above and accepted reads mapping at 80% identity and 80% length.

Acknowledgements

We thank the farmers and veterinarians that participated in the study. Most farms are anonymized. Farm NL1 is Schothorst Feed Research B.V. (www.schothorst.nl) and the forest piglets (FST) were collected from Boeren in het Bos (www.boereninhetbos.nl).

Availability of data and materials

All microbiome sequencing data generated by this study is deposited in the NCBI BioProject portal under accession number PRJNA854341. Genome assemblies are available under accession numbers PRJNA849547 and PRJNA849577.

Authors' contributions

VA, FCF, PB, and JW conceived the study. SF, CNI, IHP, XG, JD, IFO, and MLF performed field sampling and sample processing. SF, FCF and JB performed bioinformatic and statistical analysis. All authors contributed to editing the manuscript and read and approved the final version.

References

1. Gottschalk M, Segura M. The pathogenesis of the meningitis caused by Streptococcus suis: the unresolved questions. Veterinary Microbiology. 2000;76:259–72.

2. Lun Z-R, Wang Q-P, Chen X-G, Li A-X, Zhu X-Q. Streptococcus suis: an emerging zoonotic pathogen. The Lancet Infectious Diseases. 2007;7:201–9.

3. Neila-Ibáñez C, Casal J, Hennig-Pauka I, Stockhofe-Zurwieden N, Gottschalk M, Migura-García L, et al. Stochastic Assessment of the Economic Impact of Streptococcus suis-Associated Disease in German, Dutch and Spanish Swine Farms. Frontiers in Veterinary Science. 2021;8.

4. Vötsch D, Willenborg M, Weldearegay YB, Valentin-Weigand P. Streptococcus suis – The "Two Faces" of a Pathobiont in the Porcine Respiratory Tract. Front Microbiol. 2018;9.

5. Aarestrup FM, Rasmussen SR, Artursson K, Jensen NE. Trends in the resistance to antimicrobial agents of Streptococcus suis isolates from Denmark and Sweden. Veterinary Microbiology. 1998;63:71–80.

6. Hadjirin NF, Miller EL, Murray GGR, Yen PLK, Phuc HD, Wileman TM, et al. Large-scale genomic analysis of antimicrobial resistance in the zoonotic pathogen Streptococcus suis. BMC Biol. 2021;19:191.

7. Huang J, Ma J, Shang K, Hu X, Liang Y, Li D, et al. Evolution and Diversity of the Antimicrobial Resistance Associated Mobilome in Streptococcus suis: A Probable Mobile Genetic Elements Reservoir for Other Streptococci. Front Cell Infect Microbiol. 2016;6.

8. Arends JP, Hartwig N, Rudolphy M, Zanen HC. Carrier rate of Streptococcus suis capsular type 2
in palatine tonsils of slaughtered pigs. Journal of Clinical Microbiology. 1984;20:945–7.

9. Clifton-Hadley F, Alexander T. The carrier site and carrier rate of Streptococcus suis type II in
pigs. Veterinary Record. 1980;107:40–1.

10. Lowe BA, Marsh TL, Isaacs-Cosgrove N, Kirkwood RN, Kiupel M, Mulks MH. Defining the
"core microbiome" of the microbial communities in the tonsils of healthy pigs. BMC Microbiol.
2012;12:20.

11. Murase K, Watanabe T, Arai S, Kim H, Tohya M, Ishida-Kuroki K, et al. Characterization of pig
saliva as the major natural habitat of Streptococcus suis by analyzing oral, fecal, vaginal, and
environmental microbiota. PLOS ONE. 2019;14:e0215983.

12. Kernaghan S, Bujold AR, MacInnes JI. The microbiome of the soft palate of swine. Animal
Health Research Reviews. 2012;13:110–20.

13. Liang Z, Wu H, Bian C, Chen H, Shen Y, Gao X, et al. The antimicrobial systems of
Streptococcus suis promote niche competition in pig tonsils. Virulence. 2022;13:781–93.

14. Pena Cortes LC, LeVeque RM, Funk J, Marsh TL, Mulks MH. Development of the tonsillar
microbiome in pigs from newborn through weaning. BMC Microbiology. 2018;18:35.

15. Pena Cortes LC, LeVeque RM, Funk JA, Marsh TL, Mulks MH. Development of the Tonsil
Microbiome in Pigs and Effects of Stress on the Microbiome. Front Vet Sci. 2018;5.

16. Niazy M, Hill S, Nadeem K, Ricker N, Farzan A. Compositional analysis of the tonsil microbiota
in relationship to Streptococcus suis disease in nursery pigs in Ontario. Animal Microbiome.
2022;4:10.

17. Mathieu-Denoncourt A, Letendre C, Auger J-P, Segura M, Aragon V, Lacouture S, et al. Limited
Interactions between Streptococcus Suis and Haemophilus Parasuis in In Vitro Co-Infection Studies.
Pathogens. 2018;7:7.

18. Obradovic MR, Segura M, Segalés J, Gottschalk M. Review of the speculative role of co-
infections in Streptococcus suis-associated diseases in pigs. Veterinary Research. 2021;52:49.

19. Thanawongnuwech R, Brown GB, Halbur PG, Roth JA, Royer RL, Thacker BJ. Pathogenesis of
Porcine Reproductive and Respiratory Syndrome Virus-induced Increase in Susceptibility to
Streptococcus suis Infection. Vet Pathol. 2000;37:143–52.

20. Segura M, Calzas C, Grenier D, Gottschalk M. Initial steps of the pathogenesis of the infection
caused by Streptococcus suis: fighting against nonspecific defenses. FEBS Lett. 2016;590:3772–99.

21. Roodsant TJ, Van Der Putten BCL, Tamminga SM, Schultsz C, Van Der Ark KCH. Identification
of Streptococcus suis putative zoonotic virulence factors: A systematic review and genomic meta-
analysis. Virulence. 2021;12:2787–97.

22. Weinert LA, Chaudhuri RR, Wang J, Peters SE, Corander J, Jombart T, et al. Genomic signatures
of human and animal disease in the zoonotic pathogen Streptococcus suis. Nat Commun. 2015;6:1–
10.

23. Wileman TM, Weinert LA, Howell KJ, Wang J, Peters SE, Williamson SM, et al. Pathotyping the
Zoonotic Pathogen Streptococcus suis: Novel Genetic Markers To Differentiate Invasive Disease-

Associated Isolates from Non-Disease-Associated Isolates from England and Wales. Journal of
Clinical Microbiology. 2019;57:e01712-18.

24. Callahan BJ, McMurdie PJ, Rosen MJ, Han AW, Johnson AJA, Holmes SP. DADA2: High-
resolution sample inference from Illumina amplicon data. Nature Methods. 2016;13:581–3.

25. Wirbel J, Zych K, Essex M, Karcher N, Kartal E, Salazar G, et al. Microbiome meta-analysis and
cross-disease comparison enabled by the SIAMCAT machine learning toolbox. Genome Biology.
2021;22:93.

26. Bortolaia V, Kaas RS, Ruppe E, Roberts MC, Schwarz S, Cattoir V, et al. ResFinder 4.0 for
predictions of phenotypes from genotypes. J Antimicrob Chemother. 2020.
https://doi.org/10.1093/jac/dkaa345.

27. Opriessnig T, Giménez-Lirola LG, Halbur PG. Polymicrobial respiratory disease in pigs. Animal
Health Research Reviews. 2011;12:133–48.

28. De Witte C, Demeyere K, De Bruyckere S, Taminiau B, Daube G, Ducatelle R, et al.
Characterization of the non-glandular gastric region microbiota in Helicobacter suis-infected versus
non-infected pigs identifies a potential role for Fusobacterium gastrosuis in gastric ulceration.
Veterinary Research. 2019;50:39.

29. Wolff B, Boutin S, Lorenz H-M, Ueffing H, Dalpke A, Wolff D. FRI0698 Prevotella and
alloprevotella species characterize the oral microbiome of early rheumatoid arthritis. Annals of the
Rheumatic Diseases. 2017;76 Suppl 2:754–754.

30. Correa-Fiz F, Santos JMG dos, Illas F, Aragon V. Antimicrobial removal on piglets promotes
health and higher bacterial diversity in the nasal microbiota. Sci Rep. 2019;9:1–9.

31. Hall CW, Mah T-F. Molecular mechanisms of biofilm-based antibiotic resistance and tolerance in
pathogenic bacteria. FEMS Microbiology Reviews. 2017;41:276–301.

32. Ricker N, Trachsel J, Colgan P, Jones J, Choi J, Lee J, et al. Toward Antibiotic Stewardship:
Route of Antibiotic Administration Impacts the Microbiota and Resistance Gene Diversity in Swine
Feces. Frontiers in Veterinary Science. 2020;7:255.

33. Zhang L, Huang Y, Zhou Y, Buckley T, Wang HH. Antibiotic Administration Routes
Significantly Influence the Levels of Antibiotic Resistance in Gut Microbiota. Antimicrobial Agents
and Chemotherapy. 2013. https://doi.org/10.1128/AAC.00670-13.

34. Law K, Lozinski B, Torres I, Davison S, Hilbrands A, Nelson E, et al. Disinfection of Maternal
Environments Is Associated with Piglet Microbiome Composition from Birth to Weaning. mSphere.
2021;6:e00663-21.

35. Holman DB, Gzyl KE, Mou KT, Allen HK. Weaning Age and Its Effect on the Development of
the Swine Gut Microbiome and Resistome. mSystems. 2021;6:e00682-21.

36. Choudhury R, Middelkoop A, Boekhorst J, Gerrits WJJ, Kemp B, Bolhuis JE, et al. Early life
feeding accelerates gut microbiome maturation and suppresses acute post-weaning stress in piglets.
Environmental Microbiology. 2021;23:7201–13.

37. Vo N, Tsai TC, Maxwell C, Carbonero F. Early exposure to agricultural soil accelerates the
maturation of the early-life pig gut microbiota. Anaerobe. 2017;45:31–9.

38. Megahed A, Zeineldin M, Evans K, Maradiaga N, Blair B, Aldridge B, et al. Impacts of
environmental complexity on respiratory and gut microbiome community structure and diversity in
growing pigs. Sci Rep. 2019;9:13773.

39. Munk P, Knudsen BE, Lukjancenko O, Duarte ASR, Van Gompel L, Luiken REC, et al.
Abundance and diversity of the faecal resistome in slaughter pigs and broilers in nine European
countries. Nat Microbiol. 2018;3:898–908.

40. Mencía-Ares O, Cabrera-Rubio R, Cobo-Díaz JF, Álvarez-Ordóñez A, Gómez-García M, Puente
H, et al. Antimicrobial use and production system shape the fecal, environmental, and slurry
resistomes of pig farms. Microbiome. 2020;8:164.

41. Zou G, Zhou J, Xiao R, Zhang L, Cheng Y, Jin H, et al. Effects of Environmental and
Management-Associated Factors on Prevalence and Diversity of Streptococcus suis in Clinically
Healthy Pig Herds in China and the United Kingdom. Applied and Environmental Microbiology.
2018;84:e02590-17.

42. Correa-Fiz F, Blanco-Fuertes M, Navas MJ, Lacasta A, Bishop RP, Githaka N, et al. Comparative
analysis of the fecal microbiota from different species of domesticated and wild suids. Sci Rep.
2019;9:13616.

43. Huang J, Zhang W, Fan R, Liu Z, Huang T, Li J, et al. Composition and functional diversity of
fecal bacterial community of wild boar, commercial pig and domestic native pig as revealed by 16S
rRNA gene sequencing. Arch Microbiol. 2020;202:843–57.

44. Petrelli S, Buglione M, Maselli V, Troiano C, Larson G, Frantz L, et al. Population genomic,
olfactory, dietary, and gut microbiota analyses demonstrate the unique evolutionary trajectory of feral
pigs. Molecular Ecology. 2022;31:220–37.

45. Correa-Fiz F, Fraile L, Aragon V. Piglet nasal microbiota at weaning may influence the
development of Glässer's disease during the rearing period. BMC Genomics. 2016;17.

46. Corsaut L, Misener M, Canning P, Beauchamp G, Gottschalk M, Segura M. Field Study on the
Immunological Response and Protective Effect of a Licensed Autogenous Vaccine to Control
Streptococcus suis Infections in Post-Weaned Piglets. Vaccines. 2020;8:384.

47. Curtis J, Bourne FJ. Half-lives of immunoglobulins IgG, IgA and IgM in the serum of new-born
pigs. Immunology. 1973;24:147–55.

48. Morris CR, Gardner IA, Hietala SK, Carpenter TE, Anderson RJ, Parker KM. Persistence of
passively acquired antibodies to Mycoplasma hyopneumoniae in a swine herd. Preventive Veterinary
Medicine. 1994;21:29–41.

49. Procházka Z, Fránek M, Krejĉí J. Duration of the Persistence of 131I-labelled Colostral and
Serum IgG in the Blood of Newborn Piglets. Zentralblatt für Veterinärmedizin Reihe B. 1979;26:366–
70.

50. Vandeputte J, Too HL, Ng FK, Chen C, Chai KK, Liao GA. Adsorption of colostral antibodies
against classical swine fever, persistence of maternal antibodies, and effect on response to vaccination
in baby pigs. Am J Vet Res. 2001;62:1805–11.

51. Vigre H, Ersbøll AK, Sørensen V. Decay of Acquired Colostral Antibodies to Actinobacillus
pleuropneumoniae in Pigs. Journal of Veterinary Medicine, Series B. 2003;50:430–5.

52. Bolger AM, Lohse M, Usadel B. Trimmomatic: a flexible trimmer for Illumina sequence data.
Bioinformatics. 2014;30:2114–20.

53. Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, et al. SPAdes: A New
Genome Assembly Algorithm and Its Applications to Single-Cell Sequencing. Journal of
Computational Biology. 2012;19:455–77.

54. Seemann T. Prokka: rapid prokaryotic genome annotation. Bioinformatics. 2014;30:2068–9.

55. Martin M. Cutadapt removes adapter sequences from high-throughput sequencing reads.
EMBnet.journal. 2011;17:10–2.

56. Quast C, Pruesse E, Yilmaz P, Gerken J, Schweer T, Yarza P, et al. The SILVA ribosomal RNA
gene database project: improved data processing and web-based tools. Nucleic Acids Res.
2013;41:D590–6.

57. Steinegger M, Söding J. MMseqs2 enables sensitive protein sequence searching for the analysis of
massive data sets. Nat Biotechnol. 2017;35:1026–8.

58. McMurdie PJ, Holmes S. phyloseq: An R Package for Reproducible Interactive Analysis and
Graphics of Microbiome Census Data. PLOS ONE. 2013;8:e61217.

59. Oksanen J, Kindt R, Legendre P, O'Hara B, Stevens MHH, Oksanen MJ, et al. The vegan
package. Community ecology package. 2007;10:631–7.

60. Wood DE, Lu J, Langmead B. Improved metagenomic analysis with Kraken 2. Genome Biol.
2019;20:257.

61. Uritskiy GV, DiRuggiero J, Taylor J. MetaWRAP—a flexible pipeline for genome-resolved
metagenomic data analysis. Microbiome. 2018;6:158.

62. Chaumeil P-A, Mussig AJ, Hugenholtz P, Parks DH. GTDB-Tk: a toolkit to classify genomes
with the Genome Taxonomy Database. Bioinformatics. 2020;36:1925–7.