Genomic evidence of natural selection for ash dieback resistance

Carey L. Metheringham[1,2], William J. Plumb[1,2,3], Jonathan J. Stocks[1,2], Laura J. Kelly[1,2], Miguel Nemesio Gorriz[3], Justin Moat[2], Richard J. A. Buggs[1,2], Richard A. Nichols[1]

[1] School of Biological and Behavioural Sciences, Queen Mary University of London, London, UK.
[2] Royal Botanic Gardens, Kew, Richmond upon Thames, UK.
[3] Forestry Development Department, Teagasc, Dublin, Republic of Ireland

Correspondence: r.buggs@kew.org; r.a.nichols@qmul.ac.uk

Abstract

Evolutionary responses to sudden changes in the environment can, in theory, be particularly rapid if they involve small shifts in allele frequencies at many loci. However, such responses are hard to characterise in wild populations. The invasive alien fungal pathogen *Hymenoscyphus fraxineus* represents a strong novel selective challenge to European ash (*Fraxinus excelsior*). A degree of quantitative resistance to the fungus exists in natural populations. A genome-wide association study on field trials identified many single nucleotide polymorphism (SNP) loci associated with resistance. Here, using whole genome resequencing of an infected natural ash tree population in England, we survey the presence of alleles associated with resistance at 10,000 loci. Of these, 7,985 were variable within the population. We used SNP effect sizes inferred in field trials to calculate the average genetic merit of each tree, as a Genomic Estimated Breeding Value (GEBV). The GEBV of trees established after the start of the epidemic was significantly higher than that of related adults from the pre-epidemic generation. To produce a GEBV shift of this magnitude, would require truncation selection eliminating at least 13% of the juvenile population. The changes in allele frequencies that we observe correspond to their effect sizes. Thus, we document a heritable micro-evolutionary adaptive change involving very many loci, occurring over a single generation. This rapid rate of change could be accelerated by a breeding programme informed by genomic selection.

Main

Responses to selection, due to a sudden change in the environment, can theoretically occur
rapidly via small shifts in allele frequencies at multiple genetic loci[1–6]. It is possible to
demonstrate that the appropriate additive genetic variance exists, by analysing resemblance
among relatives (having established relatedness by recording pedigrees, or by exploiting
genetic markers). In wild populations these patterns can be used to infer a response to
selection, for example, in long term studies of bird and deer populations, although the
anticipated response to selection does not always occur, or continue over multiple generations
[7–10]. Demonstrating which allele frequencies have changed in response to selection is more
challenging. The task is particularly difficult in the case of rapid selection from standing genetic
variation at multiple loci. The effects on allele frequencies at the selected site and adjacent loci
can be weak, compared to the much stronger signals expected when a single rare allele or de-
novo mutation sweeps through the population[11]. Consequently, much more convincing
evidence for the genetic basis of evolutionary responses has been shown empirically for one or
a few newly arisen loci of large effect[12,13] than for responses involving many loci of small effect.

Population-level whole genome re-sequencing can be used in a strategy to detect the potentially
more widespread polygenic responses to selection if two requirements are met: (1) many single
nucleotide polymorphisms (SNP) are identified associated with a trait, and (2) temporal change
in their frequencies under selective pressure for the trait can be shown. Genomic prediction
approaches, developed for agricultural breeding, using genome-wide SNP data to predict
individuals' genetic merit for a quantitative trait of interest, could enable us to meet the first
requirement[14,15]. To meet the second requirement we must know genome-wide allele
frequencies before and after the arrival of a new selective pressure[16] or in different age classes
at a single time point[8,17–21]. Trees are a promising study system for the latter approach due to
their large numbers of progeny and long lifespans which preserve multiple generations in place.
Under strong selective pressures an adaptive response is of practical as well as theoretical
interest, as it could lead to evolutionary rescue, allowing population persistence or even the
reversal of population declines[22].

The possibility of such a rapid response is of particular interest in the case of the ash dieback
epidemic that has swept across Europe in the past three decades, caused by the fungus
Hymenoscyphus fraxineus, an alien invasive from East Asia[23]. Epidemics of novel invasive
pathogens provide particularly stark changes in conditions, expected to drive rapid changes in

allele frequencies if heritable variation relating to resistance exists[24]. Numerous studies based

on planted trials suggest that heritable variation in susceptibility to the fungus exists within

European ash (*Fraxinus excelsior*) populations[25–28]. The intensity of selection on viability may be

stronger in smaller, younger ash trees, since they are observed to die more rapidly from ash

dieback infection than larger older trees[29,30]. Rapid juvenile mortality may occur because the

fungus can more quickly encircle the main stem and because smaller trees are closer to the leaf

litter where *Hymenoscyphus fraxineus* apothecia are produced. Recruitment of the next

generation may also be impacted by reduced reproduction by adult trees damaged by not yet

killed by the fungus[31,32].

We may therefore hypothesise[25] that mortality of susceptible juvenile trees and reduced

reproduction by susceptible adult trees will drive changes in allele frequencies leading to an

increase in disease resistance in the new generation of ash. A previous study of allele

frequencies in ~7 year old ash trees with high and low ash dieback damage in large trials

identified thousands of loci putatively affecting damage level in these trees[28]. These data have

been used to train genomic prediction models that estimate the genomically estimated breeding

values (GEBVs) of individual trees[28]. If natural selection is acting on ash under high disease

pressure, then we would expect to see GEBVs increase in the younger generation of trees

which have been exposed to infection since germination, and shifts in allele frequencies to

correlate with their effect sizes. Here we sought to test this hypothesis in a woodland in SE

England.

Results

Phenotypic and genomic characterisation of an ash population

Our study site, Marden Park wood, is an ancient semi-natural woodland dominated by *F.*

excelsior where the pathogen *H. fraxineus* is thought to have been present since 2012.

Phenotypic assessments of this woodland in 2019, when we collected ash tissue samples for

sequencing, found *H. fraxineus* symptoms on the majority of trees. Damage from the fungus

had further increased by 2021 especially in juvenile trees (Figure 1). This fits with widespread

documentation of the progress of the ash dieback epidemic throughout Europe[33,34].

We generated short read sequence data for 580 individuals (128 pre-epidemic adults and 452

post-epidemic juveniles) and 30 technical replicates at approximately 11X whole genome

coverage and from these data estimated allele frequencies at SNP loci. In a previous study,

based on 38,784 ash trees from diverse provenances growing in ash dieback field trials, we had

sequenced the 623 least-damaged trees and 627 highly damaged trees and identified 10,000

SNP loci associated with these phenotypes[28]. In that study these SNPs were used to train

genomic prediction models to estimate the Genomic Estimated Breeding Values (GEBVs) of

individual trees. The 10,000 effect sizes used to calculate the GEBVs were validated by being

used to successfully predict the resilience of trees in a test population[28]. Of these 10,000 SNPs,

7,985 were variable in the Marden Park dataset and passed allele frequency and quality

thresholds. This smaller number of polymorphic SNPs reflects the lower genetic diversity

present in Marden Park wood than in the planted trials of trees chosen from diverse accessions.

Of the 2,015 SNPs that were not variable in the Marden Park population, 1,055 were fixed for

the allele associated with low ash dieback damage in the planted trials and 960 for the allele

associated with high damage.

We calculated GEBVs for the Marden Park trees from the 7,985 polymorphic SNPs using the

parameters of the genomic prediction model trained on the field trials[28]. A visual assessment of

ash dieback damage showed only a weak relationship with GEBV scores (Extended Data

Figure 3 and 4), possibly because of the different phenotype used in adults (canopy cover) and

the reduced control of age and environment for both juveniles and adults, compared to the field

trials. Rather than using correlations with phenotypes, which may be only indirectly related to

survival, we sought more direct evidence for the action of natural selection by comparing the

GEBV scores of adults and juveniles.

Evidence for allele shifts due to natural selection

We found a shift in mean GEBV between adult trees and juvenile trees in our population. The

mean GEBV of juvenile trees (μ = 0.22, n = 452) was higher than the mean GEBV of adult trees

(μ= 0.15, n = 128) (Figure 2A), as we had predicted. The statistical significance of this increase

in score cannot be evaluated by treating each juvenile tree as an independent observation,

since some adults have left multiple offspring. Instead we compared the GEBV score of each

individual with that predicted from its ancestry. If the surviving juveniles have atypically high

GEBV scores, then on average their scores should be higher than predicted from their ancestry.

To capture this ancestry, we characterised relatedness between juveniles and the adults by

selecting 5,793 loci that were not in close linkage with the 7,985 GEBV sites or with one

another, had no missing data and had minor allele frequencies of over 0.3. We expected these

loci to be near-neutral in their effect on ash dieback susceptibility but able to predict the GEBV

in juveniles on the basis of relatedness alone. By way of illustration, genomic prediction using

these 5,793 loci trained on the juvenile GEBV scores produced a perfect correlation with

juvenile GEBVs (r=1, Extended Data Figure 5). To evaluate the change in GEBV across

generations, the model was trained on the adult trees and used to predict the juvenile scores.

These predictions were significantly lower (p = 0.0004) than the actual juvenile GEBVs

(regression in Figure 2B). Thus, juvenile trees had higher GEBVs than predicted on their

relationship to the adult trees, in the absence of selection. The observed shift is 0.058, slightly

lower than the raw estimate of 0.07 from the difference of adult and parent means.

The observed GEBV shift of 0.058 is 15% of the standard deviation of GEBV distribution in the

juvenile trees. To characterise the intensity of selection required to produce this shift, we can

calculate the level of truncation selection that would produce such a shift in GEBV. If we

assume that the narrow sense heritability of ADB susceptibility is 0.4, as previously estimated

by several field-trial studies[25], it would require selection eliminating the most susceptible 13% of

the juvenile population. This level of selection appears plausible, given the 85% mortality seen

in some mature ash plantations due to ash dieback over a 20 year period[34,33], of which only a

small portion would need to be conditional on the loci underlying our GEBV score to have

produced the response that we have detected. If heritability in the woodland population is lower

than the field-trial estimates, due to greater environmental variability and lower genetic diversity,

the estimate of selection would be correspondingly larger. For heritabilities of 0.2 or 0.1, the

estimates would be 29% or 60% elimination respectively.

If, as we propose, this shift in GEBV is due to selection, then we would predict a pattern in

which the change in allele frequency depends on the effect size and the frequency of the

beneficial allele. We detected this pattern (Figure 3) in a regression of allele frequency change

on effect-size x p(1-p), where p is the minor-allele frequency. The relationship was highly

significant (r = 0.0015, p = 0.00038, F = 12.64 on 1 and 7983 DF). The loci previously

estimated to be of large effect (the largest 25%) also showed a stronger association with our

visual assessments of dieback phenotype than randomly selected loci in both adults and

juveniles (P < 10^{-14}, combined over 100 random selections using Fisher's method).

A second test of our interpretation of the shift in GEBV can be conducted in the case of the 121

juveniles for which we were able to assign at least one parent with confidence (using the

sequoia R package[35] on 1000 SNPs). In these cases we could predict the allele frequencies

expected by random transmission. The observed frequencies were significantly different from

these random expectations (q < 0.05, Wilcoxon rank sum test with Bonferroni correction). The

advantageous alleles, associated with low susceptibility to ash dieback, were over-represented,

the excess frequency being correlated with the effect size (r = 0.0595, p < 10^{-6}). This pattern

again suggests that the offspring carrying these alleles had a selective advantage, making them

more likely to survive to become part of our sample (compared to siblings that did not).

For the 121 juveniles for which we could identify parents, we compared parental phenotypes

and parental GEBVs as means of predicting the health of juvenile offspring. We found no

significant correlation between our health phenotypes of the parent trees and the mean

phenotypic health score of their likely offspring (r = 0.112, p = 0.54, n = 37, Figure 4). In

contrast, we found significant correlation between the GEBV of the parent trees and the health

score of their offspring (r = 0.368 , p = 0.029, n = 37, Figure 4). This provides further evidence

that our GEBVs of adult trees are better assessments of their genetic value for resistance to ash

dieback than our leaf canopy phenotypes.

Discussion

We have documented a heritable micro-evolutionary adaptive change occurring over a single

generation. This selection may have occurred partly by reduced seed or pollen production from

adult trees damaged by ash dieback[31,32] and partly through rapid death of young trees infected

by ash dieback, constituting a missing fraction that did not survive. To have demonstrated such

a shift in heritable variation without genome-sequencing would have involved producing large

numbers of clones or offspring from many adult and juvenile trees in the woodland, and growing

these in uniform conditions for several years under ash dieback inoculum pressure. Our

approach was possible because we could build on the estimates of the effect size at 7,985 loci

obtained from large field trials involving 38,784 individuals raised in standardised conditions.

The direction and magnitude of the observed changes in allele frequency correspond to these

effect size estimates, showing that they are broadly applicable in the semi-natural conditions of

Marden Wood.

Few other studies have thus far sought evidence for micro-evolutionary change via shifts in

GEBVs[14], though several studies in animals have examined shifts in predicted breeding values

based on pedigrees[36]. A recent study showed an increase in GEBVs calculated using genomic

prediction algorithms for adult weight in a 35 year data set for Soay sheep which aligned with

changes in predicted breeding values (from pedigree based studies), though phenotypic body

weight decreased over this time span[15]. In the present study, by contrast, we used the

difference between predicted GEBV (based on relatedness) and observed GEBV (based on loci

associated with the trait of interest) to infer that changes in allele frequency could be attributed

to selection. We were able to take advantage of pre-existing estimates[25] of the genetic effects of

alleles contributing genetic merit of these ash trees, which could extend to a large number of

loci because of the large number of trees in the previous study (38,784) from which the

individuals with extreme phenotypes were selected, and also combined information across

different accessions, each of which had been exposed to the fungus. Similarly Lai *et al.* were

successful in demonstrating adaptation at 24 loci from standing variation in a songbird, by

exploiting adaptation in several independent populations to the selective challenge of living at

high altitude[37]. These strategies could be extended to other organisms in the search for highly

polygenic allelic shifts associated with a trait under selection, to produce neo-Darwinian

evolution.

The action of natural selection in the wild for increased health of European ash under ash

dieback is good news, but whether the rate of change will be sufficient for evolutionary rescue to

occur[22], and whether its end point will be a fully resistant tree, is as yet unknown. The

population we studied is fixed for alleles associated with high susceptibility at more than 900

loci; at these loci gene flow from other populations will be essential to introduce beneficial

alleles. Our parentage analyses suggest ongoing migration of pollen and/or seed into the

population, but this may decrease as ash becomes more scarce in the landscape. As low

susceptibility is highly polygenic, the speed of evolution may be inhibited by linkage among

loci[38,39]. As the epidemic progresses, selective pressure may decrease as the pathogen follows

the host demographics to become more scarce in the landscape. This will diminish the speed of

evolution, especially given that heritability may be low in highly variable woodland environments.

In many British ash woodlands, natural regeneration of ash is scarce due to deer browsing, so

deer management will be essential for natural selection by *H. fraxineus* to be effective. Further

interventions could enhance the rate of evolutionary change, for example, by introducing genetic

diversity from other seed sources or from trees selected, or bred, for high GEBV.

Several breeding programmes for ash dieback resistance are currently underway in Europe,

with parental trees selected based on their ash dieback damage phenotypes. Here, we have

shown that a more effective approach may be to select parental trees based on their GEBV. A

breeding programme for increased disease resistance based on GEBVs would allow more

accurate selection of parents with high breeding values, and would also allow parental pairs with

complementary sets of alleles associated with resistance to be selected for crossing.

Online Methods

Site Selection

For this study we required an accessible ash-dominated woodland site in the south east of

England, where *Hymenoscyphus fraxineus* has been present since 2012. We sought a site with

the following characteristics: containing ash trees of a wide range of ages and sizes, including

the presence of young naturally regenerated saplings; the older ash trees should be (as far as

could be ascertained) naturally seeded rather than planted; the owners should have conducted

few management interventions since 2012, and herbivore populations should be low enough to

allow ongoing natural regeneration. We examined potential sites that fit these criteria, with

Woodland Trust and Kent County Council. In August 2019 we selected a site within Stubbs

Copse in Marden Park Wood in Surrey (51.2707, -0.0400) that has belonged to the Woodland

Trust since 1994. This is on a chalk plateau of the North Downs at approximately 244m above

sea level. It is in the Surrey Hills Area of Outstanding Natural Beauty and is a Site of Special

Scientific Interest. It is an ancient semi-natural woodland site where mature and pole stage ash

is dominant; with beech, whitebeam and oak as rarer components. It approximates to National

Vegetation Classification (NVC) W12a - beech/oak/ash with a dog's mercury sub-community.

When the Woodland Trust surveyed the site in 2019 they found that ash dieback was prevalent

and only a "handful" of trees showed no signs of the disease with others showing "advanced

stages of decline, especially in the younger, pole-staged trees". The 50 years plus management

objectives for the wood are to increase its resilience to pests and diseases and maximise its

biodiversity. The only management interventions planned from 2019-2024 are the removal of

non-native species and the coppicing of some hazel in the understorey. Ash will only be felled

where they pose a risk to site visitors and footpaths.

Sample and Phenotype Collection

From 16-20 September 2019 we surveyed and collected tissue samples from 784 ash trees in Stubbs Copse. We sought to exhaustively sample all ash trees, moving outwards from a central point. For each ash tree we recorded its size, level of ash dieback damage, and GPS coordinates. We labelled each tree with a plastic tag. We took a leaf sample from each tree, and when leaves were unavailable or inaccessible we took buds or twigs. Failing the availability or accessibility of any of these tissues, we took a cambial sample from the trunk using a hammer and chisel, cutting out an area of bark approximately 2cm x 8cm. Sampled tissues were immediately placed into plastic bags containing silica gel, to dry. For trees less than approximately breast height, we measured tree size as height from ground level, and we scored ash dieback damage using a scale similar to that of Pliura[40] with the difference that we had fewer categories of dead trees. Trees we scored as 1 were completely dead, 2 were mostly dead but with small sections of the main stem still living, 3 had infection which had spread into the main stem, 4 had minor infections and 5 were healthy trees with no signs of infection in the stem. For larger trees we measured size as diameter at breast height (DBH) and scored ash dieback damage as an estimate of percentage leaf canopy loss from visual expectation from the ground. On October 16 2019 and 21 and 22 January 2020 we re-collected tissue samples from trees where the first DNA extraction attempt had failed, and as far as possible added aluminium tags to the trees included in the study. On 4 and 12 March 2021 positions of trees were collected using an RTK GPS (Bad Elf Pro), using NTRIP (Networked Transport of RTCM via Internet Protocol) connection to the Imperial University unit (LICCOOGBRO), approximately 27 km from the site. On 18 and 18 August 2021, the trees were again re-scored for ash dieback damage.

Analysis of phenotypic and spatial data

In 2021, 126 of the adult trees and 355 of the juvenile trees were re-phenotyped, and mapped in detail (Extended Data Fig. 1). The phenotypes showed damage due to *H. fraxineus* in juvenile trees increased between 2019 and 2021, with the mean health score decreasing from 3.6 (SE = 0.1, n = 642) in 2019 to 2.5 (SE = 0.1, n = 418) in 2021 (Figure 1); see methods section for scoring system. The percentage of juvenile trees observed to be dead increased from 5.5% in 2019 to 40.9% in 2021. For adult trees, mean percentage canopy coverage was 22% (SE = 2.9, n = 160) in 2019 and 20% (SE = 2.3, n = 126) in 2021 (Figure 1). The lower number of trees re-

301 phenotyped in 2021 was most likely due to loss of labels and accidental loss of juveniles due to

302 trampling or browsing. The 224 juvenile trees that were missing in 2021 had the same average

303 health score (3.6, SE = 0.1) in 2019 as all juvenile trees scored in 2019. Thus it seems unlikely

304 that their disappearance was mainly due to mortality caused by ash dieback. Moran's I (Moran,

305 1950) was calculated in order to estimate the degree of spatial auto-correlation within the

306 phenotypic data of trees which had their positions collected in 2021. The 2019 health scores of

307 the juvenile trees which were mapped in 2021 showed significant spatial auto-correlation

308 (Moran's I = 0.127, p = 0) but this was not the case in 2021 (Moran's I = 0.0135, p = 0.327). The

309 health of the adult trees did not show significant spatial auto-correlation in either year (2019:

310 Moran's I = 0.0546, p = 0.0538; 2021: Moran's I = 0.0452, p = 0.137). Spatial autocorrelation

311 among juvenile trees may be a product of local biotic and abiotic factors, or of relatedness

312 between juveniles in close proximity.

314 **DNA extraction and sequencing**

316 DNA was extracted from dried plant material using Qiagen DNeasy Plant Pro Kits or an adapted

317 CTAB extraction procedure[41]. DNA for 755 trees, including 52 replicates was sent to Novogene

318 for sequencing. Library preparation with NEBNext DNA Library Prep Kit and paired end Illumina

319 sequencing on NovaSeq 6000 was carried out by Novogene (Germany) with a total amount of

320 1.0μg DNA per sample. Genomic DNA was randomly fragmented to a size of 350bp by

321 shearing, then DNA fragments were end polished, A-tailed, and ligated with the NEBNext

322 adapter for Illumina sequencing, and further PCR enriched by P5 and indexed P7 oligos. PCR

323 products were purified using the AMPure XP system and resulting libraries were analysed for

324 size distribution by Agilent 2100 Bioanalyzer, quantified using real-time PCR, then sequenced.

326 **Sequence analysis**

328 We generated 0.71 to 19.92 Gb of read data per tree (μ = 9.93 Gb), giving 0.81 - 22.64 X

329 coverage (μ = 11.28) of the whole genome (assuming a genome size of 877Mbp[42]) for 649

330 individuals and 30 technical replicates. Reads containing adapters, more than 10%

331 undetermined bases or with more than 50% low quality (Qscore<= 5) bases were filtered out

332 with SAMtools (V1.9)[43] and remaining reads were aligned against the BATG0.5 *F. excelsior*

333 genome using the bwa mem function in the Burrows Wheeler Aligner (V0.7.17)[44] with minimum

334 seed length of 19, matching score of 1, mismatch penalty of 4, gap open penalty of 6 and gap

extension penalty of 1. For each sample, 3.0% to 98.9% of reads mapped to the *F. excelsior*

reference genome (μ = 86.3%). Samples with less than 50% of reads aligning to the reference

genome were excluded from further analysis.

SNP calling was performed using HaplotypeCaller, GenomicsDBImport and GenotypeGVCFs in

the Genome Analysis Toolkit (GATK V4.1.4.0)[45]. Sites with greater than 25% missing data were

removed and hard filtering was performed on remaining SNP variants using VariantFiltration in

GATK (V4.2.1.0) with the following parameters; read depth ≥ 10 and <200, variant confidence ≥

20, QD (variant confidence normalised by read depth) ≥ 4, mapping quality ≥ 40, FS ≤ 60 and

minor allele frequency ≥ 0.01. After excluding samples with low alignment and filtering our read

alignments for quality, coverage and indels, we called SNPs at 28,499,977 variable sites in 600

samples, including 25 replicates. Linkage disequilibrium pruning was performed using PLINK

1.9[46] with an independent pairwise r^2 threshold of 0.5, window size of 50 and step size of 5

using the option --indep-pairwise, resulting in a reduced set of 4,287,210 SNPs.

In a PCA of the SNP data performed in PLINK, the first and second principal components

explained 12.1% and 9.2% of the genetic variance respectively (Extended Data Fig. 2). The

identity of replicate samples was confirmed by calculating the genomic distance between

samples using the –distance function in PLINK, with genomic distance recorded as 1 minus

identity-by-state. The 25 technical replicate pairs had genomic distances of between 0.051 and

0.094 (μ = 0.063) (Supplementary Table 1) whereas all other samples (which likely include

some trees with full sibling and parent-offspring relationships) had genomic distances of

between 0.051 and 0.235 (μ = 0.214). The latter included 13 pairs of trees with genomic

distances less than 0.1, which were likely to be close relatives (Supplementary Table 2). The

genomic distance between our technical replicates suggests that we can expect 5-9% of

genotypes to be miscalled, likely due to low sequence coverage. The sequence set with the

highest coverage per replicate pair was used in all subsequent analyses.

Genomic Prediction of Susceptibility

A genetic estimated breeding value (GEBV) for each tree was calculated using effect sizes

obtained from field trial data as described in Stocks (2019)[28]. Briefly, the trials comprised ash

trees from 10 UK native seed zone (NSZ) provenances - including the zone in which Marden

Park is found - together with one from Ireland and one from Germany. The trial trees were of

similar age (~7 years) planted in randomised plots, with even spacing, on former agricultural land. For the genome-wide association study (GWAS) 38,784 trees were scored in the trials for ash dieback damage using a health assessment method very similar to the one we used on the juvenile trees in Marden Park wood. Only 792 fully healthy trees were found in the trials (1.96% of all trees), and of these 623 were used for GWAS and to train the GP model, together with 627 highly damaged trees, giving a total of 1250 trees selected from the extremes of ash dieback damage phenotypes[28]. An rrBLUP model was trained using pool-seq data from these 1250 trees, based on 10,000 loci that had the most significant association with disease susceptibility in a pool-seq GWAS[28]. These 10,000 were able to predict the disease classification of individually sequenced test trees with 68% accuracy[28].

7,985 of these 10,000 "predictive" SNPs were variable within the Marden Park population. Effect sizes from this model, characterised as the additive effect of each copy of the alternate allele , were used to calculate GEBV of the Marden Park trees as $GEBV_j = \Sigma(g_j f_{ij}) + 1$, where GEBV$_j$ is the genomic estimated breeding value of individual j, g_j is the effect size at site i (Supp. Data 2 - from Stocks et al (2019)), and $f_{ij} \epsilon \{0,1,2\}$ is the frequency of the alternate alleles at site i in individual j (Supp. Data 3).

We assessed the relationship between GEBVs and tree health phenotypes by correlation of trees' GEBVs with the health scores of juvenile trees and the percentage canopy cover of the adult trees (Extended Data Fig. 3). The mean difference in health phenotypes of the trees in equal numbers of trees with the highest and lowest GEBVs were compared in increments of one tree and plotted in terms of percentage of trees included (Extended Data Fig. 4), finding a weak trend in the expected direction, which was not formally significant.

Detecting shifts in breeding value between adult and juvenile trees

We wished to investigate whether juvenile trees had higher GEBVs than we would have expected, based on the GEBVs of their adult relatives. All else being equal we would expect closely related individuals to have similar GEBV values. We should therefore be able to predict the GEBV of a tree from a sufficient number of SNPs that are unlinked to the GEBV sites and to each other, but which capture the pattern of relatedness. We therefore selected 5,793 such loci with a minor allele frequency of at least 0.3 (to ensure they were informative). To test whether that selection was sufficient to estimate GEBV on the basis of relatedness alone we fitted an

rrBLUP(V4.6.1)[47] model on the juvenile trees as: $GEBV_J = \mu + X_J\beta_J + \varepsilon$, where GEBV$_j$ is the

vector of breeding values of juvenile trees, μ is the mean, X_J is a matrix of juvenile genotypes ($\in$

{0,1,2}, giving the number of copies of the alternate allele) for the 5,793 SNPs, β_J is a vector of

allelic effects (treated as normally distributed random effects) and the residual variance is

$Var[\varepsilon]$. Effect sizes from this model were then used to calculate rGEBV, the prediction based on

relatedness as: $rGEBV_J = \mu + \Sigma(g_j f_{ij})$, where g_j the effect size at site i, and f$_{ij}$ is the genotype

at site i in individual j. This overfitted model predicted GEBV perfectly (Extended Data Figure 5),

showing that this selection of SNPs are sufficient to predict GEBV, and could be used to make

predictions across generations (where over-fitting will not occur because the juveniles are not

included in the training dataset). To predict GEBV across generations we then trained the same

model on the adults, and then used the X_J of the juveniles to predict their GEBV values.

The level of truncation selection required to produce the observed shift in GEBV was found by

solving the equation Δu = m(s, σ^2) for s, where m is the function giving the mean of the normal

distribution with variance σ^2 with the lowest area, s, removed and Δu is the required shift in

population mean given by the breeders equation: 0.15/h^2. The values of σ^2 was set to 1/h^2 to

correspond to the phenotypic variance unit additive genetic variance and h^2=0.4.

Allelic shifts at causative loci

If allele shifts are due to natural selection we would expect the magnitude of shifts to be related

to the effect size of the allele and the frequency of the allele in the population. The expected

change in allele frequency under selection, from *f* to *f'*, is given by the standard equation *f'* = (*f*^{2}

+ *f(1-f)(1-s))/w,* where s is the selective effect and w the mean fitness.

The value, Δf = *f'-f,* is therefore proportional to *sf*(1-*f*). Under the assumption that the selective

effect was proportional to the estimated effect sizes g_A, we plotted the observed change in

minor allele frequency, Δf, against g_A*f*(1-*f*) in Figure 3. To visualise the trend more clearly we

grouped the loci into 200 bins (quantiles of g_A*f*(1-*f*)). The means of each bin are plotted in with

an area of each point being inversely proportional to the variance in Δf, to convey the relative

precision of the mean Δf estimate. The line is the fitted linear regression (carried out on the

individual Δf values).

To test if the sites identified as having large effects in the field trial[28] contributed to the visually

assessed dieback damage at Marden Park, a new genomic prediction was carried out using the

BGLR (V1.1)[48] package's implementation of the BayesB algorithm[49] (50,000 iterations with a

burn in of 2000). The genotypes comprised the sites in the top 25% (n=1996) of absolute effect

size, estimated from the field trial. These were combined with an equal number of control

genotypes randomly selected from across the genome (that were not part of the GEBV set).

This design was to discern any biological signal from the effect sizes that would be estimated

when fitting a model with a large number of loci to a relatively small number of observations.

The large effect loci had significantly larger estimated effect sizes in this new genomic prediction

model (characterised by their variance) across multiple selections of control loci. (Adults: Pr(>F)

< 1.1e-16, Juveniles: (Pr(>F) < 1.1e-16, combined over 100 random selections using Fisher's

method).

Assignment of parentage

Parentage assignment was performed with the sequoia R package (V2.3.1) [35], using a randomly

selected set of 1,000 SNPs having read depth > 20, minor allele frequency > 0.4, and an

estimated error rate < 0.01. Parentage assignment was run using the hermaphrodite "B" mode.

DBH was used to create a proxy for the birth year of adult trees (birth year = 100 - DBH) and

juvenile trees were assigned a birth year of 100. This allowed for larger, and therefore

presumably older, adult trees to be assigned as parents of younger adult trees, while

disallowing juveniles from being considered as parent trees. The maximum age of parents was

set as 99, allowing all adult trees to be considered as parents of the juveniles. The proxy years

were not intended to be an accurate estimate of tree age. Confidence of parentage assignment

was estimated within sequoia by simulating genotype data based on the estimated pedigree,

recalculating parentage assignment based upon simulated data and comparing the recalculated

pedigree to the original pedigree. The proportion of correct assignments over multiple runs of

the process gives an estimate of confidence in our parentage assignments.

Parentage was assigned to 153 trees, with a confidence value of 0.72. For 125 trees we could

assign a single parent and for 28 trees we could assign two parents. Of trees with assigned

parentage, 121 were classified as juveniles and 16 as adults. We identified 36 adult trees as

likely parents, with a mean of 4.9 offspring per parent tree. Twelve of the trees were only

assigned one likely offspring and 18 trees were assigned at least three offspring

(Supplementary Table 3). The largest set of offspring (n = 29) came from tree 51, which was

both large and centrally located (Extended Data Fig. 1), accounting for 21% of all trees with

assigned parentage. Of these 29 offspring, seven had tree 10 (located in close proximity to tree

51) as the other parent, making them full siblings.

For trees with assigned parents the expected frequency of the minor allele was calculated as;

$\frac{P1 + P2}{2}$, where $P1$ was the frequency of the allele in parent one and $P2$ was the frequency in

parent two. Where only one parent was identified the mean frequency of the minor allele across

the whole of the population was used as $P2$. The 7,985 sites of interest were tested for

significant deviation between expected and observed allele frequencies on a per site basis

using a two sided Wilcoxon rank sum test with continuity correction, with p-values corrected for

multiple testing using the Bonferroni correction. Correlation between effect size and the mean

difference between observed and expected allele frequency in offspring at each site was tested

using Pearson's product-moment correlation.

To test if parental GEBV was a better predictor of offspring health than parental phenotype,

GEBV and percentage canopy cover of parent trees assigned by sequoia were each fitted

against the mean health score of their likely offspring in linear models, weighted according to

the number of likely offspring assigned to each parent tree (Figure 4). The fit of each model was

summarised as the Pearson's correlation coefficient.

Data Availability

Raw reads for all sequenced samples are available on the ENA short read archive under

accession PRJEB44697 (ERP128769). R scripts are available at

https://github.com/CareyMetheringham/MardenPark

References

1. Hermisson, J. & Pennings, P. S. Soft sweeps: molecular population genetics of adaptation

from standing genetic variation. *Genetics* **169**, 2335–2352 (2005).

2. Messer, P. W. & Petrov, D. A. Population genomics of rapid adaptation by soft selective

sweeps. *Trends Ecol. Evol.* **28**, 659–669 (2013).

3. Barton, N. H. & Keightley, P. D. Understanding quantitative genetic variation. *Nat. Rev.*

Genet. **3**, 11–21 (2002).

4.   Höllinger, I., Pennings, P. S. & Hermisson, J. Polygenic adaptation: From sweeps to subtle
frequency shifts. *PLoS Genet.* **15**, e1008035 (2019).

5.   Fisher, R. A. *The genetical theory of natural selection*. (The Clarendon Press, 1930).

6.   Falconer, D. S. *Introduction to Quantitative Genetics*. (Oliver and Boyd, 1962).

7.   Pujol, B. *et al.* The Missing Response to Selection in the Wild. *Trends Ecol. Evol.* **33**, 337–
346 (2018).

8.   Endler, J. A. *Natural Selection in the Wild*. (Princeton University Press, 1986).

9.   Merilä, J., Sheldon, B. C. & Kruuk, L. E. Explaining stasis: microevolutionary studies in
natural populations. *Genetica* **112-113**, 199–222 (2001).

10.   Charmantier, A., Garant, D. & Kruuk, L. E. B. *Quantitative Genetics in the Wild*. (OUP
Oxford, 2014).

11.   Przeworski, M., Coop, G. & Wall, J. D. The signature of positive selection on standing
genetic variation. *Evolution* **59**, 2312–2323 (2005).

12.   Linnen, C. R., Kingsley, E. P., Jensen, J. D. & Hoekstra, H. E. On the origin and spread of
an adaptive allele in deer mice. *Science* **325**, 1095–1098 (2009).

13.   Van't Hof, A. E. *et al.* The industrial melanism mutation in British peppered moths is a
transposable element. *Nature* **534**, 102–105 (2016).

14.   McGaugh, S. E., Lorenz, A. J. & Flagel, L. E. The utility of genomic prediction models in
evolutionary genetics. *Proc. Biol. Sci.* **288**, 20210693 (2021).

15.   Hunter, D. C. *et al.* Using genomic prediction to detect microevolutionary change of a
quantitative trait. *Proc. Biol. Sci.* **289**, 20220330 (2022).

16.   Messer, P. W., Ellner, S. P. & Hairston, N. G., Jr. Can Population Genetics Adapt to Rapid
Evolution? *Trends Genet.* **32**, 408–418 (2016).

17.   Berry, R. J. & Crothers, J. H. Stabilizing selection in the dog-whelk (Nucella lapillus). *J.*
Zool. **155**, 5–17 (2009).

18.   Hiorns, R. W. & Harrison, G. A. Sampling for the detection of natural selection by age group
genetic differences. *Human biology* **42**, 53–64 (1970).

19.   Weldon, W. F. R. A First Study of Natural Selection in Clausilia Laminata (Montagu).
Biometrika **1**, 109–124 (1901).

20.   Berry, R. J., Bonner, W. N. & Peters, J. Natural selection in House mice (Mus musculus)
from South Georgia (South Atlantic Ocean). *J. Zool.* **189**, 385–398 (2009).

21.   Dowdeswell, W. H. Experimental studies on natural selection in the butterfly, Maniola
jurtina. *Heredity* vol. 16 39–52 (1961).

22. Carlson, S. M., Cunningham, C. J. & Westley, P. A. H. Evolutionary rescue in a changing
world. *Trends Ecol. Evol.* **29**, 521–530 (2014).

23. McMullan, M. *et al.* The ash dieback invasion of Europe was founded by two genetically
divergent individuals. *Nat Ecol Evol* (2018) doi:10.1038/s41559-018-0548-9.

24. Novembre, J. & Han, E. Human population structure and the adaptive response to
pathogen-induced selection pressures. *Philos. Trans. R. Soc. Lond. B Biol. Sci.* **367**, 878–
886 (2012).

25. Plumb, W. J. *et al.* The viability of a breeding programme for ash in the British Isles in the
face of ash dieback. *Plants People Planet* **58**, 227 (2019).

26. Lobo, A., McKinney, L. V., Hansen, J. K., Kjær, E. D. & Nielsen, L. R. Genetic variation in
dieback resistance in Fraxinus excelsior confirmed by progeny inoculation assay. *For.*
Pathol. **45**, 379–387 (2015).

27. McKinney, L. V. *et al.* The ash dieback crisis: genetic variation in resistance can prove a
long-term solution. *Plant Pathol.* **63**, 485–499 (2014).

28. Stocks, J. J., Metheringham, C. L., Plumb, W. & Lee, S. J. Genomic basis of European ash
tree resistance to ash dieback fungus. *Nature Ecology & Evolution* **3**, 1686–1696 (2019).

29. Erfmeier, A. *et al.* Ash dieback and its impact in near-natural forest remnants - A plant
community-based inventory. *Front. Plant Sci.* **10**, 658 (2019).

30. Coker, T. L. R. *et al.* Estimating mortality rates of European ash (Fraxinus excelsior) under
the ash dieback (Hymenoscyphus fraxineus) epidemic. *Plants People Planet* **in press**,
(2018).

31. Semizer-Cuming, D., Finkeldey, R., Nielsen, L. R. & Kjær, E. D. Negative correlation
between ash dieback susceptibility and reproductive success: good news for European ash
forests. *Ann. For. Sci.* **76**, 1–9 (2019).

32. Semizer-Cuming, D., Chybicki, I. J., Finkeldey, R. & Kjær, E. D. Gene flow and reproductive
success in ash (Fraxinus excelsior L.) in the face of ash dieback: restoration and
conservation. *Ann. For. Sci.* **78**, 1–15 (2021).

33. George, J.-P., Sanders, T. G. M., Timmermann, V., Potočić, N. & Lang, M. European-wide
forest monitoring substantiate the neccessity for a joint conservation strategy to rescue
European ash species (Fraxinus spp.). *Sci. Rep.* **12**, 4764 (2022).

34. Coker, T. L. R. *et al.* Estimating mortality rates of European ash (Fraxinus excelsior) under
the ash dieback (Hymenoscyphus fraxineus) epidemic. *Plants, People, Planet* **1**, 48–58
(2019).

35. Huisman, J. Pedigree reconstruction from SNP data: parentage assignment, sibship

clustering and beyond. *Mol. Ecol. Resour.* **17**, 1009–1024 (2017).

36. Postma, E. Implications of the difference between true and predicted breeding values for

the study of natural selection and micro-evolution. *J. Evol. Biol.* **19**, 309–320 (2006).

37. Lai, Y.-T. *et al.* Standing genetic variation as the predominant source for adaptation of a

songbird. *Proc. Natl. Acad. Sci. U. S. A.* **116**, 2152–2157 (2019).

38. Hill, W. G. & Robertson, A. The effect of linkage on limits to artificial selection. *Genet. Res.*

8, 269–294 (1966).

39. Comeron, J. M., Williford, A. & Kliman, R. M. The Hill-Robertson effect: evolutionary

consequences of weak selection and linkage in finite populations. *Heredity* **100**, 19–31

(2008).

40. Pliura, A., Lygis, V., Suchockas, V. & Bartkevicius, E. Performance of twenty four European

Fraxinus excelsior populations in three Lithuanian progeny trials with a special emphasis on

resistance to Chalara fraxinea. *Baltic For.* **17**, 17–34 (2011).

41. Wang, N. *et al.* Genome sequence of dwarf birch (Betula nana) and cross-species RAD

markers. *Mol. Ecol.* **22**, 3098–3111 (2013).

42. Sollars, E. S. A. *et al.* Genome sequence and genetic diversity of European ash trees.

Nature **541**, 212–216 (2017).

43. Danecek, P. *et al.* Twelve years of SAMtools and BCFtools. *Gigascience* **10**, (2021).

44. Li, H. Aligning sequence reads, clone sequences and assembly contigs with BWA-MEM.

arXiv [q-bio.GN] (2013).

45. Poplin, R. *et al.* Scaling accurate genetic variant discovery to tens of thousands of samples.

bioRxiv 201178 (2018) doi:10.1101/201178.

46. Chang, C. C. *et al.* Second-generation PLINK: rising to the challenge of larger and richer

datasets. *Gigascience* **4**, 7 (2015).

47. Whittaker, J. C., Thompson, R. & Denham, M. C. Marker-assisted selection using ridge

regression. *Genet. Res.* **75**, 249–252 (2000).

48. Pérez, P. & de los Campos, G. Genome-Wide Regression and Prediction with the BGLR

Statistical Package. *Genetics* **198**, 483–495 (2014).

49. Meuwissen, T. H. E., Hayes, B. J. & Goddard, M. E. Prediction of Total Genetic Value

Using Genome-Wide Dense Marker Maps. *Genetics* **157**, 1819–1829 (2001).

Acknowledgements

Whole genome sequencing of trees in Marden Park wood was funded by a Future Proofing Plant Health grant to R.J.A.B. at RBG Kew from the Department for Environment Food and Rural Affairs (Defra). C.L.M.'s PhD studentship was funded by Defra. Post-doctoral research by C.L.M. was funded by Defra under the Centre for Forest Protection. W.J.P.'s PhD studentship was funded by Defra and Teagasc. We thank the Woodland Trust for access to Marden Park wood, and Tim Hodges and Kate Harvey for their support and assistance.

This research utilised Queen Mary's Apocrita HPC facility, supported by QMUL Research-IT. http://doi.org/10.5281/zenodo.438045

Author contributions

R.J.A.B., R.A.N. and C.L.M. conceived the project and designed the analysis. R.J.A.B. selected the field site. R.A.N. and C.L.M. performed the bioinformatic and statistical analyses. C.L.M, W.J.P., J.J.S., L.J.K., M.N.G., R.A.N. and R.J.A.B. conducted the fieldwork. W.J.P. and C.L.M. extracted DNA. J.M. and C.L.M. mapped the trees. L.J.K. advised on analyses. R.J.A.B., R.A.N., L.J.K.and C.L.M. wrote the paper. R.J.A.B. and R.A.N. supervised the project.

Competing interests

The authors declare no competing interests.

    Display Items

630
**Figure 1. Phenotypic estimates of ash dieback damage as a percentage of trees scored
each year.** A) Health scores (where 5 is the most healthy) of juvenile trees as a percentage of
the total trees scored for each age group in 2019 and 2021 and B) Percentage canopy coverage
estimates of adult trees

638
639 **Figure 2. Shift in GEBV between adult and juvenile trees** (A) Genomic estimates of breeding
640 values for 128 adult trees and 452 juvenile trees calculated from 7,985 SNP loci. (B) Juvenile
641 GEBV plotted against the predicted value. The predictions were based on their ancestry using a
642 rrBLUP model trained on the adult trees. The dashed black line indicates the 1:1 line and blue
643 line the fitted regression.
644
645

Figure 3. Mean change in allele frequency between adults and juveniles

Mean change in allele frequency plotted against the product of effect size and minor allele frequency. The values for 7,985 sites have been combined into 200 points. Each point represents the average of 39 or 40 sites (adjacent values of Effect size x pq, binned), with area representing the precision of the mean change (1/var(Δf)). The red line is the fitted linear regression.

654
655 **Figure 4.** Correlation of A) Canopy cover of likely parent trees and B) GEBV of likely parent
656 trees with the mean phenotypic health score of juvenile offspring as assigned in sequoia.
657 Weighted by the number of offspring trees assigned to each parent.

Extended Data

Extended Data Fig. 1. Map of samples showing location of trees at the Marden Park site.
Collected in March 2021 using an RTK GPS (Bad Elf Pro), using NTRIP (Networked Transport
of RTCM via Internet Protocol) connection to the Imperial University unit (LICCOOGBRO),
approximately 27 km from the site.

675
676 **Extended Data Fig. 2.** Principal genetic components of all adult and juvenile ash trees,
677 including technical replicates, based on 4,287,210 SNPs.
678
679

680
681 **Extended Data Fig. 3**. **Relationship between genomically estimated breeding values**
682 **(GEBV) and phenotypic estimates of ash dieback damage.** (A) GEBV of 128 adult trees
683 plotted against percentage canopy cover as scored in 2019 and (B) GEBV of 452 juvenile trees
684 plotted against 2019 phenotypic scores on a 1-5 scale where 1 is most severely affected.
685 GEBVs were calculated from 7,985 SNP loci.
686

688
Extended Data Fig. 4. **Difference in phenotypic scores of ash dieback damage for trees in**
equally sized upper and lower extremes of the GEBV distribution. (A) Percentage canopy
cover of adult trees (n = 128) and (B) Health scores of juvenile trees (n = 452). GEBVs were
calculated from 7,985 SNP loci. Bars represent the standard error on the cumulative mean
phenotypes and points in black show a significant difference (p <0.05) in mean phenotype
between the trees in the upper and lower extremes of the GEBV distribution.
695

697
Extended Data Fig. 5. Prediction of juvenile GEBV scores using an rrBLUP model trained on 5,793 sites not used in original GEBV calculations produced a perfect correlation (r=1) with the juvenile GEBV scores.

 Supplementary information

 Supplementary Tables

 Supplementary Table 1 | Technical replicate identification. Genomic distance (1 minus
 identity-by-state) between replicate samples

Sample 1	Sample 2	Genomic Distance
S31C	S31R	0.067
S54Q	S54R	0.051
S67C	S67R	0.059
S76C	S76R	0.055
S82C	S82R	0.057
S223C	S223R	0.067
S332C	S332Q	0.053
S410C	S410Q	0.061
S415C	S415R	0.059
S571C	S571Q	0.064
S655Q	S655R	0.065
S868C	S868R	0.058
S885C	S885R	0.076
S895M	S895R	0.062
S901C	S901Q	0.067
S925C	S925Q	0.094
S928C	S928Q	0.090
S939C	S939Q	0.061
S954C	S954R	0.078
S987C	S987Q	0.056
A	S612Q	0.059
B	S873Q	0.057

C	121*	
D	5*	
E	S165C	0.063
F	968*	
G	277*	
H	S439R	0.052
I	999*	
J	S202C	0.54

722 *For half the blind replicates their duplicate did not pass read alignment thresholds and was
723 therefore not included in SNP calling. In these cases the replicate was relabelled to act as a
724 replacement sample.
725
726

727 **Supplementary Table 2 | Potential Close Relatives.** Genomic distance (1 minus identity-by-
728 state) between similar samples not labelled as replicates
729

Sample 1	Sample 2	Genomic Distance
7	17	0.074
16	110	0.059
44	951	0.053
156	160	0.061
207	209	0.069
281	282	0.074
281	283	0.073
282	283	0.065
458	461	0.058
470	560	0.069
472	877	0.063
473	476	0.064
533	549	0.064
903	906	0.064

 Supplementary Table 3 | Parentage assignments. Showing the number and mean (where
 number of offspring > 1) health scores of offspring assigned to each likely parent tree

Parent Tree	DBH of parent (cm)	% canopy cover in parent (2019)	Parental GEBV	Number of likely offspring	Mean score of offspring (2019)
51	80.0	90	0.8720	29	4.3
10*		5	0.0087	20	4.5
889	42.2	5	-0.3587	19	3.9
39	37.0	75	0.1908	15	3.5
71	49.8	5	0.2481	12	4.1
53	48.1	80	0.5806	8	3.7
890	33.0	10	-0.3074	8	3.0
937	45.0	10	-0.3026	7	2.8
853	39.3	30	0.3898	6	3.0
860	22.3	0	0.3718	5	4.0
945	28.3	60	0.3248	5	3.8
850	34.5	40	-0.4849	4	2.3
898	78.4	85	-0.5119	4	2.0
901	55.5	0	-0.2281	4	5.0
872	24.7	5	0.7402	3	2.3
902	64.0	10	0.0427	3	4.0
907	66.0	80	-0.5879	3	4.7
920	34.0	5	-0.0741	3	5.0
38	37.5	5	-0.3661	2	3
55	50.1	50	0.3625	2	4.5
884	89.8	70	-0.3513	2	3
886	40.0	5	0.0052	2	2
908	4.5	0	0.5565	2	5
910	4.5	20	0.3279	2	3.5

Parent Tree	DBH of parent (cm)	% canopy cover in parent (2019)	Parental GEBV	Number of likely offspring	Mean score of offspring (2019)
51	80.0	90	0.8720	29	4.3
10*		5	0.0087	20	4.5
52	35.0	5	-0.2242	1	NA
712	28.0	NA	0.4506	1	5
852	29.6	10	-0.2466	1	2
896	14.2	1	0.0870	1	2
903	40	0	0.2233	1	NA
927	35.0	1	0.4155	1	5
928	47.0	30	-0.2549	1	2
934	26.0	5	0.1546	1	3
973	6.8	10	0.4924	1	5
978	13.2	0	-0.3005	1	4
988	16.8	0	-0.0603	1	NA
989	4.1	70	0.8080	1	5

* Accurate DBH measurement missing for tree 10

Supplementary Data 1. Phenotypic data for all trees scored or sampled at Marden Park
(attached as excel file)
Marden_Park_Data

Supplementary Data 2. Effect sizes of 7,985 sites used to calculate GEBV
Effect Sizes

Supplementary Data 3. Genotype frequency matrix for 7,985 sites used to calculate GEBV
genotype.matrix.csv