1 **Activation of hTREK-1 by polyunsaturated fatty acids does not only involve**
2 **membrane tension**
3

4 Emilie BECHARD[1], Elodie AREL[1], Jamie BRIDE[1], Julien LOURADOUR[1], Xavier BUSSY[1],
5 Anis ELLOUMI[2], Claire VIGOR[2], Pierre SOULE[4], Camille OGER[2], Jean-Marie
6 GALANO*[2], Thierry DURAND*[2], Jean-Yves LE GUENNEC*[1], Hamid MOHA-OU-
7 MAATI*[3], Marie DEMION*[$,1].
8
[1]: PhyMedExp, Université de Montpellier, Inserm U1046, UMR CNRS 9412, Montpellier
(France)
[2]: IBMM, Université de Montpellier, UMR CNRS 5247, ENSCM, Montpellier (France)
[3]: IGF, Université de Montpellier, Inserm 1191, UMR CNRS 5203, Montpellier (France)
Present adress INM, Inserm U1298, Montpellier (France)
[4] : NanoTemper Technologies GmbH, Munich (Germany)
* : last co-authors
16
[$]corresponding author
Marie DEMION
PhyMedExp, Inserm U1046, CNRS UMR9214
CHU Arnaud de Villeneuve
Bâtiment Craste de Paulet
370 avenue du Doyen Gaston Giraud
34290 MONTPELLIER Cedex 05
France

25
26
27 **Running title** : hTREK-1 is directly activated by PUFAs
28

ABSTRACT

TREK-1 is a mechanosensitive channel also activated by polyunsaturated fatty acids (PUFAs). In this study, we compared the effect of multiple fatty acids and ML402. First, we showed a variable TREK-1 activation by PUFAs related to the variable constitutive activity of TREK-1. Then, we observed no correlation between TREK-1 activation and acyl chain length or number of double bonds suggesting that the bilayer-couple hypothesis cannot explain by itself the activation of TREK-1 by PUFAs. The membrane fluidity measurement is not modified by PUFAs at 10 µM. The spectral shift analysis in TREK-1-enriched microsomes indicates a $K_{D,TREK1}$ at 44 µM of C22:6 n-3. PUFAs display the same activation and reversible kinetics than the direct activator ML402 and activate TREK-1 in both whole-cell and inside-out configurations of patch-clamp suggesting that the binding site of PUFAs is accessible from both sides of the membrane, as for ML402. Finally, we proposed a two steps mechanism for TREK-1 activation by PUFAs: first, insertion into the membrane, without fluidity or curvature modifications, and then interaction with TREK-1 channel to open it.

INTRODUCTION

Potassium channels have a crucial role in the electrical activity of excitable cells such as neurons and cardiomyocytes. Some of these potassium channels are common drug targets modulating the action potential shape and consecutively organs functions. Among the K^+ channels, members of the two-pore domains K^+ channels family (K2P) are involved in the repolarization phase of action potential and in the resting membrane potential (Kelly *et al*, 2006). K2P family includes 15 members classified in 6 functional subfamilies: TWIK (Tandem of pore domains in a Weak Inward rectifying K channel), TREK (TWIK-Related K channel), TASK (TWIK-related Acid Sensitive K channel), TALK (TWIK-related Alkaline pH-activated K channel), THIK (Tandem pore domain Halothane-Inhibited K channel),

TRESK (TWIK-Related Spinal Cord K Channel). K2P channels share common structural

features, each subunit containing two-pore domains (P1, P2) and four putative transmembrane

segments (M1-M4) (Honoré, 2007). The dimerization of K2P channels allows the formation

of the canonical K^+ selective pore domain. TREK-1 channel, one of the three members of the

TREK subfamily with TREK-2 and TRAAK, has been discovered in 1996 by Fink *et al*. The

gating of TREK-1 channel is poly-modulated by a wide range of physical and chemical

stimuli including mechanical stretch, temperature, voltage, pH changes, pharmacological

agents and polyunsaturated fatty acids (PUFAs). This channel is widely studied since its

activation is involved in neuroprotection (Lamas & Fernández-Fernández, 2019),

cardioprotection (Kamatham *et al*, 2019), analgesia (Li & Toyoda, 2015) and reduced

epilepsy crisis (Heurteaux *et al*, 2004). In the majority of these diseases, the protection

afforded by TREK-1 activation is due to the hyperpolarization of the membrane potential

(Djillani *et al*, 2019)

Among the wide diversity of TREK-1 modulators, PUFAs have been shown to behave as

strong activators (Patel *et al*, 1998; Danthi *et al*, 2003). PUFAs are amphipathic molecules

with a hydrophilic carboxyl head and a long hydrophobic chain of carbons and multiple

double bonds. The two main classes of PUFAs are n-3-PUFAs and n-6-PUFAs based on the

position of the first double bond from the carbon ω starting at the methyl extremity.

Numerous studies suggest that n-3-PUFAs, such as docosahexaenoic acid (DHA) and

eicosapentaenoic acid (EPA), exert antiarrhythmic properties through the modulation of ionic

channels and consequent membrane hyperpolarization (Kang & Leaf, 1994, 1996). Such

hyperpolarization could be consecutive to TREK-1 activation as it is largely expressed in the

myocarde(Wiedmann *et al*, 2021; Bechard *et al*, 2022). Since TREK channels are mechano-

sensitives, the comparison of the effects of different PUFAs on TRAAK (Fink *et al*, 1996)

suggests that PUFAs can insert inside the membrane inducing an increase in membrane

fluidity (Leifert *et al*, 1999) that in turn modifies membrane curvature and tension (Sheetz and

Singer., 1974). In fact, TRAAK is very likely to be activated by PUFAs through this

mechanosensitive pathway since the activation is related to the acyl chain length and to the

number of double bonds (Fink et al., 1998; Maingret *et al*, 1999; Patel et al., 2001; Honoré,

2007). However, while it is known that PUFAs also activate TREK-1 channel, no study has

compared the effects of different PUFAs on TREK-1 to determine a potential mechanism of

action. Here, we propose a study that allows a better understanding of which features of

PUFAs are essential for their effects on TREK-1. We performed a thorough comparison of the

effects of 9 PUFAs (from 18 to 22 carbons with 2 to 6 double bonds) and 2 other C18 FA

(mono-unsaturated and saturated) on the TREK-1 current (I_{TREK-1}). We report that there is no

correlation between the acyl chain length and TREK-1 activation, as for the PUFAs-induced

membrane fluidity. Then, the comparison of the effects of PUFAs and ML402, a direct

activator of TREK-1 binding within a cryptic pocket behind the selectivity filter (Lolicato *et*

al, 2017), suggests that there is a at least one binding site for PUFAs on TREK-1 channel.

This hypothesis is reinforced by the affinity protein-PUFA test performed in TREK-1

enriched-microsomes using the nanotemper technology, but the precise binding site remains

to be determined in further studies.

RESULTS

TREK-1 channel activation by PUFAs does not depend of acyl chain length

As previously described,TREK-1 channel is activated by PUFAs, such as arachidonic acid (AA, C20:4 n-6) (Patel *et al*, 1998), LA (Danthi *et al*, 2003) or DHA (Ma & Lewis, 2020). In our hands, the average membrane potential (Em) of HEK hTREK-1 cells was -69.2 ± 0.7 mV (mean ± SEM) (**Fig 1D**, n=130), with a variable initial current density of 15.3 ± 1.2 pA/pF (n=130) at 0 mV. During PUFA superfusion, the current density elicited by voltage ramp from -100 mV to +30 mV progressively increased until a steady-state was reached (**Fig 1A and 1B**). When TREK-1 is activated, Em hyperpolarized to -81.7 ± 0.3 mV (**Fig 1D**, n=130), close to the theorical equilibrium potential of K^+ ions (E_K = -86.5 mV). In order to determine wether the variability of the initial current density (I_0) was related to a variability of TREK-1 channel activity in initial condition, we superfused an inhibitor of TREK-1 channel, Norfluoxetine (NrFlx) at 10 µM. As the initial current was significantly decrease from 8.8 ± 2.2 pA/pF to 3.7 ± 0.9 pA/pF (p-value=0.02; n=5) when Norfluoxetine was applied and as the characteristic outward rectification of I_{TREK-1} was lost, we concluded that I_0 was carried out mostly by TREK-1 channel (**Fig 1C**).

To investigate the importance of the acyl chain length and double bounds, we compared the response of 9 PUFAs with various chain lengths, from 18 to 22 carbons with different number of double bounds, from 2 to 6 unsaturations (**Table 1**). We plotted the relationship between current density at 0 mV in the presence of PUFA (I_{PUFA}) normalized to the initial current density (I_0)(this normalized parameter corresponds to the current fold-increase (I_{PUFA}/I_0)) and the number of carbons on the acyl chain. As shown in **Fig 1E**, there is no correlation between PUFA activation I_{PUFA}/I_0 and the acyl chain length (Spearman correlation test : p-value = 0.19), as for the current density of TREK-1 channel after PUFA perfusion (Spearman correlation test : p-value = 0.10) (**Fig 1G**). We then plotted the relationship between I_{PUFA}/I_0

at 0 mV or current density in response of PUFA perfusion and the number of double bounds

in the acyl chains and also observed no correlation (**Fig 1F and Fig 1H**; Spearman correlation

tests: p-value = 0.63 and p-value = 0.56 respectively). These data reveal that there is no

relationship between the effect of PUFAs on TREK-1 and the acyl chain length or the number

of double bonds. Indeed, C22:6 n-3 tends to have a stronger effect than C22:5 n-3 on TREK-1

current, while they share the same number of carbons (**Fig 1E-H** and **Table 1**). Accordingly,

the most potent activators were both one of the shortest one, C18:2 n-6 (I_{PUFA}/I_0 = 24.8 ± 3.3;

current density = 353.7 ± 22.4 pA/pF), and one of the longest one, C22:6 n-3 (I_{PUFA}/I_0 = 29.8

± 4.4; current density = 276.8 ± 24.5 pA/pF). Conversely, C18:2 n-6 is one of the most potent

activator but C18:3 n-3 failed to activate TREK-1 channel (I_{PUFA}/I_0 = 1.4 ± 0.1; Current

density = 20.7 ± 4.2 pA/pF). Like C18:3 n-3, the saturated stearic acid (C18:0) had no effect

on TREK-1 while the mono-unsaturated C18:1 n-9 produced a 7.4 ± 1.9-fold increase of

I_{TREK-1} (n=12, **Table 1**). However, statistical analysis failed to discriminate the PUFA's

effects (I_{PUFA}/I_0 parameters compared with a nonparametric kruskall-wallis test) probably due

to the important variability of the effects.

The variability in PUFA responses is related to the variable initial current density

Despite an important number of cells studied, we observed a large variability of the TREK-1

current activation by PUFAs illustrated in **Fig 2A**. The severity of the inclusion criteria (see

Material and Methods section) suggests that the variability observed in PUFAs responses is

inherent to TREK-1 channel.

Fig 2A illustrates the large variability of PUFA effects based on the fold-increase analysis

(I_{PUFA}/I_0).We demonstrated that the variability of the I_{PUFA}/I_0 parameter resulted from the

variability of the initial current I_0 but not from the current density at steady-state after the

application of the PUFAs (I_{PUFA}) (**Fig 2B**). Indeed, the calculation of the coefficient of

variation (CV = $\frac{SD}{Mean}$) confirmed that the dispersion of the I_{PUFA}/I_0 calculation is higher than

the dispersion of the I_{PUFA} at steady-state of the activation (**Fig 2C**). Thus, CV modification is

in accordance with the hypothesis that the variability of I_0 is responsible of the variability of

the I_{PUFA}/I_0 parameter.

To better characterize the relationship between I_0 and I_{PUFA}/I_0, we plotted the fold-increase of

TREK-1 current (I_{PUFA}/I_0, Y axis) as a function of the initial current (I_0, X axis). As shown in

Fig 3A, there is a non-linear relationship between I_{PUFA}/I_0 and I_0. This relationship can be

linearized by log-transforming the data (Log10) (**Fig 3B**). Thus, the effects of all PUFAs but

C22:5 n-3 depends on I_0, independently of the absolute amplitude of I_{PUFA}. Therefore, there is

a negative relationship between $Log10(I_{PUFA}/I_0)$ and $Log10(I_0)$: C18:2 n-6 (R^2=0.82, p-

value=0.0001), C20:2 n-6 (R^2=0.66, p-value=0.0271), C20:4 n-3 (R^2=0.80, p-value=0.0001),

C20:4 n-6 (R^2=0.50, p-value=0.0023), C20:5 n-3 (R^2=0.53, 0.0008), C22:5 n-6 (R^2=0.84, p-

value=0.0001) and C22:6 n-3 (R^2=0.68, p-value=0.0001) (**Fig 3B, Table 2**). To determine if

the observed variability is due to the cellular model that we use, or not, we also used two

other models: another stable model of TREK-1 overexpression (Andharia *et al*, 2017) and

transiently transfected HEK 293T cells with TREK-1 (pIRES2 *KCNK2* WT) (**Figure 3C**).

The I_{PUFA}/I_0 variability observed can be explained by the variety of constitutively active

TREK-1 channels at resting condition.

Thanks to the linear regression analysis, we obtained the Y-intercept which reflects the fold-

activation of TREK-1 current for an unitary current (Log(1)=0). The Y-intercept of DPA n-3

cannot be calculated since there was no correlation between I_{PUFA}/I_0 and I_0. For the 7 others

PUFAs, the Y-intercept values allow their separation into 3 groupes with the following

activation sequence : C22:6 n-3, C22:5 n-6, C18:2 n-6 > C20:5 n-3, C20:4 n-6 > C20:4 n-3,

C20:2 n-6 (**Table 2**). Then, plotting the relationship between Y-intercept (Y axis) and the

number of carbons (X axis) reveled once again that there is no correlation between the fold-

activation of TREK-1 and the acyl chain length of PUFAs (**Fig 3D**). There is also no

correlation with the number of double bounds (**Fig 3E**). In conclusion, TREK-1 activation by

PUFAs is definitively not dependent of the acyl chain length and the number of double bond

(**Fig 1D-G** and **Fig 3D-E**).

Variable activation rate suggests different binding affinities of PUFAs for TREK-1

channel

To explore a new way of action of PUFAs on TREK-1 channel, we analyzed and compared

the kinetics of activation of TREK-1 perfusing PUFAs or ML402, a binding activator of

TREK-1. **Fig 4A to C** represent the mean ± SEM of the normalized current densities

$(\frac{I-I_0}{I_{PUFA}-I_0})$ over the time in response to PUFA n-6, PUFA n-3 and ML402, respectively.

Representative traces of current densities activation are presented on the right panels.

Activation kinetics were fitted with a sigmoid equation (see material and method

equation(3)) and the activation rate (min^{-1}) was calculated as the inverse of the slope of the

sigmoid (**Fig 4D**). We were able to distinguish at least two types of activation rate, a fast one

above 3 min^{-1} (C18:2 n-6, C22:6 n-3 and ML402) and a slow one less than 3 min^{-1} (C20:2 n-

6, C20:4 n-3, C20:4 n-6, C20:5 n-3, C22:5 n-6 and C22:5 n-3). As reported in **Table 3**, the

averaged half-activation of TREK-1 channel was smaller for C18:2 n-6 and C22:6 n-3, both

having comparable kinetics to those observed for ML402. Although there were no significant

differences between these three compounds and C22:5 n-6 and C20:5 n-3, the kinetics of the

latters appeared slightly slower (**Fig 4D, Table 3**). In contrast, C20:4 n-3, C20:4 n-6 and

C22:5 n-3 had slower kinetics with an averaged half-activation close to 4 min (**Table 3**).

Since C18:2 n-6 and C22:6 n-3 have the same fast activation rates and C22:5 n-3 has the

slowest one, we assumed that the activation rate of the TREK-1 by PUFAs does not depend

on the acyl chain length. However, among the PUFAs, there is a positive correlation between the activation rate (min^{-1}) and the fold-increase of TREK-1 current (I_{PUFA}/I_0) (Spearman correlation test : p-value = 0.007 and r = 0.88; **Fig 4E**). As the stronger activators are the faster activators of TREK-1, we proposed that some PUFAs, as C18:2 n-6 and C22:6 n-3 have a higher binding affinity for TREK-1 which would allow them to activate it faster and stronger.

Activation of TREK-1 channel by PUFAs is fully reversible.

To see if the activation of TREK-1 channel by PUFA is due to an insertion and thus a modification of membrane tension, we looked at the washout kinetic with extracellular medium free of Bovin Serum Albumin (BSA). We focused on C18:2 n-6, C20:5 n-3 and C22:6 n-3, the most potent activators of C18, C20 and C22 PUFAs, respectively (**Table 1**). ML402 activation reversed immediately and 50% of washout occurred in less than 1 min (**Fig 5A and 5B**). C20:5 n-3, had a kinetic of washing (washout 50%: 0.9 ± 0.1 min) comparable to ML402 (**Fig 5A and 5B**). Even though the washout of C18:2 n-6 and C22:6 n-3 was slower than ML402 (washout 50%, mean ± SEM: 2.4 ± 0.2 min, 3.7 ± 0.3 min and 0.4 ± 0.04 min, respectively) , PUFAs effect were also fully reversed under washing. Once again, there is no correlation between the acyl chain length (**Fig 5C**, Spearman correlation tes: p-value > 0.99) or the number of double bounds (**Fig 5D**, Spearman correlation test: p-value > 0.99) and the time needed to reverse TREK-1 activation. C18:2 n-6 and C22:6 n-3, that activated TREK-1 at least twice more than ML402 (I/I_0: 24.8 ± 3.3, 29.8 ± 4.4 and 9.6 ± 0.9, respectively), had a total reversibility in few minutes. At this point, we cannot exclude a membrane insertion of PUFAs, but we assume that the main effects of PUFAs on TREK-1 activation could be a direct and reversible interaction of PUFAs with the channel, as ML402, or as it is well known for KCNQ1 (Liin *et al*, 2015) and the Shaker H4 Kv channel (Börjesson *et al*, 2008)

Alteration of membrane curvature or fluidity did not explain activation of TREK-1 channel by PUFAs.

In order to evaluate the membrane curvature and tension effects on the PUFAs-induced TREK-1 activation, we performed experiments in the inside-out configuration of the patch-clamp technique (+30 mV, symmetrical condition: 145 mM KCl). In this configuration, we were able to superfuse molecules at the inner face of the membrane and PUFAs must induce a curvature of the membrane opposite to the one obtained in the whole-cell configuration. As shown in **Figure 6A**, ML402 superfusion at the inner face of the membrane induced a reversible increase of TREK-1 current (**Table 4**). Kinetics of activation and washout were comparable to those obtained in the whole-cell configuration (**Fig 4** and **Fig 5**) suggesting that the ML402-binding site is accessible from the outer and the inner leaflet of the membrane. Interestingly, a comparable reversible activation of TREK-1 channel was obtained for C18:2 n-6 and C22:6 n-3 5 μM, suggesting that the membrane curvature is not involved in the activation of TREK-1 channel by PUFAs (**Fig 6B** and **6C, Table 4**).

Then, we assessed the membrane fluidity changes during PUFAs application with a pyrenedecanoic acid probe (PDA), analog to lipids. By measuring the ratio of PDA monomer to excimer fluorescence (405nm/470nm ratio), a quantitative assesment of the membrane fluidity can be obtained at different time points by following the ratio modification over time (F/F_0-1). We focus our experiments on C18:2 n-6 and C22:6 n-3 which are the stronger activators of TREK-1 and C18:3 n-3 that failed to activate TREK-1. These 3 PUFAs at 10 μM did not modify the membrane fluidity even after 50 minutes of application, while at 100 μM they induced a decreased of F/F_0-1 from T_0 and compared to the control condition (basic extracellular medium). These results indicate that membrane fluidity is not modified by PUFAs at 10 μM (**Fig 7A-C**), at least within 50 minutes of application. In addition, given that

TREK-1 activation starts at 1 minute of perfusion of C18:2 n-6 and C22:6 n-3 10 µM (**Fig 4**)

it is unlikely that PUFA effects on TREK-1 activation are due to an increase in membrane

fluidity. Altogether, these data suggest that at least both C18:2 n-6 and C22:6 n-3 PUFAs

activate TREK-1 channel by direct interaction with TREK-1 protein and not by a modification

of the membrane fluidity.

DHA interacts directly with TREK-1 channel protein in TREK-1 enriched microsomes

In order to assess a potential direct PUFA-TREK-1 interaction, we purified microsomes from

hTREK-1/ HEK and native HEK 293T cells and labeled lysine residues of the total proteins to

perform affinity measurements using Spectral Shift (SpS). At first glance, we observed

similar affinity from C22:6 n-3 in these two type of microsomes : $K_{d,TREK-1} \sim 50$ µM and

$K_{d,HEK} \sim 100$ µM highlighting a similar mode of association of C22:6 n-3 within the

microsomes. However, the SpS signal displayed subtle differences according to whether or

not microsomes were enriched in TREK-1 protein. Knowing SpS signal rises from

fluorescence recorded by two individual channel, and having a strong reproducibility from

each condition, we can derive the following assumption:

(1) $I_\lambda(TREK1, HEK) = I_\lambda(TREK1) + I_\lambda(HEK)$

263
Where for each individual wavelength (λ), the fluorescence recorded for the TREK-1-

enriched microscomes correspond to both the fluorescence from the labelled empty

microsomes and from the labelled TREK-1.

Being able to isolate the specific fluorescence associated to TREK-1 ($I_\lambda(TREK1)$) within the

TREK-1-enriched microsomes ($I_\lambda(TREK1, HEK)$), we can use individual channel for further

calculation using **equation 2**, isolating the SpS signal associated only to TREK-1 by

normalizing out the background signal coming for the free microsomes.

(2) $R(TREK1) = \dfrac{I_{670}(TREK1,HEK) - I_{670}(HEK)}{I_{650}(TREK1,HEK) - I_{650}(HEK)}$

When performing so, the specific TREK-1 dose response to C22:6 n-3 gives a $K_d = 44\ \mu M$

(**Fig 8**). Despite displaying similar affinities, the TREK-1-enriched microsomes shows a

statistically better affinity than the microsomes themselves. However, it is worth noticing only

a 2-fold increase of affinity, which may highlight a similar binding mode for both

interactions. Altogether, this suggests an interaction mediated by the lipid bilayer, such as a

membrane insertion followed by interaction with TREK-1 channel.

DISCUSSION

TREK-1 channel activation by PUFAs does not involve mechano-sensitivity but a direct

interaction

In this study, we reported that TREK-1 channel is reversibly activated by polyunsaturated

fatty acids (PUFAs), as already shown in different studies, each being focused mainly on one

PUFA : AA (Patel *et al*, 1998); LA (Danthi *et al*, 2003); DHA (Ma & Lewis, 2020). Our

study is the first to compare the effects of different PUFAs having between 18 to 22 carbon

atoms and 2 to 6 double bonds on TREK-1 channel. We demonstrate that C22:6 n-3 and

C18:2 n-6 are the most potent activators of TREK-1 with an activation as fast as the one of

the direct activator ML402 and a fully reversibility.

TREK subfamily of K2P channels that includes TREK-1, TREK-2 and TRAAK, is

characterized by a mechano-sensitivity and therefore, channels could feel changes in the

membrane curvature induced by PUFAs insertion as for CPZ (Patel et al., 1998). As PUFAs

are anionic amphipath compounds, with a hydrophilic carboxyl group and a lipophilic tail,

they preferentially insert into the outer leaflet of the membrane which is positively charged

(Martinac *et al*, 1990; Sheetz & Singer, 1974). Thus, the longer the lipophilic carbon chain is,

the more the PUFAs will be inserted into the membrane, modifying the local membrane

elastic properties (curvature and fluidity) (Leifert *et al*, 1999). Also, for a given carbon chain

length, the fluidity increases with the number of double bonds. According to the bilayer-

couple hypothesis, PUFA effects on TREK subfamily channels were supposed to be due to a

modification of elastic properties of the membrane leading to an increase of the tension

transmitted to the channels (Patel *et al*, 1998). The study of the TRAAK channel activation by

PUFAs (C18:2 n-6, C20:4 n-6, C20:5 n-3, C22:6 n-3) in the excised patch configuration

shows that TRAAK activation is positively correlated with the carbon chain length of PUFAs

and the number of double bonds (Fink *et al*, 1998; Patel *et al*, 2001). However, we found no

correlation between the acyl chain length, the number of double bonds and the potentiation of

I_{TREK-1}. In the opposite, C18:2 n-6 and C22:6 n-3, respectively the shortest and the longest

PUFA tested, are the most potent activators of TREK-1 channel. It is worth to note that C18:3

n-3, which differs only by one double bond from C18:2 n-6, failed to activate TREK-1

channel. Also, C22:6 n-3 having the same number of carbons than C22:5 n-3 is more than

twice as effective in activating TREK-1. PUFAs having intermediate acyl chain length

produce intermediate activation of TREK-1, in the same range as the direct activator ML402,

independently of the double bonds number. In that respect, the bilayer-couple hypothesis

suggesting an activation of TREK-1 by PUFAs-induced mechanosensitive pathway is not

appropriate. Similar results were obtained in the literature on TREK-2 channel study, C20:4

n-6 being less efficient than C22:6 n-3 and C18:2 n-6 (Lesage *et al*, 2000).

To better characterize the mechanism of action of C22:6 n-3 and C18:2 n-6 on TREK-1, we

compared their kinetics of activation and reversibility with those of ML402. ML402 is a direct

activator of TREK-1, binding within a cryptic pocket behind the selectivity filter that directly

stabilize the C-type gate (Lolicato *et al*, 2017). The activation kinetic of TREK-1 by C22:6 n-

3, C18:2 n-6 and ML402 are comparable and faster than the other PUFAs. This suggests a

possible interaction of C22:6 n-3 and C18:2 n-6 with the channel like the activator ML402

(Lolicato *et al*, 2017) and the inhibitor norfluoxetine (Dong *et al*, 2015). This hypothesis is

reinforced by the inside-out experiments where C22:6 n-3, C18:2 n-6 and ML402 were

applied on the inner leaflet of the membrane. Although the PUFA insertion must induce

opposite curvartures while they insert from the inner (inside-out configuration) or the outer

leaflet (whole-cell configuration), they still activate TREK-1 channel. We finally propose that

at least C22:6 n-3 and C18:2 n-6, like ML402, interact with the channel on an accessible site

from both the inner and the outer leaflet of the cell, and suggesting a binding site accessible

via the lipid bilayer. Studies have already hypothesized that arachidonic acid (C20:4 n-6)

could act directly by interacting with the channel (Maingret *et al*, 1999, 2000) and others have

already shown that free PUFAs can directly interact with ionic channels. Indeed, C22:6 n-3

and C18:3 n-6 interact with K_v7.1 (*KCNKQ1*)(Liin *et al*, 2015; Yazdi *et al*, 2021). PUFAs

also interact with Shaker H4 Kv channel closed to the voltage-sensor domaine through the

negatively charged carboxyl group (Börjesson *et al*, 2010; Börjesson & Elinder, 2011;

Börjesson *et al*, 2008). As TREK-1 lacks a canonical voltage-sensor domain, we can

hypothesize that there is another lipophilic binding site in TREK-1 interacting with the

carboxyl head of PUFAs. This hypothetical PUFA binding site on TREK-1 does not

correspond to the binding site of ML402 since its mutation does not prevent activation by

C20:4 n-6 (Lolicato et al., 2017). Moreover, since the steady-state of TREK-1 activation by

PUFA and ML402 was reached in a range of the minutes, we suggest that the PUFAs binding-

sites have a limited access (Maingret *et al*, 2000). It has to be noted that bovine serum

albumin (BSA) is not required to get a reversal effect during the washout as it is supposed to

be when the effects are due to membrane insertion of the PUFAs (Kang & Leaf, 1994; Leifert

et al, 1999).

Finally, in both whole-cell and inside-out configurations, the washout kinetics of PUFAs were

immediate and the initial current recovered in few minutes. Despite a total reversibility of the

TREK-1 activation under washout, we cannot definitively exclude a membrane insertion of

PUFAs owing to their lipophilic properties. Nevertheless, if PUFAs insert into the membrane,

they do not modify the biophysic properties of the membrane (fluidity, curvature, tension) at

10 µM over a 50 min periode whereas TREK-1 is fully activated by PUFAs in few minutes.

Affinity measurements between PUFAs and TREK-1-enriched microsomes effectively

indicates a binding with the lipid bilayer as observed for the empty microsomes. However,

once the signal due to PUFA interaction with the bilayer is subtracted from the total signal, a

direct binding of PUFAs to TREK-1 is measurable. This specific interaction displays a

stronger affinity ($K_{D,TREK-1} \sim 44$ µM) than the simple signal involving PUFA binding with the

bilayer of the microsomes. Therefore, we propose that two mechanisms act together to

increase I_{TREK-1}: (1)PUFA insertion into the membrane, with no modification of its elastic

properties, is probably recquired to reach (2) a lipophilic binding site on TREK-1 channel

accessible via the lipid bilayer as for the Shaker channel (Börjesson & Elinder, 2011)

Initial TREK-1 variability influence PUFAs response

An unexpected result of this study is the observation that the PUFA effects depend on the

TREK-1 initial current. This variability in initial I_{TREK-1} might have many origins such as

different levels of post-translationnal modifications, different channels recruitment into the

membrane or even the presence of 2 different conductances of TREK-1 channel at the single

channel level (Xian Tao Li *et al*, 2006; Andharia *et al*, 2017). A comparable variability in the

current-fold increase induced by PUFAs was already observed for TREK-1 activation by AA

(Maingret *et al*, 2000), TRAAK (Fink *et al*, 1998; Patel *et al*, 2001) and TREK-2 (Bang *et al*,

2000; Lesage *et al*, 2000) but never explained.

It is important to note that all these studies were performed on COS-7 cells or HEK-293 cells

where mammalian post-translational modifications exist. In the study of Ma and Lewis in

2020, whole-cell recording of TREK-1 and TREK-2 by arachidonic acid were performed in

oocytes (*Xenopus Laevis*) and no such variability in the current fold-increase was observed. It

is known that in this heterologous expression model, post-translational modification are

different (Mantegazza *et al*, 2010). Therefore, the possibility that constitutive TREK-1 current

varies according to the phosphorylation level (or other modifications) and influence the

effects of PUFAs should be taken in consideration. It is known that TREK-1 channel is

modulated by intracellular pathways, related to PKA, PKC and PKG signalizations (Koh *et al*,

2001; Murbartián *et al*, 2005; Honoré, 2007), but to the best of our knowledge, no study

between PUFA activation of TREK-1 and phosphorylation levels has been performed so far.

As a consequence of this variability, to study the effect of an activator of I_{TREK-1}, we need a

sufficient number of cells to apply the Y-intercept of the regression line, probably the best

indicator of the degree of activation of TREK-1. The existence of such variability should be

taken into consideration in pathophysiological studies associated with variability of

expression or response, potentially demultiplying the heterogeneity of TREK-1 response to

activators.

Surprisingly, one of the most efficient activator of TREK-1 channel is C18:2 n-6. As TREK-1

is expressed in cardiomyocytes (Li & Toyoda, 2015; Kelly *et al*, 2006; Decher *et al*, 2017)

C18:2 n-6 should modulate action potential shape and resting membrane potential.This

suggests that C18:2 n-6 could display potent cardio as well as neuroprotective effects. If anti-

arrhythmic properties of omega-3 PUFAs have been thoroughly studied (Kang & Leaf, 1994,

1996; Siscovick *et al*, 2017), the potential anti-arrhythmic effect of omega-6 PUFAs such as

C18:2 n-6 has never been considered. C18:2 n-6 can be found in vegetarian and

Mediterranean diets, both diets having cardioprotective vertues (de Lorgeril *et al*, 1994;

Barnard *et al*, 2019). In both cases the beneficial effects are attributed to the presence of anti-

oxidant molecules and C18:3 n-3 but a possible beneficial effect of C18:2 n-6 has never been

explored.

CONCLUSION

To conclude, there is no relationship between the carbon number of the acyl chain, the

number of double bonds and the activation of TREK-1 channel. Its most potent activators are

C18:2 n-6 (linoleic acid) and C22:6 n-3 (docosahexaenoic acid) and kinetics analysis suggest

a direct interaction of PUFAs on TREK-1 through a lipophilic binding site. This direct

activation of PUFAs in TREK-1 could require the membrane insertion of PUFAs to facilitate

the access to the binding site in the channel.

MATERIEL AND METHODS

Cell culture

We used a HEK-293T cell line for spectral shift mesurement and two HEK/hTREK-1 cell

lines that stably overexpress the human TREK-1 channel subunit (Moha ou Maati *et al*, 2011;

Andharia *et al*, 2017) for electrophysiological experiments and spectral shift mesurement.

Cells were grown in an atmosphere of 95% air/5% CO2 in Dulbecco's modified Eagle's

medium and Glutamax (Invitrogen, Cergy- Pontoise, France) supplemented with 10% (v/v)

heat inactivated fetal bovine serum and 0.5 mg/mL G418 to maintain a selection pressure in

the HEK/hTREK-1 cell line.

The transient expression of TREK-1 was performed in the HEK 293T cell line. The pIRES2

plasmid in which the coding sequence of TREK-1 (pIRES2 *KCNK2* WT) was inserted, was

transfected in HEK 293T cells using the jetPEI® kit (Ozyme) and following the manufacturer

protocol. Briefly, 48h before electrophysiological experiments, cells were transfected with a

mix of the water-soluble polymer jetPEI® with DNA at 6 ng/mL and then seeded in 35x10

mm dishes (Falcon) in presence of 0.5 mg/mL of G418

Electrophysiology

Currents were recorded from hTREK-1/HEK-cells using the patch-clamp technique in whole-

cell recording (WCR) configuration and in inside-out (IO) configuration. Patch pipette having

2.5-4 MΩ resistances (WCR) and 5-7 MΩ resistances (IO) were obtained from borosilicate

glass capillaries by using a two-stage vertical puller (PC- 10, Narishige, London, UK).

Current acquisition was performed with an Axopatch 200B amplifier (Axon Instrument,

Sunnyvale, CA, USA) and low-pass filtered at 5 kHz (WCR) and 2 kHz (IO). Data were

digitalized with a digidata 1550B (Axon Instrument, Sunnyvale, CA, USA) at 10 kHz. The

pClamp10.7 (Axon Instrument) software was used to impose stimulations protocols and

record TREK-1 current. In WCR, cells were kept for experiments if the series resistance were

lower than 8 MΩ, the membrane capacitance between 20 and 35 pF.

The experiments were performed at room temperature (~22 °C). Cells were continuously

superfused with a extracellular medium and modulatory compounds at a rate of 1-1.5 mL/min.

The dishes volume was kept constant at 2.5-3 mL using a vacuum system, connected to a

peristaltic pump (ISMATEC). Using different extracellular potassium concentrations (5 mM

KCl or 15 mM KCl), changes in membrane potential were measured in current-clamp

allowing to determine the time to change completely the medium around the patched-cells (Le

Guennec & Noble, 1994). A complete change of the extracellular medium around the

patched-cell occurred in less than 15 seconds.

Whole-cell recording : the extracellular medium consisted of (in mM): 150 NaCl, 5 KCl, 3

$MgCl_2$, 1 $CaCl_2$ and 10 HEPES pH adjusted to 7.4 with NaOH. The pipette solution consisted

of (in mM): 155 KCl, 3 $MgCl_2$, 5 EGTA and 10 HEPES, pH adjusted to 7.2 with KOH. Cells

were clamped at a holding potential of -80 mV and superfused with the extracellular medium

for over 1 min before the initial current (I_0) recording. Cells were hyperpolarized to -100 mV

for 50 ms and then the macroscopic outward TREK-1 current was elicited with an 800 ms

voltage ramp protocol from -100 mV to +30 mV every ten seconds (see the ramp protocole in

Fig 1A). Thus, the evolution of the current amplitude changes during compound superfusion

was followed in real time (**Fig 1B**). A minimum of 3 min superfusion was applied even when

the tested compound showed no effect and the superfusion condition was switched once a

steady-state was reached. In the case where the tested compound had no effect, a positive

control of the TREK-1 activation was performed with 10 µM docosahexaenoic acid (DHA,

C22:6 n-3).

Data analysis (WCR): Once at steady-state, the current amplitude at 0 mV during the voltage

ramp was measured from the average of the last 3 sweeps over a delta of 1 mV (-0,5 mV to

0,5 mV) to get out of the noise. Current amplitudes are expressed in current densities (pA/pF)

to reduce the variability due to cell size. Activation an washout kinetics were followed at 0

mV. The activation kinetics from each cell of each compound (PUFAs and ML402) was fitted

with a sigmoidal equation (**Fig 5**):

(3) $I_t = I_0 + \dfrac{(I_{PUFA} - I_0)}{1 + e^{\frac{t_{50} - t}{slope}}}$

where I_0 is the initial current and I_{PUFA} is the current caused by PUFA or ML402 at 10 µM

concentration. The activation rate (min^{-1}) was then calculated as : activation rate $= \dfrac{1}{Slope}$.

Inside-out configuration: bath medium contained (in mM) : 140 NaCl, 4.8 KCl, 1.2 $MgCl_2$,

10 glucose, 10 HEPES, pH adjusted to 7.4 with NaOH. After excising the membrane patch,

the bath medium was replaced by the a medium identical to the pipette medium consisted of

(mM): 145 KCl, 1.2 $MgCl_2$, 10 glucose, 10 HEPES, pH adjusted to 7.2 with KOH. Cells were

clamped at a holding potential of 0 mV, the theorical equilibrium potential for K^+ ions in this

condition. Then, the amplitude current was followed at +30 mV in real time during the

perfusion of compounds and the washout.

Membrane fluidity experiments

Membrane fluidity was assessed using the pyrenedecanoic acid probe (PDA), a probe analog

to lipids that incorporates into the cell membrane. The probes in the membrane form

monomers and excimers, with a rate of excimer proportional to the membrane fluidity. Under

PUFAs (C18:2 n-6, C18:3 n-3, C22:6 n-3) application at 10 µM and 100 µM, we measured

the emission spectrum of the PDA: 470 nm for the excimers and 400 nm for the the

monomers. The monitoring of the fluorescent ratio 470/400 nm shift over time with PUFAs

allowed a quantitative monitoring of the membrane fluidity changes due to PUFAs insertion

into the membrane. Briefly, HEK/hTREK-1 cells were grown in culture in glass bottom

culture dishes (MatTek, Ashland, U.S.A). Before fluorescent experiment, adherent

HEK/hTREK-1 cells were incubated 1h at room temperature in the dark with 5 µM of the

fluorescent lipid reagent probe in buffer provided in the membrane fluidity kit (Abcam). After

incubation, the unicorporated probes were removed by washing cells twice with the same

extracellular medium that used for WCR experiments. Epifluorescence microscopic

experiments were performed on a 40X lens using excitation light filter with a 370-mm

dichroic filter (Zeiss). Emitted light were taken at 480±15 nm and 405±10 nm and the

480/405 nm ratio was calculated at intial time before PUFA application (T_0) and T_2 (2 min),

T_4, T_6, T_8, T_{10}, T_{20}, T_{30}, T_{40}, T_{50} and T_{60}. The 470/400 nm ratio was calculated as the average

ratio overs 5 s. Then, the 470/400 nm ratio was normalized as (F/F_0)-1 and represented over

time (**Fig 7**) with F the ratio of fluorescence corresponding to a time T_t and F_0 the ratio

corresponding to the T_0.

Microsomes preparation

Microsomes were obtained from the HEK/hTREK-1 cell line and from the HEK 293T cell

line as control. Briefly, cells were cultured in T75 flasks until confluence.Then, cells were

washed with PBS twice and centrifugated 5 min at 15,000 RPM. The pellet was lysed in a 20

mM PIPES buffer (300 mM Sucrose, 20 mM PIPES, pH 7 with NaOH). Membranes were

then mechanically braked up using insulin syringe. Samples were centrifuged 20 min at

10,000 g at 4°C. Lysis protocol was repeated twice. The pellet was discarded and the

supernatant containing plasma membrane and thus transmembrane proteins was

ultracentifugated 1h at 32,000 RPM at 4°C (Optima^{TM}L-90K Ultracentrifuge, Beckamn

Coulter; Rotor SW60Ti). The pellet containing microsomes was resuspended in 5 mM PIPES

(300 mM Sucrose, 5 mM PIPES, pH 7.4 with NaOH) at a protein concentration adjusted to 25

mg/mL and conserved at -20°C 2 days before the spectral shift measurment.

Spectral shift measurement

Microsomes of HEK 293T cell lin and HEK/hTREK-1 cell line were used to evaluate the

TREK-1-PUFAs interaction with HEK 293T microsomes as control. Briefly, the proteins

contained in the microsomes were labeled on the lysine residues with a fluorescent dye

(MO_L011 RED-NHS 2[nd] generation, NanoTemper Tehnologies GmbH) following the

manufacturer protocol. The 20 µL at 600 µM of dye were used to labelled 200 µL of

microsomes, thus leading to a final incubation volume of 200 + 20 µL. After 20 min of

incubation, the 220 µL of labeled microsomes were applied to a gravity size exclusion B-

column supplied in the kit, following the manufacturer protocol until etution. Labeled

microsomes were eluted with 5 x 200 µL of the 5mM PIPES buffer. 5 fractions of 200 µL

were collected in a clean Eppendorf. The first elution fraction contains the labeled

microsomes while the last fractions contain more free dye. The fluorescence intensity of

fraction 1, 2 and 3 were verified and fractions with a fluorescence count between 400 and

2,000, using 100% excitation power, were used for the experiment after a 10 min

centrifugation at 16,000 g to avoid homogeneity problem. A serial dilution of C22:6 n-3 over

16 points following 1:1 pattern was set up and subsequently mixed with labeled microsomes

and loaded into standard capillaries (NanoTemper Technologies GmbH). The total proteins

concentration in microsomes was kept around 2,5 µg/mL whereas C22:6 n-3 was titrated from

1 mM to 30,5 nM. The read out was performed on a Monolith X instrument from

NanoTemper Technologies GmbH. The samples were subjected to SpS and MST

measurments and only SpS was analyzed due to the normalization requirments. The changes

in the maxima emission wavelength following a ratiometric measurement upon protein-C22:6

n-3 complex formation was used to generate a binding curve as a function of C22:6 n-3

concentrations (Langer *et al*, 2022). The data were pulled from 9 and 5 individual repeat for

the TREK-enriched microsomes and from HEK 293T microsomes respectively. Data

processing and analysis were carried out using the MO.Control 3 software from NanoTemper

Technologies followed by normalization as described in the results section. All the data points

presenting signs of irregularity were automatically discarded upon merging the data, as

documented in the software.

Chemicals

In this study we tested one direct activator of TREK-1, ML402, one saturated fatty acid,

stearic acid, one monounsaturated fatty acid, oleic acid, 9 different PUFAs having between 18

and 22 carbons and 2 to 6 double bonds (**Table 1**). All reagents are summarized in the

reagents and tools table. 10 mM stock solutions were prepared by dissolving PUFAs in

absolute ethanol, ML402 in DMSO. PUFAs and ML402 were stored at -80 °. 10 µM solutions

were obtained by diluting an aliquot of stock solution in the extracellular medium just before

use for WCR experiments. Cells for current recordings were first superfused with

extracellular medium containing ethanol at 1/1000 as a control.

Reagent/resource	Source	Identifier or catalog number
Experimental model		
HEK hTREK-1 cell line	(Moha ou Maati *et al*, 2011)	
HEK 293T/hTREK-1 cell line	(Andharia *et al*, 2017)	
HEK-293 T cell line		
Cell culture		
DMEM + Glutamax	ThermoFisher	31966-021
Fetal bovine serum	Sigma-Aldrich	F7524-500ML
jetPEI®	Ozyme	POL101000053
Trypsin-EDTA	ThermoFisher	25300-054
G418 (geneticin)	Sigma-Aldrich	4727878001
Chemicals		
Linoleic acid	Sigma-Aldrich	L1376-500MG
cis-11,14-Eicosadienoic acid	Sigma-Aldrich	E3127-25MG
Oleic acid	Sigma-Aldrich	O1008
Stearic acid	Sigma-Aldrich	S4751-1G
Cis-5,8,11,14,17-Eicosapentaenoic acid	Sigma-Aldrich	E2011-10MG
Cis-4,7,10,13,16,19-Docosahexaenoic acid	Sigma-Aldrich	D2534-100MG
Alpha-linolenic acid	Sigma-Aldrich	L-039-1ML
Docosapentaenoic acid (cis-7,10,13,16,19)	Sigma-Aldrich	D-120-1ML
Cis-4,7,10,13,16-Docosapentaenoic acid	Sigma-Aldrich	18566-10MG
Arachidonic acid	Sigma-Aldrich	A36-11-100MG
Omega-3 Arachidonic acid	Santa Cruz	sc229695
ML402	MedChemExpress	HY-104027
NanoTemper experiment		

Protein labeling kit	NanoTemper	MO-L011
Capillaries	NanoTemper	MO-K022
Monolith™ X instrument	NanoTemper	
Membrane fluidity experiment		
Membrane fluidity kit	Abcam	ab189819
Glass Bottom culture dishes	MatTek	P35G-0-14-C

Statistical analysis

All set of experiments were performed on at least three different batches (congelation and/or

passage) of cells. All descriptive statistics are displayed from **Table 1** to **Table 4**. Statistical

analysis were performed using Prism software (GraphPad Prism 9, Inc., USA). Spearman

correlation test were performed and the p-value indicates the significance of the correlation

and the r parameter indicates the direction of the correlation (negative or positive). For the

linear regression, the p-value indicates the significance of the relationship, the R^2 indicates the

goodness of the fit an the error bars represent the 90 % confidence interval. Kruskal-Wallis

test followed by a *post hoc* Dunn's test was used for multiple comparison of unmatched data.

A p-value of 0.05 or less was considered as statistically significant. The differences between

more than 2 groups are displayed by the letter code, where groups that do not share the same

letter are significantly different.

AUTHORS CONTRIBUTION

EB carried out the experiments (WCR and IO), set up experiments and protocols, conceived the original idea and wrote the manuscript.
EA carried out the experiments (WCR) and discussed the results.
JB carried out the experiments (WCR) and discussed the results.
JL carried out the experiments(WCR) and discussed the results.
XB carried out the experiments (IO) and discussed the results.
AE participated to the PUFAs structure discussion.
CV helped supervise the project on chemistry of PUFAs and discussed the results.
PS helped in the design and result analyses of affinity tests.
TD helped supervise the project on chemistry of PUFAs and discussed the results.
JMG helped supervise the project on chemistry of PUFAs and discussed the results.
CO helped supervise the project on chemistry of PUFAs and discussed the results.
JYL set up experiments and protocols, conceived the original idea, wrote the manuscript and supervised the project.
HMM provided HEK hTREK-1 cells
MD set up experiments and protocols, conceived the original idea, wrote the manuscript and supervised the project.

ACKNOWLEDGEMENT

This work is supported by the Agence Nationale de la Recherche : acronym TA-BOOm (ANR-20-CE44-0001), Région Occitanie Pyrénées-Méditerranée (MNE-AGPI03), Key Initiative Montpellier Université d'Excellence (KIM Biomarkers and Therapies) and the CBS2 doctoral school. We thank Pr OKADA for the kind gift TR-1 cells: a 293 T cell line that stably expresses TREK-1 channel.

REFERENCES

Andharia N, Joseph A, Hayashi M, Okada M & Matsuda H (2017) Involvement of intracellular transport in TREK-1c current run-up in 293T cells. *Channels Austin Tex* 11: 224–235

Bang H, Kim Y & Kim D (2000) TREK-2, a new member of the mechanosensitive tandem-pore K+ channel family. *J Biol Chem* 275: 17412–17419

Barnard ND, Goldman DM, Loomis JF, Kahleova H, Levin SM, Neabore S & Batts TC (2019) Plant-Based Diets for Cardiovascular Safety and Performance in Endurance Sports. *Nutrients* 11: E130

Bechard E, Bride J, Le Guennec J-Y, Brette F & Demion M (2022) TREK-1 in the heart: Potential physiological and pathophysiological roles. *Front Physiol* 13: 1095102

Börjesson SI & Elinder F (2011) An electrostatic potassium channel opener targeting the final voltage sensor transition. *J Gen Physiol* 137: 563–577

Börjesson SI, Hammarström S & Elinder F (2008) Lipoelectric modification of ion channel voltage gating by polyunsaturated fatty acids. *Biophys J* 95: 2242–2253

Börjesson SI, Parkkari T, Hammarström S & Elinder F (2010) Electrostatic tuning of cellular excitability. *Biophys J* 98: 396–403

Danthi S, Enyeart JA & Enyeart JJ (2003) Modulation of native TREK-1 and Kv1.4 K+ channels by polyunsaturated fatty acids and lysophospholipids. *J Membr Biol* 195: 147–164

Decher N, Ortiz-Bonnin B, Friedrich C, Schewe M, Kiper AK, Rinné S, Seemann G, Peyronnet R, Zumhagen S, Bustos D, *et al* (2017) Sodium permeable and 'hypersensitive' TREK-1 channels cause ventricular tachycardia. *EMBO Mol Med* 9: 403–414

Djillani A, Mazella J, Heurteaux C & Borsotto M (2019) Role of TREK-1 in Health and Disease, Focus on the Central Nervous System. *Front Pharmacol* 10: 379

Dong YY, Pike ACW, Mackenzie A, McClenaghan C, Aryal P, Dong L, Quigley A, Grieben M, Goubin S, Mukhopadhyay S, *et al* (2015) K2P channel gating mechanisms revealed by structures of TREK-2 and a complex with Prozac. *Science* 347: 1256–1259

Fink M, Duprat F, Lesage F, Reyes R, Romey G, Heurteaux C & Lazdunski M (1996) Cloning, functional expression and brain localization of a novel unconventional outward rectifier K+ channel. *EMBO J* 15: 6854–6862

Fink M, Lesage F, Duprat F, Heurteaux C, Reyes R, Fosset M & Lazdunski M (1998) A neuronal two P domain K+ channel stimulated by arachidonic acid and polyunsaturated fatty acids. *EMBO J* 17: 3297–3308

Heurteaux C, Guy N, Laigle C, Blondeau N, Duprat F, Mazzuca M, Lang-Lazdunski L, Widmann C, Zanzouri M, Romey G, *et al* (2004) TREK-1, a K+ channel involved in neuroprotection and general anesthesia. *EMBO J* 23: 2684–2695

Honoré E (2007) The neuronal background K2P channels: focus on TREK1. *Nat Rev Neurosci* 8: 251–261

Kamatham S, Waters CM, Schwingshackl A & Mancarella S (2019) TREK-1 protects the heart against ischemia-reperfusion-induced injury and from adverse remodeling after myocardial infarction. *Pflugers Arch* 471: 1263–1272

Kang JX & Leaf A (1994) Effects of long-chain polyunsaturated fatty acids on the contraction of neonatal rat cardiac myocytes. *Proc Natl Acad Sci U S A* 91: 9886–9890

Kang JX & Leaf A (1996) Evidence that free polyunsaturated fatty acids modify Na+ channels by directly binding to the channel proteins. *Proc Natl Acad Sci U S A* 93: 3542–3546

Kelly D, Mackenzie L, Hunter P, Smaill B & Saint DA (2006) Gene expression of stretch-activated channels and mechanoelectric feedback in the heart. *Clin Exp Pharmacol Physiol* 33: 642–648

Koh SD, Monaghan K, Sergeant GP, Ro S, Walker RL, Sanders KM & Horowitz B (2001)
TREK-1 regulation by nitric oxide and cGMP-dependent protein kinase. An essential role in
smooth muscle inhibitory neurotransmission. *J Biol Chem* 276: 44338–44346
Lamas JA & Fernández-Fernández D (2019) Tandem pore TWIK-related potassium channels
and neuroprotection. *Neural Regen Res* 14: 1293–1308
Langer A, Lüdecke A, Bartoschik T, Cehlar O, Duhr S, Baaske P & Streicher W (2022) A
New Spectral Shift-Based Method to Characterize Molecular Interactions. *ASSAY Drug Dev*
Technol 20: 83–94
Le Guennec JV & Noble D (1994) Effects of rapid changes of external Na+ concentration at
different moments during the action potential in guinea-pig myocytes. *J Physiol* 478 Pt 3:
493–504
Leifert WR, McMurchie EJ & Saint DA (1999) Inhibition of cardiac sodium currents in adult
rat myocytes by n-3 polyunsaturated fatty acids. *J Physiol* 520 Pt 3: 671–679
Lesage F, Terrenoire C, Romey G & Lazdunski M (2000) Human TREK2, a 2P domain
mechano-sensitive K+ channel with multiple regulations by polyunsaturated fatty acids,
lysophospholipids, and Gs, Gi, and Gq protein-coupled receptors. *J Biol Chem* 275: 28398–
28405
Li X-Y & Toyoda H (2015) Role of leak potassium channels in pain signaling. *Brain Res Bull*
119: 73–79
Liin SI, Silverå Ejneby M, Barro-Soria R, Skarsfeldt MA, Larsson JE, Starck Härlin F,
Parkkari T, Bentzen BH, Schmitt N, Larsson HP, *et al* (2015) Polyunsaturated fatty acid
analogs act antiarrhythmically on the cardiac IKs channel. *Proc Natl Acad Sci U S A* 112:
5714–5719
Lolicato M, Arrigoni C, Mori T, Sekioka Y, Bryant C, Clark KA & Minor DL (2017) K2P2.1
(TREK-1)-activator complexes reveal a cryptic selectivity filter binding site. *Nature* 547:
364–368
de Lorgeril M, Renaud S, Mamelle N, Salen P, Martin JL, Monjaud I, Guidollet J, Touboul P
& Delaye J (1994) Mediterranean alpha-linolenic acid-rich diet in secondary prevention of
coronary heart disease. *Lancet Lond Engl* 343: 1454–1459
Ma R & Lewis A (2020) Spadin Selectively Antagonizes Arachidonic Acid Activation of
TREK-1 Channels. *Front Pharmacol* 11: 434
Maingret F, Fosset M, Lesage F, Lazdunski M & Honoré E (1999) TRAAK is a mammalian
neuronal mechano-gated K+ channel. *J Biol Chem* 274: 1381–1387
Maingret F, Patel AJ, Lesage F, Lazdunski M & Honoré E (2000) Lysophospholipids open
the two-pore domain mechano-gated K(+) channels TREK-1 and TRAAK. *J Biol Chem* 275:
10128–10133
Mantegazza M, Rusconi R, Scalmani P, Avanzini G & Franceschetti S (2010) Epileptogenic
ion channel mutations: from bedside to bench and, hopefully, back again. *Epilepsy Res* 92: 1–
29
Martinac B, Zhu H, Kubalski A, Zhou XL, Culbertson M, Bussey H & Kung C (1990) Yeast
K1 killer toxin forms ion channels in sensitive yeast spheroplasts and in artificial liposomes.
Proc Natl Acad Sci U S A 87: 6228–6232
Moha ou Maati H, Peyronnet R, Devader C, Veyssiere J, Labbal F, Gandin C, Mazella J,
Heurteaux C & Borsotto M (2011) A human TREK-1/HEK cell line: a highly efficient
screening tool for drug development in neurological diseases. *PloS One* 6: e25602
Murbartián J, Lei Q, Sando JJ & Bayliss DA (2005) Sequential phosphorylation mediates
receptor- and kinase-induced inhibition of TREK-1 background potassium channels. *J Biol*
Chem 280: 30175–30184
Patel AJ, Honoré E, Maingret F, Lesage F, Fink M, Duprat F & Lazdunski M (1998) A
mammalian two pore domain mechano-gated S-like K+ channel. *EMBO J* 17: 4283–4290

693 Patel AJ, Lazdunski M & Honoré E (2001) Lipid and mechano-gated 2P domain K(+)
694 channels. *Curr Opin Cell Biol* 13: 422–428
695 Sheetz MP & Singer SJ (1974) Biological membranes as bilayer couples. A molecular
696 mechanism of drug-erythrocyte interactions. *Proc Natl Acad Sci U S A* 71: 4457–4461
697 Siscovick DS, Barringer TA, Fretts AM, Wu JHY, Lichtenstein AH, Costello RB, Kris-
698 Etherton PM, Jacobson TA, Engler MB, Alger HM, *et al* (2017) Omega-3 Polyunsaturated
699 Fatty Acid (Fish Oil) Supplementation and the Prevention of Clinical Cardiovascular Disease:
700 A Science Advisory From the American Heart Association. *Circulation* 135: e867–e884
701 Wiedmann F, Rinné S, Donner B, Decher N, Katus HA & Schmidt C (2021)
702 Mechanosensitive TREK-1 two-pore-domain potassium (K2P) channels in the cardiovascular
703 system. *Prog Biophys Mol Biol* 159: 126–135
704 Xian Tao Li null, Dyachenko V, Zuzarte M, Putzke C, Preisig-Müller R, Isenberg G & Daut
705 J (2006) The stretch-activated potassium channel TREK-1 in rat cardiac ventricular muscle.
706 *Cardiovasc Res* 69: 86–97
707 Yazdi S, Nikesjö J, Miranda W, Corradi V, Tieleman DP, Noskov SY, Larsson HP & Liin SI
708 (2021) Identification of PUFA interaction sites on the cardiac potassium channel KCNQ1. *J*
709 *Gen Physiol* 153: e202012850

715 **FIGURE LEGENDS**

716 **Figure 1: Activation of TREK-1 channel by PolyUnsaturated Fatty Acids.**
717 **A.** Illustration of the voltage ramp protocol to record I_{TREK-1} between -100 mV and +30 mV.
718 **B.** A representative traces of current recordings during C22:6 n-3 (DHA) application is shown
719 in one trace *per* 10s (middle). The inset on the right shows the characteristic outward
720 rectifying initial current of TREK-1 (I_0: initial current density). **C.** Effect of 10µM of
721 Norfluoxetine (NrFlx on I_0. **D.** Bar graph showing the initial membrane potential Em_0 (mV)
722 for HEK 293T and HEK hTREK-1 cell lines and the membrane potential Em_{PUFA} of HEK
723 hTREK-1 cells after the PUFA perfusion. **E.** Lack of correlation between the acyl chain
724 length and I/I_0 and **F.** the current density (pA/pF), in response to 10 µM PUFAs at 0 mV. **G.**
725 Lack of correlation between the number of double bounds and I/I_0 and **H.** the current density
726 (pA/pF), in response to 10 µM PUFAs at 0 mV. (Purple: C18:2 n-6, blue: C18:3 n-3, light
727 green : C20:2 n-6, brown: C20:4 n-3, red : C20:4 n-6, green: C20:5 n-3, light green : C22:5 n-
728 3, pink: C22:5 n-6, orange: C22:6 n-3).

729
730 **Figure 2: Variability of TREK-1 activation by PUFAs**
731 **A.** Boxplot of the fold increase (I_{PUFA}/I_0) of TREK-1 and **B.** the current density at steady-state
732 (I_{PUFA}, pA/pF) at 0 mV for each PUFA at 10 µM. Boxplots represent the median (lines) with
733 max and min values (error bars). Groups were compared with a Kruskal-Wallis test followed
734 with the *post-hoc* Dunn's test. Two bars having the same letter are not significantly different.
735 **C.** Changes in the coefficient of variation (SD/mean) between I_{PUFA}/I_0 and I_{PUFA} at 0 n☐ (
736 : coefficient of variation for I_{PUFA}/I_0☐ : coefficient of variation for I_{PUFA}). Both coefficient
737 of variation are superimposed.

738
739 **Figure 3: Variability in TREK-1 activation by PUFAs depends on the initial current I_0.**
740 **A.** Scatter plot representations of I_{PUFA}/I_0 as a function of I_0. **B.** Linearization of the I_{PUFA}/I_0 *vs*
741 I_0 relationship following log-transformation for each PUFA. The simple linear regressions (

) are represented with the 90% confidence interval (IC; dotted line⋯⋯) used to discard one of the C20:4 n-3 and C20:4 n-6 data points (⊠). A point is discarded only if it is out by more than twice the 90% IC. **C.** Relationship between $Log10(I_{C22:6\ n-3}/I_0)$ and $Log10(I_0)$ at 0 mV in different cell lines: HEK hTREK-1 stable cell line #1 (orange), HEK hTREK-1 stable cell line #2 (red) and HEK 293T cells transiently transfected with TREK-1 channel (pIRES2 KCNK2 WT 6 ng/µL)(purple).   **D.** Lack of correlation between the Y-intercept of the linear regressions obtained before and the acyl chain length, **E.** and the number of double bounds, for each PUFA at 10 µM.

Figure 4: Activation kinetics of TREK-1 channel by PUFAs.
A-C. Time course showing the effects of 10 µM PUFAs and ML402 on I_{TREK-1} at 0 mV. PUFAs and ML402 were superfused until the steady-state was reached. *Insets* show the representative current densities recorded in control medium and then under PUFAs or ML402 application. **D.** Bar graphs of the activation rate + SEM to reach the steady-state of I_{PUFA}. Groups were compared with a Kruskal-Wallis test followed with the *post-hoc* Dunn's test. Two bars having the same letter are not significantly different. **E.** Relationship between the fold-increase of TREK-1 (I_{PUFA}/I_0) and the activation rate (min^{-1}) (R^2=0.60; p-value=0.02; error barres show the SEM of the I_{PUFA}/I_0 and the activation rate).

Figure 5: Washout kinetics of LA, EPA, DHA and ML402.
A. Representative traces of the reversibility of the current density with one trace per minute (grey: initial current I_0; colors: PUFA-activated current density at steady-state; black: washout current density). **B.** Normalized current-time curve of the reversibility for each compound (mean ± SEM) with 1 point every 10s. The current was normalized as $(I-I_0)/(I_{PUFA}-I_0)$. The inset shows the time to reach 50% of the reversibility of TREK-1 activation. **C.** Lack of correlation between the time for 50 % of washout and the acyl chain length and **D.** the number of double bounds.

Figure 6: PUFAs activate TREK-1 in inside-out configuration of patch-clamp.
A-C. Representative traces showing the activation and reversibility in inside-out configuration of patch-clamp technique for **A.** ML402 10 µM **B.** C18:2 n-6 5 µM and **C.** C22:6 n-3 5 µM superfused at the inner face of the membrane. Membrane potential was held at +30 mV. The expanded current traces where extracted at the time indicated by the arrows.

Figure 7: Membrane fluidity is not altered by PUFAs at 10 µM
A-C. Membrane fluidity is not changed over time by 10 µM of PUFA but 100 µM :**A.** C18:3 n-3 (10 µM: n=3; 100 µM n=6; CTL n=4), **B.** 18:2 n-6 (10 µM: n=6; 100 µM: n=7; CTL n=6), **C.** C22 :6 n-3 (10 µM: n=5; 100 µM: n=5; CTL n=5),

Figure 8 : **Effect C22:6 n-3 on SpS signal in TREK-1-enriched microsomes and non-enriched microsomes**
(I_λ(TREK1, HEK) represents the binding affinity of C22:6 n-3 (from 1 mM to 30.5 nM) for TREK-1-enriched microsomes from 1 mM to 30.5 nM (n=5-9 experiments).

805 **TABLES**

	IUPAC Name		Tail length	Unsaturation number	ω	I_{PUFA}/I_0	I_{PUFA} (pA/pF)
C18:0	Stearic acid	18:0	18	0		1,0 ± 0,1	18.4 ± 5.5
C18:1 n-9	Oleic acid (9)	18:1 Δ 9	18	1	ω-9	7.4 ± 1.9	113,1 ± 24.6
C18:2 n-6	Linoleic acid (9,12)	18:2 Δ 9,12	18	2	ω-6	24.8 ± 3.3	353.7 ± 22.4
C18:3 n-3	α-Linolenic acid (9,12,15)	18:3 Δ 9,12,15	18	3	ω-3	1.4 ± 0.1	20.72 ± 4.2
C20:2 n-6	Eicosadienoic acid (11,14)	20:2 Δ 11,14	20	2	ω-6	10.1 ± 1.2	166.1 ± 16.2
C20:4 n-3	Eicosatetraenoic acid (8,11,14,17)	20:4 Δ 8,11,14,17	20	4	ω-3	10.9 ± 2.0	119.2 ± 15.5
C20:4 n-6	Eicosatetraenoic acid (5,8,11,14)	20:4 Δ 5,8,11,14	20	4	ω-6	15.7 ± 2.1	199.5 ± 23.1
C20:5 n-3	Eicosapentaenoic acid (5,8,11,14,17)	20:5 Δ 5,8,11,14,17	20	5	ω-3	17.4 ± 2.8	211.7 ± 21.0
C22:5 n-6	Docosapentaenoic acid (4,7,10,13,16)	22:5 Δ 4,7,10,13,16	22	5	ω-6	26.1 ± 8.3	245.3 ± 21.4
C22:5 n-3	Docosapentaenoic acid (7,10,13,16,19)	22:5 Δ 7,10,13,16,19	22	5	ω-3	13.3 ± 2.9	147.1 ± 26.6
C22:6 n-3	Docosahexaenoic acid (4,7,10,13,16,19)	22:6 Δ 4,7,10,13,16,19	22	6	ω-3	29.8 ± 4.4	276.8 ± 24.5

Table 1: Fatty acids nomenclature, characteristics and effects on the TREK-1 current in whole-cell configuration of patch-clamp. Values of I_{PUFA}/I_0 and I_{PUFA} are expressed as mean ± SEM.

	n	p-value	R²	Equation	Y-intercept ± SD
C18:2 n-6	20	0.0001	0.82	Y = -0.9378*X + 2.343	2.3 ± 0.1
C20:2 n-6	7	0.027	0.66	Y = -0.5264*X + 1.621	1.6 ± 0.2
C20:4 n-3	12	0.0001	0.80	Y = -0.6305*X + 1.672	1.7 ± 0.1
C20:4 n-6	16	0.002	0.50	Y = -0.8315*X + 2.083	2.1 ± 0.3
C20:5 n-3	17	0.0008	0.53	Y = -0.7172*X + 1.958	1.9 ± 0.2
C22:5 n-6	11	0.0001	0.84	Y = -1.073*X + 2.462	2.5 ± 0.2
C22:5 n-3	12	0.371	0.08	Y = -0.3221*X + 1.378	1.4 ± 0.4
C22:6 n-3	20	0.0001	0.68	Y = -0.7378*X + 2.133	2.1 ± 0.1

808
809 **Table 2: Parameters of the linear regression $Log10 (I_{PUFA}/I_0)=f(Log10(I_0))$.**
810 TREK-1 current was recorded in whole-celle configuration of patch-clamp.The p-value
811 indicates the significativity of the relationship, the R^2 indicates the goodness of the fit and the
812 Y-intercept reflects the fold-activation of TREK-1 current for an unitary current as Log(1)=0.

	n	Half-activation (min) Mean ± SEM	Steady-state (min) Mean ± SEM	Activation rate (min^{-1}) Mean ± SEM
C18:2 n-6	20	1.9 ± 0.2	3.7 ± 0.3	4.5 ± 0.7
C20:2 n-6	7	4.6 ± 0.3	7.4 ± 0.3	0.9 ± 0.1
C20:4 n-3	12	4.6 ± 0.2	7.1 ± 0.2	0.9 ± 0.1
C20:4 n-6	16	3.9 ± 0.3	6.7 ± 0.3	1.1 ± 0.1
C20:5 n-3	17	3.2 ± 0.3	5.7 ± 0.3	2.0 ± 0.3
C22:5 n-6	11	3.1 ± 0.3	5.3 ± 0.4	1.5 ± 0.2
C22:5 n-3	12	4.2 ± 0.4	5.9 ± 0.4	0.9 ± 0.1
C22:6 n-3	20	2.3 ± 0.2	4.3 ± 0.3	3.4 ± 0.4
ML402	19	2.1 ± 0.2	3.9 ± 0.3	3.8 ± 0.5

831
832 **Table 3: Parameters of the activation kinetic of TREK-1 by PUFAs and ML402 in**
833 **whole-cell configuration of patch-clamp.** Values indicate the time (min) needed to reach the
834 half-activation and the steady-state of the activation of TREK-1. The activation rate (min^{-1})
835 derived from the sigmoidal fit as : $I_t = I_0 + \frac{(I_{PUFA} - I_0)}{1 + e^{\frac{t_{50} - t}{slope}}}$ and activation rate = $\frac{1}{slope}$.

836
837

	n	Concentration (µM)	Steady-state (min) Mean ± SEM	50% of washout (s) Mean ± SEM
C18:2 n-6	4	5	4.6 ± 0.4	18.3 ± 6.3
C22:6 n-3	6	5	2.9 ± 0.3	18.3 ± 4.2
ML402	6	10	2.6 ± 0.6	15.5 ± 7.6

838
839 **Table 4**: **Descriptive statistics for TREK-1 activation and washout kinetics in inside-out**
840 **configuration of patch-clamp.** Values indicate the time needed to reach the steady-state
841 (min) and the time to reach 50% of washout (s).
842
843
844
845