Discovery of unknown quorum-sensing reprogramming factors in *Pseudomonas aeruginosa* using

a dual-acting-gene-blocking approach

Wenjie Cai[1,2], Huiming Liao[1,2], Mingqi Lu[1,2], Xiangting Zhou[1,2], Xiaoyan Cheng[1,2], Weijun Dai[1,2]

[1]Guangdong Laboratory for Lingnan Modern Agriculture, South China Agricultural University,

Guangzhou, 510642, China

[2]College of Plant Protection, Integrative Microbiology Research Center, South China Agricultural

University, Guangzhou, 510642, China

Send correspondence to: Weijun Dai

Integrative Microbiology Research Center, South China of Agricultural University, Guangzhou, 510642,

China

daiweijun@scau.edu.cn

31 **Running title**: identification of QS-reprogramming determinants
32
33
34
Key words: quorum sensing, LasR, *Pseudomonas aeruginosa*

ABSTRACT
LasR-null, QS-active mutants commonly appear in *Pseudomonas aeruginosa* isolates from chronically
infected cystic fibrosis (CF) lung patients. However, underlying QS-reprogramming factors in CF
isolates have yet to be uncovered. Here, we developed an unconventional screening approach, named
dual-acting-gene-blocking (DAGB), to identify unknown QS-reprogramming determinants. Mutations
in *mexT* have been linked to QS-reprogramming in the laboratory PAO1 LasR-null mutant. To block the
inactivation of MexT, we engineered a recombinant LasR-null strain by introducing another copy of
wild-type *mexT* into the neutral site of the genome. This strategy presumably ensures at least one copy
of *mexT* is functioning if another one is inactive during the QS-required evolution, which thus blocks the
inactivating MexT-mediated QS activation. Using DAGB, we identified a point mutation in *rpoA*
(T262->A, RpoA*), encoding the α-subunit of RNA polymerase (RNAP), which could rewire QS
activity in the LasR-null background. Like the MexT mutant, RpoA* also activated QS by suppressing
the expression of the MexEF-OprN efflux pump. In contrast to the MexT mutant, RpoA* caused milder
repression of the MexEF-OprN pump, resulting in an intermediate QS-active phenotype and a slower
decline rate of growth fitness when treated with chloramphenicol. With iterative applications of the
DAGB-based approach, we further discovered other non-*mexT* determinants capable of rewiring QS. Our
study first reports non-MexT QS-reprogramming regulators in the LasR-null mutant, shedding light on
understanding the QS activation mechanism in those *P. aeruginosa* LasR-null, QS-active clinical isolates.
Our findings demonstrate that DAGB is a productive avenue to rapidly identify unknown genetic
determinants in *P. aeruginosa*. Most importantly, the established DAGB-based framework is
generalizable and could potentially be extended to uncover untapped pathways in other research areas as
well as other organisms.

Introduction
Pseudomonas aeruginosa is an opportunistic pathogen that causes severe acute and chronic human
infections in cystic fibrosis (CF) patients with compromised immune systems (Gellatly and Hancock,
2013; Klockgether and Tümmler, 2017). A cohort of *P. aeruginosa* virulence factors is under the control
of the quorum-sensing (QS) system (Lee and Zhang, 2015; Papenfort and Bassler, 2016). QS is a bacterial
cell-cell communication system that regulates the expression levels of hundreds of genes in a cell density-
dependent manner (Lee and Zhang, 2015). *P. aeruginosa* has two acyl-homoserine lactone (AHL) QS
systems, Las and Rhl QS systems. In the Las QS system, LasI catalyzes the diffusible QS signal *N*-3-
oxododecanoyl homoserine lactone (3OC12-HSL), while LasR binds to the 3OC12-HSL forming a
protein-signal complex. This active 3OC12-HSL-LasR complex further activates Las-regulon genes.
Similarly, Rhl QS contains an RhlI-RhlR pair, with RhlI synthesizing the butyryl-HSL (C4-HSL) and
the C4-HSL-bound RhlR activating the Rhl-regulon. These two AHL QS systems further interact with
another Pseudomonas Quinolone Signal (PQS) system, forming a QS hierarchy. In general, Las QS is
atop the QS hierarchy, and disruption of either LasI or LasR diminishes the expression of QS-activated
genes.

P. aeruginosa genome encodes several resistance-nodulation-division (RND) efflux pumps that transport
small molecules into the extracellular environment. MexEF-OprN efflux pumps regulate the QS system
by exporting the signaling molecule PQS (Köhler et al., 2001). Overexpression of MexEF-OprN was
found in the *nfxC*-type mutants, displaying strongly reduced QS-related phenotypes (Ko¨ hler et al., 1997;
Köhler et al., 2001). As a LysR-type transcriptional regulator, MexT is encoded upstream of the *mexEF-*

oprN operon and positively regulates the expression of this operon (Köhler et al., 1999). Overexpression
of MexT substantially decreases QS-controlled products, including C4-HSL, hydrogen cyanide, elastase,
pyocyanin and rhamnolipids (Köhler et al., 2001; Tian et al., 2009b). MexT itself is negatively regulated
by its neighboring gene *mexS* (Richardot et al., 2016; Sobel et al., 2005). In addition to the *mexEF-oprN*
operon, MexT also globally regulates more than 40 genes (Tian et al., 2009a; Tian et al., 2009b).
Mutations in the *mexT* gene were identified in isolates obtained from chronic CF lungs and other
infections (Gilbert et al., 2012; Smith et al., 2006), and they appeared in *P. aeruginosa* laboratory strains
(Cheng et al., 2022; Luong et al., 2014).
88
LasR mutants are commonly identified in CF chronic infections (Feltner et al., 2016; Hoffman et al.,
2009; Smith et al., 2006). Intriguingly, many LasR-null clinical isolates remain QS activity, producing
QS molecules and QS-controlled virulence products, such as C4-HSL and pyocyanin (Bjarnsholt et al.,
2010; Chen et al., 2019; Cruz et al., 2020; Feltner et al., 2016). Inactivation of MexT was found to
reprogram QS activity in the laboratory strain PAO1 LasR-null background (Kostylev et al., 2019; Oshri
et al., 2018) by diminishing the expression of the MexEF-OprN efflux pump that regulates QS activity
(Köhler et al., 2001; Tian et al., 2009b). Likewise, disruption of the MexEF-OprN pump also produces
similar QS-active phenotypes (Kostylev et al., 2019). Since inactivation of MexT produces active Rhl
and PQS QS activities (Kostylev et al., 2019; Oshri et al., 2018), deletion of RhlR abolished the QS
activity. However, the deletion of PqsA and/or PqsE only partially attenuated the QS activity (Kostylev
et al., 2019). Although mutations in *mexT* were shown to be critical for QS-reprogramming in the
laboratory strain, they were rare in clinical isolates (Smith et al., 2006). For example, a LasR-null isolate
from CF chronic infection showed QS-active phenotypes, while containing a functioning MexT (Cruz et
al., 2020). Therefore, in those LasR-null, QS-active clinical isolates, it is not MexT but other unknown
factors that control QS-reprogramming.
104
To identify genetic determinants in those LasR-null, QS-active isolates, we here developed a dual-acting-
gene-blocking (DAGB) approach. We hypothesized that if the inactivation of MexT is not available,
alternative pathways of the bacterium could be selected to operate for QS-reprogramming. Based on this
hypothesis, we generated a recombinant LasR-null mutant bearing two copies of functioning *mexT*. We
assumed that either copy of two *mexT* genes being mutated would still leave another functioning one
during the evolution. Using the DAGB approach, our study identified a RpoA mutant (T262->A, RpoA*)
capable of activating QS in the LasR-null background. RNA-seq analysis revealed a QS-active
transcriptome profile in the LasR-null-RpoA* mutant, showing elevated expression levels in Rhl- and
PQS-related genes. Compared to the MexT mutation, RpoA* had a modest decrease in the expression of
the MexEF-OprN efflux pump, resulting from a weaker binding of RNAP to the efflux pump promoter.
The RpoA*-mediated modulation of the efflux pump led to an intermediate level of QS phenotypes and
a slower decline rate of growth fitness in the presence of chloramphenicol. With further iterative
applications of the DAGB-based approach, our study additionally identified GshA and MexE as QS-
reprogramming factors. Our work first established a powerful framework for uncovering unknown QS-
reprogramming factors in *P. aeruginosa*. Moreover, DAGB developed here could potentially be broadly
applied to reveal unknown fields in a given stress environment as well as in different organisms.
121
Results
***mexT* and *lasR* mutations are not statistically associated in *P. aeruginosa* chronical CF isolates**

LasR mutants are common in *P. aeruginosa* CF isolates from chronic infection patients, which often
contain an active Rhl QS (Bjarnsholt et al., 2010; Feltner et al., 2016). Through an in vitro evolution
approach, *mexT* mutations were found to be responsible for the active Rhl QS in the laboratory PAO1
LasR-null mutant strain (Kostylev et al., 2019; Oshri et al., 2018). However, it is unclear whether this
laboratory-based finding applies to clinical isolates. To address this concern, we re-analyzed previously
published whole genome sequencing (WGS) data of chronical CF infection isolates to investigate a
possible correlation between *lasR* and *mexT* mutations. Among 29 CF patients examined, *lasR* mutations,
including nonsynonymous single nucleotide substitutions and insertions, were identified in isolates from
22 (75.9%) patients. However, only two (9.1%) of those isolates simultaneously had nonsynonymous
mutations in *lasR* and *mexT* (Fisher's exact test, $P = 0.23$, odds ratio = 0.27). Moreover, *mexT* mutations
were additionally found in some isolates that contain an intact *lasR* (Table S1). Recently, Cruz et al.
reported a chronical CF infection isolate (strain E90) (Cruz et al., 2020). This LasR-null isolate E90
exhibits typical Rhl-active phenotypes, while containing a complete, functioning MexT. Combined with
above findings, *mexT* mutations are not statistically colocalized with *lasR* mutations in those chronically
infected CF isolates. Therefore, inactivating MexT might not account for the rewired QS activity at least
in some LasR-null, Rhl-active clinical isolates.
140
Development of a systematic approach for identifying none-*mexT* QS-reprograming factors
To explore uncharacterized QS factors responsible for QS hierarchy reprogramming in the LasR-null
background, we first tried the traditional transposon mutagenesis approach. A *mariner*-based transposon
(pBT20) (Kulasekara et al., 2005) was used to construct the insertion mutant library. This library was
screened on the skim milk plate for the visualization of the QS-controlled proteolytic activity.
Transposon-inserted genes encoding QS negative regulators, such as *mexT*, would be expected to give
rise to QS-active phenotypes. A total of 40, 000 mutant colonies, corresponding to a transposon density
of about 156 bp per insertion in the PAO1 genome, were examined. By this way, we could only obtain
one transposon colony exhibiting a protease-positive phenotype. This colony has a transposon insertion
in the *mexF* gene, which codes for an essential component of the MexEF-OprN efflux pump. Given that
MexT-mediated QS-reprogramming mostly relies on the MexEF-OprN efflux pump (Kostylev et al.,
2019), it is not surprising to see that the disruption of MexF resulted in a QS-active phenotype. No other
QS-reprogramming genetic elements were identified using this approach in our study.
154
To circumvent the limitations of transposon mutagenesis, we developed a dual-acting-gene-blocking
(DAGB) approach (Fig. 1). Briefly, another *mexT* DNA fragment was integrated into the neutral site of
the genome, forming the dual-acting-*mexT*. Under QS-required evolution conditions, *P. aeruginosa*
rapidly and spontaneously mutates *mexT* to activate QS in the LasR-null mutant (Kostylev et al., 2019;
Oshri et al., 2018). Our DAGB strategy could ensure at least one functioning *mexT* if another copy of
mexT is mutated in the course of stress evolution. Considering *P. aeruginosa* has an intriguing capacity
for adapting to environments through frequent genome variations (Huang et al., 2019), we hypothesized
that DAGB would encourage the bacterium to switch to alternative regulatory pathways, i.e. none-*mexT*
QS-regulators, to achieve rapid QS activation. Through the evolution of the dual-acting-*mexT*-containing
LasR-null mutant and the screening of QS-active single colonies, previously unknown none-*mexT*
genetic elements could be identified through localization of the spontaneous mutations.
166
Evolution of the recombinant LasR-null mutant and screening of QS-active colonies

The recombinant LasR-null mutant bearing dual-acting *mexT* was routinely evolved in the casein-based
broth, in which QS is activated to produce extracellular proteases required for cell growth (Diggle et al.,
2007; Sandoz et al., 2007). Evolving cells were isolated and screened for QS-active phenotypes. Clear
evolving cells started to turn turbid at 25d, implying the event of genome adaptation that allows for the
production of QS-controlled proteases (Fig. 2). QS activation was confirmed by the elevated pyocyanin
in separated protease-positive colonies. Sequencing of *mexT* in those colonies also validated that at least
one copy of *mexT* is intact. Therefore, it was not the inactivation of MexT, but other unknown factors
that triggered the QS activation. Our results also suggest that *mexT* is not the unique determinant for QS-
reprogramming.

The mutation in *rpoA* leads to QS activation
To identify occurring mutations in those protease-positive colonies, we selected three of them for whole-
genome re-sequencing (WGS). The WGS result revealed that they all bear a common single nucleotide
mutation (T4754641->C, T262->A) in *rpoA* (Table S2 and S3), which encodes an α-subunit of RNA
polymerase (RNAP). We therefore focused on *rpoA* in the following study. Since *rpoA* is functionally
essential for cell viability in *P. aeruginosa* (Lee et al., 2015), in-frame deletion of *rpoA* is not available.
To elucidate the roles of *rpoA* in QS regulation, we introduced the same point mutation (T262 ->A) into
the LasR-null mutant, generating the LasR-RpoA-T262A mutant (hereafter designated as LasR-RpoA*).
The incorporation of RpoA* activated QS in the LasR-null background, showing QS-active phenotypes,
including elevated productions of pyocyanin, C4-HSL and PQS as well as cell killing activity (Fig. 3,
Fig. S2 and Fig. S3). We noted that QS activity detected in the LasR-RpoA* mutant is higher than the
LasR-null mutant, while lower than the LasR-MexT mutant. On the other hand, complementation of an
episomal copy of wild-type *rpoA* diminished the QS-controlled proteolysis and pyocyanin yield in the
LasR-RpoA* mutant (Fig. S1). Through DAGB, our study first reports RpoA as a novel QS-
reprogramming regulator.

The LasR-RpoA* mutant exhibits a QS-active transcriptome profile
To investigate the regulatory basis of RpoA* in reprogramming QS, we performed an RNA-seq analysis.
Overall, global expression changes were induced in the LasR-RpoA* mutant, with a total of 182 genes
differentially expressed (at least a 2-fold change, $P < 0.05$) (Fig. 4 and Table S4). As for the QS-related
genes, *rhlI* had a mild induction and the *pqs* operon (including *pqsA/B/C/D/E*) had a strong up-regulation
(Table S4), suggesting RooA* induced the activation of both Rhl and PQS. Similar to the inactivating
MexT, RpoA* also substantially decreased the expression of the *mexEF-oprN* operon encoding the RND
efflux pump. Suppression of the MexEF-OprN pump caused by *mexT* mutations has been reported to
lead to QS reprogramming in the LasR-null background (Kostylev et al., 2019; Lamarche and Déziel,
2011; Oshri et al., 2018). We next compared the transcriptome profile between the LasR-MexT mutant
and the LasR-RpoA* mutant. Given the most up-regulated genes caused by *mexT* mutations are MexEF-
OprN pump-dependent (Tian et al., 2009a), we selected induced genes with at least 4-fold up-regulation
in the LasR-MexT mutant for further analysis. Genome-wide gene expression analysis revealed that those
genes strongly up-regulated in the LasR-MexT mutant were also enriched in the LasR-RpoA* mutant (P
$= 0.01$, one-sided KS test), but not for the genes down-regulated ($P = 0.07$, one-sided KS test). Taken
together, our results showed that RpoA* induced QS activation in the LasR-null background and a
decreased expression of the MexEF-OprN efflux pump, and suggested an efflux pump-induced QS-
active transcriptome profile analogous to the inactivating MexT.

RpoA* directly represses the transcription of the MexEF-OprN efflux pump

We used a transcriptional fusion GFP reporter (P*mexEF*-GFP) to monitor the RpoA*-mediated transcriptional change of the *mexEF-oprN* operon. In line with the result of RNA-seq, RpoA* caused a sharp reduction of the P*mexEF*-GFP (Fig. 5A), indicating a repressed MexEF-OprN effluxe pump. Furthermore, as the inactivating MexEF-OprN pump stimulates the bacterial swarming motility (Lamarche and Déziel, 2011), RpoA* consistently expanded the swarming movement (Fig. S5). Of note, the LasR-RpoA* mutant showed an intermediate expression of the efflux pump between the LasR-null mutant and the LasR-MexT mutant. Because the MexEF-OprN pump negatively regulates QS activity (Lamarche and Déziel, 2011), a weak QS phenotype was observed in the LasR-RpoA* mutant relative to the LasR-MexT mutant (Fig. 3). We next sought out to investigate how RpoA* attenuates the activity of the MexEF-OprN pump. As RpoA is a transcriptional governor, we hypothesized that RpoA* may directly interfere with the binding of RNAP to the promoter of the *mexEF-oprN* operon. To test this hypothesis, we performed an electrophoretic mobility shift assay (EMSA) assay. We recombinantly expressed and purified RNAPs from strain PAO1 either containing the wild-type RpoA or the point-mutated RpoA*. Both types of RNAPs were tested for their binding to the probe DNA that harbors the core promoter of *mexEF-oprN* (Fig. 5B). These RNAPs efficiently bound to the assayed probe (Fig. 5C). In contrast to the wild-type RpoA, RpoA* reduced the RNAP-probe interaction. We therefore concluded that RpoA* down-regulates the activity of the MexEF-OprN efflux pump by reducing the direct contact of RNAP to the promoter DNA.

PQS transport contributes to RpoA*-mediated QS activation

Previous findings showed that inactivation of the MexEF-OprN efflux pump up-regulates the QS activity by reducing the export of the PQS precursor (Köhler et al., 2001; Lamarche and Déziel, 2011). We inferred that RpoA* may activate QS in a similar manner. This was supported by the strong induction of the *pqs* operon shown in RNA-seq (Fig. 4B). We asked the question whether PQS accumulation is responsible for the RpoA*-mediated QS-reprogramming. Given that *pqsA* codes for a coenzyme ligase essential for PQS biosynthesis (Déziel et al., 2004), we deleted *pqsA* from the LasR-RpoA* mutant and evaluated the resulting QS phenotypes. The *pqsA* deletion greatly but not completely diminished the QS-controlled protease activity (Fig. S6). The effects of PQS observed in the LasR-RpoA* mutant are reminiscent of those in the LasR-MexT mutant. Deletion of *pqsA* from the LasR-MexT mutant also did not fully eliminate QS activity (Kostylev et al., 2019). Therefore, inactivating MexT and RopA* both partially rely on PQS to rewire QS activity. Taken together, our data demonstrated that PQS transport contributes to the RpoA*-mediated QS-reprogramming.

The LasR-RpoA* mutant is more resistant to chloramphenicol

We next investigated the clinical meanings of the LasR-RpoA* background, in which a higher expression of the MexEF-OprN efflux pump and a lower QS activity relative to the LasR-MexT mutant. Since the active MexEF-OprN efflux pump promotes antibiotics resistance (Köhler et al., 1997; Ko¨ hler et al., 1997), we hypothesized that in the presence of antibiotics, the LasR-RpoA* mutant would be fitter than the LasR-MexT mutant. To test this hypothesis, we used the LasR-null mutant to compete against either the LasR-MexT mutant or the LasR-RpoA* mutant in the casein-based broth with or without antibiotic treatment. In the absence of antibiotics, the LasR-MexT mutant outcompeted the LasR-null mutant (Fig. 6), in agreement with the previous report that the functioning QS in the LasR-MexT mutant provides the

competitive advantage (Oshri et al., 2018). Likewise, the LasR-RpoA* mutant exhibited a growth advantage over the LasR-null mutant. We next conducted this competition assay exposed to varying concentrations of chloramphenicol, a substrate modulated by the MexEF-OprN pump (Maseda et al., 2000). In the presence of chloramphenicol, this efflux pump is important for the bacterium to acquire antibiotic resistance. Our study found that with increased concentrations of chloramphenicol, the growth advantage of either the LasR-MexT mutant or the LasR-RpoA* mutant over the LasR-null mutant was quickly eliminated. In particular, the growth fitness declined slower in the LasR-RpoA* mutant than in the LasR-MexT mutant. This competition results showed that in the QS-required condition, the LasR-RpoA* mutant is more resistant to chloramphenicol than the LasR-MexT mutant when cocultured with the LasR-null mutant.

Further discovery of QS-reprogramming regulators

To further identify unknown QS-reprogramming genetic determinants, we conducted another round of evolution with the same LasR-null recombinant strain bearing dual-*mexT*. In this batch experiment, we found that QS-active colonies had mutations in *gshA*, which encodes a glutamate-cysteine ligase. We further illustrated that like RpoA*, the GshA mutant could also reprogram the QS hierarchy in the LasR-null background in a MexEF-OprN-dependent manner (detailed description in the accompanying manuscript). This novel functional role of GshA in QS-reprogramming has not yet been realized before.

Iterative application of the DAGB approach

We are curious and aim to dig out more QS-reprogramming genetic determinants. To this end, we iteratively applied the DAGB approach. We introduced another copy of *rpoA* into the dual-*mexT*-containing strain, generating a new recombinant LasR-null strain simultaneously bearing dual-*mexT* and dual-*rpoA*. With this strategy, we hypothesized that spontaneous mutations for inactivating either *mexT* or *rpoA* would be blocked during evolution, allowing for quick identification of QS-reprogramming determinants that are none-*mexT* and none-*rpoA*. The new recombinant LasR-null strain was likewise evolved and screened for QS-active phenotypes. Our study found that QS-active colonies had mutations in *mexE*, coding for a membrane fusion protein of the MexEF-OprN pump (Ko¨hler et al., 1997). Like MexT, disruption of MexE could also induce QS-reprogramming in the LasR-null mutant (Kostylev et al., 2019). Combining our findings, MexT, RpoA, GshA and MexE are either positive regulators or direct components of the MexEF-OprN efflux pump (Fig. 7), suggesting the critical role of the MexEF-OprN regulatory circuit in QS-reprogramming. Taken together, using DAGB-based iteration our research further identified genetic determinants capable of reprogramming QS in the LasR-null background. More importantly, our study set up an example for using DAGB-based iteration to identify underlying genetic factors in other research fields in a broader spectrum of organisms (Fig. 8).

Discussion

Our study offers an innovative approach, named DAGB, for discovering unknown pathways or pathway components in functional studies. Using DAGB, we first uncovered critical non-MexT regulators that could reprogram the QS hierarchy in the *P. aeruginosa* LasR-null mutant. In a comparison of current approaches, DAGB can rapidly identify untapped pathways or related components. For example, the QS-reprogramming function of RpoA* could not be revealed if using the traditional transposon insertional mutagenesis. Because *rpoA* is a housekeeping gene essential for cell viability (Lee et al., 2015), it could not be arbitrarily inserted by the transposon element. As a starting point, the present study employed

DAGB to investigate unknown QS-associated factors in *P. aeruginosa*. We believe that the DAGB-based approach has great potential to be broadly applicable to a wealth of related research fields in other organisms. The application of DAGB would generate new opportunities for scientists to gain insight into dark fields and yield fundamental discoveries.

Our study uncovers a novel QS-reprogramming role for RpoA*, deepening our functional understanding of RpoA as a central transcriptional governor. Adaptive mutations in *rpoA* have been found to be beneficial for bacterial growth in long-term evolution experiments in different organisms (Le Gac et al., 2013; Rajaraman et al., 2016). Previous studies showed that *rpoA* mutations in *B. subtilis* and *E. coli* promote the utilization of secondary carbon sources and ribosome synthesis (Murayama et al., 2015). The QS-related function of RpoA* here implicates that specific residue substitution leads to a distinct functionality configuration of RpoA. Our finding is in agreement with prior studies that certain residues in the α-CTD domain of RpoA are important for promoter contact, while a subset of residues are required for the interaction with transcriptional effectors (Finney et al., 2002; Gaal et al., 1996; Murakami et al., 1996). This is demonstrated by our observation that RpoA* (T262->A) reduces the contacting of the RNAP to the promoter of the *mexEF-oprN* operon (Fig. 5). All these combined findings suggest that adjustment of the performance of RpoA through specific amino acid substitution allows for bacterial quick adaptation to challenging environments.

Inactivation of the MexEF-OprN pump has been reported to lead to QS-reprogramming in the LasR-MexT mutant (Kostylev et al., 2019; Oshri et al., 2018). Likewise, the RpoA* mutation identified in our study could also activate QS activity in a MexEF-OprN-dependent manner. Transcriptome analysis, EMSA and QS-controlled phenotypes collectively suggested that RpoA* affects the QS network through the MexEF-OprN pump. MexEF-OprN-dependent QS activation relies on the transport of the PQS precursor (Köhler et al., 2001; Lamarche and Déziel, 2011). However, deletion of key components of the PQS system, such as PqsA and/or PqsE (Déziel et al., 2004; Diggle et al., 2006), only partially reduced QS activity in the LasR-MexT mutant (Kostylev et al., 2019). How the MexEF-OprN pump shapes the QS circuit awaits further classification. Similarly, the transport of the PQS molecule could not completely account for RpoA*-mediated QS activation (Fig. S6).

RND family multidrug-resistance pumps promote *P. aeruginosa* resistance to a variety of antibiotics (Schweizer, 2003). The *mexEF-oprN* operon codes for a typical RND efflux pump, which renders the bacterium resistant to chloramphenicol, fluoroquinolones and trimethoprim (Lister et al., 2009). Overexpression of the MexEF-OprN reduces QS activity (Köhler et al., 2001) due to the exportation of the PQS precursor (Köhler et al., 2001; Lamarche and Déziel, 2011). Thus, the expression of the efflux pump ultimately affects QS-controlled bacterial virulence. Previous studies found LasR-null, QS-active isolates in the lungs of chronically infected CF patients (Bjarnsholt et al., 2010; Chen et al., 2019; Feltner et al., 2016), where antibiotic treatments are rather common (Winstanley et al., 2016). This raises an open question that how the bacterium solves the conflict between antibiotic resistance and QS activation. In such a scenario, expression of the efflux pump is required for antibiotic resistance while impairing the QS circuits that control the production of virulence factors. Our findings provide an insight into how the bacterium solves this conflict. A single nucleotide substitution in the transcriptional governor RpoA could postpone this conflict by fine-tuning the expression of the MexEF-OprN pump. This investment tradeoff between antibiotic resistance and QS activation gave rise to a slower decline of growth fitness

in the RpoA* mutant when treated with chloramphenicol (Fig. 6). Therefore, a certain degree of compromise between antibiotic resistance and QS regulation could be achieved by adjusting the expression of the RND efflux pump.

P. aeruginosa has a complicated regulatory network (Huang et al., 2019) and undergoes evolutionary changes in the process of chronic infections (Winstanley et al., 2016). This is reflected in our findings that when inactivating MexT is not immediately available, the bacterium is able to quickly switch to alternative pathways for achieving rapid QS activation. Given that the MexEF-OprN-independent regulon of MexT is potentially important for bacterium-host interactions (Tian et al., 2009a; Tian et al., 2009b), it is not hard to imagine that maintaining a wild-type MexT might be critical in certain clinical environments. That perhaps explains previous observations that numerous LasR-null, QS-active isolates from CF chronic infections remain functioning MexT (Cruz et al., 2020; Smith et al., 2006). Our findings uncover non-MexT regulators that control the expression level of the MexEF-OprN efflux pump and account for QS-reprogramming. Future studies need to classify the link between those non-MexT regulators and specific clinical conditions.

Materials and methods

Bacterial strains and growth

Pseudomonas aeruginosa wild-type strain PAO1 and LasR-null mutant were obtained from E. Peter Greenberg (University of Washington, USA). The PAO1 strain and the mutant derivatives were grown in Luria Bertani (LB) broth containing 10 mg/ml tryptone, 5 mg/ml yeast extract, 10 mg/ml NaCl at 37°C. LB broth was buffered with 50mM 3-(N-morpholino) propanesulfonic acid, pH 7.0 (LB-Mops broth). The photosynthetic medium (PM) (Kim and Harwood, 1991) supplemented with 1% sodium caseinate (PM-casein medium)(Cat. No. 9005-46-3, Sigmaaldrich, USA) as the sole carbon source was used for evolution culture at 37°C. Unless otherwise specified, *P. aeruginosa* strains were cultured in 14-mm FALCON tubes (Corning, USA) containing 3 ml medium, with shaking (220 RPM) at 37°C. *Escherichia coli* was grown in LB broth at 37°C. Bacterial strains and plasmids used in this study are listed in Supplementary Table S5.

Construction of *P. aeruginosa* mutants

Either point mutation or full gene knocking out was based on the homologous recombination exchange approach as described previously (Rietsch et al., 2005). Briefly, about 500 ~ 1000 bp of DNA flanking the targeted single nucleotide substitution or full length of gene of interest were PCR-amplified and cloned into pEXG-2 vector (Gentamycin resistance, Gm) (Rietsch et al., 2005; Stover et al., 2000) with the Vazyme ClonExpress II One Step Cloning kit (Vazyme Biotech, Nanjing, China), generating pEXG-flanking constructs. The pEXG-flanking construct was mobilized into *P. aeruginosa* strain by triparental mating with the help of *E. coli* PRK2013 strain (Kanamycin resistance, Km). Point mutation or deletion mutants were first selected on *Pseudomonas* Isolation agar (PIA) containing 100 μg/ml gentamycin (Gm) and further selected on LB agar containing 10% sucrose. All mutants were confirmed by PCR amplification and subsequent DNA Sanger sequencing. The primers used in this study are listed in supplementary TableS6.

Complementation of the RpoA* mutant

The *rpoA* gene was cloned into PJN105 expression vector with the Vazyme ClonExpress II One Step cloning kit (Vazyme Biotech, Nanjing, China), generating plasmid PJN105-*rpoA*. The plasmid PJN105-*rpoA* was mobilized into RpoA* mutant. The expression of RpoA in the complementation strain was induced by adding 1mM L-arabinose and grown in LB-Mpos broth at 37°C for 20 h.

Evolution experiment

Three independent colonies of recombinant LasR-null mutant bearing a miniTn7-MexT construct at the neutral site of the genome were separately inoculated into 3 ml LB-Mops broth for overnight culture. Cell evolution was initiated by transferring 100 μL bacteria grown in LB-Mops into 3 ml PM medium in 14-mm FALCON tubes (Corning, USA). Passage was performed every 5 days by exchanging 100 μL bacteria into 3 ml fresh PM medium, and added sterile water replenish evaporated water in halfway. To identify protease-positive variants, evolved colonies were isolated on LB agar and spotted onto the skim milk plates at 37°C for 24h to monitor the protease-catalyzed zones.

Pyocyanin measurement

Overnight cultures of *P. aeruginosa* grown in LB-mops broth were diluted into 4 ml *Pseudomonas* P broth (20g/L pancreatic digest of gelatin,1.4g/L magnesium chloride,10g/L potassium sulfate) to reach a starting $OD_{600} \approx 0.02$ and cultured at 37°C for 20 h. The cells were spin down at 16,000 g × 2 min and supernatants were measured at OD_{695} nm. Pyocyanin production was determined by plotting the OD_{695} nm / OD_{600} nm over time.

Cyanide measurement

We used a cyanogenic test paper method to detect cyanide. The test paper is a circle of Whatman 3MM chromatography paper soaked in the HCN detection reagent: copper(II) ethyl acetoacetate (5mg) and 4,4'-methylenebis-(N, N-dimethylaniline) (5 mg) in chloroform (1 to 2 ml). Briefly, cells grown overnight in LB-Mops broth were used to spot-inoculate 2% peptone agar plates, and after 12 h at 37°C, the plates were overlaid with test paper and incubated an additional 12 h at 37°C. The paper turns blue when exposed to cyanide. The Image J software was used to quantify the production of hydrogen cyanide.

Skim milk assay

Total proteolytic activities of *P. aeruginosa* strains were evaluated through skim milk assay, in which the tested strains form a zone of clearing on skim milk agar plate. Individual colonies were spotted on the skim milk agar plates (25% LB, 4% skim milk, 1.5% agar). The protease-catalyzed zones were photographed after incubation at 37°C for 24 h.

Swarming motility assay.

PAO1 strains were cultured in LB-Mops broth overnight to $OD_{600} \approx 1.8$. Then bacterial suspensions (1.5 μL; OD_{600} adjusted to 0.5) were added to agar plate (0.5% Bactor-agar, 8g/liter nutrient broth, and 5 g/liter glucose). Bacteria were incubated at 37°C for 16-18 h.

Labelling of miniTn7-Gm

The pUC18T-mini-Tn7T-Gm (NCBI accession number: AY599232) (Choi and Schweizer, 2006a) was integrated into the neutral site of genome of PAO1 strains together with the transformation of helper plasmid pTNS2 (NCBI accession number: AY884833). Integration event was confirmed by PCR

432 amplification and DNA sequencing. The excision of Gm resistance was performed with pFLP2 plasmid
433 (NCBI accession number: AF048702) (Choi and Schweizer, 2006b) and selected on LB agar containing
434 5% sucrose.
435

Reporter assay
P*rhlA*-GFP (Feltner et al., 2016) and P*pqsA*-GFP were used to quantified the RhlR- and PQS-responsive
activities, respectively. P*mexE*-GFP was used to assess the expression of the *mexEF-oprN* operon.
Reporter plasmids were mobilized into PAO1 strains and selected on LB agar plate (Gm). PAO1 strains
bearing QS reporter plasmids were grown in LB-Mops broth containing 50 mg/ml gentamycin for 12 h,
then diluted to LB-Mops broth (containing 50 mg/ml gentamycin) ($OD_{600} \approx 0.02$) and grown 18 h to
stationary phase and subsequently transferred to 96-well plates (200 ul/well) with three technique
replicates. The fluorescence (excitation 488 nm, emission 525 nm) and optical density (OD_{600}) of the
samples were recorded using a microplate reader machine (Synergy H1MF, BioTek Instruments,
Winooski, VT, USA).
446

Quantification of C4-HSL and PQS
P. aeruginosa strains were cultured overnight in LB-Mops broth. Bacteria with a starting $OD_{600} \approx 0.02$
were then grown in 4 ml LB broth at 37°C for 18 h. AHLs were extracted with equal amount of ethyl
acetate. Reporter *E. coli* strains containing pECP61.5 (Feltner et al., 2016) was used to quantify C4-HSL
in the ethyl acetate extract. Precise quantification of PQS was performed by LC/MS, as described
previously (Déziel et al., 2004). 10 μl of the ethyl acetate extraction sample was subjected into liquid
chromatography-mass spectrometry (LC-MS) detection system (Q-Exactive Focus/Ultimate 3000,
Thermo Scientific, USA) equipped with a 100*2.1mm,1.7 um ACQUITY UPLC HSS T3
chromatographic column (Waters, USA). 1% acidified water/methanol gradient was used as the mobile
phase at a flow rate of 0.4 ml min at 40°C. PQS was quantified by measuring the relative chromatographic
area of PQS molecular ions in different samples and multiplying by the concentration of the added
internal standard.
459

Production and purification of RpoA variant protein
Full-length of His-tagged RpoA or RpoA* variant was cloned into PJN105 expression vector and
overexpressed in the LasR-RpoA* mutant induced by 1mM L-arabinose at 37°C for 18 h. Cells were
pelleted at 13,000 g × 5 min and resuspended in lysis buffer (PBS supplemented with 1 ×Protease
Inhibitor Cocktail (Bimake, US), 1mg/ml lysozyme). Resuspended cells were lysed in high pressure
homogenizer machine (parameter 420 MPa, 4200 bar, Stansted Fluid power ltd, United Kingdom) and
pelleted at 13,000 g × 5 min at 4°C. The soluble fraction was incubated with Ni-NTA beads (Qiagene,
Shanghai, China) at 4°C for 2h. The mixture was loaded on a chromatographic column and washed with
low concentration of imidazole buffer (500 mM NaCl, 20 mM NaH_2PO4, 20mM imidazole, pH 8.0),
eluted with high concentration of imidazole buffer (500 mM NaCl, 20 mM NaH_2PO4, 250mM imidazole,
pH 8.0). Peak fractions were assessed by SDS PAGE analysis. Pooled fractions were concentrated with
ultrafiltration device (Millipore, USA) and eluted with PBS buffer, and stored at -80°C.
472

Electrophoretic mobility shift assay (EMSA)
The promoter sequence of the *mexEF-oprN* operon (P*mexE*) was used as probe for EMSA. The probe
DNA was labeled with biotin Biotin-11-UTP (Jena Bioscience, Germany) through PCR amplification

using T4 DNA polymerase (Novoprotein, China). The labeled probe (30 ng) was incubated with 0.2 μg

purified His-tagged RpoA or RpoA* in binding buffer (10 mM Tris-HCl, 50 mM KCl, 1 mM DTT, 1%

glycerin,10mM $MgCl_2$,1 μg/μL Poly (dI•dC), pH 7.5) at room temperature for 30min. DNA-protein

complex was subjected to electrophoresis on 5% polyacrylamide gel in 0.5×Tris-borate-EDTA (TBE)

buffer at 100 V for 180 min. DNA subsequently was transferred to a nylon membrane and incubated with

milk blocking buffer (50 g/L milk powder in TBST buffer, pH 7.5) for 30 min with shaking at room

temperature. The membrane was then incubated with streptavidin-horseradish peroxidase conjugate

(Thermo Scientific, USA) in blocking buffer for 30 min at room temperature. The membrane was

subsequently washed with TBST buffer (50 mM Tris, 150mM NaCl, 0.05% Tween-20, pH 7.5) for 4

time, 5 min each time. DNA was visualized using the Immobilon Western kit (Millipore, USA) and

photographed by Tanon 5200 Multi-Imaging System (Tanon, Shanghai, China).

Mammalian cell cytotoxicity assay

A549 cells were cultured in DMEM medium (GIBCO, USA) and F-12 medium (GIBCO, USA)

supplemented 10% FBS (GIBCO, USA), respectively, and maintained in 5% CO_2 at 37°C. Exponentially

growing bacteria (OD600≈1.0) in LB broth were washed with PBS and next were subjected to near-

confluent A549 cells at a starting multiplicity of infection (MOI) of 10: 1 and incubated at 37℃ for 8 h.

The extent of cell killing as determined by quantifying the release of lactate dehydrogenase in cell

supernatants with the LDH Cytotoxicity Detection Kit (Beyotime, Nantong, China).

Competition experiment

All assayed strains were grown in PM medium supplemented with 0.5% casein amino acid (Sangon,

Shanghai, China) for overnight and adjusted to OD_{600}≈0.02 in PM-Casein medium with or without

chloramphenicol. The strain tagged with pUC18T-mini-Tn7T-Gm cocultured with the unlabeled strain

at a starting ratio of 5:95. The mixed cells was then grown at 37°C for 48h. Cell counts were determined

by spreading bacteria onto LB agar plate with or without Gm-resistance. The relative fitness is calculated

as the ratio of Malthusian growth parameters $(w) = \ln(X_1/X_0)/\ln(Y_1/Y_0)$, according to previous definition

(Lenski et al., 1991).

RNA extraction and sequencing

PAO1 strains and derivatives were grown in LB-MOPS broth at 37°C for 18 h (OD_{600} ≈ 3.8-4.0). Total

RNA was isolated using TRIzol™ Reagent (ThermoFisher, HongKong, China). Total extracted quantity

was measured using NanoDropTM 1000 spectrophotometer (Thermo, US). Extracted RNA was DNase

treated (RQ1 RNase-free DNase I, Promega, Beijing, China) to remove genomic DNA contamination.

The obtained total RNA (2 independent RNA samples per strain) was sent for stranded paired-end

mRNA-seq sequencing in Novagen (Tianjin, China) using the Illumina Novaseq Platform.

Quantitative real-time PCR (qRT-PCR)

Total RNA of PAO1 strains and derivatives were reverse transcribed using the HiScript II 1st Strand

cDNA Synthesis Kit (Vazyme Biotech, Nanjing, China) following the manufacturer's instructions. The

obtained cDNA was used for qRT-PCR. Quantification reactions containing SYBR qPCR Master Mix

(Vazyme Biotech, Nanjing China) were prepared in 96-well plates and run in StepOnePlus Real-Time

PCR System (Applied Biosystems, USA) as recommended. Primers used for qRT-PCR are listed in Table

S6. The expression of targets of interest was normalized to the expression level of the *proC* gene.

Reactions were performed in triplicate.
521
Transcriptome analysis
RNA-seq short reads were filtered with a customized Perl script to remove Illumina adapters and low
quality of reads (Phred Score, Q < 30). The quality of filtered reads was assessed by FastQC. Trimmed
reads were then mapped to the *P. aeruginosa* PAO1 reference genome (NC_002516.2) using Hisat2. The
SAM files were further converted to BAM files by Samtools. Reads aligned to the reference genome
were counted by HTSeq. The generated count files were further processed un with the R package DESeq2.
Genes were regarded as "differentially expressed" if a fold change $\geqslant 2.0$ and a *P* value $\leqslant 0.05$.
529
Gene expression profile analysis
531 Genes showing expression 4-fold changes in strains were compared. *P* values were computed using a
532 one-sided KS test (Lim et al., 2005).
533
Used Software
The following software were used in this study.
BWA software, version 0.7.15-r1140 (http://bio-bwa.sourceforge.net) (Li and Durbin, 2009);
Cutadapt software, version 1.16 (https://github.com/mafdrcelm/cutadapt/) (Martin, 2011);
Samtools software, version 1.5 (http://samtools.sourceforge.net)(Li et al., 2009);
FastQC, version fastqc_v0.11.5 (https://www.bioinformatics.babraham.ac.uk/projects/fastqc/);
Hisat2, version 2.1.0 (https://daehwankimlab.github.io/hisat2/) (Kim et al., 2015);
HTSeq, version 0.11.1 (https://htseq.readthedocs.io/en/release_0.11.1/count.html) (Anders et al., 2015);
DESeq2 (Love et al., 2014);
Perl software, version v5.22.1 (https://www.perl.org/);
Python software, version v3.8.2 (https:// www. python. org/);
R software, version v3.6.1 (http://www.R-project.org/);
GraphPad Prism software, version 5 (https://www.graphpad.com/);
Image J software, version 1.8.0_172 (http://imagej.nih.gov/ij/).
548
Statistical analysis
550 Statistical analyses were performed using Excel and R software (http://www.R-project.org/).
551
Data Availability
553 RNA-seq data sets have been deposited in the NCBI SRA database (www.ncbi.nlm.nih.gov/sra) under
554 the accession number PRJNA863086.
555
Authors' contributions
557 W.C., H.L., M.L., X.Z., X.C. and W.D. designed experiments. W.C., H.L., M.L., X.Z., X.C. and W.D.
558 conducted the experiments. W.D. performed bioinformatics analyses. W.C. and W.D. wrote the
559 manuscript. All authors reviewed the manuscript.
560
Competing interests
562 The authors declare that they have no competing interests.
563

Funding

This work was supported by the National Natural Science Foundation of China (31771341).

Acknowledgements

We are grateful to Eugene Makeyev (King's College London, UK) for providing useful assistances in bioinformatics analyses. We thank E. Peter Greenberg (University of Washington, USA) reading this manuscript and giving useful comments. We want to express our gratitude to lab members for their assistances and supports with respect to many aspects of this work.

References

Anders, S., Pyl, P.T., and Huber, W. (2015). HTSeq—a Python framework to work with high-throughput sequencing data. *Bioinformatics 31*, 166-169.

Bjarnsholt, T., Jensen, P.Ø., Jakobsen, T.H., Phipps, R., Nielsen, A.K., Rybtke, M.T., Tolker-Nielsen, T., Givskov, M., Høiby, N., and Ciofu, O. (2010). Quorum sensing and virulence of *Pseudomonas aeruginosa* during lung infection of cystic fibrosis patients. *PloS one 5*, e10115.

Chen, R., Déziel, E., Groleau, M.-C., Schaefer, A.L., and Greenberg, E.P. (2019). Social cheating in a *Pseudomonas aeruginosa* quorum-sensing variant. *PNAS 116*, 7021-7026.

Cheng, X., Lu, M., Qiu, H., Li, Y., Huang, L., and Dai, W. (2022). Spontaneous quorum-sensing hierarchy reprogramming in *Pseudomonas aeruginosa* laboratory strain PAO1. *AMB Express 12*, 6.

Choi, K.-H., and Schweizer, H.P. (2006a). mini-Tn7 insertion in bacteria with single attTn7 sites: example *Pseudomonas aeruginosa*. *Nature protocols 1*, 153.

Choi, K.-H., and Schweizer, H.P. (2006b). mini-Tn 7 insertion in bacteria with single att Tn 7 sites: example *Pseudomonas aeruginosa*. *Nature protocols 1*, 153.

Cruz, R.L., Asfahl, K.L., Van den Bossche, S., Coenye, T., Crabbé, A., and Dandekar, A.A. (2020). RhlR-regulated acyl-homoserine lactone quorum sensing in a cystic fibrosis isolate of *Pseudomonas aeruginosa*. *MBio 11*.

Déziel, E., Lépine, F., Milot, S., He, J., Mindrinos, M.N., Tompkins, R.G., and Rahme, L.G. (2004). Analysis of *Pseudomonas aeruginosa* 4-hydroxy-2-alkylquinolines (HAQs) reveals a role for 4-hydroxy-2-heptylquinoline in cell-to-cell communication. *PNAS 101*, 1339-1344.

Diggle, S.P., Cornelis, P., Williams, P., and Cámara, M. (2006). 4-quinolone signalling in *Pseudomonas aeruginosa*: old molecules, new perspectives. *Int J Med Microbiol 296*, 83-91.

Diggle, S.P., Griffin, A.S., Campbell, G.S., and West, S.A. (2007). Cooperation and conflict in quorum-sensing bacterial populations. *Nature 450*, 411-414.

Feltner, J.B., Wolter, D.J., Pope, C.E., Groleau, M.-C., Smalley, N.E., Greenberg, E.P., Mayer-Hamblett, N., Burns, J., Déziel, E., and Hoffman, L.R. (2016). LasR variant cystic fibrosis isolates reveal an adaptable quorum-sensing hierarchy in *Pseudomonas aeruginosa*. *MBio 7*, e01513-01516.

Finney, A.H., Blick, R.J., Murakami, K., Ishihama, A., and Stevens, A.M. (2002). Role of the C-terminal domain of the alpha subunit of RNA polymerase in LuxR-dependent transcriptional activation of the lux operon during quorum sensing. *J Bacteriol 184*, 4520-4528.

Gaal, T., Ross, W., Blatter, E.E., Tang, H., Jia, X., Krishnan, V.V., Assa-Munt, N., Ebright, R.H., and Gourse, R.L. (1996). DNA-binding determinants of the alpha subunit of RNA polymerase: novel DNA-binding domain architecture. *Genes Dev 10*, 16-26.

Gellatly, S.L., and Hancock, R.E. (2013). *Pseudomonas aeruginosa*: new insights into pathogenesis and host defenses. *Pathogens and disease 67*, 159-173.

Gilbert, J., OLIVAS, A., SHOGAN, B.D., VALUCKAITE, V., ZABORIN, A., BELOGORTSEVA, N.,
MUSCH, M., MEYER, F., TRIMBLE, W.L., and AN, G. (2012). Intestinal tissues induce an SNP
mutation in *pseudomonas aeruginosa* that enhances its virulence: possible role in anastomotic leak. *PloS*
One 7.

Hoffman, L.R., Kulasekara, H.D., Emerson, J., Houston, L.S., Burns, J.L., Ramsey, B.W., and Miller,
S.I. (2009). *Pseudomonas aeruginosa* lasR mutants are associated with cystic fibrosis lung disease
progression. *Journal of Cystic Fibrosis 8*, 66-70.

Huang, H., Shao, X., Xie, Y., Wang, T., Zhang, Y., Wang, X., and Deng, X. (2019). An integrated
genomic regulatory network of virulence-related transcriptional factors in *Pseudomonas aeruginosa*.
Nature communications 10, 1-13.

Köhler, T., Epp, S.F., Curty, L.K., and Pechère, J.-C. (1999). Characterization of MexT, the regulator of
the MexE-MexF-OprN multidrug efflux system of *Pseudomonas aeruginosa*. *J Bacteriol 181*, 6300-
6305.

Köhler, T., Michea-Hamzehpour, M., Plesiat, P., Kahr, A.-L., and Pechere, J.-C. (1997). Differential
selection of multidrug efflux systems by quinolones in *Pseudomonas aeruginosa*. *Antimicrobial agents*
and chemotherapy 41, 2540-2543.

Köhler, T., Van Delden, C., Curty, L.K., Hamzehpour, M.M., and Pechere, J.-C. (2001). Overexpression
of the MexEF-OprN multidrug efflux system affects cell-to-cell signaling in *Pseudomonas aeruginosa*.
J Bacteriol 183, 5213-5222.

Kim, D., Langmead, B., and Salzberg, S.L. (2015). HISAT: a fast spliced aligner with low memory
requirements. *Nat Methods 12*, 357-360.

Kim, M.-K., and Harwood, C.S. (1991). Regulation of benzoate-CoA ligase in *Rhodopseudomonas*
palustris. *FEMS Microbiol Lett 83*, 199-203.

Klockgether, J., and Tümmler, B. (2017). Recent advances in understanding *Pseudomonas aeruginosa*
as a pathogen. *F1000Research 6*.

Ko ̈ hler, T., Michéa‑Hamzehpour, M., Henze, U., Gotoh, N., Kocjancic Curty, L., and Pechère, J.C.
(1997). Characterization of MexE–MexF–OprN, a positively regulated multidrug efflux system of
Pseudomonas aeruginosa. *Mol Microbiol 23*, 345-354.

Köhler, T., Van Delden, C., Curty, L.K., Hamzehpour, M.M., and Pechere, J.-C. (2001). Overexpression
of the MexEF-OprN multidrug efflux system affects cell-to-cell signaling in *Pseudomonas aeruginosa*.
J Bacteriol *183*, 5213-5222.

Kostylev, M., Kim, D.Y., Smalley, N.E., Salukhe, I., Greenberg, E.P., and Dandekar, A.A. (2019).
Evolution of the *Pseudomonas aeruginosa* quorum-sensing hierarchy. *PNAS 116*, 7027-7032.

Kulasekara, H.D., Ventre, I., Kulasekara, B.R., Lazdunski, A., Filloux, A., and Lory, S. (2005). A novel
two‑component system controls the expression of *Pseudomonas aeruginosa* fimbrial cup genes. *Mol*
Microbiol 55, 368-380.

Lamarche, M.G., and Déziel, E. (2011). MexEF-OprN efflux pump exports the *Pseudomonas* quinolone
signal (PQS) precursor HHQ (4-hydroxy-2-heptylquinoline). *PloS one 6*, e24310.

Le Gac, M., Cooper, T.F., Cruveiller, S., Médigue, C., and Schneider, D. (2013). Evolutionary history
and genetic parallelism affect correlated responses to evolution. *Mol Ecol 22*, 3292-3303.

Lee, J., and Zhang, L. (2015). The hierarchy quorum sensing network in *Pseudomonas aeruginosa*.
Protein & cell 6, 26-41.

Lee, S.A., Gallagher, L.A., Thongdee, M., Staudinger, B.J., Lippman, S., Singh, P.K., and Manoil, C.
(2015). General and condition-specific essential functions of *Pseudomonas aeruginosa*. *PNAS 112*,

5189-5194.

Lenski, R.E., Rose, M.R., Simpson, S.C., and Tadler, S.C. (1991). Long-term experimental evolution in

Escherichia coli. I. Adaptation and divergence during 2,000 generations. *The American Naturalist 138*,

1315-1341.

Li, H., and Durbin, R. (2009). Fast and accurate short read alignment with Burrows–Wheeler transform.

Bioinformatics 25, 1754-1760.

Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., Marth, G., Abecasis, G., and Durbin,

R. (2009). The sequence alignment/map format and SAMtools. *Bioinformatics 25*, 2078-2079.

Lim, L.P., Lau, N.C., Garrett-Engele, P., Grimson, A., Schelter, J.M., Castle, J., Bartel, D.P., Linsley,

P.S., and Johnson, J.M. (2005). Microarray analysis shows that some microRNAs downregulate large

numbers of target mRNAs. *Nature 433*, 769-773.

Lister, P.D., Wolter, D.J., and Hanson, N.D. (2009). Antibacterial-resistant *Pseudomonas aeruginosa*:

clinical impact and complex regulation of chromosomally encoded resistance mechanisms. *Clin*

Microbiol Rev 22, 582-610.

Love, M.I., Huber, W., and Anders, S. (2014). Moderated estimation of fold change and dispersion for

RNA-seq data with DESeq2. *Genome Biology 15*, 550.

Luong, P.M., Shogan, B.D., Zaborin, A., Belogortseva, N., Shrout, J.D., Zaborina, O., and Alverdy, J.C.

(2014). Emergence of the P2 phenotype in *Pseudomonas aeruginosa* PAO1 strains involves various

mutations in *mexT* or *mexF*. *J Bacteriol 196*, 504-513.

Martin, M. (2011). Cutadapt removes adapter sequences from high-throughput sequencing reads.

EMBnet journal 17, 10-12.

Maseda, H., Yoneyama, H., and Nakae, T. (2000). Assignment of the substrate-selective subunits of the

MexEF-OprN multidrug efflux pump of *Pseudomonas aeruginosa*. *Antimicrobial agents and*

chemotherapy 44, 658-664.

Murakami, K., Fujita, N., and Ishihama, A. (1996). Transcription factor recognition surface on the RNA

polymerase alpha subunit is involved in contact with the DNA enhancer element. *The EMBO Journal 15*,

4358-4367.

Murayama, S., Ishikawa, S., Chumsakul, O., Ogasawara, N., and Oshima, T. (2015). The role of α-CTD

in the genome-wide transcriptional regulation of the *Bacillus subtilis* cells. *PloS one 10*, e0131588.

Oshri, R.D., Zrihen, K.S., Shner, I., Omer Bendori, S., and Eldar, A. (2018). Selection for increased

quorum-sensing cooperation in *Pseudomonas aeruginosa* through the shut-down of a drug resistance

pump. *ISME J 12*, 2458-2469.

Papenfort, K., and Bassler, B.L. (2016). Quorum sensing signal–response systems in Gram-negative

bacteria. *Nat Rev Microbiol 14*, 576.

Rajaraman, E., Agarwal, A., Crigler, J., Seipelt-Thiemann, R., Altman, E., and Eiteman, M.A. (2016).

Transcriptional analysis and adaptive evolution of *Escherichia coli* strains growing on acetate. *Appl*

Microbiol Biotechnol 100, 7777-7785.

Richardot, C., Juarez, P., Jeannot, K., Patry, I., Plésiat, P., and Llanes, C. (2016). Amino acid

substitutions account for most MexS alterations in clinical nfxC mutants of *Pseudomonas aeruginosa*.

Antimicrobial agents and chemotherapy 60, 2302-2310.

Rietsch, A., Vallet-Gely, I., Dove, S.L., and Mekalanos, J.J. (2005). ExsE, a secreted regulator of type

III secretion genes in *Pseudomonas aeruginosa*. *PNAS 102*, 8006-8011.

Sandoz, K.M., Mitzimberg, S.M., and Schuster, M. (2007). Social cheating in *Pseudomonas aeruginosa*

quorum sensing. *PNAS 104*, 15876-15881.

Schweizer, H.P. (2003). Efflux as a mechanism of resistance to antimicrobials in *Pseudomonas*
aeruginosa and related bacteria: unanswered questions. *Genet Mol Res 2*, 48-62.

Smith, E.E., Buckley, D.G., Wu, Z., Saenphimmachak, C., Hoffman, L.R., D'Argenio, D.A., Miller, S.I.,
Ramsey, B.W., Speert, D.P., and Moskowitz, S.M. (2006). Genetic adaptation by *Pseudomonas*
aeruginosa to the airways of cystic fibrosis patients. *PNAS 103*, 8487-8492.

Sobel, M.L., Neshat, S., and Poole, K. (2005). Mutations in PA2491 (*mexS*) promote MexT-dependent
mexEF-oprN expression and multidrug resistance in a clinical strain of *Pseudomonas aeruginosa*. *J*
Bacteriol 187, 1246-1253.

Stover, C.K., Pham, X.Q., Erwin, A., Mizoguchi, S., Warrener, P., Hickey, M., Brinkman, F., Hufnagle,
W., Kowalik, D., and Lagrou, M. (2000). Complete genome sequence of *Pseudomonas aeruginosa*
PAO1, an opportunistic pathogen. *Nature 406*, 959-964.

Tian, Z.-X., Fargier, E., Mac Aogain, M., Adams, C., Wang, Y.-P., and O'Gara, F. (2009a).
Transcriptome profiling defines a novel regulon modulated by the LysR-type transcriptional regulator
MexT in *Pseudomonas aeruginosa*. *Nucleic Acids Res 37*, 7546-7559.

Tian, Z.-X., Mac Aogain, M., O'Connor, H.F., Fargier, E., Mooij, M.J., Adams, C., Wang, Y.-P., and
O'Gara, F. (2009b). MexT modulates virulence determinants in *Pseudomonas aeruginosa* independent
of the MexEF-OprN efflux pump. *Microb Pathog 47*, 237-241.

Winstanley, C., O'Brien, S., and Brockhurst, M.A. (2016). *Pseudomonas aeruginosa* evolutionary
adaptation and diversification in cystic fibrosis chronic lung infections. *Trends Microbiol 24*, 327-337.

Figure legends

Figure 1. Scheme illustrating strategy used for the identification of QS-reprogramming genetic
determinants.
Current strategy identified *mexT* as a QS-reprogramming regulator (top). Dual-acting-gene-blocking
(DAGB) strategy developed in our study (bottom), illustrating the steps to discover none-*mexT* QS-
regulators responsible for QS-reprogramming in the LasR-null background. QS-, inactive QS; QS+,
active QS.

Figure 2. Evolution of the recombinant LasR mutant bearing dual-acting *mexT*.
(A) Frequency of protease-positive colonies in the evolving population cells. Three independent
recombinant LasR-null mutants containing dual-acting *mexT* were initiated for evolution in the casein-
based broth. (B) Pyocyanin production in selected protease-positive colonies. Pyocyanin production was
quantified at OD_{695} divided by OD_{600} of each assayed strain, respectively. LasR-null, LasR-null mutant
as a negative control; LasR-1/2/3/4, evolved recombinant LasR mutant displaying protease-positive.
Data are presented as means $\pm$ SD ($n = 3$). $*P < 0.05$, $**P < 0.01$, $***P < 0.001$ (t-test).

Figure 3. Increased QS activity in the LasR-RpoA* mutant.
(A-B) Rhl- (A) and PQS-responsive (B) QS activities in indicated strains. Rhl- and PQS-responsive QS
activities were reflected by the fluorescence levels of the reporter P*rhlA*-GFP and P*pqsA*-GFP,
respectively. Fluorescence expression were recorded as relative fluorescence unit (RFU) after 18h
incubation of indicated strains. (C-D) Relative concentration of C4-HSL(C) and PQS (D) in the assayed
strains. Productions of AHLs in wild-type (WT) PAO1 was set to 100%. (E) Pyocyanin productions in
assayed strains. Pyocyanin level was quantified at OD_{695} divided by OD_{600}. (F) Cyanide levels in

designated strains. Image J software was used to compare the amount of produced cyanide. Cyanide-
sensitive filter papers were photographed after post inoculation of bacteria 18h in the 6-well plate at 37
°C. The amount of cyanide produced in wild-type PAO1 was set to 100%. The experiment was repeated
three times and obtained results were similar. Data are presented as means ± SD ($n = 3$, t-test). *$P <$
0.05, **$P < 0.01$, ***$P < 0.001$ (t-test).

Figure 4. Differentially expressed genes in the LasR-RpoA* mutant.
(A) Volcano plots showing the magnitude of the differential gene expression induced by the RpoA*
mutation. Each dot represents one annotated sequence with detectable expression. Thresholds for
defining a gene significantly differentially expressed (log2(FoldChange) ≧|1.0|, P value ≦0.05) are
shown as dashed lines. Red dots, up-regulated genes; Blue dots, down-regulated genes; Green dots,
undifferentially expressed genes. (B-C) The expression levels of genes in QS genes (B) and the *mexEF-*
oprN operon (C). FoldChange in the expression of each indicated gene was calculated. Data are presented
as mean ± SD.

Figure 5. RpoA* modulates the MexEF-OprN efflux pump.
(A) RpoA* reduces the transcription of the *mexEF-oprN* operon. The transcriptional level of the *mexEF-*
oprN operon was reflected by the P*mexE*-GFP, a GFP reporter fused with the promoter sequence of *mexE*.
(B) Illustration of the probe DNA used for the Electrophoretic mobility shift assay (EMSA). (C) EMSA
assay showing RNAP containing wild-type (WT) RpoA or RpoA* bound to the probe DNA. The
unbound probe DNA and the DNA-protein complex are indicated by arrows. Data are presented as mean
± SD. *$P < 0.05$, **$P < 0.01$, ***$P < 0.001$ (t-test).

Figure 6. Effects of chloramphenicol on the growth fitness of the LasR-RpoA* mutant.
The LasR-RopA* mutant or the LasR-MexT mutant labeled with Gm-resistance (pUC18T-mini-Tn7T-
Gm) was co-cultured with the LasR-null mutant at a start ratio of 5:95 and subjected to PM-casein
medium containing indicated concentrations of chloramphenicol for 48h. Colonies were counted and
relative fitness was computed as the ratio of Malthusian growth parameters (*w*).

Figure 7. Graphic representation of the regulatory circuit of the MexEF-OprN efflux pump
Illustration of the regulatory circuit of the MexEF-OprN efflux pump, containing the known MexT
transcriptional regulator and two new regulators identified in our study, RpoA (α-subunit of the RNAP)
and GshA.

Figure 8. Iterative application of DAGB
Schematic representation of the iterative application of the DAGB approach for identification of
unknown pathway components.

Figure S1. Complementation of a wild-type RpoA into the LasR-RpoA* mutant reduces the QS-
controlled activity.
Complementation of an episomal copy of wild-type *rpoA* gene in the LasR-RpoA* mutant diminished
the QS-controlled proteolytic activity (A) and the pyocyanin production (B). (A) Proteolytic activity
shown by the skim milk plate assay. Photographs were taken after inoculation of equal amount of bacteria
onto the skim milk plate and incubation for 24 hours at 37 °C. (B) Pyocyanin production of indicated

strains. The LasR-null mutant or the LasR-RpoA* mutant was either transformed with an empty vector
PJN105 (Ctr) or the RpoA-expressing vector PJN105-RpoA (RpoA). The RpoA expression was induced
by supplementation of L-arabinose (1 mg/ml). Data are from means $\pm$ SD ($n = 3$). *$P < 0.05$, **$P <$
0.01, ***$P < 0.001$ (*t*-test).

Figure S2. Constructed LasR-RpoA* mutant exhibits elevated proteolytic activity.
Proteolytic activity of designated strains. Equal amount of tested strains were spotted on to the skim milk
plate (A) and photographed after 24h of incubation. (B) Quantification of proteolytic zones in (A). Data
are from means $\pm$ SD ($n = 3$). *$P < 0.05$, **$P < 0.01$, ***$P < 0.001$ (*t*-test).

Figure S3. RpoA* mutation significantly enhances host cell cytotoxicity.
Human cell killing by mutant strains. Human lung carcinoma A549 were treated with indicated strains
in equal amounts. After 6h of inoculation, the relative cytosolic lactate dehydrogenase (LDH) was
quantified. The amount of LDH in wild-type PAO1 was set to 100%. Data are presented as means $\pm$
SD ($n = 3$, *t*-test). *$P < 0.05$, **$P < 0.01$, ***$P < 0.001$ (*t*-test).

Figure S4. qRT-PCR validation of differentially expressed genes obtained from the transcriptomic
analysis.
Indicated genes were validated by qRT-PCR. Relative expression was normalized to the *proC* gene. Used
primers are shown in Table S6. Data are means $\pm$ SD (3 independent RNA extractions; $n = 3$). *P*-values
were obtained from *t*-tests. *$P < 0.05$, **$P < 0.01$, ***$P < 0.001$ (*t*-test).

Figure S5. Swarming motility in the LasR-RpoA* mutant.
(A)Bacterial swarming motility was assayed on 0.5% agar plate. Plates were photographed after 18h
incubation at 37 °C and representative photos are shown. (B) quantification of diameters in colonies. The
RpoA expression was induced by adding L-arabinose (1 mg/ml). LasR-null-Ctr, the LasR-null mutant
carrying an empty PJN105 vector; LasR-RpoA*-Ctr, the LasR-RpoA* mutant having an empty PJN105
vector; LasR-RpoA*-RpoA, the LasR-RpoA* mutant containing a PJN105-RpoA construct expressing
the wild-type RpoA. Data are means $\pm$ SD from a representative experiment ($n = 3$). *P* values were
obtained from *t*-test. *$P < 0.05$, **$P < 0.01$, ***$P < 0.001$. The experiment was repeated twice with
similar results.

Figure S6. Deletion of PqsA reduces RpoA*-mediated QS activation.
(A) Proteolytic activity was measured through spotting the same amount of indicated strains on the skim
milk plate. Plates were photographed after 24h incubation at 37 °C and representative photos are shown.
(B) quantification of proteolytic zone in (A). Data are from means $\pm$ SD ($n = 3$). *$P < 0.05$, **$P < 0.01$,
***$P < 0.001$ (*t*-test).

Supplementary Tables

Table S1. Mutations in lasR and mexT of isolates from chronically infected CF patients.
Table S2. Reads mapping statistics of whole-genome re-sequencing data.
Table S3. Identification of mutations in evolved protease-positive colonies.
Table S4. Genes differentially expressed in the LasR-RpoA* mutant.

Table S5. Strains and plasmids used in this study.

Table S6. Oligonucleotides used in this study.

Current laboratory evolution strategy

Dual-acting-gene-blocking (DAGB) evolution strategy

Fig. 1

Fig.2

Fig.3

A

B QS genes

C Efflux pump genes

Fig. 4

Fig. 5

Fig. 6

Fig.7

Fig. 8