Estimating temporally variable selection intensity from ancient DNA data

Zhangyi He[a,b,1,*], Xiaoyang Dai[c,1], Wenyang Lyu[d], Mark Beaumont[e], Feng Yu[d]

[a]*Cancer Research UK Beatson Institute, Glasgow G61 1BD, United Kingdom*
[b]*Department of Computer Science, University of Oxford, Oxford OX1 3QD, United Kingdom*
[c]*The Blizard Institute, Barts and The London School of Medicine and Dentistry, Queen Mary University of London, London E1 2AT, United Kingdom*
[d]*School of Mathematics, University of Bristol, Bristol BS8 1UG, United Kingdom*
[e]*School of Biological Sciences, University of Bristol, Bristol BS8 1TQ, United Kingdom*

Abstract

Novel technologies for recovering DNA information from archaeological and historical specimens have made available an ever-increasing amount of temporally-spaced genetic samples from natural populations. These genetic time series permit the direct assessment of patterns of temporal changes in allele frequencies, and hold the promise of improving power for inference of selection. Increased time resolution can further facilitate testing hypotheses regarding the drivers of past selection events like plant and animal domestication. However, studying past selection processes through ancient DNA (aDNA) still involves considerable obstacles such as postmortem damage, high fragmentation, low coverage and small samples. To address these challenges, we introduce a novel Bayesian approach for the inference of temporally variable selection based on genotype likelihoods instead of allele frequencies, thereby enabling us to account for sample uncertainties resulting from the damage and fragmentation of aDNA molecules. Also, our method permits the reconstruction of the underlying mutant allele frequency trajectory of the population through time, which allows for a better understanding of the drivers of selection. We evaluate its performance through extensive simulations and illustrate its utility with an application to the ancient horse samples genotyped at the loci for coat colouration.

Keywords: Ancient DNA, Natural selection, Demographic history, Sampling uncertainty, Two-layer hidden Markov model, Particle marginal Metropolis-Hastings

---

[*]Corresponding author.
  Email address: `z.he@beatson.gla.ac.uk` (Zhangyi He)
[1]These authors contributed equally to this work.

1. Introduction

Natural selection is the primary mechanism of adaptive evolution within natural populations (Darwin, 1859). Evolutionary and population geneticists usually study selection and adaptation based on temporally spaced genetic samples, which can provide much more valuable information regarding selection since expected changes in allele frequencies over time are closely related to selection and its strength and timing of changes. The commonest data sources have been evolve and resequencing studies combining experimental evolution under controlled laboratory or field mesocosm conditions with next-generation sequencing technology, which however are typically limited to the species with small evolutionary timescales (*e.g.*, Turner & Miller, 2012; Bosshard et al., 2017; Good et al., 2017). Recent advances in technologies for obtaining DNA molecules from ancient biological material have given rise to massive increases in time serial samples of segregating alleles from natural populations (*e.g.*, Mathieson et al., 2015; Librado et al., 2017; Fages et al., 2019), which offer unprecedented opportunities to study the chronology and tempo of selection across evolutionary timescales (see Dehasque et al., 2020, for a review).

As the number of published ancient genomes is growing rapidly, a range of statistical methods that estimate selection coefficients and other population genetic parameters from ancient DNA (aDNA) data have been developed over the last fifteen years (see Malaspinas, 2016, for a review). Most existing methods are based on the hidden Markov model (HMM) framework developed by Williamson & Slatkin (1999), where the underlying population allele frequency is modelled as a hidden state following the Wright-Fisher model introduced by Fisher (1922) and Wright (1931), and the observed sample allele frequency drawn from the underlying population at each sampling time point is modelled as a noisy observation of the underlying population allele frequency. To remain computationally feasible, these methods typically work with the diffusion approximation of the Wright-Fisher model (*e.g.*, Bollback et al., 2008; Malaspinas et al., 2012; Steinrücken et al., 2014; Schraiber et al., 2016; Ferrer-Admetlla et al., 2016; He et al., 2020a,b; Lyu et al., 2022). The diffusion approximation enables us to efficiently integrate over all possible underlying allele frequency trajectories, therefore producing substantial reductions in computational cost of the likelihood calculation. These approaches have already been successfully applied in aDNA studies (*e.g.*, Ludwig et al., 2009; Sandoval-Castellanos et al., 2017; Ye et al., 2017; Wutke et al., 2018). Moment-based approximations of the Wright-Fisher model, as tractable alternatives, are usually

used in the existing methods tailored to experimental evolution (*e.g.*, Lacerda & Seoighe, 2014; Feder et al., 2014; Terhorst et al., 2015; Paris et al., 2019) owing to their poor performance for large evolutionary timescales (He et al., 2021).

Although the field of aDNA is experiencing an exponential increase in terms of the amount of available data, accompanied by an increase in available tailored statistical approaches, data quality remains a challenge due to postmortem damage, high fragmentation, low coverage and small samples. To our knowledge, none of the existing methods are tailored to model two main characteristics of aDNA, *i.e.*, the high error rate resulting from the damage of aDNA molecules and the high missing rate caused by the fragmentation of aDNA molecules, with the exception of He et al. (2020a) in which aDNA samples with missing genotypes are allowed. Moreover, most existing approaches assume that the selection coefficient is fixed over time, which can be easily violated in aDNA studies. Most scenarios of adaptation in natural populations generally involve adaptation to ecological, environmental and cultural changes, where it is no longer appropriate for the selection coefficient to remain constant through time. More recently, Mathieson (2020) introduced a novel approach to infer selection and its strength and timing of changes from allele frequency time series data with an application to aDNA studies. However, this approach is still built upon the assumption of no missing genotypes and sequencing errors in aDNA data.

To address these challenges, we introduce a novel Bayesian approach for estimating temporally variable selection intensity from aDNA data while modelling sample uncertainties resulting from postmortem damage, high fragmentation, low coverage and small samples. Unlike existing methods, our approach is built upon a two-layer HMM framework, where the first hidden layer characterises the underlying frequency trajectory of the mutant allele in the population through time, the second hidden layer denotes the unobserved genotype of the individual in the sample, and the third observed layer represents the data on aDNA sequences. Such an HMM framework allows for genotype likelihoods as input instead of allele frequencies, therefore modelling sample uncertainties caused by the damage and fragmentation of aDNA molecules. Our posterior computation is carried out through the particle marginal Metropolis-Hastings (PMMH) algorithm developed by Andrieu et al. (2010), which permits a joint update of the selection coefficient and population mutant allele frequency trajectory. The reconstruction of the underlying frequency trajectory of the mutant allele in the population allows for a better understanding of the drivers

of selection.

We test our approach through extensive simulations, in particular when samples are sparsely distributed in time with small sizes and in poor quality (*i.e.*, high missing rate and error rate). We analyse the ancient horse samples from Wutke et al. (2016) that were genotyped at the loci encoding coat colouration to illustrate the applicability of our method on aDNA data.

2. Materials and Methods

In this section, we begin with an introduction of the Wright-Fisher diffusion for a single locus evolving under selection over time. We describe our Bayesian procedure for inferring temporally variable selection from the data on aDNA sequences while modelling sample uncertainties caused by the damage and fragmentation of aDNA molecules.

2.1. Wright-Fisher diffusion

Let us consider a diploid population of N randomly mating individuals at a single locus $\mathcal{A}$, which evolves subject to selection under the Wright-Fisher model (see, *e.g.*, Durrett, 2008). We assume discrete time, non-overlapping generations and non-constant population size. Suppose that at locus $\mathcal{A}$ there are two possible allele types, labelled $\mathcal{A}_0$ and $\mathcal{A}_1$, respectively. We attach the symbol $\mathcal{A}_0$ to the ancestral allele, which originally exists in the population, and the symbol $\mathcal{A}_1$ to the mutant allele, which arises in the population only once. We let selection take the form of viability selection and set per-generation relative viabilities of the three possible genotypes $\mathcal{A}_0\mathcal{A}_0$, $\mathcal{A}_0\mathcal{A}_1$ and $\mathcal{A}_1\mathcal{A}_1$ to be 1, $1 + hs$ and $1 + s$, respectively, where $s \in [-1, +\infty)$ is the selection coefficient and $h \in [0, 1]$ is the dominance parameter.

We now consider the standard diffusion limit of the Wright-Fisher model with selection. We measure time in units of $2N_0$ generations, denoted by t, where N_0 is an arbitrary reference population size fixed through time, and assume that the population size changes deterministically, with $N(t)$ being the number of diploid individuals in the population at time t. In the diffusion limit of the Wright-Fisher model with selection, as the reference population size N_0 goes to infinity, the scaled selection coefficient $\alpha = 2N_0 s$ is kept constant and the ratio of the population size to the reference population size $N(t)/N_0$ converges to a function $\beta(t)$. As demonstrated in Durrett (2008), the mutant allele frequency trajectory through time converges to the diffusion

limit of the Wright-Fisher model with the reference population size N_0 approaching infinity. We refer to this diffusion limit as the Wright-Fisher diffusion with selection.

We let X denote the Wright-Fisher diffusion with selection, which models the mutant allele frequency evolving in the state space $[0, 1]$ under selection. Many existing approaches define the Wright-Fisher diffusion X in terms of the partial differential equation (PDE) that characterises its transition probability density function (*e.g.*, Bollback et al., 2008; Steinrücken et al., 2014; He et al., 2020b). Instead, like Schraiber et al. (2016), He et al. (2020a) and Lyu et al. (2022), we characterise the Wright-Fisher diffusion X as the solution to the stochastic differential equation (SDE)

$$dX(t) = \alpha X(t)(1 - X(t))(h + (1 - 2h)X(t))dt + \sqrt{\frac{X(t)(1 - X(t))}{\beta(t)}}dW(t) \qquad (1)$$

for $t \geq t_0$ with initial condition $X(t_0) = x_0$, where W represents the standard Wiener process.

2.2. Bayesian inference of selection

Suppose that the available data are always drawn from the underlying population at a finite number of distinct time points, say $t_1 < t_2 < \ldots < t_K$, measured in units of $2N_0$ generations. At the sampling time point t_k, there are N_k individuals sampled from the underlying population, and for individual n, let $\boldsymbol{r}_{n,k}$ represent, in this generic notation, all of the reads at the locus of interest. The population genetic parameters of interest in this work are the selection coefficient s and dominance parameter h, and for ease of notation, we set $\boldsymbol{\vartheta} = (s, h)$ in what follows.

2.2.1. Hidden Markov model

Our method is built upon the HMM framework introduced by Bollback et al. (2008), where the underlying population evolves under the Wright-Fisher diffusion with selection in Eq. (1), and the observed sample is made up of the individuals independently drawn from the underlying population at each sampling time point. To model sample uncertainties resulting from the damage and fragmentation of aDNA molecules, our approach infers selection from raw reads rather than called genotypes. We let $\boldsymbol{x}_{1:K} = \{x_1, x_2, \ldots, x_K\}$ represent the frequency trajectory of the mutant allele in the underlying population at the sampling time points $\boldsymbol{t}_{1:K}$, and the posterior probability distribution for the population genetic parameters $\boldsymbol{\vartheta}$ and population mutant allele

frequency trajectory $\boldsymbol{x}_{1:K}$ (up to proportionality) can be formulated as

$$p(\boldsymbol{\vartheta}, \boldsymbol{x}_{1:K} \mid \boldsymbol{r}_{1:K}) \propto p(\boldsymbol{\vartheta})p(\boldsymbol{x}_{1:K} \mid \boldsymbol{\vartheta})p(\boldsymbol{r}_{1:K} \mid \boldsymbol{x}_{1:K}), \tag{2}$$

where $\boldsymbol{r}_{1:K} = \{\boldsymbol{r}_1, \boldsymbol{r}_2, \ldots, \boldsymbol{r}_K\}$ with $\boldsymbol{r}_k = \{\boldsymbol{r}_{1,k}, \boldsymbol{r}_{2,k}, \ldots, \boldsymbol{r}_{N_k,k}\}$. In Eq. (2), $p(\boldsymbol{\vartheta})$ is the prior probability distribution for the population genetic parameters and can be taken to be a uniform prior over the parameter space if prior knowledge is poor, $p(\boldsymbol{x}_{1:K} \mid \boldsymbol{\vartheta})$ is the probability distribution for the mutant allele frequency trajectory of the underlying population at all sampling time points, and $p(\boldsymbol{r}_{1:K} \mid \boldsymbol{x}_{1:K})$ is the probability of observing the reads of all sampled individuals given the mutant allele frequency trajectory of the population.

With the Markov property of the Wright-Fisher diffusion, we can decompose the probability distribution $p(\boldsymbol{x}_{1:K} \mid \boldsymbol{\vartheta})$ as

$$p(\boldsymbol{x}_{1:K} \mid \boldsymbol{\vartheta}) = p(x_1 \mid \boldsymbol{\vartheta}) \prod_{k=1}^{K-1} p(x_{k+1} \mid x_k; \boldsymbol{\vartheta}),$$

where $p(x_1 \mid \boldsymbol{\vartheta})$ is the prior probability distribution for the starting population mutant allele frequency, taken to be a uniform distribution over the state space $[0, 1]$ if the prior knowledge is poor, and $p(x_{k+1} \mid x_k; \boldsymbol{\vartheta})$ is the transition probability density function of the Wright-Fisher diffusion X between two consecutive sampling time points for $k = 1, 2, \ldots, K-1$, satisfying the Kolmogorov backward equation (or its adjoint) resulting from the Wright-Fisher diffusion.

To calculate the probability $p(\boldsymbol{r}_{1:K} \mid \boldsymbol{x}_{1:K})$, we introduce an additional hidden layer into our HMM framework to denote the latent genotypes of all sampled individuals (see Figure 1 for the graphical representation of our two-layer HMM framework). We let $\boldsymbol{g}_{1:K} = \{\boldsymbol{g}_1, \boldsymbol{g}_2, \ldots, \boldsymbol{g}_K\}$ be the genotypes of the individuals drawn from the underlying population at the sampling time points $\boldsymbol{t}_{1:K}$ with $\boldsymbol{g}_k = \{g_{1,k}, g_{2,k}, \ldots, g_{N_k,k}\}$, where $g_{n,k} \in \{0, 1, 2\}$ denotes the number of mutant alleles in individual n at sampling time point t_k. We then have

$$p(\boldsymbol{r}_{1:K} \mid \boldsymbol{x}_{1:K}) = \prod_{k=1}^{K} p(\boldsymbol{r}_k \mid x_k) = \prod_{k=1}^{K} \prod_{n=1}^{N_k} \sum_{g_{n,k}=0}^{2} p(g_{n,k} \mid x_k)p(\boldsymbol{r}_{n,k} \mid g_{n,k}),$$

where $p(g_{n,k} \mid x_k)$ represents the probability distribution for the genotype $g_{n,k}$ of individual n in the sample given the population mutant allele frequency x_k, and $p(\boldsymbol{r}_{n,k} \mid g_{n,k})$ represents the

probability of observing reads $\boldsymbol{r}_{n,k}$ of individual n in the sample given the genotype $g_{n,k}$, known as the genotype likelihood. Under the assumption that all individuals in the sample are drawn from the underlying population in their adulthood (*i.e.*, the stage after selection but before reproduction in the life cycle, see He et al. (2017)), we have

$$p(g_{n,k} \mid x_k) = \begin{cases} \dfrac{(1-x_k)^2}{(1-x_k)^2 + (1+hs)2x_k(1-x_k) + (1+s)x_k^2}, & \text{if } g_{n,k} = 0 \\[3mm] \dfrac{(1+hs)2x_k(1-x_k)}{(1-x_k)^2 + (1+hs)2x_k(1-x_k) + (1+s)x_k^2}, & \text{if } g_{n,k} = 1 \\[3mm] \dfrac{(1+s)x_k^2}{(1-x_k)^2 + (1+hs)2x_k(1-x_k) + (1+s)x_k^2}, & \text{if } g_{n,k} = 2. \end{cases} \tag{3}$$

Genotype likelihoods are typically calculated from aligned reads and quality scores in the process of determining the genotype for each individual through genotype calling software (*e.g.*, Li et al., 2009a,b; DePristo et al., 2011), which is an essential prerequisite for most aDNA studies, thereby assuming that these numerical values are available.

Figure 1: Graphical representation of our two-layer HMM framework for the data on aDNA sequences.

2.2.2. Particle marginal Metropolis-Hastings

Since the posterior $p(\boldsymbol{\vartheta}, \boldsymbol{x}_{1:K} \mid \boldsymbol{r}_{1:K})$ is not available in a closed form, we resort to the PMMH algorithm introduced by Andrieu et al. (2010) in this work, which has already been successfully applied in population genetic studies (see, *e.g.*, He et al., 2020b; Lyu et al., 2022). The PMMH algorithm calculates the acceptance ratio with the estimate of the marginal likelihood $p(\boldsymbol{r}_{1:K} \mid \boldsymbol{\vartheta})$ in the Metropolis-Hastings procedure and generates a new candidate of the population mutant allele frequency trajectory $\boldsymbol{x}_{1:K}$ from the approximation of the smoothing distribution $p(\boldsymbol{x}_{1:K} \mid \boldsymbol{r}_{1:K}, \boldsymbol{\vartheta})$, which can both be achieved through the bootstrap particle filter developed by Gordon

et al. (1993). Such a setup permits a joint update of the population genetic parameters $\boldsymbol{\vartheta}$ and population mutant allele frequency trajectory $\boldsymbol{x}_{1:K}$.

A key ingredient of the PMMH algorithm is to construct a bootstrap particle filter that can target the smoothing distribution $p(\boldsymbol{x}_{1:K} \mid \boldsymbol{r}_{1:K}, \boldsymbol{\vartheta})$. More specifically, at the sampling time point t_{k+1}, our objective is to generate a sample from the filtering distribution $p(x_{k+1} \mid \boldsymbol{r}_{1:k+1}, \boldsymbol{\vartheta})$. Up to proportionality, the filtering distribution $p(x_{k+1} \mid \boldsymbol{r}_{1:k+1}, \boldsymbol{\vartheta})$ can be formulated as

$$p(x_{k+1} \mid \boldsymbol{r}_{1:k+1}, \boldsymbol{\vartheta}) \propto p(\boldsymbol{r}_{k+1} \mid x_{k+1})p(x_{k+1} \mid \boldsymbol{r}_{1:k}, \boldsymbol{\vartheta}),$$

where

$$p(x_{k+1} \mid \boldsymbol{r}_{1:k}, \boldsymbol{\vartheta}) = \int_0^1 p(x_{k+1} \mid x_k; \boldsymbol{\vartheta})p(x_k \mid \boldsymbol{r}_{1:k}, \boldsymbol{\vartheta}) \, dx_k \tag{4}$$

is the predictive distribution, but not available in a closed form. From Eq. (4), the predictive distribution $p(x_{k+1} \mid \boldsymbol{r}_{1:k}, \boldsymbol{\vartheta})$ can be approximated with a set of particles $\boldsymbol{x}_k^{1:M} = \{x_k^1, x_k^2, \ldots, x_k^M\}$ generated from the filtering distribution $p(x_k \mid \boldsymbol{r}_{1:k}, \boldsymbol{\vartheta})$ with each particle x_k^m being assigned a weight $w_k^m = 1/M$, thereby

$$\hat{p}(x_{k+1} \mid \boldsymbol{r}_{1:k}, \boldsymbol{\vartheta}) = \sum_{m=1}^{M} \frac{1}{M} p(x_{k+1} \mid x_k^m; \boldsymbol{\vartheta}), \tag{5}$$

where superscript represents the particle label. By substituting Eq. (5) into Eq. (4), the filtering distribution $p(x_{k+1} \mid \boldsymbol{r}_{1:k+1}, \boldsymbol{\vartheta})$ (up to proportionality) can be approximated by

$$\hat{p}(x_{k+1} \mid \boldsymbol{r}_{1:k+1}, \boldsymbol{\vartheta}) \propto p(\boldsymbol{r}_{k+1} \mid x_{k+1}) \sum_{m=1}^{M} \frac{1}{M} p(x_{k+1} \mid x_k^m; \boldsymbol{\vartheta}). \tag{6}$$

From Eq. (6), the approximation of the filtering distribution $p(x_{k+1} \mid \boldsymbol{r}_{1:k+1}, \boldsymbol{\vartheta})$ can be sampled with importance sampling, where we generate a set of particles $\boldsymbol{x}_{k+1}^{1:M}$ from the predictive distribution $\hat{p}(x_{k+1} \mid \boldsymbol{r}_{1:k}, \boldsymbol{\vartheta})$ with each particle x_{k+1}^m being assigned a weight $w_{k+1}^m \propto p(\boldsymbol{r}_{k+1} \mid x_{k+1}^m)$ (Gordon et al., 1993). We resample M particles with replacement amongst the set of particles $\boldsymbol{x}_{k+1}^{1:M}$ with probabilities proportional to weights $\boldsymbol{w}_{k+1}^{1:M}$. For clarity, we write down the bootstrap particle filter algorithm:

Step 1: Initialise the particles at the sampling time point t_1:

    Step 1a: Draw $x_1^m \sim p(x_1 \mid \boldsymbol{\vartheta})$ for $m = 1, 2, \ldots, M$.

Step 1b: Set $w_1^m = p(\boldsymbol{r}_1 \mid x_1^m)/\sum_{m=1}^{M} p(\boldsymbol{r}_1 \mid x_1^m)$ for $m = 1, 2, \ldots, M$.

Step 1c: Resample M particles with replacement amongst $\boldsymbol{x}_1^{1:M}$ with $\boldsymbol{w}_1^{1:M}$.

Repeat Step 2 for $k = 2, 3, \ldots, K$:

Step 2: Update the particles at the sampling time point t_k:

Step 2a: Draw $x_k^m \sim p(x_k \mid x_{k-1}^m; \boldsymbol{\vartheta})$ for $m = 1, 2, \ldots, M$.

Step 2b: Set $w_k^m = p(\boldsymbol{r}_k \mid x_k^m)/\sum_{m=1}^{M} p(\boldsymbol{r}_k \mid x_k^m)$ for $m = 1, 2, \ldots, M$.

Step 2c: Resample M particles with replacement amongst $\boldsymbol{x}_k^{1:M}$ with $\boldsymbol{w}_k^{1:M}$.

With the procedure for the bootstrap particle filter described above, the smoothing distribu-

tion $p(\boldsymbol{x}_{1:K} \mid \boldsymbol{r}_{1:K}, \boldsymbol{\vartheta})$ can be sampled by uniformly drawing amongst the set of particles $\boldsymbol{x}_{1:K}^{1:M}$,

and the marginal likelihood $p(\boldsymbol{r}_{1:K} \mid \boldsymbol{\vartheta})$ can be estimated by

$$\hat{p}(\boldsymbol{r}_{1:K} \mid \boldsymbol{\vartheta}) = \hat{p}(\boldsymbol{r}_1 \mid \boldsymbol{\vartheta}) \prod_{k=1}^{K-1} \hat{p}(\boldsymbol{r}_{k+1} \mid \boldsymbol{r}_{1:k}, \boldsymbol{\vartheta}) = \prod_{k=1}^{K} \sum_{m=1}^{M} \frac{1}{M} p(\boldsymbol{r}_k \mid x_k^m).$$

We can therefore write down the PMMH algorithm:

Step 1: Initialise the population genetic parameters $\boldsymbol{\vartheta}$ and population mutant allele frequency

trajectory $\boldsymbol{x}_{1:K}$:

Step 1a: Draw $\boldsymbol{\vartheta}^1 \sim p(\boldsymbol{\vartheta})$.

Step 1b: Run a bootstrap particle filter with $\boldsymbol{\vartheta}^1$ to yield $\hat{p}(\boldsymbol{r}_{1:K}, \mid \boldsymbol{\vartheta}^1)$ and $\boldsymbol{x}_{1:K}^1$.

Repeat Step 2 until a sufficient number of samples of the population genetic parameters $\boldsymbol{\vartheta}$ and

population mutant allele frequency trajectory $\boldsymbol{x}_{1:K}$ have been obtained:

Step 2: Update the population genetic parameters $\boldsymbol{\vartheta}$ and population mutant allele frequency

trajectory $\boldsymbol{x}_{1:K}$:

Step 2a: Draw $\boldsymbol{\vartheta}^i \sim q(\boldsymbol{\vartheta} \mid \boldsymbol{\vartheta}^{i-1})$.

Step 2b: Run a bootstrap particle filter with $\boldsymbol{\vartheta}^i$ to yield $\hat{p}(\boldsymbol{r}_{1:K} \mid \boldsymbol{\vartheta}^i)$ and $\boldsymbol{x}_{1:K}^i$.

Step 2c: Accept $\boldsymbol{\vartheta}^i$ and $\boldsymbol{x}_{1:K}^i$ with

$$A = \frac{p(\boldsymbol{\vartheta}^i)}{p(\boldsymbol{\vartheta}^{i-1})} \frac{\hat{p}(\boldsymbol{r}_{1:K} \mid \boldsymbol{\vartheta}^i)}{\hat{p}(\boldsymbol{r}_{1:K} \mid \boldsymbol{\vartheta}^{i-1})} \frac{q(\boldsymbol{\vartheta}^{i-1} \mid \boldsymbol{\vartheta}^i)}{q(\boldsymbol{\vartheta}^i \mid \boldsymbol{\vartheta}^{i-1})},$$

196   otherwise set $\boldsymbol{\vartheta}^i = \boldsymbol{\vartheta}^{i-1}$ and $\boldsymbol{x}_{1:K}^i = \boldsymbol{x}_{1:K}^{i-1}$.

Note that superscript represents the iteration in our procedure described above. In the sequel, we adopt random walk proposals for each component of the population genetic parameters $\boldsymbol{\vartheta}$ unless otherwise specified.

Once enough samples of the population genetic parameters $\boldsymbol{\vartheta}$ and population mutant allele frequency trajectory $\boldsymbol{x}_{1:K}$ have been obtained, we can compute the posterior $p(\boldsymbol{\vartheta} \mid \boldsymbol{r}_{1:K})$ with nonparametric density estimation techniques (see Izenman, 1991, for a review) and obtain the maximum *a posteriori* probability (MAP) estimates for the population genetic parameters $\boldsymbol{\vartheta}$. Alternatively, we can yield the minimum mean square error (MMSE) estimates for the population genetic parameters $\boldsymbol{\vartheta}$ through their posterior means. Similarly, we can take the posterior mean of the samples of the population mutant allele frequency trajectory to be our estimate for the population mutant allele frequency trajectory $\boldsymbol{x}_{1:K}$.

Our method allows the selection coefficient s to be piecewise constant over time. For example, we let the selection coefficient $s(t) = s^-$ if $t < \tau$, otherwise $s(t) = s^+$, where τ is the time of an event that might change selection, *e.g.*, the times of plant and animal domestication. Our procedure can then be directly applied to estimate the selection coefficients s^- and s^+ for any prespecified time τ. The only modification required is to simulate the underlying mutant allele frequency trajectory of the population through the Wright-Fisher diffusion with the selection coefficient s^- for $t < \tau$ and s^+ for $t \geq \tau$, respectively. In this case, our method also provides a procedure to estimate and test the change in the selection coefficient, denoted by $\Delta s = s^+ - s^-$, at time τ by computing the posterior $p(\Delta s \mid \boldsymbol{r}_{1:K})$ from the samples of the selection coefficients s^- and s^+. This is a highly desirable feature in aDNA studies as it enables us to test hypotheses about whether the shift in selection is linked to specific ecological, environmental and cultural drivers. We will evaluate the performance of our procedure applied to this scenario in Section 3.

3. Results

In this section, we evaluate the performance of our approach through extensive simulations and show the applicability of our approach with the data on aDNA sequences from earlier studies of Ludwig et al. (2009), Pruvost et al. (2011) and Wutke et al. (2016), where they sequenced a total of 201 ancient horse samples for eight loci that determine horse coat colouration. In what follows, we base our inference of selection on the MAP estimate and assume only a single event that might change selection.

 3.1. Performance evaluation

We run forward-in-time simulations of the Wright-Fisher model with selection (*e.g.*, Durrett,

2008) and test our approach through simulated datasets of genotype likelihoods to evaluate its

performance. We let the only event that might change selection occur in generation 350, thereby

taking the selection coefficient to be $s(k) = s^-$ for $k < 350$ otherwise $s(k) = s^+$. We uniformly

draw the selection coefficients s^- and s^+ from $[-0.05, 0.05]$ and pick the dominance parameter of

$h = 0.5$ (*i.e.*, additive gene action). To mimic the demographic history of the horse population

estimated by Der Sarkissian et al. (2015), we adopt a bottleneck demographic history, where the

population size $N(k) = 32000$ for $k < 200$, $N(k) = 8000$ for $200 \leq k < 400$, and $N(k) = 16000$

for $k \geq 400$. The initial population mutant allele frequency x_1 is uniformly drawn from $[0.1, 0.9]$.

For clarity, we write down the procedure of generating the dataset of genotype likelihoods:

Repeat Step 1 until $x_K \in (0, 1)$:

Step 1: Generate s^-, s^+ and $\boldsymbol{x}_{1:K}$.

Step 1a: Draw s^-, s^+ from a uniform distribution over $[-0.05, 0.05]$ and x_1 from a uniform

distribution over $[0.1, 0.9]$.

Step 1b: Simulate $\boldsymbol{x}_{1:K}$ with s^-, s^+ and x_1 through the Wright-Fisher model with selection.

Repeat Step 2 for $k = 1, 2, \ldots, K$:

Step 2: Generate $p(\boldsymbol{r}_{n,k} \mid g)$ for $g = 0, 1, 2$ and $n = 1, 2, \ldots, N_k$:

Step 2a: Draw $g_{n,k} \sim p(g \mid x_k)$ in Eq. (3).

Step 2b: Draw $p(\boldsymbol{r}_{n,k} \mid g)$ for $g = 0, 1, 2$ from a Dirichlet distribution of order 3 with $\alpha_0^{n,k}$,

$\alpha_1^{n,k}$ and $\alpha_2^{n,k}$.

We take the parameter $\alpha_{g_{n,k}}^{n,k}$ to be $\phi\psi$ and the other two to be $(1-\phi)\psi/2$, where ϕ and ψ are the

parameters introduced to control the quality of the simulated dataset in terms of the missing rate

and error rate with a common threshold for genotype calling (*i.e.*, 10 times more likely, see Kim

et al., 2011). By following this procedure, we simulate a total of 801 generations starting from

generation 0 and draw a sample of 10 individuals every 40 generations, 210 sampled individuals

in total (*i.e.*, nearly the size of the ancient horse samples), in our simulation studies.

We run a group of simulations to assess how our method performs for different data qualities,

where we vary the parameters $\phi \in \{0.75, 0.85, 0.95\}$ and $\psi \in \{0.5, 1.0\}$, respectively, giving rise

to six possible combinations of the missing rate and error rate (see Table 1 where the mean and standard deviation of missing rates and error rates are computed with 1000 replicates for each combination). For each scenario listed in Table 1, we consider nine possible combinations of the selection coefficients s^- and s^+ (see Table 2), and for each combination we repeatedly run the procedure described above until we get 200 datasets of genotype likelihoods. Thus, in summary, we consider 200 replicates for each of 54 combinations of the data quality and selection scenario.

| Scenario | ϕ | ψ | Missing rate | | Error rate | |
			Mean	Std. Dev.	Mean	Std. Dev.
A	0.75	0.50	0.3095	0.0323	0.1392	0.0278
B	0.75	1.00	0.4542	0.0357	0.0711	0.0229
C	0.85	0.50	0.2037	0.0292	0.0673	0.0192
D	0.85	1.00	0.3029	0.0319	0.0275	0.0140
E	0.95	0.50	0.0753	0.0182	0.0172	0.0094
F	0.95	1.00	0.1108	0.0216	0.0059	0.0057

Table 1: Summary of data qualities across different combinations of the parameters ϕ and ψ.

Scenario	s^-	s^+	Description
1	$s^- < 0$	$s^+ < s^-$	negatively selected mutation following a negative change
2	$s^- < 0$	$s^+ = s^-$	negatively selected mutation (no change)
3	$s^- < 0$	$s^+ > s^-$	negatively selected mutation following a positive change
4	$s^- = 0$	$s^+ < s^-$	effectively neutral mutation following a negative change
5	$s^- = 0$	$s^+ = s^-$	effectively neutral mutation (no change)
6	$s^- = 0$	$s^+ > s^-$	effectively neutral mutation following a positive change
7	$s^- > 0$	$s^+ < s^-$	positively selected mutation following a negative change
8	$s^- > 0$	$s^+ = s^-$	positively selected mutation (no change)
9	$s^- > 0$	$s^+ > s^-$	positively selected mutation following a positive change

Table 2: Summary of selection scenarios across different combinations of the selection coefficients s^- and s^+.

For each replicate, we choose a uniform prior over $[-1, 1]$ for both selection coefficients s^- and s^+, and adopt the reference population size $N_0 = 16000$. We run 10000 PMMH iterations with 1000 particles, where each generation is partitioned into five subintervals in the Euler-Maruyama scheme. We discard the first half of the total PMMH samples as burn-in and thin the remaining by keeping every fifth value. See Figure 2 for our posteriors for the selection coefficients s^- and s^+ produced from a simulated dataset of genotype likelihoods (see Supplementary Information, Table S1), including our estimate for the underlying frequency trajectory of the mutant allele in the population. Evidently, in this example our approach can accurately infer temporally variable selection from genetic time series in genotype likelihood format. The true underlying frequency

trajectory of the mutant allele in the population fluctuates slightly around our estimate and is completely covered in our 95% highest posterior density (HPD) intervals.

Figure 2: Posteriors for the selection coefficients and the underlying frequency trajectory of the mutant allele in the population produced through our method from a dataset of genotype likelihoods generated with the selection coefficients $s^- = 0.0198$ and $s^+ = -0.0081$. (a) Joint posterior for the selection coefficients s^- and s^+. (b) Marginal posteriors for the selection coefficients s^- and s^+. (c) Posterior for the underlying trajectory of the mutant allele frequency. (d) Posterior for the selection change Δs.

3.1.1. Performance in estimating selection coefficients

To assess how our method performs for estimating selection coefficients, we show the boxplot results of our estimates across different data qualities in Figure 3, where the tips of the whiskers are the 2.5%-quantile and the 97.5%-quantile, and the boxes denote the first and third quartiles with the median in the middle. We summarise the bias and root mean square error (RMSE) of our estimates in Supplementary Information, Table S2. As illustrated in Figure 3, our estimates for both selection coefficients are nearly median-unbiased across different data qualities although a slightly large bias can be found when the data quality is poor such as scenario A (about 31.0% missing rate and 13.9% error rate). The bias completely vanishes as the data quality improves

 like scenario F.

Figure 3: Empirical distributions for the bias in the estimation of the selection coefficients across different data qualities and selection scenarios. Data qualities (scenarios A–F) are described in Table 1, and selection scenarios (scenarios 1–9) are described in Table 2. (a)–(i) Boxplots of the bias for scenarios 1–9.

To evaluate how our method performs for different selection coefficients, especially weak selection, we run an additional group of simulations with an example of additive gene action, where we fix the parameters $\phi = 0.85$ and $\psi = 1$. We assume no event that changed selection and vary the selection coefficient $s \in [-0.05, 0.05]$, which is divided into nine subintervals $[-0.05, -0.01)$, $[-0.01, -0.005)$, $[-0.005, -0.001)$, $[-0.001, 0)$, $\{0\}$, $(0, 0.001]$, $(0.001, 0.005]$, $(0.005, 0.01]$ and $(0.01, 0.05]$. For each subinterval, *e.g.*, $[-0.01, -0.005)$, we uniformly choose the selection coeffi-

cient s from $[-0.01, -0.005)$, with which we generate a dataset of genotype likelihoods. Repeat this procedure until we obtain 200 simulated datasets for each subinterval. We show the boxplot results of our estimates across different selection coefficients in Figure 4 and summarise the bias and RMSE of our estimates in Supplementary Information, Table S3.

Boxplot of bias in estimation of selection coefficient

Figure 4: Empirical distributions for the bias in the estimation of the selection coefficient across different ranges of the selection coefficient s with the parameters $\phi = 0.85$ and $\psi = 1$ (*i.e.*, scenario D in Table 1).

We observe from Figure 4 that our estimates are nearly median-unbiased for weak selection, but the bias gets larger with an increase in the strength of selection (*i.e.*, $|s|$). This bias could be caused by ascertainment arising from the procedure that we use to generate simulated datasets. In our simulation studies, only the simulated datasets in which no fixation event has occurred in the underlying population are kept. This setting means that the Wright-Fisher model in our data generation process is equivalent to that conditioned on no fixation event occurred, which does not match that in our approach for estimating the selection coefficient. Such a mismatch could bring about the underestimation of the selection coefficient (*i.e.*, the simulated datasets that are retained correspond to a biased sample of the underlying population mutant allele frequency trajectories that reach loss or fixation more slowly, and therefore the estimates for the strength of selection are more likely to be smaller than their true values), especially for strong selection, since fixation events are more likely (see the histograms of the loss probability of mutations and the histograms of the fixation probability of mutations in Supplementary Information, Figure S1 for each subinterval, where the loss probability and the fixation probability for each simulated

dataset are calculated based on 1000 replicates). We find that the simulated datasets with large

loss probabilities are all generated with the selection coefficient $s \in [-0.05, -0.01)$ whereas those

with large fixation probabilities are all generated with the selection coefficient $s \in (0.01, 0.05]$,

which correspond to the subintervals with significant bias in Figure 4. The bias resulting from

the mismatch can be fully eliminated by conditioning the Wright-Fisher diffusion to survive (He

et al., 2020b).

3.1.2. Performance in testing selection changes

To evaluate how our approach performs for testing selection changes, we produce the receiver

operating characteristic (ROC) curves across different data qualities in Figure 5, where the true-

positive rate (TPR) and false-positive rate (FPR) are computed for each value of the posterior

probability for the selection change that is used as a threshold to classify a locus as experiencing

a shift in selection, and the ROC curve is produced by plotting the TPR against the FPR. We

compute the area under the ROC curve (AUC) to summarise the performance. From Figure 5,

we see that even though data qualities vary across different scenarios, all curves are very concave

and close to the upper left corner of the ROC space with their AUC values varying from 0.89 to

0.94. This suggests that our method has superior performance in testing selection changes even

for the datasets of up to approximately 31.0% missing rate and 13.9% error rate (see scenario

A). Although these ROC curves almost overlap with each other, we can still see that improved

data quality yields better performance (see, *e.g.*, scenario F).

To assess how our approach performs for different selection changes, especially small changes,

we run an additional group of simulations with an example of additive gene action, where we fix

the parameters $\phi = 0.85$ and $\psi = 1$. We vary the selection change $\Delta s \in [-0.05, 0.05]$, which is di-

vided into nine subintervals $[-0.05, -0.01)$, $[-0.01, -0.005)$, $[-0.005, -0.001)$, $[-0.001, 0)$, $\{0\}$,

$(0, 0.001]$, $(0.001, 0.005]$, $(0.005, 0.01]$ and $(0.01, 0.05]$. For each subinterval, *e.g.*, $[-0.01, -0.005)$,

we uniformly draw the selection change Δs from $[-0.01, -0.005)$ and the selection coefficient s^-

from $[\max\{-0.05, -0.05 - \Delta s\}, \min\{0.05, 0.05 - \Delta s\}]$, and therefore have the selection coefficient

$s^+ = s^- + \Delta s$. We generate a dataset of genotype likelihoods with the selection coefficients s^-

and s^+. We repeat this procedure until we obtain 200 simulated datasets for each subinterval.

Similarly, we run the ROC analysis, and the resulting ROC curves for all subintervals are shown

in Figure 6 with their AUC values.

Figure 5: ROC curves for testing a change in selection across different data qualities and selection scenarios. The AUC value for each curve is summarised. Data qualities (scenarios A–F) are described in Table 1, and selection scenarios (scenarios 1–9) are described in Table 2. ROC curves for (a) a negative change in selection and (b) a positive change in selection.

Figure 6: ROC curves for testing a change in selection across different ranges of the selection change Δs with the parameters $\phi = 0.85$ and $\psi = 1$ (*i.e.*, scenario D in Table 1). The AUC value for each curve is summarised. ROC curves for (a) a negative change in selection and (b) a positive change in selection.

By examining the plots of the ROC curves in Figure 6, we find that the performance becomes significantly better with the increase in the degree of the change in the selection coefficient (*i.e.*, $|\Delta s|$) as expected. Even for small changes $|\Delta s| < 0.001$, the AUC value is still larger than 0.70, and for large changes $|\Delta s| > 0.01$, the AUC value is up to 0.98. Our results illustrate that our

method has strong discriminating power of testing selection changes, even though such a change is small.

3.2. Horse coat colouration

We employ our approach to infer selection acting on the *ASIP* and *MC1R* genes associated with base coat colours (black and chestnut) and the *KIT13* and *TRPM1* genes associated with white coat patterns (tobiano and leopard complex) based on the data on aDNA sequences from earlier studies of Ludwig et al. (2009), Pruvost et al. (2011) and Wutke et al. (2016), which were found to be involved to ecological, environmental and cultural shifts (Ludwig et al., 2009, 2015; Wutke et al., 2016). Given that only called genotypes are available in Wutke et al. (2016), we use the following procedure to generate genotype likelihoods for each gene in the same format as those produced by GATK (McKenna et al., 2010): if the genotype is called, we set the genotype likelihood of the called genotype to 1 and those of the other two to 0, and otherwise, all possible (ordered) genotypes are assigned equal genotype likelihoods that are normalised to sum to 1. See genotype likelihoods for each gene in Supplementary Information, Table S4.

Due to the underlying assumption of our approach that mutation occurred before the initial sampling time point, in our following analysis we exclude the samples drawn before the sampling time point that the mutant allele was first found in the sample for each gene. We adopt the horse demographic history estimated by Der Sarkissian et al. (2015) (see Supplementary Information, Figure S2) with the average length of a generation of the horse being eight years and choose the reference population size of $N_0 = 16000$ (*i.e.*, the most recent size of the horse population). We take the dominance parameter h to be the value reported in Wutke et al. (2016) for each gene.

In our PMMH procedure, we run 20000 iterations with a burn-in of 10000 iterations, and the other settings are the same as we adopted in Section 3.1, including those in the Euler-Maruyama approach. The estimates of the selection coefficient and its change for each gene with their 95% HPD intervals are summarised in Supplementary Information, Table S5.

3.2.1. Selection of horse base coat colours

Base coat colours in horses are primarily determined by *ASIP* and *MC1R*, which direct the type of pigment produced, black eumelanin (*ASIP*) or red pheomelanin (*MC1R*) (Corbin et al., 2020). More specifically, *ASIP* on chromosome 22 is associated with the recessive black coat,

and *MC1R* on chromosome 3 is associated with the recessive chestnut coat. Ludwig et al. (2009) found that there was a rapid increase in base coat colour variation during horse domestication (starting from approximately 3500 BC) and provided strong evidence of positive selection acting on *ASIP* and *MC1R*. Fang et al. (2009) suggested that such an increase was directly caused by human preferences and demands. We apply our approach to test their hypothesis that selection acting on *ASIP* and *MC1R* was changed when horse became domesticated and estimate their selection intensities. The resulting posteriors for *ASIP* and *MC1R* are shown in Figures 7 and 8, respectively.

Figure 7: Posteriors for the selection coefficients of the *ASIP* mutation before and from horse domestication (starting from 3500 BC) and the underlying frequency trajectory of the *ASIP* mutation in the population. The samples drawn before 12500 BC are excluded. DOM stands for domestication. (a) Joint posterior for the selection coefficients s^- and s^+. (b) Marginal posteriors for the selection coefficients s^- and s^+. (c) Posterior for the underlying frequency trajectory of the *ASIP* mutation. (d) Posterior for the selection change Δs.

Our estimate of the selection coefficient for the *ASIP* mutation is 0.0018 with 95% HPD interval $[-0.0012, 0.0066]$ before domestication and 0.0003 with 95% HPD interval $[-0.0022, 0.0031]$ after horses were domesticated. The 95% HPD interval contains 0 for the selection coefficient s^-,

Figure 8: Posteriors for the selection coefficients of the *MC1R* mutation before and from horse domestication (starting from 3500 BC) and the underlying frequency trajectory of the *MC1R* mutation in the population. The samples drawn before 4300 BC are excluded. DOM stands for domestication. (a) Joint posterior for the selection coefficients s^- and s^+. (b) Marginal posteriors for the selection coefficients s^- and s^+. (c) Posterior for the underlying frequency trajectory of the *MC1R* mutation. (d) Posterior for the selection change Δs.

but there is still some evidence showing that the *ASIP* mutation was most probably favoured by selection before domestication since the posterior probability for positive selection is 0.904. The posterior for the selection coefficient s^+ is approximately symmetric about 0, which indicates that the *ASIP* mutation was effectively neutral after horse domestication started. Our estimate of the change in the selection acting on the *ASIP* mutation when horses became domesticated is -0.0021 with 95% HPD interval $[-0.0071, 0.0031]$, and the posterior probability for such a negative change is 0.779. Our estimate for the underlying *ASIP* mutation frequency trajectory illustrates that the *ASIP* mutation frequency rises substantially in the pre-domestication period and then keeps approximately constant in the post-domestication period.

Our estimate of the selection coefficient for the *MC1R* mutation is -0.0300 with 95% HPD interval $[-0.0928, 0.0997]$ before horses became domesticated and 0.0116 with 95% HPD inter-

val $[0.0079, 0.0174]$ after domestication started. Our estimates reveal that the *MC1R* mutation was effectively neutral or selectively deleterious in the pre-domestication period (with posterior probability for negative selection being 0.670) but became positively selected after horse domestication started (with posterior probability for positive selection being 1.000). Our estimate of the change in the selection acting on the *MC1R* mutation from a pre- to a post-domestication period is 0.0439 with 95% HPD interval $[-0.0891, 0.1058]$, and the posterior probability for such a positive change is 0.748. We observe a slow decline in the *MC1R* mutation frequency before domestication (even though the evidence of negative selection before domestication is weak) and then a significant increase after horse domestication started in our estimate for the underlying *MC1R* mutation frequency trajectory.

3.2.2. Selection of horse white coat patterns

Tobiano is a white spotting pattern in horses characterised by patches of white that typically cross the topline somewhere between the ears and tail. It is inherited as an autosomal dominant trait that was reported in Brooks et al. (2007) to be associated with a locus in intron 13 of the *KIT* gene on chromosome 3. Wutke et al. (2016) observed that spotted coats in early domestic horses revealed a remarkable increase, but medieval horses carried significantly fewer alleles for these traits, which could result from the shift in human preferences and demands. We apply our method to test their hypothesis that selection acting on *KIT13* was changed when the medieval period began (in around AD 400) and estimate their selection intensities. We show the resulting posteriors for *KIT13* in Figure 9.

Our estimate of the selection coefficient for the *KIT13* mutation is 0.0039 with 95% HPD interval $[-0.0010, 0.0103]$ before the medieval period, which shows that the *KIT13* mutation was positively selected before the Middle Ages (*i.e.*, the posterior probability for positive selection is 0.935). Our estimate of the selection coefficient for the *KIT13* mutation is -0.0284 with 95% HPD interval $[-0.0627, 0.0015]$ during the medieval period, which demonstrates that the *KIT13* mutation became selectively deleterious during the Middle Ages (*i.e.*, the posterior probability for negative selection is 0.969). Our estimate of the change in the selection acting on the *KIT13* mutation when the Middle Ages started is -0.0326 with 95% HPD interval $[-0.0700, 0.0008]$, and the posterior probability for such a negative change is 0.969. We observe from our estimate for the underlying *KIT13* mutation frequency trajectory that the *KIT13* mutation experienced

Figure 9: Posteriors for the selection coefficients of the *KIT13* mutation before and from the Middle Ages (starting from AD 400) and the underlying frequency trajectory of the *KIT13* mutation in the population. The samples drawn before 3645 BC are excluded. EMA stands for Early Middle Ages. (a) Joint posterior for the selection coefficients s^- and s^+. (b) Marginal posteriors for the selection coefficients s^- and s^+. (c) Posterior for the underlying frequency trajectory of the *KIT13* mutation. (d) Posterior for the selection change Δs.

a gradual increase after horse were domesticated and then a marked decrease during the Middle Ages.

Leopard complex is a group of white spotting patterns in horses characterised by a variable amount of white in the coat with or without pigmented leopard spots, which is inherited by the incompletely dominant *TRPM1* gene residing on chromosome 1 (Terry et al., 2004). The first genetic evidence of the leopard complex coat could date back to the Pleistocene (Ludwig et al., 2015). Ludwig et al. (2015) found shifts in the selection pressure for the leopard complex coat in domestic horses but did not explore whether *TRPM1* undergone a shift in selection from a pre- to a post-domestication period. We employ our approach to test the hypothesis that selection acting on *TRPM1* was changed when horses became domesticated and estimate their selection intensities. The resulting posteriors for *TRPM1* are illustrated in Figure 10.

Figure 10: Posteriors for the selection coefficients of the *TRPM1* mutation before and from horse domestication (starting from 3500 BC) and the underlying frequency trajectory of the *TRPM1* mutation in the population. The samples drawn before 14500 BC are excluded. DOM stands for domestication. (a) Joint posterior for the selection coefficients s^- and s^+. (b) Marginal posteriors for the selection coefficients s^- and s^+. (c) Posterior for the underlying frequency trajectory of the *TRPM1* mutation. (d) Posterior for the selection change Δs.

Our estimate of the selection coefficient for the *TRPM1* mutation is -0.0005 with 95% HPD interval $[-0.0042, 0.0029]$ before horses were domesticated and -0.0078 with 95% HPD interval $[-0.0139, -0.0005]$ after domestication started. Our estimates provide little evidence of negative selection in the pre-domestication period (with posterior probability for negative selection being 0.617) but strong evidence of negative selection in the post-domestication period (with posterior probability for negative selection being 0.980). Our estimate of the change in the selection acting on the *TRPM1* mutation when horses became domesticated is -0.0066 with 95% HPD interval $[-0.0142, 0.0025]$. The 95% HPD interval for the change in the selection coefficient contains 0, which however still provides enough evidence to support that a negative change took place in selection when horses became domesticated (*i.e.*, the posterior probability for such a negative change is 0.942). Our estimate for the underlying trajectory of the *TRPM1* mutation frequency

₄₄₄ displays a slow decrease in the *TRPM1* mutation frequency during the pre-domestication period
₄₄₅ with a significant decline after horse domestication started.

4. Discussion

₄₄₇ In this work, we introduced a novel Bayesian approach for estimating temporally changing
₄₄₈ selection intensity from aDNA samples. To our knowledge, most earlier methods ignored sample
₄₄₉ uncertainties resulting from the damage and fragmentation of aDNA molecules, which however
₄₅₀ are the main characteristics of aDNA. Our approach to circumvent this problem is that we based
₄₅₁ our inference of selection on genotype likelihoods rather than called genotypes, which facilitates
₄₅₂ the incorporation of genotype uncertainties. We introduced a novel two-layer HMM framework,
₄₅₃ where the top hidden layer models the underlying frequency trajectory of the mutant allele in
₄₅₄ the population, the intermediate hidden layer models the unobserved genotype of the individual
₄₅₅ in the sample, and the bottom observed layer denotes the data on aDNA sequences. By working
₄₅₆ with the PMMH algorithm, where the marginal likelihood was approximated through particle
₄₅₇ filtering, our method enables us to reconstruct the underlying mutant allele frequency trajectory
₄₅₈ of the population. Furthermore, our procedure provides the flexibility of modelling time-varying
₄₅₉ demographic histories.

₄₆₀ The performance of our method was evaluated through extensive simulations, which showed
₄₆₁ that our procedure could produce accurate selection inference from aDNA data across different
₄₆₂ evolutionary scenarios even though samples were sparsely distributed in time with small sizes
₄₆₃ and in poor quality. Note that in our simulations studies, we only tested our method in the case
₄₆₄ of a bottleneck demographic history and additive gene action, but in principle, the conclusions
₄₆₅ we draw here hold for other demographic histories and gene actions. Demographic histories have
₄₆₆ been demonstrated to have little effect on the inference of selection from time series genetic data
₄₆₇ (Jewett et al., 2016), and additional simulation studies for the cases of the mutant allele being
₄₆₈ recessive ($h = 0$) and dominant ($h = 1$) can be found in Supplementary Information, Figures S3
₄₆₉ and S4, respectively, as well as in Table S6.

₄₇₀ We illustrated the utility of our approach with an application to ancient horse samples from
₄₇₁ previous studies of Ludwig et al. (2009), Pruvost et al. (2011) and Wutke et al. (2016), which
₄₇₂ were genotyped at the loci (*e.g.*, *ASIP*, *MC1R*, *KIT13* and *TRPM1*) for horse coat colouration.
₄₇₃ Our findings are compatible with earlier studies that the coat colour variation in the horse is a

domestic trait that was subject to early selection by humans (Hunter, 2018), *e.g.*, *ASIP*, *MC1R* and *TRPM1*, and human preferences have changed greatly over time and across cultures (Wutke et al., 2016), *e.g.*, *KIT13*.

Our results for the base coat colour are consistent with previous studies that the shift in horse coat colour variation in the early stage of domestication could be caused by relaxed selection for camouflage alleles (Hunter, 2018). More specifically, the *ASIP* mutation was positively selected in the pre-domestication period, but the *MC1R* mutation was not. From Sandoval-Castellanos et al. (2017), forest cover was growing as a result of global warming during the Late Pleistocene, which pushed horses into the forest full of predators. Dark-coloured coats could help horses avoid predators through better camouflage, therefore improving their chances of survival. After horses were domesticated, the *ASIP* mutation was no longer selectively advantageous, but the *MC1R* mutation became favoured by selection. The shift in the horse coat colour preference from dark to light could be probably explained by the fact that light-coloured horses were no longer needed to protect against predation due to domestication, and light-coloured coats could facilitate horse husbandry since it was easier to keep track of the horses that were not camouflaged (Fang et al., 2009).

Our results for the tobiano coat pattern demonstrate that the *KIT13* mutation was favoured by selection from domestication till the Middle Ages and then became negatively selected, which confirm the findings of Wutke et al. (2016). Such a negative change in selection of the tobiano coat could result from pleiotropic disadvantages, a lower religious prestige, a reduced need to separate domestic horses from their wild counterparts or novel developments in weaponry during the medieval period (see Wutke et al., 2016, and references therein).

Our results for the leopard complex coat pattern illustrate that the *TRPM1* mutation was negatively selected from the Late Pleistocene onwards. Our evidence of negative selection acting on the *TRPM1* mutation in the pre-domestication period is not strong enough, but we can still see a slow drop in the *TRPM1* mutation frequency over time. The *TRPM1* mutation is the most common cause of congenital stationary night blindness (CSNB) (Bellone et al., 2013), which could reduce the chance to survive in the wild since vision is key for communication, localisation, orientation, avoiding predators and looking for food (Murphy et al., 2009). The weak intensity of negative selection could be explained as resulting in part from that only horses homozygous for

the leopard complex coat pattern are influenced by CSNB, which however remarkably increased when horses were domesticated. In the post-domestication period, horses were harnessed mainly for power and transportation, *e.g.*, they were used to pull wheeled vehicles, chariots, carts and wagons in the early stage of domestication and later used in war, in hunting and as a means of transport, which all strongly rely on the ability to see. Moreover, night-blind horses are nervous and timid in human care, and difficult to handle at dusk and darkness (Rebhun et al., 1984).

Compared to existing methods (*e.g.*, Bollback et al., 2008; Malaspinas et al., 2012; Steinrücken et al., 2014; Schraiber et al., 2016; Ferrer-Admetlla et al., 2016; He et al., 2020a,b), our Bayesian procedure enables the selection coefficient to vary in time (*i.e.*, piecewise constant) although the event that might change selection is required to be prespecified. However, this is still important in aDNA studies as adaptation in natural populations typically involve adaptation to ecological, environmental and cultural shifts. We run our procedure with the same settings as we adopted in Section 3.2 on the ancient horse samples presented in Supplementary Information, Table S4, except that the selection coefficient is fixed through time (see Supplementary Information, Figure S5 for the resulting posteriors and Supplementary Information, Table S7 for the estimates of the selection coefficients with their 95% HPD intervals). We find for example that the *KIT13* mutation was effectively neutral during the post-domestication period, which contradicts the archaeological evidence and historical records that spotted horses were subject to early selection by humans, but the preference changed during the medieval period (see Wutke et al., 2016, and references therein). Compared to the results presented in Figures 7–10, we see the necessity of modelling temporally variable selection in aDNA. Our approach lends itself to being extended to the analysis of multiple prespecified events that might change selection (see Section 2.2). To guarantee computational efficiency, a feasible solution is to adopt an adaptive strategy that can automatically tune the selection coefficients during a run (see Luengo et al., 2020, for a review). A potential direction for future research is the inference of selection and its strength and timing of changes from aDNA data (Shim et al., 2016; Mathieson, 2020).

In aDNA studies, the samples with missing genotypes are usually filtered, and the remaining samples are grouped into a small number of sampling time points, *e.g.*, ancient horse samples were grouped into six sampling time points in Ludwig et al. (2009) or nine sampling time points in Wutke et al. (2016). Grouping has been shown to largely bias the inference of selection from

time series data in He et al. (2020b). Our method provides an alternative that can address the issue caused by the procedure of sample filtering and grouping (see Supplementary Information, File S3 for additional simulation studies, where we compare our results produced with genotype likelihoods to those based on called genotypes). Our simulation studies show that the commonly used procedure of processing aDNA data for the inference of selection could significantly alter the result, in particular for poor data quality, and suggest that the approach based on genotype likelihoods could be a more promising alternative for future aDNA studies.

One fundamental limitation of our method applied for the inference of selection from aDNA data is that it assumes that all samples have drawn after the mutant allele was created. However, allele age is not always available, and as a result, in our analysis we have to remove the samples drawn before the time that the mutant allele was first found in the sample, which might alter the result of the inference of selection. To address this challenge, we can extend our approach to jointly estimate the allele age as in Malaspinas et al. (2012), Schraiber et al. (2016) and He et al. (2020b), and the foreseeable challenge is how to resolve particle degeneracy and impoverishment issues in our PMMH-based procedure that result from low-frequency mutant alleles at the early stage facing a higher probability of being lost. Furthermore, our approach assumes that the gene of our interest is independent of others, but in practice this assumption can be easily violated once there exist interactions between genes like epistasis and linkage, *e.g.*, *MC1R* is epistatic to *MC1R* (Rieder et al., 2001), and *KIT13* is tightly linked to *KIT16* (Dumont & Payseur, 2008), which might bias our result of the inference of selection. To extend our procedure to the scenario of multiple genes with epistasis and/or linkage, we need to find a good approximation of the Wright-Fisher model that characterises multiple genes evolving subject to selection with epistasis and linkage through time, which becomes challenging with an increase in the number of genes. We will discuss how to extend our approach to the scenario of two genes with epistasis and/or linkage in our upcoming work, and the extension for the scenario of multiple interacting genes (≥ 3) will be the topic of future investigation.

Acknowledgements

This work was carried out using the computational facilities of the Advanced Computing Research Centre, University of Bristol - `http://www.bristol.ac.uk/acrc/`.

References

Andrieu, C., Doucet, A., & Holenstein, R. (2010). Particle Markov chain Monte Carlo methods. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)*, *72*, 269–342.

Bellone, R. R., Holl, H., Setaluri, V., Devi, S., Maddodi, N. et al. (2013). Evidence for a retroviral insertion in *TRPM1* as the cause of congenital stationary night blindness and leopard complex spotting in the horse. *PLoS One*, *8*, e78280.

Bollback, J. P., York, T. L., & Nielsen, R. (2008). Estimation of $2N_e s$ from temporal allele frequency data. *Genetics*, *179*, 497–502.

Bosshard, L., Dupanloup, I., Tenaillon, O., Bruggmann, R., Ackermann, M. et al. (2017). Accumulation of deleterious mutations during bacterial range expansions. *Genetics*, *207*, 669–684.

Brooks, S. A., Lear, T. L., Adelson, D. L., & Bailey, E. (2007). A chromosome inversion near the *KIT* gene and the Tobiano spotting pattern in horses. *Cytogenetic and Genome Research*, *119*, 225–230.

Corbin, L. J., Pope, J., Sanson, J., Antczak, D. F., Miller, D. et al. (2020). An independent locus upstream of *ASIP* controls variation in the shade of the bay coat colour in horses. *Genes*, *11*, 606.

Darwin, C. (1859). *On the Origin of Species by Means of Natural Selection, or the Preservation of Favoured Races in the Struggle for Life*. London: John Murray.

Dehasque, M., Ávila-Arcos, M. C., Díez-del Molino, D., Fumagalli, M., Guschanski, K. et al. (2020). Inference of natural selection from ancient DNA. *Evolution Letters*, *4*, 94–108.

DePristo, M. A., Banks, E., Poplin, R., Garimella, K. V., Maguire, J. R. et al. (2011). A framework for variation discovery and genotyping using next-generation DNA sequencing data. *Nature Genetics*, *43*, 491–498.

Der Sarkissian, C., Ermini, L., Schubert, M., Yang, M. A., Librado, P. et al. (2015). Evolutionary genomics and conservation of the endangered Przewalski's horse. *Current Biology*, *25*, 2577–2583.

Dumont, B. L., & Payseur, B. A. (2008). Evolution of the genomic rate of recombination in mammals. *Evolution*, *62*, 276–294.

Durrett, R. (2008). *Probability Models for DNA Sequence Evolution*. New York: Springer-Verlag.

Fages, A., Hanghøj, K., Khan, N., Gaunitz, C., Seguin-Orlando, A. et al. (2019). Tracking five millennia of horse management with extensive ancient genome time series. *Cell*, *177*, 1419–1435.

Fang, M., Larson, G., Soares R, H., Li, N., & Andersson, L. (2009). Contrasting mode of evolution at a coat color locus in wild and domestic pigs. *PLoS Genetics*, *5*, e1000341.

Feder, A. F., Kryazhimskiy, S., & Plotkin, J. B. (2014). Identifying signatures of selection in genetic time series. *Genetics*, *196*, 509–522.

Ferrer-Admetlla, A., Leuenberger, C., Jensen, J. D., & Wegmann, D. (2016). An approximate Markov model for the Wright-Fisher diffusion and its application to time series data. *Genetics*, *203*, 831–846.

Fisher, R. A. (1922). On the dominance ratio. *Proceedings of the Royal Society of Edinburgh*, *42*, 321–341.

Good, B. H., McDonald, M. J., Barrick, J. E., Lenski, R. E., & Desai, M. M. (2017). The dynamics of molecular evolution over 60,000 generations. *Nature*, *551*, 45–50.

Gordon, N. J., Salmond, D. J., & Smith, A. F. M. (1993). Novel approach to nonlinear/non-Gaussian Bayesian state estimation. *IEE Proceedings F (Radar and Signal Processing)*, *140*, 107–113.

He, Z., Beaumont, M. A., & Yu, F. (2017). Effects of the ordering of natural selection and population regulation mechanisms on Wright-Fisher models. *G3: Genes, Genomes, Genetics*, *7*, 2095–2106.

He, Z., Dai, X., Beaumont, M. A., & Yu, F. (2020a). Detecting and quantifying natural selection at two linked loci from time series data of allele frequencies with forward-in-time simulations. *Genetics*, *216*, 521–541.

He, Z., Dai, X., Beaumont, M. A., & Yu, F. (2020b). Estimation of natural selection and allele age from time series allele frequency data using a novel likelihood-based approach. *Genetics*, *216*, 463–480.

He, Z., Lyu, W., Beaumont, M. A., & Yu, F. (2021). Moment-based approximations for the Wright-Fisher model of population dynamics under natural selection at two linked loci. *bioRxiv*, (p. 424882).

Hunter, P. (2018). The genetics of domestication: Research into the domestication of livestock and companion animals sheds light both on their "evolution" and human history. *EMBO Reports*, *19*, 201–205.

Izenman, A. J. (1991). Recent developments in nonparametric density estimation. *Journal of the American Statistical Association*, *86*, 205–224.

Jewett, E. M., Steinrücken, M., & Song, Y. S. (2016). The effects of population size histories on estimates of selection coefficients from time-series genetic data. *Molecular Biology and Evolution*, *33*, 3002–3027.

Kim, S. Y., Lohmueller, K. E., Albrechtsen, A., Li, Y., Korneliussen, T. et al. (2011). Estimation of allele frequency and association mapping using next-generation sequencing data. *BMC Bioinformatics*, *12*, 1–16.

Lacerda, M., & Seoighe, C. (2014). Population genetics inference for longitudinally-sampled mutants under strong selection. *Genetics*, *198*, 1237–1250.

Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J. et al. (2009a). The sequence alignment/map format and SAMtools. *Bioinformatics*, *25*, 2078–2079.

Li, R., Li, Y., Fang, X., Yang, H., Wang, J. et al. (2009b). SNP detection for massively parallel whole-genome resequencing. *Genome Research*, *19*, 1124–1132.

Librado, P., Gamba, C., Gaunitz, C., Der Sarkissian, C., Pruvost, M. et al. (2017). Ancient genomic changes associated with domestication of the horse. *Science*, *356*, 442–445.

Ludwig, A., Pruvost, M., Reissmann, M., Benecke, N., Brockmann, G. A. et al. (2009). Coat color variation at the beginning of horse domestication. *Science*, *324*, 485–485.

Ludwig, A., Reissmann, M., Benecke, N., Bellone, R., Sandoval-Castellanos, E. et al. (2015). Twenty-five thousand years of fluctuating selection on leopard complex spotting and congenital night blindness in horses. *Philosophical Transactions of the Royal Society B: Biological Sciences*, *370*, 20130386.

Luengo, D., Martino, L., Bugallo, M., Elvira, V., & Särkkä, S. (2020). A survey of Monte Carlo methods for parameter estimation. *EURASIP Journal on Advances in Signal Processing*, *2020*, 1–62.

Lyu, W., Dai, X., Beaumont, M. A., Yu, F., & He, Z. (2022). Inferring the timing and strength of natural selection and gene migration in the evolution of chicken from ancient DNA data. *Molecular Ecology Resources*, *22*, 1362–1379.

Malaspinas, A.-S. (2016). Methods to characterize selective sweeps using time serial samples: an ancient DNA perspective. *Molecular Ecology*, *25*, 24–41.

Malaspinas, A.-S., Malaspinas, O., Evans, S. N., & Slatkin, M. (2012). Estimating allele age and selection coefficient from time-serial data. *Genetics*, *192*, 599–607.

Mathieson, I. (2020). Estimating time-varying selection coefficients from time series data of allele frequencies. *bioRxiv*, (p. 387761).

Mathieson, I., Lazaridis, I., Rohland, N., Mallick, S., Patterson, N. et al. (2015). Genome-wide patterns of selection in 230 ancient Eurasians. *Nature*, *528*, 499–503.

McKenna, A., Hanna, M., Banks, E., Sivachenko, A., Cibulskis, K. et al. (2010). The genome analysis toolkit: a mapreduce framework for analyzing next-generation dna sequencing data. *Genome Research*, *20*, 1297–1303.

Murphy, J., Hall, C., & Arkins, S. (2009). What horses and humans see: a comparative review. *International Journal of Zoology*, *2009*, 721798.

Paris, C., Servin, B., & Boitard, S. (2019). Inference of selection from genetic time series using various parametric approximations to the Wright-Fisher model. *G3: Genes, Genomes, Genetics*, *9*, 4073–4086.

Pruvost, M., Bellone, R., Benecke, N., Sandoval-Castellanos, E., Cieslak, M. et al. (2011). Genotypes of predomestic horses match phenotypes painted in Paleolithic works of cave art. *Proceedings of the National Academy of Sciences*, *108*, 18626–18630.

Rebhun, W. C., Loew, E. R., Riis, R. C., & Laratta, L. J. (1984). Clinical manifestations of night blindness in the Appaloosa horse. *The Compendium on Continuing Education for the Practicing Veterinarian*, *6*, S103–S106.

Rieder, S., Taourit, S., Mariat, D., Langlois, B., & Guérin, G. (2001). Mutations in the agouti (*ASIP*), the extension (*MC1R*), and the brown (*TYRP1*) loci and their association to coat color phenotypes in horses (*Equus caballus*). *Mammalian Genome*, *12*, 450–455.

Sandoval-Castellanos, E., Wutke, S., Gonzalez-Salazar, C., & Ludwig, A. (2017). Coat colour adaptation of post-glacial horses to increasing forest vegetation. *Nature Ecology & Evolution*, *1*, 1816–1819.

Schraiber, J. G., Evans, S. N., & Slatkin, M. (2016). Bayesian inference of natural selection from allele frequency time series. *Genetics*, *203*, 493–511.

Shim, H., Laurent, S., Matuszewski, S., Foll, M., & Jensen, J. D. (2016). Detecting and quantifying changing selection intensities from time-sampled polymorphism data. *G3: Genes, Genomes, Genetics*, *6*, 893–904.

Steinrücken, M., Bhaskar, A., & Song, Y. S. (2014). A novel spectral method for inferring general diploid selection from time series genetic data. *The Annals of Applied Statistics*, *8*, 2203–2222.

Terhorst, J., Schlötterer, C., & Song, Y. S. (2015). Multi-locus analysis of genomic time series data from experimental evolution. *PLoS Genetics*, *11*, e1005069.

Terry, R. B., Archer, S., Brooks, S., Bernoco, D., & Bailey, E. (2004). Assignment of the appaloosa coat colour gene (*LP*) to equine chromosome 1. *Animal Genetics*, *35*, 134–137.

Turner, T. L., & Miller, P. M. (2012). Investigating natural variation in *Drosophila* courtship song by the evolve and resequence approach. *Genetics*, *191*, 633–642.

Williamson, E. G., & Slatkin, M. (1999). Using maximum likelihood to estimate population size from temporal changes in allele frequencies. *Genetics*, *152*, 755–761.

Wright, S. (1931). Evolution in Mendelian populations. *Genetics*, *16*, 97–159.

Wutke, S., Benecke, N., Sandoval-Castellanos, E., Döhle, H.-J., Friederich, S. et al. (2016). Spotted phenotypes in horses lost attractiveness in the Middle Ages. *Scientific Reports*, *6*, 38548.

Wutke, S., Sandoval-Castellanos, E., Benecke, N., Döhle, H., Friederich, S. et al. (2018). Decline of genetic diversity in ancient domestic stallions in Europe. *Science Advances*, *4*, eaap9691.

Ye, K., Gao, F., Wang, D., Bar-Yosef, O., & Keinan, A. (2017). Dietary adaptation of *FADS* genes in Europe varied across time and geography. *Nature Ecology & Evolution*, *1*, 1–11.

Data Accessibility Statement

The authors state that all data necessary for confirming the conclusions of the present work are represented fully within the article. Source code implementing the approach described in this work is available at `https://github.com/zhangyi-he/WFM-1L-DiffusApprox-PMMH/`.

Author Contributions

Z.H. designed the project and developed the method; Z.H., X.D. and W.L. implemented the method; X.D. and W.L. analysed the data under the supervision of Z.H., M.B. and F.Y.; Z.H. and X.D. wrote the manuscript; W.L., M.B. and F.Y. reviewed the manuscript.