# Harmful self-pollination drives gynodioecy in European chestnut, a self-incompatible tree

Clément Larue[a,b,*], Rémy J. Petit[a*]

[a]Univ. Bordeaux, INRAE, BIOGECO, 33610 Cestas, France
[b]INVENIO, Maison Jeannette, 24140 Douville, France

**ORCID**

Clément Larue : 0000-0003-2931-345X

Rémy J Petit : 0000-0002-4809-1453

**\*Corresponding authors**

Clément Larue, INRAE, 69 route d'Arcachon, 33610 Cestas, France.

Email: clement.larue@inrae.fr

Rémy Petit, INRAE, 69 route d'Arcachon, Cestas, France
Email: remy.petit@inrae.fr

Number of words: 5870 (Introduction: 1555, Mat & Meth: 1951, Results: 609, Discussion: 1755)

Number of tables: 5

Number of figures: 3

Supporting information: 5

## SUMMARY

- Gynodioecy is a rare sexual system in which two genders, cosexuals and females, coexist. It provides the opportunity to compare individuals having both sexual functions with individuals lacking the male function, an ideal situation to understand how sexes interact within individual plants.

- We report gynodioecy in the European chestnut, an outcrossing self-incompatible Fagaceae tree species. This finding was unexpected because gynodioecy is often interpreted as an outbreeding mechanism. To understand how female chestnuts compensate for their lack of siring capacity, we compared key female fitness components between genders and performed emasculation experiments.

- Genders have similar basal area and number of flowers but different fruit set. Following the removal of nectar-producing catkins on branches or entire trees, fruit set increased in cosexual trees but decreased in female trees.

- These results show that self-pollination impairs fruit set in cosexual trees, a likely effect of self-pollen interference caused by late-acting self-incompatibility and by early inbreeding depression. Female trees escape from self-pollen interference but continue to attract pollinators thanks to their sterile but rewarding male catkins, resulting in a much higher fruit set than cosexuals. This demonstrates that even entirely outcrossed plants can benefit from the cessation of self-pollination.

**Key-words:** *Castanea*, emasculation experiment, female advantage, gynodioecy, late-acting self-incompatibility, ovule discounting, sexual interference, selfing avoidance

## INTRODUCTION

In angiosperms, most species have retained the ancestral hermaphrodite floral organization (Yampolsky, 1922). Starting with this sexual system well adapted for animal pollination (Sauquet *et al.*, 2017), flowering plants have further evolved, resulting in an amazing variety of reproductive systems at the flower, inflorescence or whole plant level. For instance, in gynodioecy, a rare but widespread dimorphic sexual system, both cosexual (typically hermaphrodite) and unisexual (female) individuals are present. The evolutionary mechanisms leading to such gender

polymorphism in plants have fascinated researchers ever since Darwin (1877). Studying these

mechanisms should also help understand why the vast majority of flowering plants are cosexual

(Bawa, 1984). To explore the consequences of the coexistence of both sexual functions within

individuals, gynodioecious plants are better models than dioecious plants: a comparison between

their two genders can help identify negative and positive interactions between sexual functions

taking place within cosexual individuals (Webb, 1999). In gynodioecious species, to persist,

females must compensate for their lack of male fitness (Lewis, 1941), which requires rather

stringent conditions (Charlesworth & Charlesworth, 1978). This compensation, or 'female

advantage', is the reproductive advantage of females over cosexuals due to increased life-long

female fecundity through higher seed quantity, higher seed quality, or both. In recently established

male-sterile mutants that lack the refined secondary characters of long-established specialist

females, female advantage should be mostly attributable to the release from negative sexual

interactions (Charlesworth & Charlesworth, 1978; Lloyd, 1982).

Mechanisms of female advantage can be classified in two categories, genetic and ecological

(Givnish, 1982). The most cited genetic mechanism of female advantage is the outbreeding

advantage, attributed to the escape from inbreeding depression in selfed offspring (Darwin, 1876;

Mather, 1940; Baker, 1959; Lloyd, 1975). Other genetic mechanisms that can affect cosexuals but

not females include pollen tube competition between selfed and outcrossed pollen (Shykoff, 1992)

and self-pollen recognition and discrimination in the styles. The most cited ecological mechanism

of female advantage is reallocation of resources to the female function from an abandoned male

function (Darwin, 1877). Other ecological mechanisms of female advantage exist, such as reduction

of antagonistic biotic interactions (reviewed in Ashman, 2002). In the 1980s, a heated debate took

place on the nature of mechanisms driving separate sexes (Freeman *et al.*, 1997), with supporters of

the central importance of outbreeding advantage (Thomson & Barrett, 1981; Lloyd, 1982) being

challenged by opponents emphasizing mostly ecological mechanisms (Willson, 1979; Bawa, 1980,

1982; Givnish, 1982). Forty years afterwards, we still have only a limited understanding of the

relative importance of the different mechanisms (genetic or ecological) of female advantage (Dufaÿ

& Billard, 2012; Pannell, 2018).

There are two main reasons why selection for outcrossing was considered by some to be the most

important driver of the evolution of separate sexes. First, its principle is straightforward (Mather,

1940), and second, it has considerable predictive power. In particular, it predicts that, if

hermaphrodites are partially self-fertilizing and produce lower-quality offspring because of

inbreeding depression, females, but not males, will be selected. This prediction is supported by the

much greater prevalence of gynodioecy over androdioecy (Lloyd, 1975; Charlesworth & Charlesworth, 1978). However, a more recent review of the origin of gender differences in gynodioecious species has shown that reduced selfing is not always the cause of female advantage (Dufaÿ & Billard, 2012). Moreover, other mechanisms than unisexuality can promote outcrossing, such as self-incompatibility (Willson, 1979). Clearly, the question of the relative importance of genetic and ecological mechanisms in the evolution of separate sexes remains open.

In females, outcrossing is enforced. However, it would be erroneous to consider that selfing avoidance is the only female advantage derived from pollen emission breakdown. Reduced seed set after self-pollination is widespread among hermaphrodite flowering plants (Burbidge & James, 1991). The cause is not merely postzygotic. In plant species with late-acting (ovarian) self-incompatibility (Seavey & Bawa, 1986), ovules are disabled by self-pollen tubes (Charlesworth, 1985; Sage *et al.*, 1994; Gibbs, 2014; Johnson *et al.*, 2019). Barrett *et al.* (1996) introduced the concept of ovule discounting to describe '*the situation where ovules are excluded from cross-fertilization because they are rendered non-functional by self-pollen tubes as a consequence of prior self-pollination*'. This represents a special case of male interference with the female function (Barrett, 2002; Duffy & Johnson, 2014). Self-pollen interference with ovules closely parallels inbreeding depression acting on embryos, seeds and seedlings: both are genetic mechanisms that ultimately limit inbreeding. In principle, if strong enough, self-pollen interference and ovule discounting could drive the evolution of separate sexes, alone or in combination with selfing and inbreeding depression. Hence, a more encompassing view would be to consider 'self-pollination avoidance' instead of 'selfing avoidance' as an explanation for any female advantage resulting from mating system change in females.

Regardless of the mechanisms involved, some circumstances can affect the success of the establishment of females in cosexual populations. First, the transmission mode of male-sterility can greatly affect female establishment. Specifically, in case of maternal inheritance of male sterility, male-sterility mutations will benefit from exclusive transmission via ovules (e.g. Lewis, 1941; Lloyd, 1975; Gouyon & Couvet, 1987). Second, the broader ecological context is also highly relevant. In particular, the establishment of females is easier when pollen is not limiting (e.g. Lloyd, 1974; Dornier & Dufaÿ, 2013; Lahiani *et al.*, 2015; Stone & Olson, 2018). Third, not all sexual interactions are negative. Pollinators often prefer hermaphrodite over female flowers, due to male-biased pollinator attractiveness (Darwin, 1877; Lloyd, 1975; Wise et al., 2011; van Etten and Chang, 2014). Such positive sexual interactions can inhibit the evolution of females in cosexual populations.

Gynodioecy evolves when male-sterile mutants are able to establish. For this, the mutants must

benefit not only from favourable circumstances but also from an immediate and tangible advantage

(Webb, 1999). A simple but powerful method to identify mechanisms capable to bring such an

instantaneous advantage is emasculation of cosexual individuals. By removing the source of pollen,

we can simulate the effect of the apparition of a male-sterile mutant. Hence, relying on

emasculation to investigate female advantage should help clarifying the early steps of the evolution

of gynodioecy by ruling out possible pleiotropic effects of the male-sterility genes themselves as

well as the effects of modifier genes selected after the establishment of sexual dimorphism (Sun &

Ganders, 1986). However, one possible issue with emasculation experiments is that, by removing

stamens, we remove not only the source of self-interference (pollen) but also a potential reward

(pollen) as well as possible visual clues (stamens) for pollinators (e.g. Duffy and Johnson, 2011).

Another possible issue with emasculation experiments that applies to hermaphrodite but not to

monoecious plants is the potential effect of the experimentation on corolla development, due to

corolla's physiological interdependence with stamens (Gibson & Diggle, 1998). This illustrates the

need for careful selection of model species (Charlesworth, 1993) and of emasculation controls to

account for possible confounding effects. Surprisingly, given their great potential, emasculation

experiments involving gynodioecious plants remain very scarce. We found only four previous

studies that have explored the consequences of emasculation on female fitness in gynodioecious

plants (Kikuzawa, 1989; Pettersson, 1992; Alonso & Herrera, 2001; Wang *et al.*, 2021), none of

which had attempted to control for an effect of the treatment on pollinator attractiveness.

Focusing on gynodioecious plants with unusual life histories or breeding systems could help

understand possible limits to generalization (Freeman *et al.*, 1997). Gynodioecious species tend to

be herbaceous rather than woody (Caruso *et al.*, 2016), hermaphrodite rather than monoecious

(Lloyd, 1975), and self-compatible rather than self-incompatible (Dufaÿ and Billard, 2012). Trees

are typically outcrossed (Petit & Hampe, 2006), and monoecy and self-incompatibility both

promote outcrossing. Gynodioecious species with these unusual features could help explore how

gynodioecy can evolve when outcrossing rates are already maximum.

In this study, to test if self-pollination avoidance is sufficient to explain the persistence of females

in outbred species, we selected a long-lived Fagaceae tree, European chestnut (*Castanea sativa*).

Spontaneous male-sterile variants have been reported in this species (Kaul, 1988; Soylu, 1992).

Chestnut trees are monoecious, self-incompatible, and duodichogamous, all mechanisms frequently

interpreted as adaptations to promote outcrossing (Lloyd & Webb, 1986; Webb & Lloyd, 1986;

Bertin & Newman, 1993; Routley *et al.*, 2004; Koelling & Karoly, 2007). In a previous study, we

found self-fertilization rates of 5% in chestnut trees with fully developed stamens and 1% in trees with dysfunctional stamens (Larue *et al.*, 2022). Even with strong inbreeding depression, this difference cannot explain the maintenance of gender dimorphism. After attempting to establish gynodioecy in the wild in European chestnut, we therefore studied if female chestnut trees have some kind of female advantage other than increased outcrossing rate. Specifically, we compared five key components of female fitness between genders in a chestnut experimental stand: basal area (a proxy for tree size), flower density, burr set, fruit set and fruit weight. Both genders harbour nectar-producing male catkins that attract pollinators; the only visible difference is that the stamens are aborted or greatly reduced in female trees (Larue *et al.*, 2021a, 2022). We took advantage of this feature to assess the importance of self-pollen avoidance in an experiment where we removed male catkins not only from cosexual but also from female chestnut trees. This unique situation with female flowers associated either with fertile or with sterile but rewarding male catkins should help to disentangle the positive and negative effects of male function on female fitness.

## MATERIALS AND METHODS

## Chestnut floral biology

Chestnuts (*Castanea spp.*) are insect-pollinated Fagaceae tree species (Larue *et al.*, 2021a; Petit and Larue, 2022). Chestnut trees are characterized by massive blooming, huge pollen production and the largest pollen:ovule ratio ever reported in plants (~10-30 millions; Larue *et al.*, 2021a). They have multiple mechanisms limiting self-pollination. First, they are monoecious, with separate male and female flowers distributed in two types of inflorescences: unisexual male catkins and bisexual catkins composed of one or two female inflorescences associated with a single male catkin (Larue *et al.*, 2021a). Second, they have a complex phenology called duodichogamy, characterized by two peaks of pollen emission, which reduces the risk of self-pollination without eliminating it completely (Hasegawa *et al.*, 2017). Unisexual male catkins bloom first and release huge amounts of pollen, about 97% of the total (Larue *et al.*, 2021a). Non-rewarding female flowers then become receptive. Finally, male catkins from bisexual inflorescences start to emit pollen, resulting in a second much smaller pollen emission peak. Third, chestnuts are self-incompatible (Xiong *et al.*, 2019).

**<u>Figure 1:</u>** Degrees of male-sterility observed. (a) Fully male-sterile (i.e. female) tree. Most stamens are aborted and do not produce pollen. (b) Mostly male-sterile tree. Stamens do not protrude from glomerules. (c) Partly male-sterile tree. Stamens slightly protrude from glomerules. (d) Fully male-fertile (i.e. cosexual) tree. Stamen filaments are long and largely protrude from glomerules. These trees produce large amounts of pollen.

In European chestnuts and in its hybrids, two genders can be distinguished: cosexual trees, which are fully male-fertile; and female trees, which are completely or largely male-sterile (Solignat and Chapa, 1975; Larue *et al.*, 2022; Fig. **1**). These male-sterile trees have dysfunctional staminate flowers with fully aborted stamens or with short stamen filaments producing scarce amounts of mostly non-functional pollen (Bounous *et al.*, 1992). Interestingly, male-sterile catkins continue to

produce nectar and attract insects such as flies and beetles, but not pollen-seeking insects such as

hoverflies and bees (Larue *et al.*, 2021a). All flowers on a tree and all trees from a clone have the

same type of male flowers (Larue *et al.*, 2021b, 2022). In interspecific crosses, male sterility is

expressed only in some cytoplasmic backgrounds (Bolvanský & Mendel, 1999; Sisco *et al.*, 2014;

Larue, 2021). In particular, in crosses between European chestnut (*C. sativa*) and Japanese chestnut

(*C. crenata*), only crosses with European chestnut as mother generate male sterile individuals

(Larue, 2021). However, within European chestnuts, character inheritance studies point to strict

nuclear inheritance of male sterility involving a major recessive gene modified by one or more other

genes (Soylu, 1992; Bolvanský & Mendel, 1999), not to nucleo-cytoplasmic inheritance, probably

because cytoplasmic variation is often lacking within populations of European chestnut (Fineschi *et*

*al.*, 2000; Larue, 2021).

Chestnut trees, despite their long lifespan and large sizes, are good models for studying female

advantage. Fruit set is easy to estimate in chestnuts. A female inflorescence is typically composed

of three female flowers located side by side. The inflorescence develops into a spiny infrutescence

called a burr. In each burr, each flower, if pollinated, gives a nut (a fruit with a single seed

surrounded by a closed pericarp), and if not pollinated, an empty fruit. Multi-seeded nuts are very

rare (< 1%, unpublished data). Hence, the proportion of developed fruits per burr should provide an

indication on pollination success. Moreover, it is technically feasible to emasculate trees despite the

tiny size of flowers, as male flowers are packed together into inflorescences that can be removed as

one unit (Fig. **2**).

**Figure 2:** Chestnuts have two types of catkins: numerous unisexual male catkins that flower first
and few bisexual catkins located at the tip of the branches that flower last.

## Gender polymorphism in chestnut forests

223  We visited 14 naturally regenerated populations of European chestnut, one in Spain (described in
224  Larue, 2021) and the rest in France, and estimated the proportion of trees with different types of
225  stamens using descriptions of Solignat & Chapa (1975) as reference (Fig. **1**). Stands were selected
226  to be as natural as possible, avoiding ancient plantations and coppices, and focusing on regions
227  identified as potential glacial refugia for chestnut (Krebs *et al.*, 2019).

## Female advantage

**Study site**

For this work, we used the INRAE chestnut germplasm collection located in Villenave d'Ornon
(44.788319 N, −0.577062 E) (Larue *et al.*, 2021b). Most trees investigated come from a 3.5 ha-
large experimental unreplicated orchard planted in 1990. It includes 211 grafted trees (ramets)
corresponding to 83 different genotypes (clones) assigned to different chestnut species: the
European chestnut (*C. sativa*; 50% of the trees), the Japanese chestnut (*C. crenata*; 9%), the
Chinese chestnut (*C. mollissima*; 8%) and their interspecific hybrids, mostly *C. sativa* × *C. crenata*

hybrids (27%). The trees are grafted on three different hybrid rootstocks: 'Marsol' (CA07; 56% of

the trees), 'Maraval' (CA74; 27%), or 'Marlhac' (Ca118; 3%). The remaining trees (14%) grow on

their own roots. Among the clones, 55 are cosexual and 28 are partly or completely male-sterile

(thereafter called female), corresponding to 137 (65%) cosexual and 74 (35%) female ramets

distributed rather evenly throughout the plot (Supplementary Material 1). This even-aged collection

includes genotypes of forest trees as well as some clones selected for fruit production, mostly

superior trees selected in nature or first generation hybrids, as selective breeding is very recent.

**Female fitness components in the study plot and in a subset of trees**

In 2019, we systematically compared several components of female fitness: basal area, fruit set and

fruit weight, controlling for species status and rootstock identity. For basal area, we measured the

diameter of the stem (or for multi-stemmed individuals, of each stem) at breast height and derived

the cross-sectional area. To estimate fruit set, we targeted at least 30 burrs per tree. We statistically

corrected fruit set to account for the possibility of premature abortion of empty burrs as explained

below in the section *Statistical analyses*. To estimate fruit weight, we collected 50 developed fruits

per tree and weighted them. In 2020, we studied two additional female fitness components on a

subset of 18 trees, nine from each gender, belonging either to European chestnut (five clones, each

with two ramets except for one clone) or to *C. sativa* × *C. crenata* hybrids (five clones, each with

two ramets except for one clone). On 10 marked branches per tree, we counted the number of

female inflorescences in the spring, the number of burrs formed in the summer, and the number of

mature burrs harvested in the fall. We estimated burr set by dividing the number of burrs counted in

August by the number of female inflorescences counted in June during flowering time. To

investigate inter-annual fluctuations of fruit set between genders, we gathered fruit set estimates for

these trees in 2018, 2019, 2020 and 2022. In view of the composition of this large collection and of

the use of both female and cosexual clones in chestnut cultivation, systematic gender-biased effects

of artificial selection on phenotypic traits were deemed unlikely.

Emasculation experiment

**Plant material**

We performed the emasculation experiment in 2019 in three orchards of the INVENIO experimental

station in Douville (45.019723 N, 0.614637 W). We selected six hybrid clones, three female ones

('Bellefer', 'OG19' and 'Bouche de Bétizac') and three cosexual ones ('Florifer', 'Maraval' and

'Marigoule'). These trees are planted in three nearby orchards. The first orchard is composed of 8-year-old trees belonging to 'Bellefer', 'OG19', 'Florifer' and 'Maraval' clones. The second orchard is composed of 20 m-high trees belonging to 'Bouche de Bétizac' and 'Marigoule' clones. We replicated the experiment on 'Marigoule' clone in a third orchard, because this clone is very susceptible to cynips (*Dryocosmus kuriphilus*), which complicates fruit set measurements. Indeed, following cynips attacks, female flowers tend to proliferate, resulting in atypical inflorescences with more than three flowers that are difficult to use to estimate fruit set.

**Modalities**

To remove male flowers for the emasculation treatment, we cut all unisexual male catkins and the male part of bisexual catkins with a scissor (Fig. **2**). As these two types of catkins flower at different times, we proceeded in three steps before trees started to flower. At the first pass, in the end of May, we removed all unisexual male catkins. At the second pass, we checked if no unisexual male catkins remained and we removed the male part of bisexual catkins. At the third and last pass, we removed the male flowers of bisexual catkins that remained.

We also removed the male-sterile catkins of female clones to evaluate the decrease in insect attractiveness or possible reallocation of resources in the female function following their removal. For younger trees, for each modality, we selected five 'OG19' trees, six 'Bellefer' trees, five 'Florifer' trees and two 'Maraval' trees. Hence, there were 18 control and 18 emasculated chestnuts. We emasculated entirely each tree. For older trees, we applied the treatment to single branches. In the first orchard, we selected five 'Bouche de Bétizac' and five 'Marigoule' trees. For each tree, we selected 20 branches and used ten for each modality. We replicated these two modalities on three more 'Marigoule' trees from another orchard.

For young trees, we also performed partial emasculations experiments on two trees per clone for the two most abundant clones, i.e. 'Bellefer' and 'Florifer'. In the 'unisexual modality', we removed all unisexual male catkins but left the male part of bisexual catkins. On the contrary, in the 'bisexual modality', we removed the male part of bisexual catkins but left all unisexual male catkins.

Statistical analyses

**Data analysis**

We performed all analyses with R software (v3.6.6; R Core Team, 2013). The corrected fruit set was calculated with basic functions implemented in R and the violin plot is computed with ggplot2 ( v3.6.3; Wickham, 2016a) and ggthemes (v4.2.4; Arnold, 2016) packages.

**Corrected fruit set**

To estimate the probability of a female flower to give a fruit, we statistically corrected fruit set to account for the possibility of premature abortion of empty burrs, i.e. burrs with no developed fruit. Not accounting for these missing empty burrs when estimating fruit set in the fall would result in an overestimation of fruit set (strong overestimation when fruit set is low, slight overestimation when it is high). Using a zero-deflated binomial distribution, we obtained unbiased estimates of fruit set for the large-scale fruit set study of 2019, using only information from burrs with at least one developed fruit (Larue *et al.*, 2022). For a given tree, typical burrs developed from 3-flowers inflorescences with one, two, or three developed fruits are noted $x_1$, $x_2$, and $x_3$ while the total number of harvested burrs is noted $x_{tot}$. Based on the definition of multinomial distribution, a maximum likelihood estimator reflecting pollination probability $\hat{p}$ is obtained (Annex 1):

$$\hat{p} = \frac{3}{2} - \sqrt{3\frac{x_{tot}}{x_1 + 2x_2 + 3x_3} - \frac{3}{4}}$$

All these estimates are automatically computed for all trees in R using the *apply()* function.

In clones susceptible to cynips, inflorescences with more than three flowers can form following cynips attacks, resulting in burrs with up to nine fruits instead of the standard case with three flowers (Breisch, 1995). We excluded these rare abnormal burrs as well as even rarer cases with less than three fruits per burr from fruit set estimations.

**Estimation of female advantage**

To compare female fitness components between genders across the study plot (basal area, fruit set and fruit weight) as a function of species and rootstock, we checked ANOVA application conditions, i.e. independence, normality and homogeneity of residuals, using a Durbin-Watson test, a Shapiro test and a Bartlett test. The normality and homogeneity of the residuals were also visually inspected using the 'Normal Q-Q' and 'Scale-Location' plots. Since these conditions were not satisfied, we used non-parametric permutational analysis of variance (PERMANOVA) using *aovp()* function from lmPerm package (Wheeler & Torchiano, 2016).

We tested the differences in flower density and burr set measured in 2020 between genders and

between species (European chestnut and *C. sativa × C. crenata* hybrids) on a subset of 18 trees

using a Student *t* test (unilateral for gender effect, and bilateral for species effect).

We also tested the differences in fruit set between genders and between species during four years on

the same subset of 18 trees using the same approach.

**Emasculation experiment**

We tested the significance of the difference between the modalities (control and complete

emasculation) with a Fisher exact test.

## RESULTS

### Gender polymorphism in chestnut forests

338 We monitored gender variation in 430 trees from 14 populations (26 to 42 trees sampled per

339 population) (Supplementary Material 2). Male-fertile trees were more frequent in all but one

340 population. On average, 15% of the trees were at least partly male-sterile (11% fully or mostly

341 male-sterile and 4% partly male-sterile). Four populations were monomorphic, harbouring

342 only male-fertile trees.

### Female advantage in chestnut

*Basal area, fruit set and fruit weight*

We found no difference in basal area between genders but significant differences among species and

rootstocks (Tables 1 and 2, Supplementary Material 3). In contrast, fruit set and fruit weight measured

in 2019 differed between genders but not among species or rootstocks. Fruit set was much higher in

females than in cosexual trees for both European chestnut and for *C. sativa × C. crenata* hybrids. In

contrast, fruit weight was slightly lower in female than in cosexual trees in both taxa. Fruit set and

fruit weight were negatively correlated (Spearman rank-correlation coefficient: -0.15, *p*-value =

0.04).

Table 1: Summary table for the PERMANOVA testing of the effects of factors (gender, species and rootstock) on each female fitness component

| Fitness component | Source | Df | R Sum Sq | R Mean Sq | Iterations | Pr(Prob) | |
|---|---|---|---|---|---|---|---|
| Basal area | Gender | 1 | 8500 | 8500 | 51 | 1.0 | |
| | Species | 4 | 960000 | 240000 | 5000 | $<10^{-15}$ | *** |
| | Rootstock | 3 | 470000 | 160000 | 5000 | 0.006 | ** |
| | Residuals | 202 | 8600000 | 43000 | | | |
| Fruit set 2019 | Gender | 1 | 2.5 | 2.5 | 5000 | $<10^{-15}$ | *** |
| | Species | 4 | 0.05 | 0.013 | 160 | 1.0 | |
| | Rootstock | 3 | 0.13 | 0.04 | 319 | 0.3 | |
| | Residuals | 167 | 5.4 | 0.03 | | | |
| Fruit weight | Gender | 1 | 273 | 270 | 5000 | $<10^{-15}$ | *** |
| | Species | 4 | 141 | 35 | 717 | 0.3 | |
| | Rootstock | 3 | 83 | 28 | 1201 | 0.4 | |
| | Residuals | 167 | 4000 | 24 | | | |

Table 2: Gender effect on female fitness components

| Fitness component | #♂[1] | #♀[1] | Mean ♂[2] | Mean ♀[2] | Test[3] |
|---|---|---|---|---|---|
| Basal area (cm²) | 137 | 74 | 308 (257) | 290 (194) | ns (Permanova) |
| Flower density | 9 | 9 | 51 (34) | 67 (80) | ns (Student *t* test) |
| Burr set | 9 | 9 | 0.77 (0.15) | 0.87 (0.14) | ns (Student *t* test) |
| Fruit set | 115 | 61 | 0.57 (0.21) | 0.84 (0.11) | $< 10^{-15}$ (Permanova) |
| Fruit weight (g) | 115 | 61 | 10.5 (5.2) | 7.7 (4.3) | 0.0013 (Permanova) |

[1]Number of cosexual (♂) and female (♀) trees

[2]Mean value for cosexual and female trees (standard deviation)

[3]Difference significance (test performed)

*Flower density and burr set*

Flower density was highly variable among trees and there was no significant difference between genders or between species (Tables 2 and 3, Supplementary material 4). Similarly, there were no difference in burr set between genders nor between species in our sample.

*Fruit set across years*

For female trees, average fruit set fluctuated during the four studied years, ranging from 0.69 in 2020 to 0.83 in 2018 (Table 3, Fig. **3**, Supplementary material 4). For cosexual trees, average fruit

370 set fluctuated more drastically, ranging from 0.34 in 2022 to 0.64 in 2018. During all four fruiting

371 episodes, fruit set was higher in females: the female/cosexual fruit set ratio ranged from 1.3 in 2018

372 and 1.6 in 2019, to 2.0 in 2020 and 2.5 in 2022.

374 **Figure 3:** Inter-annual variation in fruit set in cosexual and female chestnut trees.

376 Table 3: Fruit set measured during four years and details for 2020

|  | Fruit set | | | | Details for 2020 | | |
|---|---|---|---|---|---|---|---|
| Gender | 2018 | 2019 | 2020 | 2022 | #♀-inflo[1] | #Burrs[2] | Burr set[3] |
| Female | 0.83 | 0.77 | 0.69 | 0.86 | 67.1 | 59.9 | 0.86 |
| Cosexual | 0.64 | 0.49 | 0.34 | 0.34 | 50.8 | 35.9 | 0.77 |
| **Ratio F/C** | **1.30** | **1.56** | **2.01** | **2.54** | **1.32** | **1.67** | **1.12** |
| *t-test* | 0.003 | 0.002 | 0.004 | $<10^{-6}$ | ns | ns | ns |

383 [1]Mean number of female inflorescences per tree (10 branches per tree)

384 [2]Number of burrs counted in August

385 [3]Burr set calculated by dividing the number of burrs by the number of female inflorescences

## Emasculation experiment

For older trees, the emasculation treatment resulted in a significant increase in fruit set compared to controls for the cosexual clone (Table 4). Fruit set in control trees was 52% whereas it was 67% for emasculated trees (Fisher exact test, $p < 10^{-5}$). In contrast, the emasculation treatment resulted in a significant decrease in fruit set compared to control for the female clone. Fruit set in control trees was 95% whereas it was 85% for emasculated trees ($p < 10^{-7}$).

Table 4: Fruit set of control and emasculated adult trees

| Gender[1] | Clone | Modality | #replicates | Mean (sd) | Fisher exact test |
|---|---|---|---|---|---|
| F | Betizac | Control | 5 ramets × 10 branches | 0.95 (0.04) | $10^{-6}$ |
| F | Betizac | Emasculated | 5 ramets × 10 branches | 0.85 (0.07) | |
| C | Marigoule | Control | 8 ramets × 10 branches | 0.52 (0.08) | $10^{-5}$ |
| C | Marigoule | Emasculated | 8 ramets × 10 branches | 0.66 (0.07) | |

[1]F: Female (male-sterile); C: cosexual (male-fertile, monoecious)

For younger trees, which were entirely emasculated, the emasculation treatment resulted in an increase in fruit set compared to controls for both cosexual clones (Table 5). For 'Florifer', fruit set in control trees was 46% whereas it was 63% for emasculated trees, a significant increase ($p < 10^{-10}$). For 'Maraval', fruit set of control trees was 73% compared to 78% for emasculated trees, a non-significant increase ($p > 0.16$). In contrast, the emasculation treatment resulted in a significant decrease in fruit set compared to control for one of the two studied female clones. For 'Bellefer', fruit set in control trees was 91% whereas it was 81% for emasculated trees, a significant decrease ($p < 10^{-5}$). However, for 'OG19', fruit set in control trees was 77% whereas it was 82% for emasculated trees, a non-significant increase ($p > 0.8$).

409    Table 4: Fruit set of control and emasculated adult trees

| Gender[1] | Clone | Modality | #replicates | Mean (sd) | Fisher exact test |
|---|---|---|---|---|---|
| F | Betizac | Control | 5 ramets × 10 branches | 0.95 (0.04) | $10^{-6}$ |
| F | Betizac | Emasculated | 5 ramets × 10 branches | 0.85 (0.07) | |
| C | Marigoule | Control | 8 ramets × 10 branches | 0.52 (0.08) | $10^{-5}$ |
| C | Marigoule | Emasculated | 8 ramets × 10 branches | 0.66 (0.07) | |

410    [1]F: Female (male-sterile); C: cosexual (male-fertile, monoecious)
411
412
413
414    Table 5: Fruit set of control and emasculated young trees

| Gender[1] | Clone | Modality | #trees | Mean (sd) | Fisher exact test |
|---|---|---|---|---|---|
| F | Bellefer | Control | 5 | 0.91 (0.04) | $10^{-5}$ |
| F | Bellefer | Emasculated | 5 | 0.81 (0.11) | |
| F | OG19 | Control | 5 | 0.77 (0.14) | ns |
| F | OG19 | Emasculated | 5 | 0.82 (0.05) | |
| C | Maraval | Control | 2 | 0.73 (0.09) | ns |
| C | Maraval | Emasculated | 2 | 0.78 (0.06) | |
| C | Florifer | Control | 6 | 0.46 (0.14) | $10^{-10}$ |
| C | Florifer | Emasculated | 6 | 0.63 (0.14) | |

[1]F: Female (male-sterile); C: cosexual (male-fertile, monoecious)

We performed partial emasculations experiments on two clones (Supplementary Table 5). For

'Florifer', a cosexual clone, fruit set increased compared to the control after removing all male

catkins or only catkins from unisexual inflorescences. On the other hand, when we removed catkins

from bisexual inflorescences, fruit set decreased compared to the control. For 'Bellefer', a female

clone, fruit set following partial emasculations were intermediate between the control (higher fruit

set) and the complete emasculation treatment (lower fruit set), with more reduced fruit set after

removing male catkins from bisexual inflorescences.

# DISCUSSION

Finding gynodioecy in European chestnut was unexpected. While dioecy and monoecy are particularly frequent in trees, stable gynodioecy is rare (Olson *et al.*, 2016; but see Steyn & Robbertse, 1990; Ellis & Sedgley, 1993; Gibson & Diggle, 1998; Penagos Zuluaga *et al.*, 2020). Moreover, gynodioecy is unknown in Fagaceae, a widespread species-rich family (Petit *et al.*, 2013). Chestnut is an ecologically and economically important genus cultivated for its nutritious nuts. The fact that gynodioecy had remained unnoticed in chestnuts seems therefore surprising. In gynodioecious plants, the average female frequency is 25% (Varga & Soulsbury, 2020). If gender variation is studied in few populations, evidence for gynodioecy might be missed (Dufaÿ *et al.*, 2014; Caruso *et al.*, 2016). Our survey illustrates this point, as we found a rather low (15%) but variable (0-53%) proportion of females, so that half of the populations have less than 10% of female trees. We also found some partly male-sterile trees, a frequent feature in gynodioecious species (Koelewijn & van Damme, 1996; Schultz, 2002). Because spontaneous male sterile mutants have been reported in other chestnut species (Kaul, 1988), exploring gender variation in other chestnut species would be worthwhile.

To explore female advantage in chestnut, we investigated for the first time in a tree several key female fitness components covering the complete life cycle (Philipp, 1980). Using a grafted clonal collection established for conservation purposes, we found no difference in basal area and flower production between genders. In grafted trees, the rootstock partly controls a number of phenotypic traits, including growth, but scions are known to influence tree canopy height in chestnut (Camisón *et al.*, 2023), so the lack of gender effect on scion growth is remarkable. This finding suggests that there is no significant reallocation of resources to the female function from the lost male function at this stage. We also found that females have lighter, not heavier, fruits than cosexual trees. Fruit weight is negatively correlated with fruit set, a likely effect of competition for resources between fruits within female inflorescences (Larue, 2021). Therefore, females do not appear to reallocate significant resources to fruit growth. Long-lived perennial plants need to allocate more resources than short-lived plants for the maintenance of vegetative structures, so intersexual competition for resources plays a relatively smaller role in long-lived than in short-lived plants (Obeso, 2002). Furthermore, reallocation of resources from male toward female function is necessarily limited in gynodioecious trees because of the higher cost for female than for male reproduction in woody plants (Obeso, 2002; Thomas, 2011).

The single female advantage identified in chestnut is increased fruit set. This advantage is large and holds for both European chestnut and for its hybrids. Japanese and Chinese chestnuts, for which

evidence for gynodioecy is lacking, had fruit sets similar to those of cosexual European chestnut trees and hybrids. To our knowledge, gender effect on fruit set has not been investigated before. However, chestnut cultivation in Europe relies heavily on female clones (Furones-Pérez & Fernández-López, 2009) and some researchers have noticed that female chestnuts tend to have higher yields (e.g. Pereira-Lorenzo & Ramos-Cabrer, 2004). We observed higher and more stable fruit set in females than in cosexual trees during four years. We tentatively attribute the less stable fruit set of cosexual trees to Jensen's inequality (Jensen, 1906; Garibaldi *et al.*, 2011): as fruit set decelerates with increased pollen receipt, the deficit of a bad pollination episode is necessarily higher than the benefit of a good pollination episode. This asymmetrical response is more pronounced at lower pollination rates where small changes translate into large differences. What matters for long-term female maintenance is the difference in geometric lifetime fitness between genders (Eckhart, 1992). Hence, episodes in which cosexual individuals have particularly reduced fruit set compared to females will accentuate long-term female advantage. Altogether, our experimental data support the hypothesis that female advantage is sufficient for female maintenance, as it is close to the two-fold threshold needed in the case of nuclear inheritance of male sterility (Lewis, 1941).

The emasculation experiments comfort the hypothesis that increased fruit set is the main female advantage in chestnut. When we removed male catkins (the sources of self-pollen) in cosexual trees, fruit set increased. In contrast, when we removed male-sterile catkins from female trees, fruit set decreased or remained unchanged. These contrasting effects suggest that the presence of pollen in anthers is decisive. On the contrary, reallocation of resources from the male catkins following emasculation is unlikely, as it should affect both female and cosexual trees. We thus attribute the opposite effects of emasculation on fruit set in the two genders to decreased self-pollination and associated ovule and seed discounting in emasculated cosexual trees and to reduced attractiveness to pollinators in 'emasculated' female trees. Together, these findings suggest that, in cosexual trees, the positive effect of emasculation caused by the elimination of the source of self-pollination overcomes the negative effect of decreased pollinator attractiveness caused by the elimination of the rewarding catkins. In a previous emasculation experiment involving Chinese chestnut, Zhao and Liu (2009) described increased fruit yield by up to 39%, indicating that self-pollination also negatively affects fruit production in this species not known to be gynodioecious. The finding that a single main fitness component (fruit set) determines female advantage and that this advantage can be reproduced in cosexual trees by removing the source of pollen is compatible with the hypothesis of a relatively recent origin of gynodioecy in European chestnut. Other studies have reported increased seed set following emasculation of self-incompatible hermaphrodite plant species (e.g., Waser & Price, 1991;

Vaughton & Ramsey, 2010; Duffy *et al.*, 2013, 2021), suggesting that self-pollen often interferes with

female fitness in outcrossing plants.

What is the cause of the large difference in fruit set between genders? Self-pollination in cosexual

chestnut trees is massive, a likely consequence of huge pollen production and frequent geitonogamy

in this large mass-flowering tree. In the study site, we previously estimated, using a spatially explicit

mixed mating model, that the self-pollination rate is 74% (Larue et al. 2022). Hasegawa *et al.* (2009)

found even higher values of self-pollination in Japanese chestnut by genotyping pollen grains directly

collected on stigmas. The model-based estimate of the proportion of ovules that are eventually self-

fertilised is 48%, as cross-pollen is somehow favoured over self-pollen. However, according to the

model, 95% of the remaining self-fertilized ovules abort, meaning that 46% of ovules are lost in

cosexual trees, a very large proportion (Larue et al. 2022). Cytological studies and pollination

experiments have revealed both prezygotic late-acting self-incompatibility and early-acting

inbreeding depression in chestnut (Xiong et al. 2019). Self-pollen tubes grow well in the styles but

many ovules abort when pollen tubes reach the ovary. If self-fertilization nevertheless takes place,

the resulting embryos stop growing before a mature seed is formed due to early inbreeding depression

(Xiong et al. 2019). Early inbreeding depression is an important but often neglected component of

inbreeding depression (Lande *et al.*, 1994), particularly in outcrossing species (Husband & Schemske,

1996). An obvious advantage of early elimination of self-pollinated ovules and selfed embryos is the

prevention of wasteful provisioning of low-quality progeny (Johnson *et al.*, 2019; Larue *et al.*, 2022).

Hence, in chestnuts, gynodioecy should not be considered as a superfluous, redundant outcrossing

mechanism (Baker, 1963; Krohne *et al.*, 1980). Little attention has been given so far to self-pollen

interference as a source of female advantage (but see Kikuzawa, 1989). We suggest that it is time to

reconsider the evidence, as late-acting self-incompatibility is widespread, especially in woody plants

(Seavey & Bawa, 1986; Sage *et al.*, 1994; Gibbs, 2014; Johnson *et al.*, 2019).

We used female trees as 'natural controls' in the emasculation study, removing their rewarding but

sterile male catkins. This decreased their fruit set, showing that even sterile male catkins promote

pollination. Our pilot partial emasculation experiment further suggests that it is the late-flowering

male catkins physically associated with female inflorescences that attract insects close to female

flowers (van de Pijl, 1978; Larue et al., 2021a). We previously showed that insect visits to female

trees are reduced compared to visits to cosexual trees, but only for insects feeding on pollen, such as

syrphid flies and bees (Larue *et al.*, 2021a). However, the scarcity of these insects on female trees has

no adverse consequence for pollination because they fail to visit female flowers. In contrast, beetles

and calyptrate flies, the putative pollinators of chestnut, visit both genders equally. Hence, in

chestnuts, the breeding system (monoecy) and main reward type (nectar) is not an obstacle to the evolution of gynodioecy.

The emasculation experiments involved either single branches or entire trees, resulting in both cases in increased fruit set in cosexual trees. Beetles are particularly abundant in chestnut groves, walking back and forth on the crown, presumably bringing much self-pollen from nearby male catkins on female inflorescences (Larue *et al.*, 2021a). By emasculating branches rather than entire trees, we reduce self-pollination brought about by the behaviour of these insects, not by the behaviour of insects flying among branches of the same tree, such as flies. This suggests that insects that tend to walk on inflorescences rather than to fly from one place to another are major causes of self-pollination. Greater abundance of those pollinators responsible for insect-assisted geitonogamous mating could widen the fitness gap between genders and further promote gynodioecy.

In European chestnut, female and cosexual trees are equally outcrossed, ruling out outbreeding advantage as possible cause of female advantage. According to Baker (1959), '*one might expect that in any hermaphrodite taxon where an incompatibility system is already established (or even a strongly effective outbreeding system of some other sort) there will be little likelihood of dioecism arising through direct natural selection. Any exceptions to this rule would be in the nature of unstoppable accidents with special causes*'. Our study, focusing precisely on one of these exceptions, illustrates that the 'special causes' mentioned by Baker are no accidents after all. Rather, they involve important early acting genetic mechanisms of plant reproduction. Because females avoid self-pollination and its deleterious consequences (Charlesworth, 1993), they are more effectively outcrossed than cosexual individuals, if outcrossing success is measured in numbers of outcrossed progenies rather than in proportion of outcrossed progenies. Considering 'self-pollination avoidance' rather than 'selfing avoidance' as key female advantage could clarify the debate on the relative importance of genetic versus ecological mechanisms for the evolution of unisexuality (Freeman *et al.*, 1997). The world of possibilities for the evolution of female advantages is vast but some pathways might be more important than others, compounded by plant species life histories.

## ACKNOWLEDGEMENTS

We thank Teresa Barreneche (chestnut germplasm collection administrator) for her invaluable support, and Catherine Bodénès for her collaboration and for allowing us to report the first results of her study on gender variation in natural chestnut populations from southern France. This study was

possible thanks to the continuous management of the experimental plots by INRAE experimental unit Vigne Bordeaux (UEVB) and by Xavier Capdevielle of UMR Biogeco. We thank Yannick Mellerin, Céline Lalanne, Chantal Helou Bagate and our many great students for their invaluable help in the field. We also thank the INVENIO chestnut team and the numerous seasonal workers who helped with the emasculation experiment. Many thanks to Olivier Lepais and to Etienne K. Klein for developing a fruit set model and to Josefina Fernández López for sharing her knowledge on chestnut reproduction and guiding us in the field. Mathilde Dufaÿ comments helped improve an earlier version of this manuscript.

This paper was part of the PhD and postdoc of CL. The Cifre thesis (Conventions Industrielles de Formation par la Recherche) was supported by the ANRT (Association Nationale de la Recherche de la Technologie, convention Cifre N° 2018/0179). It was funded by Invenio, the Région Nouvelle-Aquitaine (Regina project N° 22001415-00004759 and chestnut pollination project N° 2001216-00002632), by INRAE (Institut National de Recherche pour l'Agriculture et l'Environnement) and by ANR (project ANR-21-PRRD-0008-01).

## AUTHOR CONTRIBUTIONS

CL, RJP and Biogeco colleagues estimated fruit set in the INRAE chestnut germplasm collection. CL and Invenio colleagues carried out the emasculation experiments in Invenio orchards. CL and RJP estimated fruit set in INVENIO orchards. CL corrected fruit set thanks to the statistical correction developed by Etienne K. Klein. CL and RJP carried out statistical analyses. CL wrote the first draft of this article and RJP revised it.

## DATA AVAILABILITY

The data that support the findings of this study are openly available at:
https://entrepot.recherche.data.gouv.fr/dataset.xhtml?persistentId=doi:10.57745/MOTPXM

They can be cited as: Larue, Clément, 2022, "DATA: Self-interference and female advantage in chestnut", https://doi.org/10.57745/MOTPXM, Recherche Data Gouv, V1.

The data on gender variation that support the findings of this study are openly available at:

582 https://entrepot.recherche.data.gouv.fr/dataset.xhtml?persistentId=doi:10.57745/LFZFT2

583 They can be cited as: Bodénès, Catherine, 2023, "Conservation des ressources génétiques de
584 châtaignier", Recherche Data Gouv, V1.

## COMPETING INTEREST

587 The authors declare that they have no known competing financial interests or personal relationships
588 that could have appeared to influence the work reported in this paper.

## REFERENCES

**Alonso C, Herrera CM**. **2001**. Neither vegetative nor reproductive advantages account for high
frequency of male-steriles in southern Spanish gynodioecious *Daphne laureola* (Thymelaeaceae).
*American Journal of Botany* **88**: 1016–1024.

**Arnold JB**. **2019**. ggthemes: extra themes, scales and geoms for 'ggplot2'. R package version 4.2.
0.

**Ashman T-L**. **2002**. The role of herbivores in the evolution of separate sexes from
hermaphroditism. *Ecology* **83**: 1175–1184.

**Baker HG**. **1959**. Reproductive methods as factors in speciation in flowering plants. *Cold Spring
Harbor Symposia on Quantitative Biology* **24**: 177–191.

**Baker HG**. **1963**. Evolutionary mechanisms in pollination biology: Origins and functions of floral
systems are being elucidated by genetical and ecological studies. *Science* **139**: 877–883.

**Barrett SCH**. **2002**. Sexual interference of the floral kind. *Heredity* **88**: 154–159.

**Barrett SCH, Lloyd DG, Arroyo J**. **1996**. Stylar polymorphisms and the evolution of heterostyly
in *Narcissus* (Amaryllidaceae). In: Lloyd DG, Barrett SCH, eds. Floral Biology. Boston, MA:
Springer US, 339–376.

**Bawa KS**. **1980**. Evolution of dioecy in flowering plants. *Annual Review of Ecology and
Systematics* **11**: 15–39.

**Bawa KS**. **1982**. Outcrossing and the incidence of dioecism in island floras. *The American
Naturalist* **119**: 866–871.

**Bawa KS**. **1984**. The evolution of dioecy-Concluding remarks. *Annals of the Missouri Botanical
Garden* **71**: 294–296.

**Bertin RI, Newman CM**. **1993**. Dichogamy in angiosperms. *The Botanical Review* **59**: 112–152.

**Bolvanský M, Mendel L**. **1999**. Inheritance of some fruit and flower characteristics in full-sib progenies of chestnut. *Acta Horticulturae*: 339–344.

**Bounous G, Paglietta R, Peano C**. **1992**. Methods for observing chestnut pollen viability, germinability and pollen tube growth. *Proceedings of the International Chestnut Conference, Morgantown, USA*: 76–78.

**Breisch H**. **1995**. *Châtaignes et Marrons*. Paris, France: Centre technique interprofessionnel des fruits et légumes.

**Burbidge AH, James SH**. **1991**. Postzygotic seed abortion in the genetic system of *Stylidium* (Angiospermae: Stylidiaceae). *Journal of Heredity* **82**: 319–328.

**Camisón Á, Martín MÁ, Sánchez-Bel P, Flors V, Cubera E, Solla A**. **2023**. Effect of grafting on phenology, susceptibility to *Phytophthora cinnamomi* and hormone profile of chestnut. *Scientia Horticulturae* **311**: 111789.

**Caruso CM, Eisen K, Case AL**. **2016**. An angiosperm-wide analysis of the correlates of gynodioecy. *International Journal of Plant Sciences* **177**: 115–121.

**Charlesworth D**. **1985**. Distribution of dioecy and self-incompatibility in angiosperms. In: Greenwood PJ, Slatkin M, eds. Evolution: Essays in honour of John Maynard Smith. Cambridge: Cambridge Univ. Press, 237–268.

**Charlesworth D**. **1993**. Why are unisexual flowers associated with wind pollination and unspecialized pollinators? *The American Naturalist* **141**: 481–490.

**Charlesworth B, Charlesworth D**. **1978**. A model for the evolution of dioecy and gynodioecy. *The American Naturalist* **112**: 975–997.

**Darwin C**. **1876**. *The effects of cross and self fertilization in the vegetable kingdom*. London: John Murray.

**Darwin C**. **1877**. *The different forms of flowers on plants of the same species*. London: John Murray.

**Dornier A, Dufaÿ M**. **2013**. How selfing, inbreeding depression, and pollen limitation impact nuclear-cytoplasmic gynodioecy: A model. *Evolution* **67**: 2674–2687.

**Dufaÿ M, Billard E**. **2012**. How much better are females? The occurrence of female advantage, its proximal causes and its variation within and among gynodioecious species. *Annals of Botany* **109**: 505–519.

**Dufaÿ M, Champelovier P, Käfer J, Henry JP, Mousset S, Marais GAB**. **2014**. An angiosperm-wide analysis of the gynodioecy–dioecy pathway. *Annals of Botany* **114**: 539–548.

**Duffy KJ, Johnson SD**. **2011**. Effects of pollen reward removal on fecundity in a self-incompatible hermaphrodite plant: Pollen reward removal and plant fecundity. *Plant Biology* **13**: 556–560.

**Duffy KJ, Johnson SD**. **2014**. Male interference with pollination efficiency in a hermaphroditic orchid. *Journal of Evolutionary Biology* **27**: 1751–1756.

**Duffy KJ, Mdlalose ZM, Johnson SD**. **2021**. Sexual conflict in hermaphroditic flowers of an
African *Aloe*. *International Journal of Plant Sciences* **182**: 238–243.

**Duffy KJ, Patrick KL, Johnson SD**. **2013**. Emasculation increases seed set in the bird-pollinated
hermaphrodite *Kniphofia linearifolia* (Xanthorrhoeaceae): Evidence for sexual conflict? *American*
*Journal of Botany* **100**: 622–627.

**Eckhart VM**. **1992**. Resource compensation and the evolution of gynodioecy in *Phacelia linearis*
(Hydrophyllaceae). *Evolution* **46**: 1313–1328.

**Ellis MF, Sedgley M**. **1993**. Gynodioecy and male sterility in *Eucalyptus leucoxylon* F. Muell.
(Myrtaceae). *International Journal of Plant Sciences* **154**: 314–324.

**van Etten ML, Chang S-M**. **2014**. Frequency-dependent pollinator discrimination acts against
female plants in the gynodioecious *Geranium maculatum*. *Annals of Botany* **114**: 1769–1778.

**Fineschi S, Taurchini D, Villani F, Vendramin GG**. **2000**. Chloroplast DNA polymorphism
reveals little geographical structure in *Castanea sativa* Mill. (Fagaceae) throughout southern
European countries. *Molecular Ecology* **9**: 1495–1503.

**Freeman DC, Doust JL, El-Keblawy A, Miglia KJ, McArthur ED**. **1997**. Sexual specialization
and inbreeding avoidance in the evolution of dioecy. *The Botanical Review* **63**: 65–92.

**Furones-Pérez P, Fernández-López J**. **2009**. Morphological and phenological description of 38
sweet chestnut cultivars (*Castanea sativa* Miller) in a contemporary collection. *Spanish Journal of*
*Agricultural Research* **7**: 829–843.

**Garibaldi LA, Aizen MA, Klein AM, Cunningham SA, Harder LD**. **2011**. Global growth and
stability of agricultural yield decrease with pollinator dependence. *Proceedings of the National*
*Academy of Sciences* **108**: 5909–5914.

**Gibbs PE**. **2014**. Late-acting self-incompatibility – the pariah breeding system in flowering plants.
*New Phytologist* **203**: 717–734.

**Gibson JP, Diggle PK**. **1998**. Flower development and male sterility in *Ocotea tenera* (Lauraceae):
A gynodioecious tropical tree. *International Journal of Plant Sciences* **159**: 405–417.

**Givnish TJ**. **1982**. Outcrossing versus ecological constraints in the evolution of dioecy. *The*
*American Naturalist* **119**: 849–865.

**Gouyon PH, Couvet D**. **1987**. A conflict between two sexes, females and hermaphrodites. In:
Experientia Supplementum. The evolution of sex and its consequences. Birkhäuser, Basel, 245–261.

**Hasegawa Y, Suyama Y, Seiwa K**. **2009**. Pollen donor composition during the early phases of
reproduction revealed by DNA genotyping of pollen grains and seeds of *Castanea crenata*. *The*
*New Phytologist* **182**: 994–1002.

**Hasegawa Y, Suyama Y, Seiwa K**. **2017**. Flowering phenology of a duodichogamous self-
incompatible tree species, *Castanea crenata*. *Japanese Journal of Ecology* **67**: 31–39.

**Husband BC, Schemske DW**. **1996**. Evolution of the magnitude and timing of inbreeding
depression in plants. *Evolution* **50**: 54–70.

**Jensen JLWV**. **1906**. Sur les fonctions convexes et les inégalités entre les valeurs moyennes. *Acta*
*Mathematica* **30**: 175–193.

**Johnson SD, Butler HC, Robertson AW**. **2019**. Breeding systems in *Cyrtanthus*
(Amaryllidaceae): Variation in self-sterility and potential for ovule discounting. *Plant Biology* **21**:
1008–1015.

**Kaul MLH**. **1988**. *Male sterility in higher plants*. Berlin: Springer.

**Kikuzawa K**. **1989**. Floral biology and evolution of gynodioecism in *Daphne kamtchatica var.*
*jezoensis*. *Oikos* **56**: 196–202.

**Koelewijn HP, van Damme JMM**. **1996**. Gender variation, partial male sterility and labile sex
expression in gynodioecious *Plantago coronopus*. *New Phytologist* **132**: 67–76.

**Koelling VA, Karoly K**. **2007**. Self-pollen interference is absent in wild radish (*Raphanus*
*raphanistrum* , Brassicaceae), a species with sporophytic self-incompatibility. *American Journal of*
*Botany* **94**: 896–900.

**Krebs P, Pezzatti GB, Beffa G, Tinner W, Conedera M**. **2019**. Revising the sweet chestnut
(*Castanea sativa* Mill.) refugia history of the last glacial period with extended pollen and
macrofossil evidence. *Quaternary Science Reviews* **206**: 111–128.

**Krohne DT, Baker I, Baker HG**. **1980**. The maintenance of the gynodioecious breeding system in
*Plantago lanceolata* L. *American Midland Naturalist* **103**: 269–279.

**Lahiani E, Touzet P, Billard E, Dufaÿ M**. **2015**. When is it worth being a self-compatible
hermaphrodite? Context-dependent effects of self-pollination on female advantage in gynodioecious
*Silene nutans*. *Ecology and Evolution* **5**: 1854–1862.

**Lande R, Schemske DW, Schultz ST**. **1994**. High inbreeding depression, selective interference
among loci, and the threshold selfing rate for purging recessive lethal mutations. *Evolution* **48**: 965–
978.

**Larue C**. **2021**. *De la pollinisation à la formation des graines : le cas du châtaignier*. Université de
Bordeaux.

**Larue C, Austruy E, Basset G, Petit RJ**. **2021a**. Revisiting pollination mode in chestnut
(*Castanea* spp.): An integrated approach. *Botany Letters* **168**: 348–372.

**Larue C, Barreneche T, Petit RJ**. **2021b**. An intensive study plot to investigate chestnut tree
reproduction. *Annals of Forest Science* **78**: 90.

**Larue C, Klein E, Petit R**. **2022**. Sexual interference revealed by joint study of male and female
pollination success in chestnut. *Molecular Ecology*.

**Lewis D**. **1941**. Male sterility in natural populations of hermaphrodite plants. The equilibrium
between females and hermaphrodites to be expected with different types of inheritance. *New*
*Phytologist* **40**: 56–63.

**Lloyd DG**. **1974**. Theoretical sex ratios of dioecious and gynodioecious angiosperms. *Heredity* **32**:
11–34.

**Lloyd DG**. **1975**. The maintenance of gynodioecy and androdioecy in angiosperms. *Genetica* **45**: 325–339.

**Lloyd DG**. **1982**. Selection of combined versus separate sexes in seed plants. *The American Naturalist* **120**: 571–585.

**Lloyd DG, Webb CJ**. **1986**. The avoidance of interference between the presentation of pollen and stigmas in angiosperms I. Dichogamy. *New Zealand Journal of Botany* **24**: 135–162.

**Mather K**. **1940**. Outbreeding and separation of the sexes. *Nature* **145**: 484–486.

**Obeso JR**. **2002**. The costs of reproduction in plants. *New Phytologist* **155**: 321–348.

**Olson MS, Hamrick JL, Moore R**. **2016**. Breeding systems, mating systems, and genomics of gender determination in angiosperm trees. In: Groover A, Cronk Q, eds. Comparative and Evolutionary Genomics of Angiosperm Trees. Cham: Springer International Publishing, 139–158.

**Pannell JR**. **2018**. Transitions between combined and separate sexes in flowering plants. In: Leonard JL, ed. Transitions between sexual systems. Cham: Springer.

**Penagos Zuluaga JC, Queenborough SA, Comita LS**. **2020**. Flowering sex ratios and costs of reproduction in gynodioecious *Ocotea oblonga* (Lauraceae). *Biological Journal of the Linnean Society* **131**: 344–355.

**Pereira-Lorenzo S, Ramos-Cabrer AM**. **2004**. Chestnut, an ancient crop with future. In: Dris R, Jain SM, eds. Production practices and quality assessment of food crops. Preharvest practice. Dordrecht: Springer, 105–161.

**Petit RJ, Carlson J, Curtu AL, Loustau M, Plomion C, González-Rodríguez A, Sork V, Ducousso A**. **2013**. Fagaceae trees as models to integrate ecology, evolution and genomics. *New Phytologist* **197**: 369–371.

**Petit RJ, Hampe A**. **2006**. Some evolutionary consequences of being a tree. *Annual Review of Ecology, Evolution, and Systematics* **37**: 187–214.

**Petit RJ, Larue C**. **2022**. Confirmation that chestnuts are insect-pollinated. *Botany Letters* **169**: 370–374.

**Pettersson MW**. **1992**. Advantages of being a specialist female in gynodioecious *Silene vulgaris* s.l. (Caryophyllaceae). *American Journal of Botany* **79**: 1389–1395.

**Philipp M**. **1980**. Reproductive biology of *Stellaria longipes* Goldie as revealed by a cultivation experiment. *New Phytologist* **85**: 557–569.

**van der Pijl L**. **1978**. Reproductive integration and sexual disharmony in floral functions. In: Richards A, ed. The pollination of flowers by insects. London: Academic Press, 79–88.

**R Core Team**. **2013**. R: A language and environment for statistical computing. *R Foundation for Statistical Computing, Vienne, Autriche*.

**Routley MB, Bertin RI, Husband BC**. **2004**. Correlated evolution of dichogamy and self-incompatibility: A phylogenetic perspective. *International Journal of Plant Sciences* **165**: 983–993.

**Sage TL, Bertin RI, Williams EG**. **1994**. Ovarian and other late-acting self-incompatibility systems. In: Williams EG, Clarke AE, Knox RB, eds. Advances in Cellular and Molecular Biology of Plants, vol. 2. Genetic control of self-incompatibility and reproductive development in flowering plants. Dordrecht: Springer, 116–140.

**Sauquet H, Von Balthazar M, Magallón S, Doyle JA, Endress PK, Bailes EJ, Barroso De Morais E, Bull-Hereñu K, Carrive L, Chartier M, *et al.* 2017**. The ancestral flower of angiosperms and its early diversification. *Nature Communications* **8**: 16047.

**Schultz ST**. **2002**. Partial male sterility and the evolution of nuclear gynodioecy in plants. *Genetical Research* **80**: 187–195.

**Seavey SR, Bawa KS**. **1986**. Late-acting self-incompatibility in angiosperms. *The Botanical Review* **52**: 195–219.

**Shykoff JA**. **1992**. Sex polymorphism in *Silene acaulis* (Caryophyllaceae) and the possible role of sexual selection in maintaining females. *American Journal of Botany* **79**: 138–143.

**Sisco PH, Neel TC, Hebard FV, Craddock JH, Shaw J**. **2014**. Cytoplasmic male sterility in interspecific hybrids between American and Asian *Castanea* species is correlated with the American D chloroplast haplotype. *Acta Horticulturae* **1019**: 215–222.

**Solignat G, Chapa J**. **1975**. *La biologie florale du châtaignier*. Paris: Institut National de Vulgarisation pour les Fruits Légumes et Champignons.

**Soylu A**. **1992**. Heredity of male sterility in some chestnut cultivars (*Castanea sativa* Mill.). *Acta Horticulturae* **317**: 181–186.

**Steyn EMA, Robbertse PJ**. **1990**. Fruit production in a morphologically gynodioecious population of *Bequaertiodendron magalismontanum*. *South African Journal of Botany* **56**: 6–10.

**Stone JD, Olson MS**. **2018**. Pollination context alters female advantage in gynodioecious *Silene vulgaris*. *Journal of Evolutionary Biology* **31**: 111–122.

**Sun M, Ganders FR**. **1986**. Female frequencies in gynodioecious populations correlated with selfing rates in hermaphrodites. *American Journal of Botany* **73**: 1645–1648.

**Thomas SC**. **2011**. Age-related changes in tree growth and functional biology: The role of reproduction. In: Meinzer FC, Lachenbruch B, Dawson TE, eds. Size- and Age-Related Changes in Tree Structure and Function. Dordrecht: Springer, 33–64.

**Thomson JD, Barrett SCH**. **1981**. Selection for outcrossing, sexual selection, and the evolution of dioecy in plants. *The American Naturalist* **118**: 443–449.

**Varga S, Soulsbury CD**. **2020**. Environmental stressors affect sex ratios in sexually dimorphic plant sexual systems. *Plant Biology* **22**: 890–898.

**Vaughton G, Ramsey M**. **2010**. Floral emasculation reveals pollen quality limitation of seed output in *Bulbine bulbosa* (Asphodelaceae). *American Journal of Botany* **97**: 174–178.

**Wang H, Barrett SCH, Li X, Niu Y, Duan Y, Zhang Z, Li Q**. **2021**. Sexual conflict in protandrous flowers and the evolution of gynodioecy. *Evolution* **75**: 278–293.

**Waser NM, Price MV**. **1991**. Reproductive costs of self-pollination in *Ipomopsis aggregata*
(Polemoniaceae): Are ovules usurped? *American Journal of Botany* **78**: 1036–1043.

**Webb CJ**. **1999**. Empirical studies: Evolution and maintenance of dimorphic breeding systems. In:
Geber MA, Dawson TE, Delph LF, eds. Gender and Sexual Dimorphism in Flowering Plants.
Berlin: Springer, 61–95.

**Webb CJ, Lloyd DG**. **1986**. The avoidance of interference between the presentation of pollen and
stigmas in angiosperms II. Herkogamy. *New Zealand Journal of Botany* **24**: 163–178.

**Wheeler B, Torchiano M**. **2016**. lmPerm: Permutation Tests for Linear Models. *R package version*
*2.1. 0*.

**Wickham H**. **2016**. Programming with ggplot2. In: Use R! ggplot2. Cham: Springer, 241–253.

**Willson MF**. **1979**. Sexual selection in plants. *The American Naturalist* **113**: 777–790.

**Wise MJ, Vu JV, Carr DE**. **2011**. Potential ecological constraints on the evolution of gynodioecy
in *Mimulus guttatus* : Relative fecundity and pollinator behavior in a mixed-sex population.
*International Journal of Plant Sciences* **172**: 199–210.

**Xiong H, Zou F, Guo S, Yuan D, Niu G**. **2019**. Self-sterility may be due to prezygotic late-acting
self-incompatibility and early-acting inbreeding depression in Chinese chestnut. *Journal of the*
*American Society for Horticultural Science* **144**: 172–181.

**Yampolsky C**. **1922**. Distribution of sex forms in the phanerogamic flora (H Yampolsky and C
Yampolsky, Eds.). *Bibliotheca Genetica* **3**: 1–62.

**Zhao Z, Liu K**. **2009**. Effect of chemical thinning catkins on Chinese chestnut yield and quality.
*Acta Horticulturae* **844**: 457–460.

## LIST OF FIGURES

**Figure 1:** Degrees of male-sterility observed. A) Fully male-sterile (i.e. female) tree. Most stamens
are aborted and do not produce pollen. B) Mostly male-sterile tree. Stamens do not protrude from
glomerules. C) Partly male-sterile tree. Stamens slightly protrude from glomerules. D) Fully male-
fertile (i.e. cosexual) tree. Stamen filaments are long and largely protrude from glomerules. These
trees produce large amounts of pollen.

**Figure 2:** Chestnuts have two types of catkins: numerous unisexual male catkins that flower first
and few bisexual catkins located at the tip of the branches that flower last.

**Figure 3:** Inter-annual variation in fruit set in cosexual and female chestnut trees.

# SUPPLEMENTARY MATERIALS

Supplementary material 1: Map of the chestnut experimental plot 1.

○ Female
● Cosexual

832    Supplementary material 2: Estimation of gender polymorphism in chestnut forests

| Population name | Male sterile | Partly male sterile | Male fertile | #Trees | MS[1] | PMS[2] | MF[3] |
|---|---|---|---|---|---|---|---|
| Argelès Massane | 0 | 0 | 30 | 30 | 0.00 | 0.00 | 1.00 |
| Amorochilo | 0 | 0 | 32 | 32 | 0.00 | 0.00 | 1.00 |
| Bidegainea | 0 | 0 | 30 | 30 | 0.00 | 0.00 | 1.00 |
| Caminarteko Erreka | 0 | 0 | 26 | 26 | 0.00 | 0.00 | 1.00 |
| Forêt des Cars | 2 | 1 | 27 | 30 | 0.07 | 0.03 | 0.90 |
| Forêt Beauregard | 1 | 0 | 29 | 30 | 0.03 | 0.00 | 0.97 |
| Foret privée près de Marval | 4 | 0 | 26 | 30 | 0.13 | 0.00 | 0.87 |
| Foret privée de Beaussac | 1 | 2 | 27 | 30 | 0.03 | 0.07 | 0.90 |
| Ceret | 6 | 1 | 23 | 30 | 0.20 | 0.03 | 0.77 |
| Puig de l'Esterel | 3 | 4 | 23 | 30 | 0.10 | 0.13 | 0.77 |
| Saint Marsal | 6 | 1 | 23 | 30 | 0.20 | 0.03 | 0.77 |
| Fragas do Eume | 2 | 4 | 36 | 42 | 0.05 | 0.10 | 0.86 |
| Massif Albères | 11 | 5 | 14 | 30 | 0.37 | 0.17 | 0.47 |
| Montagne Noire Castans | 9 | 1 | 20 | 30 | 0.30 | 0.03 | 0.67 |
| Total | 45 | 19 | 366 | 430 | 0.11 | 0.04 | 0.85 |

833    [1]Proportion of male-sterile trees in the studied population

834    [2]Proportion of partly male-sterile trees in the studied population

835    [3]Proportion of male-fertile trees in the studied population

836
837

839     Supplementary material 3: Gender effect on female fitness components by species

| Fitness component | Species | #♀[1] | #♀[1] | Cosexual[2] | Female[2] | Test[3] |
|---|---|---|---|---|---|---|
| Basal area (cm²) | *C. sativa* | 56 | 49 | 283 (186) | 261 (225) | ns (Permanova) |
| | hybrids | 38 | 20 | 341 (212) | 479 (275) | ns (Permanova) |
| Flower density | *C. sativa* + hybrids | 9 | 9 | 51 (34) | 67 (80) | ns (Student) |
| Burr set | *C. sativa* + hybrids | 9 | 9 | 0.77 (0.15) | 0.87 (0.14) | ns (Student) |
| Fruit set | *C. sativa* | 46 | 38 | 0.58 (0.22) | 0.86 (0.10) | $< 10^{-15}$ (Permanova) |
| | hybrids | 35 | 19 | 0.59 (0.18) | 0.83 (0.11) | $< 10^{-15}$ (Permanova) |
| Fruit weight (g) | *C. sativa* | 46 | 38 | 10.5 (3.8) | 8.6 (4.2) | 0.03 (Permanova) |
| | hybrids | 35 | 19 | 12.0 (7.3) | 6.1 (4.3) | 0.06 (Permanova) |

840     [1]Number of cosexual (♀) and female (♀) trees

841     [2]Mean value for cosexual and female trees (standard deviation)

842     [3]Difference significance (test performed)

844 Supplementary material 4: Fruit set measured during four consecutive years and details for 2020

| Clone | Ramet[1] | Species[2] | Gender[3] | 2018 | 2019 | 2020 | 2022 | #♀-inflo[4] | #Burrs[5] | Burr set[6] |
|---|---|---|---|---|---|---|---|---|---|---|
| | | | | \multicolumn Fruits set | | | | \multicolumn Details for 2020 | | |
| Dauphine | A_64_C | S | F | 0.63 | 0.42 | 0.23 | 0.62 | 42 | 33 | 0.79 |
| Dauphine | E_76_I | S | F | 0.76 | 0.93 | 0.67 | 0.85 | 5 | 3 | 0.60 |
| Précoce des Vans | E_40_R | S | F | 0.96 | 0.92 | 0.95 | 0.95 | 57 | 46 | 0.81 |
| Précoce des Vans | E_72_S | S | F | 0.89 | 0.82 | 0.92 | 0.82 | 25 | 22 | 0.88 |
| Merle | A_45_E | S | F | 0.85 | 0.50 | 0.26 | NA | | | |
| Bétizac | A_58_O | SC | F | 0.97 | 0.92 | 0.89 | 0.94 | 16 | 17 | 1.06 |
| Bétizac | E_80_E | SC | F | NA | 0.92 | 0.94 | NA | 42 | 40 | 0.95 |
| Marlhac | A_66_Q | SC | F | 0.80 | 0.75 | 0.62 | 0.91 | 254 | 230 | 0.91 |
| Marlhac | E_73_D | SC | F | 0.77 | 0.75 | 0.72 | 0.89 | 96 | 88 | 0.92 |
| 381 | E_43_O | S | C | 0.63 | 0.40 | 0.28 | 0.18 | 46 | 40 | 0.87 |
| 381 | E_52_J | S | C | 0.67 | 0.47 | 0.29 | 0.28 | 32 | 28 | 0.88 |
| Despont n3 | A_65_F | S | C | 0.58 | 0.47 | 0.39 | 0.35 | 49 | 35 | 0.71 |
| Despont n3 | E_78_E | S | C | 0.67 | 0.57 | 0.62 | 0.35 | 15 | 13 | 0.87 |
| Maridonne | A_60_Q | SC | C | 0.64 | 0.51 | 0.28 | 0.56 | 128 | 63 | 0.49 |
| Maridonne | E_50_P | SC | C | 0.73 | 0.75 | 0.62 | NA | 63 | 47 | 0.75 |
| Marigoule | A_42_N | SC | C | 0.34 | 0.29 | 0.17 | 0.27 | 44 | 42 | 0.95 |
| Marigoule | A_44_Q | SC | C | 0.63 | 0.59 | 0.29 | 0.36 | 29 | 19 | 0.66 |
| Vignols | A_55_S1 | SC | C | 0.84 | 0.40 | 0.14 | NA | | | |
| | | | Female | 0.83 | 0.77 | 0.69 | 0.86 | 67.1 | 59.9 | 0.86 |
| | | | Cosexual | 0.64 | 0.49 | 0.34 | 0.34 | 50.8 | 35.9 | 0.77 |
| | | | **Ratio F/C** | **1.30** | **1.56** | **2.01** | **2.54** | **1.32** | **1.67** | **1.12** |
| | *t-test* | | *Gender* | **0.003** | **0.002** | **0.004** | **<10⁻⁶** | **ns** | **ns** | **ns** |
| | | | *Species* | **ns** | **ns** | **ns** | **ns** | **ns** | **ns** | **ns** |

[1]Identification code of each studied tree (ramet)

[2]Species: *C. sativa* (S) / *C. sativa* × *C. crenata* (SC)

[3]Gender: female (F) / cosexual (C)

[4]Number of female inflorescences per tree (10 branches per tree)

[5]Number of burrs per tree counted in August (10 branches per tree)

[6]Burr set calculated by dividing the number of burrs by the number of female inflorescences

Supplementary material 5: Fruit set following partial and complete emasculation

| Gender[1] | Clone | Modality | #trees[2] | Mean |
|---|---|---|---|---|
| F | Bellefer | Control | 5 | 0.91 |
| F | Bellefer | Bisexual | 2 | 0.83 |
| F | Bellefer | Unisexual | 2 | 0.87 |
| F | Bellefer | Complete | 5 | 0.81 |
| C | Florifer | Control | 6 | 0.46 |
| C | Florifer | Bisexual | 2 | 0.39 |
| C | Florifer | Unisexual | 2 | 0.49 |
| C | Florifer | Complete | 6 | 0.62 |

855    [1]Gender of studied trees: F = Female, C = cosexual
856    [2]Number of studied trees in each modality
857