1

Diversity, distribution, and expression of opsin genes in freshwater lakes

Shaomei He[1,2], Alexandra M. Linz[1], Sarah L.R. Stevens[1], Patricia Q. Tran[1,3], Francisco Moya-

Flores[4], Ben O. Oyserman[4], Jeffrey R. Dwulit-Smith[1,5], Katrina T. Forest[1,5], Katherine D.

McMahon[1,4]

[1]Department of Bacteriology, University of Wisconsin-Madison, Madison, WI, USA

[2]Department of Geoscience, University of Wisconsin-Madison, Madison, WI, USA

[3]Department of Integrative Biology, University of Wisconsin-Madison, Madison, WI, USA

[4]Department of Civil and Environmental Engineering, University of Wisconsin-Madison,

Madison, WI, USA

[5]Program in Biophysics, University of Wisconsin-Madison, Madison, WI, USA

Corresponding Author

Katherine D. McMahon (trina.mcmahon@wisc.edu)

Running Title: Opsin genes in freshwater

ABSTRACT

Microbial opsin genes are widely distributed in aquatic environments and may significantly contribute to phototrophy and energy budgets in global oceans. However, the study of freshwater opsin genes has been largely limited. Here, we explored Type I opsin gene diversity, ecological distribution, and expression in lakes with contrasting physicochemical and optical characteristics. Using metagenomes and derived draft genomes from a clearwater lake and a humic bog, we recovered opsin genes from Actinobacteria, Proteobacteria, Bacteroidetes, Verrucomicrobia, Planctomycetes, Chloroflexi, and Candidatus Saccharibacteria. Most were predicted to encode green light-absorbing proton pumps within the proteorhodopsin and xanthorhodopsin groups, with the humic bog harboring much less diversity. Notably, viral opsin and novel actinobacterial opsin genes phylogenetically significantly different from the typical actinorhodopsin were recovered. Analyses of the surface layer of 33 freshwater systems revealed an inverse correlation between opsin gene abundance and lake dissolved organic carbon (DOC). In humic water with high terrestrial DOC and light-absorbing humic substances, opsin gene abundance was low and dramatically declined within the first few meters, whereas the abundance remained relatively high along the bulk water column in clearwater lakes with low DOC. These results suggest opsin gene distribution is influenced by lake optical properties and DOC. Gene expression analysis confirmed the significance of opsin-based phototrophy in clearwater lakes and revealed different diel expressional patterns among major phyla. Overall, our analyses recovered novel diversity, revealed distribution patterns, confirmed opsin gene expression, and suggested the significance of rhodopsin-based phototrophy in freshwater energy budgets, especially in clearwater lakes.

INTRODUCTION

Microbial rhodopsin is a retinal-activated photosystem. In marine environments, microbial rhodopsins are found in many diverse prokaryotes and exhibit a rich genetic and functional diversity (McCarren & DeLong 2007; Olson *et al.* 2018; Rozenberg *et al.* 2021; Rusch *et al.* 2007). As microbial rhodopsins often occur in more than half of the cells in a wide range of marine samples (Finkel *et al.* 2013; Maresca *et al.* 2018; Olson *et al.* 2018; Sieradzki *et al.* 2018), rhodopsin-based phototrophy has been suggested to be a prevalent form of phototrophy and may significantly influence energy budgets in the global oceans (Finkel *et al.* 2013; Gomez-Consarnau *et al.* 2019), although whether the harvested sunlight is used for growth, survival or other physiological and ecological benefits may differ among different bacteria (Pinhassi *et al.* 2016). Contrasting to the rich knowledge regarding marine microbial rhodopsins, the diversity, ecological distribution, and significance of rhodopsins in freshwater have been much less studied, particularly rhodopsins in humic aquatic environments with high terrestrially-derived allochthonous organic carbon, dimly lit water columns, and often acidic pH, are largely unexplored.

Being a simple photosystem, a rhodopsin consists of an opsin, which is an integral membrane protein with seven transmembrane helices, and a covalently bound retinal as a light-absorbing chromophore. Microbial (also termed Type I, as compared to Type II in animals) rhodopsins play important roles in microbial physiology, with known functions including light-driven ion transporters (e.g. proton, H^+, chloride, Cl^-, and sodium, Na^+ pumps), cation channels, and phototaxis sensors (Rozenberg *et al.* 2021). More recently, a novel family of microbial

rhodopsins distinct from Type I rhodopsins was discovered through functional metagenomics

and named heliorhodopsin (HeR) (Pushkarev *et al.* 2018). As HeRs are only distantly related to

Type I rhodopsins and their functions are largely unknown, we solely focused on Type I

rhodopsins in our study, although we note that HeRs have been found in diverse monoderm

bacteria in freshwater environments (Chazan *et al.* 2022; Flores-Uribe *et al.* 2019).

The first microbial opsin genes in freshwater were discovered through metagenomic studies,

where the predominant opsin genes were recovered from Actinobacteria and therefore named

"actinorhodopsin" (ActR) (Sharma *et al.* 2009; Sharma *et al.* 2008). Phylogenetically, ActR

belongs to the xanthorhodopsin group (Vollmers *et al.* 2013) and functions as an outward H^+

pump (Dwulit-Smith *et al.* 2018; Keffer *et al.* 2015a). In addition to ActR, opsin gene fragments

were also recovered from a few other phyla in freshwater lakes (Atamna-Ismaeel *et al.* 2008;

Galachyants *et al.* 2021; Martinez-Garcia *et al.* 2012; Sharma *et al.* 2009).

Rhodopsin has been shown to enhance bacterial growth and survival under carbon-limited

conditions when exposed to light (Gómez-Consarnau *et al.* 2010; Gómez-Consarnau *et al.* 2007;

Johnson *et al.* 2010; Steindler *et al.* 2011; Yoshizawa *et al.* 2014), suggesting the importance of

light and carbon availability in the regulation of this photoreceptive protein. In freshwater, lake

optical properties and organic carbon quality/quantity are often correlated, and the former can be

greatly impacted by the latter (Maizel *et al.* 2017; Read & Rose 2013). Therefore, an interesting

yet unexplored question is whether the distribution of opsin genes in freshwater varies with

different lake optical properties and organic carbon quality/quantity.

In this study, we surveyed two contrasting freshwaters, a eutrophic clearwater lake and a humic

brown-water bog lake with distinct physicochemical characteristics. In particular, the humic bog

has much higher concentrations of dissolved organic carbon (DOC), most of which is

allochthonous (entering the aquatic system from its catchment). Such terrestrially-derived DOC

comprises light-absorbing humic substances which stain the water brown, and thus the bog

exhibits very different optical properties from the eutrophic lake, which is "clear" as opposed to

"stained" (Maizel *et al.* 2017). To investigate the opsin gene diversity, we used metagenomic

sequences and derived draft genomes obtained from the co-assembly of a large number of

metagenomes collected from a time-series study of these two lakes (Linz *et al.* 2018). To further

study the relationship between opsin gene distribution and freshwater DOC and optical

properties, we included metagenomes from a total of 33 freshwater lakes around the globe that

span a wide range of DOC levels and clarity. We estimated the relative opsin gene abundance in

the surface layer of these lakes, as well as depth-resolved opsin abundance profiles from a subset

of these lakes. Lastly, we explored opsin gene expression and diel regulation in the epilimnion of

two clearwater lakes and the humic bog. Overall, our analysis recovered a greater than

previously appreciated diversity of freshwater opsin genes, revealed the importance of DOC and

optical properties in opsin distribution, and suggested the significance of rhodopsin-based

phototrophy in clearwater lakes.

MATERIALS AND METHODS

Lake Mendota and Trout Bog metagenomes. Two contrasting temperate lakes in Wisconsin,

USA were studied in detail. Lake Mendota (43.10^{o}N, 89.41^{o}W) is a eutrophic clearwater lake

with an average DOC of 5.0 mg/L. Trout Bog (46.01^{o}N, 89.61^{o}W) is a small humic lake, with an

average DOC of 20.0 mg/L (data from NTL-LTER, averaged over the course of the study). More

detailed lake physicochemical and limnological characteristics are listed in **Table S1**. For Lake

Mendota, 0 to 12 m depth-integrated water samples (mostly consisting of the epilimnion layer)

were collected at 94 time points from 2008 to 2012, and samples are referred to as "ME". For

Trout Bog, the integrated epilimnion layer and the integrated hypolimnion layer were collected at

45 time points respectively from 2007 to 2009, and samples were referred to as "TE" and "TH",

respectively.

Details of metagenome sequencing, assembly, and binning were previously described (Bendall *et*

al. 2016; Linz *et al.* 2018). Briefly, shotgun metagenome libraries were constructed from each of

the 184 DNA samples. Assembly was performed on individual metagenomes. In addition, three

"combined" assemblies were generated by co-assembling reads from all metagenomes within the

ME, TE, and TH groups, respectively. Binning was conducted on the three combined assemblies

to obtain metagenome-assembled genomes (MAGs), and this resulted in 486 genome bins. By

further fine-tuning to remove outlier contigs, and excluding bins with high contamination (>10%)

or low completeness (<50%), a total of 202 high-quality MAGs (referred to henceforth as MAGs

to distinguish them from the remaining bins which have lower qualities) were generated (Linz *et*

al. 2018). Taxonomic assignment of MAGs and bins was conducted using PhyloSift based on 37

conserved phylogenetic marker genes (Darling *et al.* 2014). The 184 individual metagenome

assemblies, the three combined assemblies, and recovered MAGs/bins were submitted to the

DOE Joint Genome Institute's Integrated Microbial Genome (IMG) database for gene prediction

and function annotation (Markowitz *et al.* 2013).

Opsin phylogenetic tree reconstruction and inference of taxonomy. Opsin sequences were identified by using pfam01036 (http://pfam.xfam.org). Representative opsin reference sequences were chosen by clustering bacterial and archaeal opsin protein sequences from isolate genomes at IMG based on 90% amino acid sequence identity using the UCLUST algorithm provided by the USEARCH tool (Edgar 2010). Representative sequences from each cluster, opsins from previously sequenced freshwater single-amplified genomes (SAGs) and MAGs, and viral opsin sequences from the study by Yutin and Koonin (Yutin & Koonin 2012) were included in the phylogenetic analysis, together with the 317 opsin sequences longer than 200 amino acids (which is ~80% of rhodopsin full length) in our three combined assemblies (including sequences in MAGs/bins and the remaining un-binned). Amino acid sequences were aligned using MUSCLE (Edgar 2004), and the alignment was trimmed to exclude columns that contained gaps for more than 30% of the included sequences. Poorly aligned positions and divergent regions were further eliminated using Gblocks (Castresana 2000). A maximum likelihood phylogenetic tree was constructed using PhyML 3.0 (Guindon *et al.* 2010), with the LG substitution model (Le & Gascuel 2008) and the gamma distribution parameter estimated by PhyML. Bootstrap values were calculated based on 100 replicates. After constructing this phylogenetic tree, sequences shorter than 200 amino acids in the three combined assemblies were added to the existing tree using PPlacer (Matsen *et al.* 2010). The resulting phylogenetic tree was used for taxonomic classification of un-binned metagenome opsin sequences at the phylum level.

Estimation of opsin gene relative abundance. The abundance of an individual gene in an assembled metagenome was estimated using its average per-base coverage, which indicates how many times a base was sequenced and was determined by mapping metagenome reads to

assembled contigs. This estimate is available in the IMG database as "read depth", and we

referred to it as "coverage-weighted abundance" in this study to reflect the abundance of the

population that contributed to this gene in the metagenome. To compare opsin gene abundance

among different metagenomes, the opsin gene relative abundance was estimated by the sum of

the coverage-weighted abundance of all opsin genes in the metagenome, and further normalized

(i.e. divided) by the average of the coverage-weighted abundance of three single-copy conserved

housekeeping genes (represented by pfam00121, pfam00573, pfam00687, http://pfam.xfam.org)

in the same metagenome. The three single-copy genes were chosen because they have

comparable hidden Markov model (HMM) lengths to opsin genes to minimize the bias in gene

abundance estimates associated with gene lengths (He *et al.* 2015).

Estimation of opsin gene abundance for additional lakes. To further study opsin gene

distribution, we included metagenomes from 31 additional freshwater lakes for which organic

carbon concentrations were available. Among these additional lakes, 18 are from the

stratfreshDB deposited at European Nucleotide Archive (Buck *et al.* 2021) and 13 are from

publicly available metagenomes at IMG, including two unpublished from our lab and 11 from

published studies (Dam *et al.* 2020; Denef *et al.* 2016; Linz *et al.* 2020; Podowski *et al.* 2021;

Tran *et al.* 2018; Tran *et al.* 2021) (**Tables S2 and S3**). Together with Mendota and Trout Bog,

the 33 lakes span a wide range of DOC levels and exhibit different clarity (examples of clarity

data are listed in **Table S1**). Some of these lakes have samples collected from different depths,

allowing the study of opsin gene vertical distribution (**Table S4**). For metagenomes available at

IMG, the relative abundance of opsin genes was estimated as described above. For metagenomes

from the stratfreshDB, we performed gene prediction using Prodigal and then HMMsearch

(HMMER version 3.3) using the HMM models for pfam01036, pfam00121, pfam00573, and

pfam00687 to identify opsin and the three single-copy conserved genes respectively. Coverage

information was obtained by mapping metagenome reads to corresponding metagenome

assemblies using bowtie2 (version 2.2.2) with default settings. Coverage-weighted abundance

from these genes was used to estimate opsin gene abundance as described earlier. The in-house

bioinformatics workflow that we used on the stratfreshDB metagenomes, when used on a subset

of the included IMG metagenomes, yielded results very comparable to results obtained from the

IMG pipeline. Therefore, no systemic bias was associated with different bioinformatic

workflows.

Analysis of opsin gene expression and diel trends. We used metatranscriptomes from the

GEODES (Gene Expression in Oligotrophic, Dystrophic, and Eutrophic Systems) dataset (Linz

et al. 2020) to study the expression of opsin genes. This dataset includes triplicate samples

collected every four hours for two days from Lake Mendota and Sparking Lake, an oligotrophic

clearwater lake in Northern Wisconsin (**Table S1**). Triplicate samples were similarly collected

from Trout Bog but samples from the second day did not yield sufficient RNA for sequencing.

Notably, an internal standard, an *in vitro* transcribed pFN18A of a known concentration was

added to each sample after the bead beating step during the RNA extraction (Linz *et al.* 2020).

This type of internal standard enables absolute quantification of transcript abundance as copies

of transcripts per litter of sample and thus allows comparing transcript abundance across

different samples (Gifford *et al.* 2011). As two metagenomes were collected along with the

metatranscriptomes for each lake, we first clustered opsin sequences from the two replicate

metagenomes with 97% nucleotide identity using the UCLUST algorithm provided by the

USEARCH tool (Edgar 2010) to remove redundancy, followed by removing sequences shorter

than 600 bp. The resulting opsin sequences from Sparkling Lake were used as references for

mapping Sparkling metatranscriptome reads. For Mendota and Trout Bog, the resulting

GEODES opsin sequences from the above steps were compared to opsin sequences longer than

600 bp from the Mendota and Trout Bog combined assembly respectively by BLASTN to

identify GEODES sequences not represented in the corresponding combined assembly as defined

by <97% nucleotide identity. The identified GEODES sequences were added to the >600 bp

opsin sequences from the corresponding combined assembly to generate the mapping reference

for Mendota and Trout Bog respectively. Metatranscriptome reads processed to remove rRNA

sequences from Linz *et al*. (2020) were mapped to corresponding opsin references with the

internal standard sequence by using bowtie2 (version 2.2.2) with default settings.

Metatranscriptomes with low read counts for the internal standard were discarded according to

Linz *et al.* (2020). Opsin transcript read counts were normalized using the read count from the

internal standard and converted to transcript concentrations. RAIN test (Thaben & Westermark

2014) was conducted to detect genes with diel cyclic trends with *p*-values of less than 0.05 for

cyclic genes.

RESULTS

1. Diversity of freshwater opsin genes

Overview

Here, we studied opsin gene diversity using three combined metagenome assemblies, generated

by co-assembling reads from all metagenomes within the ME (Mendota Epilimnion, 94

metagenomes), TE (Trout Bog Epilimnion, 45 metagenomes), and TH (Trout Bog Hypolimnion,

45 metagenomes) groups (Linz *et al.* 2018). Altogether, from the three combined assemblies,

opsin genes were recovered in 42 MAGs (**Table 1**) and 50 bins that did not pass our criteria for

quality MAGs. Genome completeness/contamination assessments and taxonomic classification

of these MAGs and bins, and their opsin gene IMG IDs are provided in **Table S5**. At the phylum

level, the classification of these opsin genes through classifying their resident MAG/bins was

largely consistent with the IMG classification of opsin gene-containing contigs based on the

consensus BLASTP hits for all genes on the contig (**Table S5**). Therefore, we assumed that

opsin genes from these MAGs/bins were suitable phylogenetic anchors to classify metagenome

opsin sequences in un-binned contigs to at least the phylum level.

We constructed a phylogenetic tree using references from microbial isolates and previously

published freshwater SAGs, together with opsin genes longer than 200 amino acids in the three

combined assemblies, which include sequences from MAGs/bins and un-binned contigs (**Figure

1**, with a rectangular view of this tree with more detailed gene information in **Figure S1**).

Metagenome opsin sequences shorter than 200 amino acids were aligned and added to the

phylogenetic tree (**Figure S2**). Based on the coverage-weighted opsin gene abundance in the

assemblies, we estimated the percent distribution of opsin genes among different taxa and

rhodopsin types in each lake/layer (**Figure S3**).

The phylogenetic analysis suggested Mendota opsin genes were present in all major freshwater

phyla except Cyanobacteria, including Actinobacteria, Proteobacteria, Bacteroidetes,

Planctomycetes, Verrucomicrobia, and Chloroflexi, and the first three phyla were the major

contributors to opsin gene abundance (**Figure S3**). Whereas in Trout Bog, opsin genes were

restricted to fewer taxa, mainly from Actinobacteria (acI-B2 in particular) and some from

Proteobacteria and *Candidatus* Saccharibacteria (TM7) (**Figure S3**).

The phylogenetic analysis also indicated that opsin genes mostly belong to the proteorhodopsin

(PR) and xanthorhodopsin (XR) branches, and genes outside of these two branches had very low

abundances in Mendota and Trout Bog (**Figures 1 and S3**). With some exceptions (discussed in

Supplementary Text), they have the conserved DTE (aspartic acid, threonine, glutamic acid)

motif corresponding to Positions 85, 89, and 96 (BR numbering) for H^+ accepting and donation

(Beja & Lanyi 2014). Both PR and XR are outward H^+ pumps and subgroup I of XR also has a

secondary chromophore, a noncovalently bound carotenoid functioning as a light-harvesting

antenna, in addition to the retinal (Balashov *et al.* 2005; Béjà *et al.* 2000; Imasheva *et al.* 2009;

Vollmers *et al.* 2013). XR sequences in the two lakes are mainly ActR, and nearly all ActR

sequences recovered in our study contain the signature glycine that replaces a bulky residue on

helix 5, which permits binding of the carotenoid antenna salinxanthin in the well-characterized

XR from *Salinibacter ruber* (Balashov *et al.* 2005). Whether any individual ActR-encoding

microbe synthesizes a carotenoid antenna and whether particular ActR exemplars covalently

bind a secondary carotenoid may need to be addressed on a case-by-case basis but the presence

of the signature glycine residue and the genetic machinery for a secondary carotenoid

biosynthesis have been noted (Dwulit-Smith *et al.* 2018), and in at least one case the binding of a

native actinobacterial carotenoid to an ActR has been demonstrated (Chuon *et al.*, 2021).

Opsin gene prevalence and diversity in freshwater Actinobacteria

Actinobacteria was the phylum possessing the largest number of opsin genes, accounting for 46-

61% of all opsin gene coverage in each of the three combined assemblies (**Figure S3**). As

Actinobacteria is one of the best studied freshwater lineages, with many SAGs and MAGs

available, their opsin genes were classified beyond the phylum level (**Figure 2**). In ME, opsin

genes were widely distributed in nearly all lineages/clades of Actinobacteria that were previously

known to harbor opsin genes. The acI lineage, particularly Ca. Planktophila (acI-A) and

Nanopelagicus (acI-B), was the most prominent contributor to opsin genes in ME (**Figure S3**),

and this was expected as members of this lineage are known to be the most abundant bacterial

group in ME (Garcia *et al.* 2017; Newton & McMahon 2011). Although acI was also the most

abundant contributor to opsin genes in TE and TH, their ActR genes were confined within a

branch of acI-B, likely from acI-B2. This branch exclusively contained opsin sequences from

humic lakes, consistent with the finding that acI-B2 prefers humic lakes as compared to acI-B1

(Garcia *et al.* 2014; Newton *et al.* 2006).

Previously, a rhodopsin group sufficiently phylogenetically distinct from the typical ActR was

discovered in both freshwater and marine actinobacterial MAGs belonging to the order of

Acidimicrobiales and was named acidirhodopsin (short as "acidiR" in our study in contrast to

ActR) (Ghai *et al.* 2014; Mizuno *et al.* 2015). According to the freshwater taxonomy, the acidiR-

containing freshwater MAG (Ghai *et al.* 2014) belongs to the Iluma-B clade (alternative name

acIV-B). We also found acidiR in the Iluma-B clade in ME (**Figures 1** and **2)**, accounting for ~5%

of all opsin gene coverage in the combined ME assembly (**Figure S3**). In addition, we identified

a new branch of opsin genes belonging to the Iluma-A clade (alternative name acIV-A) in ME,

accounting for ~6% of total metagenome reads that mapped to opsin genes. The Iluma-A and

Iluma-B branches did not form a monophyletic group, and instead, the Iluma-A branch is outside

of the larger branch formed by ActR, other known XR, and Iluma-B (**Figures 1 and S1**).

Therefore, we use "acidiR (Iluma-A)" and "acidiR (Iluma-B)" to distinguish the two acidiR

groups. Both acidiR clades have the conserved DTE motif associated with H^+ pumps. These two

clades as well as the Chloroflexi clade which is next to Iluma-A do not contain any known XR

sequences (**Figures 1 and S1**). However, as they are phylogenetically closer to the branch

containing known XR than to any other rhodopsin branches, we tentatively classify them as XR.

Notably, only the Chloroflexi clade and about half of the sequences in the Iluma-A clade have

the signature glycine on helix 5 which is necessary (but not sufficient) for binding the carotenoid

antenna, and therefore rhodopsins in these clades may not have the carotenoid antenna, similar to

the subgroup II members of XR (Vollmers *et al.* 2013).

MAGs from acV (also within Acidimicrobiales) were exclusively from Trout Bog. No opsin

genes (nor retinal biosynthesis genes) were recovered from any of the eight acV MAGs,

probably indicating the absence of rhodopsin-based phototrophy in this humic-specific

Actinobacteria lineage. For more results and discussion about the diversity of actinobacterial

opsin genes, especially genes outside of the ActR-XR branch, see Supplementary Text.

Wider than previously thought opsin gene distribution in Bacteroidetes

Actinobacteria have long been regarded as the major phylum containing opsin genes among

freshwater bacteria. For example, among single cells recovered from freshwater lakes, the six

opsin-containing Sphingobacteriia SAGs were dwarfed by the large number (~100) of opsin-

containing Actinobacteria SAGs (Martinez-Garcia *et al.* 2012). In our study, opsin genes from

Bacteroidetes were exclusively within the PR group, and correspond to MAGs placed in the

Sphingobacteriia, Flavobacteriia, and Cytophagia classes (**Table 1** and **Figure S1**). They were

mostly from low-abundance populations in ME (**Figure 1**), yet when added together, they

accounted for ~13% of all opsin gene coverage in the combined ME assembly (**Figure S3**).

Notably, 14 out of the 35 Actinobacteria MAGs and 21 out of the 45 Bacteroidetes MAGs

contained opsin genes (**Table S6**), suggesting that, similar to Actinobacteria, Bacteroidetes also

widely possess opsin genes, and therefore, could be a previously overlooked photoheterotrophic

lineage in freshwater.

***Candidatus* Fonsibacter (LD12), the major contributor to proteobacterial opsins**

Proteobacteria are abundant and prevalent in freshwater, which is also reflected by the total of 54

Proteobacteria MAGs recovered from the combined assemblies. Yet compared to Actinobacteria

and Bacteroidetes, relatively fewer opsin sequences were recovered from this phylum (**Table S6**).

Proteobacterial opsin genes in our metagenomes belong to the PR and XR groups (**Figures 1** and

S3). Notably, several un-binned metagenome sequences in the PR branch, including one

occurring at an extremely high abundance, fall within the freshwater *Candidatus* Fonsibacter

(LD12) (**Figures 1** and **S1**). LD12 is a sister clade of marine SAR11, which is the major

contributor to rhodopsins in the ocean (Maresca *et al.* 2018; Olson *et al.* 2018; Sieradzki *et al.*

2018). As LD12 are very abundant in Mendota, opsin genes in the LD12 clade alone contributed

to nearly 11% of all opsin gene coverage in the combined ME assembly. The lack of recovered

MAGs affiliated with LD12 may reflect its unique and high within-population microdiversity

(Garcia et al 2018).

Opsin genes in other bacterial taxa

Opsin genes belonging to Planctomycetes, Verrucomicrobia, and Chloroflexi were all from ME and were distributed in the PR and XR groups (**Figure 1**). PR sequences from Planctomycetes and Verrucomicrobia form a cluster, which is part of a larger branch that also includes Firmicutes and Actinobacteria Luna1 and Luna2 opsins (**Figures 1** and **S1**). Sequences within this branch have the DTK variation of the H^+-translocating motif originally identified in the firmicute *Exiguobacterium* spp. (Balashov *et al.* 2013). Our sequences expand the diversity within this unusual PR group.

Opsin genes belonging to *Candidatus* Saccharibacteria (previously candidate division TM7) were recovered from un-binned contigs in Trout Bog. Their sequences are highly similar (~93% amino acid identity) to the opsin gene in a genome bin classified as TM7 (TM7 FNED7-bin-18, IMG genome ID 2648501560) recovered in an enrichment culture originating from the bog-like basin of Grosse Fuchskuhle in Germany (**Figure S1**) (Garcia *et al.* 2014). These opsin genes do not belong to either PR or XR, rather they form a cluster basal to the sensory xenorhodopsin (XeR) group (**Figure S1**), and lack any discernable motif associated with H^+-translocation or the diagnostic conserved aspartic acid to proline substitution in the seventh helix for XeR (Ugalde *et al.* 2011). Therefore they may represent a novel clade and may be of interest to the field of optogenetics in search of new light-responsive proteins for desired properties.

Viral opsin genes

Viral opsin genes were first discovered in giant viruses that infect marine unicellular eukaryotes,

such as Phycodnaviruses and *Phaeocystis globosa* virus in Organic Lake, an Antarctic lake with

extremely salty water (Yutin & Koonin 2012). They were later detected in the Red Sea and were

referred to as "VR" for viral rhodopsins (Philosof & Beja 2013). Protein structure

characterization and functional analyses revealed that they encode light-driven H^+-pumps

(Needham *et al.* 2019) or light-gated ion channels (Bratanov *et al.* 2019; Zabelskii *et al.* 2020).

In our study, we recovered a large opsin gene cluster basal to all bacterial and archaeal branches

in the phylogenetic tree (**Figures 1** and **S1**). Three viral bins (see **Supplementary Text 2** and

Tables S7 and **S8** for their classification as viruses) recovered from ME each contain one to two

copies of opsin genes belonging to this cluster (**Tables 1** and **S5**). This cluster also harbored

opsin sequences from algal viruses previously recovered from marine or salty water (**Figure S1**),

supporting the viral origin of this cluster. These freshwater viral opsin genes also form two

distinct sub-clusters, Viral Group I and II as previously defined (Yutin & Koonin 2012), each

harboring both freshwater and marine sequences (**Figure S1**). The function of these freshwater

viral rhodopsins, their roles in virus-host interactions, and their influences on host physiology

and ecology are yet to be studied.

2. Retinal biosynthesis

Retinal is a photosensitive chromophore cofactor of rhodopsin. One important question is

whether the opsin gene-containing genomes we recovered also have the genetic potential for

retinal biosynthesis to form functional rhodopsin. For this, we checked key genes responsible for

the last few steps in retinal biosynthesis, including phytoene synthesis, phytoene desaturation,

lycopene cyclization, and β-carotene cleavage. We found many opsin-containing MAGs also

have the genetic potential for retinal biosynthesis (**Table 1**). Due to incompleteness, it is difficult

to conclude the lack of genes based on MAGs. However, some patterns are consistent. For

example, all viral bins lacked retinal biosynthesis genes, and therefore, they might rely on their

hosts or the environment for the chromophore. In addition, among Actinobacteria, retinal

biosynthesis genes were mostly found within acI, particularly acI-A and acI-B. This trend is

supported by a total of 74 freshwater Actinobacteria genomes, including SAGs and MAGs from

other studies that span most freshwater Actinobacteria lineages (**Table S9**). Although opsin

genes are widely distributed among these lineages (except acV), oxygenase genes responsible for

β-carotene cleavage are only present in some tribes within the acI-A and acI-B clades (**Table S9**).

Thus, Actinobacteria without the retinal biosynthesis capability need to take up retinal from the

environment if their opsins are to be photoactive. For example, the freshwater Actinobacterium

Rhodoluna lacicola WH-Ta8[T] conducted light-induced H[+] pumping only after exogenous retinal

was provided (Keffer *et al.* 2015a).

3. Predicted light-absorbing spectra

Light absorbing spectra of rhodopsin can be predicted based on the spectral tuning residue at

Position 105 (according to eBAC31A08 numbering), which is hydrophobic residues (e.g.

methionine, M or leucine, L) for absorbing green light, and hydrophilic glutamine (Q) for blue

light (Gómez-Consarnau *et al.* 2007; Man *et al.* 2003). Except for five Proteobacteria sequences

that contain Q, all bacterial and viral opsin sequences in our metagenomes have M or L at this

position, and therefore likely absorb green light. Consistent with such a bioinformatic prediction

for ActR, *in vitro* synthesis and *in vivo* expression of ActR indicated peak absorbance in the

green range (Dwulit-Smith *et al.* 2018; Keffer *et al.* 2015b). The predominance of green-

absorbing rhodopsins agrees with previous studies in freshwater rhodopsins (Atamna-Ismaeel *et*

al. 2008; Ghai *et al.* 2014).

4. Ecological distribution of opsin genes.

We estimated the relative abundance of opsin genes in metagenomes by comparing the coverage-

weighted abundance of opsin genes to single-copy essential housekeeping genes that have

comparable lengths to opsin genes. As opsin genes in our MAGs are mostly single-copy (**Table**

S5) and viral contribution to the total opsin genes is relatively small (**Figure S3**), such estimation

largely reflects the fraction of bacteria that possess opsin genes in the total community. We

performed this estimation using individual metagenomes generated from each time point from

ME, TE, and TH. Comparison between ME and TE revealed that ME had a higher frequency of

opsin gene encoding MAGs within all MAGs from the corresponding co-assembly than TE

(**Figure 3A**), and opsin gene overall abundance was also higher in ME (~70%) than in TE (~14%)

(**Figure 3B**). We therefore hypothesized that opsin gene distribution could be influenced by lake

DOC and optical properties, in particular, that high DOC and light-absorbing humic substances

and thus low light availabilities can lead to the low occurrence of opsin genes, whereas low DOC

and good light penetration may enrich for opsin genes.

Comparison among the surface layers of 33 lakes

To test this hypothesis, we included a total of 33 freshwater systems for which metagenome data

and DOC levels are publicly available (**Table S2**). These lakes range from oligotrophic,

mesotrophic, and eutrophic clearwater (not stained brown) lakes, to dystrophic brown-water

humic lakes, spanning a wide range of DOC levels and exhibiting different clarities. Examples of

lake DOC levels, clarity as reflected in the Secchi depth and light diffuse attenuation coefficient,

and other physicochemical and limnological characteristics are listed in **Table S1**. As opsin gene

abundance can be influenced by water depth as discussed later, only metagenomes collected

from the surface layer were included in this analysis to reduce variables. For lakes from the

stratfreshDB, which largely consists of small boreal lakes, metagenomes collected within the

first meter were included. An inverse correlation (r = -0.77) between opsin gene abundance and

DOC was observed (**Figure 4**). Opsin gene abundance was >50% in clearwater lakes irrespective

of their trophic status. Especially, the abundance was extremely high (~90%) in the Laurentian

Great Lakes (Podowski *et al.* 2021) and Lake Lanier (Dam *et al.* 2020) where DOC levels are

low (~2 mg/L). Opsin genes were much less prominent in humic lakes with high terrestrial DOC.

Extremely low abundance (<10%) was observed in the surface water of Lake Mekkoja☐rvi, a

highly humic boreal lake with ~30 mg/L of DOC (Buck *et al.* 2021).

Comparison at different depths within lakes

Within Trout Bog, the epilimnion had a higher frequency of opsin gene encoding MAGs within

all MAGs from the corresponding co-assembly than the hypolimnion (**Figure 3A**), and

comparing individual metagenomes sampled on the same day also indicates the epilimnion had a

higher opsin gene abundance than the hypolimnion, and the difference between layers is

statistically significant (paired student T-test p-value < 1e-8, n = 29) (**Figure 3B**). Indeed,

analyses of depth-discrete metagenome profiles show a steep decline of opsin gene abundance

within the first few meters in humic lakes, leading to extremely low opsin gene levels in the

deeper layer (**Figure 5**). By contrast, such a sharp decrease was not observed in clearwater lakes,

where a depth-associated change in opsin gene abundance was much more moderate and slower

(**Figure 5**).

5. Opsin gene expression and diel trends

We used metatranscriptomes previously generated from the epilimnia of Mendota, Trout Bog,

and Sparkling Lake by Linz *et al.* (2020) to study opsin gene expression and diel cyclic rhythms.

From metatranscriptome mapping outputs in the original analyses by Linz *et al.* (2020), the total

opsin transcripts are about 0.5%, 0.3%, and 0.02% of the total mRNA for Mendota, Sparkling

Lake, and Trout Bog respectively when averaged from all time points, and notably, a few opsin

genes were among the top 500 most highly expressed genes within the entire metatranscriptomes

in Mendota and Sparkling Lake. This indicates that opsin genes were highly expressed in these

two clearwater lakes.

In our current analysis with more tailored and curated references for opsin genes, we found that

201 out of the 396 opsin genes in the Mendota opsin reference set had expression levels higher

than the cutoff, and these numbers were 105 out of 173 and 18 out of 69 for Sparkling Lake and

Trout Bog respectively. The total opsin transcripts per liter of lake water averaged from all time

points showed that the total expression in the two clearwater lakes was about a magnitude higher

than that in the humic bog (**Figure 6**). Bacterial phyla contributing to the most opsin expression

were Actinobacteria, followed by Bacteroidetes and Proteobacteria in the two clearwater lakes,

whereas opsin expression from Proteobacteria was minimal in the humic bog.

Ranking individual opsin genes by their average transcript abundances indicated that opsin from

Actinobacteria (mostly acI-A and acI-B), Bacteroidetes, and Proteobacteria (LD12 in particular)

are among the most highly expressed opsins in the two clearwater lakes, whereas the most highly

expressed opsins in Trout Bog were mainly from Actinobacteria acI-B (**Table S10**). Notably, the

novel actinobacterial opsins acidiR from acIV-A (Iluma-A) and acIV-B (Iluma-B) were

expressed in the two clearwater lakes.

Among the expressed opsin genes, some exhibited diel trends. Depending on the

photosynthetically active radiation (PAR) measured just below the surface of lake water, samples

were categorized into "day" and "night" groups (Linz *et al.* 2020). Some taxa-specific diel

patterns emerged. Cyclic genes from Actinobacteria tend to have peak expression at night, and

Bacteroidetes opsins tend to peak at daytime in both Mendota and Sparkling Lake (**Figure 7**). In

Trout Bog, only eight genes were identified as cyclically transcribed. Seven of them belong to

Actinobacteria and showed increased expression during the night and peaked before sunrise

(**Figure S4**). However, Trout Bog results need further verification as only one day of data is

available for this lake (Linz *et al.* 2020).

DISCUSSION

Diversity of freshwater opsin genes

Opsin gene taxonomic diversity in freshwater has been previously studied (Atamna-Ismaeel *et al.*

2008; Galachyants *et al.* 2021; Martinez-Garcia *et al.* 2012; Sharma *et al.* 2009). However, some

of these studies were constrained by PCR primer limitations or difficulties for accurate

phylogenetic inference when lacking closely related freshwater reference sequences. In addition,

opsin genes in brown-color, acidic humic lakes had not been explored. In our study, we

overcame these limitations by employing metagenomes and derived MAGs/bins collected from

both clearwater and humic lakes.

Opsin genes were found in more bacterial taxa in the clearwater lake than in the humic bog.

Many opsin genes were recovered from freshwater taxa previously known to harbor opsin genes,

mostly predicted encoding proton pumps belonging to the XR and PR groups. We also found

opsin genes from freshwater viruses and recovered humic lake-specific TM7 opsin sequences

which may represent a novel rhodopsin clade. Among the known taxa with opsin genes, we

found wider than previously thought distribution within freshwater Bacteroidetes and identified

novel actinobacterial opsins that are significantly different from the typical actinorhodopsin in

the acIV-A (Iluma-A) and acTH2 (Myco) clades. Therefore, our analysis expands the taxonomic

spectra and reveals novel diversity of freshwater opsin genes.

Significance of rhodopsins in clearwater lakes

The retinal-activated rhodopsin is a much simpler photosystem compared to multi-component

photochemical reaction centers, such as oxygenic photosystems I and II and anoxygenic reaction

center types I and II, and thus requires less cost to maintain and is expected to be more likely

subjected to horizontal gene transfer (Frigaard *et al.* 2006). Consistent with this, opsin genes are

widely distributed in marine environments where they often occur in more than half of the cells

in a wide range of samples (Finkel *et al.* 2013; Maresca *et al.* 2018; Olson *et al.* 2018). Some

have suggested that opsin genes are more abundant in marine than in freshwater systems (Finkel

et al. 2013; Oh *et al.* 2011). In particular, opsin gene abundance was positively correlated to

salinity (Brindefalk *et al.* 2016; Maresca *et al.* 2018). However, a very limited number of

freshwater systems were included in these studies. The distribution of opsin genes in freshwater,

especially in humic lakes had been essentially unstudied.

We found that in contrast to previous predictions opsin genes are indeed prevalent in clearwater

lakes with low DOC and good light penetration. Opsin gene abundance was usually higher than

60% in lakes with DOC less than 5 mg/L. In particular opsin gene abundance is remarkably high

(~90%) in the Laurentian Great Lakes, where DOC is around 2 mg/L and the terrestrial influence

is small due to the large surface area. This suggests that rhodopsin-based phototrophy is an

important light-harvesting mechanism and its significance may have been underestimated in such

lakes. Contrasting to that, opsin genes appear to be less important overall in humic lakes which

are often associated with a large terrestrial influence, high allochthonous DOC, brown water

color, and a shallow photic zone as discussed in more detail later.

Opsin gene expression and diel regulation

In Mendota and Sparkling Lake, the majority of bacterial opsin genes were expressed, the total

opsin transcripts accounted for 0.3-0.5% of all mRNA, and a few opsin genes were among the

most highly expressed genes in these lakes, supporting their importance in bacterial

ecophysiology in the surface water of clearwater lakes. Although rhodopsins may also be

important in populations that express opsin genes in the bog, the higher total expression of opsin

genes in the two clearwater lakes compared to in the humic bog suggests the overall greater

significance of rhodopsin-based phototrophy in clearwater lakes.

Opsin gene regulation in freshwater had been largely unexplored. In marine systems, some

studies showed that opsin gene expression exhibited diel rhythms, with maximal expression at

the darker periods instead of peak sunlight (e.g. pre-sunrise, post-sunrise, pre-sunset, and post-

sunset) (Ottesen *et al.* 2014; Zehnpfennig *et al.* 2022), whereas others suggested opsin genes

may be constitutively expressed (Riedel *et al.* 2010; Shi *et al.* 2011). In freshwater, an earlier

study of Actinorhodopsin expression indicated their maximal expression at dawn and minimal

expression at dusk for samples collected every six hours (Wurzbacher *et al.* 2012). In our study,

samples were collected every four hours for two days. From both clearwater lakes, we observed

consistent patterns showing diel cyclic behaviors differing among major taxa. Specifically,

Actinobacterial opsin genes tend to peak at darker periods of the day, and Bacteroidetes tend to

peak during the daytime, especially at peak sunlight.

Taxa-specific diel patterns might suggest differences in a circadian-like clock among bacteria.

As opsin gene expression can be regulated by light (Lami *et al.* 2009), carbon and nutrient

availability (Kwon *et al.* 2013; Wang *et al.* 2012), the taxa-specific patterns might also suggest

dual regulation of opsin genes by both light regime and carbon/nutrient availabilities, as the latter

can also experience diel cascade changes following photosynthesis in the lake (Linz *et al.* 2020).

We should note that only a portion of the expressed opsin genes exhibited a statically significant

diel pattern, contributing to about half of all opsin transcripts in the two clearwater lakes (**Figure**

6). It is not clear whether those without cyclic trends are constitutively expressed as some studies

suggested (Keffer *et al.* 2015a; Riedel *et al.* 2010) or if their diel changes are too small to be

detected given the noise in the metatranscriptome data.

Influence of DOC and optical properties on opsin gene distribution

Lake optical properties are strongly influenced by DOC quality/quantity (Maizel *et al.* 2017;

Read & Rose 2013), and in particular, a linear relationship between the light attenuation

coefficient for PAR and lake DOC was previously revealed (Read & Rose 2013). Humic lakes

often have higher DOC levels than clearwater lakes, as they receive large amounts of

terrestrially-derived allochthonous DOC, most of which are in the forms of humic and fulvic

acids. These carbon molecules are light-absorbing chromophoric dissolved organic matter

(CDOM), mainly absorbing light at wavelengths below 500 nm, including UV-A, UV-B, and the

blue light spectrum of photosynthetically active radiation (400-700nm) (Markager & Vincent

2000). In particular, the high concentrations of gilvin and tripton (mostly dissolved and

particulate humic substances, respectively) in peat lakes were shown to rapidly absorb blue and

green light (Stomp *et al.* 2007). The light spectral distribution can influence phototroph

distribution as revealed previously, as increasing humic substance concentration in freshwater

increased phototrophs absorbing red light and decreased those absorbing the central part of the

PAR spectrum (500–600 nm) (Vila *et al.* 1998). Therefore, as expected, we found that the

inferred lower availability of green light strongly limited opsin gene occurrence in humic lakes

since the encoded rhodopsins in our analysis were predicted to absorb green light.

Light availability also changes with water depth and such changes differ between clearwater and

humic lakes. Opsin gene abundance similarly varied with depth. Depth-associated decreases in

opsin gene abundance were relatively moderate and gradual in clearwater lakes. An extreme

example is from Lake Michigan, one of the Laurentian Great Lakes with low DOC (~2 mg/L),

where opsin genes remained abundant throughout the water column to 110 m deep. In Lake

Tanganyika, the opsin gene abundance estimate was around 30% even at 1100 m. These findings

are similar to marine studies, where sunlight penetrates deep into the open ocean, and opsin

genes and rhodopsin pigments were abundant in the photic zone (the upper 200 m) (Gomez-

Consarnau *et al.* 2019; Olson *et al.* 2018) and opsin genes were still present and expressed at

deeper depths (down to 1000 meters, albeit at low abundances) (Olson *et al.* 2018). Therefore, in

very large and clear lakes like Michigan and Tanganyika, rhodopsin-based phototrophy may

contribute significantly to the total phototrophy and energy budgets within the lake, similar to its

counterpart in the open ocean. By contrast, opsin gene abundance quickly declined to very low

levels within the first few meters in humic lakes. Therefore, humic lakes not only have lower

opsin abundance in the surface layer, but also have a much shallower opsin-containing zone,

both contributing to a lower significance of rhodopsin-enabled phototrophy in such environments.

The consistency between light climate and opsin gene profiles with depth suggests the

importance of lake optical properties in opsin gene distribution. At the same time, the inverse

correlation between opsin gene abundance and DOC levels for samples collected near the surface

suggests a strong influence of DOC on opsin gene occurrence. However, as DOC, especially

CDOM, and optical properties are generally correlated, it is difficult to tease apart their

individual influence on opsin gene distribution from our data. It is plausible that CDOM

indirectly influences opsin gene distribution by impacting light availability. In addition, we also

postulate a direct influence of DOC on opsin gene distribution, as low DOC may select for

opsin-containing bacteria as discussed later.

We should note that DOC not only influences lake optical properties but also influences lake

water pH and dissolved oxygen (DO). Humic lakes often have slightly acidic pH (~5-6, **Table**

S1), due to the high levels of humic and fulvic acids. In addition, DO usually decreases sharply

along the depth due to lower oxygenic photosynthesis and higher biological oxygen demand as

compared to clearwater lakes, leading to a shallow oxygenated zone. Therefore, compared

among lakes, acidic pH seems to be associated with low opsin abundance, and compared within

a lake, low DO is associated with low opsin abundance. Given that rhodopsin is a light-sensitive

protein, the pH and DO influence on opsin genes is likely less important than light availability,

although we cannot exclude their potential impacts based on the available data.

Ecological roles and opsin gene selection in freshwater

Light availability may be one of the main reasons for the wide distribution of opsin genes in

clearwater lakes. At the same time, we speculate that competitive advantages for surviving

provided by rhodopsins may favor their selection in clearwater lakes. Nearly all bacterial opsin

genes that we analyzed encode H^+-pumps, generating proton motive force (PMF), which

contributes to bacterial energy metabolism and renders physiological advantages, especially

when bacteria are under stress or carbon-limited conditions as discussed below.

As many vitamins are vulnerable to UV light, vitamins in surface water are subjected to

photodegradation in the photic zone. This may impose pressure on microbes to quickly take up

vitamins as soon as they are available, particularly in clearwater lakes. Proteorhodopsin gene

expression was linked to the light-enhanced growth of two marine vitamin-B_1 auxotrophic

Flavobacteria strains by providing PMF to power the Ton-B dependent acquisition of vitamin-B_1

(Gomez-Consarnau *et al.* 2016). Therefore, bacteria with H^+-pumping rhodopsins may compete

better for vitamin uptake compared to bacteria without this additional PMF-generating revenue,

and are selected for in clearwater lakes where bacteria are subjected to more intense UV

radiation than in brown-water humic bogs.

Rhodopsin-containing bacteria are photoheterotrophs relying on organic carbon for growth.

Timely carbon substrate scavenging is a key for heterotrophs to succeed under substrate-limited

conditions. Due to the additional PMF/ATP generated from light energy, bacteria with H^+-

pumping rhodopsins may be able to enhance the active uptake of organic substrates. More

importantly, the energy captured from the sunlight may reduce their reliance on respiration. For

example, when organic carbon became limiting, the respiration rate of '*Ca.* Pelagibacter ubique'

was lower under light as compared to in the dark while maintaining comparable cell densities

(Steindler *et al.* 2011). Therefore, possessing rhodopsins may be more advantageous in carbon-

limited than in carbon-rich environments. For example, some rhodopsin-possessing bacteria

exhibited light-promoted growth on low organic carbon media or light-enhanced survival under

carbon starvation (Gómez-Consarnau *et al.* 2010; Gómez-Consarnau *et al.* 2007; Johnson *et al.*

2010; Steindler *et al.* 2011; Yoshizawa *et al.* 2014). As lakes with high DOC, particularly humic

bogs, are not carbon-limited, whereas many clearwater lakes are, rhodopsin-containing bacteria

may be strongly selected for in clearwater lakes. This may partly and directly contribute to the

inverse correlation between opsin gene abundance and lake DOC levels, in addition to the

indirect impact of DOC through its influence on the availability of rhodopsin-absorbing spectra

and UV light.

Rhodopsin-based phototrophy in carbon and energy fluxes in freshwaters

We propose that rhodopsin-based phototrophy influences the carbon and energy fluxes in freshwaters, and the significance of this influence differs between clearwater and humic lakes, as illustrated in **Figure 8**. In clearwater lakes, a large fraction of heterotrophs are actually photoheterotrophs using rhodopsins to harvest sunlight while respiring DOC. As DOC may be limited by photoautotrophic primary producers or present in low levels in clearwater lakes, rhodopsin-phototrophy may be a significant pathway for harvesting energy in such lakes. By contrast, in humic lakes, a large amount of light energy is absorbed by CDOM and particulate humic substances, and such photochemical reactions generate labile DOC, which can be used for heterotrophic respiration (Cory & Kling 2018). As such, although the photodegradation process leaves a small amount of light for rhodopsin-phototrophy, photodegradation provides bacteria with a steady labile organic substrate supply for energy generation. Therefore, compared to clearwater lakes, the contribution of heterotrophy is larger, whereas the contribution of light harvesting by rhodopsins is smaller in the total energy budget in humic lakes. This further influences carbon flux, as the greater reliance on organic carbon based respiration increases the net emission of CO_2 from humic lakes. It was shown that CO_2 emission and net heterotrophy were associated with allochthonous DOC concentrations, and were higher in humic than in clearwater lakes (Jonsson *et al.* 2003). From our study, it is possible that rhodopsin-based phototrophy may be a previously overlooked factor contributing to such a difference in CO_2 emission from these two lake types.

SUMMARY

In this study, by analyzing opsin genes recovered in metagenomes collected from lakes with contrasting optical properties and DOC quantity/quality, we revealed differences in opsin diversity between two major freshwater types and recovered novel diversity within freshwater opsin genes. Many of these opsin genes were expressed, and some of them exhibited diel cyclic rhythms that differ among taxa. Further, we found opsin gene abundance is associated with lake DOC and depth in the water column, suggesting their ecological distribution can be influenced by organic matter and lake optical properties. In particular, opsin genes are widely present and highly expressed in clearwater lakes, and may be strongly selected for due to the higher light availability, lower DOC, and thus greater competitive advantages provided by rhodopsins in such environments. Therefore, we assert that rhodopsin-based phototrophy is a significant light harvesting mechanism, substantially influencing energy and carbon budgets in clearwater lakes.

ACKNOWLEDGEMENTS

We thank the North Temperate Lakes Microbial Observatory 2007-2012 field crews, UW-Trout Lake Station, the UW Center for Limnology, and the Global Lakes Ecological Observatory Network for field and logistical support. We acknowledge efforts by many McMahon Lab undergrads and technicians related to sample collection and DNA extraction. We thank the individual program directors and leadership at the National Science Foundation for their commitment to the continued support of long-term ecological research. The project was supported by funding from the United States National Science Foundation Microbial Observatories program (MCB-0702395), the Long-Term Ecological Research program (NTL-LTER DEB-1440297) and an INSPIRE award (DEB-1344254). This material is also based upon

work supported by the National Institute of Food and Agriculture, U.S. Department of

Agriculture (Hatch Project 1002996). The work conducted by the U.S. Department of Energy

Joint Genome Institute, a DOE Office of Science User Facility, is supported by the Office of

Science of the U.S. Department of Energy under Contract No. DE-AC02-05CH11231.

REFERENCES

Atamna-Ismaeel N, Sabehi G, Sharon I, *et al.* (2008) Widespread distribution of
proteorhodopsins in freshwater and brackish ecosystems. *ISME J* **2**, 656-662.
Balashov SP, Imasheva ES, Boichenko VA, *et al.* (2005) Xanthorhodopsin: A Proton Pump
with a Light-Harvesting Carotenoid Antenna. *Science* **309**, 2061-2064.
Balashov SP, Petrovskaya LE, Imasheva ES, *et al.* (2013) Breaking the carboxyl rule: lysine
96 facilitates reprotonation of the Schiff base in the photocycle of a retinal protein
from Exiguobacterium sibiricum. *J Biol Chem* **288**, 21254-21265.
Béjà O, Aravind L, Koonin EV, *et al.* (2000) Bacterial Rhodopsin: Evidence for a New Type of
Phototrophy in the Sea. *Science* **289**, 1902-1906.
Beja O, Lanyi JK (2014) Nature's toolkit for microbial rhodopsin ion pumps. *Proc Natl Acad*
Sci U S A **111**, 6538-6539.
Bendall ML, Stevens SLR, Chan L-K, *et al.* (2016) Genome-wide selective sweeps and gene-
specific sweeps in natural bacterial populations. *ISME J*.
Bratanov D, Kovalev K, Machtens JP, *et al.* (2019) Unique structure and function of viral
rhodopsins. *Nature Communications* **10**.
Brindefalk B, Ekman M, Ininbergs K, *et al.* (2016) Distribution and expression of microbial
rhodopsins in the Baltic Sea and adjacent waters. *Environ Microbiol* **18**, 4442-4455.
Buck M, Garcia SL, Fernandez L, *et al.* (2021) Comprehensive dataset of shotgun
metagenomes from oxygen stratified freshwater lakes and ponds. *Scientific Data* **8**.
Castresana J (2000) Selection of conserved blocks from multiple alignments for their use in
phylogenetic analysis. *Mol Biol Evol* **17**.
Chazan A, Rozenberg A, Mannen K, *et al.* (2022) Diverse heliorhodopsins detected via
functional metagenomics in freshwater Actinobacteria, Chloroflexi and Archaea.
Environ Microbiol **24**, 110-121.
Chuon K, Kim SY, Meas S, *et al.* (2021) Assembly of Natively Synthesized Dual
Chromophores Into Functional Actinorhodopsin. *Front Microbiol* **12**:652328.
Cory RM, Kling GW (2018) Interactions between sunlight and microorganisms influence
dissolved organic matter degradation along the aquatic continuum. *Limnology and*
Oceanography Letters **3**, 102-116.
Dam P, Rodriguez-R LM, Luo CW, *et al.* (2020) Model-based Comparisons of the Abundance
Dynamics of Bacterial Communities in Two Lakes. *Scientific Reports* **10**.

Darling AE, Jospin G, Lowe E, *et al.* (2014) PhyloSift: phylogenetic analysis of genomes and
metagenomes. *PeerJ* **2**, e243.
Denef VJ, Mueller RS, Chiang E, Liebig JR, Vanderploeg HA (2016) Chloroflexi CL500-11
Populations That Predominate Deep-Lake Hypolimnion Bacterioplankton Rely on
Nitrogen-Rich Dissolved Organic Matter Metabolism and C(1) Compound Oxidation.
Applied and Environmental Microbiology **82**, 1423-1432.
Dwulit-Smith JR, Hamilton JJ, Stevenson DM, *et al.* (2018) acI Actinobacteria Assemble a
Functional Actinorhodopsin with Natively Synthesized Retinal. *Applied and
Environmental Microbiology* **84**.
Edgar RC (2004) MUSCLE: a multiple sequence alignment method with reduced time and
space complexity. *BMC Bioinformatics* **5**, 113-113.
Edgar RC (2010) Search and clustering orders of magnitude faster than BLAST.
Bioinformatics **26**, 2460-2461.
Finkel OM, Beja O, Belkin S (2013) Global abundance of microbial rhodopsins. *ISME J* **7**,
448-451.
Flores-Uribe J, Hevroni G, Ghai R, *et al.* (2019) Heliorhodopsins are absent in diderm
(Gram-negative) bacteria: Some thoughts and possible implications for activity.
Environ Microbiol Rep **11**, 419-424.
Frigaard N-U, Martinez A, Mincer TJ, DeLong EF (2006) Proteorhodopsin lateral gene
transfer between marine planktonic Bacteria and Archaea. *Nature* **439**, 847-850.
Galachyants AD, Krasnopeev AY, Podlesnaya GV, *et al.* (2021) Diversity of Aerobic
Anoxygenic Phototrophs and Rhodopsin-Containing Bacteria in the Surface
Microlayer, Water Column and Epilithic Biofilms of Lake Baikal. *Microorganisms* **9**.
Garcia SL, McMahon KD, Grossart H-P, Warnecke F (2014) Successful enrichment of the
ubiquitous freshwater acI Actinobacteria. *Environ Microbiol Rep* **6**, 21-27.
Garcia SL, Stevens SLR, Crary B, *et al.* (2017) Contrasting patterns of genome-level diversity
across distinct co-occurring bacterial populations. *The ISME Journal*.
Ghai R, Mizuno CM, Picazo A, Camacho A, Rodriguez-Valera F (2014) Key roles for
freshwater Actinobacteria revealed by deep metagenomic sequencing. *Molecular
Ecology* **23**, 6073-6090.
Gifford SM, Sharma S, Rinta-Kanto JM, Moran MA (2011) Quantitative analysis of a deeply
sequenced marine microbial metatranscriptome. *Isme Journal* **5**, 461-472.
Gómez-Consarnau L, Akram N, Lindell K, *et al.* (2010) Proteorhodopsin Phototrophy
Promotes Survival of Marine Bacteria during Starvation. *PLoS Biology* **8**, e1000358.
Gómez-Consarnau L, Gonzalez JM, Coll-Llado M, *et al.* (2007) Light stimulates growth of
proteorhodopsin-containing marine Flavobacteria. *Nature* **445**, 210-213.
Gomez-Consarnau L, Gonzalez JM, Riedel T, *et al.* (2016) Proteorhodopsin light-enhanced
growth linked to vitamin-B1 acquisition in marine Flavobacteria. *ISME J* **10**, 1102-
1112.
Gomez-Consarnau L, Raven JA, Levine NM, *et al.* (2019) Microbial rhodopsins are major
contributors to the solar energy captured in the sea. *Science Advances* **5**.
Guindon S, Dufayard JF, Lefort V, *et al.* (2010) New algorithms and methods to estimate
maximum-likelihood phylogenies: assessing the performance of PhyML 3.0. *Syst Biol*
59, 307-321.

He S, Malfatti SA, McFarland JW, *et al.* (2015) Patterns in wetland microbial community
composition and functional gene repertoire associated with methane emissions.
mBio **6**, e00066-00015.
Imasheva ES, Balashov SP, Choi AR, Jung K-H, Lanyi JK (2009) Reconstitution of
Gloeobacter violaceus Rhodopsin with a Light-Harvesting Carotenoid Antenna.
Biochemistry **48**, 10948-10955.
Johnson ET, Baron DB, Naranjo B, *et al.* (2010) Enhancement of survival and electricity
production in an engineered bacterium by light-driven proton pumping. *Appl*
Environ Microbiol **76**, 4123-4129.
Jonsson A, Karlsson J, Jansson M (2003) Sources of Carbon Dioxide Supersaturation in
Clearwater and Humic Lakes in Northern Sweden. *Ecosystems* **6**, 224-235.
Keffer JL, Hahn MW, Maresca JA (2015a) Characterization of an Unconventional Rhodopsin
from the Freshwater Actinobacterium Rhodoluna lacicola. *J Bacteriol* **197**, 2704-
2712.
Keffer JL, Sabanayagam CR, Lee ME, *et al.* (2015b) Using Total Internal Reflection
Fluorescence Microscopy To Visualize Rhodopsin-Containing Cells. *Applied and*
Environmental Microbiology **81**, 3442-3450.
Kwon SK, Kim BK, Song JY, *et al.* (2013) Genomic makeup of the marine flavobacterium
Nonlabens (Donghaeana) dokdonensis and identification of a novel class of
rhodopsins. *Genome Biol Evol* **5**, 187-199.
Lami R, Cottrell MT, Campbell BJ, Kirchman DL (2009) Light-dependent growth and
proteorhodopsin expression by Flavobacteria and SAR11 in experiments with
Delaware coastal waters. *Environ Microbiol* **11**, 3201-3209.
Le SQ, Gascuel O (2008) An improved general amino acid replacement matrix. *Mol Biol Evol*
25, 1307-1320.
Linz AM, Aylward FO, Bertilsson S, McMahon KD (2020) Time-series metatranscriptomes
reveal conserved patterns between phototrophic and heterotrophic microbes in
diverse freshwater systems. *Limnology and Oceanography* **65**, S101-S112.
Linz AM, He SM, Stevens SLR, *et al.* (2018) Freshwater carbon and nutrient cycles revealed
through reconstructed population genomes. *Peerj* **6**.
Maizel AC, Li J, Remucal CK (2017) Relationships Between Dissolved Organic Matter
Composition and Photochemistry in Lakes of Diverse Trophic Status. *Environmental*
Science & Technology **51**, 9624-9632.
Man D, Wang W, Sabehi G, *et al.* (2003) Diversification and spectral tuning in marine
proteorhodopsins. *The EMBO Journal* **22**, 1725-1731.
Maresca JA, Miller KJ, Keffer JL, Sabanayagam CR, Campbell BJ (2018) Distribution and
Diversity of Rhodopsin-Producing Microbes in the Chesapeake Bay. *Applied and*
Environmental Microbiology **84**.
Markager S, Vincent WF (2000) Spectral light attenuation and the absorption of UV and
blue light in natural waters. *Limnology and Oceanography* **45**, 642-650.
Markowitz VM, Chen I-MA, Palaniappan K, *et al.* (2013) IMG 4 version of the integrated
microbial genomes comparative analysis system. *Nucleic Acids Research*.
Martinez-Garcia M, Swan BK, Poulton NJ, *et al.* (2012) High-throughput single-cell
sequencing identifies photoheterotrophs and chemoautotrophs in freshwater
bacterioplankton. *ISME J* **6**.

Matsen FA, Kodner RB, Armbrust EV (2010) pplacer: linear time maximum-likelihood and
Bayesian phylogenetic placement of sequences onto a fixed reference tree. *BMC*
Bioinformatics **11**, 538.
McCarren J, DeLong EF (2007) Proteorhodopsin photosystem gene clusters exhibit co-
evolutionary trends and shared ancestry among diverse marine microbial phyla.
Environ Microbiol **9**, 846-858.
Mizuno CM, Rodriguez-Valera F, Ghai R (2015) Genomes of Planktonic Acidimicrobiales:
Widening Horizons for Marine Actinobacteria by Metagenomics. *mBio* **6**.
Needham DM, Yoshizawa S, Hosaka T*, et al.* (2019) A distinct lineage of giant viruses brings
a rhodopsin photosystem to unicellular marine predators. *Proc Natl Acad Sci U S A*
116, 20574-20583.
Newton RJ, Kent AD, Triplett EW, McMahon KD (2006) Microbial community dynamics in a
humic lake: differential persistence of common freshwater phylotypes. *Environ*
Microbiol **8**, 956-970.
Newton RJ, McMahon KD (2011) Seasonal differences in bacterial community composition
following nutrient additions in a eutrophic lake. *Environ Microbiol* **13**, 887-899.
Oh S, Caro-Quintero A, Tsementzi D*, et al.* (2011) Metagenomic Insights into the Evolution,
Function, and Complexity of the Planktonic Microbial Community of Lake Lanier, a
Temperate Freshwater Ecosystem. *Applied and Environmental Microbiology* **77**,
6000-6011.
Olson DK, Yoshizawa S, Boeuf D, Iwasaki W, DeLong EF (2018) Proteorhodopsin variability
and distribution in the North Pacific Subtropical Gyre. *Isme Journal* **12**, 1047-1060.
Ottesen EA, Young CR, Gifford SM*, et al.* (2014) Multispecies diel transcriptional oscillations
in open ocean heterotrophic bacterial assemblages. *Science* **345**, 207-212.
Philosof A, Beja O (2013) Bacterial, archaeal and viral-like rhodopsins from the Red Sea.
Environ Microbiol Rep **5**, 475-482.
Pinhassi J, DeLong EF, Beja O, Gonzalez JM, Pedros-Alio C (2016) Marine Bacterial and
Archaeal Ion-Pumping Rhodopsins: Genetic Diversity, Physiology, and Ecology.
Microbiology and Molecular Biology Reviews **80**.
Podowski JC, Paver SF, Newton RJ, Coleman ML (2021) Genome streamlining,
proteorhodopsin, and organic nitrogen metabolism in freshwater nitrifiers. *bioRxiv*,
2021.2001.2019.427344.
Pushkarev A, Inoue K, Larom S*, et al.* (2018) A distinct abundant group of microbial
rhodopsins discovered using functional metagenomics. *Nature* **558**, 595-+.
Read JS, Rose KC (2013) Physical responses of small temperate lakes to variation in
dissolved organic carbon concentrations. *Limnology and Oceanography* **58**, 921-931.
Riedel T, Tomasch J, Buchholz I*, et al.* (2010) Constitutive Expression of the
Proteorhodopsin Gene by a Flavobacterium Strain Representative of the
Proteorhodopsin-Producing Microbial Community in the North Sea. *Applied and*
Environmental Microbiology **76**, 3187-3197.
Rozenberg A, Inoue K, Kandori H, Beja O (2021) Microbial Rhodopsins: The Last Two
Decades. In: *Annual Review of Microbiology, Vol 75, 2021* (ed. Gottesman S), pp. 427-
447.
Rusch DB, Halpern AL, Sutton G*, et al.* (2007) The Sorcerer II Global Ocean Sampling
expedition: northwest Atlantic through eastern tropical Pacific. *PLoS Biol* **5**.

Sharma AK, Sommerfeld K, Bullerjahn GS, *et al.* (2009) Actinorhodopsin genes discovered
in diverse freshwater habitats and among cultivated freshwater Actinobacteria.
ISME J **3**, 726-737.
Sharma AK, Zhaxybayeva O, Papke RT, Doolittle WF (2008) Actinorhodopsins:
proteorhodopsin-like gene sequences found predominantly in non-marine
environments. *Environ Microbiol* **10**, 1039-1056.
Shi YM, Tyson GW, Eppley JM, DeLong EF (2011) Integrated metatranscriptomic and
metagenomic analyses of stratified microbial assemblages in the open ocean. *Isme*
Journal **5**, 999-1013.
Sieradzki ET, Fuhrman JA, Rivero-Calle S, Gomez-Consamau L (2018) Proteorhodopsins
dominate the expression of phototrophic mechanisms in seasonal and dynamic
marine picoplankton communities. *Peerj* **6**.
Steindler L, Schwalbach MS, Smith DP, Chan F, Giovannoni SJ (2011) Energy starved
Candidatus Pelagibacter ubique substitutes light-mediated ATP production for
endogenous carbon respiration. *PLoS One* **6**, e19725.
Stomp M, Huisman J, Stal LJ, Matthijs HC (2007) Colorful niches of phototrophic
microorganisms shaped by vibrations of the water molecule. *ISME J* **1**, 271-282.
Thaben PF, Westermark PO (2014) Detecting Rhythms in Time Series with RAIN. *Journal of*
Biological Rhythms **29**, 391-400.
Tran P, Ramachandran A, Khawasik O, *et al.* (2018) Microbial life under ice: Metagenome
diversity and in situ activity of Verrucomicrobia in seasonally ice-covered Lakes.
Environ Microbiol **20**, 2568-2584.
Tran PQ, Bachand SC, McIntyre PB, *et al.* (2021) Depth-discrete metagenomics reveals the
roles of microbes in biogeochemical cycling in the tropical freshwater Lake
Tanganyika. *ISME J* **15**, 1971-1986.
Ugalde JA, Podell S, Narasingarao P, Allen EE (2011) Xenorhodopsins, an enigmatic new
class of microbial rhodopsins horizontally transferred between archaea and bacteria.
Biol Direct **6**, 52.
Vila X, Cristina XP, Abella CA, Hurley JP (1998) Effects of gilvin on the composition and
dynamics of metalimnetic communities of phototrophic bacteria in freshwater
North-American lakes. *J Appl Microbiol* **85 Suppl 1**, 138S-150S.
Vollmers J, Voget S, Dietrich S, *et al.* (2013) Poles apart: Arctic and Antarctic
Octadecabacter strains share high genome plasticity and a new type of
xanthorhodopsin. *PLoS One* **8**, e63422.
Wang Z, O'Shaughnessy TJ, Soto CM, *et al.* (2012) Function and Regulation of Vibrio
campbellii Proteorhodopsin: Acquired Phototrophy in a Classical
Organoheterotroph. *PLoS One* **7**.
Wurzbacher C, Salka I, Grossart HP (2012) Environmental actinorhodopsin expression
revealed by a new in situ filtration and fixation sampler. *Environ Microbiol Rep* **4**,
491-497.
Yoshizawa S, Kumagai Y, Kim H, *et al.* (2014) Functional characterization of flavobacteria
rhodopsins reveals a unique class of light-driven chloride pump in bacteria. *Proc*
Natl Acad Sci U S A **111**, 6732-6737.
Yutin N, Koonin EV (2012) Proteorhodopsin genes in giant viruses. *Biology Direct* **7**, 34-34.
Zabelskii D, Alekseev A, Kovalev K, *et al.* (2020) Viral rhodopsins 1 are an unique family of
light-gated cation channels. *Nature Communications* **11**.

Zehnpfennig JR, Hansel CM, Wankel SD, *et al.* (2022) Diel Patterns in Marine Microbial
Metatranscriptomes Reflect Differences in Community Metabolic Activity Over
Depth on the Continental Shelf of the North Atlantic. *Frontiers in Marine Science* **9**.

926
927

Data Accessibility and Benefit-Sharing

All metagenome data are publicly available at the DOE Joint Genome Institute's Integrated

Microbial Genome (IMG) database or European Nucleotide Archive, with their Accession No.

listed in Tables S2, S3, and S4. All metatranscriptome data are from the study by Linz *et al.*

(2020). Amino acid sequences of opsin genes used in this study are listed in Supplementary File

1.

Author Contributions

K.D.M and S.H. designed and initiated the study; A.M.L. generated the metatranscriptome and

part of the metagenome data; S.S. created the metagenome bins; P.T. generated part of the

metagenome data; F.M.-F., B.O.O., J.R.D.-S., K.T.F., K.D.M., S.H., A.M.L. and S.S.

participated in study discussion and metatranscriptome sampling; S.H. performed data analysis

and interpreted the results; S.H. and K.D.M. wrote the manuscript. All authors contributed to the

discussion and improvement of the manuscript.

Table 1 and Figure Legends

Table 1. MAGs containing opsin genes and the occurrence of key genes in their retinal
biosynthesis pathway

IMG genome ID	Genome name	Phylum	Class/Lineage	Phytoene synthesis	Phytoene desaturation	Lycopene cyclization	β-carotene cleavage
2582580528	ME1363	Actinobacteria	acI-A	+	+	+	-
2582580550	ME2297	Actinobacteria	acI-A	+	+	+	+
2582580564	ME3119	Actinobacteria	acI-A	+	+	+	+
2582580555	ME283	Actinobacteria	acI-B1	+	+	+	-
2582580575	ME4252	Actinobacteria	acI-B4	+	+	+	+
2582580630	TE3207	Actinobacteria	acI-B	+	+	+	+
2582580635	TE4208	Actinobacteria	acI-B	+	+	+	+
2582580657	TH2152	Actinobacteria	acI-B	+	+	+	+
2582580572	ME3864	Actinobacteria	acI-C	-	+	-	-
2582580553	ME2538	Actinobacteria	Luna1-A	-	+	-	-
2582580513	ME10152	Actinobacteria	Myco	+	+	+	-
2582580520	ME11576	Actinobacteria	acIV	-	-	+	-
2582580530	ME14260	Actinobacteria	acIV	-	-	-	-
2582580539	ME1719	Actinobacteria	Iluma-B1	-	-	-	-
2582580603	ME7484	Bacteroidetes	Cytophagia	+	+	+	+
2582580526	ME13165	Bacteroidetes	Flavobacteriia	+	+	+	+
2582580549	ME2278	Bacteroidetes	Flavobacteriia	+	+	-	+
2582580568	ME34551	Bacteroidetes	Flavobacteriia	+	+	+	+
2582580592	ME6180	Bacteroidetes	Flavobacteriia	+	+	+	+
2582580602	ME7257	Bacteroidetes	Flavobacteriia	+	+	+	-
2582580532	ME14726	Bacteroidetes	Sphigobacteriia	+	+	+	+
2582580533	ME15216	Bacteroidetes	Sphigobacteriia	+	+	+	+
2582580535	ME15824	Bacteroidetes	Sphigobacteriia	+	+	+	+
2582580541	ME17509	Bacteroidetes	Sphigobacteriia	+	+	+	+
2582580563	ME3085	Bacteroidetes	Sphigobacteriia	+	+	+	+
2582580571	ME3713	Bacteroidetes	Sphigobacteriia	+	+	+	+
2582580585	ME5493	Bacteroidetes	Sphigobacteriia	+	+	+	-
2582580587	ME5895	Bacteroidetes	Sphigobacteriia	+	+	+	+
2582580596	ME6777	Bacteroidetes	Sphigobacteriia	+	+	+	+
2582580601	ME7198	Bacteroidetes	Sphigobacteriia	+	+	+	+
2582580610	ME9147	Bacteroidetes	Sphigobacteriia	+	+	+	+
2582580614	ME9830	Bacteroidetes	Sphigobacteriia	+	+	+	+
2582580640	TE5144	Bacteroidetes	Sphigobacteriia	+	+	+	+
2582580669	TH3063	Bacteroidetes	Sphigobacteriia	-	+	+	+
2582580613	ME9495	Bacteroidetes	Unclassified	+	+	+	-
2582580574	ME422	Chloroflexi	Unclassified	+	+	-	-
2582580581	ME5338	Proteobacteria	Alphaproteobacteria	+	+	+	+
2582580650	TE9727	Proteobacteria	Alphaproteobacteria	-	-	-	-
2582580561	ME3080	Proteobacteria	Betaproteobacteria	+	+	+	+
2582580558	ME2907	Planctomycetes	Planctomycetia	+	+	+	+
2582580606	ME7803	Planctomycetes	Planctomycetia	+	+	-	+
2556921353[a]	MEE2912	Planctomycetes	Unclassified	+	+	+	+
2556921396[a]	MEE3553	Verrucomicrobia	Unclassified	+	+	-	+
2556921562[a]	MEE8025	Verrucomicrobia	Unclassified	+	+	-	+
2556921572[a]	MEE8561	Verrucomicrobia	Unclassified	+	+	-	-
2582580545	ME19941	Unclassified	Unclassified	+	+	+	+
2556921260[a]	MEE15704	Virus		-	-	-	-
2556921414[a]	MEE3901	Virus		-	-	-	-
2556921518[a]	MEE6525	Virus		-	-	-	-

947 [a]Although we mostly restricted this analysis to high quality MAGs, a few bins were also reported here, because
948 these taxa were not represented in the high quality MAGs.

Figure Legends

951
952
Figure 1. Phylogenetic tree of opsin protein sequences. This tree was viewed and annotated
using the Interactive Tree of Life tool (http://itol.embl.de). Opsin gene names are colored as
following: blue for ME, red for TE and TH metagenomes, and black for references from the
database. For sequences in MAGs/bins (highlighted with bold fonts), gene names start with
"ME", "MEE", "TE", and "TH" respectively. For the rest (un-binned) metagenome sequences,
gene names start with "ME-MG", "TE-MG", and "TH-MG", respectively. The inner circle color
strip indicates opsin gene types, and the outer circle color strip indicates taxon. Opsin types and
taxon present in our metagenomes were indicated with "*" in the legend. The outmost bar
indicates normalized coverage-weighted abundance of opsin genes in the metagenome (blue for
ME, and red for TE and TH). Some branches were labeled to highlight points discussed in the
manuscript. Rhodopsin types include ActR (actinorhodopsin), XR (xanthorhodopsin), acidiR
(acidirhodopsin), NaR (sodium-pumping rhodopsins), ClR (chloride-pumping rhodopsins), PR
(proteorhodopsin), HR (halorhodopsin), BR (bacteriorhodopsin), SR (sensory rhodopsin), XeR
(Xenorhodopsin), TM7 (novel rhodopsins from Candidatus Saccharibacteria phylum, previously
candidate division TM7), MycoR (novel rhodopsins from *Mycobacterium* spp. within the acTH2
lineage of Actinobacteria. See Supplementary Text for details about "mycoR"), and VR (viral
rhodopsin).

970
Figure 2. Subtree of opsin protein sequences belonging to freshwater Actinobacteria. Opsin
genes from our metagenomes were colored (blue for ME and red for TE and TH). Sequences in
MAGs/bins from our study were highlighted with bold fonts, with their taxonomic classification
indicated and their opsin sequence IMG gene IDs listed in the bracket. For the remaining
metagenome sequences (un-binned), gene names are their sequence Locus ID in IMG. Locus ID
starting with "B570J40625", "TBL_comb48_EPIDRAFT", and "TBL_comb47_HYPODRAFT"
are from ME, TE and TH metagenomes, respectively.

978
Figure 3. The significance of opsin genes in ME, TE and TH. (A) Percentage of opsin gene
containing MAGs among the total of 103, 36, and 63 bacterial MAGs recovered in ME, TE and
TH metagenomes respectively. (B) Average of opsin gene relative abundance in metagenomes
estimated by opsin gene coverage normalized by the coverage of single-copy genes. Error bars
indicate standard deviations derived from the 95, 29 and 29 individual metagenomes generated
from ME, TE and TH (listed in Table S3).

985
Figure 4. Opsin gene relative abundance in the surface layer of the 33 lakes versus the lake DOC
levels. The 33 lakes, DOC levels, Accession No. of metagenomes included in this analysis and
corresponding references for published studies were listed in Table S2. For lakes with multiple
metagenomes (derived from different dates, depths, or studies), opsin gene abundance was
estimated from individual metagenomes from the surface layer and averaged for that lake.
Among the 33 lakes, four small boreal lakes from Buck *et al.* (2021) only had total organic
carbon (TOC) measurements. As the difference between DOC and TOC was generally small for
lakes in that study, we use TOC in place of DOC for those four lakes on this plot.
994

995   **Figure 5.** Depth profiles of opsin gene relative abundance. Lakes on the top panels are
996   clearwater lakes, and on the bottom panels are humic lakes. For Lake Tanganyika, we only
997   plotted abundances from the top 250 meters. Opsin gene abundance for samples at the depths of
998   300 m and 1200 m was 0.28 and 0.30 respectively. Metagenomes included in this analysis and
999   corresponding reference papers were listed in Table S4.
1000
1001 **Figure 6.** Total opsin transcripts per liter of lake water averaged from all time points included in
1002 this analysis. The values from Lake Mendota and Sparkling Lake are about an order of
1003 magnitude higher than that in Trout Bog.
1004
Figure 7. Diel cyclic trends of opsin gene expression. Cyclic genes were determined based on
expression data collected from two consecutive days using the RAIN test with the *p*-value of <
0.05. For both lakes, the top panel shows all cyclic genes, and the bottom three panels show
cyclic genes in Actinobacteria, Bacteroidetes and Proteobacteria respectively. Read counts in
transcripts per liter were z-score transformed and are shown on y-axis. Genes were colored by
the peak hour of their expression as determined in the RAIN test, with warm and cold colors for
peaks at "day" and "night" (as determined by photosynthetically active radiation), respectively.
1012
1013 **Figure 8.** Difference on the significance of rhodopsin-based phototrophy in carbon metabolism
1014 and energy flux between clearwater lakes (such as Lake Mendota) and humic lakes (such as
1015 Trout Bog). Prevalent of reactions is qualitatively indicated by the thickness of lines.

Inner Circle: Opsin Type

- * ActR
- * Other XR
- * acidiR (Iluma-B)
- * acidiR (Iluma-A)
- NaR
- ClR
- * PR
- HR
- BR
- SR
- XeR
- * TM7
- * MycoR
- * VR
- * Unknown

Outer Circle: Taxon

- * Actinobacteria
- * Proteobacteria
- * Verrucomicrobia
- * Planctomycetes
- * Bacteroidetes
- Deinococcus
- Cyanobacteria
- * Chloroflexi
- Firmicutes
- Euryarchaeota
- * TM7
- * Viral group I
- * Viral group II
- * Other viral
- * Unclassified

acMyco
TM7

Virus

Firmicutes PR
Actinobacteria PR
Planctomycetes
(Verrucomicrobia)
PR with DTK motif

LD12

Taxon Type

ActR

Bacteroidetes

Planctomycetes
(Verrucomicrobia)

acidiR
(Iluma-B)

acidiR
(Iluma-A)

Lake Mendota

Clearwater lake with autochthonous DOC

Trout Bog

Humic lake with allochthonous DOC

hν ∿∿➤ Sunlight radiation

CDOM: chromophoric dissolved organic matter;   **LDOC**: Labile dissolved organic carbon

● Collection of photoautotrophic bacteria

● Collection of heterotrophic bacteria with a high propotion of cells containing proton-pumping rhodopsins

● Collection of heterotrophic bacteria with a low propotion of cells containing proton-pumping rhodopsins