Polymorphic short tandem repeats make widespread contributions to blood and serum
traits

Jonathan Margoliash[1], Shai Fuchs[2], Yang Li[1,3], Xuan Zhang[3], Arya Massarat[4], Alon Goren[3,*],

Melissa Gymrek[1,3,*]

[1]Department of Computer Science and Engineering, University of California San Diego, La Jolla,

CA

[2]Pediatric Endocrine and Diabetes Unit, Sheba Medical Center, Edmond and Lily Safra Children's

Hospital, Tel-Hashomer, Israel.

[3]Department of Medicine, University of California San Diego, La Jolla, CA

[4]Bioinformatics and Systems Biology Program, University of California San Diego, La Jolla, CA

*Correspondence should be addressed to agoren@ucsd.edu and mgymrek@ucsd.edu.

Abstract

Short tandem repeats (STRs), genomic regions each consisting of a sequence of 1-6 base pairs repeated in succession, represent one of the largest sources of human genetic variation. However, many STR effects are not captured well by standard genome-wide association studies (GWAS) or downstream analyses that are mostly based on single nucleotide polymorphisms (SNPs). To study the involvement of STRs in complex traits, we imputed genotypes for 445,720 autosomal STRs into genotype array data from 408,153 White British UK Biobank participants and tested for association with 44 blood and serum biomarker phenotypes. We used two fine-mapping methods, SuSiE and FINEMAP, to identify 119 high-confidence STR-trait associations across 93 unique STRs predicted as causal variants under all fine-mapping settings tested. Using these results, we estimate that STRs account for 5.2-7.6% of causal variants identifiable from GWAS signals for these traits. Our high confidence STR-trait associations implicate STRs in some of the strongest hits for multiple phenotypes, including a CTG repeat in *APOB* associated with circulating apolipoprotein B levels, a CGG repeat in the promoter of *CBL* associated with multiple platelet traits and a poly-A repeat in *TAOK1* associated with mean platelet volume. Replication analyses in additional population groups and orthogonal expression data further support the role of a subset of the candidate STRs we identify. Together, our study suggests that polymorphic tandem repeats make widespread contributions to complex traits, provides a set of stringently selected candidate causal STRs, and demonstrates the need to routinely consider a more complete view of human genetic variation in GWAS.

Introduction

Genome-wide association studies (GWAS) have become an indispensable tool for identifying which genes and non-coding regions in the genome influence complex human traits. While GWAS routinely identify tens to hundreds of genomic regions associated with individual traits, biological interpretation of GWAS results remains challenging[1]. A major limitation is that typical GWAS pipelines only consider a subset of common genetic variants. The majority of GWAS have been based on common single nucleotide polymorphisms (SNPs) and short insertions or deletions (indels) either genotyped using microarrays or imputed from population reference databases based on whole genome sequencing (WGS) data. However, detailed follow-up of individual GWAS signals has often revealed complex variants that were absent from the original analysis, such as repeats[2,3] or structural variants[4,5], to be the causal drivers of those signals. Indeed, a recent study showed that polymorphic protein-coding variable number tandem repeats (VNTRs) are likely causal drivers of some of the strongest GWAS signals identified to date for multiple traits[2].

Short tandem repeats (STRs) are a type of complex variant that consist of repeat units between 1-6bp duplicated many times in succession. Over one million STRs occur in the human genome[6], each spanning from tens to thousands of base pairs. STRs undergo frequent mutations resulting in gain or loss of repeat units[7], with per-locus mutation rates several orders of magnitude higher than average rates for SNPs[8] or indels[9]. Large repeat expansions at STRs are known to result in Mendelian diseases such as Huntington's disease, muscular dystrophies, hereditary ataxias and intellectual disorders[10,11]. Further, recent evidence suggests that more modest but highly prevalent variation at multi-allelic non-coding STRs can also be functionally relevant. We and others have found associations between STR length and both gene expression[3,12,13] and splicing[14,15]. The impact of non-coding STRs on gene expression is hypothesized to be mediated by a variety of mechanisms including modulating nucleosome positioning[16], altering methylation[12,17], affecting transcription factor recruitment[3] and impacting the formation of non-canonical DNA[18,19] and RNA[20,21] secondary structures. Together, these suggest that STRs potentially play an important role in shaping complex traits in humans.

Despite this potential, STRs are not well-captured by current GWAS. Because STRs are not directly genotyped by microarrays and are challenging to analyze from WGS, STRs have been largely excluded from widely used reference haplotype panels[22–24] and downstream GWAS analyses. While some STRs are in high linkage disequilibrium (LD) with nearby SNPs, many

highly multi-allelic STRs can only be imperfectly tagged by individual common SNPs, which are

typically bi-allelic. Thus effects driven by variation in repeat length have likely not been fully

captured, especially at those highly polymorphic STRs which cannot be well approximated by a

bi-allelic indel or well-tagged by bi-allelic SNPs.

Recent technological advances can now enable incorporation of STRs into GWAS. We and others

have created a variety of bioinformatic tools to genotype STRs directly from WGS by statistically

accounting for the noise inherent in STR sequencing[6,25–29]. We previously used these tools to

develop a reference haplotype panel consisting of both SNP and STR genotypes that allows for

imputation of STRs from genotype array data[30] in samples for which WGS is unavailable. In that

study we found that all but the most highly polymorphic STRs are amenable to imputation in

European cohorts, with an average per-locus imputation concordance of 97% with genotypes

obtained from WGS.

Here, we leverage our SNP-STR reference haplotype panel to impute genome-wide STRs into

SNP array data from 408,153 White British individuals obtained from the UK Biobank (UKB) for

which deep phenotype information is available[31]. Whereas a recent publication studied the effects

of protein-coding VNTRs (118 total VNTRs with repeat units of 7+ base pairs; total length of up

to several kilobases) on complex traits[2], our study focuses on a distinct set of repeats (namely

445,720 STRs with repeat units of 1bp to 6bp) which are mostly non-coding. We test for

association between imputed STR lengths and 19 blood cell count and 25 biomarker traits. These

traits provide multiple advantages: they are broadly and reliably measured, continuous, highly

polygenic and have variants with relatively large effect sizes, thus enabling well-powered

association testing.

We performed fine-mapping on these associations and estimate that STRs account for 5.2-7.6%

of signals identified by GWAS for the traits we studied. We observed that fine-mapping results in

some instances are substantially influenced by the choice of fine-mapper, and additionally are

moderately sensitive to data-processing choices and fine-mapper instabilities, and thus require

careful interpretation. After restricting to the signals which were consistently fine-mapped across

multiple fine-mappers and fine-mapping settings, we identified 93 unique STRs strongly predicted

to be causal for at least one trait. We highlight multiple STRs in this set which we predict contribute

to some of the strongest hits for many traits, including apolipoprotein B, platelet crit and mean

platelet volume. Overall, our study demonstrates the widespread role of polymorphic tandem

repeats and the need to consider a broad range of variant types in GWAS and downstream
analyses such as fine-mapping.

Results

Performing genome-wide STR association studies in 44 traits

We imputed genotypes for 445,720 autosomal STRs into phased genotype array data from
408,153 White British individuals from the UKB using Beagle[32] in combination with our published
SNP-STR reference haplotype panel[30] (**Methods**; **Fig. 1a**; **Supplementary Fig. 1**; **URLs**). We
find that this imputation yields broadly similar genotypes to WGS (see below). Compared to
common SNPs, which are typically bi-allelic, many of the imputed STRs are highly multi-allelic
(**Fig. 1b**). We tested STRs for association with 44 quantitative blood cell count and other
biomarker traits (**Supplementary Table 1**) for which phenotype information was available for
between 304,658-335,585 genetically unrelated subsets of individuals. To facilitate this and other
STR association studies, we developed associaTR (**URLs**), an open-source software package
for identifying associations between STR lengths (as measured by the number of repeat units)
and phenotypes.

For each STR-trait pair, we used associaTR to test for a linear association between STR dosage
(the sum of the imputed allele length dosages of both chromosomes) and the measured trait value
(**Fig. 1c-d**). For comparison, we used plink[33] to perform similar association tests using 70,698,786
SNP and short indel variants that were imputed into the same individuals. For all associations
(STR, SNP and indel), we included as covariates SNP-genotype principal components, sex, and
age (**Methods**). Additional covariates were included on a per-trait basis (**Supplementary Table
1**). We compared the output of our SNP analysis pipeline to previous results reported by Pan
UKBB[34] and found that our pipeline produced similar results, although had slightly weaker p-
values overall. (**Supplementary Fig. 2**).

We then compared signals identified by SNPs and indels to those identified by STRs. For each
trait we defined peaks as non-overlapping 250kb intervals centered on the lead genome-wide
significant variant (a SNP, indel or STR with p<5e-8) in that interval (**Methods**). We identified an
average of 389 peaks per trait, with blood cell count traits generally more polygenic than other
biomarkers (**Fig. 1e**). Of these peaks, 65.9% contained both a significant STR and a significant
SNP or indel, 32.5% contained only significant SNPs or indels, and 1.7% contained only
significant STRs. The majority of strong peaks (containing any variant with p<1e-100) were

identified by both STRs and SNPs and/or indels, in that they contain both an STR and a SNP or

indel with p<1e-80. No new strong peaks were identified only by STRs (**Fig. 1f**), which is

unsurprising given that SNP and indel genotypes were used to impute the STRs. Overall, p-values

of the lead SNP or indel and lead STR were similar for most peaks. Thus, we focused on fine-

mapping to determine which variants might be causally driving the identified signals.

Figure 1: Genome-wide association tests identify STRs, SNPs and indels associated with blood and biomarker
traits in the UKB. (a) Schematic overview of this study. STRs are imputed into phased hard-called variants obtained
from genotype arrays. GWAS is performed on SNPs and STRs in parallel. Regions with significant signals are identified
and then fine-mapped by two independent methods each under multiple scenarios, resulting in candidate causal STRs.
(b) Distribution of the number of common alleles at each imputed STR. Common alleles are defined as alleles
with estimated frequency ≥1% (**Methods**). For clarity we omitted from this figure the 237 imputed STRs with only a
single imputed allele with frequency ≥1%. **(c-d) Representative association results.** Manhattan plots are shown for
phenotypes **(c)** total bilirubin (an example moderately polygenic trait) and **(d)** platelet count (an example highly
polygenic trait). Large diamonds represent the lead variants (pruned to include at most one lead variant per 10Mb for
visualization). -log10 p-values are truncated at 100. Blue=SNPs and indels; orange=STRs. **(e) Summary of signals**
identified for each trait. Bars show the number of peaks per phenotype. Blue denotes peaks only containing genome-

wide significant SNPs and indels, purple denotes peaks containing both significant SNPs/indels and STRs. The number

of peaks only containing significant STRs is too small to be visible in this display. **(f) Comparison between lead SNP**

and indel and STR p-values at each peak. If there are no STRs in a peak, the y coordinate is set to zero (same for

SNPs and indels). p-values are capped at 1e-300, the maximum precision of our pipeline. The shade represents the

number of peaks falling at each position on the graph. The bottom left tile (which only contains peaks whose lead

SNP/indel and STR variants fall in the least significant bin) has been removed so as not to not skew the scale of the

color bar.

Fine-mapping suggests 5.2-7.6% of significant signals are driven by STRs

We applied statistical fine-mapping to identify candidate causal variants that may be driving the

GWAS signals detected above. We used two fine-mapping methods: SuSiE[35] and FINEMAP[36].

These methods differ in their modeling assumptions and thus provide partially orthogonal

predictions. For each trait we divided its genome-wide significant variants (SNPs, indels and

STRs) and nearby variants into non-overlapping regions of at least 250kb (**Methods**). This

resulted in 14,491 fine-mapping trait-regions (**Supplementary Table 2**), with some trait-regions

containing multiple nearby peaks. To compare outputs between fine-mappers in downstream

analyses, we defined the causal probability (CP) of each variant to be a number between 0 and

1 that indicates the variant's chance of causality. For FINEMAP we defined a variant's CP to be

the FINEMAP posterior inclusion probability (PIP) calculated for that variant. For SuSiE we

defined a variant's CP to be the maximal SuSiE alpha value for that variant across pure credible

sets in the region (**Supplementary Figs. 3-4**). We explain the rationale behind this choice in

Supplementary Note 1.

We used two approaches to study the contribution of STRs vs. SNPs and indels to fine-mapped

signals. First, we focused on the genome-wide significant variants (STR, SNP, or indel) with CP

≥ 0.8. (These accounted for a minority of the total 21,045 pure signals detected by SuSiE and the

total 33,756 signals detected by FINEMAP). SuSiE identified 4,494 such variants and FINEMAP

identified 5,170. Of these, 7.4% (range 1.3-13.0% across traits; SuSiE) and 7.6% (range 1.4-

14.0%; FINEMAP) are STRs. Among the subset of variants identified by both methods (3,961),

5.4% (range 1.0-11.1%) are STRs. Second, we considered the sum of CPs from all genome-wide

significant variants, thereby taking into account the many signals which were not resolved to a

single variant. STRs make up 5.2% (range 1.1-6.8%) of the total SuSiE CP sum and 7.4% (range

2.9-9.0%) of the total FINEMAP CP sum. A potential limitation of this second metric is that variants

with small CPs (CP ≤ 0.1) represent a large fraction (29.3% for SuSiE, 35.1% for FINEMAP) of

these totals (**Supplementary Fig. 5**). Additionally, our results below suggest that a sizable subset

of variant CPs are either discordant between fine-mappers or unstable, particularly for STRs

(**Supplementary Notes 2-3**), impacting the totals in both metrics. Nevertheless, the results above

suggest that between 5.2-7.6% of causal variants identifiable from GWAS can be attributed to an
STR, regardless of the fine-mapping method or metric used. This is in line with the total
percentage of non-major alleles per person and roughly half the total percentage of base pair
variation per person accounted for by STR lengths compared to other variant types in our study
(**Supplementary Table 3**). We report all 511 genome-wide significant STR associations across
409 distinct STRs with either FINEMAP or SuSiE CP ≥ 0.8 in **Supplementary Table 4**, and a
subset of those which pass more stringent thresholding in **Supplementary Table 5** (see below).

To evaluate the reliability of our approach for determining the relative contributions of STRs vs
SNPs and indels, we performed fine-mapping simulation analyses assuming a simple additive
model. We used two strategies for simulating phenotypes, in each case simulating only causal
SNPs and indels, and assessed to what extent STRs were incorrectly identified by SuSiE or
FINEMAP as contributing to the underlying signals. For the first strategy, we chose between one
and three causal SNPs or indels at random, weighting by minor allele frequencies, for a total of
1644 simulations. For the second, we chose causal variants to be those indicated by SuSiE as
being potentially causal for a representative trait (platelet count), in an attempt to more closely
capture properties of truly causal variants during simulation, for a total of 1374 simulations. These
simulations procedures and rationales are described in more detail in the **Methods**,
Supplementary Table 6 and **Supplementary Fig. 6**.

When simulating from one to three randomly chosen causal SNPs or indels, of genome-wide
significant variants that SuSiE or FINEMAP assigned CP ≥ 0.8, between 0% and 0.46% were
STRs. In contrast, using phenotypes simulated from the second strategy, between 1.4% and 3.2%
were STRs (**Supplementary Table 7**). Across all genome-wide significant variants, of the total
CP assigned by SuSiE or FINEMAP, 0.50% to 0.95% and 3.1% to 3.2% were assigned to STRs
in the first and second simulation strategies, respectively (**Supplementary Table 8**). These
numbers are uniformly lower than the 5.2-7.6% STR contributions identified above. This suggests
that if the 44 traits studied here have genetic architectures similar to the phenotypes we simulated,
the results above are unlikely to be fully explained by systematic bias of these fine-mappers in
favor of STRs. However, we expect there are complexities to the genetic architecture of blood
traits that are not captured by our simulations, and we cannot rule out the possibility that they
could result in such bias. These results also suggest that some of the fine-mapped STRs may be
false positives, and thus that our estimates of STR contributions could be somewhat inflated. On
the other hand, we also observed that a large fraction (66-81%) of simulated causal SNPs and
indels are not assigned CP ≥ 0.8 by fine-mapping, and we see a similar lack of sensitivity in limited

simulations performed with causal STRs (**Supplementary Table 7**). We expect this low sensitivity

is a greater source of uncertainty regarding the relative contribution of each variant type than the

false positive rates reported here.

We evaluated imputation quality at the 409 STRs in **Supplementary Table 4** by comparing

imputed genotypes to genotypes obtained from recently released WGS data for 200,025 UKB

individuals. At each locus, we computed the Pearson r^2 between length dosages from imputation

and summed lengths from WGS hard-calls, in addition to other per-locus metrics (**Supplementary**

Table 4). For 78.7% of these STRs, per-locus r^2 values are greater than 0.9 and for 92.7% they

are greater than 0.8. Other measurements of imputation concordance perform comparably well

(**Supplementary Fig. 7; Methods**). Overall, they suggest that fine-mapping results from imputed

data are unlikely to systematically differ from fine-mapping of hard-called genotypes for these loci.

Unless otherwise stated, all results reported below are based on imputed genotypes.

Identifying and characterizing confidently fine-mapped STRs

We performed additional analyses to identify a high confidence set of causal STR candidates.

First, we noticed that while SuSiE and FINEMAP tended to output similar results, they assigned

highly discordant CPs to a subset of variants (**Supplementary Note 2**; **Supplementary Figs. 8-**

10). Thus we conservatively restricted our follow-up analyses to the 167 candidate STR

associations with association p-values < 1e-10 and with CP ≥ 0.8 in both FINEMAP and SuSiE.

Second, to confirm that the default fine-mapping settings did not appreciably dictate our results,

we reran SuSiE and FINEMAP under a range of alternative settings (**Methods**), including using

best-guess STR genotypes instead of dosages and varying the prior distribution of effect sizes.

These additional fine-mapping runs tended to produce concordant results, but again for a subset

of STRs, produced highly inconsistent CPs (**Supplementary Figs. 11-14**), which we mostly

attribute to imputation uncertainty and FINEMAP instability (**Supplementary Note 3**). Thus, we

further restricted to the 118 (70.7%) of the 167 STR-trait associations which also maintained CP

≥ 0.8 across these additional fine-mapping runs. We refer to the STR-trait associations meeting

these criteria as confidently fine-mapped STR associations. Lastly, we added an association with

an STR in the *APOB* gene to this set as this association only failed to meet the above criteria

because the STR was simultaneously represented in both our STR reference panel and in the

SNP and indel set generated by the UKB team (**Supplementary Note 4**). In sum, we were left

with 119 confidently fine-mapped STR-trait associations corresponding to 93 distinct STRs, which

we display in **Fig. 2** and **Supplementary Table 5**.

We evaluated these fine-mapping results by measuring their replication rates in populations
besides White British individuals, with the expectation that causal associations will replicate at
higher frequencies in other populations than non-causal associations due to having common
biological functionality. The UKB includes self-identified groups of 8,043 Black, 7,952 South
Asian, 1,568 Chinese, 12,957 Irish and 16,051 Other White participants that passed quality
control, with about 40% of each population having WGS data (**Methods**). Using that WGS data
we validated the imputed genotypes of those populations for the STRs in **Supplementary Table**
4 and found that 78.2% and 93.9% of per-locus dosage r^2 values are greater than 0.8 and 0.6 in
the South Asian population, respectively, 45.5% and 84.8% in the Black population, and 64.3%
and 84.8% in the Chinese population (**Supplementary Fig. 7**). These metrics are weaker in the
non-White populations; this is expected given the largely European origin of our reference
haplotype panel. Nevertheless, our results suggest imputed genotypes are sufficiently accurate
across these groups for downstream analysis.

For each of these populations we performed follow-up STR association testing. Specifically, for
each trait, for each fine-mapping region for that trait identified among White British individuals, we
tested each STR in that region for association with that trait in each of the other populations
(**Supplementary Table 9**; data for individual loci in **Supplementary Tables 4** and **5**). As
expected, signals replicate at a higher rate in groups most closely related to our discovery cohort
(i.e. Irish and Other White). Encouragingly, fine-mapped associations replicate at higher rates
than non-fine-mapped associations in the Black, South Asian, and Chinese populations, even
after stratifying by the discovery p-value (**Fig. 3**, **Supplementary Fig. 15**). To quantitatively
measure this trend, for each population we fit a logistic regression model using whether signals
replicated in that population as the outcome, the fine-mapping status of those associations as the
independent variable, and their $-\log_{10}$(p-value) in the discovery cohort as a covariate
(**Supplementary Table 10**). This analysis further supports the conclusion that fine-mapped
associations replicate at higher rates. Additionally, the model consistently predicts that confidently
fine-mapped STR associations replicate at higher rates than STR associations fine-mapped by
either fine-mapper alone, although only a subset of those predictions reached nominal
significance, likely due to the small number of fine-mapped STR associations.

Next, we sought to characterize the set of confidently fine-mapped STRs. This set contains 62
poly-A repeats, 11 poly-AC, 5 poly-CCG, and 15 repeats with other units. Twelve of these STRs
overlap coding or untranslated regions (UTRs) (**Table 1**; **Supplementary Table 11;** the two
protein-coding repeats are further described in **Supplementary Note 4**; **Supplementary Fig. 16**).

Compared to all genome-wide significant STRs, confidently fine-mapped STRs were more likely

to be exonic trinucleotide STRs, in 5' UTR regions or in genes that are not protein-coding (two-

sided two-sample test of difference between proportions p=2e-26, 1e-3 and 2e-4 respectively).

No other annotation categories that we tested showed significant enrichment or depletion after

multiple hypothesis correction, likely due to low power arising from the small number of confidently

fine-mapped STRs (**Methods**; **Supplementary Fig. 17**). Lastly, we observed that 18 of these

confidently fine-mapped STRs are significant *cis* expression quantitative trait loci (QTLs) and 12

are significant *cis* DNA methylation QTLs in the Genotype-Tissue Expression (GTEx) dataset[37]

(**Fig. 2; Supplementary Tables 12-14**; **Supplementary Fig. 18**; **Methods**). We note that the

expression and methylation analyses were underpowered because of low sample sizes,

particularly for relevant tissue types (e.g. kidney, liver).

STR coordinate (hg19 chr:pos)	Reference allele	Repeat unit	Trait	Association p-value	Association Z-score	Gene (annotation)	Transcription direction
1:204527033	$(TAA)_9$	AAT	platelet crit	5.76e-17	-8.37	MDM4 (3' UTR)	+
2:21266752	$(CAG)_6(CGCAGGCAG)[CGC(CAG)_2]_2CGC$	CTG (poly-leucine)	Apolipo-protein B	1.37e-279	-35.76	APOB (Coding)	-
2:106510441	$(AC)_6GTG(CA)_{10}C(TA)_7T$	AC	mean platelet volume	6.93e-29	-11.15	NCK2 (3' UTR)	+
2:111878544	$(CGC)(CGCTGC)_2(CGC)_{13}C$	CCG	eosinophil count	4.96e-58	+16.06	BCL2L11 (5' UTR)	+
			eosinophil percent	5.88e-75	+18.32		
2:204311891	$T_4CT_4CT_3CT_{18}$	T	IGF-1	3.97e-11	-6.61	ABI2 (3' UTR*) ENST0000029 5851.10 (1)	+
6:90121977	$(TC)_7$	TC	cystatin C	1.24e-16	+8.28	RRAGD (3' UTR)	-
11:119077000	$(CGG)_{11}C$	CGG	mean sphered cell volume	6.88e-16	-8.07	CBL (5' UTR*) ENST0000063 4586.1 (5)	+
			platelet count	3.77e-83	+19.32		
			platelet crit	6.07e-103	+21.55		
15:40312923	T_{16}	T	red blood cell count	2.27e-20	+9.25	EIF2AK4 (Not protein coding*) ENST0000055 8743.1 (2)	+
16:67229794	$(CAG)_{13}(CAA)(CAG)(TAA)(CAG)_3$	AGC (poly-serine)	mean sphered cell volume	3.07e-23	+9.93	E2F4 (Coding)	+
			red blood cell count	1.08e-13	-7.43		
			mean corpuscular haemoglobin	2.83e-23	+9.94		
			mean corpuscular volume	9.27e-26	+10.49		
17:30469471	$(CCG)_{16}CC$	CCG	red blood cell distribution width	6.57e-13	+7.19	RHOT1 (5' UTR)	+
17:33871548	T_{17}	A	mean platelet volume	4.30e-62	-16.63	SLFN14 (3' UTR)	-
20:32971954	A_{20}	A	shbg	6.37e-15	-7.80	Y RNA ENST0000036 4628.1 (3)	-

Table 1: Confidently fine-mapped STRs are identified in coding regions and untranslated regions (UTRs). Imputed alternate alleles and rsIDs are provided in **Supplementary Table 11.** Repeat units here are calculated as described in the **Methods**, except that they are required to be on the strand in the direction of transcription of the overlapping gene. We denote with asterisks the three STRs in this list which only appear in non-canonical transcripts

294   for their genes from Ensembl release 106. Additionally, two STRs in this list only appear in transcripts or genes which
295   are not protein coding. For all those STRs, we provide Ensembl transcript numbers followed by parentheses containing
296   the Ensembl transcript support level, a number from 1 to 5, with 1 being the highest level of evidence and 5 being the
297   lowest. Additional analysis and discussion of the potential impact of the protein-coding repeats in *APOB* and *E2F4* are
298   given in **Supplementary Note 4**.

Figure 2: STRs are confidently fine-mapped to causally impact many traits. Only STRs with at least one confidently fine-mapped association are shown. Each triangle represents an STR-trait association with association p-value < 1e-10. Black=confidently fine-mapped, red-brown=CP ≥ 0.8 in either initial FINEMAP or SuSiE run, light-tan=all other associations with p-values < 1e-10. Triangle direction (up or down) indicates the sign of the association between STR length and the trait. Triangle size scales with association p-value. Similar traits are grouped on the x-axis by white and light-grey bands. STRs are grouped on the y-axis according to the traits they were confidently fine-mapped to.

STRs are labeled by the genes they reside in (protein coding genes preferred) or by chromosomal location and the
nearest gene for intergenic STRs. *CCDC26* and *TFDP2* each contain two confidently fine-mapped STRs and appear
twice. Light blue rows indicate (from left to right): whether each STR is associated with expression of a nearby gene in
the GTEx cohort (adjusted p<0.05; **Supplementary Table 12**), associated with the methylation of a nearby CpG island
(**Supplementary Table 14**), replicates with the same direction of effect in other populations (adjusted p<0.05;
Methods), repeat unit, and the number of common alleles for each STR (as defined in **Fig. 1**; see scale beneath).
Additionally, we mark the STRs in *TAOK1* and *RHOT1* as being expression QTLs even though they did not pass WGS
call rate filters in the GTEx dataset, as when we imputed the *TAOK1* STR into the GTEx dataset it was associated with
TAOK1 expression (**Methods**) and the *RHOT1* STR was associated with *RHOT1* expression in the Geuvadis dataset
(**Methods**). The data summarized here is available in **Supplementary Tables 4**, **5**, **12** and **14**.

Figure 3: Concordance of White British STR effect directions in Black, South Asian and Chinese populations.
The y-axis gives the fraction of STR associations measured in the discovery cohort that have the same direction of
effect when measured in the replication population regardless of p-value. Brackets beneath the x-axis denote the
binning of discovery -log$_{10}$(p-values). Brown=genome-wide significant associations (discovery p<5e-8),
orange=FINEMAP fine-mapped STR associations (discovery p<5e-8 and FINEMAP CP ≥ 0.8), teal=SuSiE fine-mapped
STR associations (discovery p<5e-8 and SuSiE CP ≥ 0.8) and purple=confidently fine-mapped STR associations.
Annotations above each bar indicate the number of STR-trait associations considered. We required confidently fine-
mapped STR associations to have p-value < 1e-10, thus they do not appear in the left-most bin. The trends in these
figures are somewhat sensitive to the choice of p-value bin boundaries so we additionally analyze this data using logistic
models (**Supplementary Table 10**).

Fine-mapped STRs capture known associations

We identified multiple fine-mapped STRs that were previously demonstrated to have functional
roles, providing supporting evidence for the validity of our pipeline. For instance, our confidently
fine-mapped set implicates a protein-coding CTG repeat (**Supplementary Table 11**) to be the
causal variant for one of the strongest signals for apolipoprotein B (p=1e-279; in one of the four
peaks for this trait with minimal p-value exceeding our numeric precision) which forms the
backbone of LDL cholesterol lipoproteins[38]. This locus is also one of the strongest signals for LDL
cholesterol levels (two-sided association t-test p-value = 6e-236; fifth most-significant genome-
wide peak) and the STR was marked as causal in 8 of 9 fine-mapping conditions. This repeat is
bi-allelic in the UKB cohort with an alternate allele corresponding to deletion of three residues
(Leu-Ala-Leu) in the signal peptide coded in the first exon of the apolipoprotein B (*APOB*) gene[39].
This deletion occurs in an imperfect region of the CTG repeat, with sequence CTGGCGCTG. In
agreement with a previous study[40], we found that the short allele is associated with higher levels
of both analytes. We discuss this locus further in **Supplementary Note 4**.

As another example, our initial fine-mapping implicates a multi-allelic AC repeat (**Supplementary**

Table 11) 6bp downstream of exon 4 of *SLC2A2* (also known as *GLUT2*, a gene that is most

highly expressed in liver) as causally impacting bilirubin levels (p=9e-18)*.* However, this repeat

was not included in the final confidently fine-mapped set due to its FINEMAP CP of 0.61 not

passing our 0.8 threshold, despite its SuSiE CP of 0.99. The potential link between GLUT2 and

bilirubin is described in **Supplementary Note 5.** Previous studies in HeLa and 293T cells showed

that inclusion of exon 4 of *SLC2A2* is repressed by the binding of mRNA processing factor hnRNP

L to this AC repeat[41,42], implicating this STR in *SLC2A2* splicing. Notably, these studies did not

investigate the impact of varying repeat copy number. We examined this STR in GTEx liver

samples and did not find a significant linear association between repeat count and the splicing of

exon 4, though we did find evidence for association with the splicing of exon 6 (**Supplementary**

Fig. 19).

A trinucleotide repeat in **CBL** *regulates platelet traits*

Most of the confidently fine-mapped STR associations identified by our pipeline have, to our

knowledge, not been previously reported. For example, this set includes positive associations

between the length of a highly polymorphic CGG repeat in the promoter of the gene *CBL* and both

platelet count (p=4e-83) and platelet crit (p=6e-103; 11[th] most-significant platelet crit peak;

Supplementary Table 11; **Fig. 4a-b**; **Supplementary Fig. 20**). This finding fits with observations

at other loci, as CG-rich repeats in promoter and 5' UTR regions have been strongly implicated in

transcriptomic regulation[13], often via epigenomic regulation[43,44]. This repeat in *CBL* is also

confidently fine-mapped to an association with mean sphered cell volume (p=7e-16;

Supplementary Fig. 21), but that is comparatively weaker and we do not discuss it here. For

both the platelet crit and platelet count phenotypes, SuSiE and FINEMAP identify two genome-

wide significant signals in this region, one of which they both localize to this STR. After

conditioning on a lead variant from the other signal (rs2155380), this STR becomes the lead

variant in the region by a wide margin (**Fig. 4c-d**). Conditioning on both rs2155380 and this STR

accounts for all the signal in the region (**Fig. 4e**). This supports the fine-mappers' prediction that

there are two signals in this region, one of which is driven by this STR.

This STR contains a common imperfection (rs7108857, which changes the second CGG copy to

TGG). That variant is in weak linkage disequilibrium (LD) with the length of the STR (r[2] in imputed

genotypes ranging between 0.023 (White British) and 0.175 (Chinese)) (**Fig. 4a**) and in strong LD

with the lead variant of the other signal in this region (rs2155380, White British r[2]=97.8%). While

rs7108857 is more strongly associated with the platelet traits than the STR's length (platelet count
p=9e-86, platelet crit p=4e-98) and is associated with CBL expression in the GTEx cohort
(minimum p=2.04e-18 in Esophagus Muscularis), given the fine-mappers' results that the STR
length association is an independent signal, it is unsurprising that the STR-length association
remains after stratifying on the presence of this imperfection (**Fig. 4f**). This suggests that
imperfections and repeat lengths are different characteristics of repeats that may have distinct
associations.

The imputation of this STR displays more modest levels of concordance with WGS data than less
polymorphic loci (r^2=0.582 between length dosages from imputation and summed lengths from
WGS; additional statistics reported in **Supplementary Table 5**). Yet reassuringly, hard-called
genotypes from WGS show similar trends with both platelet traits (**Fig. 2b**, **Supplementary Fig.**
20a,c-d). Further, population-specific allele length distributions at this STR observed in UKB are
highly concordant with those observed in the 1000 Genomes Project (**Fig. 4a**, **Supplementary**
Fig. 22).

CBL codes for a protein in the RING finger subfamily of E3 ubiquitin ligases – a class of proteins,
each with specific target molecule(s), that ubiquitinate their targets and thus prime them for
downstream degradation. CBL targets the thrombopoietin receptor MPL[45], thereby
downregulating thrombopoietin signaling[46]. As thrombopoietin is the primary positive regulator of
platelet production[47], this implicates *CBL* as a negative regulator of platelet production. As further
evidence of this link, controlled experiments in mice demonstrate that loss of *CBL* function in
megakaryocytes, the bone marrow platelet progenitor cells, results in increased platelet counts[48].
Further, we observed that increased CCG length was negatively associated with *CBL* expression
in three tissues in the GTEx cohort[37] (each with p-value < 0.05 after multiple hypothesis correction;
Supplementary Table 12; **Fig. 4g**). This association replicated (p=0.007) in European individuals
in the Geuvadis cohort[49] (**Fig. 4h**). This data fits with the STR association with platelet traits and
the connection between *CBL* and platelet production, leading to an overall hypothesis that longer
CCG repeat alleles contribute to increased platelet count by decreasing CBL expression (**Fig. 4i**).

Figure 4: A highly polymorphic CGG repeat in the promoter of *CBL* influences platelet traits. (a) Distribution of STR alleles across populations. The x-axis gives STR length (in number of full repeat unit lengths, using WGS data) and y-axis gives the population frequency. The hatched portion of each bar corresponds to the alleles of that length that include a "TGG" imperfection at the second repeat (rs7108857). The ultra-rare alleles with T imperfections at other locations we label as 'perfect' for the sake of these analyses. Colors denote different UKB populations. Extreme allele lengths 3-6, 21-33, 36 and 37 each have frequency less than 1% in all populations and have been omitted. **(b) STR length vs. platelet count.** STR length sums were calculated from WGS data on (potentially related) White British participants that passed quality control. Error bars correspond to 95% confidence intervals. Only allele length sums with a population frequency of 0.1% or greater are displayed. Rounded population-wide counts are displayed for each sum. **(c-e) Association of variants at the *CBL* locus with platelet crit.** Association plots in the White British population are shown before conditioning **(c)**, after conditioning on rs2155380 **(d)**, and after conditioning on both rs2155380 and STR length **(e)**. Light blue=SNPs and indels; orange=STRs. Red line=genome-wide significance threshold, black circles=the (CGG)$_n$ STR and rs2155380. **(f) STR length vs. platelet count conditioned on the TGG imperfection rs7108857.** STR length sums were calculated from WGS data on (potentially related) White British individuals that passed quality control. Blue=individuals homozygous for no imperfection (n=86,974); orange=individual homozygous for the imperfection (n=11,778). For each category, only length sums with a frequency of 0.2% or greater in that category are displayed. **(g-h) STR length vs. *CBL* expression.** Associations are shown for Cultured Fibroblasts from GTEx (n=393) **(g)** and LCLs from Geuavdis (n=447) **(h).** Yellow=African, blue=European. For **(g-h)** only allele length sums with at least 5 corresponding participants are displayed. **(i) Proposed pathway for effect of STR length on platelet traits.** The arrow denotes a positive association, the capped lines denote negative associations. Interactions are captioned by their information sources.

Additional examples of confidently fine-mapped STR-trait associations

We observe another 5' UTR CCG repeat in *BCL2L11* (also known as *BIM*) that is confidently fine-mapped to eosinophil percentage (p=6e-75) and eosinophil count (p=5e-58) (**Supplementary Table 11**). This repeat is the most strongly associated variant in the region for both traits, and conditioning on it accounts for the entire signal in this region (**Supplementary Fig. 23**). Proteins in the BCL-2 family are known to act as anti- or pro-apoptotic regulators. *BCL2L11* in particular is pro-apoptotic and is required in the tightly regulated lifespan of myeloid lineage cells[50], which include eosinophils. One mouse-model study showed that loss of repression of *BCL2L11* is associated with lowered eosinophil counts[51] and another showed that *BCL2L11* knockout increases counts of granulocytes, a class of cells including eosinophils[52]. This implicates *BCL2L11* in the regulation of eosinophil count, supporting the connection we observe between eosinophil count and the length of this STR.

While exonic repeats are potentially easier to interpret, a majority of our confidently fine-mapped STRs fall in intronic regions. We resolve one of the strongest signals for mean platelet volume (p<1e-300; one of twelve such peaks with p-values exceeding our numeric precision) to a multi-allelic poly-A STR in an intron of the gene *TAOK1* (**Supplementary Table 11**; **Supplementary Fig. 24a**). Conditioning on the length of this STR demonstrates that it explains the majority of the signal in this region (**Supplementary Fig. 24b**). The same STR also shows a strong association with platelet count, with p=2e-181 and a SuSiE CP of 1.

TAOK1 is a protein kinase that plays a role in regulating microtubule dynamics[53] and microtubule function is known to be critical to platelet generation[54]. The STR is in an intron of the canonical *TAOK1* transcript but lies immediately downstream of a non-protein coding transcript of *TAOK1* (ENST00000577583; which contains a retained intron) and is approximately 2.4kb upstream of a differentially spliced exon. The STR also bears the hallmarks of a regulatory element: it is located in a DNase hypersensitivity cluster and overlaps a transcription factor binding site for ESR1 (**Methods**). Although this STR was filtered from our initial GTEx callset due to low call rate (11%), we used our haplotype panel to impute it into available SNP data for that cohort (**Methods**). While we did not identify significant associations between repeat length and splicing of any nearby exons, STR lengths showed modest but significant negative correlation with *TAOK1* expression in 5 tissues (strongest p-value 8e-6 in thyroid; **Supplementary Fig. 24c-d**). The repeat also showed associations with the expressions of nearby genes *ANKRD13B* (negative association;

strongest p-value 5e-05 in heart) and *TP53I13* (positive association; strongest p-value 6e-5 in

tibial nerve), although their potential role in platelet regulation is less clear.

In another confidently fine-mapped example we identify a previously unreported association

between a GTTT repeat in an intron of estrogen receptor beta (*ESR2*) and haemoglobin

concentration (p=1e-24), red blood cell count (p=3e-24) and haematocrit (p=1e-26), where

additional repeat copies correspond to lower measurements of all three traits (**Supplementary**

Table 11; Supplementary Note 5; Supplementary Fig. 25). Despite the relatively weak

discovery p-value and differing allele distributions between the White British and Black

populations (**Supplementary Fig. 25c**), all three associations replicate in the Black population

with p-values < 0.05. Consistent with these associations, *ESR2* ligand 17β-estradiol has been

implicated in the regulation of red blood cell production[55,56]. We found a significant negative

association between STR length and *ESR2* expression in two tissues in GTEx (each with p-value

< 0.05 after multiple hypothesis correction; **Supplementary Table 12**). While evidence suggests

a link between *ESR2* and red blood cell production, the expected direction of effect is unclear

given the highly tissue-specific isoform usage and functions of this gene (**Supplementary Note**

5). Nevertheless, our results support a role of this STR in red blood cell production through

regulation of *ESR2*.

We observed many additional associations of interest amongst the confidently fine-mapped

STRs. For example, we find multiple AC repeats in our set that are significantly associated with

expression of nearby genes. This includes a polymorphic AC repeat located in the 3' UTR of

NCK2 which is associated with mean platelet volume (p=7e-29; **Supplementary Table 11;**

Supplementary Fig. 26). This repeat overlaps a binding site for the transcription factor PABPC1

and has a significant negative association with *NCK2* expression in multiple GTEx tissues

(strongest p=5e-7; **Supplementary Table 12**). We also find a highly polymorphic CCG repeat in

the 5' UTR of *RHOT1* that is associated with red blood cell distribution width (p=7e-13;

Supplementary Table 11). WGS data shows our imputation of this locus is poor, which is

unsurprising given the highly polymorphic nature of this STR. Nonetheless, the effect of this STR

is biologically plausible – this repeat overlaps a CTCF binding site, is located within a nucleosome

depleted region of a H3K27ac peak in LCLs, and shows a strong association with the expression

of *RHOT1* in LCLs from the Geuvadis dataset (p=2e-44 in Europeans, p=0.035 in Africans;

Supplementary Fig. 27). Finally, many STRs in our fine-mapped set consist of poly-A repeats.

While traditionally these have been particularly challenging to genotype[57], many such STRs,

including poly-A repeats in *MYO9B*, *DENND4A*, and *NRG4*, show strong statistical evidence of

causality and replicate across multiple population groups (**Fig. 2**). Taken together, these loci

exemplify the large number of confidently fine-mapped STRs our analysis provides for future

study.

Discussion

In this study, we imputed 445,720 STRs into the genomes of 408,153 participants in the UK

Biobank and associated their lengths with 44 blood cell and other biomarker traits. Using fine-

mapping, we estimate that STRs account for 5.2-7.6% of causal variants for these traits that can

be identified by GWAS. We stringently filtered the fine-mapping output to produce 119 confidently

fine-mapped STR-trait associations with strong evidence for causality across 93 distinct STRs

including some of the strongest signals for apolipoprotein B, platelet crit and mean platelet

volume. These confidently fine-mapped STRs replicated in the Black, South Asian and Chinese

UKB populations at higher rates than non-fine-mapped STRs (p<0.02 in each). A subset of these

STRs were associated with expression of nearby genes, providing evidence for their impact on

regulatory processes and explanations for their effects on the studied traits.

Broadly, our study highlights the importance of considering a more complete set of genetic

variants in complex trait analysis. It has been proposed that STRs may represent an important

source of the "missing heritability" associated with SNP-based GWAS[58,59]. Indeed, STRs, as well

as other complex variant types such as VNTRs[2], copy number variants[4], HLA types[60], and some

structural variants[61] are often highly multi-allelic and only imperfectly tagged by individual common

SNPs. This suggests that omitting these variants from analysis pipelines may overlook important

sources of causal variants and heritability. Further, we expect incorporation of this additional

source of causal variants, which we observe often exhibit population-specific allele distributions,

will improve downstream applications such as polygenic risk scores, particularly in constructing

scores that are more applicable across diverse populations.

While our results uncover many novel candidate causal STR variants, these findings are not

exhaustive. Our fine-mapping procedure was exceptionally conservative and excluded hundreds

of STR-trait associations strongly predicted to be causal in some but not all settings tested.

Further, whereas here we performed association tests with a fixed effects model, analysis with a

linear mixed model would increase power to detect additional associations. Additionally, compute

constraints limited our analysis to a small number of traits. In follow up studies we plan to extend

this analysis to a wide variety of medically-actionable traits.

Another limitation of our study is that it is based on imputed genotypes. Our SNP-STR reference

panel only included 27.5% of the 1.6 million STRs in the original HipSTR reference panel (**URLs**),

due to the exclusion of the many STRs with low heterozygosity, but also due to the exclusion of

non-autosomal STRs, most long repeats such as those implicated in pathogenic expansion

disorders, and many STR alleles that are common only in non-European populations[30]. Further,

imputed genotypes are inherently subject to some amount of noise, especially in non-European

populations. Despite these limitations, analysis of WGS data released for 200,025 UKB

participants[62] during the course of this study validated associations seen in imputed data.

Subsequently, popSTR calls at 2.5 million STRs were released for 150,000 of the participants

with WGS data[62]. Future studies which perform STR-based GWAS solely using WGS datasets

such as this should ameliorate the limitations mentioned above, enabling many future discoveries.

Methodological advances are also needed to support the study of STRs. Here we developed

associaTR, an open-source reproducible pipeline that enables future studies to conduct STR

length-based association tests. However, we envision that integrating support for STR length-

based tests and other complex variant associations into widely used GWAS toolkits would enable

more routine analysis of the full spectrum of human genetic variation. Improvements to our

association testing models are also likely to reveal new insights. In this study we only modeled

linear associations between STR lengths and trait values. Visualization of some of the

associations we identify suggests that these linear models only partially approximate those

signals. Many STR effects may be best described by non-linear models, such as quadratic or

sigmoid relationships between repeat copy numbers and traits. Yet fitting non-linear models would

require modeling the effects of the two alleles at each locus separately while simultaneously

controlling for the possibility of over-fitting, and is a topic of ongoing work. We also only tested

traits for association with the number of repeat units. However, inspection of individual loci reveals

that complex repeat structures are common (**Table 1**). Systematic evaluation of the potential for

epistasis between repeat imperfections and STR lengths, as well as between the lengths of

neighboring repeats, would potentially enable better understanding of the phenotypic impact of

STRs.

Importantly, our results highlight current challenges in performing statistical fine-mapping. We

found that fine-mapping results were in some cases highly sensitive to the choice of fine-mapping

tool, and to a lesser extent to data-processing choices and fine-mapper instabilities, where one

fine-mapping run would identify a variant as highly likely to be causal but a second would identify

it as having no causal impact. Further, our simulations suggest that fine-mappers have low

sensitivity rates even when sample sizes are large and all model assumptions are met. This
suggests statistical fine-mapping results should be interpreted with caution and evaluated for
sensitivity to model choices, and that further work is needed to make the process of fine-mapping
more robust.

Although fine-mapping inconsistencies were identified for SNPs and indels as well as STRs, they
were most prevalent for STRs. While this may in part be due to issues with imputing STR
genotypes, more research is needed to further evaluate the performance of current fine-mapping
tools on regions containing STRs. Additionally, there is a need for fine-mapping tools that can
model effects of multi-allelic variants. Existing fine-mapping frameworks in theory can accurately
model linear repeat-length associations, but we hypothesize that more detailed modeling of LD
between SNPs and individual STR alleles may enable more accurate model fitting procedures.
Similarly, during model fitting, existing tools often compare models which trade one causal variant
for another variant in close LD, but greater accuracy may be obtained by comparing models which
trade off a single, potentially causal, multi-allelic variant for multiple simultaneously-causal bi-
allelic variants.

Overall, our study provides a statistical framework for incorporating hundreds of thousands of
tandem repeat variants into GWAS studies, either via imputation, or using STR genotypes called
from WGS such as the newly released[62] UKB callset. Our study identifies dozens of novel
candidate variants for future mechanistic studies and demonstrates that STRs likely make a
widespread contribution to complex traits.

Methods

Selection of UK Biobank participants

We downloaded the fam file and sample file for version 2 of the phased SNP array data (referred
to in the UKB documentation as the 'haplotype' dataset) using the ukbgene utility (ver Jan 28
2019 14:09:15 - using Glibc2.28(stable)) described in UKB Data Showcase Resource ID 664
(**URLs**). The IDs from the sample file already excluded 968 individuals previously identified as
having excessive principal component-adjusted SNP array heterozygosity or excessive SNP
array missingness after call-level filtering[31] indicating potential DNA contamination. We further
removed withdrawn participants, indicated by non-positive IDs in the sample file as well as by IDs
in email communications from the UKB access management team. After the additional filtering,
data for 487,279 individuals remained.

We downloaded the sample quality control (QC) file (described in the sample QC section of UKB

Data Showcase Resource ID 531 (**URLs**)) from the European Genome-Phenome Archive

(accession EGAF00001844707) using pyEGA3[63]. We subsetted the non-withdrawn individuals

above to the 408,870 (83.91%) participants identified as White-British by column

`in.white.British.ancestry.subset` of the sample QC file. This field was computed by the

UKB team to only include individuals whose self-reported ethnic background was White British

and whose genetic principal components were not outliers compared to the other individuals in

that group[31]. In concordance with previous analyses of this cohort[31] we additionally removed data

for:

- 592 ● 2 individuals with an excessive number of inferred relatives, removed due to plausible
- 593 SNP array contamination (participants listed in sample QC file column
- 594 `excluded.from.kinship.inference` that had not already been removed by the UKB
- 595 team prior to phasing)

- 596 ● 308 individuals whose self-reported sex did not match the genetically inferred sex,
- 597 removed due to concern for sample mislabeling (participants where sample QC file
- 598 columns `Submitted.Gender` and `Inferred.Gender` did not match)

- 599 ● 407 additional individuals with putative sex chromosome aneuploidies removed as their
- 600 genetic signals might differ significantly from the rest of the population (listed in sample
- 601 QC file column `putative.sex.chromosome.aneuploidy`)

Following these additional filters the data for 408,153 individuals remained (99.82% of the White

British individuals considered above).

SNP and indel dataset preprocessing

We obtained both phased hard-called and imputed SNP and short indel genotypes made

available by the UKB. These variants were provided in reference genome hg19 coordinates, and

all analyses in this study, unless otherwise specified, were performed with hg19 coordinates.

Phased hard-called genotypes: We downloaded the bgen files containing the hard-called SNP

and indel haplotypes (release version 2) and the corresponding sample and fam files using the

ukbgene utility (UKB Data Showcase Resource 664 (**URLs**)). These variants had been genotyped

using microarrays and phased using SHAPEIT3[64] with the 1000 genomes phase 3 reference

panel[23]. Variants genotyped on the microarray were excluded from phasing and downstream
analyses if they failed QC on more than one microarray genotyping batch, had overall call-
missingness rate greater than 5% or had minor allele frequency less than 0.01%. Of the resulting
658,720 variants, 99.5% were single nucleotide variants, 0.2% were short indels (average length
1.9bp, maximal length 26bp), and 0.2% were short deletions (average length 1.9bp, maximal
length 29bp).

Imputed genotypes: We similarly downloaded imputed SNP data using the ukbgene utility
(release version 3). Variants had been imputed with IMPUTE4[31] using the Haplotype Reference
Consortium panel[22], with additional variants from the UK10K[24] and 1000 Genomes phase 3[23]
reference panels. The resulting imputed variants contain 93,095,623 variants, consisting of 96.0%
single nucleotide variants, 1.3% short insertions (average length 2.5bp, maximum length 661bp),
2.6% short deletions (average length 3.1bp, maximum length 129bp). This set does not include
the 11 classic human leukocyte antigen alleles imputed separately.

We used bgen-reader[65] 4.0.8 to access the downloaded bgen files in python. We used plink2[33]
v2.00a3LM (build AVX2 Intel 28 Oct 2020) to convert bgen files from both hard-called and imputed
SNPs to the plink2 format for downstream analyses. For hard-called genotypes, we used plink to
set the first allele to match the hg19 reference genome. Imputed genotypes already matched the
reference. Unless otherwise noted, our pipeline worked with imputed genotypes as non-reference
allele dosages, i.e. $\Pr(\text{heterozygous}) + 2 * \Pr(\text{homozygous alternate})$ for each individual.

STR imputation

We previously published a reference panel containing phased haplotypes of SNP variants
alongside 445,720 autosomal STR variants in 2,504 individuals from the 1000 Genomes
Project[23,30] (**URLs**). This panel focuses on STRs ascertained to be highly polymorphic and well-
imputed in European individuals. Notably, this excludes many STRs known to be implicated in
repeat expansion diseases, STRs that are primarily polymorphic only in non-European
populations, or STRs that are too mutable to be in strong linkage disequilibrium (LD) with nearby
SNPs.

The IDs listed in the 'str' column of Supplementary Table 2 at that URL describe which variants
in the reference panel are STRs and which are other types of variants. That produces a list of
445,715 unique variant IDs and 5 IDs which are each assigned to four separate variants in the
reference panel VCFs. For the IDs with multiple assignments, we selected the variant that

appeared first in the VCF and discarded the others, leaving 445,720 unique STR variants each

with unique IDs.

While our analyses with these STRs were performed using hg19 coordinates unless otherwise

stated, we also provide hg38 reference coordinates for these STRs in the supplementary tables.

We obtained those coordinates using liftover (**URLs**) which resulted in identical coordinates as in

HipSTR's[6] hg38 STR reference panel (**URLs**). All STRs successfully lifted over to hg38

coordinates.

To select shared variants for imputation, we note that 641,582 (97.4%) of SNP and indel variants

that were hard-called and phased in the UKB participants were present in our SNP-STR reference

panel. As a quality control step, we filtered variants that had highly discordant minor allele

frequencies between the 1000 Genomes European subpopulations (**URLs**) and White British

individuals from the UKB. We first took a maximal unrelated set of the White British individuals

(see **Phenotype Methods** below) and then visually inspected the alternate allele frequency of

the overlapping variants (**Supplementary Fig. 1**) and chose to remove the 110 variants with an

alternate allele frequency difference of more than 12%.

We used Beagle[32] v5.1 (build 25Nov19.28d) with the tool's provided human genetic maps (**URLs**)

and non-default flag `ap=true` to impute STRs into the remaining 641,472 SNPs and indels from

the SNP-STR panel into the hard-called SNP haplotypes. Though we performed the above

comparison between reference panel Europeans and UKB White British individuals, we

performed this STR imputation into all UKB participants using all the individuals in the reference

panel. We chose Beagle because it can handle multi-allelic loci. Due to computational constraints,

we ran Beagle per chromosome on batches of 1000 participants at a time with roughly 18GB of

memory. We merged the resulting VCFs across batches and extracted only the STR variants.

Lastly, we added back the `INFO` fields present in the SNP-STR reference panel that Beagle

removed during imputation.

Unless otherwise noted, our pipeline worked with these genotypes as length dosages for each

individual, defined as the sum of length of each of the two alleles, weighted by imputation

probability. Formally, $dosage = \sum_{a \in A} len(a) * [Pr(hap_1 == a) + Pr(hap_2 == a)]$, where A is the

set of all possible STR alleles at the locus, $len(a)$ is the length of allele a, and $Pr(hap_i == a)$ is

the probability that the allele on the ith haplotype is a, output by Beagle in the `AP1` and `AP2`

`FORMAT` fields of the VCF file.

Estimated allele frequencies (**Fig. 1b**) were computed as follows: for each allele length L for each

STR, we summed the imputed probability of the STR on that chromosome to have length L over

both chromosomes of all unrelated participants. That sum is divided by the total number of

chromosomes considered to obtain the estimated frequency of each allele.

Inferring repeat units

Each STR in the SNP-STR reference panel was previously annotated with a repeat period - the

length of its repeat unit - but not the repeat unit itself. We inferred the repeat unit of each STR in

the panel as follows: we considered the STR's reference allele and given period. We then took

each k-mer in the reference allele where k is the repeat period, standardized those k-mers, and

took their counts. We define the standardization of a k-mer to be the sequence produced by

looking at all cyclic rotations of that k-mer and choosing the first one lexicographically. For

example, the standardization of the k-mer CAG would be AGC. If the most common standardized

k-mer was less than twice as frequent as the second most common standardized k-mer, we did

not call a repeat unit for that STR (11,962 STRs; 2.68%). Otherwise, the most common

standardized k-mer was labeled as the forward-strand (based on the reference genome) repeat

unit for that STR. To infer the strand-independent repeat unit for the STR we looked at all rotations

of the forward-strand repeat unit in both the forward and reverse-complement directions and

chose whichever comes first lexicographically. For example the repeat unit for the STR

TGTGTGTG would be AC, while the forward-strand repeat unit would be GT. In the large majority

of cases the repeat unit identified by this approach is the unit which is duplicated or deleted in

alternate alleles, but this method of identifying repeat units does not consider alternate alleles and

so does not make that guarantee.

Phenotypes and covariates

IDs listed in this section refer to the UKB Data Showcase (**URLs**).

We analyzed a total of 44 blood traits measured in the UKB. 19 phenotypes were chosen from

Category Blood Count (Data Field ID 100081) and 25 from Category Blood Biochemistry (Data

Field ID 17518). We refer to them as blood cell count and biomarker phenotypes respectively.

The blood cell counts were measured in fresh whole blood while all the biomarkers were

measured in serum except for glycated haemoglobin which was measured in packed red blood

cells (details in Resource ID 5636). The phenotypes we analyzed are listed in **Supplementary**

Table 1, along with the categorical covariates specific to each phenotype that were included
during association testing.

We analyzed all the blood cell count phenotypes available except for the nucleated red blood cell,
basophil, monocyte, and reticulocyte phenotypes. Nucleated red blood cell percentage was
omitted from our study as any value between the bounds of 0% and 2% was recorded as exactly
either 0% or 2% making the data inappropriate for study as a continuous trait. Nucleated red blood
cell count was omitted similarly. Basophil and monocyte phenotypes were omitted as those cells
deteriorate significantly during the up-to-24-hours between blood draw and measurement. This
timing likely differed consistently for different clinics, and different clinics drew from distinct within-
White British ancestry groups, which could lead to confounding with true genetic effects. See
Resource ID 1453 for more information. Reticulocytes were excluded from our initial pipeline. This
left us with 19 blood cell count phenotypes. For each blood cell count phenotype we included the
machine ID (1 of 4 possible IDs) as a categorical covariate during the association tests to account
for batch effects.

Biomarker measurements were subject to censoring of values below and above the measuring
machine's reportable range (Resource IDs 1227, 2405). **Supplementary Table 1** includes the
range limits and the number of data points censored in each direction. Five biomarkers (direct
bilirubin, lipoprotein(a), oestradiol, rheumatoid factor, testosterone) were omitted from our study
for having >40,000 censored measurements across the population (approximately 10% of all
data), since those would require analysis with models that take censoring into account. The
remaining biomarkers had less than 2,000 censored measurements. We excluded censored
measurements for those biomarkers from downstream analyses as they consisted of a small
number of data points. For each serum biomarker we included aliquot number (0-3) as a
categorical covariate during association testing as an additional step to mediate the dilution issue
(described in Resource ID 5636). Glycated haemoglobin was not subject to the dilution issue,
being measured in packed red blood cells and not serum, so no aliquot covariate was published
in the UKB showcase or included in our analysis.

For each phenotype we took the subset of the 408,153 individuals above that had a measurement
for that phenotype during the initial assessment visit or the first repeat assessment visit,
preferentially choosing the measurement at the initial assessment when measurements were
taken at both visits. We include a binary categorical covariate in association testing to distinguish
between phenotypes measured at the initial assessment and those measured at the repeat

assessment. Each participant's age at their measurement's assessment was retrieved from Data
Field ID 21003.

The initial and repeat assessment visits were the only times the biomarkers were measured. The
blood cell count phenotypes were additionally measured for those participants who attended the
first imaging visit. We did not use those measurements and for each phenotype excluded the
<200 participants whose only measurement for that phenotype was taken during the first imaging
visit as we could not properly account for the batch effect of a group that small (**Supplementary**
Table 1).

No covariate values were missing. Before each association test we checked that each category
of each categorical covariate was obtained by at least 0.1% of the tested participants. We
excluded the participants with covariate values not matching this criterion, as those quantities
would be too small to properly account for batch effects. In practice, this meant that for each
biomarker phenotype we excluded the <100 participants that were measured using aliquot 4, and
that for 8 of the biomarker phenotypes we additionally excluded the ≤125 participants that were
measured using aliquot 3 (**Supplementary Table 1**).

For each phenotype we then selected a maximally-sized genetically unrelated subset of the
remaining individuals using PRIMUS[66] v1.9.0. When multiple such maximal subsets existed (for
instance, wherever a single individual needed to be chosen from a family of two), one subset was
chosen arbitrarily, thus introducing some lack of reproducibility. Precomputed measures of
genetic relatedness between participants (described in UKB paper supplement section 3.7.1[31])
were downloaded using ukbgene (Resource ID 664). We ran PRIMUS with non-default options –
-no_PR -t 0.04419417382 where the t cutoff is equal to 0.5^9, chosen so that two individuals
are considered to be related if they are relatives of third degree or closer. This left between
304,658 and 335,585 unrelated participants per phenotype (**Supplementary Table 1**).

Sex and ancestry principal components (PCs) were included as covariates for all phenotypes.
Participant sex was extracted from fam file available with hard called genotypes (see above). The
top 40 ancestry PCs were extracted from the corresponding columns of the sample QC file (see
the Participants **Methods** section above).

We then rank-inverse-normalized phenotype values for association testing. The remaining
unrelated individuals for each phenotype were ranked by phenotype value from least to greatest
(ties broken arbitrarily) and the phenotype value for association testing for each individual was

taken to be $normal\ quantile\left(\frac{sample\ rank\ +0.5}{n\ samples}\right)$. We use rank-inverse normalization as it is standard

practice, though it does not have a strong theoretical foundation[67] and only moderate empirical

support[68–71].

For each phenotype and its remaining unrelated individuals we standardized all covariates to

have mean zero and variance one for numeric stability.

Association testing

We performed STR and SNP association testing separately. We developed associaTR to

streamline performing association tests between STR length and quantitative traits. While our

approach relies on a standard linear model, linear mixed models based on STR length dosages

would likely result in increased power and will be considered in future studies. As our downstream

analyses required STR and SNP associations to be comparable, we also used a standard linear

model for SNP association testing.

For STR association testing, the imputed VCFs produced by Beagle were accessed in python

with cyvcf2[72] 0.30.14 and v4.2.1 of our TRTools library[73]. In line with plink's recommendation for

SNP GWAS (**URLs**), 6 loci with non-major allele dosage < 20 were filtered. For each STR, we fit

the linear model $\vec{y} = \vec{g} * \beta_g + C * \overrightarrow{\beta_C} + \vec{\epsilon}$ where $\vec{y}$ is the vector of rank-inverse-normalized

phenotype values per individual, $\vec{g}$ is the vector of STR length dosage genotypes per individual,

β_g is the effect size of this STR, C is the matrix of standardized covariates, $\overrightarrow{\beta_C}$ is the vector of

covariate effect sizes, and $\vec{\epsilon}$ is the vector of errors between the model predictions and the

outcomes. Models were fit using the `regression.linear_model.OLS` function of the Python

statsmodels library v0.13.2 (**URLs**). Per GWAS best-practices, we used imputation dosage

genotypes instead of best-guess genotypes[74].

We used plink2[33] v2.00a3LM (build AVX2 Intel 28 Oct 2020) for association testing of imputed

SNPs and indels. For each analysis, plink first converts the input datasets to its pgen file format.

To avoid performing this operation for every invocation of plink, we first used plink to convert the

SNP and indel bgen files to pgen files a single time. We invoked plink once per chromosome per

phenotype. We used the plink flag `--mac 20` to filter loci with minor allele dosage less than 20

(**URLs**). Plink calculates minor allele counts across all individuals before subsetting to individuals

with a supplied phenotype, so this uniformly filtered 22,396,837 (24.1%) of the input loci from

each phenotype's association test leaving 70,698,786 SNPs and indels. Plink fit the same linear

model described above in the STR associations, except that $\vec{g}$ is the vector of dosages of the

non-reference SNP or indel allele.

For conditional regressions, we fit the model $\vec{y} = \vec{g} * \beta_g + \vec{f} * \beta_f + C * \overrightarrow{\beta_C} + \vec{\epsilon}$ where all the terms

are as described above, except $\vec{f}$ is the vector of per-individual genotypes of the variant being

conditioned on, and β_f is its effect size.

Comparison with Pan-UKB pipeline

We compared the results of our pipeline to results available on the Pan UKBB[34] website (see

URLs) using bilirubin as an example trait. We matched variants between datasets on

chromosome, position, reference and alternate alleles, excluding variants not present in both

pipelines. We found our pipeline produced largely similar but somewhat less significant p-values

than those reported for European participants in Pan UKBB (**Supplementary Fig. 2**).

Defining significant peaks

Given a peak width w (bp), we selected variants to center peaks on in the following manner:

1. Order all variants (of all types) from most to least significant. For variants which exceed

our pipeline's precision (p<1e-300), order them by their chromosome and base pair from

first to last. (These variants will appear at the beginning of the list of all variants).

2. For each variant: If the variant has p-value ≥ 5e-8, break. If there is a variant in either

direction less than w bp away which has a lower p-value, continue. Otherwise, add this

variant to the list of peak centers.

We define peaks to be the w (base pair) width regions centered on each selected variant. The

statistics given in the **Results** are calculated using $w = 250kb$. The identification of peaks in **Fig.**

1c-d was made with $w = 20mb$ for visualization purposes. Note that peaks within w bp of the end

of a chromosome will necessarily be smaller than w bp in width.

Identifying indels which are STR alleles

Some STR variant alleles are represented both as alleles in our SNP-STR reference panel and

as indel variants in the UKB imputed variants panel. We excluded the indel representations of

those alleles from fine-mapping, as they represent identical variants and could confound the fine-

mapping process. For each STR we constructed the following interval:

825
$$\begin{cases} (start - 3, end + 3), & period = 1 \\ (start - 2 * period, end + 2 * period), & period > 1 \end{cases}$$

where $period$ is the length of the repeat unit. and $start$ and end give the coordinates of the STR

in base pairs. We call an indel an STR-indel if it only represents either a deletion of base pairs

from the reference or an insertion of base pairs into the reference (not both), overlaps only a

single STR based on the interval above, and represents an insertion or deletion of full copies of

that STR's repeat unit. We conservatively did not mark any STR-indels for STRs whose repeat

units were not called (see above) or for which the insertion or deletion was not a whole number

of copies of any rotation of the repeat unit.

Fine-mapping

For each phenotype, we selected contiguous regions to fine-map in the following manner:

1. Choose a variant (SNP or indel or STR) with p-value < 5e-8 not in the major
histocompatibility complex (MHC) region (chr6:25e6-33.5e6).

2. While there is a variant (SNP or indel or STR) with p-value < 5e-8 not in the MHC region
and within 250kb of a previously chosen variant, include that variant in the region and
repeat.

3. This fine-mapping region is (min variant bp – 125kb, max variant bp + 125kb).

4. Start again from step 1 to create another region, starting with any variant with p-value <
5e-8 not already in a fine-mapping region.

This is similar to the peak selection algorithm above but is designed to produce slightly wider

regions so that we could fine-map nearby peaks jointly. We excluded the MHC because it is known

to be difficult to effectively fine-map. Note that peaks within 125kb of the end of a chromosome

will necessarily be smaller than the minimum 125kb width in that direction.

This produced 14,494 trait-regions. Due to computational challenges during fine-mapping (see

below), we excluded three regions (urate 4:8165642-11717761, total bilirubin 12:19976272-

22524428 and alkaline phosphatase 1:19430673-24309348) from downstream analyses (see

below), leaving 14,491 trait-regions.

We used two fine-mapping methods to analyze each region:

SuSiE[35]: For each fine-mapping trait-region, for each STR and SNP and indel variant in that region

that was not filtered before association testing, was not an STR-indel variants (see above) and

had p-value ≤ 5e-4 (chosen to reduce computational burden), we loaded the dosages for that

variant from the set of participants used in association testing for that phenotype. For those

regions we also loaded the rank-inverse-normalized phenotype values and covariates

corresponding to that phenotype. We separately regressed the covariates out of the phenotype

values and out of each variant's dosages and streamed the residual values to HDF5 arrays using

h5py v3.6.0 (**URLs**). We used rhdf5 v2.38.0 (**URLs**) to load the h5 files into R. We used an R

script to run SuSiE v0.11.42 on that data with non-default values `min_abs_corr=0` and

`scaled_prior_variance=0.005`. `min_abs_corr=0` forced SuSiE to output all credible sets

it found so that we could determine the appropriate minimum absolute correlation filter threshold

in downstream analyses. We set `scaled_prior_variance` to 0.005 which we considered is a

more realistic guess of the per-variant percentage of signal explained than the default of 20%,

although we determined that this parameter had no effect on the results (**Supplementary Note**

3). The SuSiE results for some regions did not converge within the default number of iterations

(100) or produced the default maximum number of credible sets (10) and all those credible sets

seemed plausible (minimum pair-wise absolute correlation ≥ 0.2 or size ≤ 50). We reran those

regions with the additional parameters `L=30` (maximum number of credible sets) and

`max_iter=500`. No regions failed to converge in under 500 iterations. We re-analyzed several

loci that produced 30 plausible credible sets again with `L=50`. No regions produced 50 plausible

credible sets. SuSiE failed to finish for two regions (urate 4:8165642-11717761, total bilirubin

12:19976272-22524428) in under 48 hours; we excluded those regions from downstream

analyses. A prior version of our pipeline had applied a custom filter to some SuSiE fine-mapping

runs that caused SNPs with total minor allele dosage less than 20 across the entire population to

be excluded. For consistency, any regions run with that filter which produced STRs included in

our confidently fine-mapped set were rerun without that filter. Results from the rerun are reported

in **Supplementary Table 4**.

SuSiE calculates credible sets for independent signals and calculates an alpha value for each

variant for each signal – the probability that that variant is the causal variant in that signal. We

used each variant's highest alpha value from among credible sets with purity ≥ 0.8 as its casual

probability (CP) in our downstream analyses (or zero if it was in no such credible sets). See

Supplementary Note 1.

FINEMAP[36]: We selected the STR and SNP and indel variants in each fine-mapping region that

were not filtered before association testing and had p-value < 0.05 (chosen to reduce

computational burden). We excluded STR-indels (see above). We constructed a FINEMAP input

file for each region containing the effect size of each variant and the effect size's standard error.

All `MAF` values were set to `nan` and the `ref` and `alt` columns were set to `nan` for STRs as this

information is not required. We then took the unrelated participants for the phenotype, loaded

their dosage genotypes for those variants and saved them to an HDF5 array with h5py v3.6.0

(**URLs**). To construct the LD input file required by FINEMAP, we computed the Pearson

correlation between dosages of each pair of variants. We then ran FINEMAP v1.4 with non-default

options `--sss --n-causal-snps 20`. In regions which FINEMAP gave non-zero probability

to their being 20 causal variants, we reran FINEMAP with the option `-n-causal-snps 40` and

used the results from the rerun. FINEMAP did not suggest 40 causal variants in any region.

FINEMAP caused a core dump when running on the region alkaline phosphatase 1:19430673-

24309348 so we excluded that region from downstream analyses. (For convenience, for the

regions containing no STRs, we directly ran FINEMAP with `--n-causal-snps 40`, unless those

regions contained less than 40 variants in which case we ran FINEMAP with `--n-causal-snps`

`<#variants>`).

We used the FINEMAP's PIP output for each variant in each region as its CP in downstream

analyses.

Alternative fine-mapping conditions

We reran SuSiE and FINEMAP using alternative settings on trait-regions that contained one or

more STRs with p-value < 1e-10 and CP ≥ 0.8 in both the original SuSiE and FINEMAP runs.

Each new run differed from the original run in exactly one condition. We restricted our set of high-

confidence fine-mapped STRs (**Supplementary Table 5**) to those that had p-value < 1e-10 and

CP ≥ 0.8 in the original runs and maintained CP ≥ 0.8 in a selected set of those alternate

conditions.

For SuSiE, we evaluated using best-guess genotypes vs. genotype dosages as input. For

FINEMAP, we tested varying the p-value threshold, choice of non-major allele frequency

threshold, effect size prior, number of causal variants per region, and stopping threshold.
Additionally, we reran FINEMAP with no changed settings to examine potential FINEMAP
instability.

See **Supplementary Note 3** for a more detailed discussion of these various settings and their
impact on fine-mapping results.

Fine-mapping simulations

We simulated phenotypes under additive genetic models and fine-mapped those phenotypes
separately at individual regions. Our simulations used real genotypes from White British UKB
participants and focused on regions originally identified by our GWAS to maintain realistic LD
patterns observed at regions with true signals. We used regions associated with platelet count as
it was the phenotype with the maximal number of fine-mapping regions (n=548).

Strategies for choosing causal variants and effect sizes: We applied three different strategies for
choosing causal variants from these regions and choosing their effect sizes. For each strategy,
we simulated phenotypes from those variants, ran SuSiE and FINEMAP on all SNPs and STRs
in the region against the simulated phenotypes and determined whether the fine-mappers
correctly identified the variants simulated to be causal.

For the first strategy we chose causal SNPs and indels at random, weighting by minor allele
frequency (MAF). For this strategy, we did not simulate causal STRs. To begin, we took all
SNPs/indels in all platelet count regions that had either FINEMAP CP ≥ 0.5 or SuSiE CP ≥ 0.5
and binned them by MAF (bin boundaries = [0.01%, 0.1%, 10%, 50%]), excluding all variants with
MAF < 0.01%. We assigned each bin a relative weight by the proportion of causal variants in that
bin vs in all bins as compared to the proportion of all variants in that bin vs all bins, noting that
these weights were relatively consistent across bins (within a factor of 2, **Supplementary Table
6**). Using those bin weights, for each fine-mapping region, we then drew causal SNPs/indels at
random from all SNPs/indels in the region, with each variant's chance of being drawn weighted
by the bin that its MAF corresponds to. For each bin, we also collected all observed effect sizes
of all variants falling in that bin, noting that as expected the effect sizes for common variants were
smaller than those for rarer variants (**Supplementary Fig. 6**). For each variant chosen to be
causal, we drew an effect size from the corresponding MAF bin. This strategy is designed so that
the distributions of MAFs and effect sizes of causal variants in our simulations are similar to those

observed for fine-mapped variants for the real phenotype. We repeated this strategy nine times
for each simulation region, three times each choosing sets of one, two and three causal variants.

While the first strategy allows for a wide range of simulations by drawing causal variants at
random, it may not capture systematic differences between the LD patterns of causal variants
and the LD patterns of non-causal variants in causal regions. To address this, for the second
strategy we chose variants fine-mapped by SuSiE for platelet count for simulating as causal as
these may more closely capture LD patterns of truly causal variants. Specifically, we ran SuSiE
on all the SNPs and indels in the fine-mapping region with p < 0.0005 against real platelet count
data. Note that by only running SuSiE against the SNP and indel variants in the region, we forced
SuSiE to give us the most plausibly causal set of SNPs/indels in the region under the condition
that no STRs are causal. We discarded non-pure credible sets (those with variants in less than
0.8 r^2) as we expect them to be less reliable in identifying truly causal variants. In the 458/548
regions where there were any pure credible sets remaining, we took the top variant from each of
the remaining credible sets to use as causal for simulations, using their effect sizes measured
against the real platelet count trait as their effect sizes for simulation. For each region, we used
its causal variant set to simulate three phenotypes (which are distinct due to different noise terms).

While this second strategy may capture more realistic causal LD patterns compared to choosing
causal variants at random, it has the drawback that it relies on the accuracy of fine-mapping to
choose the causal variants, which is what we are trying to assess. Strategy one relies on fine-
mappers as well, but to a much lesser extent, using them only to identify causal variant MAF and
effect size distributions. A second caveat to strategy two is that by restricting to pure credible sets,
we likely omit real signals which SuSiE could not resolve well.

For our third strategy, we paralleled our second strategy, except instead of fine-mapping platelet
count against only SNPs and indels, we fine-mapped it against all the variants in the region
(including STRs), thus allowing it to select STRs as causal for simulation. We continued with
simulations as in the second strategy for the 52/548 regions where SuSiE identified a causal STR.
This third strategy is the only strategy we performed which simulated causal STRs. As the number
of simulations performed with this third strategy was limited, we only use it to contribute briefly to
our discussion in the main text.

Simulating phenotypes: Let V represent the set of causal variants for a region. For each variant
$v \in V$ let $\vec{g_v}$ represent a vector of participant genotype dosages and β_v denote the variant's

chosen effect size. Assuming additive and independent contributions of each variant, we

simulated a vector of phenotypes $(\vec{y})$ as $\vec{y} = \sum_{v \in V} \overrightarrow{g_v} * \beta_v + \vec{\epsilon}$, where $\vec{\epsilon} \sim N(0, \text{diag}(1 -$

$\sum_{v \in V} \beta_v^2 Var[\overrightarrow{g_v}]))$ so that similarly to the real, normalized, phenotypes used for our GWAS, the

resulting phenotypes have mean 0 and variance 1.

Evaluating fine-mapping on simulated phenotypes: For each simulated phenotype and region we

performed association testing of the variants in that region using the same methods as in the main

analysis, excepting that we included no covariates and that the phenotypes were not subjected

to rank-inverse normalization. We then ran FINEMAP and SuSiE against the variants in the region

as described above (in particular, FINEMAP runs were restricted to variants with $p<0.05$, SuSiE

runs to variants with $p<0.0005$), with the difference that the fine-mapping region was not

recalculated from the simulated phenotype GWAS statistics but instead exactly matched to the

causal region determined from the platelet count GWAS. Once fine-mappers were run, we

calculated STR contribution statistics as for fine-mapping runs on the UKB blood traits

(**Supplementary Tables 7-8**).

Simulation caveats: Many choices in the design of these simulations affect the interpretation of

their results. Notably, these simulated phenotypes make standard assumptions of additive genetic

architectures, including no non-linear effects, no epistasis between variants, and that the

environmental contribution to each phenotype is both independent of an individual's genotypes

and normally distributed. These simulations also assume that there are no confounding

covariates. Additionally, these simulations choose the effect sizes of causal variants from effect

sizes calculated in our platelet count GWAS. As effect sizes calculated in the GWAS were

measured in mono-variant regressions against platelet count, they will be mis-estimated

according to the corresponding variant's LD to all causal variants in the region in which it resides.

Further, we note that not recalculating the fine-mapping regions may artificially inflate the rate at

which strategy one identifies causal variants, as when causal variants in strategy one were

randomly chosen to fall near the edges of the region, there would be fewer variants in LD with

those variants and fine-mapping them would be easier. This may contribute to the observation in

Supplementary Table 7 that both fine-mappers select STRs in simulation strategy two much

more than in simulation strategy one. We also speculate that STRs truly causal for platelet count

would contribute to that observation: if those STRs are well tagged by SNPs, strategy two's run

of SuSiE would likely select those tagging SNPs for causal simulation. Then fine-mapping of those

simulated phenotypes would have a relatively high chance of confusing those SNPs with the
STRs they tag.

Lastly, we observe that FINEMAP mostly identifies variants with low p-values, while a p-value
cutoff is necessary for accurate SuSiE results. Once a p-value cutoff is applied, we see that the
fine-mappers' results are almost entirely consistent with one another, in large distinction from how
they perform when applied to real datasets, suggesting that there are some features of the
architectures of blood traits are not captured by these simulations.

Validation of imputation results for fine-mapped STRs using WGS data

We worked with WGS CRAM files for 200,025 UKB participants on the UKB Research Analysis
Platform cloud solution provided by DNA Nexus. This data was aligned to reference genome
hg38. HipSTR was unable to load the index files for the CRAM files of 10 participants, possibly
due to file corruption. Removing those participants left us with 200,015 participants. We
inadvertently truncated the participant list, leaving 200,000 participants. From that participant list
we called genotypes of the 409 STRs in **Supplementary Table 4** using HipSTR[6] in batches of
500 participants, using the flag `--min-reads 10` and allowing HipSTR to estimate stutter-error
models from the data. We merged batches using MergeSTR[73]. We performed call level filtering
using DumpSTR[73] with the flags `--hipstr-min-call-Q 0.9 --hipstr-min-call-DP 10`
`--hipstr-max-call-DP 10000 --hipstr-min-supp-reads 2 --hipstr-max-call-`
`stutter 0.15 --hipstr-max-call-flank-indel 0.1`. After calling all 200,000
individuals we summarized their genotypes separately per population, noting that 166,638
individuals were in our set of QC'ed (potentially related) White British UKB participants,
accounting for 40.8% of the QC'ed White British participants.

We did not apply any locus-level filters, such as Hardy-Weinberg equilibrium, to our WGS results.
We report per-locus WGS call rates for QCed (potentially related) individuals in each population.
We used liftOver to lift the hg38 WGS calls to the hg19 reference genome (**URLs**). To compare
the WGS calls to the imputed STR calls, we used CompareSTR from TRTools[73] branch
`compareSTR_upgrade` using the flags `--ignore-phasing --balanced-accuracy --`
`vcf2-beagle-probabilities` . We report multiple metrics at each locus, specifically
concordance, the mean absolute summed-length difference, r^2 and dosage r^2.

For the following definitions, let X be the set of all samples, A be the set of all possible STR length
alleles at a locus, let $S = \{a_1 + a_2 | a_1, a_2 \in A\}$ be the set of all summed-lengths possible at a locus

(including the case of homozygous individuals when $a_1 = a_2$), for $x \in X$ let $s_{x,WGS}$ be the summed-

length call for sample x from WGS data, and for $x \in X, s \in S$ let $Pr_{x,imp}(s)$ be the probability that

sample x has a summed imputation length of s as output by the Beagle `AP1` and `AP2 FORMAT`

fields in the imputed VCF file.

We report (summed-length) per-locus concordances as $E_{x \in X}[Pr_{x,imp}(s_{x,WGS})]$. This metric has the

advantage of being intuitive but is biased upwards for loci with a single very common allele and

so should be interpreted cautiously for such loci. We also report mean absolute summed length

differences as $E_{x \in X}[\sum_{s \in S} Pr_{x,imp}(s) \cdot |s_{x,WGS} - s|]$. This metric has similar caveats as the

concordance metric. However, for highly multi-allelic loci where concordance is low, this metric

can help quantify how close (or not) imputed calls are to the actual genotypes. We calculated r^2

as the square of the weighted Pearson correlation between $s_{x,WGS}$ and s for each sample $x \in X$

and all possible summed-lengths $s \in S$ (so that there are $|X| \cdot |S|$ total values being correlated),

weighting by the imputation probabilities $Pr_{x,imp}(s)$. This correlation measure is more comparable

across loci with different numbers of alleles than concordance. It has the downside of being less

intuitive and of being more sensitive to the WGS-vs-imputation concordance of rare long and short

alleles than the WGS-vs-imputation concordance of common average-length alleles. We report

dosage r^2 as the square of the Pearson correlation between $s_{x,WGS}$ and the dosage

$\sum_{s \in S} s \cdot Pr_{x,imp}(s)$ for each sample $x \in X$. Dosage r^2 is strictly greater than or equal to the weighted

r^2 measure. While the weighted r^2 measure more directly measures the concordance of individual

imputation probabilities with the WGS calls, the dosage r^2 measure better estimates how analyses

like GWAS, which condense imputed probabilities into dosages, will perform.

Lastly, at each locus we report the frequency of each summed-length according to WGS calls,

and for all samples with each WGS summed-length we report the probability that imputation

concurs with that length: $E_{X|s_{x,WGS} = s}[Pr_{x,imp}(s_{x,WGS})]$.

Replication in other populations

We separated the participants not in the White British group into population groups using the self-

reported ethnicities summarized by UKB showcase data field 21000 (**URLs**). This field uses UKB

showcase data coding 1001. We defined the following five populations based on those codings

(counts give the maximal number of unrelated QC'ed participants, ignoring per-phenotype

missingness):

● Black (African and Caribbean, n=7,562, codings 4, 4001, 4002, 4003)

● South Asian (Indian, Pakistani and Bangladeshi, n=7,397, codings 3001, 3002, 3003)

● Chinese (n=1,525, coding 5)

● Irish (n=11,978, coding 1002)

● Other White (White non-Irish non-British, n=15,838, coding 1003)

Self-reported ethnicities were collected from participants at three visits (initial assessment, repeat
assessment, first imaging). The above groups also exclude participants who self-reported
ethnicity at more than one visit and where their answers corresponded to more than one
population (after ignoring 'prefer not to answer' `code=-3` responses). We did not include any
participants who were neither in the White British population nor any of the above populations.
Unlike for the determination of White British participants, genetic principal components were not
used as filters for these categories.

For the association tests in these populations we applied the same procedures for sample quality
control, unrelatedness filtering, phenotype transformations, and preparing genotypes and
covariates as in the White British group. The only changes in procedure were that (a) we removed
categorical covariate values where there were fewer than 50 participants with that value, (in which
case we also removed those participants from analysis, as that would be too few to properly
control for batch effects), whereas for White British individuals we used a cutoff of 0.1% instead
and (b) we also applied this cutoff to the visit of measurement categorical covariate, resulting in
some association tests that excluded individuals whose first measurement of the phenotype
occurred outside the initial assessment visit. See **Supplementary Table 9** for details.

STRs were marked as replicating in another population (**Fig. 2**) if any of the traits confidently fine-
mapped to that STR share the same direction of effect as the White British association and
reached association p-value < 0.05 after multiple hypothesis correction (i.e. if there are three
confidently fine-mapped traits, then an STR is marked as replicating in the Black population if any
of them has association p-value < 0.05/3 = 0.0167 in the Black population).

We validated imputation STR lengths using WGS data in these populations as was done in the
White British population, and report these results in **Supplementary Tables 4** and **5**. The number
of samples in our QC'ed set that had WGS data were 2,990 Black, 3,373 South Asian, 619

Chinese, 5,174 Irish and 6,428 Other White samples, all roughly 40% of their respective
populations.

Logistic regression analysis of replication direction

We used logistic regression to quantitatively assess the impact of fine-mapping on replication
rates while controlling for discovery p-value. For this analysis, to have sufficient sample sizes, we
defined that an STR-trait association replicates in another population if it had the same direction
of effect in that population as in the White British population, regardless of the replication p-value.

For each of the five replication populations, we compared four categories: all gwsig (genome-
wide significant associations in the discovery population, i.e. p-value < 5e-8), FINEMAP
(discovery p-value < 5e-8 and FINEMAP CP ≥ 0.8), SuSiE (discovery p-value < 5e-8 and SuSiE
CP ≥ 0.8) and confidently fine-mapped STR (STR associations in our confidently fine-mapped
set).

For each comparison, we used the function `statsmodels.formula.api.logit` from
statsmodels v0.13.2 (**URLs**) to fit the logistic regression model:

`replication_status ~ STR_in_target_category` + $\log_{10}$`(p-val)` + $\log_{10}$`(p-`
`val)`^{2}

where `replication_status` is a binary variable indicating whether or not the given STR-trait
association replicated in the other population, `p-val` is the discovery p-value, and
`STR_in_target_category` is a binary variable indicating if the STR is in the target category.

For each replication population, we considered various models:

• All gwsig STRs with either FINEMAP, SuSiE, or confidently fine-mapped STRs as the
target category.
• All FINEMAP STRs with confidently fine-mapped STRs as the target category.
• All SuSiE STRs with confidently fine-mapped STRs as the target category.

For each model, we performed a one-sided t-test for the hypothesis that the coefficient for the
covariate `STR_in_target_category` was greater than zero, i.e. testing that being in the target
category increased the predicted chance of replicating in the chosen population (**Supplementary**
Table 10).

Gene and transcription factor binding annotations

For all analyses not using GTEx data, gene annotations were based on GENCODE 38[75] (**URLs**).

Transcription factor binding sites and DNaseI hypersensitivity regions were identified by

ENCODE[76] overlapping several loci (*TAOK1*, *RHOT1* and *NCK2*) through visual inspection of the

"Txn Factor ChIP" and "DNase Clusters" tracks in the UCSC Genome Browser[77] and using the

"Load from ENCODE" feature of the Integrative Genomics Viewer[78].

Enrichment testing

We tested the following categories for enrichment in STRs identified by our association testing

pipeline:

• Genomic feature: We grouped records by feature type and restricted to features with

support level 1 or 2 except for genes which don't have a support level. We used bedtools[79]

to compute which features intersect each STR and the distance between each STR and

the nearest feature of each feature type.

• Repeat unit: unit length and standardized repeat unit were defined as described above.

Repeat units occurring in <1000 STRs were grouped by repeat length. Repeats whose

unit could not be determined were considered as a separate category.

• Overlap with expression STRs (eSTR): we tested for overlap with either all eSTRs or fine-

mapped eSTRs as defined in our previous study to identify STR-gene expression

associations in the Genotype Tissue Expression (GTEx) cohort[13].

Enrichment p-values were computed using a Chi-squared test (without Yate's continuity

correction) if all cells had counts ≥ 5. A two-sided Fisher's exact test was used otherwise. Chi-

squared and Fisher's exact tests were implemented using the `chi2_contingency` and

`fisher_exact` functions from the Python scipy.stats package v1.7.3 (**URLs**).

Expression association analysis in GTEx

We had previously analyzed associations[13] between STRs and gene expression in GTEx V7.

Here we reanalyzed those associations using GTEx V8. We obtained 30x Illumina whole genome

sequencing (WGS) data from 652 unrelated participants in the Genotype-Tissue Expression

project (GTEx)[37] through dbGaP accession number phs000424.v8.p2. WGS data was accessed

using fusera (**URLs**) through Amazon Web Services. We genotyped STRs using HipSTR[6] v0.5

with HipSTR's hg38 reference STR set (**URLs**). All individuals were genotyped jointly using
default parameters. GTEx's whole genome sequencing procedure is not PCR-free, which likely
contributed to low call rates at long poly-A and GC-rich STRs. The resulting VCFs were filtered
using DumpSTR from TRTools[73], using the parameters `--filter-hrun --hipstr-min-`
`call-Q 0.9 --hipstr-min-call-DP 10 --hipstr-max-call-DP 1000 --hipstr-`
`max-call-flank-indel 0.15 --hipstr-max-call-stutter 0.15 --min-locus-`
`callrate 0.8 --min-locus-hwep 0.00001` . We also removed STRs overlapping
segmental duplication regions (UCSC Genome Browser[80] h38.genomicSuperDups table).
Altogether, 728,090 STRs remained for downstream analysis.

The *TAOK1* STR locus was filtered from this genotyping for having an 11% call rate, so we
imputed the genotypes at that locus into the GTEx cohort. GTEx V7 SNP files were downloaded
from GTEx data portal (**URLs**). SNPs on chromosome 17 were extracted and filtered to remove
using vcftools with the parameters `--maf 0.01 --mac 3 --we 0.00001 --max-missing`
`0.8 --minQ 30` . We used Beagle v5.2 (beagle.28Jun21.220.jar) with the tool's provided human
genetic maps to impute STRs into the GTEx SNPs using the same reference panel used for
imputation in the UKB cohort above[30]. From this imputation we took the best-guess genotypes of
the *TAOK1* STR. We lifted the coordinates of the *TAOK1* STR from hg19 to hg38 using liftOver.

For each tissue, we obtained gene-level and transcript-level transcripts-per-million (TPM) values,
exon-exon junction read counts, and exon read counts for each participant from GTEx Analysis
V8 publicly available from the GTEx project website (**URLs**). Gene annotations are based on
GENCODE v26[75]. We focused on 41 tissues with expression data for at least 100 samples
(**Supplementary Table 13**). We restricted our analysis to protein-coding genes, transcripts and
exons that did not overlap segmental duplication regions.

To control for population structure, we obtained publicly available genotype data on 2,504
unrelated individuals from the 1000 Genomes project[23] genotyped with Omni 2.5 SNP genotyping
arrays. We performed the following principal components analysis jointly on that data and the
SNP genotypes based on WGS of the 652 individuals above. We removed all indels, multi-allelic
SNPs, and SNPs with minor allele frequency less than 5%. We then used plink v.1.90b3.44 to
subset these remaining SNPs to a set of SNPs in approximate linkage equilibrium with the
command `--indep 50 5 2`. We excluded any remaining SNPs with missingness rate 5% or
greater. We lastly ran principal component analysis using smartpca70[81] v.13050 with default
parameters.

We removed genes with TPM less than 1 in more than 90 percent of individuals. PEER factors[82]

were calculated using PEER v1.0 from the TPM values which remained after filtering. For each

gene, we tested for association with each STR within 100kb. For each test we performed a linear

regression between the STR's dosage (sum of allele lengths) and gene expression (TPM). We

included the loadings of the top five genotype principal components as computed above and the

top N/10 PEER factors as covariates. The number of PEER factors was chosen to maximize the

number of significant associations across a range of tissues. We did not include sex or age as

covariates.

For each STR we computed Bonferroni-adjusted p-values to control for the number of gene ×

tissue tests performed for that STR. Associations that remained with adjusted $p < 0.05$ are shown

in **Supplementary Table 12**.

We additionally used the GTEx cohort to test for an association between length of the bilirubin-

associated dinucleotide repeat identified in *SLC2A2* with splicing efficiency in liver. We obtained

exon-exon junction read counts and exon read counts from the GTEx website (**URLs**). We

calculated the percent spliced in value for each exon in the manner suggested by Schafer et al.[83].

We performed a linear regression to test between the STR's dosage and the percent spliced in of

each exon within 10kb, using the top 5 ancestry principal components as covariates.

Methylation association analysis in GTEx

This analysis used the same STR data and genotype principal components as the GTEx

expression association analysis above.

We downloaded genome-wide DNA methylation (DNAm) profiling results from the NCBI GEO

database under accession number GSE213478. This contained DNA methylation levels from the

whole blood of 47 individuals who had been genotyped, including 754,054 autosomal CpG loci

which passed quality control checks in that dataset (**URLs**)[84]. We lifted those loci from hg19 to

hg38. We performed per-locus inverse-normalization of the DNAm data prior to downstream

analysis. We calculated 5 PEER factors from the normalized DNAm data across quality-controlled

loci from all chromosomes (including sex chromosomes) using PEER v1.0[82], choosing 5 factors

to match the number of PEER factors used by the methylation study which generated this data[84].

We tested for associations between the methylation of each autosomal CpG locus and the length

of each STR located within 100kb of that locus. For each such pair, we performed a linear

regression between the STR's dosage (sum of allele lengths across both chromosomes) and the
inverse-normalized DNAm levels of that CpG locus, including the top five genotype principal
components and the 5 PEER factors as covariates. We compared the effect sizes of these
associations with those from another paper studying STR-methylation correlations in two separate
cohorts in whole blood[17] and found that they were broadly consistent (r=0.73, p<10^{-200},
Supplementary Fig. 18c).

For each STR we computed Bonferroni-adjusted p-values to control for the number of CpG tests
performed for that STR. Associations that remained with adjusted p < 0.05 are shown in
Supplementary Table 14.

Expression analysis of the* CBL *and* RHOT1 *STRs in Geuvadis

We applied HipSTR[6] v0.6.2 to genotype STRs from HipSTR's hg38 reference STR set (**URLs**) in
2,504 individuals from the 1000 Genomes Project[85] for which high-coverage WGS data was
available. Gene-level reads per kilobase per million reads (RPKM) values based on RNA-seq in
lymphoblastoid cell lines for 462 1000 Genomes participants were downloaded from the Geuvadis
website (**URLs**). Of these, 449 individuals were genotyped by HipSTR.

Similar to the GTEx analysis, we performed a linear regression between STR dosage (sum of
allele lengths) and RPKM, adjusting for the top 5 genotype principal components (computed as
above for the GTEx analysis, but only on populations included in Geuvadis and separately for
Europeans and Africans) and N/10 (45) PEER factors as covariates. PEER analysis was applied
using PEER v1.0 to the matrix of RPKM values after removing genes overlapping segmental
duplications and those with RPKM less than 1 in more than 90% of LCL samples. We performed
a separate regression analysis for African individuals (YRI) and European individuals (CEU, TSI,
FIN, and GBR). After restricting to individuals with non-missing expression data and STR
genotypes and who were not filtered as PCA outliers by smartpca[81,86] included in EIGENSOFT
v6.1.4, 447 LCL samples remained for analysis in each case (num. EUR=358, and AFR=89 for
CBL, EUR=359 and AFR=88 for *RHOT1*).

URLs

- 1000 Genomes individuals: https://www.internationalgenome.org/data-portal/sample using the "Download the list" tab
- associaTR: https://trtools.readthedocs.io/
- Beagle Human genetic maps: https://bochet.gcc.biostat.washington.edu/beagle/genetic_maps/
- fusera: https://github.com/ncbi/fusera
- GENCODE 38 (hg19): http://ftp.ebi.ac.uk/pub/databases/gencode/Gencode_human/release_38/GRCh37_mapping/gencode.v38lift37.annotation.gff3.gz
- Geuvadis: https://www.ebi.ac.uk/arrayexpress/experiments/E-GEUV-1/files/analysis_results/?ref=E-GEUV-1
- GTEx v8:
  - https://www.gtexportal.org/home/datasets
  - https://storage.googleapis.com/gtex_analysis_v8/rna_seq_data/GTEx_Analysis_2017-06-05_v8_RNASeQCv1.1.9_gene_tpm.gct.gz
  - https://storage.googleapis.com/gtex_analysis_v8/rna_seq_data/GTEx_Analysis_2017-06-05_v8_STARv2.5.3a_junctions.gct.gz
  - https://storage.googleapis.com/gtex_analysis_v8/rna_seq_data/GTEx_Analysis_2017-06-05_v8_RNASeQCv1.1.9_exon_reads.parquet
- h5py https://github.com/h5py/h5py
- HDF5: https://www.hdfgroup.org/HDF5/
- HipSTR STR reference https://github.com/HipSTR-Tool/HipSTR-references/raw/master/human/hg38.hipstr_reference.bed.gz
- liftOver:
  - https://genome.ucsc.edu/cgi-bin/hgLiftOver , accessed on 2023/03/09
  - https://hgdownload.soe.ucsc.edu/goldenPath/hg19/liftOver/hg19ToHg38.over.chain.gz
  - ftp://hgdownload.soe.ucsc.edu/goldenPath/hg38/liftOver/hg38ToHg19.over.chain.gz
- Methylation in GTEx:
  - Overview: https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE213478
  - CpG locations: https://www.ncbi.nlm.nih.gov/geo/download/?acc=GSE213478&format=file

•   NCBI protein database: https://www.ncbi.nlm.nih.gov/protein

•   Pan-UKB:

○   Overview: https://pan.ukbb.broadinstitute.org/downloads

○   manifest:

https://docs.google.com/spreadsheets/d/1AeeADtT0U1AukliiNyiVzVRdLYPkTbru

QSk38DeutU8

○   bilirubin SNP summary statistics: https://pan-ukb-us-east-

1.s3.amazonaws.com/sumstats_flat_files/biomarkers-30840-both_sexes-

irnt.tsv.bgz and https://pan-ukb-us-east-

1.s3.amazonaws.com/sumstats_flat_files_tabix/biomarkers-30840-both_sexes-

irnt.tsv.bgz.tbi

•   Plink association testing best practices: https://www.cog-

genomics.org/plink/2.0/assoc#glm

•   PyMOL: https://pymol.org/2/

•   rhdf5: https://www.bioconductor.org/packages/release/bioc/html/rhdf5.html

•   Scipy.stats: https://docs.scipy.org/doc/scipy/reference/stats.html

•   SNP-STR reference panel: https://gymreklab.com/2018/03/05/snpstr_imputation.html

•   Statsmodels: https://www.statsmodels.org/stable/index.html

•   UKB Data Showcase Search Page: https://biobank.ctsu.ox.ac.uk/crystal/search.cgi

1293

Acknowledgments

Research reported in this publication was supported in part by NIH/NHGRI grants R01HG010885

(M.G. and A.G.) and 1RM1HG011558 (M.G.). This research has been conducted using the UK

Biobank Resource under application number 46122. We thank R. Wachs for helping with

illustrations. We also thank K. Frazer and M. D'Antonio for helpful discussions and comments.

GTEx Project: The Genotype-Tissue Expression (GTEx) Project was supported by the Common

Fund of the Office of the Director of the National Institutes of Health (commonfund.nih.gov/GTEx).

Additional funds were provided by the NCI, NHGRI, NHLBI, NIDA, NIMH, and NINDS. Donors

were enrolled at Biospecimen Source Sites funded by NCI\Leidos Biomedical Research, Inc.

subcontracts to the National Disease Research Interchange (10XS170), GTEx Project March 5,

2014 version Page 5 of 8 Roswell Park Cancer Institute (10XS171), and Science Care, Inc.

(X10S172). The Laboratory, Data Analysis, and Coordinating Center (LDACC) was funded

through a contract (HHSN268201000029C) to the The Broad Institute, Inc. Biorepository

operations were funded through a Leidos Biomedical Research, Inc. subcontract to Van Andel
Research Institute (10ST1035). Additional data repository and project management were
provided by Leidos Biomedical Research, Inc. (HHSN261200800001E). The Brain Bank was
supported supplements to University of Miami grant DA006227. Statistical Methods development
grants were made to the University of Geneva (MH090941 & MH101814), the University of
Chicago (MH090951,MH090937, MH101825, & MH101820), the University of North Carolina -
Chapel Hill (MH090936), North Carolina State University (MH101819),Harvard University
(MH090948), Stanford University (MH101782), Washington University (MH101810), and to the
University of Pennsylvania (MH101822). The datasets used for the analyses described in this
manuscript were obtained from dbGaP at http://www.ncbi.nlm.nih.gov/gap through dbGaP
accession numbers phs000424.v7.p2 and phs000424.v8.p2.

Author contributions

J.M. led, designed and performed the analyses and wrote the manuscript. S.F. helped oversee
physiological interpretation of individual signals. Y.L. performed expression and methylation
analyses of the GTEx data. X.Z. performed analyses of protein-coding STRs with AlphaFold. A.M.
assisted with analysis of the *APOB* locus. A.G. and M.G. conceived the study, supervised
analyses, and wrote the manuscript. All authors read and approved this manuscript.

Competing interests

The authors have no competing financial interests to declare.

Code availability

The associaTR tool has been published as part of the TRTools[73] package and tool-suite
(https://trtools.readthedocs.io/). Code for performing most of the analyses and generating most
of the figures in this paper can be found at
https://github.com/LiterallyUniqueLogin/ukbiobank_strs/

Data Availability

Full results from our analyses can be found at the URL
https://gymreklab.com/science/2023/09/08/Margoliash-et-al-paper.html

      We are in the process of uploading imputed STR calls to the UK Biobank.
1334

References cited

1. Visscher, P. M. *et al.* 10 Years of GWAS Discovery: Biology, Function, and Translation. *Am. J. Hum. Genet.* **101**, 5–22 (2017).

2. Mukamel, R. E. *et al.* Protein-coding repeat polymorphisms strongly shape diverse human phenotypes. *Science* **373**, 1499–1505 (2021).

3. Grünewald, T. G. P. *et al.* Chimeric EWSR1-FLI1 regulates the Ewing sarcoma susceptibility gene EGR2 via a GGAA microsatellite. *Nat. Genet.* **47**, 1073–1078 (2015).

4. Sekar, A. *et al.* Schizophrenia risk from complex variation of complement component 4. *Nature* **530**, 177–183 (2016).

5. Boettger, L. M. *et al.* Recurring exon deletions in the HP (haptoglobin) gene contribute to lower blood cholesterol levels. *Nat. Genet.* **48**, 359–366 (2016).

6. Willems, T. *et al.* Genome-wide profiling of heritable and de novo STR variations. *Nat. Methods* **14**, 590–592 (2017).

7. Ellegren, H. Microsatellites: simple sequences with complex evolution. *Nat. Rev. Genet.* **5**, 435–445 (2004).

8. Sun, J. X. *et al.* A direct characterization of human mutation based on microsatellites. *Nat. Genet.* **44**, 1161–1165 (2012).

9. Lynch, M. Rate, molecular spectrum, and consequences of human mutation. *Proc. Natl. Acad. Sci.* **107**, 961–968 (2010).

10. Mirkin, S. M. Expandable DNA repeats and human disease. *Nature* **447**, 932–940 (2007).

11. Malik, I., Kelley, C. P., Wang, E. T. & Todd, P. K. Molecular mechanisms underlying nucleotide repeat expansion disorders. *Nat. Rev. Mol. Cell Biol.* **22**, 589–607 (2021).

12. Quilez, J. *et al.* Polymorphic tandem repeats within gene promoters act as modifiers of gene expression and DNA methylation in humans. *Nucleic Acids Res.* **44**, 3750–3762 (2016).

13.   Fotsing, S. F. *et al.* The impact of short tandem repeat variation on gene expression.

Nat. Genet. **51**, 1652–1659 (2019).

14.   Tw, H., Jd, G., Ce, Y. & Gr, C. A variable dinucleotide repeat in the CFTR gene

contributes to phenotype diversity by forming RNA secondary structures that alter splicing.

Proc. Natl. Acad. Sci. U. S. A. **101**, (2004).

15.   Hui, J., Stangl, K., Lane, W. S. & Bindereif, A. HnRNP L stimulates splicing of the eNOS

gene by binding to variable-length CA repeats. *Nat. Struct. Biol.* **10**, 33–37 (2003).

16.   Vinces, M. D., Legendre, M., Caldara, M., Hagihara, M. & Verstrepen, K. J. Unstable

tandem repeats in promoters confer transcriptional evolvability. *Science* **324**, 1213–1216

(2009).

17.   Martin-Trujillo, A., Garg, P., Patel, N., Jadhav, B. & Sharp, A. J. Genome-wide

evaluation of the effect of short tandem repeat variation on local DNA methylation. *Genome*

Res. **33**, 184–196 (2023).

18.   Murat, P., Guilbaud, G. & Sale, J. E. DNA polymerase stalling at structured DNA

constrains the expansion of short tandem repeats. *Genome Biol.* **21**, 209 (2020).

19.   Rothenburg, S., Koch-Nolte, F., Rich, A. & Haag, F. A polymorphic dinucleotide repeat in

the rat nucleolin gene forms Z-DNA and inhibits promoter activity. *Proc. Natl. Acad. Sci.* **98**,

8985–8990 (2001).

20.   Freudenreich, C. H. R-loops: Targets for Nuclease Cleavage and Repeat Instability.

Curr. Genet. **64**, 789–794 (2018).

21.   Niehrs, C. & Luke, B. Regulatory R-loops as facilitators of gene expression and genome

stability. *Nat. Rev. Mol. Cell Biol.* **21**, 167–178 (2020).

22.   McCarthy, S. *et al.* A reference panel of 64,976 haplotypes for genotype imputation. *Nat.*

Genet. **48**, 1279–1283 (2016).

23.   Auton, A. *et al.* A global reference for human genetic variation. *Nature* **526**, 68–74

(2015).

24.   Huang, J. *et al.* Improved imputation of low-frequency and rare variants using the UK10K

1387   haplotype reference panel. *Nat. Commun.* **6**, 8111 (2015).

1388   25.   Dashnow, H. *et al.* STRetch: detecting and discovering pathogenic short tandem repeat

1389   expansions. *Genome Biol.* **19**, 1–13 (2018).

1390   26.   Mousavi, N., Shleizer-Burko, S., Yanicky, R. & Gymrek, M. Profiling the genome-wide

1391   landscape of tandem repeat expansions. *Nucleic Acids Res.* **47**, e90 (2019).

1392   27.   Dolzhenko, E. *et al.* ExpansionHunter: a sequence-graph-based tool to analyze variation

1393   in short tandem repeat regions. *Bioinformatics* **35**, 4754–4756 (2019).

1394   28.   Tang, H. *et al.* Profiling of Short-Tandem-Repeat Disease Alleles in 12,632 Human

1395   Whole Genomes. *Am. J. Hum. Genet.* **101**, 700–715 (2017).

1396   29.   Tankard, R. M. *et al.* Detecting Expansions of Tandem Repeats in Cohorts Sequenced

1397   with Short-Read Sequencing Data. *Am. J. Hum. Genet.* **103**, 858–873 (2018).

1398   30.   Saini, S., Mitra, I., Mousavi, N., Fotsing, S. F. & Gymrek, M. A reference haplotype panel

1399   for genome-wide imputation of short tandem repeats. *Nat. Commun.* **9**, 4397 (2018).

1400   31.   Bycroft, C. *et al.* The UK Biobank resource with deep phenotyping and genomic data.

1401   *Nature* **562**, 203–209 (2018).

1402   32.   Browning, B. L., Zhou, Y. & Browning, S. R. A One-Penny Imputed Genome from Next-

1403   Generation Reference Panels. *Am. J. Hum. Genet.* **103**, 338–348 (2018).

1404   33.   Chang, C. C. *et al.* Second-generation PLINK: rising to the challenge of larger and richer

1405   datasets. *GigaScience* **4**, (2015).

1406   34.   Pan-UKB team. Pan-ancestry genetic analysis of the UK Biobank.

1407   https://pan.ukbb.broadinstitute.org/ (2020).

1408   35.   Wang, G., Sarkar, A., Carbonetto, P. & Stephens, M. A simple new approach to variable

1409   selection in regression, with application to genetic fine mapping. *J. R. Stat. Soc. Ser. B Stat.*

1410   *Methodol.* **82**, 1273–1300 (2020).

36.   Benner, C. *et al.* FINEMAP: efficient variable selection using summary data from

genome-wide association studies. *Bioinformatics* **32**, 1493–1501 (2016).

37.   THE GTEX CONSORTIUM. The GTEx Consortium atlas of genetic regulatory effects

across human tissues. *Science* **369**, 1318–1330 (2020).

38.   Berberich, A. J. & Hegele, R. A. A Modern Approach to Dyslipidemia. *Endocr. Rev.*

bnab037 (2021) doi:10.1210/endrev/bnab037.

39.   Boerwinkle, E. & Chan, L. A three codon insertion/deletion polymorphism in the signal

peptide region of the human apolipoprotein B (APOB) gene directly typed by the polymerase

chain reaction. *Nucleic Acids Res.* **17**, 4003 (1989).

40.   Niu, C. *et al.* Associations of the APOB rs693 and rs17240441 polymorphisms with

plasma APOB and lipid levels: a meta-analysis. *Lipids Health Dis.* **16**, 166 (2017).

41.   Hui, J. *et al.* Intronic CA-repeat and CA-rich elements: a new class of regulators of

mammalian alternative splicing. *EMBO J.* **24**, 1988–1998 (2005).

42.   Huang, Y. *et al.* Mediator complex regulates alternative mRNA processing via the

MED23 subunit. *Mol. Cell* **45**, 459–469 (2012).

43.   Sutcliffe, J. S. *et al.* DNA methylation represses FMR-1 transcription in fragile X

syndrome. *Hum. Mol. Genet.* **1**, 397–400 (1992).

44.   Garg, P. *et al.* A Survey of Rare Epigenetic Variation in 23,116 Human Genomes

Identifies Disease-Relevant Epivariations and CGG Expansions. *Am. J. Hum. Genet.* **107**,

654–669 (2020).

45.   Saur, S. J., Sangkhae, V., Geddis, A. E., Kaushansky, K. & Hitchcock, I. S.

Ubiquitination and degradation of the thrombopoietin receptor c-Mpl. *Blood* **115**, 1254–1263

(2010).

46.   Plo, I. *et al.* Genetic Alterations of the Thrombopoietin/MPL/JAK2 Axis Impacting

Megakaryopoiesis. *Front. Endocrinol.* **8**, 234 (2017).

47. Kaushansky, K. *et al.* Promotion of megakaryocyte progenitor expansion and

differentiation by the c-Mpl ligand thrombopoietin. *Nature* **369**, 568–571 (1994).

48. Märklin, M. *et al.* C-Cbl regulates c-MPL receptor trafficking and its internalization. *J.*

Cell. Mol. Med. **24**, 12491–12503 (2020).

49. Lappalainen, T. *et al.* Transcriptome and genome sequencing uncovers functional

variation in humans. *Nature* **501**, 506–511 (2013).

50. Shinjyo, T. *et al.* Downregulation of Bim, a Proapoptotic Relative of Bcl-2, Is a Pivotal

Step in Cytokine-Initiated Survival Signaling in Murine Hematopoietic Progenitors. *Mol. Cell.*

Biol. **21**, 854–864 (2001).

51. Kotzin, J. J. *et al.* The long non-coding RNA Morrbid regulates Bim and short-lived

myeloid cell lifespan. *Nature* **537**, 239–243 (2016).

52. Bouillet, P. *et al.* Proapoptotic Bcl-2 Relative Bim Required for Certain Apoptotic

Responses, Leukocyte Homeostasis, and to Preclude Autoimmunity. *Science* **286**, 1735–

1738 (1999).

53. Draviam, V. M. *et al.* A functional genomic screen identifies a role for TAO1 kinase in

spindle-checkpoint signalling. *Nat. Cell Biol.* **9**, 556–564 (2007).

54. Favier, R. & Raslova, H. Progress in understanding the diagnosis and molecular

genetics of macrothrombocytopenias. *Br. J. Haematol.* **170**, 626–639 (2015).

55. Azad, P., Villafuerte, F. C., Bermudez, D., Patel, G. & Haddad, G. G. Protective role of

estrogen against excessive erythrocytosis in Monge's disease. *Exp. Mol. Med.* **53**, 125–135

(2021).

56. Mukundan, H., Resta, T. C. & Kanagy, N. L. 17β-Estradiol decreases hypoxic induction

of erythropoietin gene expression. *Am. J. Physiol.-Regul. Integr. Comp. Physiol.* **283**, R496–

R504 (2002).

57. Krusche, P. *et al.* Best practices for benchmarking germline small-variant calls in human

genomes. *Nat. Biotechnol.* **37**, 555–560 (2019).

58.   Hannan, A. J. Tandem repeat polymorphisms: modulators of disease susceptibility and

candidates for 'missing heritability'. *Trends Genet.* **26**, 59–65 (2010).

59.   Press, M. O., Carlson, K. D. & Queitsch, C. The overdue promise of short tandem repeat

variation for heritability. *Trends Genet.* **30**, 504–512 (2014).

60.   D'Antonio, M. *et al.* Systematic genetic analysis of the MHC region reveals mechanistic

underpinnings of HLA type associations with disease. *eLife* **8**, e48476 (2019).

61.   Chiang, C. *et al.* The impact of structural variation on human gene expression. *Nat.*

Genet. **49**, 692–699 (2017).

62.   Halldorsson, B. V. *et al.* The sequences of 150,119 genomes in the UK Biobank. *Nature*

607, 732–740 (2022).

63.   Foix, Anna & Blachly, James. pyEGA3: EGA download client. (2021).

64.   O'Connell, J. *et al.* Haplotype estimation for biobank-scale data sets. *Nat. Genet.* **48**,

817–820 (2016).

65.   Horta, D. bgen-reader: Bgen file format reader.

66.   Staples, J., Nickerson, D. A. & Below, J. E. Utilizing Graph Theory to Select the Largest

Set of Unrelated Individuals for Genetic Analysis. *Genet. Epidemiol.* **37**, 136–141 (2013).

67.   Beasley, T. M., Erickson, S. & Allison, D. B. Rank-Based Inverse Normal

Transformations are Increasingly Used, But are They Merited? *Behav. Genet.* **39**, 580–595

(2009).

68.   Bishara, A. J. & Hittner, J. B. Testing the significance of a correlation with nonnormal

data: comparison of Pearson, Spearman, transformation, and resampling approaches.

Psychol. Methods **17**, 399–417 (2012).

69.   Zwiener, I., Frisch, B. & Binder, H. Transforming RNA-Seq Data to Improve the

Performance of Prognostic Gene Signatures. *PLOS ONE* **9**, e85150 (2014).

70.   Bishara, A. J. & Hittner, J. B. Reducing Bias and Error in the Correlation Coefficient Due

to Nonnormality. *Educ. Psychol. Meas.* **75**, 785–804 (2015).

71.   Auer, P. L., Reiner, A. P. & Leal, S. M. The effect of phenotypic outliers and non-

normality on rare-variant association testing. *Eur. J. Hum. Genet.* **24**, 1188–1194 (2016).

72.   Pedersen, B. cyvcf2: fast vcf parsing with cython + htslib.

73.   Mousavi, N. *et al.* TRTools: a toolkit for genome-wide analysis of tandem repeats.

Bioinformatics **37**, 731–733 (2021).

74.   Zheng, J., Li, Y., Abecasis, G. R. & Scheet, P. A Comparison of Approaches to Account

for Uncertainty in Analysis of Imputed Genotypes. *Genet. Epidemiol.* **35**, 102–110 (2011).

75.   Frankish, A. *et al.* GENCODE reference annotation for the human and mouse genomes.

Nucleic Acids Res. **47**, D766–D773 (2019).

76.   Dunham, I. *et al.* An integrated encyclopedia of DNA elements in the human genome.

Nature **489**, 57–74 (2012).

77.   Kent, W. J. *et al.* The Human Genome Browser at UCSC. *Genome Res.* **12**, 996–1006

(2002).

78.   Robinson, J. T. *et al.* Integrative genomics viewer. *Nat. Biotechnol.* **29**, 24–26 (2011).

79.   Quinlan, A. R. & Hall, I. M. BEDTools: a flexible suite of utilities for comparing genomic

features. *Bioinformatics* **26**, 841–842 (2010).

80.   Karolchik, D. *et al.* The UCSC Table Browser data retrieval tool. *Nucleic Acids Res.* **32**,

D493–D496 (2004).

81.   Price, A. L. *et al.* Principal components analysis corrects for stratification in genome-

wide association studies. *Nat. Genet.* **38**, 904–909 (2006).

82.   Stegle, O., Parts, L., Piipari, M., Winn, J. & Durbin, R. Using probabilistic estimation of

expression residuals (PEER) to obtain increased power and interpretability of gene

expression analyses. *Nat. Protoc.* **7**, 500–507 (2012).

83.   Schafer, S. *et al.* Alternative Splicing Signatures in RNA-seq Data: Percent Spliced in

(PSI). *Curr. Protoc. Hum. Genet.* **87**, 11.16.1-11.16.14 (2015).

84.   Oliva, M. *et al.* DNA methylation QTL mapping across diverse human tissues provides

molecular links between genetic variation and complex traits. *Nat. Genet.* **55**, 112–122

(2023).

85.   Byrska-Bishop, M. *et al.* High coverage whole genome sequencing of the expanded

1000 Genomes Project cohort including 602 trios. 2021.02.06.430068 Preprint at

https://doi.org/10.1101/2021.02.06.430068 (2021).

86.   Patterson, N., Price, A. L. & Reich, D. Population Structure and Eigenanalysis. *PLOS*

Genet. **2**, e190 (2006).

1521